Genetic and thermal variation influences adaptation to fluctuating

temperature in the seed beetle, *Callosobruchus maculatus*

Edward Ivimey-Cook[1], Claudio Piani[2], Wei-Tse Hung[2] and Elena C. Berg[2]

[1]School of Biological Sciences, University of East Anglia, Norwich Research Park,

Norwich, UK

[2]Department of Computer Science, Mathematics, and Environmental Science, The

American University of Paris, France

* Corresponding author:

e-mail: e.ivimeycook@gmail.com

orcid: 0000-0003-4910-0443

Keywords: thermal adaptation, fluctuating temperatures, stress, climate change,

Callosobruchus maculatus

Summary

The impacts of climate change on biological systems are notoriously difficult to

measure, and laboratory studies often do not realistically represent natural

fluctuations in environmental conditions. To date, most experimental studies of

thermal adaptation test populations at constant temperatures, or they make

incremental changes to an otherwise constant mean background state. To

address this, we examined the long-term effects of stressful fluctuating daily

temperature on several key life history traits in two laboratory populations of the

seed beetle, *Callosobruchus maculatus*. These populations were kept for 19

generations at either a constant control temperature, T=29°C, or at a fluctuating

daily cycle with T_{mean} = 33°C, T_{max}=40, and T_{min}=26. The latter being a simple

representation of daily temperatures in southern central India in May, where this

species originally evolved. We found that beetles that had evolved in stressful

environments were smaller in body size when switched to a constant 29°C and

had far greater reproductive fitness in comparison to beetles from both the

constant control and continuously stressful 33°C environments. This suggests

that beetles raised in environments with stressful fluctuating temperatures were

more phenotypically plastic and had greater genetic variability than control

treatment beetles and indicates that populations that experience fluctuations in

temperature may be better able to respond to short-term changes in

environmental conditions.

Introduction

An increase in mean temperature negatively impacts a multitude of life history

traits in both plants (Hedhly *et al.*, 2009; Hatfield & Prueger, 2015) and animals

(Bauerfeind & Fischer, 2014; Berger *et al.*, 2017; Chen *et al.*, 2018; Lopez &

Adair, 2019; Ivimey-Cook *et al.*, 2021). In particular, heat stress caused by high

temperatures outside of a species' thermal optimum can disrupt reproduction in

mammals (Hansen, 2009) and fish (Lopez & Adair, 2019); inhibit growth and

development in plants (Hatfield & Prueger, 2015); and reduce longevity and

survival in insects (Berger *et al.*, 2017; Ivimey-Cook *et al.*, 2021).

Elevated temperatures, one of the main outcomes of rising anthropogenic

gas emissions (Lashof & Ahuja, 1990; Pereira *et al.*, 2012; Pachauri *et al.*, 2014),

can influence populations both directly, by increasing the rate of genome wide *de

novo* mutations (Berger *et al.*, 2017, 2021), or indirectly, by triggering

phenological mismatches between species and prey (Visser *et al.*, 1998, 2012;

see also Parmesan, 2006 for an extensive review). This increase in

environmental stress leads populations to either adapt to novel conditions through

phenotypic plasticity, to shift their geographic boundaries to match thermal

tolerance, or to go extinct (Holt, 1990; Pereira *et al.*, 2012; Hetem *et al.*, 2014;

Trisos *et al.*, 2020). Species in tropical habitats are predicted to be especially

vulnerable, as they have evolved far more narrow and specialized thermal ranges

(Janzen, 1967; Deutsch *et al.*, 2008; Walters *et al.*, 2012).

However, not only are the Earth's mean temperatures rising, but they are

also becoming more variable (Folguera *et al.*, 2011; Vasseur *et al.*, 2014;

Masson-Delmotte *et al.*, 2021). This increase in environmental variability is likely

to intensify the already detrimental impacts of elevated temperatures (Vasseur *et

al.*, 2014). Building on work by Deutsch *et al.* (2008), Vasseur *et al.* (2014) found

that changes to both the mean and variation in environmental temperature

produced long-term changes to population fitness. More specifically, performance

data collected from 38 globally distributed invertebrate species indicated that

whilst increasing mean temperature had a mixture of both positive and negative

effects on performance, which were largely latitude-dependent, increased

variation was associated with more negative effects (Vasseur *et al.*, 2014). In

addition, the authors highlighted the relative contribution of mean and variance in

determining projected performance in the years 2050-2059. They found that a

change in mean temperature explained only 32% of future species performance,

whereas combining both the mean and variance simultaneously explained 93%.

This highlights the importance of understanding how species react not only to

changes in mean temperature, but also to changes in temperature *variation* in

order to form accurate predictions of a population's response to climate change.

Studying the effects of climate warming and the resulting adaptive

responses of organisms is often confined either to field studies, which normally

consist of a multitude of uncontrollable ecological variables, or to laboratory

studies, which have the benefit of increased experimental control but typically rely

on the use of constant mean temperatures with very little consideration of natural

fluctuations (Fischer *et al.*, 2011; Folguera *et al.*, 2011; Thompson *et al.*, 2013).

As a result, populations could respond in an entirely different manner when

exposed to fluctuating rather than constant thermal stress in the laboratory

(Folguera *et al.*, 2011; Bauerfeind & Fischer, 2014). For example, reaction norms

of the mosquito *Anopheles stephensi* under fluctuating thermal regimes were

fundamentally different to those under constant temperatures (Paajimans *et al.,*

2013). Similarly, in the tropical butterfly *Bicyclus anyana*, life history traits were

strongly influenced by both the mean and the amplitude of the experimental

thermoperiod (Brakefield & Mazzotta, 1995). Indeed, the use of constant

temperatures might significantly underestimate the impact of global warming on

life history evolution (Fischer *et al.* 2011). These concerns clearly highlight the

need to integrate more realistic thermal regimes into empirical studies (Niehaus *et*

al., 2012; Paaijmans *et al.*, 2013) in order to fully understand how species are

responding to climate change (specifically increases in mean temperature and

increased variability) under more natural conditions (Folguera *et al.*, 2011; Colinet

et al., 2015).

To this end, we investigated the long-term effects of thermal stress

caused by realistically fluctuating temperatures on the phenotypic plasticity and

adaptation of two laboratory populations of seed beetle, *Callosobruchus*

maculatus. Two different thermal regimes were used: in one regime, beetles were

kept at a constant control temperature of 29°C, and in the second regime we

exposed beetles to a more stressful and realistic fluctuating daily cycle with a

mean temperature of 33°C. This temperature regime was chosen because it is

representative of a May day in southern India, where the laboratory lines were

originally collected. The two beetle populations were then either kept in these two

environments for the entirety of the experiment or were *switched* to the other

environment for two generations without selection prior to assaying several key

life history traits (namely, development time, body mass and various measures of

fitness).

Previous work in the same species of beetle has highlighted the negative

effect of elevated temperatures on various life history traits (Rogell *et al.*, 2014;

Vasudeva *et al.*, 2014; Ivimey-Cook *et al.*, 2021). In addition, a series of papers in

by Hallsson and Björklund (2012) investigated how increasing temperature with

and without environmental fluctuation influenced reproductive success,

development time, and body size. Whilst predominantly focusing on the amount

of treatment-specific genetic and environmental variance for each trait, they found

that fluctuating environments had a significantly negative effect on fecundity but

little to no effect on hatching success, development time and body mass.

Importantly, our definition of fluctuating environments differs from theirs. Whilst

the term employed here represents daily sinusoidal fluctuations around a

constant mean, representative of the natural environment that a seed beetle

could experience, Hallsson and Björklund (2012) used the term to describe the

addition of thermal noise around a linear increase from 30°C to 36°C. As such,

we may expect entirely different life-history responses to higher mean

temperatures that are also fluctuating around that mean.

In particular, the heterogeneity of the environment an organism

experiences, along with the associated costs and benefits of maintaining differing

widths of thermal tolerance, can dramatically alter which genotypic strategies are

favoured by selection. For instance, stabile and homogenous environments have

been shown to lead to the evolution of specialists (Reboud & Bell, 1997) who

reduce the width of their thermal niche in order to maximise performance. The

particular direction of this specialization is largely dependent on the optimum

temperature of the organism, be that hot: "Hotter is Better" hypothesis (Kingsolver

& Huey, 2008, Fig. 1), or cold: "Colder is Better" hypothesis (Huey *et al.*, 1999,

Fig. 1). Fluctuating environments may instead favour the evolution of generalists

with increased phenotypic plasticity (or reversible phenotypic plasticity,

Ratikainen & Kokko, 2019) and enhanced genetic variation (Reboud & Bell, 1997;

see Table 1 and 2 from Kassen, 2002). However, this plasticity might carry a cost,

for example if individuals trade off increased thermal tolerance with reduced

reproductive performance (e.g. the "Jack of all Trades, Master of None"

hypothesis - Huey & Kingsolver, 1989; Berger *et al.*, 2014; Fig. 1). Therefore, we

would predict that populations evolving in fluctuating environments should

underperform (i.e., through reduced fitness), in comparison to those in more

constant benign environments. Alternatively, If phenotypic plasticity is instead

beneficial and can evolve without a cost to reproduction, then beetles evolving in

fluctuating environments should have higher performance (i.e. evolve to become

"Jack of all Trades, Master of All" - Huey & Hertz, 1984 *or* "Superior Thermal

Generalists", see Berger *et al.*, 2014; Fig. 1) than those evolving in constant

benign environments. As a result, the relative comparison between benign and

stressful environments will largely depend on the costs and benefits of phenotypic

plasticity (see Fig. 1). Lastly, individuals from a benign environment that are then

switched to a stressful environment should suffer reduced reproductive

performance owing to a lack of evolved phenotypic plasticity.

The predictions for development time and body size were more

straightforward. Increases in temperature have been shown to reduce

development time in many insects (Hagstrum & Leach, 1973; Nylin & Gotthard,

1998; Hallsson & Björklund, 2012b; a; Czarnoleski *et al.*, 2013). Since the mean

temperature of the fluctuating environment was higher in our experiment, we

predicted that individuals that evolved in, or were later placed into, the fluctuating

environment would experience shorter development times. Finally, increasing

mean temperature whilst within fluctuating environments has been shown to

decrease body size in insects (Nylin & Gotthard, 1998; Kingsolver & Huey, 2008;

Hallsson & Björklund, 2012a; b; Czarnoleski *et al.*, 2013). Many organisms,

particularly insects, experience faster growth at higher temperatures; under these

conditions, decreasing cellular size and thus body size would allow for an

optimum rate of resource exchange (Czarnoleski *et al.*, 2013). Therefore, we

predicted that body size should be smaller in the populations that evolved in, or

were placed into, the fluctuating environment.

To investigate potential population-level differences we used two different

strains of *C. maculatus*, derived from USA and Leicester. Although we had no *a*

priori predictions about if and how any population-level differences would

manifest, previous research has highlighted the substantial genotypic and

phenotypic differences that exist between strains of *C. maculatus* (Rankin &

Arnqvist, 2008). Lastly, we also investigated relative differences between the two

sexes, as males and females in this species can vary in their response to

stressful environments (Stillwell & Fox, 2007; Rogell *et al.*, 2014). Previous

research on this species has shown that sexual dimorphism in traits such as body

size at emergence decreases when beetles are gradually exposed to novel,

higher temperatures over 18 generations (Hallsson & Björklund, 2012b; 2012a).

We therefore predicted that any differences between sexes would similarly

decrease in populations exposed to more stressful, fluctuating temperatures.

Figure 1. Hypotheses regarding reproductive performance and phenotypic plasticity

and the evolution of generalist or specialist strategies.

Methods

Study system

C. maculatus is an agricultural pest that originates from Africa and Asia but is

currently found throughout the world. Adult seed beetles are facultatively aphagous –

that is, they do not require food or water to survive and acquire all the resources they

need during the larval stage within the bean. Aphagous adults can live up to two

weeks but adults with access to nutrients can live three weeks or more (Fox, 1993;

Ursprung *et al.*, 2009). Females lay their eggs on the surface of dried legumes, such

as mung beans (*Vigna radiata*) or black-eyed beans (*Vigna unguiculata*). Larvae from

the fertilized eggs burrow into the bean and eclose as adults 23-27 days later.

Females of this species start to mate and lay eggs as soon as they emerge (Beck &

Blumer, 2014).

Laboratory evolution lines and climatic conditions

In our experiment, we used two different strains of *C. maculatus,* South India (SI)

USA and SI Leicester, both derived from a population of beetles collected in

Tirunelveli, India, in 1979 (Mitchell, 1991) and then subsequently raised at the

University of Kentucky, USA, and the University of Leicester, UK (from 1992),

respectively (Fricke & Arnqvist, 2004). We obtained these strains from Uppsala

University in Sweden and maintained them at the American University of Paris for

approximately 25 generations before establishing the replicates for the current

experiment. Beetles were cultured exclusively on mung beans and kept in climate

chambers at a constant 29°C, 50% relative humidity, and a 12:12h light:dark cycle.

They were housed at aphagy in 1L jars with 250g of beans, and approximately 250-

350 newly hatched beetles were transferred to new jars with fresh beans every 24

days on a continual basis.

For 19 generations prior to the experiment, we subdivided SI USA and SI

Leicester populations (Fig. 2A) into four replicates each. Two "Control" replicates of

each were maintained at a constant temperature of 29°C, while two "Stress"

replicates were subject to daily temperature cycle (Fig. 2B). The daily cycle consists

of 12 separate 2hr periods of constant temperature T_i,

226
$$T_i = T_{mean} + \Delta\mathrm{T}\sin\left(\frac{i-12}{12}\pi\right),$$

where T_{mean} = 33 °C, ΔT = 7°C, and i = 0,1…. 11. This is a stepwise sinusoidal

temperature cycle with T_{max}=40 and T_{min}=26 and which mimics typical late spring

condition in Southern India, where this species evolved. Humidity and light cycles

remained the same for all replicates: 50% relative humidity, and a 12:12 h light:dark

cycle.

Prior to the assays of body size, development time, and reproductive fitness (see

next section), we created two replicates of two populations (SI USA and SI Leicester;

see Fig. 2A/B) for:

1.   beetles adapted to and assayed at Control conditions (**USA**-C and **LEI**-C);

2.   beetles adapted (for one year) to and assayed in Stress conditions (**USA**-S

237   and **LEI**-S);

238   3.   beetles adapted to Control conditions that were then acclimatized to Stress

239   conditions for two generations without selection prior to being assayed (**USA**-

240   C-to-S and **LEI**-C-to-S); and

241   4.   beetles adapted to Stress conditions that were acclimatized to Control

242   conditions for two generations without selection prior to being assayed (**USA**-

243   S-to-C, **LEI**-S-to-C).

To separate genetic adaptation from phenotypic plasticity – that is, to control

for the influence of parental effects – we created "acclimatized" treatments by

allowing beetles from one set of conditions to adjust to the new set of conditions for

two generations prior to assays (C-to-S and S-to-C; Fig. 2B). Critically, to remove the

possible effects of natural selection and rule out the possibility that rapid genetic

responses might influence our measurements for the "acclimatized" treatments, we

randomly paired 50 pairs of virgin beetles during each of the two generations. That is,

we removed the effects of assortative mating.

For the Stress-to-Control (USA S-to-C, LEI S-to-C) and the Control-to-Stress

(USA C-to-S, LEI C-to-S) treatments, the following steps were performed. First, we

moved fertilized beans from jars into 48-well virgin chambers. When enough males

and females had hatched (from day 21-24 after hatch), we paired 60 males and 60

females in 60 60-mm petri dishes with 100 beans in each dish and placed these

dishes in the opposing chamber. We made sure that there were sufficient number of

beans in each dish for females to lay just one egg per bean, thereby eliminating

competition among multiple larvae within a bean. To further reduce larval

competition, we removed adult beetle pairs from the petri dishes after 72 hours.

After 19 days, we then choose the 50 petri dishes with the most fertilized

beans and put 48 fertilized beans from each petri dish into individual marked 48-well

virgin chambers. One day after eclosion, we randomly selected and paired 50

females and 50 males and put them into 50mm petri dishes with 80 beans. After 72

hours, the adult males and females were removed from the petri dishes. After 19

days, fertilized beans from each petri dish were transferred into 48-well virgin

chambers. We then marked the date of hatch and sex of all offspring and conducted

assays of development time, body mass, and reproductive fitness.

For the continuous C and S treatments, we skipped the acclimatization steps

and simply moved fertilized beans to virgin chambers prior to hatch, selecting one-

day old males and females for all subsequent assays.

A

SI USA — SI LEI

Control (**USA**-C)	Replicate 1 Replicate 2	Replicate 1 Replicate 2	Control (**LEI**-C)
Stress (**USA**-S)	Replicate 1 Replicate 2	Replicate 1 Replicate 2	Stress (**LEI**-S)
Control-to-Stress (**USA**-C-to-S)	Replicate 1 Replicate 2	Replicate 1 Replicate 2	Control-to-Stress (**LEI**-C-to-S)
Stress-to-Control (**USA**-S-to-C)	Replicate 1 Replicate 2	Replicate 1 Replicate 2	Stress-to-Control (**LEI**-S-to-C)

B

Control (C)

Constant 29°C

Stress (S)

Fluctuating around 33°C

Control-to-Stress (C-to-S)

Constant 29°C **- to -** Fluctuating around 33°C

Stress-to-Control (S-to-C)

Fluctuating around 33°C **- to -** Constant 29°C

Figure 2. (A) A schematic diagram of the four temperature treatments (Control,

Stress, Control-to-Stress, and Stress-to-Control) across both the USA (SI USA) and

Leicester (SI LEI) populations. (B) A representation of the four different temperature

treatments used. The black bar in the bottom two treatments, denotes a switch from

one treatment to another in order for acclimatization to occur over two generations

prior to assaying.

Body mass assay

For each of the 16 populations, we weighed male and female beetles separately in

two batches of 12 beetles each using an Ohaus Pioneer Plus Analytical Balance

(Model PA214C). All virgin chambers were weighed empty, prior to adding the

beetles to be weighed. We then calculated the difference between the weight of the

virgin chamber before the beetles were placed inside and the weight of the virgin

chamber after the beetles were placed inside. Beetles from the body mass assay

were not used for the other assays.

Development time and fitness assays

Development time and fitness assays were done concurrently. We paired 50 1-day

old male and female beetles and placed them into 60mm petri dishes filled with 85

beans. After 24 hours, each pair was moved together into a new 60mm petri dish

filled with 75 beans, and the first dish was set aside in the same climate chamber.

The next day, this procedure was repeated. The beetles were left in the third and

final set of dishes until they died. The first dishes for each pair were marked "Day 0",

the second dishes were marked "Day 1", and the third and final dishes were marked

"Day 2+". After 19 days for each dish, all the fertilized beans were placed into virgin

chambers, which were monitored daily for the date of eclosion and the sex of all

offspring. To calculate development time, we counted the number of days between

the date that an egg was laid and the date the adult offspring eclosed. To calculate

LRS, we calculated the number of male and female offspring that emerged for each

pair, from all three sets of dishes (Days 0, 1, and 2+). To calculate individual fitness

(or λind), we took the dominant eigenvalue (using the popbio v. 2.7 package,

(Stubben & Milligan, 2007) from an age-structured Leslie matrix (Leslie, 1945) where

the top row denotes fertility and the sub-diagonal represents survival from age t to

$t+1$. 24 days of development time were added prior to day 0-2+ fertility schedule.

Statistical analyses

All statistical analyses were performed in R v. 4.1.0 (R Core Team, 2016) using

glmmTMB v. 1.0.2.9000 (Brooks *et al.*, 2017; Magnusson *et al.*, 2019) with pairwise

interactions and slopes compared using emmeans v. 1.6.1 (Lenth *et al.*, 2019).

Ggplot2 v. 3.3.4 (Wickham, 2009) was used for all graphical visualization. The overall

effects of each variable were identified using the Type 3 Anova function from car v.

3.0-10 package (Fox *et al.*, 2012).

For both development time, body mass, and individual fitness, a Gaussian

linear mixed effects model was fitted (after log-transforming development time to

reduce right-skew) with the fixed effects of Population, Treatment, Sex (for

development time and body mass), and all higher order interactions. In addition, the

fixed factor of replicate number and, for development time, the random effect of

parental ID was added to account for potential pseudoreplication. If the three-way

interaction was non-significant (identified from the Anova function), it was removed in

order to reduce model complexity. A generalized linear mixed effects model with

Poisson distribution and similar fixed effect structure was fitted for LRS and age-

specific reproduction, albeit without the variable Sex and all subsequent two/three-

way interactions. In addition to a covariate of Day (and all higher-order interactions

with Population and Treatment), a random effect of individual ID was added to the

age-specific reproduction model to account for repeated measures. Both models

were checked for overdispersion and zero-inflation (ZI) using DHARMa v. 0.4.1,

(Hartig, 2020) by simulating the residuals of a Poisson model. If significant, a variety

of error distributions with/without zero-inflated parameters (if ZI was identified) were

then fitted and the best model identified by Akaike's Information Criterion (AIC).

Results

Body mass

Individual body mass was significantly affected by both of the interactions involving

Sex (Sex and Population: $\chi^2 = 25.95$, p<0.001; Sex and Treatment: $\chi^2 = 19.18$,

p<0.001; Table S1A) but not by the remaining two-way interaction of Population and

Treatment ($\chi^2 = 2.51$, p = 0.47) or the three-way interaction ($\chi^2 = 6.37$, p = 0.09).

Females were generally heavier than males (Fig. 3) in both populations. The

difference was more pronounced in the USA population (Average body mass (g): F-

LEI = 0.087, M-LEI = 0.075, estimate = 0.013, p<0.001; F-USA = 0.089, M-USA =

0.068, estimate = 0.021, p<0.001; Fig. 3, Tables S1A-B). The females of both

populations were similar in weight, but the USA males were smaller (F-LEI = 0.087,

F-USA = 0.089, estimate = -0.002, *p* = 0.35; M-LEI = 0.075, M-USA = 0.068,

estimate = 0.007, p<0.001; Fig. 3, Tables S1A-B). The relative difference between

male and female weight was largely dependent on temperature treatment (Fig. 3).

Overall, beetles raised in the S-to-C regime were smaller for both sexes in

comparison to other treatments (all comparisons significant to p = 0.05; Tables S1A-

B). However, the difference between male and female weights was far less

pronounced in the constant stress (S) treatment (albeit all differences were still

significant to p<0.001; F-S-to-C = 0.083, F-S = 0.089, F-C-to-S = 0.091, F-C = 0.089;

M-S-to-C = 0.064, M-S = 0.079, M-C-to-S = 0.071, M-C = 0.071; Tables S1A-B).

Figure 3. Average body mass of individuals from the Leicester (Red) or USA (Blue)

populations in the C (Control), C-to-S (Control-to-Stress), S (Stress), or S-to-C

(Stress-to-Control) temperature treatments. Sex is denoted with either circle

(Female) or triangle (Male) points. Points represent mean values with accompanying

standard errors. Half violins show the distribution of raw data.

Development time

Development time was influenced by both two-way interactions involving

Temperature treatment (Treatment and Population: $\chi^2 (3) = 105.94$, $p <0.001$;

365 Treatment and Sex: χ^2 (3) = 11.76, p = 0.008). Similar to body mass, no detectable

366 three-way interaction was found (χ^2 (3) = 1.71, p = 0.635). The USA and Leicester

367 populations responded differently to the four temperature treatments, with beetles

368 from the Leicester population developing faster on average. Populations from USA

369 took significantly longer to develop in comparison to populations from Leicester in

370 both the Control and S-to-C Treatments (Average Development Time (Days): LEI-C

371 = 23.7, USA-C = 24.7, Ratio = 0.960, p<0.001; LEI-S-to-C = 23.1, USA-S-to-C =

372 24.1, Ratio = 0.961, p<0.001; Fig. 4) but not in C-to-S (LEI-C-to-S = 23.0, USA-C-to-

373 S = 23.0, Ratio = 0.999, p = 1.00) or S Treatments (LEI-S = 22.5, USA-S = 22.3,

374 Ratio = 1.01, p = 0.38; Fig. 4, Tables S2A-B). Despite these differences, both

375 populations in the S treatment developed faster compared to other temperature

376 treatments (LEI-S = 22.5, LEI-S-to-C = 23.1, LEI-C-to-S = 23.0, LEI-C = 23.7; USA-S

377 = 22.3, USA-S-to-C = 24.1, USA-C-to-S = 23.0, USA-C = 24.7). In addition, whilst

378 males typically developed faster in both populations (Fig. 4), only in the S-to-C

379 treatment was this difference significant (F-S-to-C = 23.7, M-S-to-C = 23.50, Ratio =

380 1.008, p<0.001; Fig. 4, Tables S2A-B). However, both males and females in the S

381 treatment developed faster than other temperature treatments and showed no sex

382 difference in development time (F-S = 22.4, F-S-to-C = 23.7, F-C-to-S = 23.0, F-C =

383 24.2; M-S = 22.4, M-S-to-C = 23.5, M-C-to-S = 22.9, M-C = 24.1).

Figure 4. Average development time in days of individuals from the Leicester (Red)

or USA (Blue) populations in the C (Control), C-to-S (Control-to-Stress), S (Stress),

or S-to-C (Stress-to-Control) temperature treatments. Sex is denoted with either

circle (Female) or triangle (Male) points. Points represent mean values with

accompanying standard errors. Half violins with associated jittered points show the

distribution of raw data.

Lifetime reproductive success

LRS was significantly influenced by the interaction between Population and

Treatment (χ^2(1) = 12.61, p = 0.005). Across all temperature treatments, LRS was

lower in the USA population than in the Leicester population (Average LRS: LEI =

53.2, USA = 40.2, Ratio = 1.32, *p*<0.001; Fig. 5, Tables S3A-B). The largest

differences between populations occurred in the C-to-S and C treatments (LEI-C-to-S

= 50.2, USA-C-to-S = 33.6, Ratio = 1.49, *p*<0.001; LEI-C = 52.6, USA-C = 37.2, Ratio

= 1.41, *p*<0.001; Fig. 5, Tables S3A-B). Despite these differences, LRS was highest

in the S-to-C treatment for both populations (LEI-C = 52.6, LEI-S-to-C = 69.7, Ratio =

0.75, *p*<0.001; USA-C = 37.2, USA-S-to-C = 60.9, Ratio = 0.61, *p*<0.001; Fig. 5,

Tables S3A-B).

Figure 5. Average lifetime reproductive success (LRS) of individuals from the

Leicester (Red) or USA (Blue) populations in the C (Control), C-to-S (Control-to-

Stress), S (Stress), or S-to-C (Stress-to-Control) temperature treatments. Points

represent mean values with accompanying standard errors. Half violins show the

distribution of raw data.

Age-specific reproduction

The interaction between population, treatment and day significantly influenced age-

specific reproduction (χ^2(3) = 203.54, *p*<0.001, Fig. 6). Whilst continuous C and S

treatments exhibited similar age-specific patterns across both populations (Fig. 6),

there were striking differences between S-to-C and C-to-S treatments. This was

particularly apparent on Days 1 and 3 of reproduction (Fig. 6). Individuals from both

LEI-S-to-C and LEI-C-to-S treatments exhibited far greater Day 1 reproduction than

all other treatments from both populations (including the analogous S-to-C and C-to-

S treatments). In stark contrast, USA-S-to-C exhibited increased reproduction *after*

Day 1, with the result that Day 3 of reproduction was comparable with the Day 1

reproduction for LEI S-to-C (Fig. 6). This variation in reproductive schedule resulted

in large differences in age-specific slopes between the USA S-to-C and LEI S-to-C

treatments (Average change in average offspring produced per day: LEI S-to-C = -

7.14, USA S-to-C = 8.78, estimate = -15.92, p<0.001; Fig. 6, Tables S4A-B).

Figure 6. Age-specific reproduction (LRS) of individuals from the Leicester (dash-

dot) or USA (solid) populations in the C (Control, Red), C-to-S (Control-to-Stress,

Green), S (Stress, Blue), or S-to-C (Stress-to-Control, Purple) temperature

treatments. Points with error bars represent mean values with accompanying

standard errors. Jittered points represent raw data.

Individual fitness

Whilst the individual main effects of Population and Temperature treatment

significantly influenced λind (χ^2 (1) = 81.02, *p* <0.001, χ^2 (3) = 111.23, *p* <0.001,

respectively; Fig. 7), the interaction between these two did not ($\chi^2 (3)$ = 7.64, *p* =

0.054). In a similar manner to LRS, the USA population had reduced λind over all

temperature treatments (Average λind: LEI = 1.164, USA = 1.146, estimate = 0.018,

p<0.001; Fig. 7, Tables S5A-B). Across both populations, individuals from the S-to-C

treatment had the highest λind (C = 1.154, C-to-S = 1.148, S = 1.146, S-to-C =

1.173, all comparisons p<0.001; Tables S5A-B). This was also consistent when

looking within each population (all comparisons p<0.001).

Figure 7. Fitness (Lambda) of individuals from the Leicester (Red) or USA (Blue)

populations in the C (Control), C-to-S (Control-to-Stress), S (Stress), or S-to-C

(Stress-to-Control) temperature treatments. Points represent mean values with

accompanying standard errors. Half violins show the distribution of raw data.

Figure 8. Hypotheses and subsequent results (indicated with ticks and crosses)

regarding reproductive performance and phenotypic plasticity and the evolution of

generalist or specialist strategies.

Discussion

Previous hypotheses (Lewontin, 1974; Nevo, 1978) have suggested that evolution in

fluctuating environments should select for genetically variable individuals that are

robust to a wide range of thermal conditions, often termed generalists (Kassen,

2002). In this experiment, generalists exposed to novel benign conditions (S-to-C,

Fig. 8) (although not those exposed to continuously stressful environments, S, see

below) had higher reproductive fitness (higher LRS and λind) than specialists that

were exposed to novel stressful conditions (C-to-S) or kept continuously in a benign

environment (C). These results are broadly in line with previous research conducted

in the opportunistic bacterial pathogen, *Serratia marcescens* (Ketola *et al.*, 2013)*.*

Heterogenous environments selected for bacterial clones that had faster growth and

higher yield than those grown under constant environments, even when introduced

into the same control temperature of 31°C (Ketola *et al.,* 2013). This increased

growth may be due to far greater selective pressure to remove mutations that could

otherwise accumulate in more constant benign environments (Ketola *et al.*, 2013).

However, results from a digital evolution experiment have suggested that this may

only allow for optimization in short-term rather than long-term evolvability (Canino-

Koning *et al.*, 2019).

We found that individuals within stressful fluctuating environments evolved

strategies similar to thermal-tolerance generalists as well as performance specialists

(Gilchrist, 1995). More specifically, individuals were able to tolerate and survive at

wide thermal breadth, presumably owing to the evolution of phenotypic plasticity, but

they reached maximal performance and experienced increased LRS only when

conditions were optimal (Gilchrist, 1995; Ketola *et al.*, 2013). This could in part be

due to differential gene expression as a result of exposure to a novel thermal

environment (also known as cryptic genetic variability, McGuigan & Sgro, 2009;

Paaby & Rockman, 2014). Similarly, this release of cryptic genetic variation could

explain why individuals that were predicted to perform poorly when switching from a

benign control to stressful fluctuating environment (C-to-S), suffered no fitness cost.

However, the relative merits of this form of variation have been queried and require

additional empirical testing (McGuigan & Sgro, 2009; Edwards & Yang, 2021;

Johansson *et al.*, 2021). Regardless, these results suggest that the evolution of

phenotypic plasticity in experiment did not carry substantial fitness costs.

When comparing body mass across treatments, individuals from the S-to-C

group were significantly smaller (across both populations and sexes), despite having

similar development times to those of the constant control group. Smaller body size

could therefore be a cost of adaptation (Kassen, 2002; Ketola *et al.*, 2013) to the

fluctuating conditions that is only detected in the benign environment and is

manifested in reduced body size (albeit with no negative effects on fitness). However,

whether this reduction in mass is a disadvantage is difficult to determine. Large size

provides a fitness advantage due to sexual selection (Andersson & Iwasa, 1996),

whereas small size decreases larval competition (Messina, 2004). Interestingly,

similar to Hallsson & Björklund (2012a), we found that sexual dimorphism in adult

body size decreased in the continuously stressful environment (S), although this

could be considered population-specific as individuals from USA failed to show such

a decrease.

Lastly, there were consistent population-level differences across all

environmental treatments. In general, beetles from the USA population produced

fewer offspring, took longer to develop (although this difference disappeared under

stressful conditions), and had greater sexual dimorphism in body size compared to

individuals from the Leicester population. Such genotypic variation between

populations of the same species (particularly in *Callosobruchus maculatus*) has been

found to impact larval competition (Messina, 1991), various morphometric measures

(Rankin & Arnqvist, 2008), egg-to-adult survival and offspring production (Rankin &

Arnqvist, 2008), and adult longevity and mortality (Fox *et al.*, 2004a; b). However, in

this experiment, one particular observation stood out. Whilst a switch from a stressful

environment to a benign regime (S-to-C) was associated with an increase in LRS and

λind in both USA and Leicester populations, there was a noticeable difference in the

age-specific reproductive schedules between the USA and Leicester populations.

Leicester beetles produced more offspring on the first day and fewer on days two and

three, whilst in the USA population the opposite was observed. Why such a

difference in reproductive schedule exists deserves deeper investigation. If genetic

variance differed between populations, this may explain why Leicester individuals

were able to respond more quickly, with greater Day 1 reproduction. This could also

in part explain why development time was similar across all experimental

temperature regimes for the Leicester population. Together this clearly highlights the

benefits of comparing and contrasting results from multiple strains, as the response

of species to environmental change can often be highly population-specific.

Successfully mimicking natural conditions within the laboratory is key to

understanding how species will react to climate change. As temperatures increase

and become more variable, the incorporation of more realistic conditions into

laboratory studies such as these will allow us not only to predict more accurately how

wild species will respond to climate change, specifically thermal stress, but also to

produce accurate species- and population-specific thermal reaction norms

(Paaijmans *et al.*, 2013). This study highlights the need to include both constant and

more variable conditions in laboratory studies and suggests that the evolution of

generalist and specialist strategies will be vital in explaining species-specific

responses to climate change.

Acknowledgements

Funding from the American University of Paris supported this study. Many thanks to

Sophie Bricout for helping to maintain the beetle populations and to Sophie Bricout,

Linda Martz, Shannon Monahan, Sarah Sidi, and Xaviera Steele for their assistance

with the experimental assays. Thanks also to Ram Vasudeva, Kynan Delaney, and

Joel Pick for helpful discussions.

References

Andersson, M. & Iwasa, Y. 1996. Sexual selection. *Trends Ecol. Evol.* **11**: 53–58.
Elsevier.

Bauerfeind, S.S. & Fischer, K. 2014. Simulating climate change: temperature
extremes but not means diminish performance in a widespread butterfly.
Popul. Ecol. **56**: 239–250.

Beck, C.W. & Blumer, L.S. 2014. A handbook on bean beetles, Callosobruchus
maculatus. *Natl. Sci. Found. URL Httpwww Beanbeetles Orghandbook Pdf*
Last Accessed 16 June 2015.

Berger, D., St\aangberg, J., Baur, J. & Walters, R.J. 2021. Elevated temperature
increases genome-wide selection on de novo mutations. *Proc. R. Soc. B* **288**:
20203094. The Royal Society.

Berger, D., Stångberg, J., Grieshop, K., Martinossi-Allibert, I. & Arnqvist, G. 2017.
Temperature effects on life-history trade-offs, germline maintenance and
mutation rate under simulated climate warming. *Proc. R. Soc. B Biol. Sci.* **284**:
20171721. Royal Society.

Berger, D., Walters, R.J. & Blanckenhorn, W.U. 2014. Experimental evolution for
generalists and specialists reveals multivariate genetic constraints on thermal
reaction norms. *J. Evol. Biol.* **27**: 1975–1989.

Brakefield, P.M. & Mazzotta, V. 1995. Matching field and laboratory environments:
effects of neglecting daily temperature variation on insect reaction norms. *J.*
Evol. Biol. **8**: 559–573.

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W.,
Nielsen, A., *et al.* 2017. glmmTMB balances speed and flexibility among
packages for zero-inflated generalized linear mixed modeling. *R J.*, doi:
10.32614/rj-2017-066.

Canino-Koning, R., Wiser, M.J. & Ofria, C. 2019. Fluctuating environments select for
short-term phenotypic variation leading to long-term exploration. *PLOS*
Comput. Biol. **15**: e1006445. Public Library of Science.

Chen, H., Zheng, X., Luo, M., Guo, J., Solangi, G.S., Wan, F., *et al.* 2018. Effect of
short-term high-temperature exposure on the life history parameters of
Ophraella communa.

Colinet, H., Sinclair, B.J., Vernon, P. & Renault, D. 2015. Insects in Fluctuating
Thermal Environments. *Annu. Rev. Entomol.* **60**: 123–140.

Czarnoleski, M., Cooper, B.S., Kierat, J. & Angilletta, M.J., Jr. 2013. Flies developed
small bodies and small cells in warm and in thermally fluctuating
environments. *J. Exp. Biol.* **216**: 2896–2901.

Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak,
D.C., *et al.* 2008. Impacts of climate warming on terrestrial ectotherms across
latitude. *Proc. Natl. Acad. Sci.* **105**: 6668–6672.

Edwards, C.B. & Yang, L.H. 2021. Evolved Phenological Cueing Strategies Show
Variable Responses to Climate Change. *Am. Nat.* **197**: E1–E16. The
University of Chicago Press.

Fischer, K., Kölzow, N., Höltje, H. & Karl, I. 2011. Assay conditions in laboratory
experiments: is the use of constant rather than fluctuating temperatures
justified when investigating temperature-induced plasticity? *Oecologia* **166**:
23–33.

Folguera, G., Bastías, D.A., Caers, J., Rojas, J.M., Piulachs, M.-D., Bellés, X., *et al.*
2011. An experimental test of the role of environmental temperature

variability on ectotherm molecular, physiological and life-history traits: implications for global warming. *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* **159**: 242–246.

Fox, C.W. 1993. Multiple mating, lifetime fecundity and female mortality of the bruchid beetle, Callosobruchus maculatus (Coleoptera: Bruchidae). *Funct. Ecol.* 203–208. JSTOR.

Fox, C.W., Bush, M.L., Roff, D.A. & Wallin, W.G. 2004a. Evolutionary genetics of lifespan and mortality rates in two populations of the seed beetle, Callosobruchus maculatus. *Heredity* **92**: 170–181.

Fox, C.W., Czesak, M.E. & Wallin, W.G. 2004b. Complex genetic architecture of population differences in adult lifespan of a beetle: Nonadditive inheritance, gender differences, body size and a large maternal effect. *J. Evol. Biol.* **17**: 1007–1017.

Fox, J., Weisberg, S., Adler, D., Bates, D., Baud-Bovy, G., Ellison, S., *et al.* 2012. Package 'car.' *Vienna R Found. Stat. Comput.*

Fricke, C. & Arnqvist, G. 2004. Divergence in replicated phylogenies: the evolution of partial post-mating prezygotic isolation in bean weevils. *J. Evol. Biol.* **17**: 1345–1354. Wiley Online Library.

Gilchrist, G.W. 1995. Specialists and Generalists in Changing Environments. I. Fitness Landscapes of Thermal Sensitivity. *Am. Nat.* **146**: 252–270.

Hagstrum, D.W. & Leach, C.E. 1973. Role of Constant and Fluctuating Temperatures in Determining Development Time and Fecundity of Three Species of Stored-Products Coleoptera2. *Ann. Entomol. Soc. Am.* **66**: 407–410.

Hallsson, L.R. & Björklund, M. 2012a. Selection in a fluctuating environment and the evolution of sexual dimorphism in the seed beetle Callosobruchus maculatus. *J. Evol. Biol.* **25**: 1564–1575.

Hallsson, L.R. & Björklund, M. 2012b. Selection in a fluctuating environment leads to decreased genetic variation and facilitates the evolution of phenotypic plasticity: Evolutionary response in a fluctuating environment. *J. Evol. Biol.* **25**: 1275–1290.

Hansen, P.J. 2009. Effects of heat stress on mammalian reproduction. *Philos. Trans. R. Soc. B Biol. Sci.* **364**: 3341–3350.

Hartig, F. 2020. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. R package.

Hatfield, J.L. & Prueger, J.H. 2015. Temperature extremes: Effect on plant growth and development. *Weather Clim. Extrem.* **10**: 4–10.

Hedhly, A., Hormaza, J.I. & Herrero, M. 2009. Global warming and sexual plant reproduction. *Trends Plant Sci.* **14**: 30–36. Elsevier.

Hetem, R.S., Fuller, A., Maloney, S.K. & Mitchell, D. 2014. Responses of large
mammals to climate change. *Temperature* **1**: 115–127.

Holt, R.D. 1990. The microevolutionary consequences of climate change. *Trends*
Ecol. Evol. **5**: 311–315. Elsevier.

Huey, R.B., Berrigan, D., Gilchrist, G.W. & Herron, J.C. 1999. Testing the Adaptive
Significance of Acclimation: A Strong Inference Approach. *Am. Zool.* **39**:
323–336.

Huey, R.B. & Hertz, P.E. 1984. Is a Jack-of-All-Temperatures a Master of None?
Evolution **38**: 441–444.

Huey, R.B. & Kingsolver, J.G. 1989. Evolution of thermal sensitivity of ectotherm
performance. *Trends Ecol. Evol.* **4**: 131–135.

Ivimey-Cook, E., Bricout, S., Candela, V., Maklakov, A.A. & Berg, E.C. 2021.
Inbreeding reduces fitness of seed beetles under thermal stress. *J. Evol. Biol.*
jeb.13899.

Janzen, D.H. 1967. Why Mountain Passes are Higher in the Tropics. *Am. Nat.* **101**:
233–249. [University of Chicago Press, American Society of Naturalists].

Johansson, F., Watts, P.C., Sniegula, S. & Berger, D. 2021. Natural selection
mediated by seasonal time constraints increases the alignment between
evolvability and developmental plasticity. *Evolution* **75**: 464–475.

Kassen, R. 2002. The experimental evolution of specialists, generalists, and the
maintenance of diversity. *J. Evol. Biol.* **15**: 173–190.

Ketola, T., Mikonranta, L., Zhang, J., Saarinen, K., Örmälä, A.-M., Friman, V.-P., *et*
al. 2013. FLUCTUATING TEMPERATURE LEADS TO EVOLUTION OF
THERMAL GENERALISM AND PREADAPTATION TO NOVEL
ENVIRONMENTS. *Evolution* **67**: 2936–2944. [Society for the Study of
Evolution, Wiley].

Kingsolver, J.G. & Huey, R.B. 2008. Size, temperature, and fitness: three rules. 18.

Lashof, D.A. & Ahuja, D.R. 1990. Relative contributions of greenhouse gas emissions
to global warming. *Nature* **344**: 529–531. Springer.

Lenth, R., Singmann, H., Love, J., Buerkner, P. & Herve, M. 2019. *emmeans:*
Estimated Marginal Means, aka Least-Squares Means (Version 1.3. 4).

Leslie, P.H. 1945. On the Use of Matrices in Certain Population Mathematics.
Biometrika **33**: 183–212.

Lewontin, R.C. 1974. *The genetic basis of evolutionary change.* Columbia University
Press New York.

Lopez, A.D. & Adair, T. 2019. Is the long-term decline in cardiovascular-disease
mortality in high-income countries over? Evidence from national vital
statistics net.

Magnusson, A., Skaug, H.J., Nielsen, A., Berg, C., Kristensen, K., Maechler, M., *et*
al. 2019. Package 'glmmTMB. *R Package Version 01*.

Masson-Delmotte, V., Zhai, P. & Pirani, A. 2021. *IPCC, 2021: Climate Change*
2021: The Physical Science Basis. Contribution of Working Group I to the
Sixth Assessment Report of the Intergovernmental Panel on Climate Change.

McGuigan, K. & Sgro, C.M. 2009. Evolutionary consequences of cryptic genetic
variation. *Trends Ecol. Evol.* **24**: 305–311. Elsevier.

Messina, F.J. 1991. Competitive interactions between larvae from divergent strains of
the cowpea weevil (Coleoptera: Bruchidae). *Environ. Entomol.* **20**: 1438–
1443. Oxford University Press Oxford, UK.

Messina, F.J. 2004. Predictable Modification of Body Size and Competitive Ability
Following a Host Shift by a Seed Beetle. *Evolution* **58**: 2788–2797. [Society
for the Study of Evolution, Wiley].

Mitchell, R. 1991. The traits of a biotype of Callosobruchus maculatus
(F.)(Coleoptera: Bruchidae) from South India. *J. Stored Prod. Res.* **27**: 221–
224. Elsevier.

Nevo, E. 1978. Genetic variation in natural populations: Patterns and theory. *Theor.*
Popul. Biol. **13**: 121–177.

Niehaus, A.C., Angilletta, M.J., Sears, M.W., Franklin, C.E. & Wilson, R.S. 2012.
Predicting the physiological performance of ectotherms in fluctuating thermal
environments. *J. Exp. Biol.* **215**: 694–701.

Nylin, S. & Gotthard, K. 1998. Plasticity in Life-History Traits. *Annu. Rev. Entomol.*
43: 63–83.

Paaby, A.B. & Rockman, M.V. 2014. Cryptic genetic variation: evolution's hidden
substrate. *Nat. Rev. Genet.* **15**: 247–258.

Paaijmans, K.P., Heinig, R.L., Seliga, R.A., Blanford, J.I., Blanford, S., Murdock,
C.C., *et al.* 2013. Temperature variation makes ectotherms more sensitive to
climate change. *Glob. Change Biol.* **19**: 2373–2380.

Pachauri, R.K., Allen, M.R., Barros, V.R., Broome, J., Cramer, W., Christ, R., *et al.*
2014. *Climate change 2014: synthesis report. Contribution of Working*
Groups I, II and III to the fifth assessment report of the Intergovernmental
Panel on Climate Change. Ipcc.

Parmesan, C. 2006. Ecological and Evolutionary Responses to Recent Climate
Change. *Annu. Rev. Ecol. Evol. Syst.* **37**: 637–669.

Pereira, H.M., Navarro, L.M. & Martins, I.S. 2012. Global Biodiversity Change: The
Bad, the Good, and the Unknown. *Annu. Rev. Environ. Resour.* **37**: 25–50.

R Core Team. 2016. R: A language and environment for statistical computing. *R*
Found. Stat. Comput. Vienna Austria **0**: {ISBN} 3-900051-07-0.

Rankin, D.J. & Arnqvist, G. 2008. Sexual Dimorphism Is Associated with Population
Fitness in the Seed Beetle Callosobruchus Maculatus. *Evolution* **62**: 622–630.

Ratikainen, I.I. & Kokko, H. 2019. The coevolution of lifespan and reversible
plasticity. *Nat. Commun.* **10**: 538.

Reboud, X. & Bell, G. 1997. Experimental evolution in Chiamydomonas. Ill.
Evolution of specialist and generalist types in environments that vary in space
and time. *Heredity* **78**: 507—514.

Rogell, B., Widegren, W., Hallsson, L.R., Berger, D., Björklund, M. & Maklakov,
A.A. 2014. Sex-dependent evolution of life-history traits following adaptation
to climate warming. *Funct. Ecol.* **28**: 469–478.

Stillwell, R.C. & Fox, C.W. 2007. Environmental effects on sexual size dimorphism
of a seed-feeding beetle. *Oecologia* **153**: 273–280.

Stubben, C. & Milligan, B. 2007. Estimating and analyzing demographic models
using the popbio package in R. *J. Stat. Softw.* **22**: 1–23.

Thompson, R.M., Beardall, J., Beringer, J., Grace, M. & Sardina, P. 2013. Means and
extremes: building variability into community-level climate change
experiments. *Ecol. Lett.* **16**: 799–806.

Trisos, C.H., Merow, C. & Pigot, A.L. 2020. The projected timing of abrupt
ecological disruption from climate change. *Nature* **580**: 496–501. Nature
Publishing Group.

Ursprung, C., Den Hollander, M. & Gwynne, D.T. 2009. Female seed beetles,
Callosobruchus maculatus, remate for male-supplied water rather than
ejaculate nutrition. *Behav. Ecol. Sociobiol.* **63**: 781–788. Springer.

Vasseur, D.A., DeLong, J.P., Gilbert, B., Greig, H.S., Harley, C.D.G., McCann, K.S.,
et al. 2014. Increased temperature variation poses a greater risk to species than
climate warming. *Proc. R. Soc. B Biol. Sci.* **281**: 20132612.

Vasudeva, R., Deeming, D.C. & Eady, P.E. 2014. Developmental temperature affects
the expression of ejaculatory traits and the outcome of sperm competition in
Callosobruchus maculatus. *J. Evol. Biol.* **27**: 1811–1818.

Visser, M.E., Noordwijk, A.J. van, Tinbergen, J.M. & Lessells, C.M. 1998. Warmer
springs lead to mistimed reproduction in great tits (Parus major). *Proc. R. Soc.*
Lond. B Biol. Sci. **265**: 1867–1870. Royal Society.

Visser, M.E., te Marvelde, L. & Lof, M.E. 2012. Adaptive phenological mismatches
of birds and their food in a warming world. *J. Ornithol.* **153**: 75–84.

Walters, R.J., Blanckenhorn, W.U. & Berger, D. 2012. Forecasting extinction risk of
735   ectotherms under climate warming: an evolutionary perspective. *Funct. Ecol.*
736   **26**: 1324–1338.

737   Wickham, H. 2009. *ggplot2 Elegant Graphics for Data Analysis*.

738