Evidence for alternate stable states in an Ecuadorian Andean Cloud Forest

Authors: Ana Mariscal Chávez[1], Daniel Churchill Thomas[2], Austin Haffenden[3], Rocío Manobanda[4], Laurence Duvauchelle[5], William Defas[6], Miguel Angel Chinchero[7], Danilo Simba[8], Edison Jaramillo[9], Bitty A. Roy[10], Mika Peck[11]

Institutional affiliations & emails

[1]Ana Mariscal Chávez, National Herbarium of Ecuador and

Cambugan Foundation, Quito Ecuador

[2]Daniel Church Thomas, BayCEER Institute, Bayreuth, Germany

[3]Austin Haffenden, University of Wales, Bangor

[4]Rocío Manobanda, National Herbarium of Ecuador.

[5]Laurence Duvauchelle, Amazonia Simbiosis

[6]William Defas, National Herbarium of Ecuador.

[7]Miguel Angel Chinchero, National Herbarium of Ecuador.

[8]José Danilo Simba Larco, National Herbarium of Ecuador.

[9]Edison Jaramillo, National Herbarium of Ecuador.

[10]Bitty A. Roy, Institute of Ecology and Evolution, University of Oregon, Eugene, OR, USA.

[11]Mika Peck, Ecology, Behaviour and Environment, School of Life Sciences, University of Sussex, Brighton, 27 UK.

Author Contributions:

 Conceptualization, A.M.C., D.C.T., B.A.R., and M.P.; methodology, A.M.C., D.C.T., and M.P.; formal analysis, D.C.T.; investigation, A.M.C, R.M, A.H., L.D., W.D, M.A.C, D.S, E.M.; resources, M.P.; data curation, A.M.C, D.C.T.; writing—original draft preparation, D.C.T.; writing—review and editing, A.M.C., D.C.T., B.A.R.and M.P.; visualization, D.C.T.; supervision, M.P.; project administration, M.P.; funding acquisition, M.P. All authors have read and agreed to the final version of the manuscript.

Abstract

We analyzed a set of historical data from rapid vegetation inventories in a tropical montane cloud forest in northern Andean Ecuador. Trees in plots from several types of forest were counted and measured, including: (1) primary forest, including mature and recently closed-canopy sites and naturally formed gaps, (2) abandoned pasture, and (3) intensively-farmed sites. The goal of the study was to understand in a specific period of time the similarities and differences among natural and anthropogenic disturbances and their potential long term effect on the forest plant community. We found that mature and intermediate close canopy sites are similar. Also old forest is quite resilient to gap-forming disturbance. Natural gaps are quickly colonized by old-forest-associated plant species, and return to an old-forest-type community of trees in a short time. In contrast, forests regenerating from anthropogenic disturbance appear to have multiple possible states: some regenerating forest sites where the anthropogenic disturbance were low are coming to closely resemble old-forest-type communities, but some where the anthropogenic disturbance was intense appear to be changing in a very different direction, which does not resemble any other vegetation community type currently in the forest. A major predictor of present ecological state is the type of land use before reforestation: pastures can occasionally transition back to the pre-disturbance state of forest. More intensively used sites, many of which are abandoned sugar cane plantations, do not return to a pre-disturbance ecological state, instead forming a new and different kind of forest, dominated by a different community of trees. We examined tree-seedling communities to understand the trajectory of the forest into the future, and find that new forest types may be forming that do not resemble any existing associationsintensive agricultural sites. We also found that Los Cedros is extremely diverse in tree species. We estimate approximately 500 species of tree in only the small southeastern area of the reserve that has been explored scientifically. Additionally, the forest tree community shows extremely rapid distance decay (beta-diversity), approaching near complete turn-over in the limited spatial extent of the study. This suggests that hundreds of other tree species remain to be observed in the reserve, in addition to the 350+ that are directly observed in the present study, including new observations of species with IUCN threatened-endangered status. We also highlight the conservation value of Reserva Los Cedros, which has managed to reverse deforestation within its boundaries despite a general trend of extensive deforestation in the surrounding region, and to protect large, contiguous areas of highly-endangered Andean primary cloud forest habitat.

Introduction

Succession has been a central topic of discussion in ecology for at least a century (Bray & Curtis, 1957; Bunin, 2021; Chambers et al., 2013; Chang & Turner, 2019; Clements, 1916; Connell & Slatyer, 1977; Cowles, 1899; Curtis & Greene, 1949; Gleason, 1926; Holling, 1973; Whittaker, 1956). As with many emergent properties of ecosystems, succession in ecosystems is endlessly controversial: the very existence of stable equilibria states and predictable successional (seral) stages have long been both called-into-question and defended (Bunin, 2021; Gleason, 1926; Norden et al., 2015; Richards, 1952). In recent times increasing attention has been given to models that lie within a community assembly and/or coexistence framework, to explain and predict community composition in plants (HilleRisLambers et al., 2012; T. P. Young et al., 2001). Rather than attempting to broadly predict changes in dominant species associations in the form of seral stages, community-assembly and coexistence models give greater attention to the importance of individuals, which sum to explain differences in community compositions. These models emphasize dispersal, priority, and stochastic effects on individuals, and in the case of niche-based or trait-based models, a further discussion of selection or "filtering" by biotic and abiotic factors (Fisher & Mehta, 2014; Hubbell, 2008; Keddy, 1992; Kraft et al., 2008; Laughlin & Laughlin, 2013; Ulrich et al., 2016). There is no inherent contradiction among these two broad approaches to community modeling: successional and assembly/co-existence approaches are highly complementary (Bhaskar et al., 2014; T. P. Young et al., 2001), and are sometimes packaged into a general "dynamics" framework (Pickett et al., 2009). This cultural shift in community ecology is perhaps due to a desire by ecologists to move away from the highly idiosyncratic, localized knowledge of a site that is often required to successfully predict successional patterns (Norden et al., 2009, 2015), and instead to more abstract and more universal ecological models.

This shift may also perhaps be partly due to the chaotic times in which we find ourselves. In the current era, most ecosystem types are experiencing historical rates of species loss (Barnosky et al., 2011; Pimm et al., 2014), and are already undergoing consequences of climate change (Allen et al., 2010; Carmody et al., 2016; Ciais et al., 2005; Day et al., 2008; Doney et al., 2012; Fadrique et al., 2018; Mantgem et al., 2009). Ecosystems are also continually receiving new species from human activity (Hobbs et al., 2009) and face direct modification or even wholesale elimination due to land use change (Ellis et al., 2010; Hautier et al., 2015; Vitousek et al., 1997). Few anthropogenically disturbed ecosystems seem to have been successfully "returned" to stable historical ecological states (Hobbs et al., 2009), let alone to climax states that resemble some sort of prehistoric ecological stability. In our destabilized era, the search for ancient species associations that require multiple uninterrupted, decadal journeys through intermediate ecological stages can sometimes seem quixotic.

However, the discussion of succession in forests has taken on particular and new urgency in the current era of accelerating climate change. Forests and their soils fix a significant portion of carbon every year, which has probably been essential in slowing negative effects of anthropogenic atmospheric carbon release (Pan et al., 2011). Forests and their soils are also sitting reservoirs of vast amounts of sequestered carbon in soils that are readily lost through disturbance (Lal, 2005). After many decades of underappreciation as essentially carbon-neutral ecosystems (Odum, 2014), a vibrant discussion has arisen around the role of primary forests not only as immense storehouses of carbon, but also as continuing carbon sinks (Gundersen et al., 2021; Luyssaert et al., 2008). New, proforestation-centered prescriptions for climate mitigation have therefore gained support: old forests (primary or late-stage successional) should be be protected as reserves of existing fixed carbon, and potentially high-functioning

sinks for further carbon fixation, and management of secondary forests should enhance characteristics that maximize carbon sequestration (Lewis et al., 2019; Moomaw et al., 2019; Sacco et al., 2021; Watson et al., 2018). Such characteristics are often exemplified by old forests (Harmon et al., 1990; Stephenson et al., 2014). Tropical forests are the most volatile portion of the global forest carbon sink, and their role as a sink for atmospheric CO_2, rather than a source, will depend greatly on how they are managed and protected in the coming years (Mitchard, 2018). Tropical montane forests are likely more significant carbon stores than was historically thought, given higher-than-predicted productivity and their large surface area, resulting from complex, high-angle topography (Girardin et al., 2014; Spracklen & Righelato, 2014). Thus predictive models of succession may now become essential tools in mitigating the climate crisis: an understanding of seral stages (if indeed they exist!) in forests will be necessary for reaching maximal carbon storage and protecting existing carbon reserves.

The current study was undertaken at Reserva Los Cedros, a mid-elevation cloud forest reserve located in the heart of the Tropical Andean biodiversity hotspot (cite our 2019 paper). While the tropical Andes may not display the same levels of alpha tree diversity often seen in the forests of the lower Amazon basin (Draper et al., 2019; Valencia et al., 1994), the Tropical Andes have long been recognized for a high biodiversity across multiple taxonomic groups (Brehm et al., 2016; Churchill, 2009; Gentry & Dodson, 1987; Hughes, 2016; Küper et al., 2004; Olson, 1994; Ramírez-Barahona et al., 2011, p.; Ríos-Touma et al., 2017; Smith et al., 2007). This biodiversity is characterized by endemism at very fine spatial scales (Borchsenius, 1997; K. R. Young et al., 2002), making it an extremely important conservation priority (Myers et al., 2000). The mechanisms for the high biodiversity and endemism in the tropics-at-large, including montane forests, have been debated for decades and perhaps centuries(Janzen, 1967; Von Humboldt & Bonpland, 2010). High solar energy and constant, plentiful rainfall are characteristic in many areas of the neotropics, and these traits are often associated with high biodiversity (Hawkins et al., 2003). Additionally, the neotropics that may have higher speciation rates, acting as evolutionary "cradles" and/or have higher retention of taxa, thus acting as refugia or "museums", allowing for greater accumulation of species over geological time (Arita & Vázquez-Domínguez, 2008; Jablonski et al., 2006). The tropical Andes augment these general biodiversity trends in the tropics with characteristics that promote even further the role of endemism: complex topography creates diversified habitats and environmental gradients and increased niches(Gentry, 1992; Hughes, 2016; Janzen, 1967), increasing the potential for sympatric speciation, and also adding dispersal barriers, increasing the potential for fine-scale allopatric speciation (Särkinen et al., 2012). The insular or "island" nature of cloud/mountain systems may be particularly enhanced situations for speciation events that result from neutral events, especially founder effects (Gentry, 1992). All of these factors presumably interact and sum to create the unique patterns of diversity and endemism observed in the Andes.

Here we examine the dynamics of early (<20 years) forest succession at Reserva Los Cedros, following anthropogenic or natural-gap-forming-disturbances. The landscape of the reserve is dominated by primary forest, but contains mixed-age secondary forests of varying land use histories (see results) . In regions where remnants of ancient forest ecosystems - ecosystems without a history of significant modern anthropogenic disturbance - persist on the landscape, there are advantages for both ecological analysis and restoration efforts.. First, we have information in the form of local, functioning examples of the biological complexity and reference points for primary-forest equilibrium. Second, forest recovery is facilitated by the presence of species that have co-evolved in local conditions for many millennia, available to act as nuclei for reforestation efforts (Corbin & Holl, 2012; Zahawi et al., 2013). The existence of extensive primary forest has proven particularly valuable for the study of succession and community

equilibria in neotropical forests (Bhaskar et al., 2014; Loughlin et al., 2018; Norden et al., 2009).

We also examine the spatial signature of this fine scale beta diversity of trees, with the working hypothesis that individual, steep-sided catchments may be the spatial unit of importance for understanding Andean biodiversity. Our hypothesis was that two characteristics/processes govern the behavior of community similarity in the tropical Andes: (1) large-scale spatial auto-correlation will cause short-distance comparisons to be more similar than farther comparisons, and with with distance, all site comparisons will approach complete dissimilarity, a phenomenon known as Tobler's Law (Tobler, 1970). However, (2) we predict significant "noise" around this general trend of decreasing similarity, because complex topography of the Andes causes great dissimilarity even among some very localized comparisons, and conversely, causes highly similar habitats to occur sometimes at great distances apart due to similar site conditions. We predict that in systems with extremely complex topography such as the Andes, the most informative unit for modeling community dissimilarity will not be euclidean distance (meters), but instead the number of watersheds crossed. If this hypothesis is supported, it will facilitate future understanding of tropical diversity.

Finally, Los Cedros has succeeded at both forest protection and reforestation, despite its location in a region of high deforestation and now mineral exploration. Thus we examine here the success of los Cedros in conservation of forest in a region of Ecuador that is under intense extractivist pressure, in terms of forest cover change and IUCN red-list species observed.

Methods

Site – Reserva los Cedros

All fieldwork described was performed at or directly adjacent to Reserva Los Cedros, a protected forest reserve in the western slope of the Andes, in northwestern Ecuador (00°18 0 31.0 00 N, 78°46 0 44.6 00 W), at 1000-2700 m asl. The reserve lies within the Andean Choco bioregion, one of the most biodiverse habitats on the planet (Gentry, 1992). It is also considered to be within the Tropical Andean Biodiversity hotspot (Myers et al., 2000). The reserve protects 5256 hectares of cloud forest. Definitions of cloud forest vary an can be quite complex (Stadtmüller, 1987), but following Foster (2001) and Stadtmüller (1987), we use the term cloud forest to mean a forest "whose characteristics are tied to the frequent presence of clouds and mist," and consider the terms montane rain forest and cloud forest to be synonyms. Following Grubb et al. (1963; Grubb & Whitmore, 1966), the ecotones within the reserve are probably most accurately classified as cloud forest of mostly lower montane rain forest, with some regions of higher montane cloud forest, and some Elfin forest zones in its highest, least-explored areas. The Reserve also shares a border with the 305,000 hectare nationally-protected Cotocachi-Cayapas Ecological Reserve. Rainfall averages 2903 (+/- 186 mm) per year according to onsite measurements. Humidity is typically high (~100%), and daily temperatures at the site range from 15°C to 25°C (Policha, 2014). Annual fluctuations in temperatures are minimal. Daily precipitation varies according to the time of year, with the wet season (October to May) to dry (June-September) seasons (___Ecuador weather service, Nanegalito and Garcia Moreno weather stations, accessed by WeatherAtlas__).

The rarity of primary forest in the north Andean region is due to deforestation from pre-colonial times (Loughlin et al., 2018; Wunder, 2000), and more recently in the 1960's, due to land

reform efforts that legalized and encouraged homestead-scale settlement of large government and private ("Hacienda") forest holdings (Dodson & Gentry, 1991). Los Cedros has apparently largely escaped deforestation during both eras. The land within Los Cedros and the surrounding region was inhabited by the poorly understood Yumbo indigenous group until 1690.. Their activities probably significantly altered sites but with unknown effects on the ecology and canopy cover of the forest, but their economy was likely integrated into the forested setting of the mid-elevation Andean region (R. D. Lippi, 1988; R. Lippi & Gudiño, 2010). However, the impacts of indigenous land use in South American forests has been until recently been greatly underestimated and mis-understood by scholars (Loughlin et al., 2018; Watling et al., 2017). Given the rainfall and high humidity it is unlikely that large-scale deforestation resulted from fire at Los Cedros. Other than possible small-scale pre-colonial indigenous activities, the majority of the Los Cedros forest has seen relatively few anthropogenic alterations.

Land acquisitions to build the reserve were made between 1988-1995.. Tracts purchased by Los Cedros were usually developed or deforested only in small proportions of their total area, usually for small-scale mixed ("finca") agriculture or for cattle and mule pasture. In some cases, these small-scale clearings were made for assertion of legal rights over land rather than agricultural production at scale (Jose DeCoux, pers. comm.). Following their acquisition by Los Cedros, all former agricultural and pasture sites were selectively grazed for several months by cattle, to reduce competition from graminoids for protected tree seedlings. Tree seedlings were not planted, but were allowed to reestablish by encroachment from adjacent forest and other sources during and after selective grazing. This method is known informally by some workers in the region as "reforestation by cattle rotation", and is intended to release naturally regenerated tree seedlings from intensive competition from pasture grasses (Goosem et al., 2013) without the use of herbicides.

Satellite data of forest cover from 1990 shows ~96% forest cover of los Cedros (see results), of which at least 80% is thought to have been primary forest (Jose DeCoux, pers. comm.). Non-forest land use was concentrated in the southeastern portion of the reserve, where the vegetation surveys were undertaken.

Tree survey and plant identification

Selection of sites and categorization of land use history

Primary forest was defined as those forests which have no historical record and no physical evidence of canopy removal or other large structural alteration by humans. However, forest ecosystems exist continuously as matrices of various states of non-anthropogenic gap formation and closure (Chambers et al., 2013; Espírito-Santo et al., 2014; Franklin et al., 2002; M. O. Hunter et al., 2015; Nicholas Brokaw, 1982), including montane wet forests and cloud forests (Crausbay & Martin, 2016). As such, primary forest is given three categories of land cover based on their gap characteristics:

Bosque cerrado ("BC") = Mature forest, with no physical sign of gap-forming disturbance. Canopy is closed.

Bosque secundario ("BS") = Sites with evidence of recent gap, now with a closed canopy,

Claros del bosque ("CLB") = recent, natural gaps in the forest.

In addition to primary forest, sites with histories of anthropogenic disturbance were placed into two categories:

Regeneración de fincas agricultura y ganadería ("RG") = abandoned small family farms, with land use mostly consisting of pasture maintained for cattle.

Regeneración cañaveral ("RCA") = Intensively farmed sites, used for sugar cane production.

Survey methods

Site selection - surveys were undertaken at Reserva Los Cedros from the years of 2005 and 2007. The southern area of Los Cedros was divided into 4 areas of study, which were then further divided into three sub-blocks each, from which one sub-block was randomly selected and searched for natural gaps. Once located, each natural gap was also accompanied by a BS and BC site, at a minimum of 40 m distance between survey sites. Two additional smaller blocks were added in the vicinity of previously settled areas, to increase coverage of anthropogenic disturbances. Overall 61 sites from various land use histories and elevations were sampled (see fig. 6).

Adult trees were defined as trees at the height 1.5m with a diameter of 10 cm or greater. Material from all trees fitting this description were sampled, within a circle plot of 30 m^2 radius from the centerpoint for morphological identification to species, where possible. Additionally, within each survey site of BC, RCA, and RG plots, all juvenile trees were examined. Juvenile trees were defined as trees with a maximum diameter less than 10 cm, growing from 50 cm up to 2 m, of species known to be capable of growing to adult tree size as given above, under ideal circumstances and with sufficient time. Juvenile trees were surveyed within a square subplot of size 5m x 5m, centered within the larger 30m circular plot. Nomenclature of identifications is based on the Flora of Ecuador (Persson et al., n.d.).

Statistical and informatic methods

All analyses were conducted in Python and R. Python version 3.8.10 (Python Core Team, 2018), using Pandas 1.1.3 (McKinney, 2010; Reback et al., 2020) and Matplotlib 3.1.2 (J. D. Hunter, 2007) for visualization. R version 3.6.3 (R Core Team, 2020) with in-box R plotter engine was used. All analysis was in an Ubuntu 20.04.2 LTS environment. Where Bayesian analyses were used, Python version 3.7.3 was used for compatibility with the PyMC3 package, version 3.8 (Salvatier et al., 2016). All Bayesian models used the default NUTS sampler to sample the posterior. Code for all pertinent statistical analyses have been run and recorded in a Jupyter Notebook that is stored in the affiliated github repo, and is viewable as a notebook online here.

Species accumulation curves and richness estimators.

Species accumulation curves for the entire area studied were calculated as the number of adult tree species observed per meter-squared of physically sampled area. Each subplot covered a circle with a diameter of 30m, or 0.071 ha. Additionally, a permanent tree diversity plot was established concurrently with rapid surveys in 2005. Within this, trees were sampled in a grid

format, every 10 m along a east/west axis and every 5 m along a north/south axis, in a rectangular half-hectare area (50m x 100m). Trees were identified to species where possible, and unidentified trees were grouped into species-level operational taxonomic designations (supp. table 1A). In the case of the permanent plot, tree species accumulation was modeled as a function of trees-examined. Species accumulation models and species estimators were calculated using the specaccum, specpool and associated functions with the Vegan package in R, version 2.5-6 (Jari et al., 2019). Point predictions of diversity for 1 ha area were generated using the predict function, which when used as a method of specpool model objects uses a Mao Tau rarefaction method (Colwell et al., 2012) to generate predictions.

Tree community turnover (distance decay or beta-diversity)

Turnover in tree communities was modeled and visualized using Bray-Curtis dissimilarity (Bray & Curtis, 1957; Legendre & Legendre, 2012) as a function of (1) physical distance or (2) watershed crossings mapped at small-scale. Bray-Curtis dissimilarities and physical distances matrices among subplots were calculated using the SciPy spatial submodule (Virtanen et al., 2020). Bayesian models of community dissimilarity decay were conducted using PyMC3 package in Python, with visualizations created using the companion ArViz package (Kumar et al., 2019).

Turnover by physical distance

Tree community turnover by physical distance was calculated using Bray-Curtis dissimilarity as a function of physical distance (meters). Two Bayesian models were used as alternative formulations of the above two processes of local variability and large-scale spatial autocorrelation.

1. **Asymptotic model**: In one approach, BC dissimilarity was modeled as an asymptotic function, or "Michaelis-Menten"-type function:

$$y = \frac{x}{K_m + x}$$

(eq. 1)

Where y is predicted Bray-Curtis dissimilarity, x is distance between compared sites, and K_m is the distance at which half of maximum BC dissimilarity is reached. This model honors both theoretical constraints of complete similarity at proximal sites and maximum dissimilarity. Additionally, we modeled variation around the mean as a linear variable, reducing as all distant comparisons approached complete dissimilarity (BC=1). This shrinking variance term ϵ was exponentiated to avoid negative values. As a Bayesian model this is formulated in probabilistic terms, with priors, as follows:

$$y \sim N\left(\frac{x}{K_m + x}, \varepsilon\right), \quad \varepsilon = e^{(\delta x + \gamma)}$$

(eq. 2)

(eq. 3) $K_m \sim N(\mu = 200, \sigma = 10)$

(eq. 4) $\delta \sim N(\mu = -0.0006, \sigma = 0.005)$

(eq. 5) $\gamma \sim N(\mu = 1.5, \sigma = 0.5)$

Prior distribution for K_m was loosely estimated from Draper et al. (2019), who showed that in many Amazonian forest ecosystem types, decay of tree community similarity occurs rapidly, losing more than half their similarity within 200 m. Owing to the novel formulation, priors for delta and gamma of equation 2 were not found in existing literature, and were assigned weak priors, with means intended to reflect the initial wide variation and its subsequent tightening around BC=1.

2. **Linear model**. In a second formulation, decay in tree community similarity was modeled as a simple linear equation. Variance around the mean BC dissimilarity was still allowed to vary negatively with distance, beginning with large initial spread to encompass both highly similar and dissimilar sites at close distances. Additionally, a skewed mean distribution (Ashour & Abdel-hameed, 2010; O'Hagan & Leonard, 1976) was used to describe the variance around the mean function, given the long skew that is observable in the data at most distances. As with the asymptotic model, the variance around the mean function of BC dissimilarity was described as an exponentiated linear function.

(eq. 6) $y \sim \text{skewNormal}\left(\alpha + \beta x, \; \varepsilon, \; a\right), \quad \varepsilon = e^{(\delta x + \gamma)}$

(eq. 7) $\alpha \sim N(\mu = 0.9, \sigma = 0.2)$

(eq. 8) $\beta \sim N(\mu = 0.0001, \sigma = 0.00001)$

(eq. 9) $a \sim N(\mu = -2.0, \sigma = 0.5)$

(eq. 10) $\delta \sim N(\mu = -0.0006, \sigma = 0.005)$

(eq. 11) $\gamma \sim N(\mu = -1.5, \sigma = 0.5)$

Prior distribution for β was again loosely estimated from Draper et al. (2019), using data from community turnover of Amazonian Terra firme forests. All other priors were assigned as weakly informative priors.

Model comparison - Comparison of performance between these two models were done using posterior predictive checks on the data and variance explained (Bayesian R^2), using the sample_posterior_predictive command in the PyMC3 package, and the r2_score command in the ArViz package. Bayesian p-values to indicate balanced performance were calculated using the distribution of residual differences between model predictions and observed data. To confirm Bayesian linear model results, a classic linear least-squares regression was also applied to the data, using a Wald test with a null hypothesis that the slope of community turnover was zero, as implemented with the linregress function in the SciPy Stats module. Variance defaults (bivariate normal) for the function were not modified.

Overall turnover by watershed crossings

To better understand changes in community as a function of topographic complexity, we delineated small watersheds in the area of the study area.The original digital elevation model used was the ASTER Global Digital Elevation Map version 2 (Tachikawa et al., 2011). Every subplot was assigned to a micro-watershed, and a distance matrix of watershed crossings. Watersheds were delineated using the pysheds package in Python (Bartos, 2021). Distance matrix of micro-watershed crossings was calculated using Djikstra's algorithm for shortest

paths (Dijkstra, 1959) in the NetworkX package, version 2.4 (Hagberg et al., 2008). Methods for specific watershed delineations are further explained in the statistical jupyter notebook (https://nbviewer.jupyter.org/github/danchurch/losCedrosTrees/blob/master/anaData/MariscalDataExploration.ipynb#watersheds).

Ordination by historical land-use/habitat

Structuring in adult tree communities as a function of historical land-use/habitat by was visualized using Bray-Curtis dissimilarity and non-metric multidimensional scaling (NMDS) (Legendre & Legendre, 2012), using the metaMDS function in the Vegan package in R.

GIS data and additional environmental data

Except where otherwise noted, geospatial data were explored and visualized using using tools from the GeoPandas (version 0.8.1, https://geopandas.org) and Rasterio (version 1.1.8, https://rasterio.readthedocs.io/), and QGIS (QGIS v3.4.11-Madeira, OSGEO v3.7.1). Several environmental variables were generated from ASTER Global Digital Elevation Map version 2, including slope, aspect, elevation, eastern and northern exposure, and distance-to-nearest-stream of all subplots. Stream data were digitized from Map CT-NII-C3-d of Bosque Protector Los Cedros, from the Ecuadorian Instituto Geográfica Militar. Rasters of slope and aspect were generated with the Raster Terrain Analysis tool suite in QGIS.

Hierarchical clustering of sites

To understand the current types of forests present at los Cedros, adult tree community data from all sites were partitioned into clusters using Ward's minimum variance clustering (Legendre & Legendre, 2012) using Bray-Curtis dissimilarity, as implemented in the hclust command in the Vegan package in R. Number of clusters was decided by eye from the resulting dendrogram, taking into consideration branch lengths and meaningful ecological groups.

Prediction of current ecological state by land use history and elevation

Current ecological state as defined from cluster analysis was modeled as a function of historical land-use/habitat data and elevation. Each of the two predictors (height, land-use/habitat) was considered individually, and then a combined model was created using both elevation and historical land-use/habitat. All models were Bayesian, written in python with the PyMC3 package.

For land-use/habitat, a multinomial logistic regression model was constructed with a softmax link function and the following priors:

(eq. 12) $\alpha \sim N(\mu=0, \sigma=5)$

(eq. 13) $\beta \sim N(\mu=0, \sigma=5)$

(eq. 14) $\theta = softmax(\alpha + \beta x)$

(eq. 15) $y \sim categorical(\theta \mid y_{observed})$

Where y is a vector of predicted probabilities for forest type (cluster) of a site, distributed as a categorical random variable of α, β, and x, conditioned on our observed current forest type (i.e., cluster number). x is observed historical land-use/habitat, in a dummy variable matrix format. α are y-intercepts for each current forest type, and β is the vector of slope coefficients for each of 5 types of historical land-use/habitat, for each current forest type. α and β were assigned weak, normally distributed priors centered on zero.

Elevation. A logistic regression model was constructed to model differences among forest types III and IV from the hierarchical cluster analysis. Hierarchical clustering of tree communities and NMS ordinations both indicated two distinct "natural" forest types occurring within the extent of the survey, which were designated Forest types III and IV (see results). To further understand differences between these two forest types, exploratory PERMANOVA models were used to test grouping by all available environmental predictors (see Jupyter notebook). Following this, forest type was modeled as a function of elevation, using a Bayesian logistic regression model. Priors and posterior were modeled as follows:

(eq. 16) $\alpha \sim N(\mu=0, \sigma=10)$

(eq. 17) $\beta \sim N(\mu=0, \sigma=10)$

(eq. 18) $\theta = \text{logistic}(\alpha + \beta x)$

(eq. 19) $y \sim \text{Bernoulli}(\theta \mid y_{observed})$

Where α and β are the intercept and slope controlling the boundary decision between type III and type IV forests, in terms of elevation. θ is the prior probability that a forest site will be type III forest, given its elevation, and y is a Bernoulli distribution with θ as the mean, conditioned by the observed forest type.

For the combined elevation and historical land-use/habitat model – Elevation and historical land-use/habitat predictors were combined in one linear model:

(eq. 20) $\alpha \sim N(\mu=0, \sigma=5)$

(eq. 21) $\beta \sim N(\mu=0, \sigma=5)$

(eq. 22) $\Theta = \text{softmax}(\alpha + \beta x)$

(eq. 23) $y \sim \text{categorical}(\theta \mid y_{observed})$

Where x is observed historical land-use/habitat, in a dummy variable matrix format, with an additional column vector giving the elevation of each site. α are y-intercepts are intercepts for each row of x, and β is the vector of slope coefficients for each column of x for each current forest type (= row of x). α and β were assigned weak, normally distributed priors centered on zero. y is a vector of predicted probabilities for forest type (cluster) of a site, distributed as a categorical random variable of α, β, and x, conditioned on our observed current forest type (i.e., cluster number).

Indicator species analysis

Indicator species analyses were conducted to ascertain representative species for both historical land-use/habitat types and current forest type (cluster analysis results), using the indicspecies package in R (De Cáceres et al., 2010), using the multipatt function and Pearson's phi coefficient correlation (function "r.g").

Spatial analyses

To determine important spatial scales on which tree community was changing, and to give shape to possible spatial structuring of tree community, Moran's eigenvector maps (MEM) were constructed (Dray et al., 2012), following examples as found in documentation for the ADEspatial package (https://cran.r-project.org/web/packages/adespatial/vignettes/tutorial.html). A spatially-weighted neighborhood matrix of sample sites was constructed using a Gabriel connectivity matrix with an inverse distance weighting. Abundances of our adult tree community matrix were Hellinger transformed (Legendre & Legendre, 2012), and new PCA axes of the transformed were determined with the dudi.pca command in ade4 package. Important MEMs were then selected on the basis of their p-value as explanatory terms in a linear model with the 4 most important tree community PCA axes as response variables. Important MEMS were determined using a forward model selection process, as implemented with the mem.select command in the adespatial package. The default cutoff of alpha = 0.05 was used to decide statistical significance of MEMs. All available environmental variables were checked for correlations with MEMs using standard linear regression as implemented in the linregress function in the Python SciPy Stats package. Correction for multiple testing was done with a Benjamini-Hochberg correction, as implemented in the multipletests command in the Python Statsmodels package. For exploratory purposes, the false discovery rate was set at FDR=0.1.

Juvenile communities

Juvenile tree community data were available for a subset of historical land-use/habitat types: "Bosque Cerrado" (CB), "Regeneración cañaveral" (RCA), and "Regeneración fincas agricultura y ganadería" (RG). Juvenile data were not taken for "Claro del Bosque'' (CLB) or "Bosque secundario ("BS"). Due to the difficulties associated with identifying poorly understood herbaceous taxa and/or juvenile trees, these data required extensive additional cleaning and verification, following botanical identifications. For the purposes of this study, juvenile trees were defined as undersized plants that would, in time and with proper growth conditions, reach the stature of the mature trees as defined above, namely: woody plants with a diameter-at-1.5m of 10 cm or greater. This is as compared to herbaceous plants or small woody plants (such as lianas or subshrubs) which would never or rarely reach 10 cm girth at 1.5m height.

Samples that did not receive sufficient identification to confidently conclude that the sample was indeed a juvenile tree were discarded from the analysis. This selection was done by automated and manual queries of growth form data requested from the TRY database (Kattge et al., 2020), manual checks of the Encyclopedia of life (Parr et al., 2014), and the Smithsonian Tropical Research collections site (https://stricollections.org/portal/index.php). The Gentry

manual (Gentry, 1993) was also frequently consulted. Exact species information was not often available and designation of juvenile tree status was made using genus growth form data where possible. Juvenile tree communities were compared to adult trees by subsetting adult tree community data to just those sites with juvenile data, and combining community matrices from both. These were then ordinated using Bray-Curtis dissimilarity and non-metric multidimensional scaling (NMS), using the metaMDS function in the Vegan package in R.

Deforestation in the region

To understand the extractive pressure in the region of Los Cedros, and the efficacy of Los Cedros as a conservation project, we examined land cover changes from 1990 to 2018. Data for changing land cover were accessed from the Ecuadorian Ministerio del Ambiente's ("MAE") interactive map and GIS catalogue, using the "Cobertura vegetal" layers from 1990 and 2018, available here. Vector data from the MAE were rasterized to a pixel resolution of 30 m^2 using the gdal_rasterize program from the GDAL tool suite from the OSGeo project (GDAL/OGR contributors, 2021), and visualized in Matplotlib using the rasterio package. Background deforestation rates were constrained to Cotacachi canton of the Imbabura province in which Los Cedros is located. Deforestation rates in the similarly-sized, nearby Bosque Protector el Chontal were quantified for comparison with los Cedros. BP Chontal is managed by the Cattlemen's Association of the nearby town of Brillasol (pers comm. Jose DeCoux). Administrative boundaries of BP Los Cedros and BP Chontal were supplied using official boundary layer "Bosque y Vegetación protectora", also available at MAE's interactive map website.

Results

Particularly interesting or pertinent posterior distributions are described here. Full results on posterior distributions of all models are available in the jupyter notebook.

Species accumulation curves and richness estimators.

343 adult tree species were directly observed in plots. Estimates of total adult tree species in the study area range from 404 to 566 species (fig. 1A, table 1). In the permanent 0.5 ha plot, 43 species of tree were observed, distributed among 36 genera and 25 plant families, with a range of predicted total species from 51 to 72 species in the half-hectare (fig. 1B, table 1). On average, 1 ha in the southern portion of Los Cedros is predicted to host 169 species of adult tree. Chao estimates, first and second order Jackknife, and Bootstrap estimates for the total study area and for the permanent 0.5 ha plot are given in table 1.

When examined by land-use/habitat type, more adult tree species were observed from reforested pasture sites, possibly due to deeper sampling, but natural-disturbance sites (BS, BC, and CLB) were comparable when rarified to a common sample size (fig. 1C, supp. table 2). Sites with history of intensive agricultural use were the least diverse, though depth of sampling was lowest in this group. Species estimators by land-use history are given in supp. table 2.

Tree community turnover (distance decay)

Turnover by physical distance

1. Asymptotic model

The asymptote model is visualized with 50% and 95% highest posterior densities ("HPD") for predicted BC values in fig. 2A. The posterior distribution of our asymptotic model explained a large amount of the variance in the adult tree community data ($R^2 = 0.53$ +/- 0.11). The distribution of posterior predictive values was nearly symmetric around observed Bray-Curtis values (Bayesian p-value = 0.49, supp. fig. 1). The posterior distribution of K_m was centered on 153 m (+/- 6.13 m), a shift of -47m from the prior estimate of 200m, with no overlap between posterior and prior (supp. fig. 2).

2. Linear model

The Bayesian linear model is visualized with 50% and 95% highest posterior densities ("HPD") for predicted BC values in fig. 2B. As constructed, the posterior distribution of our linear model explained only a small amount of the variance in the adult tree community data ($R^2 = 0.06$ +/- 0.01). The posterior predictive values tended to be lower than observed BC values (Bayesian p-value = 0.41, supp. fig. 1). The frequentist, least-squares linear regression model highly resembled the Bayesian model, also reporting very little variance explained ($R^2 = 0.06$ +/- 0.01, $p = <0.01$, fig. 2B).

Overall turnover by watershed crossings

When plotted as a function of watershed crossings (fig. 3), the widest range of Bray-curtis values occurred among sites within the same watershed (distance class 0), as did the lowest mean BC score (mean BC dissimilarity = 0.82, +/- 0.11). Following this, other mean BC values of comparisons of other distance class did not vary heavily, and are mostly statistically indistinguishable (Kruskal-Wallis H-test ($H(5, 1825)=73.27$, $p<0.01$, and Tukey HSD (p.adj=0.001), table 2 and table 3, fig. 3). After the first watershed crossing, further comparisons are all approximately equally dissimilar, with a mean of BC = 0.88 (+/- 0.086).

Hierarchical clustering of sites and prediction of current ecological state by land use history and elevation.

Tree communities first sort into two large groups that align well with previous land use - the first grouping contains sites that have historically experienced anthropogenic disturbances of any type, and the second grouping contains sites with no recorded history of anthropogenic disturbance, but all types of natural gap-forming disturbances (fig. 4). Each of these groups then sorts into two further groups (for a total of four clusters) . Cluster I contains all sites considered to be highly disturbed by agricultural use (RCA sites), and some sites with intermediate agricultural use as pasture (RG sites) (fig. 4). Cluster II consists entirely of RG sites, and also contains the majority of RG sites. Sites with no history of anthropogenic disturbances fall into two clusters, type III and type IV (fig. 4). Hierarchical clustering results are

congruent with NMS ordinations ([fig. 5](#), [supp. fig. 3](#)). A summary map giving both current ecological state and historical land-use/habitat is given in [fig. 6](#).

A linear model with these ecological states (forest types I-IV) as response variable, using the single variable of previous land-use/habitat as predictors, successfully predicts the sites that have been modified for pasture or agriculture, and it also well predicts non-anthropogenically sites to types III/IV ecological state (R^2 = 0.65 +/-0.04, [supp. fig. 4](#)). Posterior predictive checks were correct for 65% of sites, mostly for sites with type I and type II forests. However, this previous land-use/habitat-only model cannot distinguish among "natural forest" (III and IV) types.

Among sites with no history of anthropogenic disturbance ("natural forests", types III and IV), elevation strongly predicts the current ecological category of forest: sites with no history of anthropogenic disturbance at elevations of 1503 m (95% credible interval = 1469 m to 1537 m) or higher were categorized almost completely into type III forests, and below this elevation sites were categorized into type IV forests (bayesian logistic regression, R^2 = 0.87), [fig. 7](#), [fig. 6](#), [supp. fig. 5](#)).

A combined model of previous land-use/habitat and elevation explains most of the variance in total and predicts current ecological state very well (bayesian linear model, R^2 = 0.88 +/-0.16, [supp. fig. 6](#)), and performs much better on posterior predictions, with 87% of sites correctly predicted to their forest type.

Indicator species analysis

Indicator species were detected for all individual land-use-history/habitat types, all current ecological states (forest types), and for some combinations of site types. Indicator species analysis results are given in [table 4A](#) and [table 4B](#).

Spatial analyses

13 Moran's Eigenvector Maps variables were detected as correlated with changes in tree community ([fig. 8](#), cumulated, adjusted R^2 = 0.19, all MEMs p < 0.02 as components in forward model selection). Most were substantially correlated with 1-3 available environmental variables or a particular land-use/habitat history ([table 5](#)), with the exception of one highly localized MEM variable, MEM3. The most influential MEM variables in the model, MEM1 and MEM2 presented obvious geographic patterns. MEM1 (contributing R^2 = 0.026 to cumulative R^2, or 13.9% of explainable variance), indicates a general difference in tree communities between those of the northeastern and southwestern sides of the Los Cedros river ([supp fig. 7](#)). The second most influential MEM variable, MEM2, (contributing R^2 = 0.026 to cumulative R^2, or 13.6% of explainable variance) indicates a difference between the highest sites sampled in the study, which were situated along a ridge that runs ultimately to the highest points in the reserve, and lower elevations along this ridge system ([supp. fig. 8](#)). MEM8 also correlated with elevation, and with distance-to-nearest-stream, likely indicating a difference among "high and dry" sites and lower, wetter sites ([supp. fig. 9](#)).

Juvenile communities

Juvenile communities did not cluster within the groups presented by adult communities, either in terms of historical land-use/habitat (fig. 9A) or current ecological state of adult trees (fig. 9B). Rather, all juvenile communities radiate into new dissimilarity space from their respective adult communities, possibly indicating recruitment of new tree species and/or new species combinations at these sites. Of the 148 species of juvenile tree species observed in the study, 110 species were observed as both juvenile and mature specimens, and an additional 38 species were observed only as juvenile specimens in the study.

Juvenile tree communities change along the same axes as were observed for their respective adult tree communities. When categorized using historical land-use/habitat variable, RCA sites and related, similar RG sites are pulled in a negative direction along the NMS1 axis in fig. 9A, and non-anthropogenically disturbed sites (BC, BS, and CLB) are pulled in a positive direction along the NMS2 axis. When categorized by the current ecological state of their adult tree communities, the same pattern holds (fig. 9B): juvenile tree communities are different and outside of clusters formed by their respective adult communities, but change along the same axes as their adult communities. Two exceptions to this are observed: the juvenile tree communities of two type I forest sites have come to more closely resemble type III or forest (fig. 9B, sites 1.2 and 1.3). These sites have a land-use/habitat history of pasture-conversion (RG), and their adult tree communities cluster into a current ecological state of type I forest, the most anthropogenically disturbed group, often associated with RCA land-use history (fig. 5 and table. ___) . However their juvenile tree communities now resemble more "natural" forest types. Additionally, site 1.1 juvenile tree community was so unique as to be an outlier to the rest of the sites studied, and had to be removed for informative examination of the remaining sites (supp. fig. 10).

Deforestation in the region

Between 1990 and 2018, forest cover in the Cotacachi canton was reduced from an estimated 87,967 ha of native forest-cover in year 1990 to 71,739 ha in year 2018, an 18% reduction of total forest from 1990 levels, mostly due to conversion to agricultural land (fig. 10). In terms of total land cover, this equates to a shift from approximately 52% of total Cotacachi canton land cover being native forest to 42% of total land cover as native forest. Forest cover in Los Cedros increased from an estimated 5,094 ha of native forest-cover in year 1990 to 5,210 ha in year 2018, an 2.3% addition of total forest from 1990 levels, due mostly to the reforestation of former pasture (fig. 11). Forest cover in Bosque Protector Chontal decreased from an estimated 6,920 ha of native forest-cover in year 1990 to 6,565 ha in year 2018, a 5% reduction of total forest from 1990 levels, due mostly to conversion of forest to agricultural land uses (fig. 11).

Unlike earlier datasets, the 2018 MAE land cover dataset does not include pasture separately from other forms of agricultural land use, so it is difficult to quantify which types of agricultural land-use that are most commonly replacing forest in the region. However, personal communication and visitation with the surrounding communities, it is the authors assessment that much of the recent deforestation is undertaken to create pasture for cattle grazing.

Discussion

i. Stable states in the Andean cloud forest

In this study we examined 61 forest sites with known histories of anthropogenic disturbance or natural, gap-forming disturbance, and examined this history as a predictor of current ecological state. We can categorize the southern region of the Los Cedros reserve into four forest types, based on adult tree communities (fig. 4), and well predict these ecological states from past land-use/habitat-type and elevation. Sites that have no history of anthropogenic disturbance group readily into either forest types III or IV (fig. 4, fig. 5). These sites are characterized by endemic, forest-dependent tree species (table 4B). The difference among these "natural" forest types III and IV are strongly predicted by elevation (fig. 7, fig. 6, supp. fig. 5), exhibiting elevation-dependent ecological zonation often observed in montane tropical forests (Grubb, 1977; Leigh, 1975). Natural gap-forming disturbances do not change the species compositions of these sites to make them significantly different from other natural forest sites (fig. 5). As such, these natural forest types may represent stable equilibria with basins of attraction that give each some resilience to gap-forming disturbance (fig. 12).

There are indications that sites that have experienced intermediate disturbance (conversion to pasture, followed by reforestation) have entered a less stable, more pluripotent ecological state. Nearly all high and low elevation sites that were converted to pasture prior to reforestation (land-use/habitat type "RG") then developed into a single community type, ("forest type II"). This suggests a homogenizing effect on tree communities from this kind of disturbance, and a common successional response by the forest to after conversion to pasture and abandonment, regardless of elevation. In ordinations of community similarity (fig. 5 and supp fig. 3), Type II forest sites are situated in an intermediate position between all other states. It is therefore possible that the Type II forest type represents a low-sloped, convex unstable equilibrium that can sometimes allow sites to return to a natural forest state. In one case, we observe a site that has undergone anthropogenic disturbance and subsequently develops into a natural forest type (site 10.1, circled in fig. 4, fig. 5). Additionally, in the case of two pastured (RG) sites, their adult communities resemble the highly affected Type I forest, but their juvenile communities currently more closely resemble the "natural"type III forest (fig. 9, sites 1.2 and 1.3). In a different successional direction, several pastured (RG) sites developed into the novel Type I forest, the same forest type into which all intensive-agriculture sites developed (fig. 5).

The existence of this indeterminate, intermediate ecological state, forest type II, supports the hypothesis that primary cloud forest has some capacity to "repair" highly modified agricultural sites, but also indicates that this is not a certain outcome. This post-pasture successional trajectory appears to be very distinct from that which follows natural, gap-forming disturbances, and may be intermediate to severe agricultural conversions (cane production, followed by reforestation), which tended to drive sites to another state, forest Type I. The adult tree communities of sites which experienced intensive agriculture all developed into forest type I, suggesting that forest type I may be another stable equilibrium or novel ecosystem type (Hobbs et al., 2009). Additionally, all sites that were observed to be in Type I state are low-elevation sites, sites that presumably would have otherwise existed in a Type IV state, suggesting particular vulnerability of these lower-elevation forests to anthropogenic disturbance. However, all type I sites are also co-localized to the same area of the reserve (fig. 6), making it difficult to generalize this pattern to the rest of the study area.

Several forces may be at work in creating these equilibria we observed. Type III and type IV forests ("natural forest" types), gap-forming incidents appear to keep these sites within their basins of attraction, and the breakdown of these negative-feedbacks and ecological conditions that are likely responsible for shifts into forest type I equilibria:

Soil structure - While soil structure data are not available for this survey, researchers at field sites noted that sites that had undergone intensive, *cañaveral* agricultural use ("RCA" sites) had more compacted soils. Agricultural use can incur long-lasting legacy effects on soil structural and chemical characteristics, even after abandonment and reforestation (Sohng et al., 2017; Zhang et al., 2019). Deforestation has been shown to cause long-term changes in soil physical structure and microbial activity in soils, especially among plant-symbiotic microbes (Hartmann et al., 2012, 2014; Wang et al., 2018).

Soil seed bank depletion - In addition to soil compaction, several seasons of indiscriminate grazing or sugar cane culture probably greatly depleted the residual seed bank of forest plants (Bart & Davenport, 2015; K. R. Young et al., 1987). When soil-seed banks have been exhausted, recolonization of sites by forest plants must occur through dispersal of seeds to the reforesting site.

Large seeded plants and animal dispersal - Reserva Los Cedros hosts three species of primates: the brown-headed spider monkey (*Ateles fusciceps fusciceps*), the white-headed capuchin, (*Cebus capucinus*), and the mantled howler monkey, (*Alouatta palliata*), in addition to numerous other frugivorous birds and mammals (Roy et al., 2018). These animals likely play an essential role in closing gaps after disturbance. In recent natural gaps ('CLB'), we observed a single indicator species - *Endlicheria* sp. (table 4A). Some *Endolicheria* species have been suggested elsewhere to be primate dispersed (Bufalo et al., 2016), supplemental materials). Lower elevation natural forests (forest type IV) were characterized by copal trees (*Protium* and *Dacryodes* spp., table 4B), well known locally as primate food sources (Morelos-Juárez et al., 2015). In general, Type III and IV forest types were generally characterized by plants with larger seeds or fruits, often primate or otherwise vertebrate-dispersed, such as those in Lecythidaceae and Lauraceae, and *Garcinia* (Table 4B). This prevalence of trees with larger seeds/fruits is typical of tropical forests, where most trees produce animal-vectored seeds (Howe & Smallwood, 1982). Absent active transport of seeds by animals, colonization of pastures by forest trees is expected to be very slow (Cubiña & Aide, 2001), and to heavily favor wind dispersed seeds (Holl, 1999). Many ruderal species, conversely, are wind-dispersed (Grime, 1977) and are likely to readily colonize abandoned field sites, especially when aided by human vectors (Planchuelo et al., 2020). Given Los Cedros' relatively low elevation among montane tropical forests, it is not surprising that it's forests are heavily populated with large seeded species that likely rely on primate dispersal (Chapman et al., 2016).

Relative fluxes of local vs. exotic plant types - Most of the sites examined here are embedded in a landscape of primary forest, and may have been able to recruit seeds from forest dependent species even if they experienced seed bank depletion (Willson & Crome, 1989), while still insulated from outside seed sources. Approximately half of type I forest sites, however, were located on the edge of los Cedros, abutting neighboring farmland, probably allowing for extensive input of small seeded pioneer tree species such as *Saurauia* spp. The remaining type I sites were located along mule trails that supply the reserve, and were also therefore presumably repeatedly exposed to seeds from outside locations.

The above mechanisms are primarily "gap-filling" mechanisms, and may fail in the face of extensive anthropogenic disturbances such as habitat fragmentation and loss (for example,

Cramer et al. (2007)). It is probably important to note how dependent these processes may be on large animal dispersers, and especially on primates, as primate populations are in decline in the region (Peck et al., 2011; Roy et al., 2018). The importance of seed-dispersal services by large mammals and spider monkeys in particular can be appreciated by anticipating their loss: Peres et al. (2016) have predicted substantial loss of forest biomass in forests if large mammal populations are reduced, especially spider monkeys and tapirs. A similar positive feedback may be possible with the loss of forest-dependent birds due to deforestation. Frugivorous, forest-dwelling birds provide unique seed-dispersal service to forest trees but are highly interdependent with their habitat; their decline in diversity has been directly linked to vegetative diversity loss and deforestation (Morante‑Filho et al., 2018).

Other complex interactions may also be important: the presence of *Cecropia* spp. as indicator species in forest types with intermediate disturbance (forest type II), and appears less often in the novel type I forest. In this study they were also not observed as extremely prevalent in recent natural gaps. This is possibly important as *Cecropia* trees occupy a somewhat unique position of being an early-successional species while also producing large fruits and canopy structure characteristics useful to both primates and frugivorous birds (Morelos-Juárez et al., 2015; Navarro et al., 2018; Skutch, 1945). *Cecropia* trees may attract primate dispersers and forest-dependent birds back to a disturbed site, and therefore bring with them the heavier seeds of other forest-dependent, heavy-seeded species. At Los Cedros, the smallest primate dispersers, capuchin monkeys, are regularly observed in *Cecropia* trees. *Cecropia* spp. may therefore act as another gap-repairing negative feedback mechanism, one that plays out in cases of forest gaps of larger size or severity than the smaller natural gaps examined here. Thus the success or failure of *Cecropia* may play a deciding role that allows sites to return from the ecological state we have here observed as forest type II (resulting from intermediate disturbance), and potentially back to primary-like forest states. Note again, however, that this mechanism would depend upon the presence of large animal vectors, especially primates.

Given the relatively short period of time since disturbance prior to surveys (13-18 years), it is also very difficult to confidently project the stability of the here-proposed candidate equilibria to larger time scales, especially in times of deep ecosystem change due to climate change. Sites were surveyed in 2005, so an additional 16 years have passed since the collection of these data. Indeed, Loughlin et al. (2018) suggest that tropical Andean cloud forest in a nearby ecoregion may have a single long term ecological equilibrium. However, conditions in the Andes are changing dramatically from those that presumably sustained the long-term resiliency observed by Loughlin. Thus the candidate alternative states observed here should not be disregarded, they may mark the beginning of novel forest types in the region (Hobbs et al., 2009). For the moment, visual inspection of these sites in the present day confirms that unique tree communities continue to exist at Type I forest sites compared to surrounding older forest. Updated systematic surveys are now necessary to investigate the stability of the ecosystem types suggested here.

ii. Juvenile tree community

It is difficult to assign cause to the lack of structure observed in juvenile tree communities at Los Cedros (fig. 9). Much of the noise in these juvenile communities presumably stems from the incomplete filtering of young plants at each site (Baldeck et al., 2013; Ledo et al., 2015) - the environmental pressures that have shaped the adult tree community at each site likely had yet to totally act on the juvenile trees at time of sampling. However, it is also inevitable that the same global environmental changes that are acting on other forests throughout the world (Allen et al., 2010; Seidl et al., 2017), including cloud forests (Foster, 2001), are at play in the ancient

forests of Los Cedros, and are contributing to the reorganization of future tree communities, perhaps uncoupling ancient species associations. The disturbance regime under study (conversion to agriculture, followed by reforestation) also introduced new conditions and plant species to Los Cedros. Thus in these juvenile trees we may also be observing two sources of disturbance that could shift the forests of los Cedros out of the basins of attraction of the primary forest state, of very different scales: local land use change, and global climate change. In the terminology of Beisner et al. (2003), the former may still be considered a state variable change from which it is sometimes within the capacity of the cloud forest to rebound and recover to a primary forest state. The latter, however, is a deep shift in the parameters of the Andean cloud forest ecosystem, which will no doubt change the shape of the possible (Clerici et al., 2019; Feeley et al., 2011; Rehm & Feeley, 2015). It is not unreasonable to expect interactions between these two fundamental sources of ecological disturbance (Clerici et al., 2019; Pimm et al., 2014; Stork, 2010; Tovar et al., 2013).

iii. Beta-diversity and spatial heterogeneity in the Andean cloud forest

We examined patterns of distance decay in the tree communities of the southern region of los Cedros. When community turnover is modeled as a function of euclidean distance, a model in the form of an asymptote function that fits well to observed patterns of community-turnover. Our asymptote model suggests turnover in short distance, reporting half of maximum dissimilarity is reached in just ~150m, and a mean BC > 0.8 in comparisons with distances larger than 600m (fig. 2). When small watersheds are used as the basic spatial unit, rather than Euclidean distance, most of the decay in tree community similarity occurs with the first crossing over to a neighboring watershed (fig. 3). Following this, mean dissimilarity among sites increases slightly but remains uniformly high, and further comparisons are not statistically significantly different, meaning comparisons between sites five watersheds away were not on average more or less similar than comparisons of sites that were only two, three or four watersheds apart, because so much change in community composition has already occurred just within the first watershed crossing. This is supportive of the colloquial understanding that in the Andes, each small drainage can host an almost entirely distinctive community from its neighbors.

This may also give more insight into the processes that create the fine-scale of endemism often observed in Andean species. We hypothesized that this fine-scale, watershed-based community turnover is due to high dispersal limitation and high micro-site variability that result from the complex, dramatic topography of the Andes. This microsite variation is visible to some degree in our Moran's eigenvector maps and their environmental correlations (fig. 8, supp. figs 7-9, table 5): we see that environmental variables vary on scales that can be well understood in terms of change within and among small watersheds, and that these spatio-environmental patterns may explain up to 19% of variance in the plant community. This fraction may represent much of the environmental filtering that is occurring within the tree community at los Cedros. Dispersal limitation is harder to test, but the predominance of large-seeded species observed in our ancient forest sites suggests that many important tree species are dispersal limited to highly local scales and dependent on large animal dispersers to overcome this limitation. This preponderance of heavy-seeded species also might be well approximated by a symmetric dispersal-limited neutral model, a model which can generate significant small-scale spatial patterning in communities even without considering additional microsite environmental conditions (Hubbell, 2008).

iv. Conservation value of Los Cedros:

Since the sounding of alarm by Gentry , Myers et al., and numerous others (Gentry, 1992; Myers, 1988; Myers et al., 2000), there has been much concern about the future of the Chocó and the tropical Andean Biodiversity hotspot by conservation groups. However, there is little evidence that this call for conservation has resulted in sufficient meaningful change for the region of Los Cedros - quite the opposite. Instead, during this time frame, Cotacachi Canton has lost significant forest cover (fig. 10), as has Ecuador generally (Dodson & Gentry, 1991; Mosandl et al., 2008; Tapia-Armijos et al., 2015). In stark contrast, Los Cedros has well withstood the traditional pressures of timbering and settlement, as evidenced by its increase of forest cover during a time of net forest loss in the Cotacachi Canton. In addition to the historical habitat conversion from timber extraction and settlement, the tropical Andean Biodiversity hotspot has now found itself in a new center of metals mining exploration (Guayasamin et al., 2021; Roy et al., 2018; Vallejo Galárraga & Freslon, 2017), a new and entirely different extractive pressure on the region. Los Cedros itself is targeted by a junior mining partner (cornerstoneresources.com). Los Cedros has responded with a legal challenge that has reached Ecuador's highest courts, with implications for all Bosque Protectores in Ecuador (Guayasamin et al., 2021). Due to its proactive defensive legal efforts, Los Cedros' conservation effect is thus amplified even beyond the unusually effective physical protection of its forests.

Going forward, however, conservation of primary forest reserves such as Los Cedros will face novel challenges. It is widely understood that the high endemism of the Andean biodiversity hotspot makes it both a conservation priority and an especially difficult conservation challenge. It is less widely appreciated that forests, and especially primary forests, pose a major additional challenge to conservation. In most cases, we have only a crude understanding of the numerous ecological interactions required to maintain a primary ecosystem in its current state of biodiversity and ecosystem services, otherwise known as the ecological complexity of an ecosystem (Filotas et al., 2014; Messier et al., 2015). Ancient ecosystems are unique in large part due to their emergent complexity, and quantifying this complexity is usually difficult, requiring composite measures of numerous indices of forest physical structure, plant community, and other ecological properties (Loehle, 2004; McElhinny et al., 2005). In complex ecosystems, the loss of a species is multiplied by disruption of its interactions with other species, such as with trophic cascades (Ripple et al., 2016). The number of possible interactions among species undergoes quadratic growth as biodiversity increases linearly, so the immense biodiversity of the tropics presents a particular challenge to understanding the local ecological complexity. Primary forest fragments are also presumably subject to the well-known vulnerabilities of "island" - or insular - habitats (MacArthur & Wilson, 2001; Ross et al., 2002), namely the stochastic extinction of species without any nearby "mainland" to replenish populations. This is in addition to the uniquely dynamic, weather-dependent boundaries which make all cloud forests insular systems, primary or otherwise, and that may make them very vulnerable to shrinkage from climate change (Foster, 2001; Gentry, 1992; Rehm & Feeley, 2015). Thus in insular, highly interdependent ancient ecosystems such as primary cloud forest, these stochastic species extinctions may have greater cascading impacts than in simpler ecosystems. Even if completely protected from external changes, primary forest fragments such as los Cedros may ecologically simplify or "decay" with time if they are not sufficiently understood, buffered and interconnected. As such, the regional resiliency noted by Loughlin et al. (Loughlin et al., 2018) may be subject to novel vulnerabilities that were not so urgent historically. Michaels et al. (2020) have suggested that reforestation efforts should first attempt to diagnose the ecological conditions and feedbacks that allow a basin of attraction for a desired stable state to persist, and also find the minimum "nucleus" area needed for those

processes to proceed and grow. They and others, such as some proponents of rewilding (Fernández et al., 2017), suggest seeding landscapes with species consortiums intended to recreate ecological complexity as much as possible. Los Cedros is far from needing rewilding, but conservation efforts around Los Cedros and other ancient tropical forests will be greatly enriched by such an awareness of protecting and nurturing ecological complexity, beyond simple physical protection of reserve borders.

Given the state of fragmentation and loss of forest in the region, and the poor understanding of its numerous pollination and diaspore dispersal networks, especially the threatened state and habitat requirements of the primate dispersers of los Cedros (Peck et al., 2011), Los Cedros may actually be at or close to threshold stable patch size for maintaining its level of biodiversity. Los Cedros may therefore face an important juncture, with a choice of growing or subsiding. Further reduction or anthropogenic disturbance of cloud forest area risks non-linear, catastrophic changes. Additionally, the fragments of Andean primary cloud forest habitat that persist to date are rare and small enough that there is no economic justification for eliminating or heavily modifying them. Simply as a matter of scale, any economic benefit from the degradation of these forest fragments will not be large enough to justify their loss as reserves of genetic information, providers of ecosystem services, and protectors of species diversity. Other methods of economic development and poverty reduction are available (Balmford et al., 2002; Kocian et al., 2011). However, great opportunity for the future exists in Los Cedros: the forest of Los Cedros need not be seen as a delicate ecological island in a storm, nor is its protection an academic or moral exercise, or a desperate rear-guard conservation strategy. It is instead useful to think in the sense of Michaels (2020), supported by the long-term observations of Loughlin et al. (2018), and see the Los Cedros forest as a nucleus of primary cloud forest that has prospered for thousands of years, that has weathered the recent waves of deforestation, and that now stands ready to help reseed the new forests of the northern Andes.

Acknowledgements

The work would not have been possible without Jose DeCoux and his dedication to the conservation of Reserva Los Cedros. Several field assistants, volunteers and guides associated with Los Cedros helped with data collection: Gila Roder, Martín Obando, Manuel Moreno, Danny Cumba, Homero Sánchez, Fausto Lomas and Víctor Lomas.

Figures

Figure 1, A,B

Figure 1,C

Figure 2A

Figure 2B

Adult tree community dissimilarity with physical distance
Asymptote model fit

Figure 3

Figure 4

Figure 5

Adult tree community NMS
colored by forest type and historical land-use/habitat

Figure 6

Figure 7

Figure 8

Moran's eigenvector maps of adult tree community

Figure 9A

Adult and juvenile tree communities, with historical land-use/habitat

Figure 9B

Adult and juvenile tree communities, with adult forest type

Figure 10 A,B

1990 forest cover, Cotacachi canton

2018 forest cover, Cotacachi canton

Figure 11 A-D

Figure 12

Supplemental Figures

Supplemental figure 1

Supplemental figure 2

**Ordination of Hierarchical clustering results
(forest type)**

Supplemental figure 4

Supplemental figure 5

Supplemental figure 6

Supplemental figure 7

Supplemental figure 8

Supplemental figure 9

**Adult and juvenile tree communities,
with adult forest type, outlier retained**

Tables

Table 1 -Estimates of adult tree species diversity in survey area and permanent plot.

	sample size	species observed	chao	chao.se	jack1	jack1.se	jack2	boot	boot.se
Survey Plots	61	343	516.4	39.2	483.7	21.6	566.8	404.4	10.5
Permanent Plot (Trees)	121	43	65.4	13.8	61.8	4.3	72.7	51.4	2.2

Table 2 - Mean dissimilarity of survey sites, by number of watersheds crossed.

watersheds crossed	sample size	mean BC	stdDev BC
0	215	0.820	0.114
1	630	0.880	0.089
2	559	0.872	0.085
3	346	0.891	0.079
4	80	0.881	0.096

Table 3. Tukey's Honest Significant difference of comparisons among watershed distance classes.

group1	group2	meandiff	p-adj	lower	upper	reject
0	1	0.0601	0.001	0.0408	0.0795	TRUE
0	2	0.0516	0.001	0.0319	0.0713	TRUE
0	3	0.0708	0.001	0.0495	0.0921	TRUE
0	4	0.0609	0.001	0.0287	0.093	TRUE
1	2	-0.0085	0.4759	-0.0228	0.0057	FALSE
1	3	0.0106	0.393	-0.0058	0.0271	FALSE
1	4	0.0007	0.9	-0.0284	0.0299	FALSE
2	3	0.0192	0.0158	0.0024	0.036	TRUE
2	4	0.0093	0.9	-0.0201	0.0386	FALSE
3	4	-0.0099	0.9	-0.0404	0.0205	FALSE

Table 4A. Indicator species for historical land-use/habitat

Group BC

species	rpb.g	p.value	sig code
Moraceae.Pseudolmedia.rigida	0.423	0.0317	*
Moraceae.Ficus.subandina	0.406	0.049	*

Group BS

species	rpb.g	p.value	sig code
Rubiaceae.Psychotria.paeonia	0.478	0.0148	*
Lauraceae.Persea.pseudofasciculata	0.432	0.0351	*
Myrtaceae.Myrcia.aff..aliena	0.425	0.0377	*

Group CLB

species	rpb.g	p.value	sig code
Lauraceae.Endlicheria.cf..chalisea	0.431	0.0268	*

Group RCA

species	rpb.g	p.value	sig code
Boraginaceae.Cordia.colombiana	0.894	0.0001	***
Actinidaceae.Saurauia.sp.1	0.694	0.0003	***
Melastomataceae.Miconia.aff..brevitheca	0.683	0.0004	***
Moraceae.Ficus.caldasiana	0.667	0.0033	**
Staphyleaceae.Turpinia.occidentalis	0.663	0.0006	***
Moraceae.Clarisia.biflora	0.645	0.0022	**
Lauraceae.Nectandra.membranacea	0.61	0.0027	**
Verbenaceae.Aegiphila.alba	0.594	0.002	**
Moraceae.Ficus.andicola	0.539	0.0191	*

species	rpb.g	p.value	sig code
Lauraceae.Caryodaphnopsis.theobromifolia	0.535	0.0099	**
Lauraceae.Nectandra.aff..membranacea	0.513	0.0106	*

Group RG

species	rpb.g	p.value	sig code
Cecropiaceae.Cecropia.andina	0.662	0.0018	**
Cecropiaceae.Cecropia.ficifolia	0.58	0.0035	**
Cecropiaceae.Cecropia.sp.2	0.427	0.031	*
Melastomataceae.Meriania.tomentosa	0.415	0.0426	*

Group BC+BS

species	rpb.g	p.value	sig code
Lauraceae.Aniba.aff..hostmanniana	0.411	0.0471	*

Group RCA+RG

species	rpb.g	p.value	sig code
Solanaceae.Solanum.lepidotum	0.529	0.006	**
Solanaceae.Cestrum.megalophyllum	0.423	0.0268	*

Group BC+BS+CLB

species	rpb.g	p.value	sig code
Cyatheaceae.Alsophila.erinacea	0.42	0.0423	*

Table 4B. Indicator species for current ecological state (forest type)

Group I			
species	rpb.g	p.value	sig code
Boraginaceae.Cordia.colombiana	0.661	0.0001	***
Melastomataceae.Meriania.tomentosa	0.641	0.0001	***
Actinidaceae.Saurauia.sp.1	0.545	0.0003	***
Cyatheaceae.Cyathea.halonata	0.458	0.0033	**
Fab.Caesalpiniaceae.Senna.dariensis	0.449	0.0046	**
Moraceae.Ficus.caldasiana	0.42	0.0206	*
Melastomataceae.Leandra.subseriata	0.37	0.015	*
Melastomataceae.Miconia.aff..brevitheca	0.368	0.027	*
Piperaceae.Piper.fuliginosum	0.356	0.05	*
Staphyleaceae.Turpinia.occidentalis	0.346	0.0426	*

Group II			
species	rpb.g	p.value	sig code
Cecropiaceae.Cecropia.andina	0.811	0.0001	***
Cecropiaceae.Cecropia.sp.2	0.5	0.0009	***
Melastomataceae.sp.1	0.468	0.0051	**
Fab.Faboideae.Dussia.lehmannii	0.334	0.0444	*

Group III			
species	rpb.g	p.value	sig code
Myristicaceae.Otoba.gordoniifolia	0.524	0.0007	***

species	rpb.g	p.value	sig code
Cyatheaceae.Alsophila.erinacea	0.515	0.0015	**
Lauraceae.Persea.aff..pseudofasciculata	0.494	0.0022	**
Clusiaceae.Vismia.lauriformis	0.44	0.0065	**
Melastomataceae.Conostegia.aff..centronioides	0.418	0.0083	**
Arecaceae.Wettinia.aff..oxycarpa	0.403	0.0115	*
Lauraceae.Persea.pseudofasciculata	0.386	0.0217	*
Rubiaceae.Psychotria.paeonia	0.373	0.0264	*
Moraceae.Ficus.dulciaria	0.354	0.03	*

Group IV

species	rpb.g	p.value	sig code
Burseraceae.Dacryodes.cupularis	0.754	0.0001	***
Burseraceae.Protium.ecuadorense	0.638	0.0001	***
Clusiaceae.Garcinia.macrophylla	0.613	0.0001	***
Lauraceae.Beilschmiedia.aff..costaricensis	0.448	0.0018	**
Melastomataceae.Conostegia.superba	0.444	0.0024	**
Lauraceae.Ocotea.stenoneura	0.369	0.0454	*
Lecythidaceae.Gustavia.dodsonii	0.362	0.029	*
Styracaceae.Styrax.weberbaueri	0.361	0.0408	*
Moraceae.Pseudolmedia.rigida	0.355	0.0263	*

Group I+II

species	rpb.g	p.value	sig code
Cecropiaceae.Cecropia.ficifolia	0.52	0.0005	***
Solanaceae.Solanum.lepidotum	0.447	0.0055	**

	rpb.g	p.value	sig code
Cecropiaceae.Cecropia.reticulata	0.371	0.0244	*
Verbenaceae.Aegiphila.alba	0.348	0.0465	*
Urticaceae.Urera.caracasana	0.341	0.037	*

Group I+IV

species	rpb.g	p.value	sig code
Lauraceae.Caryodaphnopsis.theobromifolia	0.416	0.0092	**
Moraceae.Clarisia.biflora	0.403	0.0157	*

Group II+III

species	rpb.g	p.value	sig code
Annonaceae.Guatteria.megalophylla	0.411	0.0133	*
Moraceae.Ficus.cuatrecasana	0.364	0.0299	*

Group III+IV

species	rpb.g	p.value	sig code
Lauraceae.Aniba.aff..hostmanniana	0.401	0.0138	*
Capparaceae.Capparis.sp.	0.385	0.0267	*
Euphorbiaceae.Hieronyma.asperifolia	0.383	0.0226	*
Lecythidaceae.Eschweilera.integrifolia	0.379	0.0199	*
Lauraceae.Ocotea.insularis	0.366	0.0295	*
Rubiaceae.Faramea.oblongifolia	0.362	0.0291	*
Moraceae.Helicostylis.tovarensis	0.346	0.0444	*

Table 5. Moran's eigen vector maps and their statistically significant environmental correlations

	slope	DEM	aspect	exposeE	exposeN	toStream	BC	BS	CLB	RCA	RG
MEM1		-0.403								0.287	
MEM2	-0.344	-0.636								0.308	
MEM3											
MEM4					0.349						
MEM5										-0.503	
MEM6						0.306					
MEM7											0.385
MEM8		0.404				0.460					
MEM9										-0.451	
MEM10						-0.340					
MEM11						0.282					
MEM12											-0.410
MEM13							0.345				

Supplemental Table 1. Species lists

Supplemental Table 1A - Permanent plot species list

family	genus	species
Araceae	Anthurium	sp.
Begoniaceae	Begonia	parviflora
Burseraceae	Dacryodes	peruviana
Cardiopteridaceae	Citronella	cf. ilicifolia
Clusiaceae	Garcinia	madruno
Cyatheaceae	Alsophila	cuspidata
Euphorbiaceae	Acalypha	diversifolia
Fabaceae	Inga	aff. sapindoides
Fabaceae	Inga	sp. 1
Fabaceae	Inga	sp. 2
Fabaceae	Inga	sp. 3
Gesneriaceae	Gasteranthus	corallinus
Heliconiaceae	Heliconia	aff. fragilis
Lauraceae	Caryodaphnopsis	theobromifolia
Lauraceae	Nectandra	lineatifolia
Lauraceae	Persea	pseudofasciculata
Lauraceae	Rhodostemonodaphne	cf. cyclops
Malvaceae	Quararibea	casasecae
Marattiaceae	Danea	erecta
Melastomataceae	Meriania	tomentosa
Melastomataceae	Miconia	brevitheca
Melastomataceae	Miconia	cf. gracilis
Melastomataceae	Miconia	floribunda
Melastomataceae	Ossaea	micrantha
Moraceae	Clarisia	biflora
Moraceae	Ficus	cuatrecasiana
Moraceae	Helicostylis	tovarensis
Moraceae	Naucleopsis	ulei
Myrtaceae	Myrcia	fallax
Olaceae	Heisteria	concinna
Piperaceae	Piper	obliquum

Piperaceae	Piper	spoliatum
Primulaceae	Geissanthus	cf. longistamineus
Rubiaceae	Condaminea	corymbosa
Rubiaceae	Faramea	aff. oblongifolia
Rubiaceae	Palicourea	demissa
Rubiaceae	Pentagonia	sp.
Sabiaceae	Meliosma	occidentalis
Sapindaceae	Allophylus	punctatus
Sapindaceae	Allophylus	sp.
Solanaceae	Solanum	leptopdoum
Tapisciaceae	Huertea	glandulosa
Thymelaeaceae	Daphnopsis	grandis

Supplemental Table 1B - Rapid survey species list

family	genus	species
Actinidaceae	Saurauia	sp.
Actinidaceae	sp.	
Anacardiaceae	Mauria	cf. heterophylla
Anacardiaceae	Tapirira	guianensis
Annonaceae	Guatteria	cf. megalophylla
Annonaceae	Guatteria	megalophylla
Annonaceae	Guatteria	sp.
Annonaceae	Rollinia	aff. dolichopetala
Annonaceae	Rollinia	edulis
Annonaceae	sp.	
Aquifoliaceae	Ilex	cf. yurumanguinis
Aquifoliaceae	Ilex	laurina
Araliaceae	Dendropanax	aff. macrophyllum
Araliaceae	Dendropanax	cf. macrophyllum
Araliaceae	Oreopanax	confusus
Araliaceae	Oreopanax	ecuadorensis
Arecaceae	Chamaedorea	pinnatifrons
Arecaceae	Hyospathe	elegans
Arecaceae	Wettinia	aff. oxycarpa
Arecaceae	Wettinia	quinaria

Asteraceae	Criptoniosis	cf. occidentalis
Asteraceae	Critoniopsis	occidentalis
Asteraceae	Piptocoma	discolor
Asteraceae	sp.	
Asteraceae	Vernonanthura	patens
Bombacaceae	Matisia	aff. bolivarii
Bombacaceae	Matisia	castano
Bombacaceae	Matisia	sp.
Bombacaceae	Ochroma	pyramidale
Bombacaceae	Pachira	patinoi
Bombacaceae	Quararibea	aff. asterolepis
Bombacaceae	sp.	
Boraginaceae	Cordia	aff. mexiana
Boraginaceae	Cordia	colombiana
Boraginaceae	Cordia	mexicana
Boraginaceae	Cordia	sp.
Boraginaceae	Tournefortia	cf. ramosissima
Brunelliaceae	Brunellia	acostae
Brunelliaceae	Brunellia	comocladifolia
Burseraceae	Dacryodes	cf. cupularis
Burseraceae	Dacryodes	cupularis
Burseraceae	Protium	ecuadorense
Burseraceae	Protium	sp.
Burseraceae	sp.	

Capparaceae	Capparis	macrophylla
Capparaceae	Capparis	sp.
Cecropiaceae	Cecropia	aff. andina
Cecropiaceae	Cecropia	aff. reticulata
Cecropiaceae	Cecropia	andina
Cecropiaceae	Cecropia	ficifolia
Cecropiaceae	Cecropia	reticulata
Cecropiaceae	Cecropia	sp.1
Cecropiaceae	Cecropia	sp.2
Cecropiaceae	Coussapoa	contorta
Chloranthaceae	Hedyosmum	cuatrecazanum
Chrysobalanaceae	Couepia	sp.
Chrysobalanaceae	Licania	durifolia
Chrysobalanaceae	Licania	sp.
Clusiaceae	Chrysochlamys	aff. dependens
Clusiaceae	Chrysochlamys	dependens
Clusiaceae	Chrysochlamys	sp.
Clusiaceae	Clusia	aff. dixonii
Clusiaceae	Clusia	aff. hydrogera
Clusiaceae	Clusia	dixonii
Clusiaceae	Clusia	sp.
Clusiaceae	Clusia	trochimorfis
Clusiaceae	Garcinia	macrophylla
Clusiaceae	Vismia	lauriformis

Cunoniaceae	Weinmannia	balbisiana
Cyatheaceae	Alsophila	cuspidata
Cyatheaceae	Alsophila	erinacea
Cyatheaceae	Cyathea	caracasana
Cyatheaceae	Cyathea	halonata
Cyatheaceae	sp.	
Dichapetalaceae	Stephanopodium	angulatum
Dichapetalaceae	Stephanopodium	longipedicellatum
Dicksoniaceae	Dicksonia	sellowiana
Elaeocarpaceae	Sloanea	cf. grandiflora
Elaeocarpaceae	Sloanea	sp. 1
Elaeocarpaceae	Sloanea	sp. 2
Ericaceae	Psammisia	aberrans
Ericaceae	Psammisia	sodiroi
Erythroxylaceae	Erythroxylum	cf. novogranatense
Euphorbiaceae	Acalypha	aff. diversifolia
Euphorbiaceae	Acalypha	diversifolia
Euphorbiaceae	Alchornea	cf. grandis
Euphorbiaceae	Alchornea	cf. leptogyna
Euphorbiaceae	Hieronyma	alchorneoides
Euphorbiaceae	Hieronyma	asperifolia
Euphorbiaceae	Plukenetia	volubilis
Euphorbiaceae	Sapium	marmieri
Euphorbiaceae	Sapium	stylare

Euphorbiaceae	Tetrorchidium	andinum
Euphorbiaceae	Tetrorchidium	macrophyllum
Fab-Caesalpiniaceae	Senna	dariensis
Fab-Faboideae	Dussia	lehmannii
Fab-Faboideae	Erythrina	megistophylla
Fab-Faboideae	Erythrina	schimpffii
Fab-Faboideae	Erythrina	sp.
Fab-Faboideae	Lonchocarpus	sp.
Fab-Faboideae	sp.	
Fab-Mimosoideae	Abarema	sp.
Fab-Mimosoideae	Cojoba	arborea
Fab-Mimosoideae	Inga	aff. lallensis
Fab-Mimosoideae	Inga	aff. nobilis
Fab-Mimosoideae	Inga	aff. sapindoides
Fab-Mimosoideae	Inga	aff. villosissima
Fab-Mimosoideae	Inga	lallensis
Fab-Mimosoideae	Inga	oerstediana
Fab-Mimosoideae	Inga	punctata
Fab-Mimosoideae	Inga	sp.
Fab-Mimosoideae	Inga	villosissima
Fab-Mimosoideae	sp.	
Flacourtiaceae	Banara	guianensis
Flacourtiaceae	Banara	regia

Flacourtiaceae	Banara	riparia
Flacourtiaceae	Casearia	cajambrensis
Flacourtiaceae	Casearia	sylvestris
Flacourtiaceae	sp.	
Hippocrataceae	Salacia	cordata
Indeterminado	sp. 1	
Indeterminado	sp. 10	
Indeterminado	sp. 11	
Indeterminado	sp. 12	
Indeterminado	sp. 13	
Indeterminado	sp. 14	
Indeterminado	sp. 15	
Indeterminado	sp. 16	
Indeterminado	sp. 17	
Indeterminado	sp. 18	
Indeterminado	sp. 19	
Indeterminado	sp. 2	
Indeterminado	sp. 3	
Indeterminado	sp. 4	
Indeterminado	sp. 5	
Indeterminado	sp. 6	
Indeterminado	sp. 7	

Indeterminado	sp. 8	
Indeterminado	sp. 9	
Lauraceae	Aniba	aff. hostmanniana
Lauraceae	Aniba	cf. hostmanniana
Lauraceae	Beilschmiedia	aff. costaricensis
Lauraceae	Beilschmiedia	alloiophylla
Lauraceae	Beilschmiedia	sp.
Lauraceae	Caryodaphnopsis	theobromifolia
Lauraceae	Cinnamomum	triplinerve
Lauraceae	Endlicheria	cf. chalisea
Lauraceae	Endlicheria	cf. sericea
Lauraceae	Endlicheria	cf. tessmannii
Lauraceae	Endlicheria	formosa
Lauraceae	Endlicheria	klugii
Lauraceae	Endlicheria	sp.
Lauraceae	Licaria	aff. tomentosa
Lauraceae	Licaria	cf. tomentosa
Lauraceae	Nectandra	acutifolia
Lauraceae	Nectandra	aff. guadaripo
Lauraceae	Nectandra	aff. laurel
Lauraceae	Nectandra	aff. membranacea
Lauraceae	Nectandra	aff. parviflora
Lauraceae	Nectandra	aff. Purpurea
Lauraceae	Nectandra	guadaripo

Lauraceae	Nectandra	membranacea
Lauraceae	Nectandra	obtusata
Lauraceae	Nectandra	sp. 1
Lauraceae	Nectandra	sp. 2
Lauraceae	Nectandra	sp. 3
Lauraceae	Nectandra	sp.1
Lauraceae	Ocotea	aff. cernua
Lauraceae	Ocotea	aff. floccifera
Lauraceae	Ocotea	aff. heterochroma
Lauraceae	Ocotea	aff. insularis
Lauraceae	Ocotea	aff. oblonga
Lauraceae	Ocotea	cernua
Lauraceae	Ocotea	floccifera
Lauraceae	Ocotea	insularis
Lauraceae	Ocotea	macrophylla
Lauraceae	Ocotea	oblonga
Lauraceae	Ocotea	sp. 1
Lauraceae	Ocotea	sp. 2
Lauraceae	Ocotea	stenoneura
Lauraceae	Persea	aff. areolatocostae
Lauraceae	Persea	aff. pseudofasciculata
Lauraceae	Persea	povedae
Lauraceae	Persea	pseudofasciculata
Lauraceae	Persea	rigens

Lauraceae	Pleurothyrium	aff. giganthum
Lauraceae	Pleurothyrium	cinereum
Lauraceae	Pleurothyrium	giganthum
Lauraceae	Pleurothyrium	glabritepalum
Lauraceae	Pleurothyrium	sp.
Lauraceae	Rhodostemonodaphne	aff. cyclops
Lauraceae	Rhodostemonodaphne	cyclops
Lauraceae	sp. 1	
Lauraceae	sp. 2	
Lauraceae	sp. 3	
Lauraceae	sp. 4	
Lecythidaceae	Eschweilera	caudiculata
Lecythidaceae	Eschweilera	integrifolia
Lecythidaceae	Eschweilera	rimbachii
Lecythidaceae	Gustavia	dodsonii
Magnoliaceae	Talauma	cf. gilbertoi
Malphigiaceae	sp.	
Malvaceae	Wercklea	ferox
Malvaceae	Wercklea	sp.
Melastomataceae	Conostegia	aff. centronioides
Melastomataceae	Conostegia	cuatrecasii
Melastomataceae	Conostegia	superba
Melastomataceae	Leandra	subseriata

Melastomataceae	Meriania	finicola
Melastomataceae	Meriania	maxima
Melastomataceae	Meriania	tomentosa
Melastomataceae	Miconia	aff. brevitheca
Melastomataceae	Miconia	aggregata
Melastomataceae	Miconia	cf. subnodosa
Melastomataceae	Miconia	dapsiliflora
Melastomataceae	Miconia	loveyoides
Melastomataceae	Miconia	sp.
Melastomataceae	Ossaea	cf. micrantha
Melastomataceae	Ossaea	micrantha
Melastomataceae	Ossaea	robusta
Melastomataceae	sp. 1	
Melastomataceae	sp. 3	
Melastomataceae	sp.1	
Melastomataceae	sp.2	
Melastomataceae	sp.3	
Melastomataceae	Topobea	dodsonorum
Meliaceae	Guarea	aff. macrophylla
Meliaceae	Guarea	cartaguenya
Meliaceae	Guarea	cf. pubescens
Meliaceae	Guarea	cf. subandina
Meliaceae	Guarea	kunthiana
Meliaceae	Ruagea	aff. glabra

Meliaceae	Ruagea	cf. glabra
Meliaceae	Ruagea	cf. pubescens
Meliaceae	Ruagea	glabra
Meliaceae	Ruagea	pubescens
Meliaceae	Ruagea	trisperma
Meliaceae	sp. 1	
Meliaceae	sp.1	
Meliaceae	sp.2	
Meliaceae	Trichilia	aff. septentrionalis
Meliaceae	Trichilia	septentrionalis
Meliaceae	Trichilia	sp.
Meliaceae	Trichilia	sp. 1
Monimiaceae	Mollinedia	sp.
Moraceae	Clarisia	biflora
Moraceae	Ficus	andicola
Moraceae	Ficus	caldasiana
Moraceae	Ficus	cf. caldasiana
Moraceae	Ficus	cotopaxiensis
Moraceae	Ficus	cuatrecasana
Moraceae	Ficus	dulciaria
Moraceae	Ficus	macbridei
Moraceae	Ficus	maxima
Moraceae	Ficus	mutisii
Moraceae	Ficus	sibundoya

Moraceae	Ficus	sp.
Moraceae	Ficus	subandina
Moraceae	Ficus	tonduzii
Moraceae	Ficus	trianae
Moraceae	Helicostylis	tomentosa
Moraceae	Helicostylis	tovarensis
Moraceae	Morus	insignis
Moraceae	Naucleopsis	naga
Moraceae	Perebea	aff. xanthochyma
Moraceae	Pseudolmedia	macrophylla
Moraceae	Pseudolmedia	rigida
Moraceae	Sorocea	jaramilloi
Moraceae	Sorocea	steinbachii
Moraceae	sp.	
Moraceae	sp. 1	
Moraceae	Trophis	racemosa
Moraceae	vs. Perebea	angustifolia
Myristicaceae	Otoba	gordoniifolia
Myristicaceae	Virola	aff. sebifera
Myristicaceae	Virola	sp.
Myrsinaceae	Ardisia	cf. colombiana
Myrsinaceae	Cybianthus	humilis
Myrsinaceae	Geissanthus	aff. longistamineus
Myrsinaceae	Geissanthus	aff. pinchinchana

Myrsinaceae	Stylogyne	ambigua
Myrsinaceae	Weigeltia	cf. goudotiana
Myrtaceae	Calyptranthes	sp.
Myrtaceae	Eugenia	cf. florida
Myrtaceae	Eugenia	florida
Myrtaceae	Eugenia	sp.
Myrtaceae	Myrcia	aff. aliena
Myrtaceae	Myrcia	aff. fallax
Myrtaceae	Myrcia	cf. aliena
Myrtaceae	Myrcia	cf. fallax
Myrtaceae	Myrcia	fallax
Myrtaceae	Myrcia	splendens
Myrtaceae	sp.	
Ochnaceae	Ouratea	aff. weberbaueri
Ochnaceae	Ouratea	weberbaueri
Olacaceae	Heisteria	sp.
Piperaceae	Piper	aduncum
Piperaceae	Piper	aff. nudibracteatun
Piperaceae	Piper	cf. imperiale
Piperaceae	Piper	fuliginosum
Piperaceae	Piper	imperiale
Piperaceae	Piper	sp.
Rubiaceae	Agouticarpa	grandistipula
Rubiaceae	Condaminea	corymbosa

Rubiaceae	Elaeagia	cf. karstenii
Rubiaceae	Elaeagia	karstenii
Rubiaceae	Elaeagia	utilis
Rubiaceae	Faramea	aff. calyptrata
Rubiaceae	Faramea	flavicans
Rubiaceae	Faramea	fragans
Rubiaceae	Faramea	glandulosa
Rubiaceae	Faramea	oblongifolia
Rubiaceae	Faramea	sp.
Rubiaceae	Guettarda	crispiflora
Rubiaceae	Guettarda	hirsuta
Rubiaceae	Hoffmannia	cf. parviflora
Rubiaceae	Isertia	laevis
Rubiaceae	Isertia	pittieri
Rubiaceae	Palicourea	acanthacea
Rubiaceae	Palicourea	aff. chignul
Rubiaceae	Palicourea	amethystina
Rubiaceae	Palicourea	cf. stipularis
Rubiaceae	Palicourea	chimboracensis
Rubiaceae	Palicourea	gentryi
Rubiaceae	Palicourea	gibbosa
Rubiaceae	Palicourea	sp.
Rubiaceae	Palicourea	stipularis
Rubiaceae	Palicourea	thyrsiflora

Rubiaceae	Pentagonia	macrophylla
Rubiaceae	Psychotria	cf. steyermarkii
Rubiaceae	Psychotria	flaviflora
Rubiaceae	Psychotria	gentryi
Rubiaceae	Psychotria	paeonia
Rubiaceae	Psychotria	pilosa
Rubiaceae	Rondeletia	reflexa
Rubiaceae	sp. 1	
Rubiaceae	sp.1	
Rubiaceae	sp.2	
Rubiaceae	sp.3	
Rubiaceae	Tocoyena	pittieri
Rutaceae	Zanthoxylum	mauriifolium
Rutaceae	Zanthoxylum	sp.
Sabiaceae	Meliosma	aff. occidentalis
Sabiaceae	Meliosma	occidentalis
Sapindaceae	Allophyllus	floribundus
Sapindaceae	Allophyllus	incanus
Sapindaceae	Allophyllus	sp.
Sapindaceae	Cupania	aff. cinerea
Sapindaceae	Talisia	aff. nervosa
Sapindaceae	Talisia	cf. nervosa
Sapotaceae	Chrysophyllum	aff. venezuelanense
Sapotaceae	Chrysophyllum	cf. argenteum

Sapotaceae	Chrysophyllum	manaosense
Sapotaceae	Chrysophyllum	sp.
Sapotaceae	Pouteria	torta
Simaroubaceae	Simarouba	amara
Siparumaceae	Siparuna	laurifolia
Siparumaceae	Siparuna	salvioides
Siparumaceae	Siparuna	sp.
Siparunaceae	Siparuna	salvioides
Solanaceae	Cestrum	megalophyllum
Solanaceae	Markea	pavonii
Solanaceae	Solanum	aff. circinatum
Solanaceae	Solanum	aff. cucullatum
Solanaceae	Solanum	cf. circinatum
Solanaceae	Solanum	cucullatum
Solanaceae	Solanum	lepidotum
Solanaceae	Solanum	sp.
Staphyleaceae	Huertea	glandulosa
Staphyleaceae	Turpinia	occidentalis
Styracaceae	Styrax	weberbaueri
Symplocaceae	Symplocos	fuliginosa
Theaceae	Freziera	reticulata
Theophrastaceae	Clavija	eggersiana
Theophrastaceae	Clavija	membranacea
Thymelaeaceae	Schoenobiblus	cf. daphnoides

Thymelaeaceae	Schoenobiblus	daphnoides
Tiliaceae	sp.	
Ulmaceae	Trema	micrantha
Urticaceae	Boehmeria	caudata
Urticaceae	Phenax	aff. rugosus
Urticaceae	Urera	aff. caracasana
Urticaceae	Urera	baccifera
Urticaceae	Urera	caracasana
Verbenaceae	Aegiphila	alba
Violaceae	Gloeospermum	longifolium

Supplemental Table 2. Species diversity estimates for historical land-use/habitat types

Habitat/Land-use	species observed	chao	chao.se	jack1	jack1.se	jack2	boot	boot.se	n
BC	153	282.7	38.5	234.6	26.1	287.5	187.9	11.6	12
BS	151	262.5	35.3	222.5	22.3	268.6	182.0	10.0	12
CLB	146	288.0	43.2	226.7	26.8	281.1	180.2	11.5	12
RCA	52	103.0	25.3	78.3	15.5	92.8	63.6	6.9	4
RG	186	302.9	33.7	271.7	22.1	324.8	223.6	10.9	21

Works cited

Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E. H. (Ted), Gonzalez, P., Fensham, R., Zhang, Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S. W., Semerci, A., & Cobb, N. (2010). A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, *259*(4), 660–684. https://doi.org/10.1016/j.foreco.2009.09.001

Arita, H. T., & Vázquez-Domínguez, E. (2008). The tropics: Cradle, museum or casino? A dynamic null model for latitudinal gradients of species diversity. *Ecology Letters*, *11*(7), 653–663. https://doi.org/10.1111/j.1461-0248.2008.01197.x

Ashour, S. K., & Abdel-hameed, M. A. (2010). Approximate skew normal distribution. *Journal of Advanced Research*, *1*(4), 341–350. https://doi.org/10.1016/j.jare.2010.06.004

Baldeck, C. A., Harms, K. E., Yavitt, J. B., John, R., Turner, B. L., Valencia, R., Navarrete, H., Bunyavejchewin, S., Kiratiprayoon, S., & Yaacob, A. (2013). Habitat filtering across tree life stages in tropical forest communities. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1766), 20130548.

Balmford, A., Bruner, A., Cooper, P., Costanza, R., Farber, S., Green, R. E., Jenkins, M., Jefferiss, P., Jessamy, V., Madden, J., Munro, K., Myers, N., Naeem, S., Paavola, J., Rayment, M., Rosendo, S., Roughgarden, J., Trumper, K., & Turner, R. K. (2002). Economic Reasons for Conserving Wild Nature. *Science*, *297*(5583), 950–953. JSTOR.

Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O. U., Swartz, B., Quental, T. B., Marshall, C., McGuire, J. L., Lindsey, E. L., Maguire, K. C., Mersey, B., & Ferrer, E. A. (2011). Has the Earth's sixth mass extinction already arrived? *Nature*, *471*(7336), 51–57. https://doi.org/10.1038/nature09678

Bart, D., & Davenport, T. (2015). The influence of legacy impacted seed banks on vegetation

recovery in a post-agricultural fen complex. *Wetlands Ecology and Management*, *23*(3), 405–418. https://doi.org/10.1007/s11273-014-9391-1

Bartos, M. (2021). *Pysheds* (0.2.7) [Python]. https://mattbartos.com/pysheds/

Beisner, B. E., Haydon, D. T., & Cuddington, K. (2003). Alternative stable states in ecology. *Frontiers in Ecology and the Environment*, *1*(7), 376–382. https://doi.org/10.1890/1540-9295(2003)001[0376:ASSIE]2.0.CO;2

Bhaskar, R., Dawson, T. E., & Balvanera, P. (2014). Community assembly and functional diversity along succession post-management. *Functional Ecology*, *28*(5), 1256–1265. https://doi.org/10.1111/1365-2435.12257

Borchsenius, F. (1997). Patterns of plant species endemism in Ecuador. *Biodiversity & Conservation*, *6*(3), 379–399. https://doi.org/10.1023/A:1018312724137

Bray, J. R., & Curtis, J. T. (1957). An ordination of the upland forest communities of southern Wisconsin. *Ecological Monographs*, *27*(4), 325–349.

Brehm, G., Hebert, P. D. N., Colwell, R. K., Adams, M.-O., Bodner, F., Friedemann, K., Möckel, L., & Fiedler, K. (2016). Turning Up the Heat on a Hotspot: DNA Barcodes Reveal 80% More Species of Geometrid Moths along an Andean Elevational Gradient. *PLOS ONE*, *11*(3), e0150327. https://doi.org/10.1371/journal.pone.0150327

Bufalo, F. S., Galetti, M., & Culot, L. (2016). Seed Dispersal by Primates and Implications for the Conservation of a Biodiversity Hotspot, the Atlantic Forest of South America. *International Journal of Primatology*, *37*(3), 333–349. https://doi.org/10.1007/s10764-016-9903-3

Bunin, G. (2021). Directionality and community-level selection. *Oikos*, *130*(4), 489–500. https://doi.org/10.1111/oik.07214

Carmody, M., Waszczak, C., Idänheimo, N., Saarinen, T., & Kangasjärvi, J. (2016). ROS signalling in a destabilised world: A molecular understanding of climate change. *Journal of Plant Physiology*, *203*, 69–83. https://doi.org/10.1016/j.jplph.2016.06.008

Chambers, J. Q., Negron-Juarez, R. I., Marra, D. M., Di Vittorio, A., Tews, J., Roberts, D., Ribeiro, G. H., Trumbore, S. E., & Higuchi, N. (2013). The steady-state mosaic of disturbance and succession across an old-growth Central Amazon forest landscape. *Proceedings of the National Academy of Sciences*, *110*(10), 3949–3954.

Chang, C. C., & Turner, B. L. (2019). Ecological succession in a changing world. *Journal of Ecology*, *107*(2), 503–509. https://doi.org/10.1111/1365-2745.13132

Chapman, H., Cordeiro, N. J., Dutton, P., Wenny, D., Kitamura, S., Kaplin, B., Melo, F. P. L., & Lawes, M. J. (2016). Seed-dispersal ecology of tropical montane forests. *Journal of Tropical Ecology*, *32*(5), 437–454. https://doi.org/10.1017/S0266467416000389

Churchill, S. P. (2009). Moss diversity and endemism of the tropical Andes. *Annals of the Missouri Botanical Garden*, 434–449.

Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., Aubinet, M., Buchmann, N., Bernhofer, C., Carrara, A., Chevallier, F., De Noblet, N., Friend, A. D., Friedlingstein, P., Grünwald, T., Heinesch, B., Keronen, P., Knohl, A., Krinner, G., … Valentini, R. (2005). Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature*, *437*(7058), 529–533. https://doi.org/10.1038/nature03972

Clements, F. E. (1916). *Plant succession: An analysis of the development of vegetation* (Issue 242). Carnegie Institution of Washington.

Clerici, N., Cote-Navarro, F., Escobedo, F. J., Rubiano, K., & Villegas, J. C. (2019). Spatio-temporal and cumulative effects of land use-land cover and climate change on two ecosystem services in the Colombian Andes. *Science of The Total Environment*, *685*, 1181–1192. https://doi.org/10.1016/j.scitotenv.2019.06.275

Colwell, R. K., Chao, A., Gotelli, N. J., Lin, S.-Y., Mao, C. X., Chazdon, R. L., & Longino, J. T. (2012). Models and estimators linking individual-based and sample-based rarefaction, extrapolation and comparison of assemblages. *Journal of Plant Ecology*, *5*(1), 3–21. https://doi.org/10.1093/jpe/rtr044

Connell, J. H., & Slatyer, R. O. (1977). Mechanisms of Succession in Natural Communities and

    Their Role in Community Stability and Organization. *The American Naturalist*, *111*(982),

    1119–1144. https://doi.org/10.1086/283241

Corbin, J. D., & Holl, K. D. (2012). Applied nucleation as a forest restoration strategy. *Forest*

    Ecology and Management, *265*, 37–46. https://doi.org/10.1016/j.foreco.2011.10.013

Cowles, H. C. (1899). The Ecological Relations of the Vegetation on the Sand Dunes of Lake

    Michigan. Part I.-Geographical Relations of the Dune Floras. *Botanical Gazette*, *27*(2),

    95–117. https://doi.org/10.1086/327796

Cramer, J. M., Mesquita, R. C. G., & Bruce Williamson, G. (2007). Forest fragmentation

    differentially affects seed dispersal of large and small-seeded tropical trees. *Biological*

    Conservation, *137*(3), 415–423. https://doi.org/10.1016/j.biocon.2007.02.019

Crausbay, S. D., & Martin, P. H. (2016). Natural disturbance, vegetation patterns and ecological

    dynamics in tropical montane forests. *Journal of Tropical Ecology*, *32*(5), 384–403.

    https://doi.org/10.1017/S0266467416000328

Cubiña, A., & Aide, T. M. (2001). The Effect of Distance from Forest Edge on Seed Rain and

    Soil Seed Bank in a Tropical Pasture. *Biotropica*, *33*(2), 260–267.

    https://doi.org/10.1111/j.1744-7429.2001.tb00177.x

Curtis, J. T., & Greene, H. C. (1949). A Study of Relic Wisconsin Prairies by the

    Species-Presence Method. *Ecology*, *30*(1), 83–92. https://doi.org/10.2307/1932279

Day, J. W., Christian, R. R., Boesch, D. M., Yáñez-Arancibia, A., Morris, J., Twilley, R. R.,

    Naylor, L., Schaffner, L., & Stevenson, C. (2008). Consequences of Climate Change on

    the Ecogeomorphology of Coastal Wetlands. *Estuaries and Coasts*, *31*(3), 477–491.

    https://doi.org/10.1007/s12237-008-9047-6

De Cáceres, M., Legendre, P., & Moretti, M. (2010). Improving indicator species analysis by

    combining groups of sites. *Oikos*, *119*(10), 1674–1684.

    https://doi.org/10.1111/j.1600-0706.2010.18334.x

Dijkstra, E. W. (1959). A note on two problems in connexion with graphs. *Numerische Mathematik*, *1*(1), 269–271.

Dodson, C. H., & Gentry, A. H. (1991). Biological Extinction in Western Ecuador. *Annals of the Missouri Botanical Garden*, *78*(2), 273. https://doi.org/10.2307/2399563

Doney, S. C., Ruckelshaus, M., Emmett Duffy, J., Barry, J. P., Chan, F., English, C. A., Galindo, H. M., Grebmeier, J. M., Hollowed, A. B., Knowlton, N., Polovina, J., Rabalais, N. N., Sydeman, W. J., & Talley, L. D. (2012). Climate Change Impacts on Marine Ecosystems. *Annual Review of Marine Science*, *4*(1), 11–37. https://doi.org/10.1146/annurev-marine-041911-111611

Draper, F. C., Asner, G. P., Coronado, E. N. H., Baker, T. R., García‐Villacorta, R., Pitman, N. C. A., Fine, P. V. A., Phillips, O. L., Gómez, R. Z., Guerra, C. A. A., Arévalo, M. F., Martínez, R. V., Brienen, R. J. W., Monteagudo‐Mendoza, A., Montenegro, L. A. T., Sandoval, E. V., Roucoux, K. H., Arévalo, F. R. R., Acuy, Í. M., … Baraloto, C. (2019). Dominant tree species drive beta diversity patterns in western Amazonia. *Ecology*, *100*(4), e02636. https://doi.org/10.1002/ecy.2636

Dray, S., Pélissier, R., Couteron, P., Fortin, M.-J., Legendre, P., Peres-Neto, P. R., Bellier, E., Bivand, R., Blanchet, F. G., Cáceres, M. D., Dufour, A.-B., Heegaard, E., Jombart, T., Munoz, F., Oksanen, J., Thioulouse, J., & Wagner, H. H. (2012). Community ecology in the age of multivariate multiscale spatial analysis. *Ecological Monographs*, *82*(3), 257–275. https://doi.org/10.1890/11-1183.1

Ellis, E. C., Goldewijk, K. K., Siebert, S., Lightman, D., & Ramankutty, N. (2010). Anthropogenic transformation of the biomes, 1700 to 2000. *Global Ecology and Biogeography*, *19*(5), 589–606. https://doi.org/10.1111/j.1466-8238.2010.00540.x

Espírito-Santo, F. D., Keller, M. M., Linder, E., Oliveira Junior, R. C., Pereira, C., & Oliveira, C. G. (2014). Gap formation and carbon cycling in the Brazilian Amazon: Measurement using high-resolution optical remote sensing and studies in large forest plots. *Plant

Ecology & Diversity, 7(1–2), 305–318. https://doi.org/10.1080/17550874.2013.795629

Fadrique, B., Báez, S., Duque, Á., Malizia, A., Blundo, C., Carilla, J., Osinaga-Acosta, O.,

Malizia, L., Silman, M., & Farfán-Ríos, W. (2018). Widespread but heterogeneous

responses of Andean forests to climate change. *Nature*, *564*(7735), 207.

Feeley, K. J., Silman, M. R., Bush, M. B., Farfan, W., Cabrera, K. G., Malhi, Y., Meir, P., Revilla,

N. S., Quisiyupanqui, M. N. R., & Saatchi, S. (2011). Upslope migration of Andean

trees. *Journal of Biogeography*, *38*(4), 783–791.

https://doi.org/10.1111/j.1365-2699.2010.02444.x

Fernández, N., Navarro, L. M., & Pereira, H. M. (2017). Rewilding: A Call for Boosting

Ecological Complexity in Conservation. *Conservation Letters*, *10*(3), 276–278.

https://doi.org/10.1111/conl.12374

Filotas, E., Parrott, L., Burton, P. J., Chazdon, R. L., Coates, K. D., Coll, L., Haeussler, S.,

Martin, K., Nocentini, S., Puettmann, K. J., Putz, F. E., Simard, S. W., & Messier, C.

(2014). Viewing forests through the lens of complex systems science. *Ecosphere*, *5*(1),

art1. https://doi.org/10.1890/ES13-00182.1

Fisher, C. K., & Mehta, P. (2014). The transition between the niche and neutral regimes in

ecology. *Proceedings of the National Academy of Sciences*, *111*(36), 13111–13116.

https://doi.org/10.1073/pnas.1405637111

Foster, P. (2001). The potential negative impacts of global climate change on tropical montane

cloud forests. *Earth-Science Reviews*, *55*(1–2), 73–106.

https://doi.org/10.1016/S0012-8252(01)00056-3

Franklin, J. F., Spies, T. A., Pelt, R. V., Carey, A. B., Thornburgh, D. A., Berg, D. R.,

Lindenmayer, D. B., Harmon, M. E., Keeton, W. S., Shaw, D. C., Bible, K., & Chen, J.

(2002). Disturbances and structural development of natural forest ecosystems with

silvicultural implications, using Douglas-fir forests as an example. *Forest Ecology and

Management*, *155*(1), 399–423. https://doi.org/10.1016/S0378-1127(01)00575-8

GDAL/OGR contributors. (2021). *GDAL/OGR Geospatial Data Abstraction software Library*. Open Source Geospatial Foundation. https://gdal.org

Gentry, A. H. (1992). Tropical forest biodiversity: Distributional patterns and their conservational significance. *Oikos*, 19–28.

Gentry, A. H. (1993). *A field guide to the families and genera of woody plants of northwest South America (Colombia, Ecuador, Peru), with supplementary notes on herbaceous taxa*. The University of Chicago Press.

Gentry, A. H., & Dodson, C. H. (1987). Diversity and Biogeography of Neotropical Vascular Epiphytes. *Annals of the Missouri Botanical Garden*, *74*(2), 205. https://doi.org/10.2307/2399395

Girardin, C. A. J., Espejob, J. E. S., Doughty, C. E., Huasco, W. H., Metcalfe, D. B., Durand-Baca, L., Marthews, T. R., Aragao, L. E. O. C., Farfán-Rios, W., García-Cabrera, K., Halladay, K., Fisher, J. B., Galiano-Cabrera, D. F., Huaraca-Quispe, L. P., Alzamora-Taype, I., Eguiluz-Mora, L., Salinas -Revilla, N., Silman, M. R., Meir, P., & Malhi, Y. (2014). Productivity and carbon allocation in a tropical montane cloud forest in the Peruvian Andes. *Plant Ecology & Diversity*, *7*(1–2), 107–123. https://doi.org/10.1080/17550874.2013.820222

Gleason, H. A. (1926). The Individualistic Concept of the Plant Association. *Bulletin of the Torrey Botanical Club*, *53*(1), 7–26.

Goosem, S. P., Tucker, N. I. J., Queensland, & Wet Tropics Management Authority. (2013). *Repairing the rainforest*.

Grime, J. P. (1977). Evidence for the Existence of Three Primary Strategies in Plants and Its Relevance to Ecological and Evolutionary Theory. *The American Naturalist*, *111*(982), 1169–1194. https://doi.org/10.1086/283244

Grubb, P. J. (1977). Control of Forest Growth and Distribution on Wet Tropical Mountains: With Special Reference to Mineral Nutrition. *Annual Review of Ecology and Systematics*,

8(1), 83–107. https://doi.org/10.1146/annurev.es.08.110177.000503

Grubb, P. J., Lloyd, J. R., Pennington, T. D., & Whitmore, T. C. (1963). A Comparison of

Montane and Lowland Rain Forest in Ecuador I. The Forest Structure, Physiognomy,

and Floristics. *The Journal of Ecology*, *51*(3), 567. https://doi.org/10.2307/2257748

Grubb, P. J., & Whitmore, T. C. (1966). A Comparison of Montane and Lowland Rain Forest in

Ecuador: II. The Climate and its Effects on the Distribution and Physiognomy of the

Forests. *The Journal of Ecology*, *54*(2), 303. https://doi.org/10.2307/2257951

Guayasamin, J. M., Vandegrift, R., Policha, T., Encalada, A. C., Greene, N., Ríos-Touma, B.,

Endara, L., Cárdenas, R. E., Larreátegui, F., Baquero, L., Arcos, I., Cueva, J., Peck, M.,

Alfonso-Cortes, F., Thomas, D., DeCoux, J., Levy, E., & Roy, B. A. (2021). *Tipping Point*

Towards Biodiversity Conservation? Local and Global Consequences of the Application

of 'Rights of Nature' by Ecuador. https://doi.org/10.20944/preprints202108.0428.v1

Gundersen, P., Thybring, E. E., Nord-Larsen, T., Vesterdal, L., Nadelhoffer, K. J., & Johannsen,

V. K. (2021). Old-growth forest carbon sinks overestimated. *Nature*, *591*(7851),

E21–E23.

Hagberg, A., Swart, P., & S Chult, D. (2008). *Exploring network structure, dynamics, and*

function using networkx (LA-UR-08-05495; LA-UR-08-5495). Los Alamos National Lab.

(LANL), Los Alamos, NM (United States). https://www.osti.gov/biblio/960616

Harmon, M. E., Ferrell, W. K., & Franklin, J. F. (1990). Effects on Carbon Storage of Conversion

of Old-Growth Forests to Young Forests. *Science*, *247*(4943), 699–702.

https://doi.org/10.1126/science.247.4943.699

Hartmann, M., Howes, C. G., VanInsberghe, D., Yu, H., Bachar, D., Christen, R., Henrik

Nilsson, R., Hallam, S. J., & Mohn, W. W. (2012). Significant and persistent impact of

timber harvesting on soil microbial communities in Northern coniferous forests. *The*

ISME Journal, *6*(12), 2199–2218. https://doi.org/10.1038/ismej.2012.84

Hartmann, M., Niklaus, P. A., Zimmermann, S., Schmutz, S., Kremer, J., Abarenkov, K.,

Lüscher, P., Widmer, F., & Frey, B. (2014). Resistance and resilience of the forest soil microbiome to logging-associated compaction. *The ISME Journal*, *8*(1), 226–244. https://doi.org/10.1038/ismej.2013.141

Hautier, Y., Tilman, D., Isbell, F., Seabloom, E. W., Borer, E. T., & Reich, P. B. (2015). Anthropogenic environmental changes affect ecosystem stability via biodiversity. *Science*, *348*(6232), 336–340. https://doi.org/10.1126/science.aaa1788

Hawkins, B. A., Field, R., Cornell, H. V., Currie, D. J., Guégan, J.-F., Kaufman, D. M., Kerr, J. T., Mittelbach, G. G., Oberdorff, T., O'Brien, E. M., Porter, E. E., & Turner, J. R. G. (2003). Energy, Water, and Broad-Scale Geographic Patterns of Species Richness. *Ecology*, *84*(12), 3105–3117. https://doi.org/10.1890/03-8006

HilleRisLambers, J., Adler, P. B., Harpole, W. S., Levine, J. M., & Mayfield, M. M. (2012). Rethinking Community Assembly through the Lens of Coexistence Theory. *Annual Review of Ecology, Evolution, and Systematics*, *43*(1), 227–248. https://doi.org/10.1146/annurev-ecolsys-110411-160411

Hobbs, R. J., Higgs, E., & Harris, J. A. (2009). Novel ecosystems: Implications for conservation and restoration. *Trends in Ecology & Evolution*, *24*(11), 599–605. https://doi.org/10.1016/j.tree.2009.05.012

Holl, K. D. (1999). Factors Limiting Tropical Rain Forest Regeneration in Abandoned Pasture: Seed Rain, Seed Germination, Microclimate, and Soil1. *Biotropica*, *31*(2), 229–242. https://doi.org/10.1111/j.1744-7429.1999.tb00135.x

Holling, C. S. (1973). Resilience and Stability of Ecological Systems. *Annual Review of Ecology and Systematics*, *4*(1), 1–23. https://doi.org/10.1146/annurev.es.04.110173.000245

Howe, H. F., & Smallwood, J. (1982). Ecology of Seed Dispersal. *Annual Review of Ecology and Systematics*, *13*, 201–228.

Hubbell, S. P. (2008). Approaching ecological complexity from the perspective of symmetric neutral theory. *Tropical Forest Community Ecology*, 143–159.

Hughes, C. E. (2016). The tropical Andean plant diversity powerhouse. *New Phytologist*, *210*(4), 1152–1154. https://doi.org/10.1111/nph.13958

Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science & Engineering*, *9*(3), 90–95. https://doi.org/10.1109/MCSE.2007.55

Hunter, M. O., Keller, M., Morton, D., Cook, B., Lefsky, M., Ducey, M., Saleska, S., Jr, R. C. de O., & Schietti, J. (2015). Structural dynamics of tropical moist forest gaps. *PLOS ONE*, *10*(7), e0132144. https://doi.org/10.1371/journal.pone.0132144

Jablonski, D., Roy, K., & Valentine, J. W. (2006). Out of the Tropics: Evolutionary Dynamics of the Latitudinal Diversity Gradient. *Science*, *314*(5796), 102–106. https://doi.org/10.1126/science.1130880

Janzen, D. H. (1967). Why Mountain Passes are Higher in the Tropics. *The American Naturalist*, *101*(919), 233–249. https://doi.org/10.1086/282487

Jari, O., Blanchet, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H. (2019). *Vegan: Community ecology package* (2.5-6) [R]. https://cran.r-project.org/package=vegan

Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Tautenhahn, S., Werner, G. D. A., Aakala, T., Abedi, M., Acosta, A. T. R., Adamidis, G. C., Adamson, K., Aiba, M., Albert, C. H., Alcántara, J. M., C, C. A., Aleixo, I., Ali, H., … Wirth, C. (2020). TRY plant trait database – enhanced coverage and open access. *Global Change Biology*, *26*(1), 119–188. https://doi.org/10.1111/gcb.14904

Keddy, P. A. (1992). Assembly and response rules: Two goals for predictive community ecology. *Journal of Vegetation Science*, *3*(2), 157–164. https://doi.org/10.2307/3235676

Kocian, M., Batker, D., & Harrison-Cox, J. (2011). *An ecological study of Ecuador's Intag region: The environmental impacts and potential rewards of mining*. Earth Economics.

Kraft, N. J. B., Valencia, R., & Ackerly, D. D. (2008). Functional Traits and Niche-Based Tree Community Assembly in an Amazonian Forest. *Science*, *322*(5901), 580–582.

https://doi.org/10.1126/science.1160662

Kumar, R., Carroll, C., Hartikainen, A., & Martin, O. (2019). ArviZ a unified library for

exploratory analysis of Bayesian models in Python. *Journal of Open Source Software*,

4(33), 1143. https://doi.org/10.21105/joss.01143

Küper, W., Kreft, H., Nieder, J., Köster, N., & Barthlott, W. (2004). Large-scale diversity patterns

of vascular epiphytes in Neotropical montane rain forests. *Journal of Biogeography*,

31(9), 1477–1487. https://doi.org/10.1111/j.1365-2699.2004.01093.x

Lal, R. (2005). Forest soils and carbon sequestration. *Forest Ecology and Management*,

220(1), 242–258. https://doi.org/10.1016/j.foreco.2005.08.015

Laughlin, D. C., & Laughlin, D. E. (2013). Advances in modeling trait-based plant community

assembly. *Trends in Plant Science*, *18*(10), 584–593.

https://doi.org/10.1016/j.tplants.2013.04.012

Ledo, A., Cayuela, L., Manso, R., & Condés, S. (2015). Recruitment patterns and potential

mechanisms of community assembly in an Andean cloud forest. *Journal of Vegetation

Science*, *26*(5), 876–888. https://doi.org/10.1111/jvs.12287

Legendre, P., & Legendre, L. (2012). *Numerical Ecology* (Third, Vol. 24). Elsevier.

Leigh, E. G. (1975). Structure and Climate in Tropical Rain Forest. *Annual Review of Ecology

and Systematics*, *6*(1), 67–86. https://doi.org/10.1146/annurev.es.06.110175.000435

Lewis, S. L., Wheeler, C. E., Mitchard, E. T. A., & Koch, A. (2019). Restoring natural forests is

the best way to remove atmospheric carbon. *Nature*, *568*(7750), 25–28.

https://doi.org/10.1038/d41586-019-01026-8

Lippi, R. D. (1988). Paleotopography and Phosphate Analysis of a Buried Jungle Site in

Ecuador. *Journal of Field Archaeology*, *15*(1), 85–97.

https://doi.org/10.1179/009346988791974556

Lippi, R., & Gudiño, A. (2010). Inkas and Yumbos at Palmitopamba in northwestern Ecuador.

Distant Provinces in the Inka Empire. Iowa City: University of Iowa Press. p, 260–278.

Loehle, C. (2004). Challenges of ecological complexity. *Ecological Complexity*, *1*(1), 3–6.

Loughlin, N. J. D., Gosling, W. D., Mothes, P., & Montoya, E. (2018). Ecological consequences of post-Columbian indigenous depopulation in the Andean–Amazonian corridor. *Nature Ecology & Evolution*, *2*(8), 1233–1236. https://doi.org/10.1038/s41559-018-0602-7

Luyssaert, S., Schulze, E.-D., Börner, A., Knohl, A., Hessenmöller, D., Law, B. E., Ciais, P., & Grace, J. (2008). Old-growth forests as global carbon sinks. *Nature*, *455*(7210), 213–215. https://doi.org/10.1038/nature07276

MacArthur, R. H., & Wilson, E. O. (2001). *The theory of island biogeography* (Vol. 1). Princeton university press.

Mantgem, P. J. van, Stephenson, N. L., Byrne, J. C., Daniels, L. D., Franklin, J. F., Fulé, P. Z., Harmon, M. E., Larson, A. J., Smith, J. M., Taylor, A. H., & Veblen, T. T. (2009). Widespread Increase of Tree Mortality Rates in the Western United States. *Science*, *323*(5913), 521–524. https://doi.org/10.1126/science.1165000

McElhinny, C., Gibbons, P., Brack, C., & Bauhus, J. (2005). Forest and woodland stand structural complexity: Its definition and measurement. *Forest Ecology and Management*, *218*(1), 1–24. https://doi.org/10.1016/j.foreco.2005.08.034

McKinney, W. (2010). *Data structures for statistical computing in Python*. 51–56. http://conference.scipy.org/proceedings/scipy2010/mckinney.html

Messier, C., Puettmann, K., Chazdon, R., Andersson, K. P., Angers, V. A., Brotons, L., Filotas, E., Tittler, R., Parrott, L., & Levin, S. A. (2015). From Management to Stewardship: Viewing Forests As Complex Adaptive Systems in an Uncertain World. *Conservation Letters*, *8*(5), 368–377. https://doi.org/10.1111/conl.12156

Michaels, T. K., Eppinga, M. B., & Bever, J. D. (2020). A nucleation framework for transition between alternate states: Short-circuiting barriers to ecosystem recovery. *Ecology*, *101*(9), e03099. https://doi.org/10.1002/ecy.3099

Mitchard, E. T. A. (2018). The tropical forest carbon cycle and climate change. *Nature*,

559(7715), 527–534. https://doi.org/10.1038/s41586-018-0300-2

Moomaw, W. R., Masino, S. A., & Faison, E. K. (2019). Intact Forests in the United States:

Proforestation Mitigates Climate Change and Serves the Greatest Good. *Frontiers in*

Forests and Global Change, *2*. https://doi.org/10.3389/ffgc.2019.00027

Morante‑Filho, J. C., Arroyo‑Rodríguez, V., Pessoa, M. de S., Cazetta, E., & Faria, D. (2018).

Direct and cascading effects of landscape structure on tropical forest and non-forest

frugivorous birds. *Ecological Applications*, *28*(8), 2024–2032.

https://doi.org/10.1002/eap.1791

Morelos-Juárez, C., Tapia, A., Conde, G., & Peck, M. (2015). *Diet of the critically endangered*

brown-headed spider monkey (Ateles fusciceps fusciceps) in the Ecuadorian Chocó:

Conflict between primates and loggers over fruiting tree species (No. e1963). PeerJ Inc.

https://doi.org/10.7287/peerj.preprints.1574v1

Mosandl, R., Günter, S., Stimm, B., & Weber, M. (2008). Ecuador Suffers the Highest

Deforestation Rate in South America. In E. Beck, J. Bendix, I. Kottke, F. Makeschin, &

R. Mosandl (Eds.), *Gradients in a Tropical Mountain Ecosystem of Ecuador* (pp. 37–40).

Springer. https://doi.org/10.1007/978-3-540-73526-7_4

Myers, N. (1988). Threatened biotas:" hot spots" in tropical forests. *Environmentalist*, *8*(3),

187–208.

Myers, N., Mittermeier, R. A., Mittermeier, C. G., Fonseca, G. A. B. da, & Kent, J. (2000).

Biodiversity hotspots for conservation priorities. *Nature*, *403*(6772), 853–858.

https://doi.org/10.1038/35002501

Navarro, A. B., Bovo, A. a. A., Alexandrino, E. R., Oliveira, V. C., Pizo, M. A., & Ferraz, K. M. P.

M. B. (2018). Fruit availability at the individual and local levels influences fruit removal

in *Cecropia pachystachya*. *Brazilian Journal of Biology*, *79*, 758–759.

https://doi.org/10.1590/1519-6984.198339

Nicholas Brokaw. (1982). The Definition of treefall gap and Its effect on measures of forest

dynamics. *Biotropica*, *14*(2), 158–160. JSTOR. https://doi.org/10.2307/2387750

Norden, N., Angarita, H. A., Bongers, F., Martínez-Ramos, M., Cerda, I. G. la, Breugel, M. van,
Lebrija-Trejos, E., Meave, J. A., Vandermeer, J., Williamson, G. B., Finegan, B.,
Mesquita, R., & Chazdon, R. L. (2015). Successional dynamics in Neotropical forests
are as uncertain as they are predictable. *Proceedings of the National Academy of
Sciences*, *112*(26), 8013–8018. https://doi.org/10.1073/pnas.1500403112

Norden, N., Chazdon, R. L., Chao, A., Jiang, Y.-H., & Vílchez‐Alvarado, B. (2009). Resilience
of tropical rain forests: Tree community reassembly in secondary forests. *Ecology
Letters*, *12*(5), 385–394. https://doi.org/10.1111/j.1461-0248.2009.01292.x

Odum, E. P. (2014). The strategy of ecosystem development. In *The ecological design and
planning reader* (pp. 203–216). Springer.

O'Hagan, A., & Leonard, T. (1976). Bayes Estimation Subject to Uncertainty About Parameter
Constraints. *Biometrika*, *63*(1), 201–203. https://doi.org/10.2307/2335105

Olson, D. M. (1994). The Distribution of Leaf Litter Invertebrates Along a Neotropical Altitudinal
Gradient. *Journal of Tropical Ecology*, *10*(2), 129–150.

Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L.,
Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W.,
McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S., & Hayes, D. (2011). A Large and
Persistent Carbon Sink in the World's Forests. *Science*, *333*(6045), 988–993.
https://doi.org/10.1126/science.1201609

Parr, C. S., Wilson, N., Leary, P., Schulz, K., Lans, K., Walley, L., Hammock, J., Goddard, A.,
Rice, J., Studer, M., Holmes, J., Corrigan, R., & Jr. (2014). The Encyclopedia of Life v2:
Providing Global Access to Knowledge About Life on Earth. *Biodiversity Data Journal*,
2, e1079. https://doi.org/10.3897/BDJ.2.e1079

Peck, M., Thorn, J., Mariscal, A., Baird, A., Tirira, D., & Kniveton, D. (2011). Focusing
conservation efforts for the critically endangered brown-headed spider monkey (*Ateles*

fusciceps) using remote sensing, modeling, and playback survey methods. *International Journal of Primatology*, *32*(1), 134–148.

Peres, C. A., Emilio, T., Schietti, J., Desmoulière, S. J. M., & Levi, T. (2016). Dispersal limitation induces long-term biomass collapse in overhunted Amazonian forests. *Proceedings of the National Academy of Sciences*, *113*(4), 892–897. https://doi.org/10.1073/pnas.1516525113

Persson, C., Eriksson, R., Ståhl, B., & Romoleroux, K. (Eds.). (n.d.). *Flora of Ecuador | University of Gothenburg* (Vol. 1–97). University of Gothenburg. Retrieved August 19, 2021, from https://www.gu.se/en/biological-environmental-sciences/flora-of-ecuador

Pickett, S. T. A., Cadenasso, M. L., & Meiners, S. J. (2009). Ever since Clements: From succession to vegetation dynamics and understanding to intervention [*]. *Applied Vegetation Science*, *12*(1), 9–21. https://doi.org/10.1111/j.1654-109X.2009.01019.x

Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N., Raven, P. H., Roberts, C. M., & Sexton, J. O. (2014). The biodiversity of species and their rates of extinction, distribution, and protection. *Science*, *344*(6187). https://doi.org/10.1126/science.1246752

Planchuelo, G., Kowarik, I., & von der Lippe, M. (2020). Endangered Plants in Novel Urban Ecosystems Are Filtered by Strategy Type and Dispersal Syndrome, Not by Spatial Dependence on Natural Remnants. *Frontiers in Ecology and Evolution*, *8*, 18. https://doi.org/10.3389/fevo.2020.00018

Policha, T. (2014). *Pollination biology of the mushroom-mimicking orchid genus Dracula* [University of Oregon]. http://hdl.handle.net/1794/18404

Python Core Team. (2018). *Python Software Foundation*. Python.Org. https://www.python.org/

R Core Team. (2020). *R: A language and environment for statistical computing* [R]. http://www.R-project.org/

Ramírez-Barahona, S., Luna-Vega, I., & Tejero-Díez, D. (2011). Species richness, endemism,

and conservation of American tree ferns (Cyatheales). *Biodiversity and Conservation*, *20*(1), 59–72. https://doi.org/10.1007/s10531-010-9946-2

Reback, J., McKinney, W., jbrockmendel, Bossche, J. V. den, Augspurger, T., Cloud, P., gfyoung, Sinhrks, Hawkins, S., Klein, A., Roeschke, M., Tratner, J., Petersen, T., She, C., Ayd, W., MomIsBestFriend, Garcia, M., Schendel, J., Hayden, A., … Winkel, M. (2020). *Pandas 1.1.3* [Python]. Zenodo. https://doi.org/10.5281/zenodo.4067057

Rehm, E. M., & Feeley, K. J. (2015). The inability of tropical cloud forest species to invade grasslands above treeline during climate change: Potential explanations and consequences. *Ecography*, *38*(12), 1167–1175. https://doi.org/10.1111/ecog.01050

Richards, P. W. (1952). The tropical rain forest. *The Tropical Rain Forest.*

Ríos-Touma, B., Holzenthal, R. W., Huisman, J., Thomson, R., & Rázuri-Gonzales, E. (2017). Diversity and distribution of the Caddisflies (Insecta: Trichoptera) of Ecuador. *PeerJ*, *5*, e2851. https://doi.org/10.7717/peerj.2851

Ripple, W. J., Estes, J. A., Schmitz, O. J., Constant, V., Kaylor, M. J., Lenz, A., Motley, J. L., Self, K. E., Taylor, D. S., & Wolf, C. (2016). What is a Trophic Cascade? *Trends in Ecology & Evolution*, *31*(11), 842–849. https://doi.org/10.1016/j.tree.2016.08.010

Ross, K. A., Fox, B. J., & Fox, M. D. (2002). Changes to plant species richness in forest fragments: Fragment age, disturbance and fire history may be as important as area. *Journal of Biogeography*, *29*(5–6), 749–765. https://doi.org/10.1046/j.1365-2699.2002.00722.x

Roy, B. A., Zorrilla, M., Endara, L., Thomas, D. C., Vandegrift, R., Rubenstein, J. M., Policha, T., Rios-Touma, B., & Read, M. (2018). New mining concessions could severely decrease biodiversity and ecosystem services in ecuador. *Tropical Conservation Science*, *11*, 1940082918780427. https://doi.org/10.1177/1940082918780427

Sacco, A. D., Hardwick, K. A., Blakesley, D., Brancalion, P. H. S., Breman, E., Rebola, L. C., Chomba, S., Dixon, K., Elliott, S., Ruyonga, G., Shaw, K., Smith, P., Smith, R. J., &

Antonelli, A. (2021). Ten golden rules for reforestation to optimize carbon sequestration, biodiversity recovery and livelihood benefits. *Global Change Biology*, *27*(7), 1328–1348. https://doi.org/10.1111/gcb.15498

Salvatier, J., Wiecki, T. V., & Fonnesbeck, C. (2016). Probabilistic programming in Python using PyMC3. *PeerJ Computer Science*, *2*, e55. https://doi.org/10.7717/peerj-cs.55

Särkinen, T., Pennington, R. T., Lavin, M., Simon, M. F., & Hughes, C. E. (2012). Evolutionary islands in the Andes: Persistence and isolation explain high endemism in Andean dry tropical forests. *Journal of Biogeography*, *39*(5), 884–900.

Seidl, R., Thom, D., Kautz, M., Martin-Benito, D., Peltoniemi, M., Vacchiano, G., Wild, J., Ascoli, D., Petr, M., Honkaniemi, J., Lexer, M. J., Trotsiuk, V., Mairota, P., Svoboda, M., Fabrika, M., Nagel, T. A., & Reyer, C. P. O. (2017). Forest disturbances under climate change. *Nature Climate Change*, *7*(6), 395–402. https://doi.org/10.1038/nclimate3303

Skutch, A. F. (1945). The Most Hospitable Tree. *The Scientific Monthly*, *60*(1), 5–17.

Smith, S. A., Oca, A. N. M. D., Reeder, T. W., & Wiens, J. J. (2007). A Phylogenetic Perspective on Elevational Species Richness Patterns in Middle American Treefrogs: Why so Few Species in Lowland Tropical Rainforests? *Evolution*, *61*(5), 1188–1207. https://doi.org/10.1111/j.1558-5646.2007.00085.x

Sohng, J., Singhakumara, B. M. P., & Ashton, M. S. (2017). Effects on soil chemistry of tropical deforestation for agriculture and subsequent reforestation with special reference to changes in carbon and nitrogen. *Forest Ecology and Management*, *389*, 331–340. https://doi.org/10.1016/j.foreco.2016.12.013

Spracklen, D. V., & Righelato, R. (2014). Tropical montane forests are a larger than expected global carbon store. *Biogeosciences*, *11*(10), 2741–2754. https://doi.org/10.5194/bg-11-2741-2014

Stadtmüller, T. (1987). *Cloud forests in the humid tropics: A bibliographic review*. United Nations University Press.

https://archive.unu.edu/unupress/unupbooks/80670e/80670E00.htm

Stephenson, N. L., Das, A. J., Condit, R., Russo, S. E., Baker, P. J., Beckman, N. G., Coomes,
D. A., Lines, E. R., Morris, W. K., Rüger, N., Álvarez, E., Blundo, C., Bunyavejchewin,
S., Chuyong, G., Davies, S. J., Duque, Á., Ewango, C. N., Flores, O., Franklin, J. F., …
Zavala, M. A. (2014). Rate of tree carbon accumulation increases continuously with tree
size. *Nature*, *507*(7490), 90–93. https://doi.org/10.1038/nature12914

Stork, N. E. (2010). Re-assessing current extinction rates. *Biodiversity and Conservation*,
19(2), 357–371. https://doi.org/10.1007/s10531-009-9761-9

Tachikawa, T., Kaku, M., Iwasaki, A., Gesch, D. B., Oimoen, M. J., Zhang, Z., Danielson, J. J.,
Krieger, T., Curtis, B., Haase, J., Abrams, M., & Carabajal, C. (2011). *ASTER Global
Digital Elevation Model Version 2—Summary of validation results* (2nd ed., p. 27)
[Report]. USGS Publications Warehouse. http://pubs.er.usgs.gov/publication/70005960

Tapia-Armijos, M. F., Homeier, J., Espinosa, C. I., Leuschner, C., & Cruz, M. de la. (2015).
Deforestation and Forest Fragmentation in South Ecuador since the 1970s – Losing a
Hotspot of Biodiversity. *PLOS ONE*, *10*(9), e0133701.
https://doi.org/10.1371/journal.pone.0133701

Tobler, W. R. (1970). A Computer Movie Simulating Urban Growth in the Detroit Region.
Economic Geography, *46*(sup1), 234–240. https://doi.org/10.2307/143141

Tovar, C., Arnillas, C. A., Cuesta, F., & Buytaert, W. (2013). Diverging Responses of Tropical
Andean Biomes under Future Climate Conditions. *PLOS ONE*, *8*(5), e63634.
https://doi.org/10.1371/journal.pone.0063634

Ulrich, W., Zaplata, M. K., Winter, S., Schaaf, W., Fischer, A., Soliveres, S., & Gotelli, N. J.
(2016). Species interactions and random dispersal rather than habitat filtering drive
community assembly during early plant succession. *Oikos*, *125*(5), 698–707.
https://doi.org/10.1111/oik.02658

Valencia, R., Balslev, H., & Miño, G. P. Y. (1994). High tree alpha-diversity in Amazonian

Ecuador. *Biodiversity & Conservation*, *3*(1), 21–28.

Vallejo Galárraga, M. C., & Freslon, W. S. (2017). Ecuador: Mineral Policy. In G. Tiess, T.

Majumder, & P. Cameron (Eds.), *Encyclopedia of Mineral and Energy Policy* (pp. 1–8).

Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-40871-7_153-1

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,

Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M.,

Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E.,

… van Mulbregt, P. (2020). SciPy 1.0: Fundamental algorithms for scientific computing

in Python. *Nature Methods*, *17*(3), 261–272. https://doi.org/10.1038/s41592-019-0686-2

Vitousek, P. M., Mooney, H. A., Lubchenco, J., & Melillo, J. M. (1997). Human Domination of

Earth's Ecosystems. *Science*, *277*(5325), 494–499.

https://doi.org/10.1126/science.277.5325.494

Von Humboldt, A., & Bonpland, A. (2010). *Essay on the Geography of Plants*. University of

Chicago Press.

Wang, Q., He, N., Xu, L., & Zhou, X. (2018). Microbial properties regulate spatial variation in

the differences in heterotrophic respiration and its temperature sensitivity between

primary and secondary forests from tropical to cold-temperate zones. *Agricultural and

Forest Meteorology*, *262*, 81–88. https://doi.org/10.1016/j.agrformet.2018.07.007

Watling, J., Iriarte, J., Mayle, F. E., Schaan, D., Pessenda, L. C. R., Loader, N. J.,

Street-Perrott, F. A., Dickau, R. E., Damasceno, A., & Ranzi, A. (2017). Impact of

pre-Columbian "geoglyph" builders on Amazonian forests. *Proceedings of the National

Academy of Sciences*, *114*(8), 1868–1873. https://doi.org/10.1073/pnas.1614359114

Watson, J. E. M., Evans, T., Venter, O., Williams, B., Tulloch, A., Stewart, C., Thompson, I.,

Ray, J. C., Murray, K., Salazar, A., McAlpine, C., Potapov, P., Walston, J., Robinson, J.

G., Painter, M., Wilkie, D., Filardi, C., Laurance, W. F., Houghton, R. A., …

Lindenmayer, D. (2018). The exceptional value of intact forest ecosystems. *Nature*

Ecology & Evolution, *2*(4), 599–610. https://doi.org/10.1038/s41559-018-0490-x

Whittaker, R. H. (1956). Vegetation of the Great Smoky Mountains. *Ecological Monographs*,

    26(1), 1–80. https://doi.org/10.2307/1943577

Willson, M. F., & Crome, F. H. J. (1989). Patterns of seed rain at the edge of a tropical

    Queensland rain forest. *Journal of Tropical Ecology*, *5*(3), 301–308.

    https://doi.org/10.1017/S0266467400003680

Wunder, S. (2000). *Economics of Deforestation: The Example of Ecuador*. Palgrave Macmillan

    UK. https://doi.org/10.1057/9780230596696

Young, K. R., Ewel, J. J., & Brown, B. J. (1987). Seed dynamics during forest succession in

    Costa Rica. *Vegetatio*, *71*(3), 157–173. https://doi.org/10.1007/BF00039168

Young, K. R., Ulloa, C. U., Luteyn, J. L., & Knapp, S. (2002). Plant evolution and endemism in

    Andean South America: An introduction. *The Botanical Review*, *68*(1), 4–21.

Young, T. P., Chase, J. M., & Huddleston, R. T. (2001). Community Succession and Assembly

    Comparing, Contrasting and Combining Paradigms in the Context of Ecological

    Restoration. *Ecological Restoration*, *19*(1), 5–18. https://doi.org/10.3368/er.19.1.5

Zahawi, R. A., Holl, K. D., Cole, R. J., & Reid, J. L. (2013). Testing applied nucleation as a

    strategy to facilitate tropical forest recovery. *Journal of Applied Ecology*, *50*(1), 88–96.

    https://doi.org/10.1111/1365-2664.12014

Zhang, H., Deng, Q., Hui, D., Wu, J., Xiong, X., Zhao, J., Zhao, M., Chu, G., Zhou, G., &

    Zhang, D. (2019). Recovery in soil carbon stock but reduction in carbon stabilization

    after 56-year forest restoration in degraded tropical lands. *Forest Ecology and*

    Management, *441*, 1–8. https://doi.org/10.1016/j.foreco.2019.03.037