Title: Intralocus sexual conflict can maintain alternative reproductive tactics

Running Title: Sexual conflict maintains ARTs

Authors: Madilyn M. Gamble[1] and Ryan G. Calsbeek[2]

[1]Ecology, Evolution, Ecosystems, and Society Program, Dartmouth College, 78 College Street,

Hanover, NH 03755, USA. Email: mmg.gr@dartmouth.edu. ORCID ID: https://orcid.org/0000-

0003-4791-059X

[2]Ecology, Evolution, Ecosystems, and Society Program, Dartmouth College, 78 College Street,

Hanover, NH 03755, USA. Email: ryan.g.calsbeek@dartmouth.edu. ORCID ID:

https://orcid.org/0000-0002-9236-4092

Keywords: intralocus conflict, alternative reproductive tactics, selection, heritability

Abstract

Alternative reproductive tactics (ARTs) are ubiquitous throughout the animal kingdom and

widely regarded as an outcome of high variance in reproductive success. Proximate mechanisms

underlying ARTs include genetically based polymorphisms, environmentally induced

polymorphisms, and those mediated by a combination of genetic and environmental factors.

However, few ultimate mechanisms have been proposed to explain the maintenance of ARTs

over time, the most important of which have been disruptive and negative frequency-dependent

selection. Here we explore the role that intralocus sexual conflict may play in the maintenance of

sex-specific ARTs. We use a genetically explicit individual-based model in which body size

influences both female fecundity and male tactic through a shared genetic architecture. By

modeling ART maintenance under varying selection regimes and levels of sex-specific gene

expression, we explore the conditions under which intralocus sexual conflict can maintain a

hypothetical ART defined by larger (alpha) and smaller (beta) tactics. Our models consistently

revealed that sexual conflict can result in the persistence of a sex-specific polymorphism over

hundreds of generations, even in the absence of negative frequency-dependent selection. ARTs

were maintained through correlated selection when one male ART has lower fitness but produces

daughters with higher fitness. These results highlight the importance of understanding selection

on both sexes when attempting to explain the maintenance of ARTs. Our results are consistent

with a growing literature documenting genetic correlations between male ARTs and female

fitness, suggesting that the maintenance of sex-specific ARTs through intralocus sexual conflict

may be common and widespread in nature.

Introduction

Alternative reproductive tactics (ARTs) are ubiquitous throughout the animal kingdom and

are widely regarded as an outcome of high variance in reproductive success (Shuster and Wade

2003). Proximate mechanisms underlying ARTs include purely genetically based polymorphisms

(Shuster 1989; Cordero 1990; Lank et al. 1995; Sinervo and Lively 1996; Sinervo et al. 2000b),

purely environmentally induced polymorphisms (Thornhill 1981; Emlen 1994; Hunt and

Simmons 1997; Moczek and Emlen 1999), as well as those mediated by a combination of genetic

and environmental factors (Neff and Svensson 2013). However, few ultimate mechanisms have

been proposed to explain the evolutionary maintenance of ARTs over time. The most prominent

of these mechanisms are disruptive and negative frequency-dependent forms of selection

(Oliviera et al. 2008; Hughes et al. 2013), both of which have been documented in theoretical,

laboratory, and field studies (Gross 1985, 1991, 1996; Oliviera et al. 2008). These forms of

selection are particularly important in the maintenance of polymorphism because negative

frequency dependence and disruptive selection are themselves variable forms of selection, the

former varying over space and time, the latter varying over the statistical distribution of

phenotypes.

Variable selection can also occur within a population when a trait that is expressed in both

males and females has alternative sex-specific fitness optima, resulting in sexually antagonistic

selection on that trait (Lande 1980). If that trait also has a common genetic basis in males and

females, then sexually antagonistic selection will lead to intralocus sexual conflict

(Bonduriansky and Chenoweth 2009). While others have proposed that intralocus sexual conflict

may maintain additive genetic variance (Rice and Chippindale 2001; Foerster et al. 2007) and

genetic polymorphism (Connallon and Clark 2012), result in balancing selection (Connallon and

Clark 2014), and promote speciation (Bonduriansky 2011), the potential for intralocus sexual

conflict to maintain alternative reproductive tactics is comparatively understudied. There is

nevertheless reason to think that intralocus sexual conflict could be a potent force that maintains

variation in nature. Indeed, a growing body of evidence shows that selection on male ARTs and

the traits that differentiate them results in correlated evolution in females: genetic linkage

between male ARTs and female reproductive traits has been described in lizards (Sinervo and

Zamudio 2001), crickets (Heinen-Kay et al. 2020a; Richardson et al. 2021), and bulb mites

(Bielak et al. 2014; Buzatto et al. 2018; Buzatto and Clark 2020; Łukasiewicz et al. 2020).

While analytical models describe the action of selection on quantitative traits (Otto and Day

2007), correlated traits (Lande and Arnold 1983), and threshold traits including ARTs (Gross and

Repka 1998), we are unaware of any theory that simultaneously investigates evolution under

both sexually antagonistic selection on continuous quantitative traits and variable intersexual

genetic correlations. Here we take new steps towards understanding the role that intralocus

sexual conflict may play in the maintenance of sex-specific ARTs. We use a genetically explicit

individual-based model in which body size influences both female fecundity and male tactic

through a shared genetic architecture. By modeling ART maintenance under varying selection

regimes and levels of sex-specific gene expression, we explore the conditions under which

intralocus sexual conflict can maintain a hypothetical ART defined by larger (alpha) and smaller

(beta) tactics.

To evaluate the potential for sexually antagonistic selection and intralocus sexual conflict to

maintain ARTs, we allowed a theoretical population to evolve under a range of scenarios

representing three different selection regimes acting on males: no selection, directional selection

against alpha males, and negative frequency-dependent selection. Within each selection regime

we also varied the degree to which males and females could evolve independently by varying the

number of genetic loci with sex-dependent expression. We used these models to test our

hypothesis that intralocus sexual conflict may influence ART frequency over time, and that this

process might be altered by different selection regimes and/or the breakdown of intersexual

genetic correlations through sex-specific gene expression. Specifically, we predicted ARTs could

be maintained: (1) when alpha and beta males have equal probabilities of survival and mating but

intersexual heritability is low such that larger females do not necessarily produce larger (alpha)

sons, and (2) when alpha males are selected against but intersexual heritability of body size is

high such that low-fitness alpha males produce high-fitness daughters. While it is often assumed

that a male tactic that is larger, more sexually dimorphic, or that has sexually selected weapons

should have higher fitness, this need not necessarily be the case. Sexually-selected weapons

developed in fighter males are often costly to produce, requiring higher energy investment and

delayed development, both of which may contribute to higher mortality risk (Hunt and Simmons

1997). Fighter males often have higher testosterone levels than other male tactics, which can lead

to decreased survival (Sinervo et al. 2000a). In other cases, smaller males may have higher

fitness if shorter times to reach sexual maturity reduce their probability of mortality prior to

reproduction (Gross 1996).

Methods

<u>Overview</u>

In modeling the frequency of two different male ARTs – a larger "alpha" and smaller "beta"

male – we assumed that ART was a quantitative threshold trait determined by body size, which

in turn was entirely genetically controlled by 20 quantitative loci. Individual males with a body

size greater than or equal to a threshold value became alpha males, and those below the threshold

became beta males. Females, which did not express these ARTs, had a fecundity value

proportional to their body size. For simplicity we assumed a discrete population of semelparous

individuals with non-overlapping generations. We did not include linkage disequilibrium or

mutations in our model; all genetic variance stemmed from an initial random draw of allele

values in the first generation.

To test our hypothesis that intralocus sexual conflict can affect the maintenance of ARTs, we

varied both the form of selection acting on the male ARTs and the intersexual heritability for

body size. We varied the form of selection on males by changing the relative fitness of each

tactic. We exposed the population to a null selection regime in which both male ARTs were

equally likely to survive and mate. We simulated sexually antagonistic selection by assigning

lower fitness to alpha males but higher fitness (fecundity) to larger females. Finally, we explored

the role of negative frequency-dependent selection in which the rare male ART in each

generation had a fitness advantage.

We altered the intersexual heritability of body size by varying the number of body size loci

that were "general" or "sex-specific" in their expression. When all 20 loci were "general," the

effect of each allele controlling body size was the same in both sexes. This made size equally

heritable within and between the sexes, so larger (alpha) males produced larger (more fecund)

daughters than beta males. We predicted that correlated selection acting on males because of

fecundity selection might allow alpha males to persist in the population, even when directional

selection acts against them. However, as more loci become "sex-specific" – that is, allelic

expression depended on sex and genetic architecture for size was less shared between the sexes –

intersexual heritability should decline, and sire tactic should no longer influences daughter

fitness. Thus, alpha males should no longer produce fitter daughters, and we predicted that

consistent directional selection against alpha males could result in their decline or disappearance

from the population.

<u>The Model</u>

We initiated each simulation with a parent generation comprised of 1000 diploid sexually

reproducing individuals (50:50 sex ratio). Each individual in our model had two traits that made

up its phenotype: sex and size. Sex was assigned randomly, and size was a quantitative trait

determined by the sum of allele values measured across 20 loci. To allow for sex-specific allelic

effects, each allele had two possible values (*f* and *m*) that influence body size. That is, each locus

had a total of three possible values that contribute to body size, and which value was expressed

depends on whether the locus was sex-specific or general and if the former, whether the

individual was male or female. For each sex-specific locus, only the two allele values

corresponding to that individual's assigned sex contributed to the body size value. This is the

biological equivalent of an allele that has sex-specific expression (Rice and Chippindale 2001).

For general loci the value of each allele at that locus was determined by averaging the allele's

two possible values. Allele values for the parent generation were drawn randomly from a normal

distribution with mean = 0 and SD = 0.5. We calculated the size of the individual from the

genotype using the following sex-specific equations:

146
$$Female\ Size = \sum_{i=1}^{20-n} \frac{(f_{iD} + m_{iD})}{2} + \frac{(f_{iS} + m_{iS})}{2} + \sum_{i=20-n}^{20} f_{iD} + f_{iS} + C$$

147
$$Male\ Size = \sum_{i=1}^{20-n} \frac{(f_{iD} + m_{iD})}{2} + \frac{(f_{iS} + m_{iS})}{2} + \sum_{i=20-n}^{20} m_{iD} + m_{iS} + C$$

where n was the number of sex-specific loci, f and m are the female- and male-specific values,

respectively, of the two alleles at the ith locus inherited from the dam (D subscript) or the sire (S

subscript), and C is a constant. The first summation calculated the component of the phenotypic

value contributed from the general loci; the second summation calculated the contribution of the

sex-specific loci. Adding a constant to the expression ensured that size remained a positive

number with a mean of C in the parent generation; any individual whose calculated size was less

than zero was assigned a size of zero. We arbitrarily set $C = 10$ for our simulations.

Male mating tactic was determined by threshold body size: males with size greater than or

equal to 10 were alpha males and those with size less than 10 were beta males. Fecundity was

assigned to females as a function of size:

$$Fecundity = \frac{12 \times size^2}{size^2 + 100}$$

such that fecundity increased asymptotically with increasing body size to a maximum value

approaching 12 offspring at a size of 40. This set an upper limit on fecundity selection while

preserving positive directional selection on female size.

We modeled the evolution of male and female body size and ART frequency under seven

different selection regimes on male body size and five different values for the number of sex-

specific loci for a total of 35 different model scenarios (Table 1). These varied in the degree to

which selection on size was sexually antagonistic and in the intersexual heritability of size, thus

creating a spectrum of intralocus sexual conflict. We varied direct selection on males by

changing the relative combined probability of survival and mating (hereafter W_T for each tactic).

In the "no selection" model individuals of both ARTs had equal values of W. In three separate

"directional selection" models we set $W_a = 1$ and $W_b = 1.5$, 1.8, or 2. This allowed us to evaluate

how the strength of direct selection on male tactic affected ART maintenance in the face of

intralocus sexual conflict. In each directional selection model, W_a and W_b were held constant

across generations. In three separate "negative frequency-dependent selection" models, each

tactic's W varied as function of its frequency (F_T) such that the rare tactic in each generation

always had a higher mating probability (1- F_T). We ran three different negative frequency-

dependent models which ranged in starting frequency ratios of alpha and beta males from *Fa:Fb*

= 2:1, 1:1, and 1:2. This allowed us to evaluate how initial tactic frequencies affected ART

maintenance under negative frequency-dependent selection.

Table 1. Parameters and values modeled, resulting in 35 total scenarios modeled. Each scenario

varied in the form of selection on males and the number of sex-specific loci. Negative frequency

dependent models also varied in the starting frequency ratio of alpha to beta males (*Fa:Fb*);

directional selection models also varied in the ratio of alpha fitness to beta fitness ($W_a:W_b$).

Parameter	Values modeled
Selection Regime	No Selection, $W_a = W_b$
	Negative frequency dependent selection, $F_a:F_b = 1:1$
	Negative frequency dependent selection, $F_a:F_b = 1:2$
	Negative frequency dependent selection, $F_a:F_b = 2:1$
	Directional selection against alphas, $W_a:W_b = 1:1.5$
	Directional selection against alphas $W_a:W_b = 1:1.8$
	Directional selection against alphas $W_a:W_b = 1:2$
# Sex-Specific Loci	0, 5, 10, 15, 20

In all models, W_f did not differ among individual females, but the proportion of females that

survived to mate in each generation declined with population size to simulate density-dependent

population growth regulation. The proportion of females surviving to mate (M_f) was calculated

anew for each generation using the equation:

$$M_f = \frac{0.5}{1 + e^{-2+KN}}$$

where K is a constant and N is the current population size. We chose $K = 0.0015$ such that this

equation defined a sigmoidal curve that decreased from 0.44 at $N = 0$ to 0.04 at $N = 3000$. For

each generation the number of females that were allowed to reproduce (D) was calculated as:

$$D = M_f N$$

D females were randomly chosen to be dams. D males were similarly chosen to be sires but

sire selection was weighted by W_T. Dams were randomly paired with sires (no assortative

mating), and a number of offspring equivalent to the dam's fecundity were created. Each

offspring inherited alleles at all 20 loci in a Mendelian fashion such that at each locus the

offspring inherits one allele from the dam and one from the sire, and the allele inherited from

each parent is randomly drawn from the two alleles each parent possesses. Sex was assigned

randomly to offspring to achieve an approximately equal sex ratio, and size, fecundity, and

mating probability were calculated for each offspring as described above for individuals in the

parent generation. These offspring were then used as parents for the next generation, and the

model was iterated for a total of 500 generations in each model scenario. We ran 11 replicates of

the full set of the 35 model scenarios to ensure that results did not qualitatively differ among

model runs.

<u>Analysis</u>

In each of the 35 model scenarios, for each generation we calculated (1) intra- and

intersexual heritability of size (to understand the degree to which males and females could

respond independently to selection), (2) the sex-specific mean and variance of size (to

understand the degree of sexual dimorphism that evolved and the opportunity for selection), (3)

sex-specific selection gradients for size (to understand the degree to which selection was

sexually antagonistic), and (4) the frequency of ARTs (to understand how intralocus sexual

conflict affects ART maintenance). Intra- and intersexual heritabilities were calculated by

doubling the regression coefficient of the sex-specific mean offspring size regressed on sex-

specific parent size (Falconer and Mackay 1960). Sex-specific selection gradients for each

generation were calculated as the slope of the regression of relative fitness on standardized body

size. Relative fitness for each sex was calculated by dividing the number of offspring produced

by each individual by the mean number of offspring produced by that sex; standardized body

size was calculated by scaling each individual's body size to the sex-specific mean and dividing

by the sex-specific standard deviation. We then recorded how each of these factors changed

across generations (i.e., as additive genetic and phenotypic variance change), and how that

change varied with (1) the selection regime and (2) the number of loci that were sex-specific

rather than general in their effect on body size. Finally, given that intralocus sexual conflict is

known to affect intrinsic population growth rates, we also plotted population size over 500

generations for each selection regime and calculated geometric mean population growth rates

(Lewontin and Cohen 1969) for each selection regime and number of sex-specific loci. Using

data from all 11 replicates of each of the 35 model scenarios, we used linear models to determine

whether geometric mean population growth rates differed with the number of sex-specific loci in

each selection regime.

Results

Results were not qualitatively different among the 11 replicate sets of model runs. Therefore,

for all analyses except population growth rates, we report results from a single set of 35 model

runs.

Increasing the number of sex-specific loci decreased intersexual heritability. Intrasexual

heritability was approximately 1 and was constant across selection regimes, numbers of sex-

specific loci, and generations. Intersexual heritability generally decreased from 1 to 0 as the

number of sex-specific loci increased from 0 to 20 (Figure 1). This trend was consistent across

all selection regimes.

Figure 1. Box plots show mean values of inter- and intrasexual heritability averaged across 500 generations in a "no selection" simulation. Intersexual heritability declined as the number of sex-specific loci increased. Intrasexual heritability for body size did not vary with the number of sex-specific loci. Results were qualitatively similar across all other selection regimes.

Increasing the number of sex-specific loci also facilitated the evolution of sexual size

dimorphism. Sexual dimorphism evolved in all selection regimes when intersexual heritability

was less than 1 (>0 sex-specific loci; Figure 2). The rate and degree to which the sexes diverged

in body size depended on the selection regime and the number of sex-specific loci (i.e., the

intersexual heritability). In all cases, sexual dimorphism evolved faster and to a greater degree

when intersexual heritability was low (fewer sex-specific loci) and when selection was sexually

antagonistic (males experienced directional selection favoring betas). This result is consistent

with existing literature and theory on the evolution of sexual dimorphism (Cox and Calsbeek

2009).

Figure 2. Boxplots showing the final mean body size of females and males across a range of the number of sex-specific loci and three different selection regimes in (a) a no selection regime, (b) a negative frequency-dependent selection regime in which initial tactic frequencies were equal, and (c) a directional selection regime in which beta males were 1.8 x more likely to survive and mate than alpha males. Dotted horizontal line represents the size threshold separating alpha and beta males, therefore male boxplots overlapping this line indicate that both tactics persisted over 500 generations in that scenario. Results from negative frequency-dependent selection regimes with different initial frequencies and from directional selection regimes with different mating probability ratios are qualitatively the same as (b) and (c), respectively, and are not shown here.

Sex-specific selection gradients reflected the imposed selection regimes initially in all

selection regimes (Figure 3). In subsequent generations, selection gradients for both males and

females converged on zero in most models, as the population responded to selection and body

size evolved. Smaller selection gradients in later generations reflected a reduction in the variance

in male body size (as body size responds to selection), a reduction in the covariance between

female body size and fecundity (as females reach a size at which the fecundity-body size

relationship asymptotes), or both. Differences in sex-specific selection gradients were maintained

in the directional and negative frequency-dependent selection regimes in which intersexual

heritability was high (zero sex-specific loci) and ARTs were maintained.

Figure 3. Selection gradients on males and females over 500 generations and across a range of the number of sex-specific loci (nSSML) in (a) a no selection regime, (b) a negative frequency-dependent selection regime in which initial tactic frequencies were equal ($F_a=F_b$), and (c) a directional selection regime in which beta males were 1.8 x more likely to survive and mate than alpha males. Differences in sex-specific selection gradients were maintained in (b) the negative frequency-dependent and (c) directional selection regimes in which intersexual heritability is high (zero sex-specific loci), constraining the evolution of sexual dimorphism and maintaining the gender load.

Selection regimes and intersexual heritabilities affected geometric mean population growth

rates ($F_{7,377} = 23.91$, $p << 0.0001$). In most models, population size increased from 1000 in the

parent generation to an asymptote of approximately 2000 individuals (Figure 4). However, in all

the directional selection and negative frequency-dependent selection regimes, population size

and geometric mean population growth rate were greatly reduced when there were zero sex-

specific loci (i.e., when intersexual heritabilities were approximately 1). In no-selection regimes,

geometric mean population growth rates did not differ across numbers of sex-specific loci (mean

$\pm$ SD = 0.0038 $\pm$ 0.00038, $F_{1,53} = 0.038$, $p > 0.8$). In negative frequency-dependent and

directional selection regimes, geometric mean population growth rates differed across numbers

of sex-specific loci (negative frequency-dependent selection: $F_{1,163} = 111.19$, $p << 0.0001$;

directional selection: $F_{1,163} = 56.39$, $p << 0.0001$) such that they were lowest when the number of

sex-specific loci was zero. The degree to which population growth rate was reduced when there

were no sex-specific loci (compared to other values of sex-specific loci) did not vary with

different starting frequencies in the negative frequency-dependent selection models ($F_{2,30} = 2.21$,

$p = 0.13$). However, in directional selection regimes with zero sex-specific loci (i.e. models with

intralocus sexual conflict), population growth rates decreased as the strength of directional

selection against large (alpha) males increased ($F_{2,30} = 45.65$, $p \ll 0.0001$). These reductions in

population size and geometric mean population growth rate were consistent with the unresolved

gender load maintained when males and females were not able to evolve independently

(Arnqvist and Tuda 2010). The maintenance of the gender load over time was evident in the sex-

specific selection gradients for the directional selection simulations with no sex-specific loci

(Figure 3).

Figure 4. Population sizes over 500 generations and across a range of the number of sex-specific loci (nSSML) in (a) a no selection regime, (b) a negative frequency-dependent selection regime in which initial tactic frequencies were equal ($F_a=F_b$), and (c) a directional selection regime in which beta males were 1.8 x more likely to survive and mate than alpha males. In (b) the negative frequency-dependent and (c) directional selection regimes in which intersexual heritability is high (zero sex-specific loci), population size is reduced because of the gender load.

Alternative reproductive tactics were maintained under three different conditions, comprising

19 of the 35 total scenarios we modeled. Of the 19 scenarios in which ARTs were maintained, 15

were under negative frequency-dependent selection, 2 were under the no selection regime, and 2

were under the directional selection regime. First, under negative frequency-dependent selection

(i.e., the probability of mating for tactic T is 1-F_T), both tactics were maintained regardless of

initial tactic frequencies or the degree of intersexual heritability (Figure 5). This result was

consistent across all 11 replicate model runs.

Figure 5. Frequency of alpha and beta males over 500 generations and across a range of the number of sex-specific loci (nSSML) in a "frequency-dependent selection" regime in which initial tactic frequencies were equal (F_a=F_b). Tactic frequencies were not qualitatively different when initial tactic frequency ratios were 2:1 or 1:2 and are thus not shown here.

There are two general situations besides negative frequency-dependent selection, comprising

4 modeled scenarios, in which ARTs were preserved across 500 generations in our model. First,

ARTs were maintained in the no selection regime when intersexual heritability was low (e.g., 15

or 20 sex-specific loci; Figure 6). Under no selection and 15 sex-specific loci, both tactics

persisted over 500 generations in 9 of 11 model runs; under no selection and 20 sex-specific loci,

both tactics persisted over 500 generations in 10 of 11 model runs. In these same models, alpha

males comprised >75% of the population by the 500[th] generation in 10 out of 11 "no selection"

model runs with 15 sex-specific loci and 5 out of 11 "no selection" model runs with 20 sex-

specific loci. Across all 11 "no selection" model run replicates, increasing the number of sex-

specific loci resulted in a slower trend toward fixation of alpha males (Figure 6).

Figure 6. Frequency of alpha and beta males over 500 generations and across a range of the number of sex-specific loci (nSSML) in a "no selection" regime.

ARTs were also maintained by intralocus sexual conflict over body size. This occurred when

selection on size was sexually antagonistic (favoring small males and large females) and

intersexual heritability was high (e.g., zero sex-specific loci / all loci affect males and females

equally; Figure 7). This was especially evident at intermediate strengths of directional selection

(e.g., when beta males were 1.8 – 2 times more likely to survive and reproduce than alpha

males), where direct selection on males was strong enough to reduce the frequency of alphas but

weak enough not to eliminate them entirely. In this situation ARTs were maintained because

there was a balance between (a) correlated fecundity selection and high intersexual heritability

producing larger sons and (b) selection against those large sons. This outcome was consistent

across all 11 replicates of our model runs.

Figure 7. Frequency of alpha and beta males over 500 generations and across a range of the number of sex-specific loci in a "directional selection" regime in which the mating probability of beta males was (a) 1.5, (b) 1.8, and (c) 2 times that of alpha males.

Discussion

Alternative reproductive tactics abound in nature (Sinervo and Calsbeek 2010), yet few

processes have been identified that sustain them. Those that have been identified – negative

frequency-dependent selection and disruptive selection – both focus on selection acting directly

on the sex exhibiting the polymorphism (Sinervo and Calsbeek 2006). We show that selection

acting on both sexes, and intralocus sexual conflict that results when genetic architecture is

shared, may also preserve sex-specific alternative reproductive tactics over hundreds of

generations. If male ARTs differentially affect the fitness of their daughters, then they can be

maintained (or eliminated) by selection for (against) their daughters even in the absence of

negative frequency-dependent fitness.

In addition to negative frequency-dependent selection, two other processes emerged from our

model that preserved alternative tactics over 500 generations. First, both tactics persisted when

there was no direct selection on males, and most (15) or all (20) loci were sex-specific in their

expression. In this scenario there was also no correlated selection on males, because males could

not pass on their size to their daughters. Fecundity selection favored large females, but these

females could not pass on their large size to their sons because intersexual heritability was zero.

Thus, there was neither direct nor correlated selection on male size, allowing the persistence of

both tactics over 500 generations. The fact that alpha males comprised the majority of males by

the 500[th] generation in most of these model runs suggests that these scenarios could not maintain

ARTs indefinitely. However, because the trend toward an alpha male majority occurred more

slowly in these models as the number of sex-specific loci increased, low intersexual heritability

could play a role in preserving male ARTs in natural populations where selection on males is

chronically weak or temporally and spatially variable.

The second scenario in which both ARTs were maintained over 500 generations was when

beta males were 1.8 – 2 times more likely to survive and mate than alpha males, and the number

of sex-specific loci was zero. This created a scenario in which males experienced direct selection

favoring beta males and correlated selection favoring alpha males. Because no loci were sex-

specific in their expression, size was heritable between the sexes, creating intralocus sexual

conflict. When the strength of directional selection on male size was low (i.e., the fitness of betas

was just 1.5 times higher than that of alphas), correlated selection outweighed direct selection

and alpha males went to fixation. However, as the strength of directional selection increased,

direct and correlated selection balanced each other out, and both tactics persisted over 500

generations.

Persistent intralocus sexual conflict in these scenarios was evident in the maintenance of the

gender load. Selection gradients on females remained positive and those on males remained

negative throughout these simulations due to intralocus sexual conflict constraining the evolution

of sexual dimorphism. In concordance with theory, the maintenance of the gender load in these

simulations resulted in a lower population growth rate compared with simulations in which the

sexes could respond independently to selection. This finding shows that the maintenance of

ARTs, when achieved through intralocus conflict, may reduce population fitness.

Our model also showed that one male tactic could go to fixation even when males did not

experience direct selection. When females were selected to be larger, but male body size did not

experience direct selection, alpha males became fixed in the population when intersexual

heritability for body size was high. This occurred through correlated fecundity selection. This

finding, though not explicitly predicted, again highlights the importance of understanding

selection in both sexes when attempting to understand the evolutionary maintenance of sex-

specific polymorphisms.

Our models consistently revealed that sexual conflict resulted in the persistence of a sex-

specific polymorphism over hundreds of generations, even in the absence of negative frequency-

dependent or disruptive selection. This result is consistent with a growing literature documenting

genetic correlations between male ARTs and female fitness, suggesting that the maintenance of

sex-specific ARTs through intralocus sexual conflict may be common and widespread in nature.

For example, side-blotched lizards (*Uta stansburiana*) exhibit three male morphs associated with

orange, blue, and yellow throat colors (Sinervo and Lively 1996). In this system the intersexual

heritability of throat color is high, and female throat color is correlated with fecundity, such that

the genes associated with male ART have direct consequences for female fitness and population

growth rate (Sinervo and Zamudio 2001). Though negative frequency-dependent selection seems

partly responsible for ART maintenance in this system, genetic correlations between male tactic

and female fitness also have the potential to affect ART frequencies (Sinervo and Zamudio

2001).

Male Pacific field crickets (*Teleogryllus oceanicus*) in Hawaii exhibit singing and silent

reproductive tactics that stem from different wing morphologies (normal and flatwing; Zuk et al.

2006). These wing morphologies can be inherited and expressed by daughters, and flatwing

females invest less in reproductive tissue mass, experience more frequent mating failure, and are

less likely to mount and mate with males compared to normal females (Heinen Kay et al. 2019;

Heinen-Kay et al. 2020b; Richardson et al. 2021). Because flatwing males have much higher

survival (Zuk et al. 2006) and sire more offspring than normal males (Heinen-Kay et al. 2019),

while flatwing females seem to have lower fitness, intralocus sexual conflict over wing

morphology is likely driving the maintenance of both male ARTs (Heinen-Kay et al. 2020b).

Studies of the bulb mite (*Rhizoglyphus* spp.) are also consistent with ART maintenance

through intralocus sexual conflict. In artificial selection experiments, females from lines selected

for fighter males have lower fitness (fecundity and longevity) than females from lines selected

for sneakers (Bielak et al. 2014; Łukasiewicz et al. 2020). Harano et al. (2010) showed the same

pattern in flour beetles (*Gnatocerus cornutus*) using an artificial selection experiment in which

they selected for larger or smaller male mandibles. While flour beetles do not have ARTs *per se*,

they do exhibit sexual dimorphism in a sexually selected weapon, and the intralocus sexual

conflict arising from simultaneous sexually antagonistic selection and an intersexual genetic

correlation for mandible size could maintain variation in males as shown in the present paper.

While the concepts of intralocus sexual conflict and correlated selection are decades old

(Lande and Arnold 1983; Rice and Chippindale 2001), they have only just begun to penetrate the

literature on alternative reproductive tactics. Our results show that intralocus sexual conflict can

play a role in the maintenance of ARTs, and that understanding the maintenance - or elimination

- of sex-specific polymorphisms requires considering selection in both sexes. Fruitful systems in

which to look for this phenomenon in nature will include those in which the trait or traits that

differentiate ARTs are also expressed in females (e.g., body size), or those in which genetic

correlations between ARTs and female survival or reproductive success have already been

established.

Data Availability Statement: The individual-based model used in this paper will be uploaded to

Dryad upon acceptance.

Acknowledgements: We thank Mark McPeek, Suzanne Alonzo, and Michael Whitlock for

helpful discussions and suggestions on the model and the biological systems it describes. MMG

is grateful to David Noakes for inspiration in addressing this question with a model. RC was

funded by an award from the National Science Foundation ((NSF) DEB-1655092).

Author Contributions: MMG conceived the study; MMG built the model and drafted the

manuscript; both authors contributed to critically revising and approving the final manuscript.

References:

Arnqvist, G., and M. Tuda. 2010. Sexual conflict and the gender load: correlated evolution between population fitness and sexual dimorphism in seed beetles. Proceedings of the Royal Society B 277:1345–1352.

Bielak, A. P., A. M. Skrzynecka, K. Miler, and J. Radwan. 2014. Selection for alternative male reproductive tactics alters intralocus sexual conflict. Evolution 68:2137–2144.

Bonduriansky, R. 2011. Sexual Selection and Conflict as Engines of Ecological Diversification. The American Naturalist 178:729–745.

Bonduriansky, R., and S. F. Chenoweth. 2009. Intralocus sexual conflict. Trends in Ecology & Evolution 24:280–288.

Buzatto, B. A., and H. L. Clark. 2020. Selection for Male Weapons Boosts Female Fecundity, Eliminating Sexual Conflict in the Bulb Mite. Scientific Reports 10:2311.

Buzatto, B. A., H. L. Clark, and J. L. Tomkins. 2018. Morph-specific artificial selection reveals a constraint on the evolution of polyphenisms. Proceedings of the Royal Society B 285:20180335.

Connallon, T., and A. G. Clark. 2012. A general population genetic framework for antagonistic selection that accounts for demography and recurrent mutation. Genetics 190:1477–1489.

Connallon, T., and A. G. Clark. 2014. Balancing Selection in Species with Separate Sexes: Insights from Fisher's Geometric Model. Genetics 197:991–1003.

Cordero, A. 1990. The inheritance of female polymorphism in the damseifly Ischnura graellsii (Rambur) (Odonata: Coenagrionidae). Heredity 64:341–346.

Cox, R. M., and R. Calsbeek. 2009. Sexually antagonistic selection, sexual dimorphism, and the resolution of intralocus sexual conflict. The American Naturalist 173:176–187.

Emlen, D. J. 1994. Environmental Control of Horn Length Dimorphism in the Beetle

Onthophagus acuminatus (Coleoptera: Scarabaeidae). Proceedings of the Royal Society B

256:131–136. The Royal Society.

Falconer, D. S., and T. F. C. Mackay. 1960. Introduction to Quantitative Genetics.

Foerster, K., T. Coulson, B. C. Sheldon, J. M. Pemberton, T. H. Clutton-Brock, and L. E. B.

Kruuk. 2007. Sexually antagonistic genetic variation for fitness in red deer. Nature

447:1107–1110.

Gross, M. R. 1996. Alternative reproductive strategies and tactics: diversity within sexes. Trends

in Ecology and Evolution 11:92–98.

Gross, M. R. 1985. Disruptive selection for alternative life histories in salmon. Nature 313:47–

48.

Gross, M. R. 1991. Evolution of alternative reproductive strategies: frequency-dependent sexual

selection in male bluegill sunfish. Pp. 55–75 *in* R. Wooton and G. Potts, eds. Fish

reproduction: strategies and tactics. … Transactions of the Royal Society of …, London.

Gross, M. R., and J. Repka. 1998. Stability with inheritance in the conditional strategy. Journal

of Theoretical Biology 192:445–453.

Harano, T., K. Okada, S. Nakayama, T. Miyatake, and D. J. Hosken. 2010. Intralocus Sexual

Conflict Unresolved by Sex-Limited Trait Expression. Current Biology 20:2036–2039.

Heinen Kay, J. L., D. B. Strub, S. L. Balenger, and M. Zuk. 2019. Direct and indirect effects of

sexual signal loss on female reproduction in the Pacific field cricket (Teleogryllus

oceanicus). Journal of Evolutionary Biology jeb.13534-31.

Heinen-Kay, J. L., R. E. Nichols, and M. Zuk. 2020a. Sexual signal loss, pleiotropy, and

maintenance of a male reproductive polymorphism in crickets. Evolution 74:1002–1009.

Heinen-Kay, J. L., R. E. Nichols, and M. Zuk. 2020b. Sexual signal loss, pleiotropy, and

maintenance of a male reproductive polymorphism in crickets. Evolution 74:1002–1009.

Heinen-Kay, J. L., E. M. Urquhart, and M. Zuk. 2019. Obligately silent males sire more

offspring than singers in a rapidly evolving cricket population. Biology Letters

15:20190198.

Hughes, K. A., A. E. Houde, A. C. Price, and F. H. Rodd. 2013. Mating advantage for rare males

in wild guppy populations. Nature 503:108–110.

Hunt, J., and L. W. Simmons. 1997. Patterns of fluctuating asymmetry in beetle horns: an

experimental examination of the honest signalling hypothesis. Behavioral Ecology and

Sociobiology 41:109–114.

Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters.

Evolution 34:292.

Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated characters.

Evolution 37:1210.

Lank, D. B., C. M. Smith, O. Hanotte, T. Burke, and F. Cooke. 1995. Genetic polymorphism for

alternative mating behavior in lekking male ruff *Philomachus pugnax*. Nature 378:59–62.

Lewontin, R. C., and D. Cohen. 1969. On population growth in a randomly varying environment.

Proc Natl Acad Sci U S A 62:1056–1060.

Łukasiewicz, A., M. Niśkiewicz, and J. Radwan. 2020. Sexually selected male weapon is

associated with lower inbreeding load but higher sex load in the bulb mite. Evolution

74:1851–1855.

Moczek and Emlen. 1999. Proximate determination of male horn dimorphism in the beetle

Onthophagus taurus (Coleoptera: Scarabaeidae). Journal of Evolutionary Biology 12:27–

37.

Neff, B. D., and E. I. Svensson. 2013. Polyandry and alternative mating tactics. Philosophical

Transactions of the Royal Society B: Biological Sciences 368:20120045–20120045.

Oliviera, R. F., M. Taborsky, and H. J. Brockmann. 2008. Integrating mechanisms and function:

prospects for future research. Cambridge University Press, Cambridge.

Otto, S. P., and T. Day. 2007. A Biologist's Guide to Mathematical Modeling in Ecology and

Evolution. Princeton University Press.

Rice, W. R., and A. K. Chippindale. 2001. Intersexual ontogenetic conflict. Journal of

Evolutionary Biology 14:685–693.

Richardson, J., J. L. Heinen-Kay, and M. Zuk. 2021. Sex-specific associations between life-

history traits and a novel reproductive polymorphism in the Pacific field cricket. Journal

of Evolutionary Biology 0:1–9.

Shuster, S. M. 1989. Male Alternative Reproductive Strategies in a Marine Isopod Crustacean

(Paracerceis sculpta): The Use of Genetic Markers to Measure Differences in

Fertilization Success Among α-, β-, and γ-Males. Evolution 43:1683–1698. [Society for

the Study of Evolution, Wiley].

Shuster, S. M., and M. J. Wade. 2003. Mating systems and strategies. Princeton University Press,

Princeton, NJ.

Sinervo, B., and R. Calsbeek. 2010. Behavioral concepts of selection: experiments and causes of

selection on the sexes. P. *in* Evolutionary Behavior Ecology. Oxford University Press.

Sinervo, B., and R. Calsbeek. 2006. The Developmental, Physiological, Neural, and Genetical

Causes and Consequences of Frequency-Dependent Selection in the Wild. Annual

Review of Ecology, Evolution, and Systematics 37:581–610.

Sinervo, B., and C. M. Lively. 1996. The rock–paper–scissors game and the evolution of

alternative male strategies. Nature 380:240–243.

Sinervo, B., D. B. Miles, W. A. Frankino, M. Klukowski, and D. F. DeNardo. 2000a.

Testosterone, Endurance, and Darwinian Fitness: Natural and Sexual Selection on the

Physiological Bases of Alternative Male Behaviors in Side-Blotched Lizards. Hormones

and Behavior 38:222–233.

Sinervo, B., E. Svensson, and T. Comendant. 2000b. Density cycles and an offspring quantity

and quality game driven by natural selection. Nature 406:985–988.

Sinervo, B., and K. R. Zamudio. 2001. The evolution of alternative reproductive strategies:

fitness differential, heritability, and genetic correlation between the sexes. The Journal of

Heredity 92:198–205.

Thornhill, R. 1981. Panorpa (Mecoptera: Panorpidae) Scorpionflies: Systems for Understanding

Resource-Defense Polygyny and Alternative Male Reproductive Efforts. Annual Review

of Ecology and Systematics 12:355–386.

Zuk, M., J. T. Rotenberry, and R. M. Tinghitella. 2006. Silent night: adaptive disappearance of a

sexual signal in a parasitized population of field crickets. Biology Letters 2:521–524.

523