Drastic mutations can be hidden in short-read mapping: thousands of mutation clusters in human genome are explained by short-range template switching

Ari Löytynoja[1,*]

1 Institute of Biotechnology, University of Helsinki, Finland

* ari.loytynoja@helsinki.fi

Abstract

Variation within human genomes is distributed unevenly and variants show spatial clustering. DNA-replication related template switching is a poorly known mutational mechanism capable of causing major chromosomal rearrangements as well as creating short inverted sequence copies that appear as local mutation clusters in sequence comparisons. We reanalyzed haplotype-resolved genome assemblies representing 25 human populations and multinucleotide variants aggregated from 140,000 human sequencing experiments. We found local template switching to explain thousands of complex mutation clusters across the human genome, the loci segregating within and between populations with a small number appearing as de novo mutations. We developed computational tools for genotyping candidate template switch loci using short-read sequencing data and for identification of template switch events using both short-read data and genotype data. These tools will enable building a catalogue of affected loci and studying the cellular mechanisms behind template switching both in healthy organisms and in disease. Strikingly, we noticed that widely-used analysis pipelines for short-read sequencing data – capable of identifying single nucleotide changes – may miss TSM-origin inversions of tens of base pairs, potentially invalidating medical genetic studies searching for causative alleles behind genetic diseases.

Author summary

Mutations are not randomly distributed in genomes and they often appear as clusters of nearby changes. We earlier showed that a poorly known mechanism in DNA replication can create short inverted copies of nearby sequence and that these events then show as mutation clusters in sequence comparison. Using the latest DNA sequencing and variation data we show that the human genome contains thousands of mutation clusters consistent with this mechanism and that novel mutations are created at a significant rate. Strikingly we observed that widely used methods for processing DNA sequencing data may completely miss these mutations. This has significance e.g. in medical genetic studies aiming to identify mutations causing genetic diseases.

Introduction   1

Twenty years after the publication of the draft genomes [1, 2], the Telomere-to-Telomere   2
Consortium reported the first truly complete sequence of a human genome [3, 4]. Every   3
genome is unique, however, and the challenge now is to understand the global genomic   4
diversity in the human population [5, 6] and to build a comprehensive pangenome to   5
represent this variation [7]. While single nucleotide variation (SNV) and short   6
insertion-deletions (indels) have been resolved relatively accurately with classical   7
short-read sequencing [8], the characterization of structural variants (SVs) and   8
low-complexity sequences has been lacking. The Human Genome Structural Variation   9
Consortium developed a method [9] for phased diploid genome assembly with   10
combination of long-read PacBio whole-genome sequencing (WGS) [10] and Strand-seq   11
phasing [11] data. Ebert *et al.* [12] applied this method to 32 diverse human individuals   12
and produced 64 assembled haplotypes, i.e. maternal and paternal copies of the genome.   13
With the help of phased diploid genomes, they massively expanded the catalogue of   14
known SVs and, having the information of co-inherited nearby sequence differences,   15
could study in detail the different subclasses of complex variants. Such accurate   16
characterization of multiple genomes will lay the foundations for understanding the role   17
of complex mutations in human phenotypes and disease [13, 14, 15] and in evolution   18
[13, 16, 17]. More generally, the unprecedented resolution of the new genome data will   19

open novel possibilities for understanding the mutational mechanisms of complex [20] eukaryotic genomes [18, 19, 20]. [21]

While many subclasses of variants studied by Ebert *et al.* are clearly independent, [22] mutations are known to be clustered [20] and their separation in the analysis stage may [23] dismiss crucial information. We compared earlier two independent assemblies of the [24] human genome [21, 22] and found mutation clusters consistent with DNA-replication [25] related template switching [23]. The mechanisms, originally described and studied in [26] bacteria [24, 25], resembles FoSTeS (Fork Stalling and Template Switching) [26] and [27] MMBIR (Microhomology-Mediated Break-Induced Replication) [27] but is local and the [28] reciprocal switches typically take place within a region of a few tens or hundreds of base [29] pairs (bp). Unlike the chromosomal rearrangements created by FoSTeS and MMBIR, [30] the footprint of local template switch mutations (TSMs) are short inversions, either [31] happening in place or creating a reverse-complement copy of a nearby sequence region; [32] these are believed to be produced by the replication briefly switching either to the [33] complementary DNA strand or going backwards along the nascent DNA strand [24, 25] [34] (Fig. S1). Mutations compatible with the TSM mechanism appear as multiple nearby [35] sequence changes and may be represented as combinations of SNPs, MNPs (single- and [36] multi-nucleotide polymorphisms) and short indels in variant data. We showed [23] that [37] the TSM candidates identified between the two assemblies segregate in the 1000 [38] Genomes population data; that TSM-like mutation patterns can be identified in [39] genotype data; and that parent-offspring trios [28] contain consistent de novo mutations [40] (DNMs). The haplotype-resolved data of Ebert *et al.* and the variation information [41] compiled since then by massive sequencing studies [5, 6] provide an opportunity to [42] revisit the topic and study the TSM mechanism's role in generation of genomic [43] variation. As haplotype-resolved genome data are still rare, it is of interest to know how [44] reliably the TSM patterns found with de-novo-assembled genomes can be identified [45] using the old-school short-read DNA sequencing. [46]

Results

TSMs explain thousands of haplotypic mutation clusters

Utilising the variant data of Ebert *et al.* [12] (called here "HaplotypeSV data"), we combined the SNVs, indels and SVs, and for each maternal and paternal genome copy, identified the clusters of sequence differences (see Methods). For each mutation cluster locus within a haplotype, we placed the identified variants into their genomic context, creating two alleles with 150 bp of sequence context (Fig. 1A,B). Finally, having the two alleles – the region from the GRCh38 reference (called 'REF') and the alternative allele formed by the mutation cluster (called 'ALT') – we applied the four-point alignment (FPA) algorithm [23] to find the best solution involving a template switch, testing both alleles as the ancestral state (Fig. 1C,D; see Methods). Despite the attempts to evaluate TSMs under a probabilistic model [29], it is unclear when a potential TSM solution should be considered more plausible than a combination of SNVs and indels. We set an arbitrary rule that the TSM candidates need to (i) be supported by at least two sequence changes of which at least one is a base change, and as a result show a higher sequence identity across the whole region than the original forward-aligned solution, and (ii) have a ②→③ region, inferred to be copied from the other template, of at least 8 bases long. Despite seeing strong evidence for many VNTRs (variable number of tandem repeats) evolving through template switching (Fig. S3), we required the candidate ②→③ regions to be contain all four bases, thus removing hits within low-complexity sequence.

Per individual, we found 3,049–3,934 to mutation patterns consistent with the TSM mechanisms, the longest of them 318 bp in length (Table 1, 'ALL'). Many of these were in proximity of repeat elements and could potentially have been affected by gene conversion. After removing the loci masked as repetitive sequence in the reference genome, there were 872–1,121 inferred TSM events per individual (Fig. 1E), the maximum length varying from 97-180 and the mean length being 11.56 bp (Table 1, 'UNMASKED'; Fig. S4). Homozygosity of the inferred TSM loci varied from 26.7–42.3% (Fig. 1E), and the proportion of singletons from 2.1–9.0% (Table S1). Although comparisons on superpopulation level are affected by the different sample sizes, African (AFR) individuals expectedly had the lowest level of homozygosity and fixed loci, 29.2% and 0.7%, and the highest proportion of singleton loci, 7.8%, while those for Europeans

Fig 1. TSMs in HaplotypeSV data. (**A**) HaplotypeSV data contain a cluster of five SNVs that segregate as a unit among the individuals (first six shown). (**B**) Within their genomic context, REF (cyan) and ALT (orange) alleles create two haplotypes. (**C**) The FPA algorithm [23] attempts to explain the observed differences with two reciprocal template switch events. In this case, the ALT sequence (A1, A2, A3; shown in green, blue and red) can be created from the REF sequence by copying the A2 fragment in reverse-complement (RR; shown in magenta). (**D**) The template-switch solution fully explains the cluster of five base differences. (**E**) Total numbers of inferred TSM events in different HaplotypeSV samples after removal of low-complexity sequences (full height, left); proportion of homozygous loci (middle, right); and proportion of ancestral (dark shading), derived (light shading) and unpolarized (no shading) alleles (bottom, right).

(EUR, including Azkenazi NA24385) and Asians (AAS, including South and East Asians) were higher and lower, respectively (Table 1). The greater variation of African populations and the old age of the TSM patterns was reflected in the sharing of loci across the superpopulations: AFR shared 45.6% and 49.9% of its TSM loci with EUR and AAS, respectively, whereas the latter two shared 75.3–80.7% of their TSM loci with other superpopulations (Table 1). ^{78} ^{79} ^{80} ^{81} ^{82} ^{83}

A unique feature of TSMs is that often the alternative alleles can be polarized, i.e. have the ancestral and derived state determined, without an outgroup (Fig. S2). We could define the ancestral allele in 87.7–90.3% of the cases and found the proportion of loci with the ALT variant being the ancestral allele ranging from 49.7–55.3 across the individuals (Fig. 1E; Table S1). On superpopulation level, Europeans had the highest average proportion of ancestral variants (Table 1), probably reflecting the European origin of the GRCh38 reference genome and thus it capturing a greater amount of derived European TSM variation. ^{84} ^{85} ^{86} ^{87} ^{88} ^{89} ^{90} ^{91}

Table 1. TSM statistics for HaplotypeSV data.

Superpopulation		AFR	EUR	AAS
Haplotypes		20	14	22
ALL	TSMs/individual	3783	3135	3173
	unique TSMs	12,501	7,370	8,956
	maximum length	318	188	188
	average length	13.14	13.14	13.15
UNMASKED	TSMs/individual	1,074	906	919
	unique TSMs	3,133	1,899	2,246
	maximum length	180	180	180
	average length	10.61	10.56	10.55
	singletons	7.8%	2.7%	3.2%
	fixed	0.7%	1.8%	2.8%
	homozygous	29.2%	35.3%	38.5%
	ancestral	52.0%	53.9%	51.5%
	shared with AFR	–	75.3%	69.6%
	shared with EUR	45.6%	–	68.2%
	shared with AAS	49.9%	80.7%	–

Known TSM loci can be genotyped with short-read data

One individual in the HaplotypeSV data, NA12878, is the mother of a three-generation 17-member pedigree (CEPH 1463) sequenced to high coverage [30] as well as one of the reference samples of the Genome in a Bottle (GIAB) initiative [31]. We genotyped the TSM loci identified in NA12878 in different GIAB datasets and in the pedigree 1463 in order to assess their de novo vs. standing, in vitro vs. in vivo, and somatic vs. germline origin. Utilising the HaplotypeSV genotype data for NA12878, we reconstructed the genome context for the alternative alleles and, for each individual in the family, mapped the reads extracted from that particular loci against the two alternative alleles. Outside the mutation locus, the short reads should map evenly between the two alleles, but the reads overlapping the position that differ between the alleles are expected map to one allele only (Fig. 2A,B). By comparing the read coverage across the two alternative alleles, we could identify the test individual as homozygote for REF allele (R|R), heterozygote (R/A), or homozygote for ALT allele (A|A). We adopted a strategy similar to base calling and required 99% posterior probability for the genotype calls (see Methods).

We started with the Illumina Platinum [30] and 10X Genomics Chromium [32]

Fig 2. Genotyping of a TSM locus. (**A**) TSM solution for a complex mutation in chr10. (**B**) The two haplotypes with alternative central parts (blue and magenta; top) are used as the reference for re-mapping of reads extracted from the locus. NA12877 (top) is heterozygous and mapping coverage is uniform for the two alleles (gray). NA12890 (middle) and NA12889 (bottom) are homozygous for the REF (left, reads in pink) and ALT (right, reads in light blue) alleles, respectively. Genotype is inferred of the mapping coverage in the middle of the locus (blue and magenta dots and dotted lines). (**C**) NA12878 Platinum data with full and clipped reads in yellow and red, respectively, and number of clip ends as blue bars (bottom).

datasets for NA12878 and genotyped the TSM-like loci found in the HaplotypeSV data. Focusing on the 864 loci that were not masked as repeats or low-complexity sequence and had enough read coverage in both datasets, we found the two short-read datasets to provide the genotype and agree on it in 831 cases (Fig. 3A; Table S2). While 14 loci could not be genotyped with the short-read data using the 99% posterior probability cut-off, the variant pattern in the HaplotypeSV data was inconsistent in 84 cases, and the genotype of the loci could not determined. Of the 780 loci inferred to be either heterozygous or homozygous for the ALT allele in the HaplotypeSV data, the three datasets agreed in 735 cases (Table S2). Eight loci recorded variable in the HaplotypeSV data were inferred to be homozygous for the REF allele with both short-read datasets. Many of these were in repetitive sequence context and all loci were found segregating among other samples in the HaplotypeSV data ruling out the possibility of DNMs. A small number of the TSM candidate loci showed inconsistency among different datasets, possibly due to incorrectly mapped reads (Fig. S5).

To further verify the genotyping approach, we performed a similar analysis on the parents and the children of NA12878. On the parents, we genotyped the 783 loci found

Fig 3. Confirmation of HaplotypeSV TSM loci with short-read data. (**A**) Ratio of REF and ALT allele mapping coverage ($r = depth_{\text{REF}}/depth_{\text{ALT}}$) reflects the genotype: $r \approx 1$ (and thus $log_{10}(r) \approx 0$) for a heterozygote; $r \ll 1$ and $r \gg 1$ for the two types of homozygotes. The Log10-ratios agree for the NA12878 Platinum and Chromium datasets, and 95.5% of the 883 loci identified in HaplotypeSV data are called to contain at least one ALT allele with 99% posterior probability in both datasets; 17 and 10 loci are called homozygous REF with at least one dataset and by both (top-right corner). (**B**) Of the 783 variable loci in NA12878 Chromium data, all but one locus are called to contain at least one ALT allele in the parents. In the 290 loci homozygous for ALT in NA12878 (blue), both parents contain at least one ALT allele. 'Unknown' indicates inconsistent variant data or low posterior probability. One pseudo-count was added to all values to avoid divisions by zero.

in the HaplotypeSV data and called to contain at least one ALT allele in the NA12878 ₁₂₅
Chromium data. We found strong agreement between the inferred genotypes and, of the ₁₂₆
307 loci called homozygous for the ALT allele in NA12878, 290 could be genotyped with ₁₂₇
the both parental Chromium data and each of them contained at least one ALT allele ₁₂₈
(Fig. 3B; Table S3). The only locus appearing as a novel mutation according to parental ₁₂₉
data (R|R:R|R in Fig. 3B) was a false positive and in a manual verification one parent ₁₃₀
was found to contain two supporting reads. Genotyping of the loci in the eleven children ₁₃₁
(CEPH pedigree 1463) required their transfer from GRCh38 to GRCh37. The liftover ₁₃₂
worked and the parents' alignment data (Platinum/GRCh37) contained enough reads to ₁₃₃
genotype 843 of the TSM-like loci found variable in NA12878 in the HaplotypeSV data. ₁₃₄
With the exception of loci that were inferred to be heterozygous in both parents ₁₃₅
(R/A,R/A), the observed genotypes matched nearly perfectly the expectation under ₁₃₆
Hardy-Weinberg equilibrium (Table 2). An excess of heterozygotes in genotype data is a ₁₃₇
classical mark of an artefact caused by sequencing reads originating from different loci, ₁₃₈
e.g. due to duplication of genome regions. Seeing those among the CEPH 1463 children ₁₃₉
suggests that a proportion of inferred variants for NA12878 in the HaplotypeSV data ₁₄₀
are erroneous. The TSM events causing those variants may be real but they have ₁₄₁
happened in duplicated copies, not within the loci where the calls were made. ₁₄₂

Table 2. Genotypes (%) of TSM loci in CEPH 1463 children.

Parental genotypes	n	Offspring expected			Offspring observed			
		R|R	R/A	A|A	R|R	R/A	A|A	NA*
R|R,R|R	7	100	0	0	72.7	24.7	0	2.6
R|R,R/A	169	50	50	0	48.6	50.5	0.5	0.4
R|R,A|A	37	0	100	0	0	98.5	1.5	0
R/A,R/A	214	25	50	25	16.8	63.5	17.5	2.2
R/A,A|A	206	0	50	50	0.4	53.0	45.1	1.6
A|A,A|A	177	0	0	100	0	0.5	98.5	1.1

*Loci with posterior probability below 0.99 considered as unknown

Novel TSM loci can be discovered with short-read data

Many of the TSM loci detected in HaplotypeSV data had not been originally called with the short-read data despite the two alleles differing over several tens of bases. A closer look on these revealed that alternative TSM haplotypes can show nearly uniform sequencing coverage across the region if the ②→③ region of the inferred TSM process (Fig. 2A) is long enough: the mapping algorithm then produces a descent mapping for the read core by clipping the mismatching overhangs (Fig. 2C). While this is clearly undesirable behaviour, the observation suggested a strategy to identify potential TSM loci in short-read data by searching for clusters of clip sites, extracting the reads mapped to the region and then producing a local reassembly of the reads; in positive cases, the dissimilarities between the assembled sequences could be explained by a TSM event.

To test that, we took SvABA [33], a tool developed for detection of structural variants by local reassembly of short sequencing reads, and integrated our FPA algorithm [23] to search for TSM patterns in the resulting contigs. We independently identified clusters of clip sites in the Illumina Platinum [30] and 10X Genomics Chromium [32] short-read data for NA12878 and clusters of variants in the corresponding variant data, and applied SvABA-FPA on each candidate locus (see Methods). With this, we found 1,054 and 26,222 candidate TSM loci in the Platinum and Chromium data, respectively, the significantly different numbers reflecting the technological differences between the standard and the linked-reads sequencing as well as the downstream bioinformatic methods used [34, 35]. We applied rigorous filtering (see Methods) and reduced, respectively, the numbers to 214 and 757 high-confidence

Fig 4. TSM candidate identified with short-read data. (**A**) In Platinum data (left), a region in chr10 has excess of soft clips (bottom, blue bars) and called variants (SNVs in red, indels in black); the mapping coverage (top, in gray) also shows atypical patterns. (**B**) Similar signals are seen in Chromium data. (**C**) Platinum variant data contain six SNVs and six indels within 28 bp. (**D**) De novo assembly of the reads creates two locally highly dissimilar haplotypes compatible with the called variants (top). All differences can be explained with a TSM event, an inversion in place (middle). Using two haplotypes with alternative central parts (blue and magenta; bottom) as the reference, extracted reads map in full length (**A,B**; middle, right) with roughly even coverages. No variants for NA12878 were called in this region by Ebert *et al.* [12].

loci. Encouragingly, 181 of the loci were shared among the short-read datasets, and 121 and 148 of the loci overlapped with the 883 HaplotypeSV cases, the filtering thus removing 79.7 and 97.1 % of the candidate loci but only 37.0 and 34.8 % of the overlaps.

We studied in more detail the 59 cases that had the ②→③ fragment of at least 25 bases long (Data S1). Most of the cases showed the expected signal in the original alignment data that was removed when the same reads were mapped against the two alternative haplotypes (Fig. 4A,B) while in a few rare cases the variant calling had correctly captured the differences between the two TSM haplotypes (Fig. 4C,D). Some of the heterozygous loci were not called in any variant call set despite significant differences between the two haplotypes (Fig. S6, Data S1), and, even more strikingly, several loci homozygous for the ALT allele were uncalled in different variant call sets,

including the HaplotypeSV data (Fig. S7). Although nominally of the same origin, the studied sequencing datasets derive from lymphoblastoid cell lines and the differences may indicate activity of the TSM mechanism in vitro cell cultures [36]. Among the 59 cases with the longest ②→③ fragment, 17 loci appeared to be DNMs in the Chromium data whereas no unique TSM loci were inferred in the Platinum data (Data S1). While this is consistent with the lower total number of hits in Platinum data, the result may also reflect the differences in the sequencing technologies and bioinformatic pipelines [32]. Although the lower mapping coverage for ALT alleles (Fig. S8, Data S1) is expected from DNMs, we cannot exclude the possibility of artefacts in the underlying data.

TSMs explain thousands of clusters in variation databases

A significant fraction of the TSM-candidate loci were missing from the original variant call sets (Fig. 4B), but none of the manually confirmed cases supported by both short-read datasets was truly novel and absent from the dbSNP database [37] (Data S1). While dbSNP correctly lists most studied loci as multinucleotide variants (MNVs), all the loci were also present as multiple SNVs and indels and some only as multiple independent MNV or indels (Fig. S9, Data S1). The latter is slightly surprising as variants originating from a complex mutation are expected to be present in their entirety or not at all, and similar allele frequencies should allow phasing and grouping independent variants into MNVs [38].

The Genome Aggregation Database (gnomAD, v2.1.1) [5] contains variation found among 125,000 exomes and 15,700 genomes of unrelated individuals sequenced as part of various disease-specific and population genetic studies. Wang *et al.* [39] identified gnomAD SNVs appearing within proximity of 1-10 bp, and provide them as phased variant pairs. Using these data, we selected variant pairs with similar allele counts (ACs; $\Delta AC \leq 10\%$), merged the data of different variant-pair distances, and for all clusters of three variants, created the alternative alleles and searched for a TSM solution to explain the differences (see Methods). Among the 91,300 clusters of three or more SNVs, we could find a TSM solution for 4,425 loci. Among these, we had a closer look at the 192 cases where the ②→③ region was at least 25 bp long and the REF and ALT allele differed at least by four SNVs (Data S2).

Fig 5. Complex gnomAD MNV explained by a single TSM. (**A**) Wang *et al.* [39] identified 15 SNVs within a 29 bp interval in chr12. (**B**) All the differences (top, marked with asterisks) can be explained with a TSM inverting 39 bp-long fragment in place (middle, bottom).

Among the studied cases, the TSM mechanism perfectly explained MNVs consisting of of up to 15 base changes (Fig. 5) while the longest inferred TSM events had the ②→③ region of over a hundred bases (Fig. S10). Consistent with our earlier findings, 36% of the studied TSM loci contained low complexity sequence or VNTRs, but the base changes, including those in the two longest events of 119 and 125 bp (Fig. S10B,C), were not necessarily explainable by a simple slippage mechanism; within complex sequence, the longest TSM event was 90 bp long (Fig. S10A). Interestingly, we found a locus, present in a single individual, where 16 of the 19 gnomAD MNV base changes could be explained with two adjacent TSM events of 62 and 32 bp separated by a forward fragment of 16 bp (Fig. S11), indicating that also local TSMs can be chained into complex mutations [13]. As the gnomAD MNV data lack indels, no TSM patterns involving sequence length changes were included and the real number of mutation clusters explainable by the TSM mechanism is likely much higher.

We replicated the analysis with the DNM data from 33 large, three-generation CEPH families by Sasani *et al.* [40]. We did not find TSM-like patterns among the germline DNMs of the 70 second-generation individuals but saw four consistent patterns (Fig. S12) among the 24,975 de novo SNVs and small indels of the 350 third-generation individuals. As most variants were isolated and there were only 218 DNM clusters, this gives a frequency of 1.8% for TSMs among the clustered DNMs. Sasani *et al.* estimated that, in average, there are 70.1 de novo SNVs and 5.9 de novo indels per genome; the germline status of third-generation variants cannot be verified, but assuming the 20:1 ratio of non-germline to germline DNMs [18], the observed TSM-like patterns are likely to be somatic.

Discussion

Our analyses extend the previous works [23, 29] and show that human genomes have 230
thousands of mutations patterns consistent with the DNA-replication related template 231
switching [23] and that these loci segregate within and between populations. We studied 232
the haplotype-resolved genotype data by Ebert *et al.* [12], two independent WGS 233
experiments on the reference individual NA12878 and her extended family [30, 32], and 234
variant information in the gnomAD database [5, 39] and from a DNM study [40]. While 235
we found Ebert *et al.*'s HaplotypeSV data to have captured large numbers of TSM 236
patterns and mostly reflect the haplotypic differences between the chromosomes, the 237
call set was far from perfect and we could identify many additional TSM-like loci using 238
the independent short-read datasets. We identified de novo TSMs both in short-read 239
data and in data of Sasani *et al.*, though not able to confirm their germline status, and 240
saw consistent singleton TSM patterns in gnomAD variant data [5, 39]. Overall, our 241
results demonstrate that, despite their relatively low rate, the TSM mechanism explains 242
a significant fraction of MNVs seen in human variation data and thus contributes to 243
correlation in local mutation frequencies [20, 41]. 244

While the ability of TSMs to create reverse-complement copies of short sequence 245
fragments is highly interesting, the mechanistic origin of sequence differences is 246
irrelevant to many applications of genome analysis. There are anecdotal cases of human 247
genetic diseases caused by TSM-like mutations [42], however, and a striking finding of 248
our study was that the DNA sequencing pipelines capable of identifying SNVs may miss 249
inversions of a few tens of bases. The difficulty of detecting certain TSM patterns using 250
the standard analysis tools potentially invalidates e.g. medical genetic studies searching 251
for causative alleles behind genetic diseases. On the other, Seplyarskiy *et al.* [43] 252
recently proposed an approach to separate the signals of different mutation processes, 253
and among others, described signals from "asymmetric resolution of bulky DNA 254
damage" and from "asymmetric replication errors". Such asymmetric signal could be 255
created by template switching if the 10 times higher frequency of TSMs in leading 256
strand seen bacteria [44, 45] applies to other organisms, or if the mechanism is related 257
e.g. to coordination or clashes of transcription and replication systems [46, 47]. Local 258
TSMs have many similarities with the FoSTeS [26] and MBIR [27] mechanisms – and 259

possibly with chromoanasynthesis in cancer [48, 49] – but due to their short length are ²⁶⁰
expected to be more benign [13]. With our novel tools TSM patterns can be identified ²⁶¹
in different types of genome data, enabling analyses of their genome-wide distribution ²⁶²
and possible correlation with different cellular processes [43]. Such analyses should ²⁶³
bring us closer to understanding the mechanisms underlying the template switching. ²⁶⁴

Methods ²⁶⁵

Data   The Ebert *et al.* genotype data (vcf; freeze3) were downloaded from ²⁶⁶
`http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/HGSVC2/` ²⁶⁷
`release/v1.0/integrated_callset`, combining SNV, indel and SV calls. The ²⁶⁸
Chromium data (bam;vcf) were downloaded from ²⁶⁹
`http://ftp-trace.ncbi.nlm.nih.gov/giab/ftp/data/NA12878/10Xgenomics_` ²⁷⁰
`ChromiumGenome_LongRanger2.0_06202016` and `http://ftp-trace.ncbi.nlm.nih.` ²⁷¹
`gov/giab/ftp/technical/10Xgenomics_ChromiumGenome_LongRanger2.0_06202016.` ²⁷²
The Platinum data (bam;vcf) were downloaded from `http://ftp.1000genomes.ebi.` ²⁷³
`ac.uk/vol1/ftp/data_collections/illumina_platinum_pedigree/data/CEU`, via ²⁷⁴
`http://github.com/Illumina/PlatinumGenomes`, and from dbGaP ²⁷⁵
(`http://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=` ²⁷⁶
`phs001224.v1.p1`; authorized access for A.L.). The gnomAD data were downloaded ²⁷⁷
through `https:` ²⁷⁸
`//gnomad.broadinstitute.org/downloads#v2-multi-nucleotide-variants` and ²⁷⁹
the Sasani *et al.* data through `https://github.com/elifesciences-publications/` ²⁸⁰
`ceph-dnm-manuscript/tree/master/data`. The reference genomes were downloaded ²⁸¹
from `http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/HGSVC2/` ²⁸²
`technical/reference` (GRCh38) and from ²⁸³
`http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/technical/reference` (GRCh37 ²⁸⁴
and 38). ²⁸⁵

Analysis of haplotype data   The genotype data of Ebert *et al.* (i.e., HaplotypeSV ²⁸⁶
data) were processed with a custom python script (tsm_scan_SV2.py) internally utilising ²⁸⁷
bwa [34], samtools and bcftools [50]. The script detects clusters of variants and creates ²⁸⁸

alternative haplotypes by replacing parts of the reference genome sequence with the ²⁸⁹

variant bases. The reconstructed haplotypes (with 150 bp of flanking sequence) are then ²⁹⁰

compared to the reference allele and a TSM solution is searched, reciprocally, to create ²⁹¹

one of the alleles from the other. Each individual and chromosome was processed ²⁹²

independently. ²⁹³

The candidate loci in NA12878 were processed with another python script ²⁹⁴

(tsm_alleles2.py) that (i) creates the two alleles for each locus using the reference ²⁹⁵

sequence and the HaplotypeSV variant information, (ii) extracts the reads mapped to ²⁹⁶

the locus in a bam alignment file and maps these reads to the two alternative alleles, ²⁹⁷

and (iii) computes the mapping coverage statistics for the two alleles, recording the ²⁹⁸

mapping depth at the region differing between the alleles as well as immediately ²⁹⁹

upstream and downstream of the differing region. The REF allele was taken from the ³⁰⁰

reference genome and the ALT allele was created by placing the NA12878 variants (each ³⁰¹

haplotype separately) into a copy of that; the differing regions were then placed on ³⁰²

identical context with 500 bp of flanking sequence from the reference genome. ³⁰³

The genotype of the locus was inferred from the mapping coverage for the two

alternative alleles using a strategy similar to base calling. We required a minimum

coverage of 10 reads for upstream, downstream and within the differing region for either

of the alleles. Then the probabilities for the three possible genotypes were obtained as:

$$Pr\{G = R|R\} = (1 - \epsilon)^{d_r} \epsilon^{d_a}$$

$$Pr\{G = R/A\} = 0.5^{d_r} * 0.5^{d_a}$$

$$Pr\{G = A|A\} = \epsilon^{d_r} (1 - \epsilon)^{d_a}$$

where ϵ is the error rate of 0.01 for the read being mapped to a wrong allele, and d_r and

d_a are the mapping coverages for the REF and the ALT alleles. The inferred genotype

X, where $X \in \{R|R, R/A, A|A\}$ (standing for homozygote REF, heterozygote, and

homozygote ALT, respectively), was then the genotype with the highest probability if

that was at least 99% of the total probability:

$$Pr\{G = X\} = \max \begin{cases} Pr\{G = R|R\} \\ Pr\{G = R/A\} \\ Pr\{G = A|A\} \end{cases}$$

if for $Y = \{R|R, R/A, A|A\}$:

$$\frac{Pr\{G = X\}}{\sum Pr\{G = Y\}} > 0.99$$

.

Analysis of short-read data The FPA algorithm [23], written in `C++` and available   305
through `https://github.com/ariloytynoja/fpa`, was integrated in the SvABA tool   306
[33]. Clusters of more than one base mismatch or more than ten clipped reads within a   307
20-bp interval were identified in vcf and bam data using custom bash and awk scripts.   308
The candidate loci were targeted with SvABA-FPA and, for contigs showing two or   309
more base differences in comparison to the reference sequence, TSM solutions were   310
computed using both the REF and the ALT allele (i.e. SvABA-created contig) as the   311
ancestral type. The resulting TSM candidates were filtered and those overlapping with   312
repeat elements or low-complexity sequence, or having short length or low identity at   313
flanking sequences were removed; the bash/awk scripts for filtering are provided in the   314
supplementary data.   315

For the loci passing the filtering, the REF and ALT alleles were placed on identical   316
context with 500 bp of flanking sequence. Alignment data (in bam format) were   317
genotyped by extracting the reads and mapping them to the two alleles using the custom   318
python script. The genotype was inferred from the read coverage as explained above.   319

Analysis of CEPH 1463 data The coordinates of candidate loci were transferred   320
from GRCh38 to GRCh37 with CrossMap v0.2.9 [51] using the Ensembl chain file. The   321
same methods were used to genotype the loci.   322

Analysis of MNV data   The MNV data from gnomAD and by Sasani *et al.* were processed similarly to the HaplotypeSV data with a python script (tsm_scan_dnm.py). The script detects clusters of variants and creates alternative haplotypes by replacing parts of the reference genome sequence with the variant bases. It then compares the reconstructed haplotypes (with 150 bp of flanking sequence) to the reference allele and searches for a TSM solution, reciprocally, to create one of the alleles from the other. Tandem repeats within the ②→③ region of selected hits were searched with trf [52].

Data and code availability   All material will be deposited in Github. The python/awk scripts used, the source code for the SvABA-FPA tool, and the instructions for their use are provided in the supplementary material. With the exception of the CEPH 1643 children, all data utilised in this study are publicly available and the instructions for their download and analysis are provided in the supplementary data.

Ethics statement   A.L. had authorized access to the dbGaP genome data for the CEPH 1643 offspring (http://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs001224.v1.p1) under project name "Properties of de novo template switch mutations". The offspring data were analysed in a secure computer environment only accessible by the author, and only summary statistics of genotype inheritance are reported.

Acknowledgments

We thank CSC – IT Center for Science and the University of Helsinki IT Center for the computing resources and the secure analysis environment, and the UH Biodata Analytics Unit for helpful discussions. This work was enabled by the Academy of Finland grant #322681.

References

1. Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, et al. The sequence of the human genome. Science. 2001;291(5507):1304–1351.

2. Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, et al. Initial sequencing and analysis of the human genome. Nature. 2001;409(6822):860–921.

3. Nurk S, Koren S, Rhie A, Rautiainen M, Bzikadze AV, Mikheenko A, et al. The complete sequence of a human genome; 2021.

4. Aganezov S, Yan SM, Soto DC, Kirsche M, Zarate S, Avdeyev P, et al. A complete reference genome improves analysis of human genetic variation; 2021.

5. Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, Wang Q, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. Nature. 2020;581(7809):434–443.

6. Taliun D, Harris DN, Kessler MD, Carlson J, Szpiech ZA, Torres R, et al. Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. Nature. 2021;590(7845):290–299.

7. Miga KH, Wang T. The Need for a Human Pangenome Reference Sequence. Annu Rev Genomics Hum Genet. 2021;.

8. Bentley DR, Balasubramanian S, Swerdlow HP, Smith GP, Milton J, Brown CG, et al. Accurate whole human genome sequencing using reversible terminator chemistry. Nature. 2008;456(7218):53–59.

9. Porubsky D, Ebert P, Audano PA, Vollger MR, Harvey WT, Marijon P, et al. Fully phased human genome assembly without parental data using single-cell strand sequencing and long reads. Nat Biotechnol. 2021;39(3):302–308.

10. Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, et al. Real-time DNA sequencing from single polymerase molecules. Science. 2009;323(5910):133–138.

11. Falconer E, Lansdorp PM. Strand-seq: a unifying tool for studies of chromosome segregation. Semin Cell Dev Biol. 2013;24(8-9):643–652.

12. Ebert P, Audano PA, Zhu Q, Rodriguez-Martin B, Porubsky D, Bonder MJ, et al. Haplotype-resolved diverse human genomes and integrated analysis of structural variation. Science. 2021;372(6537).

13. Zhang F, Gu W, Hurles ME, Lupski JR. Copy number variation in human health, disease, and evolution. Annu Rev Genomics Hum Genet. 2009;10:451–481.

14. Weischenfeldt J, Symmons O, Spitz F, Korbel JO. Phenotypic impact of genomic structural variation: insights from and for human disease. Nat Rev Genet. 2013;14(2):125–138.

15. Sakamoto Y, Sereewattanawoot S, Suzuki A. A new era of long-read sequencing for cancer genomics. J Hum Genet. 2020;65(1):3–10.

16. Perry GH, Dominy NJ, Claw KG, Lee AS, Fiegler H, Redon R, et al. Diet and the evolution of human amylase gene copy number variation. Nat Genet. 2007;39(10):1256–1260.

17. Yan SM, Sherman RM, Taylor DJ, Nair DR, Bortvin AN, Schatz MC, et al. Local adaptation and archaic introgression shape global diversity at human structural variant loci; 2021.

18. Conrad DF, Keebler JEM, DePristo MA, Lindsay SJ, Zhang Y, Casals F, et al. Variation in genome-wide mutation rates within and between human families. Nat Genet. 2011;43(7):712–714.

19. Veltman JA, Brunner HG. De novo mutations in human genetic disease. Nat Rev Genet. 2012;13(8):565–575.

20. Ségurel L, Wyman MJ, Przeworski M. Determinants of mutation rate variation in the human germline. Annu Rev Genomics Hum Genet. 2014;15:47–70.

21. International Human Genome Sequencing Consortium. Finishing the euchromatic sequence of the human genome. Nature. 2004;431(7011):931–945.

22. Levy S, Sutton G, Ng PC, Feuk L, Halpern AL, Walenz BP, et al. The diploid genome sequence of an individual human. PLoS Biol. 2007;5(10):e254.

23. Löytynoja A, Goldman N. Short template switch events explain mutation clusters in the human genome. Genome Res. 2017;27(6):1039–1049.

24. Ripley LS. Model for the participation of quasi-palindromic DNA sequences in frameshift mutation. Proc Natl Acad Sci U S A. 1982;79:4128–4132.

25. Dutra BE, Lovett ST. Cis and trans-acting effects on a mutational hotspot involving a replication template switch. J Mol Biol. 2006;356:300–311.

26. Lee JA, Carvalho CMB, Lupski JR. A DNA replication mechanism for generating nonrecurrent rearrangements associated with genomic disorders. Cell. 2007;131(7):1235–1247.

27. Hastings PJ, Ira G, Lupski JR. A microhomology-mediated break-induced replication model for the origin of human copy number variation. PLoS Genet. 2009;5(1):e1000327.

28. Besenbacher S, Sulem P, Helgason A, Helgason H, Kristjansson H, Jonasdottir A, et al. Multi-nucleotide de novo Mutations in Humans. PLoS Genet. 2016;12(11):e1006315.

29. Walker CR, Scally A, De Maio N, Goldman N. Short-range template switching in great ape genomes explored using pair hidden Markov models. PLoS Genet. 2021;17(3):e1009221.

30. Eberle MA, Fritzilas E, Krusche P, Källberg M, Moore BL, Bekritsky MA, et al. A reference data set of 5.4 million phased human variants validated by genetic inheritance from sequencing a three-generation 17-member pedigree. Genome Res. 2017;27(1):157–164.

31. Zook JM, Catoe D, McDaniel J, Vang L, Spies N, Sidow A, et al. Extensive sequencing of seven human genomes to characterize benchmark reference materials. Sci Data. 2016;3:160025.

32. Weisenfeld NI, Kumar V, Shah P, Church DM, Jaffe DB. Direct determination of diploid genome sequences. Genome Res. 2017;27(5):757–767.

33. Wala JA, Bandopadhayay P, Greenwald NF, O'Rourke R, Sharpe T, Stewart C, et al. SvABA: genome-wide detection of structural variants and indels by local assembly. Genome Res. 2018;28(4):581–591.

34. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv. 2013;.

35. Marks P, Garcia S, Barrio AM, Belhocine K, Bernate J, Bharadwaj R, et al. Resolving the full spectrum of human genome variation using Linked-Reads. Genome Res. 2019;29(4):635–645.

36. Shirley MD, Baugher JD, Stevens EL, Tang Z, Gerry N, Beiswanger CM, et al. Chromosomal variation in lymphoblastoid cell lines. Hum Mutat. 2012;33(7):1075–1086.

37. Sherry ST, Ward MH, Kholodov M, Baker J, Phan L, Smigielski EM, et al. dbSNP: the NCBI database of genetic variation. Nucleic Acids Res. 2001;29(1):308–311.

38. Choi Y, Chan AP, Kirkness E, Telenti A, Schork NJ. Comparison of phasing strategies for whole human genomes. PLoS Genet. 2018;14(4):e1007308.

39. Wang Q, Pierce-Hoffman E, Cummings BB, Alföldi J, Francioli LC, Gauthier LD, et al. Landscape of multi-nucleotide variants in 125,748 human exomes and 15,708 genomes. Nat Commun. 2020;11(1):2539.

40. Sasani TA, Pedersen BS, Gao Z, Baird L, Przeworski M, Jorde LB, et al. Large, three-generation human families reveal post-zygotic mosaicism and variability in germline mutation accumulation. Elife. 2019;8.

41. Harris K, Nielsen R. Error-prone polymerase activity causes multinucleotide mutations in humans. Genome Res. 2014;24(9):1445–1454.

42. Menardi C, Schneider R, Neuschmid-Kaspar F, Klocker H, Hirsch-Kauffmann M, Auer B, et al. Human APRT deficiency: Indication for multiple origins of the most common Caucasian mutation and detection of a novel type of mutation involving intrastrand-templated repair. Hum Mutat. 1997;10:251–255.

43. Seplyarskiy VB, Soldatov RA, Koch E, McGinty RJ, Goldmann JM, Hernandez RD, et al. Population sequencing data reveal a compendium of mutational processes in the human germ line. Science. 2021;.

44. Rosche WA, Trinh TQ, Sinden RR. Leading strand specific spontaneous mutation corrects a quasipalindrome by an intermolecular strand switch mechanism. J Mol Biol. 1997;269(2):176–187.

45. Seier T, Padgett DR, Zilberberg G, Sutera VA Jr, Toha N, Lovett ST. Insights into mutagenesis using Escherichia coli chromosomal lacZ strains that enable detection of a wide spectrum of mutational events. Genetics. 2011;188(2):247–262.

46. Hamperl S, Bocek MJ, Saldivar JC, Swigut T, Cimprich KA. Transcription-Replication Conflict Orientation Modulates R-Loop Levels and Activates Distinct DNA Damage Responses. Cell. 2017;170(4):774–786.e19.

47. Chen YH, Keegan S, Kahli M, Tonzi P, Fenyö D, Huang TT, et al. Transcription shapes DNA replication initiation and termination in human cells. Nat Struct Mol Biol. 2019;26(1):67–77.

48. Liu P, Erez A, Nagamani SCS, Dhar SU, Kołodziejska KE, Dharmadhikari AV, et al. Chromosome catastrophes involve replication mechanisms generating complex genomic rearrangements. Cell. 2011;146(6):889–903.

49. Holland AJ, Cleveland DW. Chromoanagenesis and cancer: mechanisms and consequences of localized, complex chromosomal rearrangements. Nat Med. 2012;18(11):1630–1638.

50. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of SAMtools and BCFtools. GigaScience. 2021;10(2). doi:10.1093/gigascience/giab008.

51. Zhao H, Sun Z, Wang J, Huang H, Kocher JP, Wang L. CrossMap: a versatile tool for coordinate conversion between genome assemblies. Bioinformatics. 2014;30(7):1006–1007.

52. Benson G. Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Res. 1999;27(2):573–580.

53. Lorenz R, Bernhart SH, Höner Zu Siederdissen C, Tafer H, Flamm C, Stadler PF, et al. ViennaRNA Package 2.0. Algorithms Mol Biol. 2011;6:26.

54. Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. Brief Bioinform. 2013;14(2):178–192.

55. Hahne F, Ivanek R. Visualizing Genomic Data Using Gviz and Bioconductor. In: Mathé E, Davis S, editors. Statistical Genomics: Methods and Protocols. New York, NY: Springer New York; 2016. p. 335–351.

Fig S1. Classical vs. four-point model model of template switching. (**A**) In the classical model of Ripley [24], the replication (blue arrow) switches the template and starts going backwards either along the newly synthesized strand (intra) or along the opposite strand (inter). After a short while, the replication switches back to the original strand (return) and continues normally. As the result, the near-perfect inverted repeat (red dashed arrows) have been converted into a perfect inverted repeat (red arrows), causing mismatches between the sequences (bulge). (**B**) The four-point model of Löytynoja and Goldman [23] is a computationally tractable representation of the switch process and relaxes some of the assumptions underlying the Ripley model. The model assumes four points, ① - ④, where the replication process changes from one template to another, and copying of the fragment ②→③ in reverse-complement manner. The outcome of the switch process depends on the relative positions of points ①-④, and the type 3-2-1-4 (read from left to right) shown here is consistent with the process shown in a and can take place either intra- or inter-strand. See Löytynoja and Goldman [23] for enumeration and outcomes of all possible switch point combinations.

Fig S2. Direction of TSMs. (**A**) The complex mutation (top) is partially explained by the TSM mechanism when the REF allele is used as the reference (middle); the solution (bottom) contains two mismatches (marked with asterisks). (**B**) With the ALT allele as the reference, the complex mutation is fully explained, indicating that ALT is the ancestral state and REF is a derived state. (**C,D**) Inversions in place cannot be polarised as both arrangements fully explain the complex mutation pattern with a single TSM event.

A

Forward alignment:

```
qry TATGCTCCCTTATAATCTTATGGGACCATCATATATATATATA|TATATA----TATaTaTGTGTGTGTGTGTGTGT|GTGTGTGTGTGTGTGTGTATATGTATATATGTGTATATATATACATATATGTATATATATACACACACACACACATATATAT
ref TATGCTCCCTTATAATCTTATGGGACCATCATATATATATATA|TATATATGTGTATGTGTGTGTGTGTGTGTGTGT|GTGTGTGTGTGTGTGTGTATATGTATATATGTGTATATATATACATATATGTATATATATACACACACACACACATATATAT
```

Switch process:

```
F1: TATGCTCCCTTATAATCTTATGGGACCATCATATATATATATA
F3:                                            GTGTGTGTGTGTGTGTATATGTATATATGTGTATATATATACATATATGTATATATATACACACACACACACATATATATATATAGTGCATGACTGTATAATAAA
RF: TATGCTCCCTTATAATCTTATGGGACCATCATATATATATATATATATATGTGTATGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTATATGTATATATGTGTATATATATACATATATGTATATATATACACACACACACACATATATATATATAGTGCATGACTGTATAATAAA
RR: ATACGAGGGAATATTAGAATACCCTGGTAGTATATATATATATATATATATACACATACACACACACACACACACACACACACACAGACATATACATATATACACATATATATATGTGTATACATATATATGTGTGTGTGTGTGTGTATATATATATATATCACGTACTGACATATTATTT
F2:                                                                                                                                                     TGTGTGTGTGTGTGTGTATATATATATAT
```

Template-switch alignment:

```
qry TATGCTCCCTTATAATCTTATGGGACCATCATATATATATATA|TATATATATATATGTGTGTGTGTGTGTGT|GTGTGTGTGTGTGTGTGTATATGTATATATGTGTATATATATACATATATGTATATATATACACACACACACACATATATAT
ref TATGCTCCCTTATAATCTTATGGGACCATCATATATATATATA|TATATATATATATGTGTGTGTGTGTGT|GTGTGTGTGTGTGTGTGTATATGTATATATGTGTATATATATACATATATGTATATATATACACACACACACACATATATAT
```

B

Fig S3. Changes in a VNTR locus consistent with the TMS process. (**A**) The differences between the alternative allele (qry) and the GRCh38 copy (ref; top) at chr1:116324126-116324132 can be explained by template switching (middle, bottom). The same outcome could be obtained with two slippage mutations, however. (**B**) DNA secondary structure reveals that, in addition to runs of internally reverse-complement ATs, the locus also has runs of pairwise reverse-complementary GTs and ACs (red shading). Presence of such pairwise reverse-complementary VNTRs in the same locus can be explained by a historical template switching event. The secondary structure was inferred with RNAfold [53].

Fig S4. Lengths of inferred TMS events. A great majority of inferred TSM events are 10 bp or shorter in length. Despite the higher numbers of TSM loci identified in African samples, the length distribution of TSM events in individuals of African (NA19238) and European (NA12878) origin are similar. The inset shows lower values. Cases longer than 50 bp are grouped.

Fig S5. Inconsistency among different short-read datasets. (**A**) The TSM solution explaining a cluster of three base differences (and not explaining one nearby difference). (**B**) The Platinum data have one supporting read (left; red box) whereas the Chromium data have none; an independent dataset has multiple supporting reads (red box) but these are associated with flanking changes (blue box) creating more than two haplotypes. A plausible explanation is mismapping of reads from another locus. Data were visualised with IGV [54].

chr1:248458096-248458132

Fig S6. A heterozygous TSM event identified in NA12878 with short-read data. (**A**) Platinum data (left) and (**B**) Chromium data (left) have excess of soft clips (bottom, blue bars) and atypical patterns in the mapping coverage (top, in gray). (**C**) Two variants were called in this region by Ebert *et al.* [12] but these fail to explain the haplotype differences. (**D**) De novo assembly of the reads creates two locally highly dissimilar haplotypes (top). All but one difference can be explained with a TSM event, an inversion in place (middle). Using two haplotypes with alternative central parts (blue and magenta; bottom) as the reference, extracted reads map in full length (**A,B**; middle, right) with roughly even coverages. Data were visualised with Gviz [55].

Fig S7. A homozygous TSM event identified in NA12878 with short-read data. (**A**) Platinum data (left) and (**B**) Chromium data (left) have excess of soft clips (bottom, blue bars) and atypical patterns in the mapping coverage (top, in gray). (**C**) Fifteen variants fully explaining the differences were called in the Chromium data while no variants were indicated in this region in the Platinum data or by Ebert *et al.* [12] (**D**) De novo assembly of the reads creates two locally highly dissimilar haplotypes (top). All differences can be explained with a TSM event, an inversion in place (middle). Using two haplotypes with alternative central parts (blue and magenta; bottom) as the reference, extracted reads map in full length on the ALT haplotype only (**A,B**; middle, right). Data were visualised with Gviz [55].

chr1:170589067-170589150

Fig S8. A potential de novo TSM event identified in NA12878 Chromium short-read data. (**A**) Platinum data (left) show no signal whereas (**B**) Chromium data (left) have some soft clips (bottom, blue bars) in this region. (**C**) One or two SNVs were called in this region by different datasets. (**D**) De novo assembly of Chromium reads creates two locally highly dissimilar haplotypes (top). All differences can be explained with a TSM event (middle). Using two haplotypes with alternative central parts (blue and magenta; bottom) as the reference, a fraction of Chromium reads map in full length on the ALT haplotype (**B**; middle, right). Data were visualised with Gviz [55].

A

```
CHR  START      END        ID           REF, ALT(s)

chr1 248458100 248458100 rs1294550520  A,  C
chr1 248458100 248458126 rs1553320400  AGTGGTTTAACTATTTTAAAATTGGAG,  AG
chr1 248458101 248458101 rs945532251   G,  C
chr1 248458102 248458101 rs370077035   -,  C,  CTCCAATTTTAAAA
chr1 248458102 248458104 rs1558344009  TGG,  CTCCAA
chr1 248458102 248458109 rs774785163   TGGTTTAA,  -
chr1 248458102 248458126 rs766858754   TGGTTTAACTATTTTAAAATTGGAG,  CTCCAATTTTAAAATAGTTAAATCA
chr1 248458103 248458103 rs55778336    G,  C,  T
chr1 248458103 248458108 rs1312056438  GGTTTA,  -
chr1 248458104 248458104 rs11488140    G,  A,  C
chr1 248458105 248458104 rs75431795    -,  AA
chr1 248458108 248458108 rs61834525    A,  T
chr1 248458108 248458111 rs1558344025  AACT,  TAAA
chr1 248458109 248458109 rs1325281402  A,  C
chr1 248458110 248458110 rs78430540    C,  A
chr1 248458110 248458110 rs1283571306  C,  -
chr1 248458111 248458111 rs79278902    T,  A
chr1 248458112 248458111 rs759219409   -,  CCA,  TCA
chr1 248458112 248458112 rs1217534201  A,  C,  T
chr1 248458112 248458112 rs1408812771  A,  -
chr1 248458114 248458114 rs77716920    T,  A
chr1 248458114 248458117 rs1558344038  TTTA,  AGTT
chr1 248458115 248458115 rs78627666    T,  G
chr1 248458117 248458117 rs61834526    A,  T
chr1 248458119 248458119 rs1259249878  A,  C
chr1 248458120 248458120 rs1442840868  A,  G
chr1 248458122 248458122 rs56040868    T,  C
chr1 248458122 248458124 rs752225071   TGG,  -
chr1 248458122 248458126 rs386642020   TGGAG,  AGTTAAATCA,  CA
chr1 248458123 248458124 rs58657081    GG,  -
chr1 248458124 248458124 rs1042592330  G,  A,  C
chr1 248458126 248458126 rs866296781   G,  A
chr1 248458126 248458126 rs58743262    G,  -
chr1 248458127 248458126 rs373041964   -,  TTAAACCA,  TTAAATCA
chr1 248458127 248458128 rs1553320408  CT,  CTCCAATTTTAAAATAGTTAAATCACT
chr1 248458129 248458129 rs1003776664  T,  C
chr1 248458130 248458130 rs1208388863  A,  G
```

Alignment:

```
REF GAAGCAAAAAAACACCAAAAAAACTAAAATTATTTCCAGCTTCACTGATAAGTGGTTTAACTATTTTAAAATTGGAGCTTATGTAAAAAAGTCATTGGCTTTTGCTAGATTCTTCATATATTCCTTC
ALT GAAGCAAAAAAACACCAAAAAAACTAAAATTATTTCCAGCTTCACTGATAAGCTCCAATTTTAAAATAGTTAAATCACTTATGTAAAAAAGTCATTGGCTTTTGCTAGATTCTTCATATATTCCTTC
```

Switch process:

```
                                                              ①                              ④
F1: GAAGCAAAAAAACACCAAAAAAACTAAAATTATTTCCAGCTTCACTG··············>GTAAAAAAGTCATTGGCTTTTGCTAGATTCTTCATATATTCCTTC
F3:
RF: GAAGCAAAAAAACACCAAAAAAACTAAAATTATTTCCAGCTTCACTGATGATAAGCAAATTTTAAAATAGTTAAATCACTGTAAAAAAGTCATTGGCTTTTGCTAGATTCTTCATATATTCCTTC
F2:                           TATTCGAGGTTAAAATTTTATCAATTTGGTGAATA<
                                                              ③                              ②
```

Template-switch alignment:

```
REF GAAGCAAAAAAACACCAAAAAAACTAAAATTATTTCCAGCTTCACTG|ATAAGTGgTTTAACTATTTTAAAATTGGAGCTTAT|GTAAAAAAGTCATTGGCTTTTGCTAGATTCTTCATATATTCCTTC
ALT GAAGCAAAAAAACACCAAAAAAACTAAAATTATTTCCAGCTTCACTG|ATAAGTGATTTAACTATTTTAAAATTGGAGCTTAT|GTAAAAAAGTCATTGGCTTTTGCTAGATTCTTCATATATTCCTTC
```

B

```
CHR  START      END        ID           REF, ALT(s)

chr1 248437366 248437371 rs1282560120  TTACTT,  TT
chr1 248437373 248437383 rs1362375318  AGTTGTGGAAG,  AG
chr1 248437375 248437375 rs1403073841  T,  A,  C
chr1 248437376 248437376 rs1282373520  T,  G
chr1 248437378 248437378 rs1451470978  T,  C
chr1 248437381 248437382 rs1379895247  AA,  A
chr1 248437388 248437387 rs879403957   -,  TG
chr1 248437389 248437390 rs1234910043  CC,  -
chr1 248437390 248437391 rs879504096   CA,  -
chr1 248437396 248437397 rs369794123   CT,  CTTCCACAACT
chr1 248437398 248437397 rs1205190376  -,  TCC
chr1 248437399 248437398 rs1287619838  -,  CAACTG
```

Alignment:

```
REF AATCAAAACTTCTCTATCGTTCTAAAGATTCCCCAAGTGATTTTACTTTAGTTGTGGAAGAAAA--GCCATTTTCT---------AGAAGTGACACTTCATGTGAAAATACTGCTGGGGAGGAAACATAATGA
ALT AATCAAAACTTCTCTATCGTTCTAAAGATTCCCCAAGTGATTTTACTTTAG---------AAAATGGC--TTTTCTTCCACAACTAGAAGTGACACTTCATGTGAAAATACTGCTGGGGAGGAAACATAATGA
```

Switch process:

```
                                                          ①                     ④
F1: AATCAAAACTTCTCTATCGTTCTAAAGATTCCCCAAGTGATTTTACTT··········>GAAGTGACACTTCATGTGAAAATACTGCTGGGGAGGAAACATAATGA
F3:
RF: AATCAAAACTTCTCTATCGTTCTAAAGATTCCCCAAGTGATTTTACTTTAGAAAATGGCTTTTCTTCCACAACTAGAAGTGACACTTCATGTGAAAATACTGCTGGGGAGGAAACATAATGA
F2:                                    ATCTTTTACCGAAAAGAAGGTGTTGAT<
                                                          ③                     ②
```

Template-switch alignment:

```
REF AATCAAAACTTCTCTATCGTTCTAAAGATTCCCCAAGTGATTTTACTT|TAGTTGTGGAAGAAAAGCCATTTTCTA|GAAGTGACACTTCATGTGAAAATACTGCTGGGGAGGAAACATAATGA
ALT AATCAAAACTTCTCTATCGTTCTAAAGATTCCCCAAGTGATTTTACTT|TAGTTGTGGAAGAAAAGCCATTTTCTA|GAAGTGACACTTCATGTGAAAATACTGCTGGGGAGGAAACATAATGA
```

Fig S9. Description of TSM variants in the dbSNP database. (**A**) TSM locus is described in full with a MNV of 25 bp (cyan) but the same changes are also given as multiple SNVs and indels. (**B**) TSM haplotypes can be reconstructed from four different variants (cyan, magenta) but the region has also additional variants annotated.

Fig S10. The three longest TSM events explaining gnomAD MNVs. (**A**) The longest TSM event within complex sequence is 90 bp long and explains a MNV of four base differences. (**B,C**) The longest TSM events involving VNTRs, 119 and 125 bp in length, explain MNVs of five and four base differences.

Fig S11. Large gnomAD MNV explained by two TSMs. (**A**) A cluster of 19 base changes in gnomAD MNV data. (**B**) Of the 19 base differences (top), 16 can be explained with two TSM events (middle, bottom). The DNA replication proceeds from left to right (green); at ①, the replication jumps to ② in the opposite strand and proceeds in reverse-complement manner until ③ (blue); it then jumps to ④ in the original template and proceeds until ⑤ (magenta); the replication again changes the template (either to the opposite strand or to the newly synthesized strand), jumps to ⑥ and continues until in reverse-complement manner until ⑦ (orange); the replication makes a final jump to ⑧ in the original template and continues normally (red).

Fig S12. De novo MNVs explainable with the TSM mechanism. (**A**) Four MNVs in the third-generation individuals of Sasani *et al.* [40] that can be explained with a single TSM. (**B**) A TSM solution that can explain two of the three clustered de novo SNVs in a third-generation individual of Sasani *et al.* [40]

Table S1. Number and length of inferred TSM events as well as proportion of homozygous loci and resolved and ancestral (of former) allelic state in different samples. Only unmasked loci are included.

ID	Population	Region	n	Length (bp) average	Length (bp) maxim.	Allele (%) homozyg.	Allele (%) singleton	Polarization (%) resolved	Polarization (%) ancestral
HG02011	ACB	AFR	1046	10.67	171	30.50	7.58	89.50	49.68
NA19983	ASW	AFR	1077	10.77	171	27.48	8.41	88.03	53.74
HG03125	ESN	AFR	1073	10.43	171	29.45	7.12	88.01	51.32
HG03371	ESN	AFR	1070	10.44	171	31.40	8.43	89.09	52.57
HG02587	GWD	AFR	1097	10.81	180	26.71	9.59	89.44	51.73
HG02818	GWD	AFR	1038	10.64	100	29.38	8.07	88.07	53.01
HG03065	MSL	AFR	1060	10.48	171	30.94	8.81	89.84	53.09
HG03486	MSL	AFR	1064	10.90	180	28.85	8.39	88.17	52.08
NA19238	YRI	AFR	1121	10.44	97	28.37	8.60	90.29	51.28
NA19239	YRI	AFR	1092	10.47	171	29.30	8.91	89.03	52.46
NA12329	CEU	EUR	922	10.70	180	35.03	3.07	88.84	52.32
NA12878	CEU	EUR	921	10.49	171	36.48	2.61	89.48	54.42
HG00171	FIN	EUR	883	10.28	171	34.65	3.63	90.05	54.27
HG00096	GBR	EUR	937	10.67	171	36.07	3.11	89.23	53.17
HG01505	IBS	EUR	920	10.79	180	32.83	4.03	89.46	52.61
NA20509	TSI	EUR	882	10.47	180	35.49	4.52	89.12	55.34
NA24385	AZK	EUR	872	10.48	171	36.24	2.71	89.58	54.86
HG00864	CDX	EAS	942	10.61	171	39.17	4.33	88.67	50.42
NA18534	CHB	EAS	923	10.59	171	39.98	4.34	87.99	52.15
HG00512	CHS	EAS	917	10.52	171	39.91	2.82	89.12	50.67
HG00513	CHS	EAS	929	10.69	180	41.87	3.57	88.41	50.24
NA18939	JPT	EAS	927	10.50	171	42.29	3.16	89.76	50.30
HG01596	KHV	EAS	910	10.78	171	39.01	4.12	88.50	50.00
HG03009	BEB	SAS	878	10.45	171	35.76	3.85	88.52	52.76
NA20847	GIH	SAS	938	10.55	171	35.82	4.59	89.15	53.46
HG03732	ITU	SAS	920	10.37	171	35.87	3.57	88.15	52.77
HG02492	PJL	SAS	879	10.80	180	39.59	3.14	87.94	52.39
HG03683	STU	SAS	939	10.22	97	33.97	4.06	89.80	50.89
HG01114	CLM	AMR	923	10.64	171	33.69	3.86	87.66	53.83
NA19650	MXL	AMR	906	10.58	171	35.76	3.43	88.41	52.93
HG00731	PUR	AMR	949	10.63	171	34.56	3.09	88.55	52.91
HG00732	PUR	AMR	955	10.60	171	34.87	3.23	88.20	51.36

Table S2. Genotypes obtained with different datasets for variable TSM loci in NA12878 HaplotypeSV data.

		Platinum			
HaplotypeSV	Chromium	R|R	R/A	A|A	NA
NA	R|R	2	0	0	0
NA	R/A	0	71	1	3
NA	A|A	0	2	4	0
NA	NA	0	0	0	1
R/A	R|R	6	1	0	0
R/A	R/A	5	425	0	0
R/A	A|A	0	0	2	0
R/A	NA	1	1	0	0
A|A	R|R	2	0	0	0
A|A	R/A	0	9	7	0
A|A	A|A	0	3	310	3
A|A	NA	0	1	4	0

Table S3. Parental genotypes for HaplotypeSV TSM loci confirmed variable within NA12878 Chromium data.

| | | | NA12892 | | |
NA12878	NA12891	R|R	R/A	A|A	NA
R/A	R|R	1	80	35	4
R/A	R/A	65	113	53	14
R/A	A|A	39	42	2	2
R/A	NA	1	20	3	2
A|A	R|R	0	0	0	0
A|A	R/A	0	42	46	1
A|A	A|A	0	43	159	4
A|A	NA	0	3	6	3