Pharmacogenetic variation in Neanderthals and Denisovans and implications for human health and response to medications

Tadeusz H. Wroblewski,[1,6] Kelsey E. Witt,[2,6] Seung-been Lee,[3] Ripan S. Malhi,[4] Emilia Huerta-Sanchez,[2] Fernando Villanea,[5,7]* Katrina G. Claw[1,7]**

Affiliations

[1] Department of Biomedical Informatics, Colorado Center for Personalized Medicine, University of Colorado Anschutz Medical Campus, Aurora, CO, USA

[2] Ecology and Evolutionary Biology and Center for Computational and Molecular Biology, Brown University, Providence, RI, USA

[3] Precision Medicine Institute, Macrogen Inc., Seoul, Republic of Korea

[4] Department of Anthropology, Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign, USA

[5] Department of Anthropology, University of Colorado Boulder, Boulder CO, USA

[6] These authors contributed equally to this work

[7] Co-corresponding authors

Correspondence

* Fernando.Villanea@colorado.edu

** Katrina.Claw@cuanschutz.edu

Abstract

Modern humans carry both Neanderthal and Denisovan (archaic) genome elements that are part

of the human gene pool and affect the life and health of living individuals. The impact of archaic

DNA may be particularly evident in pharmacogenes – genes responsible for the processing of

exogenous substances such as food, pollutants, and medications. However, the health

implications and contribution of archaic ancestry in pharmacogenes of modern humans remains

understudied. Here, we explore eleven key cytochrome P450 genes (*CYP450*) involved in drug

metabolizing reactions in three Neanderthal and one Denisovan individuals as well as examine

archaic introgression in modern human populations. We infer the metabolizing efficiency of

these eleven *CYP450* genes in archaic individuals and find important phenotypic differences

relative to modern human variants. We identify several single nucleotide variants shared between

archaic and modern humans in each gene, including some potentially function-altering mutations

in archaic *CYP450* genes, which may result in altered metabolism in living people carrying these

variants. We highlight three genes which show evidence for archaic introgression into modern

humans, as well as one additional gene that shows evidence for a gene duplication found only in

Neanderthals and modern Africans.

Keywords: pharmacogenetics, archaic, genetics, ancient DNA

Introduction

The cytochrome P450 (*CYP450*) genes encode oxidase enzymes that function in metabolism of endogenous small molecules and in detoxification of exogenous (or xenobiotic) compounds. This gene family is present in all mammals, with 57 active genes and 58 pseudogenes coding for CYP450 enzymes in humans [1]. These xenobiotic-substrate enzymes have been studied extensively because of their roles in the absorption, distribution, metabolism, and excretion (ADME) of pharmaceuticals and drug development. The evolution of xenobiotic CYP450 enzymes may be driven by organisms' need to metabolically detoxify foreign compounds - often toxic chemicals produced by plants, fungi, and bacteria - in the local environment. The *CYP450* genes show evidence of positive selection and high allele frequency variation in humans [1,2]. It has been suggested that the shift from hunting and gathering to food production in humans may have profoundly changed the selective effect of certain CYP450 enzymes [2,3]. Although these genes have been studied extensively in humans and other mammals, the extent of genetic variation in archaic individuals and its relation to modern human variation has not been addressed. Genetic variation in ADME genes varies extensively across modern human populations, and identifying alleles inherited through archaic introgression may be informative about the origin of specific pharmacogenetic (i.e., genes that modulate the body's response to pharmaceuticals; PGx) variants and their phenotypes, including metabolizer status.

Throughout human evolutionary history, our species has adapted to numerous distinct and varied challenging environments, which present the need to metabolize new xenobiotic substances, such as food, pollutants, and, most recently, medications [4]. Modern humans encountered new environmental challenges as they dispersed first throughout and then outside of the African continent, but they also encountered other hominin species already adapted to those

novel environments (i.e., Neanderthals and Denisovans) [5,6]. The direct sequencing of multiple

Neanderthal and Denisovan genomes has revealed a complex history of admixture between these

archaic humans and the ancestors of modern humans [7,8]. Most modern humans carry a small but

significant portion of archaic ancestry, which has been targeted by natural selection [9]. Purifying

selection — or negative natural selection — has removed many archaic variants in functional

genomic regions [10–12], while some archaic variants may have been lost through genetic

bottlenecks or drift in modern human populations. In spite of selective pressures, there are still

functional regions for which archaic variants are found at very high frequency in living humans

through the effects of positive natural selection [13–16]. One of the most well-known examples of

adaptation through archaic admixture is high-altitude adaptation in Tibetans, a population in

which the Denisovan variant of the gene *EPAS1* is found at extremely high frequency, and has

conferred them with resistance to hypoxic stress in high-altitude environments [14].

The *CYP450* genes offer a promising target for adaptation through archaic introgression

as humans encountered novel exogenous substances as they expanded their range. Neanderthals

and Denisovans may have possessed *CYP450* variants fine-tuned to metabolizing substances

found in their native habitats of Eurasia and Siberia. As modern humans moved outside of

Africa, they likely faced novel environmental factors which may have influenced selective

pressures; therefore, advantageous archaic *CYP450* variants introduced to modern humans

through admixture may have been retained in the modern human gene pool by natural selection.

Here, we examine genetic variation and predict metabolizer phenotypes in eleven

CYP450 genes in three Neanderthal and one Denisovan individuals using publicly available

high-resolution whole genome sequencing data. These genes represent enzymes which are

responsible for up to 75% of the metabolism of commonly prescribed drugs and other xenobiotic

substrates [17–20], which could have implications to modern human health, disease, and drug safety

and efficacy. In addition, we examine the patterns of allele sharing between archaic and modern

humans to identify haplotypes that may have been introgressed. This study aims to increase our

understanding of introgressed genetic variants and the different selective regimes and

environments acting on these enzymes.

Subjects and Methods

Samples

We investigate population-specific *CYP450* gene variation by combining data from the

1000 Genomes Project [21], the Simons Genome Diversity Project [22], the complete Neanderthal

genomes [23–25], and the Denisovan genome [26]. We extracted coding variation from four archaic

human genomes: the Denisovan individual from the Denisova cave in the Altai Mountains

(~21X coverage), a Neanderthal individual from Croatia (~42X coverage), and two Neanderthal

individuals from the Altai Mountains: one from the Denisova cave (~30X coverage), and one

from the Chagyrskaya cave (~28x coverage). These individuals are estimated to be at least

50,000 years old. We additionally utilized a set of seventy published chimpanzee genomes [27],

which were used as a proxy for variation that is ancestral to modern and archaic humans.

Variant Calling

Sequencing data for the three Neanderthal individuals and one Denisovan individual are

publicly available with alignment and processing of the BAM files previously outlined [23–26]. The

BAM files were used to calculate depth of coverage in the sample's genomic regions (see section

Star Allele Calling), and then processed for variant calling. *De novo* variant calling was

performed from BAM files instead of using the previously generated variant call files, which

were likely more stringent and may have removed heterozygous sites. Sample BAM files were

separated by chromosome for easier downstream processing using samtools (version 1.10 with

hts lib 1.10) 'view' function [28]. Variant calling was performed for each *CYP450* gene of interest,

using PyPGx (version 0.1.37) 'bam2vcf' which implements the Genome Analysis Toolkit

(GATK, version 4.1.9.0) 'HaplotypeCaller' function (options -emit-ref-confidence GVCF; -

minimum-mapping-quality 10) followed by GATK 'GenotypeGVCF' to merge individual

samples [29]. Variants were called against the HumanG1Kv37 hg19 reference assembly. GATK

'VariantFiltration' was additionally utilized to annotate variants with a quality score of 50.

Variants from modern human individuals from the 1000 Genomes Project [21] included for further

analysis were called using the same procedure.

For individual SNV analysis, variant call format (VCF) files were further filtered by the

phred-scaled quality score (QUAL) > 40 and gene regions were determined as the RefSeq start

and end coordinates of the gene with 2000 base pairs upstream to account for the promoter

region using bcftools (version 1.10.2 with hts lib 1.10.2) [28,30]. Variant read depth was calculated

with the PyPGx (version 0.14.0, https://pypgx.readthedocs.io/en/latest/readme.html) pipeline

(described below). Variants were additionally filtered for polymorphic sites that had a BAM read

depth ≤ 1 for one allele or a ratio of minor allelic read depth to major allelic read depth < 0.2.

Variant annotation was conducted through ANNOVAR (-protocol refGeneWithVer,

knownGene, avsnp150, ljb26_pp2hvar) which utilizes dbSNP150, RefSeq, and the UCSC

genome browser to identify and characterize known variants; the occurrence of novel SNVs were

additionally confirmed with Gnomad (v2.1.1) [30–34]. Variant locations were identified with

ANNOVAR, which annotates gene location based on RefSeq and UCSC genome browser data.

Variants are defined as exonic, splicing, ncRNA, 3' and 5'UTR, intronic, or in the promoter

region. If the variant type differed between the RefSeq and UCSC gene annotation, the called

location is determined by an ordering precedence: exonic/splicing > ncRNA > 3'/5'UTR >

intronic > promoter region.

Potentially function-altering variants were identified using three algorithms to rank

variant deleteriousness: Combined Annotation Dependent Depletion (CADD) integrates multiple

weighted metrics to identify deleterious variants [35,36], Sorting Intolerant From Tolerant (SIFT)

predicts the functional impact of amino acid substitution caused by genetic variation [37], and

Polymorphism Phenotyping (PolyPhen) identifies the predicted structural and functional

outcome of amino acid substitution [38]. Variants considered potentially function-altering (referred

to as "damaging" or "pathogenic" by algorithms) were defined as SNVs that are in the top 1% of

deleterious variants by CADD score (Phred-normalized CADD score ≥ 20), deleterious by SIFT

prediction (SIFT score ≤ 0.05) or damaging by Polyphen2 prediction (Polyphen2 score between

0.15-1.0 as possibly damaging and 0.85-1.0 as probably damaging).

Star Allele Calling

CYP450 "star alleles" (phased haplotypes) were called using Stargazer [39,40] and PyPGx

(version 0.14.0), bioinformatics tools for identifying star alleles by detecting single nucleotide

variants (SNVs), insertion/deletions (indels), and structural variants (SVs) of pharmacogenes.

The star allele system is a nomenclature used specifically for pharmacogenes [41] – star alleles

refer to haplotype patterns in pharmacogenes and are usually associated with protein activity

levels. Phased haplotypes were called for eleven pharmacogenes that represent enzymes which

are responsible for up to 75% of the metabolism of commonly prescribed drugs [17]: *CYP1A2*,

CYP2A6, *CYP2B6*, *CYP2C19*, *CYP2C8*, *CYP2C9*, *CYP2D6*, *CYP2E1*, *CYP2J2*, *CYP3A4*, and

CYP3A5. VCF files, without additional QC filtering, along with the read depth data for each gene

were the input into the haplotype callers, which utilize the statistical phasing software BEAGLE

to phase the variants into haplotypes [42]. Star alleles were then inferred by identifying the

haplotype calls from core variants of each star allele, as well as tag variants, which define sub-

alleles of each star allele. Star allele diplotype calls were considered "non-determinant" if any

variant used in star allele detection method had a read depth ≤ 1.

SV calls were assessed with PyPGx (version 0.14.0) using three control genes: vitamin D

receptor (*VDR)*, epidermal growth factor receptor (*EGFR)*, and ryanodine receptor 1 (*RYR1)*.

Depth of coverage data was calculated using PyPGx to call the read depth at each position and

was compared to the control genes, which are stable read regions that are highly conserved and

do not fluctuate in copy number, to assess copy number variation in the eleven pharmacogenes

[43]. Final calls for star alleles were determined as the call that was consistent in the highest

number of control genes. Phenotype prediction was conducted in PyPGx by converting the called

diplotype into an activity score, which is then used to predict the gene phenotype [39,41]. Calls were

additionally confirmed with visual inspection of read depth regions for the three control genes.

Identifying archaic alleles in modern populations

To identify the presence of archaic *CYP450* variants in modern human populations, we

calculated the allele frequency of the archaic variants described above in modern humans. We

used modern human genomes from all populations sequenced for the 1000 Genomes Project [21],

as well as the Papuans sequenced as part of the Simons Genome Diversity Project [22]. We also

focused on a subset of SNVs that were more likely to be introgressed, which are found outside of

Africa but not in African populations. These SNVs had to have a frequency in African

populations of less than 1 percent and be present in at least one non-African population with an

allele frequency greater than 1 percent. For all results described in the text, we used the archaic

VCF generated using the methods in this paper for comparison with modern humans. For

comparison, we repeated these analyses with the published VCFs that are associated with the

original archaic genome sequencing studies [25,26]. Alleles were categorized as "derived" or

"ancestral" based on the ancestral allele calls identified by the 1000 Genomes Project [18]. We

additionally examined these pharmacogenes in a cohort of seventy publicly available

chimpanzee genomes [27] to determine if any of the variants shared between archaic and modern

humans might be found in chimpanzees as well, representing shared ancestral variation.

We used Haplostrips [44] to calculate the distance between haplotypes of all *CYP450* genes

in modern humans in the 1000 Genomes Panel relative to a reference haplotype. Distances in

Haplostrips are Manhattan distances - simply the number of SNVs with different alleles between

the two sequences. Haplotypes are re-ordered by decreasing similarity with the archaic reference

haplotype. In this case, the Haplostrip is polarized to a Vindija Neanderthal reference haplotype

(a consensus of the two archaic chromosomes) or a Denisovan reference haplotype, and each

subsequent haplotype is ordered by genetic similarity, from most related to least related. For this

analysis, we looked at haplotypes of each *CYP450* gene at the gene coordinates, plus 5000 base

pairs downstream and upstream to capture more linked neutral variation and compared the 1000

Genomes individual's haplotypes with archaic haplotypes.

For one pharmacogene, *CYP2B6*, we observed highly divergent haplotypes found in the

Vindija Neanderthal and a small number of African individuals ($n = 11$). To determine how the

divergent Vindija *CYP2B6* haplotype compared to the divergence between the Vindija

Neanderthal and other Neanderthals in other regions of the genome, we calculated the pairwise

distance between the Chagyrskaya Neanderthal and Vindija Neanderthal genomes with 29.1 kb

windows (*CYP2B6* gene length) across the genome. To explore SNV sharing between the

CYP2B6 haplotypes, we divided the archaic and modern humans into four groups: the Vindija

genome (containing the divergent haplotype), the other three archaic genomes, the eleven

Africans with the divergent haplotypes, and the rest of the modern Africans without the

divergent haplotypes. Each group was scored for the presence or absence of a given SNV, and

the sharing of these SNVs was summarized in a Venn Diagram (Figure S3).

ILS Calculation

While shared ancestry between archaic humans and modern humans outside of Africa

may suggest that introgression has occurred, this pattern may also represent Incomplete Lineage

Sorting (ILS), in which a haplotype that existed in an ancestral population (for example, the

shared ancestor of humans and Neanderthals) is lost from one or more descendant populations

(for example, in Africans). To attempt to distinguish between the two possibilities, we assessed

the probability of ILS occurring in a given region using a previously published equation [14]:

196
$$L = \frac{1}{R \times {}^{b}/_{G}}$$

197   Briefly, the expected length of a shared ancestral sequence haplotype (L) is calculated as the

198   inverse of regional recombination rate (R) multiplied by the branch length (b) over the

199   generation time (G) The probability (p) of seeing a shared haplotype of length (m) follows a

200   Gamma distribution with a shape (S) of 2 and a rate (r) of 1/L:

201
$$p = 1 - pGamma\left(m, S = 2, r = {}^{1}/_{L}\right)$$

Here, we determined the regional recombination rates (R) using HapMap [45], used a generation

time (G) of 29 years [46,47], and assumed a branch length (b) of 550,000 years with a 50,000 year

timeframe of Neanderthal-human interbreeding [23,47].

Data Visualization

All data visualization was conducted in R Studio Suite (R version 4.0.2, https://www.r-

project.org/) with the use of the CRAN library *ggplot2* for graphical visualization of variant

frequency, star allele diplotype calls, deleterious variant occurrence, and haplotype distance

visualization [48,49]. The R package *venneuler* was used to create the Venn diagram in

Supplemental Figure 3 (https://CRAN.R-project.org/package=venneuler).

Results

CYP450 Variation in Neanderthals and Denisovans

In this study, we investigated genetic variation in eleven *CYP450* genes: *CYP1A2,*

CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP2E1, CYP2J2, CYP3A4, and

CYP3A5. We identified a total of 1,623 single nucleotide variants (SNVs) across the eleven

CYP450 genes investigated in one Denisovan (from Denisova Cave), and three Neanderthals

(from the Vindija, Denisova Cave/Altai, and Chagyrskaya sites). Of the variants identified in the

four archaic individuals, the majority were intronic or non-coding, while 6.8% (*n*=111) were in

promoter regions and 4.7% (*n*=77) were exonic (Figure 1A). A summary table of genetic

variants across gene and location is shown in Table S1.

The Vindija Neanderthal harbored a particularly large number of SNVs in the *CYP2B6*

gene, which was an outlier relative to other Neanderthals and warranted further scrutiny. The

Vindija Neanderthal had 334 variant sites in *CYP2B6* with approximately 92% of these variant

sites being heterozygous (Figure 1B), while the other three archaic genomes had fewer than 100

variants each (Denisovan: $n=75$; Altai: $n=72$; Chagyrskaya: $n=76$). Given that *CYP2B6* shares

significant homology with the pseudogene *CYP2B7,* which is located nearby (40.6 kb) [50], we

assessed the read depth at the *CYP2B6* locus in the Vindija Neanderthal genome to ensure that

the elevated variation identified was not a result of read mis-mapping with the paralog gene

(Figure S1A). The *CYP2B6* locus in the Vindija Neanderthal was also identified as highly

divergent from other Neanderthals relative to the rest of the genome and is within the top 1% of

~30-kb windows based on divergence between the Chagyrskaya and Vindija Neanderthals

(pairwise distance = 0.00175, Figure S2).

To further assess the genomic landscape of the archaic individuals in the context of

modern humans, we compared the archaic *CYP450* genetic variation to the variation in modern

human individuals from the 1000 Genomes Project [21]. Interestingly, a subset of modern

individuals carried a haplotype related to the Vindija Neanderthal *CYP2B6* gene and present a

similar increase in variation to the Vindija Neanderthal (Figure 1B). The eleven modern human

individuals identified that carry this haplotype were all from African populations (ASW, ESN,

GWD, LWK, MSL, YRI), and the haplotype is absent outside of Africa. These individuals were

additionally assessed for read mis-mapping (Figure S1B), and demographic information is

described in Table S2. The African divergent haplotype and the Vindija Neanderthal divergent

haplotype uniquely share some SNVs, but they also each have a large number of variants that are

exclusive to their individual haplotype (Figure S3).

Across the eleven *CYP450* genes investigated, archaic variation present in modern

humans from the 1000 Genomes Project was identified as SNVs that were shared between

modern and archaic humans, while found at a frequency of less than 1% in African populations,

which we refer to as 'archaic SNVs' hereon. We identified a total of 155 archaic SNVs that fit

these criteria, including 41 found in human populations at a frequency of 10% or higher, and four

that were exonic (Figure 1C, full list of variants in Table S3). The majority of these 155 SNVs

are derived (91.0%) and only 7 of them (4.5%) were identified in the sample of chimpanzees we

used for comparison, five of which were homozygous in all chimpanzees sampled (Table

S3).While modern Papuans had fewer alleles shared with archaic humans, those that were shared

were present at a much higher frequency and many of the variants were unique to Papuans,

including all but one of the Denisovan-unique variants (Figure 1C, Table S3).

Phased Diplotypes and Predicted Metabolizer Phenotypes

259   We identified the star allele composition of the eleven *CYP450* genes in archaic

individuals using variant and depth of coverage data with previously validated methods [39,40]. The

phased diplotype for each archaic individual was identified as the two primary haplotype calls

based on the presence or absence of star allele-defining nucleotide and structural variants (Figure

2, Table S4). It is important to highlight that the haplotype callers utilized only assess nucleotide

and structural variants included in previously identified star alleles, and therefore excludes novel

pharmacogene haplotypes that may be present in these archaic humans.

266   Pharmacogene diplotypes were identified for all eleven of the pharmacogenes studied for

all four archaic individuals, with the exception of the Vindija diplotype for *CYP2C9*, which was

indeterminate due to very low read coverage at informative variant sites (Figure 2). For each

diplotype, the predicted metabolizer status phenotype was assessed as designated by the

Pharmacogene Variation Consortium (https://www.pharmvar.org/) [41]. We observed variability in

the metabolic rate of the archaic pharmacogenes, ranging from "ultrarapid" (i.e., the enzyme

would break down compounds much faster than expected) to "poor" (i.e., the enzyme is virtually

non-functional) metabolizers (Figure 2). One nuance of this analysis is that the impact of being a

slower or faster metabolizer is not necessarily universally positive or negative but is context-

dependent and depends on multiple factors, including whether breaking down a compound

activates or inactivates its function, as well as environmental influences and the genetic

background of the individual bearing the diplotype. Additionally, the metabolizer phenotypes

were established in modern humans, and so the phenotypes may slightly differ for the archaic

individuals.

280   Structural variation (SV) was identified in three of the eleven pharmacogenes

investigated– *CYP2A6, CYP2D6,* and *CYP2B6* (Figure 3). The Altai and Vindija Neanderthals

presented the homozygous partial deletion hybrid *CYP2A6*12* variant (*12/*12*, Figure 3A),

while the Chagyrskaya Neanderthal was heterozygous for the *12* haplotype (*1/*12*, Figure 3B).

No structural variation was identified for the Denisovan individual in *CYP2A6* (Figure 3C). For

CYP2D6, we identified a gene duplication event in the Altai and Vindija Neanderthals,

presenting a *2/*2x3* diplotype (Figure 3D), while no structural variation was identified in the

Chagyrskaya Neanderthal and Denisovan individuals (Figure 3E).

288   Interestingly, a gene duplication event in *CYP2B6* was identified in the Vindija

Neanderthal (*22/*22x2*, Figure 3F-G) which was previously recognized as having a large

number of heterozygous variants, as compared to the other Neanderthals and Denisovan for this

gene (Figure 1B). Inspection of read depth in this region (Figure S1A) and allelic imbalance of

CYP2B6 for the Vindija Neanderthal (i.e., an allele fraction of 0.33 and 0.66, Figure 3F) further

substantiates a gene duplication event of *CYP2B6* in the Vindija Neanderthal, as opposed to mis-

mapping of reads containing *CYP2B7*. Perhaps most surprising is the fact that the group of

eleven modern Africans with the similarly divergent haplotype (Figure 1B) also seem to share

this gene duplication. Gene duplication was identified in eight of the 11 modern African

individuals using PyPGx, and manual visual inspection suggests that the three other individuals

also have elevated copy number in this region (Figure S4A). The duplicated *CYP2B6* gene is

absent in the other archaic humans and in any non-African population in our dataset (Table S2).

Potentially Function-altering SNVs

302   While haplotype calling methods can predict genotypes predicated on known star alleles,

novel or un-classified variants are excluded from this method of analysis. Potentially function-

altering SNVs were characterized using three programs that can predict functional impact of

genetic variation: Combined Annotation Dependent Depletion (CADD v1.6), Sorting Intolerant

From Tolerant (SIFT 4G v2.0.0), and Polymorphism Phenotyping (PolyPhen v2) [35,37,38]. While

"potentially damaging" is the standard terminology for SNVs that alter enzyme function, in this

instance, these SNVs will be referred to as "potentially function-altering" as the impact of altered

pharmacogenetic enzymes is dependent both on the substrate and on the individual's genetic

background. We identified 23 potentially function-altering variants in the eleven *CYP450* genes

investigated (Figure 4), including 7 potentially novel variants that were not observed in any

variant annotation databases utilized (see Methods).

313   Of the potentially function-altering variants identified, 6 SNVs were diagnostic for star

alleles (Figure 2 and 4, Table S5). *CYP2C9* rs1799853 is the core SNV for the *2 allele and was

identified in the Altai and Chagyrskaya Neanderthal individuals, which is predicted to cause an

intermediate metabolizer phenotype (or a slower metabolic rate) in modern humans. This SNV is

also found globally in modern humans at less than 5% frequency, although European and

admixed American populations from the 1000 Genomes Project Panel have higher frequencies of

this allele, from 8-15% (Table S3). In *CYP2C19*, the primary SNV defining the *8* allele

(rs41291556) was identified in the three Neanderthal individuals and confers a poor metabolizer

status in modern humans (Figure 4, Table S5), although it is found at very low frequencies (0.5-

1.5%) in a small number of modern human populations (Table S3). Finally, the two variants that

compose the *CYP2D6*4* haplotype (rs3892097 and rs1065852) were identified as heterozygotes

in the Denisovan individual and have been shown to confer an intermediate metabolizer status in

modern humans (Figure 4). In addition to the two examples stated above, an exonic missense

variant in *CYP2D6* was identified as heterozygous in a single Papuan (rs3915951) and was

scored as likely to alter enzyme function (Figure 4, Table S3).

328   All 23 of the potentially function-altering alleles were considered derived according to

the 1000 Genomes Project [21] ancestral allele and not identified in any of the chimpanzee samples

[27] assessed (Table S5). Aside from the potentially function-altering variants identified, the

majority of archaic variants present in non-African modern humans were either intronic (*n=140*)

or exonic synonymous (*n=4*) mutations, making their impact on metabolism unclear (Figure 1C,

Table S3).

Introgression of archaic CYP450 alleles into modern humans

336   As shown in Table 1, each *CYP450* gene had a range of archaic variants found in modern

human populations, from 2 shared SNVs (*CYP1A2*) to 64 shared SNVs (*CYP2C19*). This shared

variation also varied extensively across global human populations, likely reflecting patterns of

past interbreeding with diverse archaic populations and genetic drift [9,51], but the shared archaic

SNVs were largely at low frequency across populations (Figure 1C, Table S3). To confirm if the

archaic SNVs identified in modern human populations for the *CYP450* genes corresponded to an

archaic haplotype inherited from Neanderthals, we calculated sequence divergence between

CYP450 haplotypes using Haplostrips [44]. Of the eleven *CYP450* genes studied here, the gene

with the clearest evidence for haplotype sharing between Neanderthals and modern humans was

CYP2J2, where a Neanderthal-like haplotype was present in all non-African populations at

frequencies higher than 2%, with the highest frequencies found in admixed American

populations (Figure 5, Figure S5). The Neanderthal-like *CYP2J2* haplotype in modern humans

includes 8 SNVs that are found in Neanderthals, although the three SNVs that were predicted to

be function-altering in the Neanderthal genome are absent from the modern human Neanderthal-

like haplotypes (Figure 5, Table S3). Haplostrip plots for the ten additional *CYP450* genes are

displayed in Figures S6 and S7.

Discussion

We investigated a panel of *CYP450* genes in archaic individuals which are responsible

for 75% of all drug metabolizing reactions and represent the bulk of drug metabolizing enzymes

(DME) for which therapeutic recommendations exist [17]. DMEs may be subject to evolutionary

processes (e.g., mutation, drift, natural selection) because of their role as detoxifiers of

xenobiotic substances, yet the insights from genetic variation in archaic genomes have not been

characterized to date.

The majority of the *CYP450* genes in our panel conferred a normal/intermediate or

unknown metabolizer phenotype (*CYP1A2*, *CYP2C8*, *CYP2C9*, *CYP2E1*, *CYP2J2*, *CYP3A4* and

CYP3A5), consistent with previous studies showing strong purifying selection in pharmacogenes

[52]. Genes that were predicted to have a poor/slow (*CYP2A6*, *CYP2C19*) or ultrarapid (*CYP2B6*,

CYP2D6) metabolizer status have also been known to have extreme variation in human

populations [53] and may indicate instances of natural population variation or adaptation to the

environment. However, it is important to note that the predicted phenotypes are based on modern

human phenotypes, and it is unclear how the archaic-specific variation (particularly from the

novel function-altering variants) would affect metabolism in archaic individuals. These variants

of uncertain significance would benefit from further functional tests, individually as well as in

the haplotypes found in archaic individuals, to measure both enzyme activity and protein

abundance.

The distribution of archaic variants in the eleven *CYP450* genes we studied in modern

populations is largely consistent with our general understanding of archaic introgression in

human populations. For example, the majority of shared *CYP450* variants are Neanderthal in

origin, which is consistent with the proportion of Neanderthal ancestry being significantly higher

than the proportion of Denisovan ancestry for all populations except Melanesians [9]. Papuans and

other Melanesian populations have much more Denisovan ancestry than other populations [26] and

may have had gene flow from a different Denisovan population than the one encountered by the

ancestors of Eurasians [54], and thus, it is not surprising that they have a unique set of archaic

alleles for the *CYP450* genes, many of which are Denisovan in origin.

Some of the variation shared between archaic and modern humans likely was present in

the ancestral population to humans, Neanderthals, and Denisovans. Modern and archaic humans

split relatively recently in evolutionary time and given the evolutionary constraints around

pharmacogenetic enzyme structure and function, it is to be expected that Neanderthals and

Denisovans would share similar diplotypes for humans for some of these genes. However, given

the accumulation of neutral mutations over the half-million years of divergence between these

species, particularly in introns and third codon positions, we are still able to distinguish between

Neanderthal, Denisovan, and modern human haplotypes from this mutational background, even

if functionally they present identical diplotypes.

Evidence for archaic introgression in pharmacogenes

Given that pharmacogenes can be useful for adapting to novel environments [2,3], we

searched for evidence to suggest that some of the pharmacogene variation shared between

archaic and modern humans were introduced through gene flow. Our primary criterion for

identifying putatively archaic alleles was requiring that the archaic alleles were found in Africans

at a frequency of less than 1% and found in at least one non-African population with a frequency

higher than 1%. This method likely has an increased false-discovery rate (i.e., alleles shared

between archaic and modern humans that are rare in Africa but are not introgressed) but provides

a starting point for identifying potentially introgressed sites. While the goal of this study was to

characterize variation rather than identify regions of selection or gene flow, here we suggest that

some of the pharmacogenes may in fact show evidence for archaic introgression, which should

be tested more rigorously for confirmation.

First, we infer archaic allele sharing between Europeans and Neanderthals in *CYP2C9.*

The CYP2C9 enzyme is considered one of the most important DMEs in humans and responsible

for metabolizing multiple classes of medications [55,56] with substrates including non-steroidal

anti-inflammatories [57–60], angiotensin II blockers (e.g. losartan) [61], S-warfarin [57,62], phenytoin [63],

and tolbutamide [64]. In the *CYP2C9* gene, more than 60 haplotypes have been identified through

the PharmVar consortium [65,66]. In this study, we identify multiple SNVs within *CYP2C9* that are

shared between non-African populations and Neanderthals. Most interestingly, rs1799853, the

causative SNV for the *CYP2C9*2* haplotype, is found in the Altai and Chagyrskaya Neanderthals

and at frequencies of 8-15% in European populations, as well as in admixed American

populations with a significant component of European ancestry [67]. In modern humans, the

CYP2C9*2 variant has decreased enzyme activity compared to the reference haplotype [68], and

negatively affects warfarin metabolism [62], increases the risk of hypoglycemia from sulfonylurea

treatments [69] and increases the risk of overdose on phenytoin [70]. However, it seems to also be

protective against certain cancers [71]. While some *CYP450* genetic variants (including *CYP2C19*

and *CYP2D6* variants) have a global distribution consistent with the serial bottlenecks human

populations experienced as they expanded out of Africa and into Eurasia, Oceania, and the

Americas, the *CYP2C9* variants do not show the same correlation between genetic and

geographic distance [72]. If the *CYP2C9*2* variant was indeed introgressed from Neanderthals, the

elevated frequency in European populations suggests that it may have been adaptive for

processing certain xenobiotics that human populations were exposed to in Western Eurasia.

However, at this time we cannot rule out that the elevated frequency of the Neanderthal allele in

Europeans was the result of strong founder effect. Further exploration of this haplotype would be

needed to determine whether this haplotype was introgressed and then targeted by positive

selection.

Second, we identify a haplotype in *CYP2J2* that is found in all non-African populations

but lacks the potentially function-altering variants found in Neanderthals, possibly suggesting

that purifying selection has occurred for this gene. The CYP2J2 enzyme accounts for roughly 1-

2% of hepatic CYP protein expression and is involved in the oxidation pathways of

polyunsaturated fatty acids [73] and mediates the oxidation of drugs including ebastine [74],

astemizole [75,76], terfenadine [76], and ebastine [77], but also displays high expression in the lung,

kidney, heart, placenta, salivary gland and skeletal muscle [73,78]. The majority of established

CYP2J2 polymorphisms occur at low frequencies, with the most common variant, *CYP2J2*7,*

occurring at frequencies between 1-20% across global populations [73,79–84]. A comparison of

haplotypes demonstrates that a Neanderthal-like haplotype is found in Eurasians and admixed

American populations, with the highest frequency in the Americas (~8%). Interestingly, the

Neanderthal haplotype contains additional function-altering SNVs that are not found in modern

humans. This suggests two possibilities: either the Neanderthal populations that admixed with

modern humans did not carry these SNVs, or the function-altering SNVs were introduced to

modern humans and then removed by strong purifying selection. Purifying selection has already

removed much of the archaic ancestry in the modern human gene pool [25,26,85–88]. This is likely

because archaic humans had smaller population sizes, allowing for the accumulation of weakly

deleterious alleles [9,51,54,89,90]. After archaic and modern human admixture occurred, increased

pressure from purifying selection would have removed archaic deleterious alleles, and it has been

suggested that a significant portion of Neanderthal ancestry was removed from the modern

human gene pool in just a few generations [10,91,92]. The SNVs that remain in *CYP2J2* likely

represent neutral variants that were maintained in the population or advantageous variants that

may have been under positive selection. Further exploration of these archaic *CYP450* variants

can help clarify the interplay between purifying selection removing deleterious mutations and

positive selection retaining advantageous archaic *CYP450* alleles.

Finally, we identify a structural variant in *CYP2A6* that may have been introduced to

modern humans through gene flow with Neanderthals. The *CYP2A6* gene is expressed mainly in

the liver, and represents between 1% and 10% of the total liver CYP450 protein [93]. The CYP2A6

enzyme metabolizes drugs and pro-carcinogenic compounds including tegafur [94], valproic acid

[95,96], and coumarin [97], and is the primary enzyme involved in nicotine metabolism [98]. At present,

more than 10 different allelic variants are known that cause absent or reduced enzyme activity

[99,100], suggesting that eliminating or reducing *CYP2A6* function is a recurring evolutionary

strategy. The *CYP2A6* gene is located adjacent to the inactive *CYP2A7* gene and several allelic

variants of *CYP2A6* have been created by unequal crossover and gene conversion reactions

between these genes [99]. The allele identified in Neanderthals is *CYP2A6*12*, a hybrid allele

where exon 1 and 2 originated from *CYP2A7*, and exons 3–9 originated from *CYP2A6* (Figure

4). The *CYP2A6*12* allele causes a 50% reduction in CYP2A6 protein levels and a 40% decrease

in CYP2A6 coumarin 7-hydroxylation activity [99]. This hybrid *CYP2A6*12* allele has also been

found at low frequencies in global populations, including African-American (0.4%), Canadian

First Nation (0.5%), and Japanese (0.8%) individuals [99]. In this study, we identify the

*CYP2A6*12* hybrid allele in all three Neanderthal individuals as well as modern humans in

Europe, Southeast Asia, and the Americas. This sharing of alleles, in particular the absence in

unadmixed African populations, suggests that the human *CYP2A6*12* may have been inherited

through human-Neanderthal introgression. There are alternate explanations that do not include

archaic gene flow, such as incomplete lineage sorting (ILS) or parallel evolution. In the case of

ILS, the unequal crossover event that created *CYP2A6*12* would have predated the split time of

the modern human branch and the Neanderthal-Denisovan branch, [101], but this does not explain

why the allele disappeared in Africans. Given the length of this haplotype (31.9 kb) and the

regional recombination rate (0.77 cM/Mb [45]), however, it is improbable that the 31.9 kb

*CYP2A6*12* haplotype would have been maintained by ILS in both lineages (p=0.0014). ILS was

additionally calculated for all eleven pharmacogenes investigated (Table S6). The final

alternative to explain the prevalence of *CYP2A6*12* is that this hybrid allele evolved multiple

times, independently in modern non-Africans and Neanderthals, in addition to the low frequency

of *CYP2A6*12* introgressed from Neanderthals. It has been proposed that the slow metabolizing

of some plant secondary metabolites may be adaptive as high levels of these toxins in human

tissues may act as a deterrent to parasites [102–104], perhaps explaining why this hybrid allele may

have evolved multiple times.

Potential super-archaic introgression

For *CYP2B6*, we identified highly divergent haplotypes in the Vindija Neanderthal and a

small number of African individuals (*n*=11). CYP2B6 is responsible for metabolizing drugs

including the prodrug cyclophosphamide [50,105]; efavirenz, a non-nucleoside reverse transcriptase

inhibitor [50,106,107]; the antidepressant, bupropion [50,108,109]; and ketamine [110]. Genetic

polymorphisms in *CYP2B6* have been identified to alter enzyme activity [106,111–114] and there have

been over 37 *CYP2B6* haplotypes identified [66,115]. In this study, we observe a haplotype in

CYP2B6 that is found in the Vindija Neanderthal and eleven modern African individuals and

shows evidence of gene duplication. This duplicated haplotype is highly divergent from other

archaic and modern human *CYP2B6* haplotypes, but the duplicated Neanderthal and human

haplotypes are also distinct from one another (Supplemental Figure 4). The cause of the elevated

SNV count in this duplicated haplotype is unclear. It is possible that the gene duplicate is non-

functional, and the variation accumulated because selection has been relaxed, although there is

no obvious evidence (i.e., a stop codon) to suggest that the enzyme's function has been impaired.

This gene duplication might also be an ancestral variant that has survived at low frequencies in

modern human and Neanderthal populations, but this is unlikely given our ILS calculations for

CYP2B6 (p=0.02, see Supplemental Table 6). Finally, it is also possible that the duplicated

haplotype is introgressed from an archaic hominin that is more divergent than Neanderthals and

Denisovans, also known as a "super-archaic" human. There is evidence to suggest that

Denisovans[21] and Africans[6] have small percentages of their genomes that were inherited from

super-archaic humans, and *CYP2B6* was previously identified as a candidate region for super-

archaic introgression in Africans[6]. Gene flow has also been inferred between Neanderthals and

the ancestors of modern humans before either expanded outside of Africa [116–118]. This implies

that this gene duplication may have super-archaic origins. Future work on this haplotype's

function and evolutionary history may clarify its origin, and how it came to exist in both

Neanderthals and modern humans.

Limitations

Our findings suggest that Neanderthals and Denisovans have pharmacogene diplotypes

that are similar to that of modern humans, and in some cases may reflect gene flow between

archaic and modern humans. However, some limitations to the data must be noted. First, the

archaic genomes are ancient DNA, which is degraded compared to modern DNA. This

degradation often manifests as lower read depth, skewed allele balance, shorter read lengths, as

well as transition point mutations (cytosine (C) $\rightarrow$ thymine (T)), which impact both mapping and

sequencing accuracy. In order to distinguish true polymorphism at heterozygous sites from

sequencing errors and DNA damage, we excluded base pairs that are represented in only a single

read for each position in the BAM read alignment and removed any heterozygous calls that had

an allelic read ratio less than 0.2 or greater than 0.8. The nature of ancient genomes resulted in

low read depth for some star alleles that were called, but given this limitation, we also manually

checked the sequencing reads for all diplotypes reported.

Additionally, haplotype calling on the archaic genomes required phasing of the archaic

CYP450 sequences using a human reference panel, and diplotype calls were made using current

known human star alleles. While this is the best reference panel available, it is possible that

phasing an archaic genome using modern human reference genomes will result in an inaccurate

haplotype. However, the majority of SNVs were homozygous, which is consistent with the much

lower heterozygosity of Neanderthals and Denisovans along the entire genome [23–26], and many of

the heterozygous star allele calls were composed of only a single nucleotide variant, effectively

deeming the phasing non-impactful. The concern that the archaic diplotypes may represent novel

star alleles that are not shared with humans remains, and further functional validation of the

novel potentially function-altering SNVs will be required to confirm the archaic diplotype calls.

We also used a novel read filtering method, resulting in a larger number of heterozygous

sites compared to the variant call files that are usually used to represent the archaic genomes [23–

26]. The method used to make the previously generated variant call files likely used a more

stringent method of filtering, resulting in more sites being called as homozygous. To ensure that

our results were not biased by our variant calling method, we repeated all analyses (including

calling star alleles, identifying archaic SNVs in modern humans, and comparing human and

archaic haplotypes) using the original variant call files, and found that while the number of

identified SNVs is smaller, the patterns identified (such as the divergent *CYP2B6* haplotype in

the Vindija and allele sharing between archaic and modern humans) remain.

Conclusions

Understanding the impact of archaic variation on modern human health is still in its early

stages. While a few genes have been identified for which the archaic alleles play an important

role in human adaptation to their environment, such as *EPAS1*, the vast contribution of archaic

ancestry to human fitness is less clear. For example, modern humans carry archaic variants of

medically important genes such as ABO blood group antigens, but their impact to human health

remains unknown [119]. Our results suggest that interactions between modern and archaic humans

may have resulted in the introduction of novel *CYP450* variants to modern human populations,

helping them adapt to novel environments as they expanded out of Africa. Important insights

will continue to emerge from careful inspection of pharmacologically relevant and highly studied

genes such as the *CYP450* genes investigated here. We look forward to identifying archaic

CYP450 alleles inherited by modern humans that have been adaptive in historically

underrepresented populations in genomic research such as Indigenous American populations,

which have uniquely adapted to numerous novel environments and have undergone multiple

genetic bottlenecks due to colonization and disease.

Declaration of interests

The authors declare no competing interests

Data and Code availability

The code generated during this study is available on GitHub (https://github.com/the-claw-

lab/aDNA_PGx_2021 and https://github.com/kelsey-witt/archaic_pgx).

Author contributions: THW, KEW, SBL, RSM, EHS, FV, and KGC designed research, project

conception and development. THW, KEW, SBL, RSM, EHS, FV, and KGC conducted research,

formal analysis, and manuscript writing – original draft. THW, KEW, SBL, RSM, EHS, FV, and

KGC performed manuscript review and editing. THW, KEW, SBL, RSM, EHS, FV, and KGC

provided resources and study oversight. THW, KEW, FV, and KGC have primary responsibility

for final content. All authors read and approved final manuscript

Acknowledgements: We would like to thank George Perry and Omer Gokcumen for comments

on early versions of our manuscript, and anonymous reviewers for their feedback.

Funding: THW and KGC are supported by the NHGRI grant to KGC R35HG011319. KEW and

EHS are supported by NIH 1R35GM128946-01. EHS is also supported by the Alfred P. Sloan

Foundation grant.

Figure 1. *CYP450* genetic variation in archaic individuals and shared with modern humans.

(A) The total number of SNVs identified in each archaic individual displayed as a stacked bar

plot. Each color displays the relative variant location identified using ANNOVAR with 3' & 5'

untranslated regions grouped together (UTR), and the promoter region (Promoter) characterized

as any upstream variants that are within 2000 bp of the gene. (B) A larger number of variants

were identified in *CYP2B6* in the Vindija Neanderthal. Bar plots have been faceted by the

archaic individuals, the modern individuals identified to share the Vindija *CYP2B6* haplotype,

and the average number of variants for 10 modern individuals from each super population of the

1000 Genomes Project. Variants are colored by zygosity (i.e., heterozygous if the variant site has one alternative allele and homozygous alternative if the variant site has two alternative alleles at that position). The height of each stacked bar plot represents the total number of variants for each individual and the number of variants for the modern individuals have been reported as the mean with error bars representing the standard deviation. (C) A subset of archaic variants ($n = 155$) was identified to have an allele frequency <1% in African populations and >1% in at least one non-African population. Sub-populations have been grouped as African (AFR), East Asian (EAS), European (EUR), admixed American (AMR), South Asian (SAS), and Papuan (PNG) and allele frequencies for each population are shown on the gradient heatmap. The allele dose of each archaic variant is shown in the heatmap on the bottom, with an allele dose of 0 representing homozygous reference and an allele dose of 2 representing homozygous alternative.

Figure 2. Phased diplotypes and predicted metabolizer phenotypes identified in archaic individuals. Heatmap showing the primary star allele diplotypes for each gene from the three Neanderthal and one Denisovan individuals. Each diplotype was determined as the two phased haplotypes, which are detected with the variant and read data from each gene. The fill of each tile represents the predicted phenotype determined by the combined activity score associated with the presented diplotype as identified in modern humans. Unknown phenotypes are represented with grey tiles and indicate that the functional effect of the diplotype has not been determined. A diplotype was considered non-determinant (N.D.) if there was a lack of read coverage at specific variant locations.

Figure 3. Structural variants detected in *CYP2A6*, *CYP2D6*, and *CYP2B6*. Copy number variation plots (left) demonstrating the structural variation and allele fraction plots (right) showing allelic fraction for each Neanderthal and Denisovan individual across *CYP2A6, CYP2D6,* and *CYP2B6*. The *CYP2A6/A7* structural variant, **12*, identified for the (A) Altai Neanderthal and Vindija Neanderthal which both present a **12/*12* diplotype, characterized by a partial deletion hybridization event in both gene copies. (B) The Chagyrskaya Neanderthal

demonstrated a *1/*12 diplotype and (C) the Denisovan individual presented no structural variation. In *CYP2D6*, (D) the Altai and Vindija Neanderthals presented a gene multiplication while the (E) Chagyrskaya Neanderthal and Denisovan individual presented no structural variation in *CYP2D6*. For *CYP2B6*, (F) the Vindija Neanderthal displayed a gene duplication event, while (G) no structural variation was identified for the Altai Neanderthal, Chagyrskaya Neanderthal, and Denisovan individual. Copy number plots are estimated based on read data (displayed as the copy number normalized to the *VDR* control gene region) with the orange line indicating the copy number assessment for each gene. Allele fraction plots display the allele frequency of each variant for the two haplotypes, which demonstrates the allelic decomposition after identifying structural variation. For each structural variant, there is a schematic representation of the predicted structure of each gene, with an arrow indicating the direction of transcription. The hg19 genetic coordinates are presented on the *x axis* with the gene regions indicated directly above.

Figure 4. Function-altering variants in archaic individuals. Circle plot displaying the occurrence of potentially function-altering variants for each individual in the selected *CYP450* genes. Variants are identified with rsID, if no rsID number was identified, variants are considered potentially novel and the hg19 position is displayed along with mutation observed (position: reference allele > alternative allele). Variants that compose known star alleles are highlighted in gray and denoted with the respective star allele preceding the rsID. Allele dosage is categorized as the number of alternate alleles at that position. Un-filled circles indicate that no call was made at the given position for that individual. An allele dose of 1 indicates an SNV is heterozygous for the reference and alternate allele, and an allele dose of 2 indicates an SNV is homozygous for the alternate allele. No variants were identified for *CYP2A6*.

Figure 5. Haplostrips violin plot of *CYP2J2*. Haplotype distances at *CYP2J2* (hg19 1:60355979-60395470) in modern human populations relative to a Neanderthal reference haplotype (Vindija). The kernel density on each violin plots represents the abundance of individual haplotypes ordered by distance to the reference. Most human-like haplotypes sit at the same distance (furthest from the archaic reference), giving the violin plots a "flat top". Outlier data points near lower distances represent modern human haplotypes carrying the Neanderthal-like *CYP2J2* in all populations except African populations.

Table 1. A summary of archaic variants identified in modern human populations for

CYP450 **genes.** The number of SNVs found in each population is listed, along with the median allele frequencies for all SNVs identified in the populations in the region. "East Asians", "Europeans", "admixed Americans", and "South Asians" refer to the populations from the 1000 Genomes dataset, while "Papuans" refers to the Simons Genome Diversity Project Papuan population. Variants of interest have been highlighted with a section sign (§).

| Gene | Number of Archaic Variants | | Number of SNVs Identified in Each Population | | | | |
	Total Identified	Identified in Modern Populations	East Asians	Europeans	Admixed Americans	South Asians	Papuans
CYP1A2	46	2	2§	2§	1§	1§	0
CYP2A6	69	9	0	6	6	6	3§
CYP2B6	426	20	4	14	13	14	5§
CYP2C8	165	15	15	15§	15§	15	0
CYP2C9	162	24	24	24§	24§	24§	1§
CYP2C19	332	64	52	59	3	58	5§
CYP2D6	44	3	0	0	0	0	3§
CYP2E1	84	1	0	0	0	0	1§
CYP2J2	94	11	9	8§	9§	9	1§
CYP3A4	93	6	2	1	0	6	6§
CYP3A5	108	0	*N/A*	*N/A*	*N/A*	*N/A*	*N/A*

References

1. Thomas, J.H. (2007). Rapid Birth–Death Evolution Specific to Xenobiotic Cytochrome P450 Genes in Vertebrates. PLoS Genet. *3*, e67.

2. Fuselli, S., de Filippo, C., Mona, S., Sistonen, J., Fariselli, P., Destro-Bisol, G., Barbujani, G., Bertorelle, G., and Sajantila, A. (2010). Evolution of detoxifying systems: the role of environment and population history in shaping genetic diversity at human CYP2D6 locus. Pharmacogenet. Genomics *20*, 485–499.

3. Fuselli, S. (2019). Beyond drugs: the evolution of genes involved in human response to medications. Proc. Biol. Sci. *286*, 20191716.

4. Fan, S., Hansen, M.E.B., Lo, Y., and Tishkoff, S.A. (2016). Going global by adapting local: A review of recent human adaptation. Science *354*, 54–59.

5. Bergström, A., Stringer, C., Hajdinjak, M., Scerri, E.M.L., and Skoglund, P. (2021). Origins of modern human ancestry. Nature *590*, 229–237.

6. Durvasula, A., and Sankararaman, S. (2020). Recovering signals of ghost archaic introgression in African populations. Science Advances *6*, eaax5097.

7. Browning, S.R., Browning, B.L., Zhou, Y., Tucci, S., and Akey, J.M. (2018). Analysis of Human Sequence Data Reveals Two Pulses of Archaic Denisovan Admixture. Cell *173*, 1–9.

8. Villanea, F.A., and Schraiber, J.G. (2018). Multiple episodes of interbreeding between Neanderthal and modern humans. Nature Ecology & Evolution *3*, 39–44.

9. Sankararaman, S., Mallick, S., Patterson, N., and Reich, D. (2016). The Combined Landscape

of Denisovan and Neanderthal Ancestry in Present-Day Humans. Curr. Biol. *26*, 1241–1247.

10. Petr, M., Pääbo, S., Kelso, J., and Vernot, B. (2019). Limits of long-term selection against Neandertal introgression. Proc. Natl. Acad. Sci. U. S. A. *116*, 1639–1644.

11. Zhang, X., Kim, B., Lohmueller, K.E., and Huerta-Sánchez, E. (2020). The Impact of Recessive Deleterious Variation on Signals of Adaptive Introgression in Human Populations. Genetics *215*, 799–812.

12. Schaefer, N.K., Shapiro, B., and Green, R.E. (2021). An ancestral recombination graph of human, Neanderthal, and Denisovan genomes. Science Advances *7*, eabc0776.

13. Racimo, F., Sankararaman, S., Nielsen, R., and Huerta-Sánchez, E. (2015). Evidence for archaic adaptive introgression in humans. Nat. Rev. Genet. *16*, 359–371.

14. Huerta-Sánchez, E., Jin, X., Asan, Bianba, Z., Peter, B.M., Vinckenbosch, N., Liang, Y., Yi, X., He, M., Somel, M., et al. (2014). Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. Nature *512*, 194–197.

15. Dannemann, M., Andrés, A.M., and Kelso, J. (2016). Introgression of Neandertal- and Denisovan-like Haplotypes Contributes to Adaptive Variation in Human Toll-like Receptors. The American Journal of Human Genetics *98*, 399.

16. Mendez, F.L., Watkins, J.C., and Hammer, M.F. (2012). A haplotype at STAT2 Introgressed from neanderthals and serves as a candidate of positive selection in Papua New Guinea. Am. J. Hum. Genet. *91*, 265–274.

17. Evans, W.E., and Relling, M.V. (1999). Pharmacogenomics: translating functional genomics

into rational therapeutics. Science *286*, 487–491.

18. Chen, J., Jiang, S., Wang, J., Renukuntla, J., Sirimulla, S., and Chen, J. (2019). A comprehensive review of cytochrome P450 2E1 for xenobiotic metabolism. Drug Metab. Rev. *51*, 178–195.

19. Ekins, S., and Wrighton, S.A. (1999). The role of CYP2B6 in human xenobiotic metabolism. Drug Metab. Rev. *31*, 719–754.

20. Esteves, F., Rueff, J., and Kranendonk, M. (2021). The Central Role of Cytochrome P450 in Xenobiotic Metabolism-A Brief Review on a Fascinating Enzyme Family. J Xenobiot *11*, 94–114.

21. 1000 Genomes Project Consortium, Auton, A., Brooks, L.D., Durbin, R.M., Garrison, E.P., Kang, H.M., Korbel, J.O., Marchini, J.L., McCarthy, S., McVean, G.A., et al. (2015). A global reference for human genetic variation. Nature *526*, 68–74.

22. Mallick, S., Li, H., Lipson, M., Mathieson, I., Gymrek, M., Racimo, F., Zhao, M., Chennagiri, N., Nordenfelt, S., Tandon, A., et al. (2016). The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. Nature *538*, 201–206.

23. Prüfer, K., Racimo, F., Patterson, N., Jay, F., Sankararaman, S., Sawyer, S., Heinze, A., Renaud, G., Sudmant, P.H., de Filippo, C., et al. (2014). The complete genome sequence of a Neanderthal from the Altai Mountains. Nature *505*, 43–49.

24. Mafessoni, F., Grote, S., de Filippo, C., Slon, V., Kolobova, K.A., Viola, B., Markin, S.V., Chintalapati, M., Peyrégne, S., Skov, L., et al. (2020). A high-coverage Neandertal genome from Chagyrskaya Cave. Proc. Natl. Acad. Sci. U. S. A. *117*, 15132–15136.

25. Prüfer, K., de Filippo, C., Grote, S., Mafessoni, F., Korlević, P., Hajdinjak, M., Vernot, B., Skov, L., Hsieh, P., Peyrégne, S., et al. (2017). A high-coverage Neandertal genome from Vindija Cave in Croatia. Science *358*, 655–658.

26. Meyer, M., Kircher, M., Gansauge, M.-T., Li, H., Racimo, F., Mallick, S., Schraiber, J.G., Jay, F., Prüfer, K., de Filippo, C., et al. (2012). A high-coverage genome sequence from an archaic Denisovan individual. Science *338*, 222–226.

27. Prado-Martinez, J., Sudmant, P.H., Kidd, J.M., Li, H., Kelley, J.L., Lorente-Galdos, B., Veeramah, K.R., Woerner, A.E., O'Connor, T.D., Santpere, G., et al. (2013). Great ape genetic diversity and population history. Nature *499*, 471–475.

28. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079.

29. Poplin, R., Ruano-Rubio, V., DePristo, M.A., Fennell, T.J., Carneiro, M.O., Van der Auwera, G.A., Kling, D.E., Gauthier, L.D., Levy-Moonshine, A., Roazen, D., et al. (2018). Scaling accurate genetic variant discovery to tens of thousands of samples.

30. O'Leary, N.A., Wright, M.W., Brister, J.R., Ciufo, S., Haddad, D., McVeigh, R., Rajput, B., Robbertse, B., Smith-White, B., Ako-Adjei, D., et al. (2016). Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. Nucleic Acids Res. *44*, D733-45.

31. Kent, W.J., Sugnet, C.W., Furey, T.S., Roskin, K.M., Pringle, T.H., Zahler, A.M., and Haussler, D. (2002). The human genome browser at UCSC. Genome Res. *12*, 996–1006.

32. Wang, K., Li, M., and Hakonarson, H. (2010). ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. Nucleic Acids Res. *38*, e164.

33. Sherry, S.T., Ward, M.H., Kholodov, M., Baker, J., Phan, L., Smigielski, E.M., and Sirotkin, K. (2001). dbSNP: the NCBI database of genetic variation. Nucleic Acids Res. *29*, 308–311.

34. Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang, Q., Collins, R.L., Laricchia, K.M., Ganna, A., Birnbaum, D.P., et al. (2021). Author Correction: The mutational constraint spectrum quantified from variation in 141,456 humans. Nature *590*, E53.

35. Rentzsch, P., Witten, D., Cooper, G.M., Shendure, J., and Kircher, M. (2019). CADD: predicting the deleteriousness of variants throughout the human genome. Nucleic Acids Res. *47*, D886–D894.

36. Rentzsch, P., Schubach, M., Shendure, J., and Kircher, M. (2021). CADD-Splice-improving genome-wide variant effect prediction using deep learning-derived splice scores. Genome Med. *13*, 31.

37. Vaser, R., Adusumalli, S., Leng, S.N., Sikic, M., and Ng, P.C. (2016). SIFT missense predictions for genomes. Nat. Protoc. *11*, 1–9.

38. Adzhubei, I.A., Schmidt, S., Peshkin, L., Ramensky, V.E., Gerasimova, A., Bork, P., Kondrashov, A.S., and Sunyaev, S.R. (2010). A method and server for predicting damaging missense mutations. Nat. Methods *7*, 248–249.

39. Lee, S.-B., Wheeler, M.M., Patterson, K., McGee, S., Dalton, R., Woodahl, E.L., Gaedigk, A., Thummel, K.E., and Nickerson, D.A. (2019). Stargazer: a software tool for calling star alleles from next-generation sequencing data using CYP2D6 as a model. Genet. Med. *21*, 361–372.

40. Lee, S.-B., Wheeler, M.M., Thummel, K.E., and Nickerson, D.A. (2019). Calling Star Alleles With Stargazer in 28 Pharmacogenes With Whole Genome Sequences. Clin. Pharmacol. Ther. *106*, 1328–1337.

41. Gaedigk, A., Ingelman-Sundberg, M., Miller, N.A., Leeder, J.S., Whirl-Carrillo, M., Klein, T.E., and PharmVar Steering Committee (2018). The Pharmacogene Variation (PharmVar) Consortium: Incorporation of the Human Cytochrome P450 (CYP) Allele Nomenclature Database. Clin. Pharmacol. Ther. *103*, 399–401.

42. Browning, S.R., and Browning, B.L. (2007). Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. Am. J. Hum. Genet. *81*, 1084–1097.

43. MacDonald, J.R., Ziman, R., Yuen, R.K.C., Feuk, L., and Scherer, S.W. (2014). The Database of Genomic Variants: a curated collection of structural variation in the human genome. Nucleic Acids Res. *42*, D986-92.

44. Marnetto, D., and Huerta-Sánchez, E. (2017). Haplostrips : revealing population structure through haplotype visualization. Methods Ecol. Evol. *8*, 1389–1392.

45. Myers, S., Bottolo, L., Freeman, C., McVean, G., and Donnelly, P. (2005). A fine-scale map of recombination rates and hotspots across the human genome. Science *310*, 321–324.

46. Langergraber, K.E., Prüfer, K., Rowney, C., Boesch, C., Crockford, C., Fawcett, K., Inoue, E., Inoue-Muruyama, M., Mitani, J.C., Muller, M.N., et al. (2012). Generation times in wild chimpanzees and gorillas suggest earlier divergence times in great ape and human evolution. Proc. Natl. Acad. Sci. U. S. A. *109*, 15716–15721.

47. Zeberg, H., and Pääbo, S. (2021). A genomic region associated with protection against severe COVID-19 is inherited from Neandertals. Proc. Natl. Acad. Sci. U. S. A. *118*,.

48. R Core Team (2008). R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing).

49. Wickham, H. (2009). ggplot2: Elegant Graphics for Data Analysis (Springer Science & Business Media).

50. Zanger, U.M., and Klein, K. (2013). Pharmacogenetics of cytochrome P450 2B6 (CYP2B6): advances on polymorphisms, mechanisms, and clinical relevance. Front. Genet. *4*, 24.

51. Sankararaman, S., Mallick, S., Dannemann, M., Prüfer, K., Kelso, J., Pääbo, S., Patterson, N., and Reich, D. (2014). The genomic landscape of Neanderthal ancestry in present-day humans. Nature *507*, 354–357.

52. Nelson, M.R., Wegmann, D., Ehm, M.G., Kessner, D., St Jean, P., Verzilli, C., Shen, J., Tang, Z., Bacanu, S.-A., Fraser, D., et al. (2012). An abundance of rare functional variants in 202 drug target genes sequenced in 14,002 people. Science *337*, 100–104.

53. Zhou, Y., Ingelman-Sundberg, M., and Lauschke, V.M. (2017). Worldwide Distribution of Cytochrome P450 Alleles: A Meta-analysis of Population-scale Sequencing Projects. Clinical Pharmacology & Therapeutics *102*, 688–700.

54. Vernot, B., Tucci, S., Kelso, J., Schraiber, J.G., Wolf, A.B., Gittelman, R.M., Dannemann, M., Grote, S., McCoy, R.C., Norton, H., et al. (2016). Excavating Neandertal and Denisovan DNA from the genomes of Melanesian individuals. Science *352*, 235–239.

55. Henderson, L.M., Claw, K.G., Woodahl, E.L., Robinson, R.F., Boyer, B.B., Burke, W., and Thummel, K.E. (2018). P450 Pharmacogenetics in Indigenous North American Populations. J Pers Med *8*,.

56. Miners, J.O., and Birkett, D.J. (1998). Cytochrome P4502C9: an enzyme of major importance in human drug metabolism. Br. J. Clin. Pharmacol. *45*, 525–538.

57. Yamazaki, H., Inoue, K., Chiba, K., Ozawa, N., Kawai, T., Suzuki, Y., Goldstein, J.A., Guengerich, F.P., and Shimada, T. (1998). Comparative studies on the catalytic roles of cytochrome P450 2C9 and its Cys- and Leu-variants in the oxidation of warfarin, flurbiprofen, and diclofenac by human liver microsomes. Biochem. Pharmacol. *56*, 243–251.

58. Tracy, T.S., Rosenbluth, B.W., Wrighton, S.A., Gonzalez, F.J., and Korzekwa, K.R. (1995). Role of cytochrome P450 2C9 and an allelic variant in the 4'-hydroxylation of (R)- and (S)-flurbiprofen. Biochem. Pharmacol. *49*, 1269–1275.

59. Hamman, M.A., Thompson, G.A., and Hall, S.D. (1997). Regioselective and stereoselective metabolism of ibuprofen by human cytochrome P450 2C. Biochem. Pharmacol. *54*, 33–41.

60. Miners, J.O., Coulter, S., Tukey, R.H., Veronese, M.E., and Birkett, D.J. (1996). Cytochromes P450, 1A2, and 2C9 are responsible for the human hepatic O-demethylation of R- and S-naproxen. Biochem. Pharmacol. *51*, 1003–1008.

61. Stearns, R.A., Chakravarty, P.K., Chen, R., and Chiu, S.H. (1995). Biotransformation of losartan to its active carboxylic acid metabolite in human liver microsomes. Role of cytochrome P4502C and 3A subfamily members. Drug Metab. Dispos. *23*, 207–215.

62. Rettie, A.E., Wienkers, L.C., Gonzalez, F.J., Trager, W.F., and Korzekwa, K.R. (1994).

Impaired (S)-warfarin metabolism catalysed by the R144C allelic variant of CYP2C9. Pharmacogenetics *4*, 39–42.

63. Giancarlo, G.M., Venkatakrishnan, K., Granda, B.W., von Moltke, L.L., and Greenblatt, D.J. (2001). Relative contributions of CYP2C9 and 2C19 to phenytoin 4-hydroxylation in vitro: inhibition by sulfaphenazole, omeprazole, and ticlopidine. Eur. J. Clin. Pharmacol. *57*, 31–36.

64. Miners, J.O., and Birkett, D.J. (1996). Use of tolbutamide as a substrate probe for human hepatic cytochrome P450 2C9. Methods Enzymol. *272*, 139–145.

65. Sangkuhl, K., Claudio-Campos, K., Cavallari, L.H., Agundez, J.A.G., Whirl-Carrillo, M., Duconge, J., Del Tredici, A.L., Wadelius, M., Rodrigues Botton, M., Woodahl, E.L., et al. (2021). PharmVar GeneFocus: CYP2C9. Clin. Pharmacol. Ther. *110*, 662–676.

66. Gaedigk, A., Ingelman-Sundberg, M., Miller, N.A., Steven Leeder, J., Whirl-Carrillo, M., Klein, T.E., and the PharmVar Steering Committee (2018). The Pharmacogene Variation (PharmVar) Consortium: Incorporation of the Human Cytochrome P450 (CYP) Allele Nomenclature Database. Clinical Pharmacology & Therapeutics *103*, 399–401.

67. Martin, A.R., Gignoux, C.R., Walters, R.K., Wojcik, G.L., Neale, B.M., Gravel, S., Daly, M.J., Bustamante, C.D., and Kenny, E.E. (2017). Human Demographic History Impacts Genetic Risk Prediction across Diverse Populations. Am. J. Hum. Genet. *100*, 635–649.

68. Van Booven, D., Marsh, S., and McLeod, H. (2010). Cytochrome P450 2C9-CYP2C9. Pharmacogenetics.

69. Chen, L., Li, J.H., Kaur, V., Muhammad, A., Fernandez, M., Hudson, M.S., Goldfine, A.B., and Florez, J.C. (2020). The presence of two reduced function variants in CYP2C9 influences the

acute response to glipizide. Diabet. Med. *37*, 2124–2130.

70. Ninomiya, H., Mamiya, K., Matsuo, S., Ieiri, I., Higuchi, S., and Tashiro, N. (2000). Genetic polymorphism of the CYP2C subfamily and excessive serum phenytoin concentration with central nervous system intoxication. Ther. Drug Monit. *22*, 230–232.

71. Sausville, L.N., Gangadhariah, M.H., Chiusa, M., and Mei, S. (2018). The cytochrome P450 slow metabolizers CYP2C9* 2 and CYP2C9* 3 directly regulate tumorigenesis via reduced epoxyeicosatrienoic acid production. Cancer Res.

72. Sistonen, J., Fuselli, S., Palo, J.U., Chauhan, N., Padh, H., and Sajantila, A. (2009). Pharmacogenetic variation at CYP2C9, CYP2C19, and CYP2D6 at global and microgeographic scales. Pharmacogenet. Genomics *19*, 170–179.

73. Murray, M. (2016). CYP2J2 - regulation, function and polymorphism. Drug Metab. Rev. *48*, 351–368.

74. Hashizume, T., Imaoka, S., Mise, M., Terauchi, Y., Fujii, T., Miyazaki, H., Kamataki, T., and Funae, Y. (2002). Involvement of CYP2J2 and CYP4F12 in the metabolism of ebastine in human intestinal microsomes. J. Pharmacol. Exp. Ther. *300*, 298–304.

75. Lafite, P., André, F., Zeldin, D.C., Dansette, P.M., and Mansuy, D. (2007). Unusual regioselectivity and active site topology of human cytochrome P450 2J2. Biochemistry *46*, 10237–10247.

76. Matsumoto, S., and Yamazoe, Y. (2001). Involvement of multiple human cytochromes P450 in the liver microsomal metabolism of astemizole and a comparison with terfenadine. Br. J. Clin. Pharmacol. *51*, 133–142.

77. Lee, C.A., Jones, J.P., 3rd, Katayama, J., Kaspera, R., Jiang, Y., Freiwald, S., Smith, E., Walker, G.S., and Totah, R.A. (2012). Identifying a selective substrate and inhibitor pair for the evaluation of CYP2J2 activity. Drug Metab. Dispos. *40*, 943–951.

78. Bièche, I., Narjoz, C., Asselah, T., Vacher, S., Marcellin, P., Lidereau, R., Beaune, P., and de Waziers, I. (2007). Reverse transcriptase-PCR quantification of mRNA levels from cytochrome (CYP)1, CYP2 and CYP3 families in 22 different human tissues. Pharmacogenet. Genomics *17*, 731–742.

79. Dreisbach, A.W., Japa, S., Sigel, A., Parenti, M.B., Hess, A.E., Srinouanprachanh, S.L., Rettie, A.E., Kim, H., Farin, F.M., Hamm, L.L., et al. (2005). The Prevalence of CYP2C8, 2C9, 2J2, and soluble epoxide hydrolase polymorphisms in African Americans with hypertension. Am. J. Hypertens. *18*, 1276–1281.

80. King, L.M., Ma, J., Srettabunjong, S., Graves, J., Bradbury, J.A., Li, L., Spiecker, M., Liao, J.K., Mohrenweiser, H., and Zeldin, D.C. (2002). Cloning of CYP2J2 gene and identification of functional polymorphisms. Mol. Pharmacol. *61*, 840–852.

81. King, L.M., Gainer, J.V., David, G.L., Dai, D., Goldstein, J.A., Brown, N.J., and Zeldin, D.C. (2005). Single nucleotide polymorphisms in the CYP2J2 and CYP2C8 genes and the risk of hypertension. Pharmacogenet. Genomics *15*, 7–13.

82. Polonikov, A.V., Ivanov, V.P., Solodilova, M.A., Khoroshaya, I.V., Kozhuhov, M.A., Ivakin, V.E., Katargina, L.N., and Kolesnikova, O.E. (2008). A common polymorphism G-50T in cytochrome P450 2J2 gene is associated with increased risk of essential hypertension in a Russian population. Dis. Markers *24*, 119–126.

83. Polonikov, A.V., Ivanov, V.P., Solodilova, M.A., Khoroshaya, I.V., Kozhuhov, M.A., and Panfilov, V.I. (2007). Promoter polymorphism G-50T of a human CYP2J2 epoxygenase gene is associated with common susceptibility to asthma. Chest *132*, 120–126.

84. Wang, H., Jiang, Y., Liu, Y., Lin, C., Cheng, G., Chen, X., Hao, B., Tan, W., Lin, D., and He, F. (2006). CYP2J2*7 single nucleotide polymorphism in a Chinese population. Clin. Chim. Acta *365*, 125–128.

85. Green, R.E., Krause, J., Briggs, A.W., Maricic, T., Stenzel, U., Kircher, M., Patterson, N., Li, H., Zhai, W., Fritz, M.H.-Y., et al. (2010). A Draft Sequence of the Neandertal Genome. Science *328*, 710–722.

86. Sankararaman, S., Patterson, N., Li, H., Pääbo, S., and Reich, D. (2012). The date of interbreeding between Neandertals and modern humans. PLoS Genet. *8*, e1002947.

87. Wall, J.D., Yang, M. a., Jay, F., Kim, S.K., Durand, E.Y., Stevison, L.S., Gignoux, C., Woerner, A., Hammer, M.F., and Slatkin, M. (2013). Higher levels of neanderthal ancestry in East asians than in europeans. Genetics *194*, 199–209.

88. Chen, L., Wolf, A.B., Fu, W., Li, L., and Akey, J.M. (2020). Identifying and Interpreting Apparent Neanderthal Ancestry in African Individuals. Cell *180*, 677-687.e16.

89. Castellano, S., Parra, G., Sánchez-Quinto, F.A., Racimo, F., Kuhlwilm, M., Kircher, M., Sawyer, S., Fu, Q., Heinze, A., Nickel, B., et al. (2014). Patterns of coding variation in the complete exomes of three Neandertals. Proc. Natl. Acad. Sci. U. S. A. *111*, 6666–6671.

90. Vernot, B., and Akey, J.M. (2014). Resurrecting Surviving Neandeltal Linages from Modern Human Genomes. Science *343*, 1017–1021.

91. Harris, K., and Nielsen, R. (2016). The Genetic Cost of Neanderthal Introgression. Genetics *203*, 881–891.

92. Juric, I., Aeschbacher, S., and Coop, G. (2016). The Strength of Selection against Neanderthal Introgression. PLoS Genet. *12*, e1006340.

93. Haberl, M., Anwald, B., Klein, K., Weil, R., Fuss, C., Gepdiremen, A., Zanger, U.M., Meyer, U.A., and Wojnowski, L. (2005). Three haplotypes associated with CYP2A6 phenotypes in Caucasians. Pharmacogenet. Genomics *15*, 609–624.

94. Komatsu, T., Yamazaki, H., Shimada, N., Nakajima, M., and Yokoi, T. (2000). Roles of cytochromes P450 1A2, 2A6, and 2C8 in 5-fluorouracil formation from tegafur, an anticancer prodrug, in human liver microsomes. Drug Metab. Dispos. *28*, 1457–1463.

95. Kiang, T.K.L., Ho, P.C., Anari, M.R., Tong, V., Abbott, F.S., and Chang, T.K.H. (2006). Contribution of CYP2C9, CYP2A6, and CYP2B6 to valproic acid metabolism in hepatic microsomes from individuals with the CYP2C9*1/*1 genotype. Toxicol. Sci. *94*, 261–271.

96. Tan, L., Yu, J.-T., Sun, Y.-P., Ou, J.-R., Song, J.-H., and Yu, Y. (2010). The influence of cytochrome oxidase CYP2A6, CYP2B6, and CYP2C9 polymorphisms on the plasma concentrations of valproic acid in epileptic patients. Clin. Neurol. Neurosurg. *112*, 320–323.

97. Miles, J.S., McLaren, A.W., Forrester, L.M., Glancey, M.J., Lang, M.A., and Wolf, C.R. (1990). Identification of the human liver cytochrome P-450 responsible for coumarin 7-hydroxylase activity. Biochem. J *267*, 365–371.

98. Hukkanen, J., Jacob, P., 3rd, and Benowitz, N.L. (2005). Metabolism and disposition kinetics of nicotine. Pharmacol. Rev. *57*, 79–115.

99. Oscarson, M., McLellan, R.A., Asp, V., Ledesma, M., Bernal Ruiz, M.L., Sinues, B., Rautio, A., and Ingelman-Sundberg, M. (2002). Characterization of a novel CYP2A7/CYP2A6 hybrid allele (CYP2A6*12) that causes reduced CYP2A6 activity. Hum. Mutat. *20*, 275–283.

100. Benowitz, N.L., Swan, G.E., Jacob, P., 3rd, Lessov-Schlaggar, C.N., and Tyndale, R.F. (2006). CYP2A6 genotype and the metabolism and disposition kinetics of nicotine. Clin. Pharmacol. Ther. *80*, 457–467.

101. Lin, Y.-L., Pavlidis, P., Karakoc, E., Ajay, J., and Gokcumen, O. (2015). The evolution and functional impact of human deletion variants shared with archaic hominin genomes. Mol. Biol. Evol. *32*, 1008–1019.

102. Sullivan, R.J., Hagen, E.H., and Hammerstein, P. (2008). Revealing the paradox of drug reward in human evolution. Proc. Biol. Sci. *275*, 1231–1241.

103. Hagen, E.H., Sullivan, R.J., Schmidt, R., Morris, G., Kempter, R., and Hammerstein, P. (2009). Ecology and neurobiology of toxin avoidance and the paradox of drug reward. Neuroscience *160*, 69–84.

104. Hagen, E.H., Roulette, C.J., and Sullivan, R.J. (2013). Explaining human recreational use of 'pesticides': the neurotoxin regulation model of substance use vs. the hijack model and implications for age and sex differences in drug consumption. Front. Psychiatry *4*, 142.

105. Huang, Z., Roy, P., and Waxman, D.J. (2000). Role of human liver microsomal CYP3A4 and CYP2B6 in catalyzing N-dechloroethylation of cyclophosphamide and ifosfamide. Biochem. Pharmacol. *59*, 961–972.

106. Ward, B.A., Gorski, J.C., Jones, D.R., Hall, S.D., Flockhart, D.A., and Desta, Z. (2003).

The cytochrome P450 2B6 (CYP2B6) is the main catalyst of efavirenz primary and secondary metabolism: implication for HIV/AIDS therapy and utility of efavirenz as a substrate marker of CYP2B6 catalytic activity. J. Pharmacol. Exp. Ther. *306*, 287–300.

107. Desta, Z., Saussele, T., Ward, B., Blievernicht, J., Li, L., Klein, K., Flockhart, D.A., and Zanger, U.M. (2007). Impact of CYP2B6 polymorphism on hepatic efavirenz metabolism in vitro. Pharmacogenomics *8*, 547–558.

108. Hesse, L.M., Venkatakrishnan, K., Court, M.H., von Moltke, L.L., Duan, S.X., Shader, R.I., and Greenblatt, D.J. (2000). CYP2B6 mediates the in vitro hydroxylation of bupropion: potential drug interactions with other antidepressants. Drug Metab. Dispos. *28*, 1176–1183.

109. Faucette, S.R., Hawke, R.L., Lecluyse, E.L., Shord, S.S., Yan, B., Laethem, R.M., and Lindley, C.M. (2000). Validation of bupropion hydroxylation as a selective marker of human cytochrome P450 2B6 catalytic activity. Drug Metab. Dispos. *28*, 1222–1230.

110. Desta, Z., Moaddel, R., Ogburn, E.T., Xu, C., Ramamoorthy, A., Venkata, S.L.V., Sanghvi, M., Goldberg, M.E., Torjman, M.C., and Wainer, I.W. (2012). Stereoselective and regiospecific hydroxylation of ketamine and norketamine. Xenobiotica *42*, 1076–1087.

111. Lang, T., Klein, K., Fischer, J., Nüssler, A.K., Neuhaus, P., Hofmann, U., Eichelbaum, M., Schwab, M., and Zanger, U.M. (2001). Extensive genetic polymorphism in the human CYP2B6 gene with impact on expression and function in human liver. Pharmacogenetics *11*, 399–415.

112. Ariyoshi, N., Miyazaki, M., Toide, K., Sawamura Yi, and Kamataki, T. (2001). A single nucleotide polymorphism of CYP2b6 found in Japanese enhances catalytic activity by autoactivation. Biochem. Biophys. Res. Commun. *281*, 1256–1260.

113. Kirchheiner, J., Klein, C., Meineke, I., Sasse, J., Zanger, U.M., Mürdter, T.E., Roots, I., and Brockmöller, J. (2003). Bupropion and 4-OH-bupropion pharmacokinetics in relation to genetic polymorphisms in CYP2B6. Pharmacogenetics *13*, 619–626.

114. Hofmann, M.H., Blievernicht, J.K., Klein, K., Saussele, T., Schaeffeler, E., Schwab, M., and Zanger, U.M. (2008). Aberrant splicing caused by single nucleotide polymorphism c.516G>T [Q172H], a marker of CYP2B6*6, is responsible for decreased expression and activity of CYP2B6 in liver. J. Pharmacol. Exp. Ther. *325*, 284–292.

115. Desta, Z., El-Boraie, A., Gong, L., Somogyi, A.A., Lauschke, V.M., Dandara, C., Klein, K., Miller, N.A., Klein, T.E., Tyndale, R.F., et al. (2021). PharmVar GeneFocus: CYP2B6. Clin. Pharmacol. Ther. *110*, 82–97.

116. Kuhlwilm, M., Gronau, I., Hubisz, M.J., de Filippo, C., Prado-Martinez, J., Kircher, M., Fu, Q., Burbano, H.A., Lalueza-Fox, C., de la Rasilla, M., et al. (2016). Ancient gene flow from early modern humans into Eastern Neanderthals. Nature *530*, 429–433.

117. Hubisz, M., and Siepel, A. (2020). Inference of Ancestral Recombination Graphs Using ARGweaver. Methods Mol. Biol. *2090*, 231–266.

118. Petr, M., Hajdinjak, M., Fu, Q., Essel, E., Rougier, H., Crevecoeur, I., Semal, P., Golovanova, L.V., Doronichev, V.B., Lalueza-Fox, C., et al. (2020). The evolutionary history of Neanderthal and Denisovan Y chromosomes. Science *369*, 1653–1656.

119. Villanea, F.A., Huerta-Sanchez, E., and Fox, K. (2021). ABO Genetic Variation in Neanderthals and Denisovans. Mol. Biol. Evol. *38*, 3373–3382.