Effect of host-switching on the eco-evolutionary patterns of parasites

[1*]Elvira D'Bastiani, [2]Débora Princepe, [2,3]Flavia MD Marquitti, [1,4]Walter A Boeger, [1,4]Karla M

Campião, [1,5]Sabrina LB Araujo

[1]Laboratório de Interações Biológicas, Programa de Pós-Graduação em Ecologia e

Conservação, Universidade Federal do Paraná, UFPR - Curitiba, Paraná, Brasil

[2]Instituto de Física "Gleb Wataghin", Universidade Estadual de Campinas, UNICAMP -

Campinas, São Paulo, Brasil

[3]Insituto de Biologia, Universidade Estadual de Campinas, UNICAMP - Campinas, São

Paulo, Brasil

[4]Departamento de Zoologia, Universidade Federal do Paraná, UFPR - Curitiba, Paraná, Brasil

[5]Departamento de Física, Universidade Federal do Paraná, UFPR - Curitiba, Paraná, Brasil

*Corresponding author: Elvira D'Bastiani e-mail: elviradbastiani@gmail.com

SHORT RUNNING TITLE: Host-switching influences parasite patterns

ABSTRACT

Increasing empirical evidence has revealed that host-switching are common in the history of

parasites. Still, few have explored how the evolutionary histories of hosts might influence

such switches and then the evolution of parasites. Here, we investigated how the intensity of

host-switching, assumed to depend on the phylogenetic distance between hosts, affects the

ecological and evolutionary patterns of parasite species. We developed an individual-based

model where parasites can explore and colonise hosts under variable host-switching intensity

and have evolution driven by mutation, genetic drift, and mating restriction. We hypothesised

that our model can reproduce ecological and evolutionary patterns of empirical communities,

characterised by turnover among host species and tree imbalance, respectively. We found an

optimum range of host-switching intensity that can predict similar patterns as those observed

in the empirical studies, validating our hypothesis. Our results showed that the turnover

decreased as the host-switching intensity increased with low variation among the model

replications. On the other hand, the tree imbalance had not a monotonic tendency but a wide

variation. These results revealed that while the tree imbalance is a sensitive metric to

stochastic events, the turnover may be a proxy for host switching. Furthermore, local

empirical studies corresponded to higher host-switching intensity when compared to regional

studies, highlighting that spatial scale is probably the crucial limitation of host-switching.

KEYWORDS: dispersal of parasites, opportunity and compatibility of interaction, phylogenetic

conservatism, and structure of the community of parasites.

INTRODUCTION

The dispersal of parasite individuals followed by colonisation of a new host lineage,

known as host-switching, is a common event observed during the evolutionary trajectory of

many parasite lineages (De Vienne et al. 2013). Initially, host-switching results in the increase

of the host repertoire of a parasite (Braga et al. 2021). The colonisation of the new hosts can

result in reproductive isolation, and consequently in speciation of parasite lineages,

characterising the dynamics of the Oscillation Hypothesis (Nylin and Soren 2018). Empirical

examples showing high levels of host-switching include symbiotic interactions ranging from

host-parasite (Meinilä et al. 2004; Agosta et al. 2010; Müller et al. 2018, Fecchio et al. 2019;

Boyd et al. 2022) and plant-insect systems to microbial pathogens (Woolhouse et al. 2005),

brood parasitism (Habermannová et al. 2013; Dominguez et al. 2015), plant-feeding insects,

and parasitic plants (Nylin et al. 2014). Consequently, understanding the factors influencing

the success of host-switching and subsequent speciation events is critical for understanding

parasites diversification.

A general framework that has been used to understand infectious disease, the Stockholm

Paradigm, explores the evolutionary dynamics of host-parasite associations (Brooks et al.

2014; Brooks et al. 2019). This framework suggests that parasites perform host-switching by

ecological fitting hypothesis (Agosta and Klemmens 2008; Agosta and Brooks 2020).

Ecological fitting explains how the process whereby organisms colonise and persist in novel

environments, use novel resources, or form novel associations with other species through a set

of traits/capabilities they already possess (see Agosta and Klemmens 2008; Brooks et al.

2014; Brooks et al. 2019; Agosta and Brooks 2020). The expression of these unexplored

capabilities is mediated by the opportunity of interaction (temporal and spatial) and

determines the possibility of encounters between hosts and unfamiliar parasites. After the

encounter, and if the interaction is compatible, it is followed by the resolution of subsequent

conflicts that emerge from the basic dynamics of "living together", which should result in co-

accommodation (Brooks and McLennan 2002; Araujo et al. 2015).

Ecological and life-history traits also influence the chances of parasites dispersing from

one host species to another. Characteristics of all organisms within the interaction system,

such as niche similarity among host species, modes of transmission of parasites, dietary

preferences of the vector (if there is one), and also ecosystemic characteristics as the host

community composition and shared phylogenetic history are relevant factors that define the

chances of host-switching (Bush et al. 2006; Jaramillo and Rivera-Parra 2018). Niche

similarity among host species is one fundamental element constraining the incorporation of

new host species by ecological fitting. This is because the capacity of a parasite species to use

new resources is related to the phylogenetic conservatism of the resource provided by the host

species. Phylogenetic distance between the original and new host species can represent an

adequate proxy for the nature of the resource, which is tracked by the parasite lineage

(Charleston and Robertson 2002; Agosta and Klemmens 2008; Engelstädter and Fortuna

2019). Consequently, the host phylogenetic conservatism can define the arena of possibilities

for host-switching.

Several studies have indicated the ubiquity of host-switching in nature (see Cuthill and

Charleston 2013; De Vienne et al. 2013; Engelstädter and Fortuna 2019; Fecchio et al. 2019;

Hayward et al. 2021), but (or yet) few studies have explored to which extent the switches are

constrained by inherited possibilities and limitations across hosts evolutionary histories.

Among many potential factors determining host-switching, it seems that host phylogeny and

geographic distributions are two major players (Sanaei et al. 2021). Moreover, the relation

between host-switching and the opportunity for parasite dispersal, as well as their capacity to

explore new hosts, is mostly unexplored (Brooks et al. 2019). Here, we aim to fill these

unexplored gaps by proposing a novel approach to investigate how the intensity of host-

switching affects the ecology and phylogenetic history of the parasites. For this, we assume

that compatibility and the opportunity for interaction (spatial and temporal) may be expressed

through the evolutionary histories of the hosts, and this can influence the host-switching

events.

In this study, we propose a theoretical model based on parasite individuals that can

switch among host species and speciate over time. Host-switching is mediated by

phylogenetic conservatism; that is, the probability of parasites switching hosts decreases with

increasing divergence in the evolutionary time of the hosts. The overall intensity of host-

switching is a controlled parameter of the model. Under the absence of host-switching, the

model is adjusted to parasites speciate due to limitation of host use, resulting in a pattern of

cophylogeny and in paired specialised interaction (each parasite species interacting with one

exclusive host species). We then investigate the eco-evolutionary patterns under different

host-switching intensities, hypothesising that there is an optimum range of host-switching

intensity that can result in the same eco-evolutionary patterns observed in the empirical

studies. These patterns were characterised by species interaction turnover and tree imbalance,

respectively. The model predictions were compared to nine empirical communities, validating

our hypothesis.

MATERIAL AND METHODS

The model

We performed simulations of eco-evolutionary trajectories of parasites influenced by

their host evolutionary history and host-switching events using an individual-based model

(IBM). We assumed that the evolutionary history of the host can represent a proxy for the

resources for parasite species (Agosta et al. 2010; Imrie et al. 2021), and also assumed that the

probability of host-switching decreases as the phylogenetic distance between the species of

host involved in the event (original and new host species) increases (Araujo et al. 2015;

Engelstädter and Fortuna 2019). The model assumes that parasite evolution occurs at the same

evolutionary time scale as the host, which increases possibilities for host-switching as host

speciation occurs (Fig. 1a-c).

FIGURE 1. Schematic representation of the model. **a**. The general sequence of the model

dynamics. **b**. Hypothetical host phylogeny. **c**. Probability of host-switching (P_{hs}) over time.

Each parasite individual can host-switch after the first speciation event (t_1). One host is drawn

for each parasite individual, and the probability of a successful host-switching depends on the

divergent time between the two involved hosts. At t_1 the first speciation event occurs, and the

probability of host-switching is maximum. As time goes on, this probability decreases. At t_2

another speciation event occurs, increasing the number of migration possibilities. At this time

the two younger host species (2 and 3) have the maximum probability of switching hosts (P_{hs}

2-3), but the probability of host-switching between 1 and 2 or between 1 and 3 keeps

decreasing. The colors highlight the 2', 1, 2, and 3 host lineages presented in **b**.

Parasite individuals are explicitly described by biallelic sequences of infinite sites, a

simplified form to represent their genomes and heritable trait. Individuals are monoic and

engage in sexual reproduction, with non-overlapping generations, following the model

proposed by Higgs and Derrida (1991) and Manzo and Peliti (1994). Population evolution is

driven by mutation, genetic drift, and restriction to mating in the absence of natural selection.

With a certain set of parameters, parasite speciation occurs. Each parasite individual is also

characterised by the host species that it interacts. The host species are modelled as resources

that impose a carrying capacity of K parasite individuals, analogous to islands in the Manzo-

Pelit model (Manzo and Peliti 1994), but, in our model, the islands (hosts species in our case)

emerge (as a new host species that speciate) according to a predetermined host diversification

time (i.e. based on ultrametric empirical phylogenies - an ultrametric tree is a kind of additive

tree in which the tips of the trees are all equidistant from the root of the tree). Thus, the

overall carrying capacity increases by K individuals at each new host speciation. The model

does not consider the selection pressure imposed by parasites on the evolution of the resource

(host). Therefore, we are not modelling a process of reciprocal evolution, or co-evolution.

Reproduction of parasites

Reproduction is sexual and occurs between parasite individuals that are in the same host

and that have a minimum genetic similarity, q_{min}, measured based on the Hamming distance

between genomes. In each host species, at each generation, K offspring individuals replace the

parental population, with no generation overlapping. We establish a maximum of K random

trials with reposition to find one compatible partner. The offspring is generated by *locus*

recombination of the parents and each *locus* has a probability of mutation (μ). We set q_{min} =

0.5q_0, where q_0 is the expected mean similarity within one population in equilibrium: q_0 =

$\frac{1}{1+4\mu K}$. For a detailed demonstration of the above equation see SI1. The restriction q_{min} =

0.5q_0 is arbitrary and only assures that no parasite speciation occurs when using a unique host

(i.e., avoids sympatric speciation in the context of Higgs and Derrida (1991)). Consequently,

parasite speciation only happens when more than one host species is used.

Temporal scaling

The empirical studies have evolutionary times in the order of millions of years, and to

maintain this time scale in the model would demand a high computational cost. As proposed

by Costa et al. (2019), in our approach we adopted a high value of mutation rate (μ=0.025) in

order to decrease the number of iterations (time steps or generations) necessary for speciation

to occur. Furthermore, we assumed that, due to the shorter life cycle of parasites, they have a

faster speciation rate when compared to their hosts (Dowton and Austin 1995; Light and

Hafner 2007). To satisfy these conditions, we rescaled the whole host phylogeny assuming

that the smaller branch length consists of the minimal time for parasites to speciate due to

isolation by host use (see the demonstration in SM1):

$$\tau_a = \frac{1}{4\mu} log(\frac{1}{q_{min}}). \qquad\qquad (2)$$

The minimal time for speciation decreases with q_{min}. Therefore, making the reproduction

more restricted (i.e., increasing q_{min}) facilitates the formation of parasite species in a shorter

time. The simulation starts with a clonal parasite population using a unique host species. Also,

the first host speciation occurred only after τ_a generations for the parasite populations to

accumulate genetic diversity before the first splitting event.

Host-switching events

After the first host speciation, parasite individuals in a host species may switch to

another host. For each parasite individual, we randomly selected a host species, including the

one in use. If the selected host species is not the original host (donor), we follow a probability

function for the host-switching event. This probability of host-switching events (P_{hs})

decreases over time, representing the product of opportunity for contact and compatibility of

the interaction of parasites associated with the evolutionary history of hosts (Fig. 1c). Then,

we are assuming that compatibility, the opportunity of interaction are expressed through the

evolutionary history of hosts. The probability of a parasite individual successfully migrates

(host-switching) from one host to another host species, in a given generation n, is defined as:

$P_{hs}(n)=exp[-r * (n - n_s)]$.   (3)

where r is a positive parameter that controls the decay of the host-switching probability, and

n_s is the generation that the common host ancestor had speciated (then, n-n_s is how long the

two host species had diverged). If $r = 0$, these probabilities are equal to 1 regardless of the

host divergence time, meaning that there is no restriction to host-switching. As a

consequence, parasite gene flow is continuous and speciation does not occur. At the other

extreme, for sufficiently large r values ($P_{hs}\sim0$), host-switching is absent and cospeciation

between hosts and parasites is expected. For intermediary r value, some parasite individuals

can eventually switch hosts (Fig. 2 and Fig. S1). This will increase the host repertoire of the

parasitic species, and also enable speciation by isolation (by host use), similar to the

speciation by founder's effect (Mayr 1999; Gavrilets and Hastings 1996). The effect of the

overall host-switching in a community does not depend only on r, but also on the

particularities of each host phylogeny that is used as input for the calculation of the host-

switching probability. Therefore, to better interpret the effect of parameter r on the trajectories

and compare the results between the communities, we do not present our results in terms of r,

but how much it changes the overall host-switching events. To obtain this overall metric, we

calculated the mean percentage of parasite individuals that switch hosts over the entire

simulation and we call it ***host-switching intensity*** (Fig. 2).

FIGURE 2. Relation between r (a parameter that defines the host-switching decay, Eq. 3) and

the intensity of host-switching for each simulated community. Each ID represents empirical

studies and S_H represents the host richness. ID. 1 - Birds and feather mites. ID. 2 - Mammals

and lice. ID. 3 - Wildlife and ectoparasites. ID. 4 and 5 - Rodents and fleas. ID. 6 - Fish and

Monogeneans (Gyrodactylidae). ID. 7 - Frogs and monogeneans (Polystomatidae). ID. 8 -

Frogs and lungworms (*Rhabdias* spp.). ID. 9 - Frogs and gut worms (*Oswaldocruzia* spp.).

Regional spatial scale studies are represented by salmon colour and local spatial scale studies

by blue.

Parameters of the model

For the results presented here we fixed the population size per host ($K = 250$) and

mutation probability per locus ($\mu = 0.025$). With these parameters, we can observe species

formation with reasonable computational time. Since the empirical studies varied in the

number of host species and branch size, the total number of iterations also varied (Table S1).

The parameter r varied ($0<r<1$) for each empirical study. A total of 50 replicates were

performed for each parameter combination. We have also analysed the model predictions

under other values of population size ($K = \{50, 500, 1000\}$) and mutation rate ($\mu = 0.001$)

(Table S1). Our qualitative conclusions did not change under these parameter variations (Fig.

S2-S4).

Validation with empirical data

The development of a new method to assess the host-switching intensity allowed us to

compare the results of our simulations with empirical data from different groups of parasites

and their respective hosts. This method uses information on the evolutionary history of the

host species as a proxy for resource similarity. We used nine studies from empirical studies of

parasite-host associations (Table 1) for comparative purposes. The selection criteria was that,

in addition to information on species interaction, these empirical studies essentially needed to

have phylogenies for hosts and parasites (see the details in Fig. S5-S13). We separated these

empirical studies according to the spatial scale (Table 1). Spatial scale refers to the spatial

extent of ecological processes and the spatial interpretation of the data. In this study, we

assumed that studies in the local spatial scale are essentially in a geographic radius less than

or equal to 35km, while on a regional scale they were collected essentially in a geographic

radius greater than 35km, in the original article respectively.

TABLE 1. Description of the host sample size and parasite richness for each empirical study, of

which host phylogenies were used as model parameters and host-parasite association to

validate the simulations of the model. Legend: ID = Empirical study.

ID	Host group	Host richness	Parasites group	Parasite richness	Spatial scale	Reference
1	Bird	11	Feather mites (*Trouessartia* spp.)	11	regional	Donã et al. 2017
2	Mammals	6	Lice (*Pediculus* spp. and *Pthirus* spp.)	7	regional	Reed et al. 2007
3	Wildlife	9	Arthropods*	8	regional	Becker et al. 2018
4	Rodents	129	Fleas*	202	regional	Krasnov et al. 2016
5	Rodents	11	Fleas*	19	local	Krasnov et al. 2016
6	Fish	7	Monogeneans (Gyrodactylidae)	16	local	Patella et al. 2017

7	Frogs	15	Monogeneans	13	regional	Badets et al. 2011
8	Frogs	31	Lungworms (*Rhabdias* spp.)	18	regional	Müller et al. 2018
9	Frogs	7	Gut worms (*Oswaldocruzia* spp.)	5	local	Willkens et al. 2021

Note: *Include different parasite groups

Characterization of the ecological and evolutionary patterns of parasites

We compared both the structure of turnover of parasite species (ecological pattern) and

the imbalance of parasite phylogenies (evolutionary pattern) in the empirical studies with

those resulting from the simulations. To characterise the composition of parasite species we

used the metric that gives information about the beta diversity of multiple-site dissimilarities

(β_{SOR} - Baselga 2010; 2013a, b). The beta diversity may reflect two different phenomena:

turnover (β_{SIM}) and nestedness (β_{NES}) (Baselga et al. 2007; Baselga 2010; 2013a, b). Here, we

choose only to work with the Simpson-based multiple-site dissimilarity, that is turnover

(β_{SIM}), since it is non-dependent on species richness (Baselga et al. 2007; Baselga 2010). This

refers to the replacement of some species by others as a consequence of environmental sorting

or spatial and historical constraints. In our case, we compared the variation in parasite species

composition between host species. The Simpson-based multiple-site dissimilarity is then:

$$\beta_{SIM} = \frac{\left[\sum_{i<j} min(b_{ij}, b_{ji})\right]}{2[\sum_i s_i - S_T] + \left[\sum_{i<j} min(b_{ij}, b_{ji})\right]} , \qquad (4)$$

where S_i is the total number of species in site i, S_T is the total number of species in all sites

(hosts in our case) and $min(b_{ij}, b_{ji})$ is the minimal number of species exclusive to sites i and j

in pairwise comparison (Baselga 2010).

To characterise the structure of the phylogenetic trees we used the metric that gives

information about the tree imbalance. Tree imbalance is one of the most common

phylogenetic structural patterns and measures asymmetries between the numbers of species on

each side of the tree's branches (Marquitti et al. 2020). Tree imbalance is widely measured

using the Sackin index (I) (Sackin 1972; Blum and François 2005; Frost and Volz 2013;

Dearlove and Frost 2015). The I has a dependence on the number of leaves, making it

unsuitable for comparing trees with different numbers of species. To make this comparison

possible, we use the normalised Sackin index (I_n) given by:

265              $$I_n(R) = \frac{I(R) - E[I(R)]}{\sqrt{\sigma_R^2}},\qquad\qquad (5)$$

where tree imbalance is the $I(R)$, and $E(I_n)$ and σ_R^2 the expected and variance of trees

generated by the Yule model which have the same number of leaves (species) as the observed

tree (Cardona et al. 2013; Marquitti et al. 2020). Although $I_n(R)$ would be close to zero for

trees generated with the Yule model, independent of the species richness R, different modes

of speciation may introduce important deviations from the behaviour of the Yule model

(Marquitti et al. 2020).

As each empirical study represents particular ecological and evolutionary processes, we

analysed whether there was an optimal range of host-switching intensity in our simulated

cases that retrieves information about turnover (β_{SIM}) and normalised Sackin index (I_n) of each

study. We considered that simulations that reproduced both the β_{SIM} and the I_n metrics

simultaneously (within a $\pm 5\%$ confidence interval) were the best fit to the empirical

examples. Then we compared the best fitting of host-switching intensity among the empirical

studies to understand how it varied for different evolutionary histories. Although species

extinctions occur in the model, this aspect was not included in the analyses since we do not

have information about extinctions in the empirical studies. These analyses were performed

using 'ape' (Paradis and Schliep 2019), 'betapart' (Baselga et al. 2018) 'picante' (Kembel et

al. 2010), 'phytools' (Revell 2012), and 'vegan' (Oksanen et al. 2013) R packages. See the

details in SI3.

Statistical analysis

To test whether the spatial scale modulates the best fitting host-switching intensities, a

linear mixed-effects model (LMM) was performed using the *lmer* function from the 'lme4'

package (Bates et al. 2015). We assumed the host-switching intensity as the response variable,

the spatial scale as a fixed variable, and empirical studies were treated as random variables

(intensity~ scale+(1|study)). After performing the LMM analysis, an analysis of variance

(ANOVA) was used to determine significant differences (p-value ≤ 0.01) using the Anova

function in the 'car' package (Fox and Weisberg 2019). All statistical analyses were

performed in R v.4.0.0 (R Core Team 2020) and Rstudio v.1.3.959 (RStudio Team 2020).

RESULTS

The turnover and normalised Sackin index of parasites varied according to the mean

percentage of parasite individuals that switch hosts during the entire history of the host

community (the host-switching intensity). To illustrate the turnover and normalised Sackin

index according to the host-switching intensity, we present an example of a model application

with fleas associated with rodents (ID. 5, Fig. 3a-c). As expected, turnover decreases as host-

switching intensity increases (Fig. 3a and Fig. S14). This occurs because the increase of host-

switching promotes the interaction of different host species with the same parasite species.

Additionally, for each value of host-switching intensity, there is a small variation in the

turnover (Fig. 3a and Fig. S14). The only exception was ID. 4, which resulted in a wide

variation in turnover under high host-switching intensity (Fig. S14).

FIGURE 3. Influence of host-switching events on the eco-evolutionary patterns of simulated

parasites for fleas associated with rodents (see Table 1 for details). Here we demonstrated the

relationship between: **a**. Host-switching intensity and turnover of parasite species (β_{SIM})

between host species; **b**. Host-switching intensity and parasite normalised Sackin index (I_n); **c**.

Relationship between β_{SIM}, I_n, and the host-switching intensity. **d**. Host-switching intensity

and parasite richness. The lines refer to empirical information of parasite (continuous) and

host (dotted). The colored dots are redundant with the x-axis scale of graphs (a) and (b) but

intend to guide the interpretation of (c). A total of 50 replicates were performed with 250

individuals for each configuration of the parameters of host-switching intensity.

As imposed by the model, the parasite richness ends the same as hosts in the absence of

host-switching (Fig. S16). But, for intermediary values of host-switching, parasites can

colonise the new host and then speciate, resulting in an overwhelming increase in parasite

speciation (see the dynamics in the movie available in S17, Fig. S18, and Fig. 3d).

When host-switching intensity is low (below 1%), the normalised Sackin index (I_n) for

the simulated parasite phylogenies results in the exactly same value as the one obtained from

the empirical phylogeny of the host (note the dashed line in Fig. 3b and also Fig. S15). This is

because the low host-switching intensity does not allow the establishment of the parasite in a

new host and, as a consequence, the simulated parasite phylogenies have the same normalised

Sackin index of the empirical host phylogeny. Colonisation followed by speciation is more

likely to occur under a higher host-switching intensity, in which the normalised Sackin index

varies over simulations even when they are under the same host-switching intensity (Fig. 3b

and Fig. S15). The wide variation in the normalised Sackin index for a given host-switching

intensity reveals that stochastic host-switching events, even if host-switching is more likely to

occur between closely related species, can change the structure of the resulting phylogenetic

tree. Despite not having a monotonic tendency, the normalised Sackin index tends towards

zero (balanced tree) as host-switching intensity goes to one, regardless of the community (Fig.

S15), resembling a neutral speciation scenario Yule model (Yule 1924; Aldous 2001).

For all empirical studies analysed, there is a range of host-switching intensity that

simultaneously reproduces the observed turnover and the parasite normalised Sackin index

(Fig. 4). As mentioned, both metrics are sensitive to host-switching intensity but each one

varies independently of the other (see in Fig. 4). Generally, the turnover and the parasite

normalised Sackin index obtained under high host-switching intensity (greater than 50%) are

340 far from the empirical pattern (see Fig. 4, the yellow dots rarely approach the intersection of

341 the solid lines).

FIGURE 4. The relationship between variation in the turnover of parasite species, normalised Sackin index of parasite species, and host-switching intensity for nine empirical studies: the parasites' turnover, measured by the metric turnover(β_{SIM}) on the y-axis and the

normalised Sackin index (I_n) on the x-axis. Each ID represents an empirical case. The lines refer to empirical information of parasite

(continuous) and host (dotted). Colour scales represent each percentage interval of host-switching intensity. A total of 50 runs were

performed with 250 individuals of carrying capacity, for each configuration of the parameters of host-switching intensity.

The simulated host-switching intensity that simultaneously fit parasite turnover

and normalized Sackin index recovered a range of 0.06% to 22.07% of host-switching

intensity through the analysed empirical studies. Within this range, the associations

between mammals and lice presented the lowest host-switching intensity (case ID. 2

with 0.07% - 1.13%), followed by that involving wildlife and arthropod parasites (case

ID. 3 with 0.43% - 2.69%), frogs and monogeneans (case ID. 7 with 0.22% - 3.71%),

frogs and lungworms (case ID. 8 with 1.99% - 4.94%), frogs and gut worms (case ID. 9

with 5.29% - 9.35%), birds and feather mites (case ID. 1 with 0.06% - 8.17%), fish and

monogeneans (case ID. 6 with 8.26% - 11.64%), - the highest intensities of host-

switching were observed between rodents and fleas (case ID. 5 with 14.45% - 16.87%

and case ID. 4 with 0.43% - 22.07%). We also observed that the host-switching events

are more frequent in studies conducted in a local scale (blue colour in Fig. 5) than in

regional scales (salmon colour in Fig. 3) (LMM: relationship of host-switching intensity

on spatial scale: beta= 0.08, SE= 0.01, df = 6.92, t= 5.25, p = 0.001, ANOVA: F =

27.56, p = 0.001, Fig. 5).

FIGURE 5. Calculated host-switching intensity among empirical studies. The boxplots

show the distributions of simulated host switching intensities for each empirical study.

Regional spatial scale studies are represented by salmon, and local scale studies are in

blue. The number in axis x represents the empirical studies: ID. 1 - Birds and feather

mites. ID. 2 - Mammals and lice. ID. 3 - Wildlife and ectoparasites. ID. 4 and 5 -

Rodents and fleas. ID. 6 - Fish and Monogeneans (Gyrodactylidae). ID. 7 - Frogs and

monogeneans (Polystomatidae). ID. 8 - Frogs and lungworms (*Rhabdias* spp.). ID. 9 -

Frogs and gut worms (*Oswaldocruzia* spp.).

DISCUSSION

In this study, we developed a novel methodological framework to understand how

the intensity of host-switching shapes some aspects of ecological and evolutionary

patterns of parasites, here characterised by species interaction turnover and tree

imbalance, respectively. Our three main results are 1) We found an optimum range of

host-switching intensity that can predict similar patterns as those observed in the

empirical studies, which validates our model; 2) The model showed that the increase of

host-switching intensity promoted an increase in turnover, but the tree imbalance did

not follow any monotonic tendency. Moreover, for a specific host-switching intensity,

we observed a small variation in the turnover and a wide variation in the tree imbalance;

3) The predicted values of host-switching intensity varied among the empirical studies

and those at a local spatial scale resulted in values higher than the ones at the regional

scale.

The fact that our model rebuilt the eco-evolutionary patterns of all empirical

studies supports the idea that host-switching mediated by host evolutionary proximity is

a good predictor of parasite associations. According to the framework of the Stockholm

Paradigm (Brooks et al. 2019), parasites can colonise new host species due to pre-

existing compatibility, which is expressed when there is an opportunity for contact.

Compatibility emerges greatly from the ancestral capacity in which both hosts and

parasites must be physiologically compatible to establish a long-term association

(Brooks and McLennan 2002; Kolbe et al. 2004; Brooks et al. 2019). Hence, for a given

lineage of the parasite, the closer (phylogenetically) the original and the new host

species, the greater the possibility that the adequate resource is conserved or is at least

similar. In fact, phylogenetic proximity has been widely recognized as a potential

criterion to anticipate new associations (Streicker et al. 2010; Damas et al. 2020; Filion

et al. 2022).

Another element of the Stockholm Paradigm (Brooks et al. 2019) that we

observed over the temporal dynamics of our model is the Oscillation Hypothesis (Janz

and Nylin 2008): parasites first increase their host repertoire (generalise) and then

speciate (specialise). In our model, at each time step, a parasite individual can switch

hosts, promoting the increase of host repertoire for the parasite species. However, as we

assume that the probability of host-switching decreases as hosts diverge, given time, the

probability of individuals from the same parasite species maintaining the gene flow

between those host species decreases, and parasite speciation is likely to occur (see the

dynamics in the movie available in S17 and S18). Consequently, in our model, host-

switches promote host repertoire oscillation, as hypothesised by Janz and Nylin (2008),

and favour parasites to speciate at a greater rate than their host, which is empirically

evidenced (Poulin and Morand 2000). Although the mean argument behind the

difference in speciation rate between host and parasite is the parasite's shorter life cycle,

we support the idea that frequent host exploitation is another important mechanism to

parasite diversification (Hay et al. 2020, Boeger et al. 2022).

The tree imbalance did not have a monotonic tendency and showed a wide

variation for a given host-switching intensity. This reveals that stochastic events can

change the evolutionary trajectory of parasites. Although our model assumed that host-

switching most likely occurs between closely related species, eventually, a parasite can

switch to a phylogenetically distant host, changing the diversification history

completely. This distant host-switching was observed in most of the empirical studies

presented here, where the parasites were able to colonise hosts from different genera (ID

2 and 6), families (IDs 1, 4, 5, 7, 8, and 9), and even order (ID 3). For example, in study

ID8 the *Rhabdias* lung-worm anuran parasites occurred mostly in Bufonidae hosts and

only one species in the Hylidae host (Müller et al. 2018). Species extinction is another

class of stochastic event present in our model that could contribute to the varied outputs

on parasite evolution. As we use data only of extant species, when a species goes extinct

all its history is lost, also impacting the imbalance of the tree (Costa et al. 2019,

Marquitti 2020).

Unlike the tree imbalance, our results showed that the turnover has a monotonic

tendency: it decreased as the host-switching intensity increased. This pattern was

expected since the model imposes that as host-switching intensity increases, the

limitation to use a new host decrease (Fig. 2). Moreover, we did not observe a wide

variation in the turnover for a given host-switching intensity over the model

replications. This reveals that those stochastics events mentioned before cannot produce

significant changes in the turnover. It probably occurs because when host-switching

occurs, it produces a decrease in the turnover no matter what parasite species switched

to what host species. In other words, the identities of the species are not relevant since

turnover emerges not from a given species characteristic, but from the similarities

between species, or even spatial and temporal amplitudes (Fallon et al. 2004, Baselga et

al. 2007, Baselga, et al. 2022). This reinforces the idea that species turnover is a robust

metric to compare species assemblages (Baselga, et al. 2022) and may also be a good

proxy for host-switching intensity.

The host-switching intensity varied across empirical studies and we observed that

it is higher in empirical studies of local spatial scale than regional spatial scale. This

evidence shows that the amplitude of the spatial scale is a fundamental factor in

determining the extent of host-switching. The opportunity for interaction is larger in

host empirical studies at a local scale, as this reduces the likelihood that barriers exist,

hampering the encounter of potential actors. This is evident when comparing rodent and

flea associations at regional (ID. 4) and local spatial scales (ID. 5). Similarly, since the

association of *Rhabdias* spp. and frogs (ID. 8) are defined geographically (and not by

host taxa) it was assumed that host-switching by ecological fitting was evolutionarily

more important than association with particular host taxa (Kuzmin et al. 2014; Müller et

al. 2018). Different intensities of host-switching observed in our results may also be

influenced by biological variations of the species that make up the empirical studies

analysed. For instance, these studies include a great diversity of organisms (fleas, lice,

feather mites, helminths, platyhelminthes), with profound differences in their biological

characteristics. Expanding analyses to a broader sample of empirical studies, including

variations in the type of parasitism (e.g., mono *vs.* heteroxenic, ecto *vs.* endoparasite)

and host attributes can provide important insights into key features related to the process

of incorporation of new hosts.

In nature, host-parasite systems are more complex than those modelled here.

Although the model can reconstruct eco-evolutionary patterns of empirical studies, our

model has some limitations. For example, the carrying capacity of all host species is the

same and the host's body size, their abundances, and spatial distribution were not

explicitly considered. The selective pressure is not explicitly modelled, contrary to what

we observe in nature (Krasnov et al. 2005, Krasnov et al. 2021). Furthermore, all

parasite individuals and species are equivalent, and may compete for the same

resources. Except for resource competition, our model didn't consider intra and

interspecific interactions among parasites. Finally, the phylogenies are still scarce,

especially for parasite species, which limited the number of tests with the model.

Phylogenetic data on parasites is extremely important to clarify the role of host-

switching in the ecological and evolutionary patterns of parasite lineages. Still, we

recovered compatible eco-evolutionary patterns for modelled parasites and their

respective hosts. Our model has important implications for predicting host switching,

especially in scenarios of anthropogenic change. With anthropogenic changes constantly

modifying natural environments and altering the geographic distribution of parasites,

many species that were once restricted to specific areas are now expanding their range

into new geographic locations and changing the composition of communities (see

Brooks et al. 2014). As we showed, parasites can follow different evolutionary paths,

and eventually can switch to non-related hosts, ultimately, determining the migration of

a parasite to other species (and clades), in some cases, including humans. To conclude,

we show that a model in which host-switching mediated by evolutionary proximity

between hosts is a predictor for parasitic associations over evolutionary time, as well as

for the origins of parasite diversity. We see this as an important step in our

understanding of parasite diversification processes.

SUPPLEMENTARY MATERIAL

Data SI

DATA AVAILABILITY

The model, phylogenies, and interactions of all analysed studies are available at

https://github.com/elviradbastiani/host_switching_model.

ACKNOWLEDGEMENTS

We thank the researchers who collected and reported the field data, our

institutions, and the many colleagues who helped us in different ways during this project

for their comments and suggestions, especially to Professor Dr. Marcus Aguiar. EDB is

grateful for the Ph.D. degree scholarship provided by Capes (Coordenação de

Aperfeiçoamento de Pessoal de Nível Superior). The authors acknowledge the

computational support from Professor Carlos M. de Carvalho at LFTC-DFis-UFPR.

FINANCIAL SUPPORT

EDB was supported by Brazilian Coordination for the Improvement of Higher

Education Personnel (CAPES). DP was supported by the São Paulo Research

Foundation (FAPESP), grants #2018/11187-8, #2019/24449-3, and #2016/01343-7

(ICTP-SAIFR). SBLA was supported by Conselho Nacional de Desenvolvimento

Científico e Tecnológico (CNPq: #11284/2021-3).

AUTHORS CONTRIBUTION

Conceived and designed the experiments: EDB, SBLA and DP

Performed the experiments and analysed the data: EDB, SBLA and DP

Wrote the paper: EDB, SLB, KMC, DP, WB, FMDM

Other contributions: EDB, DP, FMDM, WB, KMC and SLBA

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

Agosta S. J., Janz N., Brooks D. R. 2010. How specialists can be generalists: resolving

the "parasite paradox" and implications for emerging infectious disease.

Zoologia (Curitiba), 27(2), 151-162.

Agosta S. J., Klemens J. A. 2008. Ecological fitting by phenotypically flexible

genotypes: implications for species associations, community assembly and

evolution. Ecology Letters, 11, 1123-1134.

Agosta S. J., Brooks D. R. 2020. The major metaphors of evolution: Darwinism then

and now (Vol. 2). Springer Nature, Switzerland AG. pp. XVII - 273.

Aldous D. J. 2001. Stochastic models and descriptive statistics for phylogenetic trees,

from Yule to today. Statistical Science, 16, 23-34.3

Araujo S. B., Braga M. P., Brooks D. R., Agosta S. J., Hoberg E. P., von Hartenthal F.

W., Boeger W. A. 2015. Understanding host-switching by ecological fitting.

PLoS One, 10(10), e0139225.

Baselga A., Jiménez-Valverde A., Niccolini G. 2007. A multiple-site similarity measure

independent of richness. Biology Letters, 6, 642-645.

Baselga A. 2010. Partitioning the turnover and nestedness components of beta diversity.

Global ecology and biogeography, 19, 134-143.

Baselga A. 2013a. Separating the two components of abundance-based dissimilarity:

balanced changes in abundance vs. abundance gradients. Methods in Ecology

and Evolution, 4, 552-557.

Baselga A. 2013b. Multiple site dissimilarity quantifies compositional heterogeneity

among several sites, while average pairwise dissimilarity may be misleading.

Ecography, 36, 124-128.

Baselga A., Orme D., Villéger S., De Bortoli J., Leprieur F., Logez M., Henriques-Silva

R. 2018. Partitioning beta diversity into turnover and nestedness components.

Package 'betapart'. Version, 1(1).

Baselga A., Gómez-Rodríguez C., Araújo M. B., Castro-Insua A., Arenas M., Posada

D., Vogler A. P. 2022. Joint analysis of species and genetic variation to quantify

the role of dispersal and environmental constraints in community turnover.

Ecography, e05808.

Bates D., Maechler M., Bolker B., Walker S., Christensen R. H. B., Singmann H.

Grothendieck G. 2011. Package 'lme4'. Linear mixed-effects models using S4

classes. R package version, 1(6).

Blum M. G. François O. 2005. On statistical tests of phylogenetic tree imbalance: the

Sackin and other indices revisited. Mathematical biosciences, 195, 141-153.

Boeger W. A., Brooks D. R., Trivellone V., Agosta S., Hoberg E. 2022. Ecological

Super-Spreaders Drive Host-Range Oscillations: Omicron and Risk-Space for

Emerging infectious Disease. In press.

Boyd B. M., Nguyen N. P., Allen J. M., Waterhouse R. M., Vo K. B., Sweet A. D.,

Johnson K. P. 2022. Long-distance dispersal of pigeons and doves generated

new ecological opportunities for host-switching and adaptive radiation by their

parasites. Proceedings of the Royal Society B, 289(1970), 20220042.

Braga M. P. Janz N. 2021. Host repertoires and changing insect–plant interactions.

Ecological Entomology, 46, 1241–1253.

Brooks D. R., McLennan D. A. 2002. The nature of diversity. University of Chicago

Press.

Brooks D. R., Hoberg E. P. Boeger W. A. 2019. The Stockholm paradigm: climate

change and emerging disease. University of Chicago Press, pp. 400.

Brooks D. R., Hoberg E. P., Boeger W. A., Gardner S. L., Galbreath K. E., Herczeg D.

Dursahinhan A. T. 2014. Finding them before they find us: informatics,

parasites, and environments in accelerating climate change. Comparative

Parasitology, 81, 155-164.

Bush S. E., Sohn E. Clayton D. H. 2006. Ecomorphology of parasite attachment:

experiments with feather lice. Journal of Parasitology, 92, 25-31.

Cardona G., Mir A., Rosselló F. 2013. Exact formulas for the variance of several

balance indices under the Yule model. Journal of Mathematical Biology, 67(6),

1833-1846.

Charleston M. A. Robertson D. L. 2002. Preferential host switching by primate

lentiviruses can account for phylogenetic similarity with the primate phylogeny.

Systematic biology, 51, 528-535.

Costa C. L., Lemos-Costa P., Marquitti F. M., Fernandes L. D., Ramos, M. F.,

Schneider D. M. de Aguiar M. A. 2019. Signatures of microevolutionary

processes in phylogenetic patterns. Systematic biology, 68, 131-144.

Cuthill J. H. Charleston M. A. 2013. A simple model explains the dynamics of

preferential host switching among mammal RNA viruses. Evolution:

International Journal of Organic Evolution, 67, 980-990.

Damas J., Hughes G. M., Keough K. C., Painter C. A., Persky N. S., Corbo M. Lewin

H. A. 2020. Broad host range of SARS-CoV-2 predicted by comparative and

structural analysis of ACE2 in vertebrates. Proceedings of the National

Academy of Sciences, 117, 22311-22322.

De Vienne D. M., Refrégier G., López-Villavicencio M., Tellier A., Hood M. E. Giraud

T. 2013. Cospeciation *vs* host-shift speciation: methods for testing, evidence

from natural associations and relation to coevolution. New Phytologist, 198,

347-385.

Dearlove B. L., Frost S. D. 2015. Measuring asymmetry in time-stamped phylogenies.

PLoS computational biology, 11(7), e1004312.

Dominguez M., de la Colina M. A., Di Giacomo A. G., Reboreda J. C., Mahler B. 2015.

Host switching in cowbird brood parasites: how often does it occur? Journal of

evolutionary biology, 28(6), 1290-1297.

Dowton M. Austin A. D. 1995. Increased genetic diversity in mitochondrial genes is

correlated with the evolution of parasitism in the Hymenoptera. Journal of

Molecular Evolution, 41, 958-965.

Engelstädter J. Fortuna N. Z. 2019. The dynamics of preferential host switching: Host

601   phylogeny as a key predictor of parasite distribution. Evolution, 73, 1330-1340.

602   Fallon S. M., Ricklefs R. E., Latta S. C., Bermingham E. 2004. Temporal stability of

603   insular avian malarial parasite communities. Proceedings of the Royal Society of

604   London. Series B: Biological Sciences, 271(1538), 493-500.

605   Fecchio A., Wells K., Bell, J.A., Tkach V. V., Lutz H. L., Weckstein J. D. Clark N. J.

606   2019. Climate variation influences host specificity in avian malaria parasites.

607   Ecology Letters, 22, 547-557.

608   Filion A., Doherty J. F., Poulin R., Godfrey S. S. 2022. Building a comprehensive

609   phylogenetic framework in disease ecology. Trends in Parasitology.

610   Fox J. Weisberg S. 2019. Nonlinear Regression, Nonlinear Least Squares, and

611   Nonlinear Mixed Models in R. population, 150, 200.

612   Frost S. D., Volz E. M. 2013. Modelling tree shape and structure in viral

613   phylodynamics. Philosophical Transactions of the Royal Society B: Biological

614   Sciences, 368(1614), 20120208.

615   Gavrilets S. Hastings A. 1996. Founder effect speciation: a theoretical reassessment.

616   The American Naturalist, 147, 466-491.

617   Habermannová J., Bogusch P., Straka J. 2013. Flexible host choice and common host

618   switches in the evolution of generalist and specialist cuckoo bees (Anthophila:

619   Sphecodes). PLoS One, 8(5), e64537.

Hay E. M., Poulin R., Jorge F. 2020. Macroevolutionary dynamics of parasite

diversification: A reality check. Journal of Evolutionary Biology, 33(12), 1758-

1769.

Hayward A., Poulin R. Nakagawa S. 2021. A broadscale analysis of host-symbiont

cophylogeny reveals the drivers of phylogenetic congruence. Ecology Letters,

24, 1681–1696.

Higgs P. G., Derrida B. 1991. Stochastic models for species formation in evolving

populations. Journal of Physics A: Mathematical and General, 24(17), L985.

Imrie R. M., Roberts K. E. Longdon B. 2021. Between virus correlations in the outcome

of infection across host species: Evidence of virus by host species interactions.

Evolution letters, 5, 472-483.

Janz N., Nylin S. 2008. The oscillation hypothesis of host-plant range and speciation.

Specialization, speciation, and radiation: the evolutionary biology of herbivorous

insects, 2008, 203-215.

Jaramillo M. Rivera-Parra J. L. 2018. Host-Switching: How It Starts. In Disease

Ecology. Springer, Cham, pp. 139-156.

Kembel S.W., Cowan P.D., Helmus M. R., Cornwell W. K., Morlon H., Ackerly D. D.

Webb C. O. 2010. Picante: R tools for integrating phylogenies and ecology.

Bioinformatics, 26, 1463-1464.

Kolbe J. J., Glor R. E., Rodríguez Schettino L., Lara A. C., Larson A., Losos J. B. 2004.

Genetic variation increases during biological invasion by a Cuban lizard. Nature,

431(7005), 177-181.

Krasnov B. R., Burdelova N. V., Khokhlova I. S., Shenbrot G. I., Degen A. 2005.

Larval interspecific competition in two flea species parasitic on the same rodent

host. Ecological Entomology, 30(2), 146-155.

Krasnov B. R., Spickett A., Junker K., Van der Mescht L., Matthee S. 2021. Functional

and phylogenetic uniqueness of helminth and flea assemblages of two South

African rodents. International journal for parasitology, 51(10), 865-876.

Kuzmin Y., de Vasconcelos Melo F. T. Dos Santos J. N. 2014. A new species of

Serpentirhabdias Tkach, Kuzmin Snyder, 2014 (Nematoda: Rhabdiasidae)

parasitic in the brown ground snake *Atractus major* Boulenger (Reptilia:

Serpentes: Dipsadidae) in Brazil. Systematic Parasitology, 89, 101-106.

Light J. E. Hafner M. S. 2007. Cophylogeny and disparate rates of evolution in

sympatric lineages of chewing lice on pocket gophers. Molecular phylogenetics

and evolution, 45, 997-1013.

Manzo F., Peliti L. 1994. Geographic speciation in the Derrida-Higgs model of species

formation. Journal of Physics A: Mathematical and General, 27(21), 7079.

Marquitti F. M. D., Fernandes L. D., de Aguiar M. A. M. 2020. Allopatry increases the

balance of phylogenetic trees during radiation under neutral speciation.

Ecography, 43(10), 1487-1498.

Mayr E. 1999. Systematics and the origin of species, from the viewpoint of a zoologist.

Harvard University Press, pp. 372.

Meinilä M., Kuusela J., Ziętara M. S., Lumme J. 2004. Initial steps of speciation by

geographic isolation and host switch in salmonid pathogen *Gyrodactylus salaris*

(Monogenea: Gyrodactylidae). International Journal for Parasitology, 34(4),

515-526.

Nylin S., Slove J. Janz N. 2014. Host plant utilization, host range oscillations and

diversification in nymphalid butterflies: a phylogenetic investigation. Evolution,

68, 105-124.

Nylin S., Agosta S., Bensch S., Boeger W. A., Braga M. P., Brooks D. R. Janz, N.

2018. Embracing colonizations: a new paradigm for species association

dynamics. Trends in ecology evolution, 33, 4-14.

Oksanen J., Blanchet F. G., Kindt R., Legendre P., Minchin P. R., O'hara R. B.

Oksanen M. J. 2013. Package 'vegan'. Community ecology package, version, 2,

1-295.

Poulin R., Morand S. 2000. The diversity of parasites. The quarterly review of biology,

75(3), 277-293.

Revell L. J. 2012. phytools: an R package for phylogenetic comparative biology (and

other things). Methods in ecology and evolution, 3, 217-223.

Sackin M. J. 1972. "Good" and "bad" phenograms. Systematic Biology, 21(2), 225-226.

Sanaei E., Charlat S., Engelstädter J. 2021. *Wolbachia* host shifts: routes, mechanisms,

constraints and evolutionary consequences. Biological Reviews, 96(2), 433-453.

Streicker D. G., Turmelle A. S., Vonhof M. J., Kuzmin I. V., McCracken G. F.

Rupprecht C. E. 2010. Host phylogeny constrains cross-species emergence and

establishment of rabies virus in bats. Science, 329, 676-679.

Team R. 2020. RStudio: Integrated Development for R. RStudio, PBC, Boston, MA.

Available at: https://www.rstudio.com/. Last accessed 10 november 2021.

Team R. C. 2020. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria: Available at:

https://www. R-project. org/. Last accessed 10 november 2021.

Woolhouse M. E., Haydon D. T., Antia R. 2005. Emerging pathogens: the epidemiology

and evolution of species jumps. Trends in ecology & evolution, 20(5), 238-244.

Yule U. G. 1925. A mathematical theory of evolution, based on the conclusions of Dr.

JC Willis, FRS. Philosophical Transactions of the Royal Society of London

Series B, 213, 21-87.

[Dataset] Badets M., Whittington I., Lalubin F., Allienne J. F., Maspimby J. L., Bentz S.

Verneau O. 2011. Correlating early evolution of parasitic platyhelminths to

Gondwana breakup. Systematic Biology, 60, 762-781. Accession number:

EDB_10.1093/sysbio/syr078

[Dataset] Becker D. J., Streicker D. G. Altizer S. 2018. Using host species traits to

understand the consequences of resource provisioning for host–parasite

interactions. Journal of Animal Ecology, 87, 511-525. Accession number:

EDB_10.1111/1365-2656.12765

[Dataset] Doña J., Sweet A. D., Johnson K. P., Serrano D., Mironov S. Jovani R. 2017.

Cophylogenetic analyses reveal extensive host-shift speciation in a highly

specialized and host-specific symbiont system. Molecular phylogenetics and

evolution, 115, 190-196. Accession number:

EDB_10.1016/j.ympev.2017.08.005

[Dataset] Krasnov B. R., Shenbrot G. I., Khokhlova I. S. Degen A. A. 2016. Trait-

based and phylogenetic associations between parasites and their hosts: a case

study with small mammals and fleas in the Palearctic. Oikos, 125, 29-38.

Accession number: EDB_10.1111/oik.02178

[Dataset] Müller M. I., Morais D. H., Costa-Silva G. J., Aguiar A., Avila R.W. da Silva

R. J. 2018. Diversity in the genus *Rhabdias* (Nematoda, Rhabdiasidae):

Evidence for cryptic speciation. Zoologica Scripta, 47, 595-607. Accession

number: EDB_10.1111/zsc.12304

[Dataset] Patella L., Brooks D. R. Boeger W. A. 2017. Phylogeny and ecology

illuminate the evolution of associations under the Stockholm paradigm:

Aglaiogyrodactylus spp. (Platyhelminthes, Monogenoidea, Gyrodactylidae) and

species of Loricariidae (Actinopterygii, Siluriformes). Vie Et Milieu, 67, 91-102.

Accession number: EDB_20183235432

[Dataset] Reed D. L., Light J. E., Allen J. M. Kirchman J. J. 2007. Pair of lice lost or

parasites regained: the evolutionary history of anthropoid primate lice. Bmc

Biology, 5, 1-11. Accession number: EDB_10.1186/1741-7007-5-7

[Dataset] Willkens Y., Furtado A. P., Dos Santos J. N. de Vasconcelos Melo F. T.

2021. Do host habitat use and cospeciation matter in the evolution of

Oswaldocruzia (Nematoda, Molineidae) from neotropical amphibians? Journal

of Helminthology, 95, e33. Accession number:

EDB_10.1017/S0022149X21000250

730