Roflumilast, a Phosphodiesterase-4 Inhibitor, Ameliorates Sleep Deprivation-Induced Cognitive Dysfunction in C57BL/6J Mice

Abid Bhat[1,2], Muhammed Bishir[1,2], Seithikurippu R. Pandi-Perumal[3], Sulie Chang[4], Saravana B. Chidambaram[1,2,5*]

1. Department of Pharmacology, JSS College of Pharmacy, JSS Academy of Higher Education & Research, Mysuru – 570015, Karnataka, India
2. Centre for Experimental Pharmacology and Toxicology, JSS Academy of Higher Education & Research, Mysuru – 570015, Karnataka, India
3. Saveetha Medical College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, Tamil Nadu India.
4. Department of Biological Sciences and Institute of Neuroimmune Pharmacology, Seton Hall University, 400 South Orange Ave South Orange, NJ 07079, USA
5. Special Interest Group - Brain, Behaviour and Cognitive Neurosciences, JSS Academy of Higher Education & Research, Mysuru – 570015, Karnataka, India

[*] **Corresponding author**
Prof. Saravana Babu Chidambaram
MPharm, PhD, FASc (AW), FIC, FST
Dept. of Pharmacology, JSS College of Pharmacy &
Coordinator, Centre for Experimental Pharmacology & Toxicology
JSS Academy of Higher Education & Research
S.S. Nagar, Mysuru – 570015, Karnataka, India
Email: babupublications@gmail.com

ABSTRACT

Sleep deprivation (SD) interferes with long-term memory and cognitive functions by over-

activation of phosphodiesterase (PDEs) enzymes. PDE4, a non-redundant regulator of the

cyclic nucleotides (cAMP), is densely expressed in the hippocampus and is involved in learning

and memory processes. In the present study, we investigated the effects of Roflumilast (ROF),

a PDE4B inhibitor, on sleep deprivation induced cognitive dysfunction in a mouse model.

Memory assessment was performed using a novel object recognition task and the hippocampal

cAMP level was estimated by the ELISA method. The alterations in the expressions of PDE4B,

amyloid-beta ($A\beta$), CREB, BDNF, and synaptic proteins (Synapsin I, SAP 97, PSD 95) were

assessed to gain insights into the possible mechanisms of action of ROF using the Western blot

technique. Results show that ROF reversed SD induced cognitive decline in mice. ROF down-

regulated PDE4B and $A\beta$ expressions in the brain. Additionally, ROF improved the cAMP

level and the protein expressions of synapsin I, SAP 97, and PSD 95 in the hippocampal region

of SD mice. Taken together, these results suggest that ROF can suppress the deleterious effects

of SD-induced cognitive dysfunction via the PDE4B-mediated cAMP/CREB/BDNF signaling

cascade.

Keywords: Sleep deprivation, PDE4B, Roflumilast, Memory, cAMP, Synaptic proteins

1. INTRODUCTION

Sleep plays a regulatory role in maintaining cellular and metabolic homeostasis. Increasing evidence reveals that sleep disturbances affect higher-order brain functions that are linked to various neurological disorders [1–3]. Sleep deprivation (SD) has a detrimental influence on both social and economic well-being [4,5]. Accumulating evidence suggests that SD reduces neurogenesis and protein expression of neurotrophic factors such as CREB and BDNF, which are the crucial regulators for learning and memory and their down-regulation was reported to induce hippocampal atrophy [6–8]. Functional magnetic resonance imaging (fMRI) and behavioral studies showed that one-night sleep deprivation substantially compromises hippocampal function in humans in turn affecting memory [9]. Furthermore, sleep disturbance predisposes neuronal accumulation of amyloid-β (Aβ) [10–12] in the brain. Positron emission tomography in humans also confirmed that SD and sleep fragmentation are associated with increased deposition of Aβ in the brain [13,14].

A mechanistic research report showed that SD promotes the synthesis and impairs the neuronal clearance of Aβ protein [15]. Intriguingly, the relationship between SD and Aβ is bidirectional because the other way around increased Aβ deposition impairs slow wave sleep [16]. Furthermore, Aβ reduces the protein expression of Synapsin I, PSD-95, and SAP-102, which indicates that it eliminates synapses and disrupts neuronal networks [17–19]. This synapto-toxic effect of Aβ is linked to the reduced expression of NMDA receptors and decreased cAMP content [20,21]. The increased accumulation of Aβ and decreased levels of cAMP decreases the release of neurotrophic factors that regulate brain development and synaptic plasticity [22,23].

Phosphodiesterases (PDEs) are a diverse family of enzymes that play a pivotal role in regulating intracellular signaling [24]. An increased expression of PDE4 enzymes hydrolyse the cAMP into its inactive form 5-AMP, which has been consistently observed in Alzheimer's disease (AD) patients' brains [25], and subjects with cognitive impairment [26] and also in the hippocampal region of SD mice [27]. PDE4s isoforms such as *PDE4A*, *PDE4B*, and *PDE4D* are expressed in the brain while *PDE4C* is predominantly expressed in testis, and lacks expression in the brain[28]. PDE4B is highly expressed in rodent hippocampal CA2 and CA3 regions, parietal and piriform cortex, and the cerebellar granular layer [29]. Inhibition of PDE4 improved learning and memory in a mouse model of AD via increasing hippocampal cAMP levels [25,30]. Furthermore, inhibition of PDE4 has also restored the deficits in synaptic proteins such as synaptophysin, and PSD 95 [31,32]. Specifically, PDE4 inhibition was shown to reverse the cognitive decline induced by muscarinic receptor antagonist [33] and also by modulating NMDA receptors mediated transduction mechanisms in rat models [34]. Albeit, the NMDA does not

affect PDE4 expression directly, however, the balance between PDE4 and NMDA mediated
adenylyl cyclase plays a pivotal role in the memory process [34]. Thus, enormous reports suggest
that PDE4s are the potential target in neurological disorders drug discovery [35].

Roflumilast (ROF), a cAMP-specific PDE4B inhibitor, is approved by USFDA for use in
chronic obstructive pulmonary disease (COPD) [36]. ROF promotes hippocampal neuron
viability [37] and improves memory in rodents and monkeys at the non-emetic doses [38]. In a
clinical study, it is observed that acute administration (3 days) of ROF improves learning and
memory in healthy subjects [39]. Long-term SD produces an AD-like pathological state, wherein
increased neuronal accumulations of Aβ, decreased cAMP, and decreased synaptic protein
expressions are well established. This spurs interest to investigate whether PDE4 expression
has any correlation with Aβ, CREB, BDNF expression in SD brains, and also to study the effect
of PDE4B inhibition, using ROF, on cognitive function in sleep-deprived mice. **Figure.1**
explains the flow of the present study's experimental design.

2. RESULTS

2.1. SD up-regulated PDE4B expression in the mice hippocampal region

In the present study, we assessed the impact of SD on PDE4B expression in the hippocampal
region of mice brains. 72h of continuous SD induced a significant $(p < 0.05)$ increase in
hippocampal PDE4B expression when compared to the NSD group. The daily dose of ROF
down-regulated PDE4B expression when compared to the vehicle-treated SD group. A
significant $(p < 0.01)$ decrease in PDE4B expression was found at 3 mg/kg dose of ROF (**Fig**
2A).

Impaired sleep is linked to AD as sleep plays a role in clearing the metabolic waste from the
brain [11]. We performed a Western blotting assay to study the impact of SD on Aβ expression
in mice hippocampal region. Mice subjected to 72h of SD showed a significant $(p < 0.01)$
increase in Aβ expression when compared to the non-sleep-deprived mice. Inhibition of
PDE4B by ROF significantly $(p < 0.01)$ reduced the expression of Aβ in mice when compared
with the vehicle-treated SD group.

2.2. SD increased Aβ deposition in hippocampal CA1 and DG regions of mice brains and
roflumilast decreased Aβ deposition

An increase in Aβ expression was further confirmed by Congo red staining of SD mice brains.
We found that vehicle-treated SD mice showed multifocal and intense deposition of Aβ in CA1

and DG regions when compared to the NSD group. ROF-treated SD mice showed very mild deposition of Aβ in CA1 and DG regions when compared to vehicle-treated SD mice brains **(Fig.3)**.

2.3. Roflumilast improves hippocampal cAMP levels in sleep-deprived mice

cAMP regulates the learning and memory processes in the brain. The decline of cAMP levels in the hippocampus impairs memory consolidation [40]. To determine the impact of SD on cAMP levels in the hippocampus region of mice, we performed an ELISA assay and found that SD significantly $(p < 0.01)$ reduced cAMP levels in vehicle-treated mice when compared with the NSD group. Administration of ROF in sleep-deprived mice showed a significant $(p < 0.001)$ increase in the levels of cAMP when compared with the vehicle-treated SD group. This indicates that ROF can rescue the cAMP levels in SD mice brains. **(Fig. 4).**

2.4. Roflumilast improved the neurotrophic factors - CREB, and BDNF proteins expression in the hippocampal region of SD mice

Next, we assessed the impact of SD on the protein expression of CREB and BDNF in the hippocampal region of mice brains. We found that 72-h SD produced a significant $(p < 0.01)$ decrease in pCREB expression as compared to the NSD group. Administration of ROF restored pCREB expression when compared to the SD control group. A significant $(p < 0.001)$ increase in pCREB was observed with ROF treatment when compared with vehicle-treated SD mice **(Fig. 5A).**

CREB positively influences the expression of BDNF, which is essential in memory consolidation and synaptic functions [6]. Hence we performed Western blot analysis to detect BDNF expression in the hippocampus region of SD mice. As shown in **Fig. 5B,** SD significantly $(p < 0.001)$ decreased the BDNF protein expression when compared with NSD mice. ROF administration in sleep-deprived mice significantly $(p < 0.001)$ increased BDNF expression when compared to vehicle-treated SD mice. These data suggest that ROF administration improves the protein expression of neurotropic factors CREB and BDNF in SD mice.

2.5. Roflumilast up-regulates the expression of the synaptic proteins in sleep-deprived mice brains

Next, we investigated whether improvement in neurotrophic factors expression has any
influence on the expression of the synaptic proteins in the hippocampal region of SD mice.
Synapsin I expression decreased significantly ($p < 0.01$) following SD when compared to the
NSD control group. ROF administration significantly ($p < 0.01$) up-regulated the expression of
Synapsin I when compared to vehicle-treated SD mice **(Fig 6).** SAP 97 regulates synaptic
plasticity by controlling the distribution of NMDA- and AMPA-type glutamate receptors [41].
We found a significant decrease in the expression of SAP-97 in SD mice which was reversed
($p < 0.001$) with ROF treatment **(Fig 6).**

Concurrently, we investigated the impact of SD on the post-synaptic density protein, PSD-95,
expression. 72 h of SD caused a significant ($p < 0.01$) decrease in PSD-95 expression when
compared to NSD mice. Administration of ROF significantly up-regulated ($p < 0.01$) PSD95
expression in SD mice **(Fig 6)**. These results imply that PDE4B inhibition can improve synaptic
functions, which could be corroborated by the restoration of neurotrophic factors, at least
partly, in SD.

168
169
170
171

2.6. Roflumilast restores sleep-deprivation induced cognitive dysfunction in mice
Recognition memory in mice was assessed using NORT. Sleep-deprived mice showed a
significant ($p < 0.05$) decrease in time spent with the novel object when compared to NSD
mice. Administration of ROF significantly (1mg/kg; $p < 0.05$, 3mg/kg; $p < 0.01$) increased time
spent with the novel object by SD mice **(Fig. 7a)**. SD significantly ($p < 0.01$) decreased the
discrimination index, but ROF significantly (1mg/kg; $p < 0.05$, 3mg/kg; $p < 0.01$) increased
the discrimination index in SD mice **(Fig. 7b)**. Decreased Aβ, and increased CREB / BDNF
and synaptic proteins (synapsin-1, SAP-97, and PSD95) expression are found to corroborate to
the improved memory in SD mice.

2.7. Roflumilast prevented the morphological changes of hippocampal neurons in sleep-
deprived mice
SD induced multifocal moderate neuronal degeneration in the CA1 and DG regions of the
hippocampus. As shown in **Fig 8**, neurons in CA1 and DG regions in vehicle-treated SD mice
showed a decrease in purple-stained Nissl granules and increased pyknotic nuclei in the

perikarya when compared to NSD mice. ROF-treated mice showed reduced morphological

changes and had regularly shaped neuronal bodies in CA1 and DG regions when compared to

SD mice **(Fig 8).**

3. DISCUSSION

The present study reveals that alleviation of Aβ pathology, cAMP signaling, and synaptic

proteins expression by ROF via PDE4B inhibition is the key mechanism in cognitive

restoration in SD mice.

SD aggravates Aβ plaque levels in AD transgenic mouse model [42]. A recent study has shown

that chronic sleep restriction (3 h per day, 5 days per week, for 4 weeks) increases the

hippocampal accumulation of Aβ plaques in mice brains, which was corroborated by the

cognitive decline [43]. Aβ deposition in cortical and hippocampal regions initiates inflammatory

responses, synaptic dysfunctions [44], and neuronal apoptosis [45]. Aβ also compromises the

cAMP-response element-binding protein (CREBP) signaling in neurons suggesting that

multiple factors contribute to neuronal damage in SD [46].

On the other hand, increased expression of PDE4D is reported to associate with Aβ plaque

pathology and memory loss [47]. Studies using PDE4D knockout mice showed improved memory

and reduced neuroinflammation and β-amyloidosis [48–50]. In the present study, we found that

72h SD increased the protein expression and/or neuronal deposition of Aβ in hippocampal

neurons of mice. Earlier, in vivo studies and clinical trials have shown that alleviating cAMP

signaling through PDE4 inhibition improves cognitive functions [51,52] and reduced Aβ

expression in mice [53]. The present study reports an increased PDE4B and Aβ expression in SD

mice brains with a significant decrease in cAMP levels. Treatment with ROF reversed the

molecular changes including Aβ pathology and improved cognitive performance in SD mice

which could be related to PDE4B inhibition.

Synapse dysfunction is an immediate consequence of increased Aβ deposition [54]. PSD-95 is an

essential regulator of synaptic strength and plasticity. SAP-97 facilitates synaptic vesicles

biogenesis, and neurotransmitters release [55]. Synapsin I regulates neurotransmitters release and

its down-regulation is shown to impair neurogenesis and synaptic plasticity [56,57]. SD causes

synaptic damage in the hippocampal region by reducing the expression of presynaptic and

postsynaptic proteins [58–60]. Recently, we have shown that ROF improves synaptic protein

expression using human neurons (in vitro model) exposed to the neurotoxin quinolinic acid [31].

In the present in vivo study, we report that SD down-regulates synaptic proteins (SAP-97, Synapsin-I, and PSD-95) in the hippocampal region of mice. Nevertheless, administration of ROF increased SAP-97, Synapsin-I, and PSD-95 expression, which indicates that PDE4B plays a role in synaptic protein expression.

Inhibition of cAMP signaling in the hippocampal region is reported to impair the consolidation of long-term memory in mice [61]. SD reduces the phosphorylation of CREB in the hippocampus and affects the protein expression of the BDNF [6]. In the present study, ROF administration restores the cAMP/CREB/BDNF signaling cascade in SD mice. These results are in agreement with earlier studies that report that inhibition of PDE4 improves memory consolidation in rodent models via the cAMP/CREB/BDNF cascade [33,62].

Histopathological examination using Congo red staining showed intense deposition of Aβ plaques in the CA1 and DG regions of the hippocampus in SD mice. Further, cresyl violet staining showed neuronal death in the hippocampus region of SD mice. Recent studies have shown that SD increases cytokine production, and microglial activation and initiates neuronal apoptosis, causing lesions in the hippocampus of mice [63,64]. Increased PDE4B expression is reported to trigger inflammation and microglial reactivity [65]. Treatment with ROF reduced Aβ plaques and also increased the Nissl granules in the hippocampal regions in SD mice. Inhibition of PDE4 transiently increases cAMP and stimulates cAMP signal transduction, which in turn, reduces the levels of pro-inflammatory and early apoptosis factors [66]. This could be a possible mechanism for the observed neuroprotection with ROF in SD

4. CONCLUSION

From the current investigation, we provide molecular, behavioral, and histopathological evidence that substantiates the protective effects of Roflumilast against SD-induced cognitive dysfunction in a mouse model. Roflumilast administration improves recognition memory via PDE4B inhibition which was found to be mediated via cAMP/CREB/BDNF signaling and downregulation of Aβ pathology in SD mice. Additionally, further studies to understand the effect of Roflumilast on NMDA activity and autophagy in chronic sleep restriction are under investigation in our lab.

5. MATERIALS AND METHODS

5.1. Animals.

Male C57BL/6J mice (25-30 g) were obtained from M/s. Adita Biosys Private Limited,
Tumakuru, Karnataka, and housed in CPT, JSS Academy of Higher Education & Research,
Mysuru, Karnataka. Animals were housed in groups (5 mice/cage) in polypropylene cages
under an ambient temperature of 19-25°C and 40-65% relative humidity, with a 12-h light/dark
artificial light cycle. Animals were provided with standard rodent feed and purified water ad
libitum. Animals were acclimatized for 7 days to the laboratory conditions prior to initiation of
the experiments. Animal experiments were performed in full compliance with the guidelines
of the "Guide for the Care and Use of Laboratory Animals" National Research Council, 2011).
Institutional Animal Ethics Committee (IAEC), Central Animal Facility, JSS AHER, Mysuru,
India, has approved the study protocol (JSSAHER/CPT/IAEC/014/2020).
263
5.2. Reagents and antibodies
Roflumilast, Cresyl violet, and Congo red stains were purchased from Sigma Aldrich (India).
cAMP ELISA kit was purchased from Cayman (Ann Arbor, MI, USA). Anti-PSD95 (sc-
32290), Anti-SAP97 (sc-9961), Anti-Synapsin-I (sc-376623), Anti-BDNF (sc-65514), Anti-
CREB (sc-377154), Anti β-Amyloid (sc-28365) were procured from Santa Cruz
Biotechnology, CA, USA. Anti-PDE4B (NB100-2562) was purchased from Novus
Biologicals, United States. All other reagents and chemicals were analytical grade.
271
272 *5.3. Roflumilast treatment*
273 Roflumilast [ROF (L): 1 mg/kg and ROF (H): 3 mg/kg] was freshly prepared in 0.5% carboxy
274 methyl cellulose (CMC) and administered intraperitoneally once a day for three days. The
275 doses of ROF was selected based on earlier published report[67].
276
5.4. Sleep deprivation method
Sleep deprivation of mice was done via a modified multiple platform method[68]. Briefly, a
mouse was placed on a cylindrical platform (8.5 cm in height and 2.5 cm in diameter). Eight
platforms were kept in a cage approximately 10 cm apart inside a cage filled with water up to
2 cm from the bottom. Water was changed twice a day. The size of the platforms permits the
mouse to sit but not lie down. Furthermore, mice could easily move between the platform but
could not stretch across the platforms. This method is reported to eliminate REM sleep[69]. Non-
sleep-deprived (NSD) animals were kept in normal polypropylene cages in the same room.

During sleep deprivation, mice had access to food and drinking water. Animals were sleep-
deprived for 72 hours.

5.5. Groups and Treatments

Animals were randomized in to four groups based on the stratified bodyweight viz Group 1-:
Non-sleep-deprived (NSD) mice which received 0.5% carboxy methyl cellulose [0.5% CMC;
n=10]; Group – II: SD which received 0.5% CMC; n=10; Group III and IV: SD which received
ROF (L) (1 mg/kg b.wt; n=10) and ROF (H) (3 mg/kg b.wt; n=10), respectively.

5.6. Novel object recognition test

A novel object recognition test was performed to access the recognition memory in rodents [70].
Briefly, the mouse was individually habituated to exploring the empty apparatus for 10 min
(2nd day of treatment). During the acquisition trial (T1), two similar objects were placed inside
the apparatus and the mouse was allowed to explore the objects for 3 min. After the acquisition
trial, the mouse was transferred to its home cage. The discrimination trial (T2) was done
twenty-four hours later (4th day). Two different objects, a familiar object, and a novel object
were placed in the exploration area. The time spent by the animal exploring the two objects
during T2 was recorded and the discrimination index (DI) was calculated, as per the following
formula. DI=RI/ (Time spent in exploring novel object + Time spent in exploring familiar
object). Recognition Index = Time spent in exploring novel objects-Time spent in exploring
the familiar objects. This test was repeated for all the animals in all the cages one at a time. All
behavioral assessments were done between 10.00 am to 1.00 pm.

5.7. Measurement of cAMP content

The hippocampal region of the mouse was isolated and homogenized using 0.1M HCl (10%
tissue homogenate). Homogenates were centrifuged for 10 min at 1500× g at 4°C, and the
supernatants were stored at 4°C. cAMP levels were determined by a cAMP enzyme
immunoassay kit following the manufacturer's instructions (Cayman Chemical Co., Ann
Arbor, MI, USA)

5.8. Western blot

Following the behavioral assessment, the animals were euthanized to collect the brain and stored at -80°C. Hippocampal region was isolated, and homogenates were prepared with radioimmunoprecipitation assay buffer (RIPA) buffer (50 mM Tris, pH 7.4, 150 mM NaCl, 1% NP-40, 5 mM EDTA, 0.5% sodium deoxycholate, 0.1% SDS, 50 nM sodium fluoride, 1 mM sodium vanadate) containing a cocktail of protease inhibitor (Sigma Aldrich, MO, USA). Total protein concentrations of the samples were determined by the Pierce™ bicinchoninic acid (BCA) protein assay (Thermofisher Scientific, USA), and samples were aliquoted and stored at -80°C until further use. Sample proteins (20 µg) were separated by using 10% bis-tris -SDS-PAGE (electrophoresis). Resolved proteins in the gels were transferred onto polyvinylidene difluoride (PVDF) membranes (Biorad, USA) and electroblotted. Membranes were blocked overnight with 5% non-fat skimmed milk in Tris-Buffered Saline and Tween 20 (TBST) at 4°C. This was followed by a 4-h incubation with the primary antibodies (PDE4B (1:1000), CREB (1:1000), BDNF (1:1000), β-Amyloid (1:1000), PSD-95 (1:1000), Synapsin-I (1:1000), SAP 97 (1:1000) at room temperature. The membranes were rinsed with TBST (3 washings for 10 min each), followed by incubation with the secondary antibodies (HRP conjugated anti-mouse or anti-rabbit IgG) for 1h at room temperature and washed with TBST (3 washings for 10 minutes each). Bands were detected using SuperSignal West Pico PLUS Chemiluminescent Substrate (Thermo Scientific, USA). Densitometric measurement of bands was done using ImageJ (NIH software). For Western blot analysis, the signal intensity (integrated density value, IDV) of PDE4B, BDNF, β-Amyloid, PSD-95, Synapsin-I, and SAP 97 was normalized against the IDV of internal control β-actin, while pCREB was normalized with total CREB and histogram was plotted.

5.9. Histopathology

Whole-brain was stored in buffered 10% formalin for 48h. Coronal sections (3-5 µm) of the hippocampus region were cut using a microtome. The hippocampal region of the brain was used for histopathological analysis. Sections were mounted on a slide, washed, and dehydrated with 95% ethanol.

5.10.   Cresyl violet staining

Coronal sections (3-5 µm) of the hippocampus region were washed with xylene and followed by five times washing with water at 5 min intervals. The samples were stained using 0.2% cresyl violet dye for 30 min. The prepared slides were examined under the microscope by a pathologist, blinded to the treatment protocol, for histopathological examination.

5.11. Congo red staining

Congo red staining was performed to detect the deposition of amyloidβ plaques in mice hippocampus. Coronal sections (3-5 µm) of the hippocampus region were stained with 1% Congo Red stain for 30 min. Amyloidβ plaques were observed under a microscope.

5.12. Statistical analysis

Data are presented as mean ±SEM. The difference between time spent exploring the novel object versus a familiar object during the discrimination trial was calculated for each group and the level of significance was analyzed using a two-sided student's t-test. For other parameters, group means differences were analyzed using a one-way ANOVA test followed by Tukey's multiple comparison test as post hoc. Graphs were plotted using GraphPad Prism version 7.04 with a *p-value* ≤ 0.05 is considered significant.

ACKNOWLEDGMENTS

AB wishes to thank the Indian Council of Medical Research (ICMR, New Delhi, India) for awarding him a Senior Research Fellowship (45/07/2019/PHA/BMS) during his Ph.D. degree. The authors would like to thank Shri Panchaksharappa Gowda DH, Biostatistician, JSS College of Pharamcy, Mysuru, for his assistance with the biostatistical analysis. All of the authors thank their respective institutions for the administrative and technical assistance provided during the study duration and manuscript preparation.

CONFLICT OF INTEREST

 The authors declare no conflict of interest.

AVAILABILITY OF DATA AND MATERIAL

The data sets generated and/or analyzed during the study are available in CPT, JSSAHER.

ETHICS APPROVAL

The animal experiment protocol was approved (JSSAHER/CPT/IAEC/014/2020) by the Institutional Animal Ethics Committee, JSS AHER, Mysuru, India.

AUTHOR CONTRIBUTIONS STATEMENT

AB: Methodology, Investigation, Data curation, Formal analysis, Writing – original draft, **MB**: Methodology, Investigation, **SRP:** Writing – review & editing, **SC:** Writing – review & editing, **SBC**: Conceptualization, Project administration, – review & editing. The manuscript has been read and approved by all authors.

REFERENCES

1. Bhat, A., Pires, A. S., Tan, V., Babu Chidambaram, S., Guillemin, G. J. (2020) Effects of Sleep Deprivation on the Tryptophan Metabolism. *Int J Tryptophan Res* 13, 1178646920970902

2. Bishir, M. Bhat, A., Essa, M.M., Ekpo, O., Ihunwo, A.O., Veeraraghavan, V.P., Mohan, S.K., Mahalakshmi, A.M., Ray, B., Tuladhar, S., Chang, S., Chidambaram, S.B., Sakharkar, M.K., Guillemin, G.J., Qoronfleh, M.W., Ojcius, D.J., (2020) Sleep Deprivation and Neurological Disorders. *BioMed Research International* vol. 2020 e5764017.

3. You, J.C., Jones, E., Cross, D.E., Lyon, A.C., Kang, H., Newberg, A.B., Lippa, C.F. (2019) Association of β-Amyloid Burden With Sleep Dysfunction and Cognitive Impairment in Elderly Individuals With Cognitive Disorders. *JAMA Network Open* **2**, e1913383.

4. Chattu, V.K., Manzar, Md.D., Kumary, S., Burman, D., Spence, D.W., Pandi-Perumal, S.R. (2018) The Global Problem of Insufficient Sleep and Its Serious Public Health Implications. *Healthcare (Basel)* **7** (1), 1.

5. Wang, W., Yang, L., Liu, T., Ma, Y., Huang, S., He, M., Wang, J., Wen, A., Ding, Y.,

(2020) Corilagin ameliorates sleep deprivation-induced memory impairments by

inhibiting NOX2 and activating Nrf2. *Brain Research Bulletin* 160, 141–149.

6. Alhaider, I. A., Aleisa, A. M., Tran, T. T., Alkadhi, K. A. (2011) Sleep deprivation

prevents stimulation-induced increases of levels of P-CREB and BDNF: Protection by

caffeine. *Molecular and Cellular Neuroscience* 46 (4), 742–751.

7. Cao, Y., Li, Qinglin, Liu, L., Wu, H., Huang, F., Wang, C., Lan, Y., Zheng, F., Xing, F.,

Zhou, Q., Li, Qi, Shi, H., Zhang, B., Wang, Z., Wu, X. (2019) M ptosis in mice with

sleep deprivation. *British Journal of Pharmacology* 176 (9), 1282–1297.

8. Havekes, R., Park, A.J., Tudor, J.C., Luczak, V.G., Hansen, R.T., Ferri, S.L.,

Bruinenberg, V.M., Poplawski, S.G., Day, J.P., Aton, S.J., Radwańska, K., Meerlo, P.,

Houslay, M.D., Baillie, G.S., Abel, T. (2016) Sleep deprivation causes memory deficits

by negatively impacting neuronal connectivity in hippocampal area CA1. *eLife* 5,

e13424.

9. Yoo, S.S., Hu, P. T., Gujar, N., Jolesz, F. A., Walker, M. P. (2007) A deficit in the ability

to form new human memories without sleep. *Nature Neuroscience* 10, 385–392.

10. Ooms, S., Overeem, S., Besse, K., Rikkert, M.O., Verbeek, M., Claassen, J.A.H.R.

(2014) Effect of 1 night of total sleep deprivation on cerebrospinal fluid β-amyloid 42 in

healthy middle-aged men: a randomized clinical trial. *JAMA Neurol* 71 (8), 971–977.

11. Shokri-Kojori, E., Wang, G.-J., Wiers, C.E., Demiral, S.B., Guo, M., Kim, S.W.,

Lindgren, E., Ramirez, V., Zehra, A., Freeman, C., Miller, G., Manza, P., Srivastava, T.,

Santi, S.D., Tomasi, D., Benveniste, H., Volkow, N.D. (2018) β-Amyloid accumulation

in the human brain after one night of sleep deprivation. *PNAS* 115 (17), 4483–4488.

12. Winer, J.R., Mander, B.A., Kumar, S., Reed, M., Baker, S.L., Jagust, W.J., Walker, M.P.

(2020) Sleep Disturbance Forecasts β-Amyloid Accumulation across Subsequent Years.

Curr Biol 30 (21), 4291-4298.e3.

13. Sharma, R.A., Varga, A.W., Bubu, O.M., Pirraglia, E., Kam, K., Parekh, A., Wohlleber,

M., Miller, M.D., Andrade, A., Lewis, C., Tweardy, S., Buj, M., Yau, P.L., Sadda, R.,

Mosconi, L., Li, Y., Butler, T., Glodzik, L., Fieremans, E., Babb, J.S., Blennow, K.,

Zetterberg, H., Lu, S.E., Badia, S.G., Romero, S., Rosenzweig, I., Gosselin, N., Jean-

Louis, G., Rapoport, D.M., de Leon, M.J., Ayappa, I., Osorio, R.S. (2018) Obstructive

Sleep Apnea Severity Affects Amyloid Burden in Cognitively Normal Elderly. A

Longitudinal Study. *Am J Respir Crit Care Med* 197 (7), 933–943.

14. Sprecher, K.E., Koscik, R.L., Carlsson, C.M., Zetterberg, H., Blennow, K., Okonkwo,

O.C., Sager, M.A., Asthana, S., Johnson, S.C., Benca, R.M., Bendlin, B.B. (2017) Poor

sleep is associated with CSF biomarkers of amyloid pathology in cognitively normal

adults. *Neurology* 89 (5), 445–453.

15. Lucey, B.P., Mawuenyega, K.G., Patterson, B.W., Elbert, D.L., Ovod, V., Kasten, T.,

Morris, J.C., Bateman, R.J. (2017) Associations Between β-Amyloid Kinetics and the β-

Amyloid Diurnal Pattern in the Central Nervous System. *JAMA Neurol* 74 (2), 207–215.

16. Mander, B.A., Marks, S.M., Vogel, J.W., Rao, V., Lu, B., Saletin, J.M., Ancoli-Israel, S.,

Jagust, W.J., Walker, M.P. (2015) β-amyloid disrupts human NREM slow waves and

related hippocampus-dependent memory consolidation. *Nature Neuroscience* 18, 1051–

1057.

17. Marsh, J., Bagol, S. H., Williams, R. S. B., Dickson, G. & Alifragis, P. (2017) Synapsin

I phosphorylation is dysregulated by beta-amyloid oligomers and restored by valproic

acid. *Neurobiol Dis* 106, 63–75.

18. Proctor, D. T., Coulson, E. J., Dodd, P. R. (2010) Reduction in post-synaptic scaffolding

PSD-95 and SAP-102 protein levels in the Alzheimer inferior temporal cortex is

correlated with disease pathology. *J Alzheimers Dis* 21 (3), 795–811.

19. Walsh, D. M., Selkoe, D. J. (2007) Aβ Oligomers – a decade of discovery. *Journal of*

Neurochemistry 101, 1172–1184.

20. Chen, Y. Huang, X., Zhang, Y., Rockenstein, E., Bu, G., Golde, T.E., Golde, E., Xu, H.

(2012) Alzheimer's β-Secretase (BACE1) Regulates the cAMP/PKA/CREB Pathway

Independently of β-Amyloid. *J Neurosci.* 32 (33), 11390–11395.

21. Parihar, M. S., Brewer, G. J. (2010) Amyloid Beta as a Modulator of Synaptic Plasticity.

J Alzheimers Dis 22, 741–763.

22. Campbell, I. G., Guinan, M. J., Horowitz, J. M. (2002) Sleep Deprivation Impairs Long-

Term Potentiation in Rat Hippocampal Slices. *Journal of Neurophysiology* 88 (2), 1073–

1076.

23. Ravassard, P., Pachoud, B., Comte, J.-C., Mejia-Perez, C., Scoté-Blachon, C., Gay, N.,

Claustrat, B., Touret, M., Luppi, P.-H., Salin, P.A. (2009) Paradoxical (REM) Sleep

Deprivation Causes a Large and Rapidly Reversible Decrease in Long-Term Potentiation,

Synaptic Transmission, Glutamate Receptor Protein Levels, and ERK/MAPK Activation

in the Dorsal Hippocampus. *Sleep* 32 (2), 227–240.

24. Bhat, A., Ray, B., Mahalakshmi, A.M., Tuladhar, S., Nandakumar, D., Srinivasan, M.,

Essa, M.M., Chidambaram, S.B., Guillemin, G.J., Sakharkar, M.K. (2020)

Phosphodiesterase-4 enzyme as a therapeutic target in neurological disorders.

Pharmacological Research 105078.

25. Zhang, H., Wang, H., Zhang, F., Li, B., Zhou, Y., Yu, H., Xu, Y., O'Donnell, J. (2019)

Inhibition of phosphodiesterase-4D reverses memory deficits and depression-like effects

via cAMP signaling in mouse models of Alzheimer's disease. *The FASEB Journal* 33,

806.1-806.1.

26. Perez-Gonzalez, R., Pascual, C., Antequera, D., Bolos, M., Redondo, M., Perez, D.I.,

Pérez-Grijalba, V., Krzyzanowska, A., Sarasa, M., Gil, C., Ferrer, I., Martinez, A., Carro,

E. (2013) Phosphodiesterase 7 inhibitor reduced cognitive impairment and pathological

hallmarks in a mouse model of Alzheimer's disease. *Neurobiology of Aging* 34 (9),

2133–2145.

27. Havekes, R., Park, A.J., Tolentino, R.E., Bruinenberg, V.M., Tudor, J.C., Lee, Y.,

Hansen, R.T., Guercio, L.A., Linton, E., Neves-Zaph, S.R., Meerlo, P., Baillie, G.S.,

Houslay, M.D., Abel, T. (2016) Compartmentalized PDE4A5 Signaling Impairs

Hippocampal Synaptic Plasticity and Long-Term Memory. *J Neurosci* 36 (34), 8936–

8946.

28. Cherry, J. A., Davis, R. L. (1999) Cyclic AMP phosphodiesterases are localized in

regions of the mouse brain associated with reinforcement, movement, and affect. *J Comp

Neurol* 407 (2), 287–301.

29. Reyes-Irisarri, E., Pérez-Torres, S., Miró, X., Martínez, E., Puigdomènech, P., Palacios,

J.M., Mengod, G. (2008) Differential distribution of PDE4B splice variant mRNAs in rat

brain and the effects of systemic administration of LPS in their expression. *Synapse* 62

(1), 74–79.

30. Xu, Y., Zhu, N., Xu, W., Ye, H., Liu, K., Wu, F., Zhang, M., Ding, Y., Zhang, C., Zhang,

H., O'Donnell, J., Pan, J. (2018) Inhibition of Phosphodiesterase-4 Reverses Aβ-Induced

Memory Impairment by Regulation of HPA Axis Related cAMP Signaling. *Front. Aging

Neurosci.* 10.

31. Bhat, A., Tan, V., Heng, B., Lovejoy, D.B., Sakharkar, M.K., Essa, M.M., Chidambaram,

S.B., Guillemin, G.J. (2020) Roflumilast, a cAMP-Specific Phosphodiesterase-4

Inhibitor, Reduces Oxidative Stress and Improves Synapse Functions in Human Cortical

Neurons Exposed to the Excitotoxin Quinolinic Acid. *ACS Chem. Neurosci.*

11(24):4405-4415

32. Cui, S.-Y., Yang, M.-X., Zhang, Y.-H., Zheng, V., Zhang, H.-T., Gurney, M.E., Xu, Y.,

O'Donnell, J.M. (2019) Protection from Amyloid β Peptide–Induced Memory,

Biochemical, and Morphological Deficits by a Phosphodiesterase-4D Allosteric Inhibitor.

J Pharmacol Exp Ther 371(2): 250–259

33. Zhang, H.T., O'Donnell, J. M. (2000) Effects of rolipram on scopolamine-induced

impairment of working and reference memory in the radial-arm maze tests in rats.

Psychopharmacology 150(3),311-6**.**

34. Zhang, H.T., Crissman, A. M., Dorairaj, N. R., Chandler, L. J., O'Donnell, J. M. (2000)

Inhibition of Cyclic AMP Phosphodiesterase (PDE4) Reverses Memory Deficits

Associated with NMDA Receptor Antagonism. *Neuropsychopharmacology* 23(2):198-

204

35. Bollen, E., Prickaerts, J. (2012) Phosphodiesterases in neurodegenerative disorders.

IUBMB Life (12):965-70.

36. Rabe, K.F., Bateman, E.D., O'Donnell, D., Witte, S., Bredenbröker, D., Bethke, T.D.

(2005) Roflumilast--an oral anti-inflammatory treatment for chronic obstructive

pulmonary disease: a randomised controlled trial. *Lancet* 366(9485):563-71.

37. Peng, S., Yan, H.-Z., Liu, P.-R., Shi, X.-W., Liu, C.-L., Liu, Q., Zhang, Y. (2018)

Phosphodiesterase 4 Inhibitor Roflumilast Protects Rat Hippocampal Neurons from

Sevoflurane Induced Injury via Modulation of MEK/ERK Signaling Pathway. *Cell.*

Physiol. Biochem. 45(6):2329-2337

38. Vanmierlo, T., Creemers, P., Akkerman, S., van Duinen, M., Sambeth, A., De Vry, J.,

Uz, T., Blokland, A., Prickaerts, J. (2016) The PDE4 inhibitor roflumilast improves

memory in rodents at non-emetic doses. *Behav. Brain Res.* 303, 26–33.

39. Blokland, A., Van Duinen, M.A., Sambeth, A., Heckman, P.R.A., Tsai, M., Lahu, G., Uz,

T., Prickaerts, J. (2019) Acute treatment with the PDE4 inhibitor roflumilast improves

verbal word memory in healthy old individuals: a double-blind placebo-controlled study.

Neurobiology of Aging 77, 37–43.

40. McGirr, A., Lipina, T.V., Mun, H.-S., Georgiou, J., Al-Amri, A.H., Ng, E., Zhai, D.,

Elliott, C., Cameron, R.T., Mullins, J.G., Liu, F., Baillie, G.S., Clapcote, S.J., Roder, J.C.

(2016) Specific Inhibition of Phosphodiesterase-4B Results in Anxiolysis and Facilitates

Memory Acquisition. *Neuropsychopharmacology* 41(4):1080-92.

41. Li, D., Specht, C.G., Waites, C.L., Butler-Munro, C., Leal-Ortiz, S., Foote, J.W., Genoux,

D., Garner, C.C., Montgomery, J.M. (2011) SAP97 directs NMDA receptor spine

targeting and synaptic plasticity. *J Physiol* 589(Pt 18): 4491–4510**.**

42. Kang, J.E., Lim, M.M., Bateman, R.J., Lee, J.J., Smyth, L.P., Cirrito, J.R., Fujiki, N.,

Nishino, S., Holtzman, D.M. (2009) Amyloid-beta dynamics are regulated by orexin and

the sleep-wake cycle. *Science* 326(5955):1005-7

43. Brice, K.N., Hagen, C.W., Peterman, J.L., Figg, J.W., Braden, P.N., Chumley, M.J.,

Boehm, G.W. (2020) Chronic sleep restriction increases soluble hippocampal Aβ-42 and

impairs cognitive performance. *Physiol Behav* 226, 113128.

44. Palop, J. J., Mucke, L. (2010) Amyloid-β Induced Neuronal Dysfunction in Alzheimer's

Disease: From Synapses toward Neural Networks. *Nat Neurosci* 13(7),812-818.

45. Wang, L., Xiaokaiti, Y., Wang, G., Xu, X., Chen, L., Huang, X., Liu, L., Pan, J., Hu, S.,

Chen, Z., Xu, Y. (2017) Inhibition of PDE2 reverses beta amyloid induced memory

impairment through regulation of PKA/PKG-dependent neuro-inflammatory and

apoptotic pathways. *Scientific Reports* **7**, 12044.

46. Tong, L., Thornton, P. L., Balazs, R., Cotman, C. W. (2001) Beta -amyloid-(1-42)

impairs activity-dependent cAMP-response element-binding protein signaling in neurons

at concentrations in which cell survival Is not compromised. *J Biol Chem* 276 (20),

17301–17306.

47. Paes, D., Lardenoije, R., Carollo, R.M., Roubroeks, J.A.Y., Schepers, M., Coleman, P.,

Mastroeni, D., Delvaux, E., Pishva, E., Lunnon, K., Vanmierlo, T., van den Hove, D.,

Prickaerts, J. (2021) Increased isoform-specific phosphodiesterase 4D expression is

associated with pathology and cognitive impairment in Alzheimer's disease. *Neurobiol*

Aging 97, 56–64.

48. Chong, Y. H., Shin, S. A., Lee, H. J., Kang, J. H. L., Suh, Y.H. (2002) Molecular

mechanisms underlying cyclic AMP inhibition of macrophage dependent TNF-alpha

production and neurotoxicity in response to amyloidogenic C-terminal fragment of

Alzheimer's amyloid precursor protein. *J Neuroimmunol* 133(1-2):160-74.

49. Tibbo, A. J., Tejeda, G. S., Baillie, G. S. (2019) Understanding PDE4's function in

Alzheimer's disease; a target for novel therapeutic approaches. *Biochemical Society*

Transactions 47(5):1557-1565.

50. Li, Y.-F., Cheng, Y.-F., Huang, Y., Conti, M., Wilson, S.P., O'Donnell, J.M., Zhang, H.-

T. (2011) Phosphodiesterase-4D knock-out and RNA interference-mediated knock-down

enhance memory and increase hippocampal neurogenesis via increased cAMP signaling.

J Neurosci 31(1): 172–183.

51. Feng, H., Wang, C., He, W., Wu, X., Li, S., Zeng, Z., Wei, M., He, B. (2019) Roflumilast

ameliorates cognitive impairment in APP/PS1 mice via cAMP/CREB/BDNF signaling

and anti-neuroinflammatory effects. *Metab Brain Dis* 34(2):583-591

52. Prickaerts, J., Heckman, P. R. A., Blokland, A. (2017) Investigational phosphodiesterase

inhibitors in phase I and phase II clinical trials for Alzheimer's disease. *Expert Opin*

Investig Drugs 26(9):1033-1048.

53. Ashour, N. H., El-Tanbouly, D. M., El Sayed, N. S., Khattab, M. M. (2021) Roflumilast

ameliorates cognitive deficits in a mouse model of amyloidogenesis and tauopathy:

Involvement of nitric oxide status, Aβ extrusion transporter ABCB1, and reversal by

PKA inhibitor H89. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*

111, 110366.

54. Lacor, P.N., Buniel, M.C., Furlow, P.W., Clemente, A.S., Velasco, P.T., Wood, M.,

Viola, K.L., Klein, W.L. (2007) Aβ Oligomer-Induced Aberrations in Synapse

Composition, Shape, and Density Provide a Molecular Basis for Loss of Connectivity in

Alzheimer's Disease. *J. Neurosci.* 27 (4), 796–807.

55. Chen, F., He, Y., Wang, P., Wei, P., Feng, H., Rao, Y., Shi, J., Tian, J. (2018) Curcumin

can influence synaptic dysfunction in APPswe/PS1dE9 mice. *Journal of Traditional*

Chinese Medical Sciences 5 (2), 168–176.

56. Barbieri, R., Contestabile, A., Ciardo, M.G., Forte, N., Marte, A., Baldelli, P., Benfenati,

F., Onofri, F. (2018) Synapsin I and Synapsin II regulate neurogenesis in the dentate

gyrus of adult mice. *Oncotarget* 9(27), 18760–18774.

57. Corradi, A., Zanardi, A., Giacomini, C., Onofri, F., Valtorta, F., Zoli, M., Benfenati, F.,

(2008) Synapsin-I- and synapsin-II-null mice display an increased age-dependent

cognitive impairment. *J Cell Sci* 121 (18), 3042–3051.

58. Alhaider, I. A., Aleisa, A. M., Tran, T. T., Alzoubi, K. H., Alkadhi, K. A. (2010)

Chronic Caffeine Treatment Prevents Sleep Deprivation-Induced Impairment of

Cognitive Function and Synaptic Plasticity. *Sleep* 33 (4), 437–444.

59. Vecsey, C. G., Huang, T., Abel, T. (2018) Sleep deprivation impairs synaptic tagging in

mouse hippocampal slices. *Neurobiol Learn Mem* 154, 136–140.

60. Wadhwa, M., Sahu, S., Kumari, P., Kauser, H., Ray, K., Panjwani, U. (2015) Caffeine

and modafinil given during 48 h sleep deprivation modulate object recognition memory

and synaptic proteins in the hippocampus of the rat. *Behav Brain Res* 294, 95–101.

61. Bollen, E., Puzzo, D., Rutten, K., Privitera, L., De Vry, J., Vanmierlo, T., Kenis, G.,

Palmeri, A., D'Hooge, R., Balschun, D., Steinbusch, H.M., Blokland, A., Prickaerts, J.

(2014) Improved Long-Term Memory via Enhancing cGMP-PKG Signaling Requires

cAMP-PKA Signaling. *Neuropsychopharmacology* 39 (11), 2497–2505.

62. Wang, C., Yang, X.M., Zhuo, Y.Y., Zhou, H., Lin, H.B., Cheng, Y.F., Xu, J.P., Zhang,

H.T. (2012). The phosphodiesterase-4 inhibitor rolipram reverses Aβ-induced cognitive

impairment and neuroinflammatory and apoptotic responses in rats. *International Journal*

of Neuropsychopharmacology 15 (6), 749–766.

63. Smith, C., Trageser, K.J., Wu, H., Herman, F.J., Iqbal, U.H., Sebastian-Valverde, M.,

Frolinger, T., Zeng, E., Pasinetti, G.M. (2021) Anxiolytic effects of NLRP3

inflammasome inhibition in a model of chronic sleep deprivation. *Transl Psychiatry* 11,

1–15.

64. Xie, G., Huang, X., Li, H., Wang, P., Huang, P. Caffeine-related effects on cognitive

performance: Roles of apoptosis in rat hippocampus following sleep deprivation.

Biochemical and Biophysical Research Communications 534, 632-638

65. Reyes-Irisarri, E., Sánchez, A. J., García-Merino, J. A., Mengod, G. (2007) Selective

Induction of cAMP Phosphodiesterase PDE4B2 Expression in Experimental

Autoimmune Encephalomyelitis. *Journal of Neuropathology & Experimental Neurology*

66 (10), 923–931.

66. Wang, H., Zhang, F., Xu, Y., Fu, H., Wang, X., Wang, L., Chen, W., Xu, X., Gao, Y.,

Zhang, J., Zhang, H.T. (2020) The Phosphodiesterase-4 Inhibitor Roflumilast, a Potential

Treatment for the Comorbidity of Memory Loss and Depression in Alzheimer's Disease:

A Preclinical Study in APP/PS1 Transgenic Mice. *International Journal of*

Neuropsychopharmacology 23 (10), 700–711.

67. Jabaris, S.S.L., Sumathy, H., Girish, R., Narayanan, S., Sugumar, M., Saravana Babu, C.,

Thanikachalam, S., Thanikachalam, M. (2015) Phosphodiesterase-4 inhibitors

ameliorates cognitive deficits in deoxycorticosterone acetate induced hypertensive rats

via cAMP/CREB signaling system. *Brain Research* 1622, 279–291.

68. Patti, C.L., Zanin, K.A., Sanday, L., Kameda, S.R., Fernandes-Santos, L., Fernandes,

H.A., Andersen, M.L., Tufik, S., Frussa-Filho, R. (2010) Effects of Sleep Deprivation on

Memory in Mice: Role of State-Dependent Learning. *Sleep* 33 (12), 1669–1679.

69. Grahnstedt, S., Ursin, R. (1985) Platform sleep deprivation affects deep slow wave sleep

in addition to REM sleep. *Behav. Brain Res.* 18 (3), 233–239.

70. Prickaerts, J., van Staveren, W.C.G., Şik, A., Markerink-van Ittersum, M., Niewöhner,

U., van der Staay, F.J., Blokland, A., de Vente, J. (2002) Effects of two selective

phosphodiesterase type 5 inhibitors, sildenafil and vardenafil, on object recognition

memory and hippocampal cyclic GMP levels in the rat. *Neuroscience* 113 (2), 351–361.

Figure 1: Schematic outline of the experimental design

Figure 2: Roflumilast downregulated the protein expression of PDE4B and Aβ in the hippocampus of SD mice. (**A**) Semi-quantification of PDE4B/β-actin expression (**B**) Semi-quantification of Aβ /β-actin expression. Data are presented as the mean ± SEM, [##]denotes $p<0.01$ versus vehicle-treated NSD group, [**]denotes $p<0.01$ versus vehicle-treated SD group.

Figure 3: Effect of ROF on Aβ plaques in CA1 and DG regions of the hippocampus in sleep-deprived mice. Arrows denote amyloid-β plaques.

Figure 4: Roflumilast administration restored SD induced decrease in cAMP level in mice.
Data are presented as the mean ± SEM. [##] denotes $p<0.01$ versus the vehicle-treated NSD group, [*] and [***] denote $p<0.05$ and $p<0.001$, respectively, versus the vehicle-treated SD group.

Figure 5: Roflumilast administration increased neurotrophic factors, CREB and BDNF, in sleep-deprived mice brain. (A) Semi-quantification of pCREB/TCREB expression **(B)** Semi-quantification of BDNF /β-actin expression. Data are presented as the mean ± SEM, [##] denotes p < 0.01, [###] denotes p < 0.001 versus vehicle-treated NSD group, [**] and [***] denotes $p<0.01$, and $p<0.001$, respectively, versus vehicle-treated SD group.

Figure 6: Roflumilast upregulated the expression of synaptic proteins in SD mice brain. (A) Semi-quantification of Synapsin-I/β-actin expression. **(B)** Semi-quantification of SAP-97/β-actin expression. **(C)** Semi=quantification of PSD-95/β-actin expression. Data are presented as the mean ± SEM. # denotes $p<0.05$, ## denotes $p<0.01$ versus vehicle-treated NSD group, ** and *** denotes $p<0.01$ and $p<0.001$, respectively, versus vehicle-treated SD group, n.s. denotes non-significant.

Figure 7: Roflumilast improved recognition memory in NORT in sleep-deprived mice. (A) time exploring the familiar and novel object (B) Discrimination index. Data are presented as the mean ± SEM. [#] and [##] denotes $p < 0.05$, and $p < 0.01$ respectively, versus the NSD group and [*] and [***] denote $p < 0.05$, and $p < 0.01$, respectively, versus the SD group

Figure 8: Cresyl violet staining of CA1 and DG regions of the hippocampus of SD mice. Arrows denote pyknotic nuclei.