Theoretical Analysis of Principal Components in an Umbrella Model of Intraspecific Evolution

Maxime Estavoyer[1]   Olivier François[1,2]

Authors' affiliations:

[1]Université Grenoble-Alpes, Centre National de la Recherche Scientifique, Grenoble INP, TIMC UMR 5525, 38000 Grenoble, France.
[2] Inria Grenoble - Rhône-Alpes Inovallée 655 Avenue de l'Europe - CS 90051 38334 Montbonnot, France.

Corresponding author:

olivier.francois@univ-grenoble-alpes.fr

Keywords: Colonization Models, Genetic Drift, Principal Component Analysis, Eigenvalues, Coalescent Theory.

Abstract

Principal component analysis (PCA) is one of the most frequently-used approach to describe population structure from multilocus genotype data. Regarding geographic range expansions of modern humans, interpretations of PCA have, however, been questioned, as there is uncertainty about the wave-like patterns that have been observed in principal components. It has indeed been argued that wave-like patterns are mathematical artifacts that arise generally when PCA is applied to data in which genetic differentiation increases with geographic distance. Here, we present an alternative theory for the observation of wave-like patterns in PCA. We study a coalescent model – the umbrella model – for the diffusion of genetic variants. The model is based on a hierarchy of splits from an ancestral population without any particular geographical structure. In the umbrella model, splits occur almost continuously in time, giving birth to small daughter populations at a regular pace. Our results provide detailed mathematical descriptions of eigenvalues and eigenvectors for the PCA of sampled genomic sequences under the model. Removing variants uniquely represented in the sample, the PCA eigenvectors are defined as cosine functions of increasing periodicity, reproducing wave-like patterns observed in equilibrium isolation-by-distance models. Including rare variants in the analysis, the eigenvectors corresponding to the largest eigenvalues exhibit complex wave shapes. The accuracy of our predictions is further investigated with coalescent simulations. Our analysis supports the hypothesis that highly structured wave-like patterns could arise from genetic drift only, and may not always be artificial outcomes of spatially structured data. Genomic data related to the peopling of the Americas are reanalyzed in the light of our new theory.

Introduction

Investigating population structure from large genotypic data set is one of the most important steps of modern population genetic data analysis. This step is often carried out with a method called principal component analysis (PCA). PCA is a nonparametric approach that summarizes the main axes of genetic variation by computing the largest eigenvalues and the first eigenvectors of the sample covariance matrix [21, 24]. The use of PCA dates back to the early days of human population genetics [5, 33], and it has been shown theoretically that it can reveal information about admixture and population differentiation among samples [35, 32, 36, 16].

Because PCA is an exploratory method that makes no assumptions about population history, interpretation of results in terms of demographic events is sometimes difficult, in particular when the details of a range expansion are scrutinized [34, 32, 14, 1, 3]. More precisely, Novembre and Stephens reported that highly structured wave-like patterns that are seen in PCA when genetic data are simulated under range expansion models, are also observed under equilibrium isolation-by-distance models without range expansions [34]. Their study warned that the wave-like patterns could be mathematical artifacts that arise when PCA is applied to spatial data in which covariance between locations decays with geographic distance [9, 17]. Sinusoidal shapes similar to those observed with geographic samples are also observed with temporal samples of ancient DNA genotypes [15], and the combination of time and spatial heterogeneity in sampling further complicates the interpretation of patterns observed in PC plots [10]. Those difficulties have questioned the usefulness of PCA for species that evolved under geographic models of range expansion for which wave-like patterns were first observed [33].

Range expansions are, however, common in the evolution of species. For example, they include the worldwide expansion of monarch butterflies [38], or the invasion and the evolution of speed in toads [37]. The models have been extensively studied

theoretically and with computer simulations [2, 20, 11]. Humans are one of the most studied example of range expansion, exhibiting a positive correlation between genetic and geographic distance at the worldwide scale [40], and a proposed model for that pattern involved serial founder effects [8, 6]. In the serial founder effect model, populations migrate outward from a geographic origin through a process where each of a series of populations is formed from a subset of the previous population in the expansion [6]. Evolution under the model corresponds to a population tree having maximal imbalance (Figure 1), and several patterns of human genetic variation were explained by serial colonization out of Africa [8, 6, 7, 23].

In this study, we analyze a simple model in which small populations diverge from an ancestral population at a constant pace, through a process similar to a serial colonization model (Figure 1). The model is called the *umbrella model* in reference to a tree known as the umbrella pine, parasol pine or Italian stone pine, in which the length of a branch decreases with the distance to the root. Unlike the serial founder effect model, the samples are not necessarily localized in geographic space, and patterns of genetic variation result from temporal drift only [41]. The model describes the diffusion of genetic variants based on hierarchy of fissions from a genetically homogeneous population without any geographical structure in the sample. The samples sort according to their date of coalescence within the trunk of the tree, representing the ancestral population lineage. Removing variants that occur exactly once in the sample, the model also describes the neutral evolution of a genetically homogeneous population since the most recent common ancestor of the sample. In this case, the present-day sequences are not different of ancient DNA sequences sampled at their date of coalescence with the ancestral lineage. Thus the model is not only relevant to present-day population structure analysis, but also highlights some interesting properties of population structure analysis for ancient DNA samples.

We provide a detailed mathematical description of population structure under the

umbrella model by studying the PCA eigenvalues and eigenvectors of sampled genomic sequences. The eigenvalues and eigenvectors of the PCA will be obtained after approximating the finite-size empirical covariance matrix by an integral (Hilbertian) operator. Removing variants that are uniquely represented in the sample, the eigenvectors are defined as cosine functions of increasing periodicity, reproducing wave-like patterns observed in equilibrium isolation-by-distance models. Keeping those rare variants produces complex oscillatory eigenvectors, for which mathematical descriptions are obtained for the largest eigenvalues. The accuracy of approximation will be further investigated with coalescent simulations, and genomic data related to the settlement of the Americas will be reanalyzed in the light of the new results [31].

1 Model description and theoretical background

1.1 The umbrella model

We study the genealogical relationships of n genomic sequences obtained from haploid individuals under a serial colonization model of evolution, the umbrella model (UM).

Skeleton tree. The UM defines a skeleton tree for the coalescent genealogies of the n samples. The skeleton can be viewed as an approximation of an ancestral population that splits continuously in time. The model considers n populations of equal size N, which must be thought of as being small compared to the effective population size N_e (N is of order N_e/n). Time is measured in units of population size. Temporal evolution is defined by population splits that occur each unit of time. Examples values related to human data are around $N_e = 10,000$, $N = 100$ and $n = 100$. A splitting event gives birth to an external branch that diverges from the trunk of the tree (Figure 1A). The geographic localization of the daughter population can be arbitrary, and no relationship between geographic and genetic distance is modelled a priori. The skeleton tree of the UM has n external branches,

Figure 1. Umbrella Model. (**A**) Skeleton tree for the umbrella model: The individuals at the leaves are labelled according to the time of divergence with the ancestral lineage. Individual lineages coalesce with the ancestral lineage at their time of divergence plus one unit. (**B**) Equivalent description of the UM as a serial colonization model, in which individuals at the leaves are labelled according to their position in a linearly arranged habitat. The initial population is given the highest label. The value τ_k indicates the splitting time for individual lineage $k + 1$ (τ_1 is for lineages 1 and 2).

for which the total length is equal to $1 + n(n-1)/2$, and $n-1$ internal branches of
unit length, forming a trunk of length n. The total tree length is equal to $n(n+1)/2$.

Sample genealogy. In the UM, a single individual is sampled from each popula-
tion, so that the sample size, n, is equal to the number of populations. The average
time since the most recent common ancestor of all individuals is equal to n. Individ-
ual lineages coalesce with the ancestral lineage at their time of divergence plus one
unit. Thus the samples sort according to their date of coalescence within the trunk
of the tree. More precisely, individual lineage i ($i \geq 2$) coalesces with the ancestral
lineage at time i. The first lineage coalesces with the ancestral lineage at time two,
just like the second individual lineage. The average length of the sample genealogy
is equal to $n(n+3)/2 - 1$, which is very close to the length of the skeleton tree.

The trunk of the tree represents the evolution of a genetically unstructured ances-
tral population, and we can see that ancient DNA is not needed in order to describe
evolution under the UM, as ancient variation reflects in the present-day genetic data
directly. Ordering samples according to geographic distance from a source population,
when the source is identified, shows that the UM can be equivalently represented as a
comb tree (Figure 1B), similar to the skeleton tree of coalescent-based serial founder
effect models [8, 6, 7]. However, the umbrella model does not suggest any particular
direction of colonization, and the samples may be randomly scattered in geographic
space at arbitrary distance from the geographic origin of their ancestors in the trunk.
A command for the `ms` program simulating coalescent genealogies under the UM is
provided in Appendix A [22].

Allele frequency spectrum. Before describing population genetic structure from
observed genotypes, we provide some elementary properties of UM genealogies. For
a sample of size n, the allele frequency spectrum, ξ, defined as the frequency of
mutations in the sample, can be computed as follows. The probability of a singleton,

defined as a mutation being represented uniquely in the sample, is equal the ratio of external branch length to the total length of the genealogy

$$\text{Prob}(\xi = 1/n) \approx 1 - \frac{2}{n+3}\,.$$

Conditional on non-unique mutations, the frequency of mutation has a uniform distribution

$$\text{Prob}(\xi = k/n \,|\, \xi > 1/n) = \frac{1}{n-2}\,, \quad k = 2,\ldots,n-1\,,$$

and the unconditional distribution can be described as follows

$$\text{Prob}(\xi = k/n) \approx \frac{2}{(n-2)(n+3)}\,, \quad k = 2,\ldots,n-1.$$

Conditioning on non-unique mutations is equivalent to sampling ancient variation at ancestor nodes in the trunk of the skeleton tree, and the uniform distribution corresponds to the allele frequency spectrum for the ancient samples.

Diploid individuals. We limit our discussion to samples of haploid individuals. Diploid individuals could be included in the model by doubling the sample size, considering two individuals from each of the n populations. For large n, the two lineages in a diploid individual coalesce much before the junction with the trunk of the skeleton tree. For population k, the probability of observing an heterozygote locus can be approximated as $4/n(n+4)$, and most genotypes will be homozygote, 0 or 2. The probability of heterogyzosity does not depend on k, reflecting that the UM does not predict any decline of heterogyzosity with respect to a particular origin. In contrast the number of singletons (for haploids) and doubletons (for diploids) can be predicted to increase linearly with the age of populations or with the divergence time from the ancestral lineage.

1.2 Eigenvalues and eigenvectors of the sample covariance matrix

Principal component analysis for large sample sizes. For a sample of size n, PCA analyzes population genetic structure by computing the eigenvalues and eigenvectors of the empirical covariance matrix [24]. The eigenvectors are also called principal components (PCs). The first PCs represent the axes of genetic variation which carry the most information, and the eigenvalues represent the variances of samples along the axes. The eigenvalues (denoted λ) and their associated eigenvectors (denoted $\mathbf{u}$) solve a system of linear equations

$$\sum_{i=1}^{n} c(i,j)\mathbf{u}(j) = \lambda\mathbf{u}(i), \quad i = 1,\ldots,n, \tag{1}$$

where the $c(i,j)$'s are the coefficients of the sample covariance matrix. To find approximated solutions, functional forms of eigenvectors will be considered

$$\mathbf{u}(i) = u(i/n), \quad i = 1,\ldots,n.$$

The function $u(x)$ will be defined on the interval $(0,1)$ as an eigenfunction of an integral operator in an Hilbert space, $L^2(0,1)$. The integral operator will be obtained by considering a continuous approximation of the system of discrete linear equations (1). In the large n limit, the continuous approximation will assume that an ancestral lineage could split continuously across time, leading to small daughter populations that evolve independently of the ancestral population. The approximation will be obtained after scaling both sides of equation (1) by a factor n^2. For the models considered in our study, the solutions of the finite system of equations

$$\frac{1}{n}\sum_{i=1}^{n} c(i/n,j/n)u(j/n) = \frac{\lambda}{n^2}u(i/n), \quad i = 1,\ldots,n, \tag{2}$$

can be approximated by the continuous solutions of an integral equation

$$n^2 \int_0^1 C(x,y)u(y)dy = \lambda u(x).$$ (3)

In this equation, $C(x,y)$ needs to be determined. This will be achieved by considering a coalescent approximation of the empirical covariance matrix, as described below.

Coalescent approximation. Assuming a neutral coalescent model for the genealogy of a sample of size n, a theoretical approximation for the mathematical expectation of the empirical covariance matrix was obtained by McVean [32] as follows

$$\bar{T}c(i,j) = \bar{t}(i) + \bar{t}(j) - \bar{t} - \bar{t}(i,j), \quad i,j = 1,\dots,n,$$ (4)

where $\bar{t}(i,j)$ is the average coalescence time of lineage i and lineage j, $\bar{t}(i)$ is the average coalescence time of a random lineage with i, $\bar{t}(i) = \sum_{k=1}^n \bar{t}(i,k)/n$, $\bar{t}$ is the average coalescence time of two random lineages, $\bar{t} = \sum_{k=1}^n \bar{t}(k)/n$, and $\bar{T}$ is the average length of the coalescent tree. For the UM, the average length of the coalescent tree is approximately equal to the length of the skeleton tree. This length is equal to $\bar{T} = n$ when the singletons are removed from the data matrix, and to $\bar{T} \approx n(n+3)/2$ otherwise. According to the central limit theorem, equation (4) becomes a valid approximation of the sample covariance matrix when a large number of genomic loci are included in the data matrix. Simulations will be used to check the accuracy of this assumption.

2   Main results

The main results of our study are mathematical descriptions of the PCA eigenvalues and eigenvectors for n multilocus genotypes sampled from the umbrella model. The first components exhibit wave-like patterns as observed in Cavalli-Sforza et al.'s paper

[33]. While Novembre and Stephens showed that sinusoidal patterns arise in principal components when PCA is applied to spatial data [34], our results support the evidence that those patterns may not always be a consequence of spatially structured data. Here, waves result from the hierarchy of population splits from a genetically homogeneous ancestral population without any geographical structure in the sample.

2.1   Spectral analysis of the filtered data set

This section describes the eigenvalues and eigenvectors of a PCA for a sample of n multilocus haploid genotypes after the data matrix has been filtered out for singletons (variants that are uniquely represented in the sample). Removing those rare variants is commonly done in empirical analyses of population genetic structure, often by considering a threshold for the minor allele frequency. Here we condition on non-unique polymorphisms for two reasons. The first reason is aesthetic: the eigenvalues and the principal components have simple exact expressions in the large n limit, and they provide direct mathematical evidence for sinusoidal patterns in the PCA. The second reason is a connection to ancient DNA. Removing singletons from present-day samples in the UM amounts to studying samples from their ancestors in the internal branch of the tree. Under the UM, filtering singletons provides as much genetic information on ancestors as ancient DNA would do, without the shortcomings of the technology.

For non-singleton variants, the average length of the conditioned genealogy is equal to n. The covariance matrix is determined by the coalescence times within the ancestral population. More specifically, the average coalescence time for ancestor i and ancestor j can be described as

$$\bar{t}(i,j) = \max(i,j), \quad 1 \leq i, j \leq n.$$

The average coalescence time of ancestor i with a random ancestor is equal to

Figure 2. Approximation of PC eigenvalues and eigenvectors for filtered data from the UM. Simulation of the umbrella model performed for $n = 100$ haploid individuals and $L = 5,938$ single nucleotide polymorphism loci obtained after removing singletons. *Top panels*: Approximations of PC eigenvalues by $L\lambda_k/n = L/\pi^2 k^2$ (left) and by the numerical values obtained from the discrete system of linear equations (right). Eigenvalues are displayed on a base 10 logarithm scale. *Bottom panels*: Approximations of PC eigenvectors by their continuous approximations, $\mathbf{u}_k(x) = \sqrt{2}\cos(k\pi x)$, for $x = i/n$, $k = 1$ to 4. Eigenvectors are represented in red (most recent samples) to blue (most ancient samples) colors. Continuous approximations are represented by grey circles.

$$\bar{t}(i) = \frac{1}{n} \sum_{k=1}^{n} \max(i, k) \,,$$

and the average coalescence time of two random ancestral lineages is equal to

$$\bar{t} = \frac{1}{n^2} \sum_{k=1}^{n} \sum_{\ell=1}^{n} \max(k, \ell) \,.$$

We are now positioned to state our main result, holding for the PCA of a filtered data set. *Let n be a large integer and $k \in \{1, \ldots, n\}$. For n samples from the UM, the kth largest eigenvalue of the empirical covariance matrix can be approximated as*

$$\lambda_k \approx \frac{n}{k^2 \pi^2} \,.$$

The kth principal component can be approximated as

$$\mathbf{u}_k(i) \approx u_k(i/n) = \sqrt{2} \cos(k\pi i/n) \,, \quad i = 1, \ldots, n.$$

The proof of this result is detailed in Appendix B. It follows from differential calculus arguments based on the approximation of the discrete spectral problem by integral equations. For $x \in (0, 1)$, the continuous eigenvalue problem corresponding to equation (3) is

$$\frac{(3x^2 + 2)}{6} \int_0^1 u(y)dy + \frac{1}{2} \int_0^1 y^2 u(y)dy - \int_0^1 \max(x, y)u(y)dy = \frac{\lambda}{n} u(x) \,, \quad (5)$$

where λ and $u(x)$ are the unknowns. This is equivalent to solving differential equations of the following type

$$\frac{\lambda}{n} u''(x) + u(x) = \int_0^1 u(y)dy \,, \quad (6)$$

with boundary conditions $u'(0) = 0$, $u'(1) = 0$, and

$$\int_0^1 u(y)dy = 0\,.$$

The solutions are provided by the cosine functions and by the values of λ given above (Appendix B). The result remains valid when PCA is extended to data matrices filtered out for doubletons, tripletons or any finite number of classes in the allele frequency spectrum. Minor allele frequency filters, such as considering variants of frequency greater than 5%, could also be analyzed with integral operators. This would require modifying the bounds of the integral, and would lead to less elegant solutions than in the case of singletons.

2.2 Spectral analysis of the unfiltered data set

This section describes the eigenvalues and principal components for a sample of n multilocus haploid genotypes under the umbrella model. All variants are now represented in the sample. In the large n limit, the eigenvalues and the principal components have more complex expressions than with filtered data, but their expression still provide mathematical evidence for wave-like patterns in PCs. For present-day samples, the average coalescence time of lineage i and lineage j can be described as

$$\bar{t}(i,j) = \begin{cases} \max(i,j)\,, \text{ if } i \neq j\,, \\ 0\,, \text{ otherwise.} \end{cases}$$

The difference with the case of filtered data is that instantaneous self-coalescence leads to $\bar{t}(i,i) = 0$ for every individual i, a change that complicates the mathematical analysis of the eigenvalue problem for the UM. The average coalescence time of lineage i and a random lineage can be described as

$$\bar{t}(i) = \frac{1}{n}\sum_{k=1}^{n}\max(i,k) - \frac{i}{n}\,,$$

and the average coalescence time of two random lineages is equal to

Figure 3. Wave patterns in PC plots for filtered data. Simulation of the umbrella model performed for $n = 100$ haploid individuals and $L = 5,938$ single nucleotide polymorphism loci obtained after removing singletons. *Left column*: PC plots for the simulated data. *Right column*: Continuous plots derived from the cosine approximations for $x = i/n$ and $k = 1$ to 4. Eigenvectors are represented in red (most recent samples) to blue (most ancient samples) colors.

$$\bar{t} = \frac{1}{n^2} \sum_{k=1}^{n} \sum_{\ell=1}^{n} \max(k, \ell) - \frac{1}{n^2} \sum_{k=1}^{n} k \, .$$

Let $J_\alpha(x)$ and $Y_\alpha(x)$ denote the Bessel functions of the first and second kind for $\alpha = 1, 2$ (and for $\alpha = 0$ in a later statement). We formulate our main result for the unfiltered data set below.

Let n be a large integer value, $k \in \{1, \dots, n\}$, and let λ_k be the kth eigenvalue of the empirical covariance matrix for n samples from the UM. Assume that $\lambda_k > 2/n$, then λ_k can be approximated by the kth largest zero of the function

$$\lambda \mapsto J_2(\sqrt{2\lambda}n)Y_2(\sqrt{2(\lambda n - 2)n}) - Y_2(\sqrt{2\lambda}n)J_2(\sqrt{2(\lambda n - 2)n}) \, .$$

The kth eigenvector of the empirical covariance matrix, $\mathbf{u}_k$, can be approximated as

$$\mathbf{u}_k(i) \approx A_k\sqrt{2}\,\frac{J_1(\sqrt{2n}\sqrt{\lambda_k n - 2i/n})}{\sqrt{\lambda_k n - 2i/n}} + B_k\sqrt{2}\,\frac{Y_1(\sqrt{2n}\sqrt{\lambda_k n - 2i/n})}{\sqrt{\lambda_k n - 2i/n}} + C_k \, , \quad (7)$$

where C_k is defined as

$$C_k = -A_k\sqrt{2}\,\frac{1}{(\lambda_k n)^{3/2}}\left(J_1(\sqrt{2\lambda_k}n) - \frac{J_2(\sqrt{2\lambda_k}n)}{Y_2(\sqrt{2\lambda_k}n)}Y_1(\sqrt{2\lambda_k}n)\right) \, , \quad (8)$$

where B_k is defined as

$$B_k = -A_k\frac{J_2(\sqrt{2\lambda_k}n)}{Y_2(\sqrt{2\lambda_k}n)} \, , \quad (9)$$

and where A_k is a normalizing constant for a unitary eigenfunction (see Appendix C).

For $n = 30$, the approximation holds for the first three eigenvalues and eigenvectors according to the condition $\lambda_k > 2/n$. For $n = 100$, it describes five eigenvalues and eigenvectors, and this number grows to eight eigenvalues and eigenvectors for $n = 200$. For large n, the condition can be verified for about $\sqrt{n/\pi^2}$ eigenvalues and

Figure 4. Approximation of PC eigenvalues and eigenvectors for unfiltered data. Simulation of the umbrella model performed for $n = 100$ haploid individuals and $L = 300,000$ single nucleotide polymorphism loci. *Top panels*: Approximations of the five first PC eigenvalues by $L\lambda_k/n$, where λ_k is obtained from the continuous approximation (left), and by the numerical values obtained from the discrete eigenvalue problem using McVean's equations (right). Eigenvalues are displayed on a base 10 logarithm scale. *Bottom panels*: Approximations of PC eigenvectors by their continuous Bessel approximations, $\mathbf{u}_k(x)$, for $x = i/n$, $k = 1$ to 4. Eigenvectors are represented in red (most recent samples) to blue (most ancient samples) colors. Continuous approximations are represented by grey circles.

eigenvectors. As for the case of filtered data, the results were obtained from classical differential calculus arguments. Those arguments are based on the approximation of the discrete spectral problem by an integral equation. For $x \in (0,1)$, the integral equation is now defined as follows. For $x \in (0,1)$, we have

$$\frac{(3x^2 + 2)}{6} \int_0^1 u(y)dy + \frac{1}{2} \int_0^1 y^2 u(y)dy - \int_0^1 \max(x,y)u(y)dy = \frac{\lambda}{2}u(x) - \frac{x}{n}u(x). \quad (10)$$

The difference with the analysis of the filtered data set is with the right hand side, which now includes the term equal to $-xu(x)/n$. A few other terms were considered as being null for large n. (Those simplifications were evaluated in the simulation study). The solutions arise from a differential equation of the following type

$$\left(\frac{\lambda}{2} - \frac{x}{n}\right) u''(x) - \frac{2}{n}u'(x) + u(x) = \int_0^1 u(y)dy, \quad (11)$$

which is similar to equation (6) for large sample sizes.

The proof of the result is detailed in Supplementary Text. Although less obvious than for the filtered data, the approximate eigenvectors still provides evidence of wave-like patterns arising in PC plots. As the simulation study will show, those eigenvectors capture wave patterns more complex than those obtained with the cosine functions.

2.3 Connection to forward-in-time models

The principal component analysis of samples obtained from the backward-in-time coalescent genealogies of the umbrella model shares similarities with the forward-in-time phylogenetic analysis of allele frequencies observed at the leaves of the skeleton tree. More precisely, Cavalli-Sforza and Edwards modelled the evolution of allele frequencies as a branching Brownian motion [4]. In their study, the branching Brownian motion was considered as a simplification of more complex diffusion models proposed

in neutral population genetics [25, 26]. In the forward-in-time umbrella model, the covariance matrix of allele frequencies observed at the leaves of a tree has a relatively simple expression. The sampled individuals, $i = 1, \ldots, n$, are now numbered from the most ancient one to the most recent one, in a direction inverse of that considered in the backward-in-time analysis. A node $i = 0$ corresponds to the root of the tree, and it is not included in the sample. Thus the sample size remains equal to n. For the sample of size n, the empirical covariance matrix is given by

$$\bar{c}(i, j) = \begin{cases} \min(i, j), & \text{if } i \neq j, \\ n, & \text{otherwise.} \end{cases} \tag{12}$$

Assuming a continuous approximation for large sample size, the eigenvalue problem corresponds to the following integral operator

$$\int_0^1 \min(x, y) u(y) dy + \frac{1}{n} u(x) = \frac{\lambda}{n^2} u(x) + \frac{x}{n} u(x), \quad x \in (0, 1). \tag{13}$$

The resolution of this equation is similar to the coalescent analysis of the UM for the unfiltered data. More precisely, we obtained the following results. The eigenvalues such that $\lambda_k > n$ are solutions of the equation

$$J_1(2\sqrt{\lambda - n}) Y_0(2\sqrt{\lambda}) = Y_1(2\sqrt{\lambda - n}) J_0(2\sqrt{\lambda})$$

and their corresponding eigenfunctions can be approximated by

$$u_k(x) \approx A_k \sqrt{n} \frac{J_1 \left(2\sqrt{\lambda_k + xn - n}\right)}{\sqrt{\lambda_k + xn - n}} + A_k B_k \sqrt{n} \frac{Y_1 \left(2\sqrt{\lambda_k + xn - n}\right)}{\sqrt{\lambda_k + xn - n}}, \tag{14}$$

where B_k is defined as

$$B_k = \frac{J_0 \left(2\sqrt{\lambda_k}\right)}{Y_0 \left(2\sqrt{\lambda_k}\right)},$$

and A_k is defined such that

$$\int_0^1 u_k^2(x)dx = 1\,.$$

293   Removing the $1/n$ term in equation (13) does not change the approximation
294 results. Removing both $1/n$ and x/n amounts to solve the eigenvalue problem for
295 the Wiener process

$$\int_0^1 \min(x,y)u(y)dy = \frac{\lambda}{n^2}u(x)\,, \quad x \in (0,1), \tag{15}$$

which is very similar to the case of unfiltered data for the UM. According to the
Karhunen-Loeve theorem [29], the continuous eigenvectors are given by $u_k(x) =$
$\sqrt{2}\sin((k-1/2)\pi x)$, and their eigenvalues are given by $\lambda_k = n^2/\pi^2((k-1/2)^2)$, for
$k = 1, \ldots, n$.

In the forward-in-time approach, the eigenvectors exhibit shapes that are very
similar to those obtained under the coalescent approach. The difference with the
backward-in-time approach comes from the interpretation of the root. In the forward-
in-time approach, the root may not correspond to the most recent common ancestor of
all lineages. Instead it may correspond to the latest common ancestor of the sample.
For application to data analysis, the forward-in-time approach would thus require
considering a strictly positive age for the first lineage, and the above integral equation
would include a strictly positive lower bound (see [15]). A last comment is that the
forward-in-time theory is not restricted to modelling the drift of allele frequencies.
The results predict that wave-like patterns also arise in PC plots obtained from
multiple phenotypic values when phenotypes evolve by neutral mutation [30, 12].

3   Simulations and data analysis

312 In order to check the accuracy of continuous approximations of PCA eigenvalues and
313 eigenvectors, we performed coalescent simulations of haploid genotypes under the

umbrella model. In this section, the results were reported for $n = 100$ populations and samples. Results for $n = 30$ populations (and samples) could be found in the supplementary materials. The number of populations ($n = 30, 100$) was chosen to be of the same order of magnitude as in our analysis of human genotypes from the Simons Genome Diversity Project [31]. Simulations of the UM were performed with the program `ms`. PCA was performed in the `R` programming language with the command `prcomp`. First, we compared the eigenvalues and eigenvectors of a PCA for genotypes filtered out for singletons. We recall that an interesting property of the model is that removing singletons amounts to observing ancestral variants, corresponding to the evolution of a single ancestral population sampled at regular time points. Then we compared the eigenvalues and eigenvectors of the PCA for the unfiltered data set, in which the number of singletons in a particular individual sample reflects the age of its ancestor in the internal branch of the skeleton tree.

Filtered data set. PCA eigenvalues, corresponding to the variance of projected samples along each PC (σ_k^2), were compared to their approximation (λ_k) obtained from equation (6). According to our theoretical result, σ_k^2 should be approximately equal to $L/\pi^2 k^2$, where L is the number of loci in the genotype matrix. We found that the approximations of PC eigenvalues were highly accurate, and they were also close to numerical eigenvalues obtained by solving the discrete system of linear equations (3) based on McVean's covariance matrix (Figure 2, top panels). The results provided evidence that the empirical covariance matrix was well-approximated by its expected value – a result that holds by the law of large numbers –, and that the integral approximation was also accurate. Next we checked that PCs exhibited wave-like patterns as predicted by the infinite-size theory (Figure 2). For the four first PCs, we found that the simulated PCs were hardly divergent from their continuous approximations by cosine functions, $\mathbf{u}_k(i) = \sqrt{2}\cos(k\pi i/n)$, for $i = 1, \ldots, n$. The PC plots, showing PC_{k+1} as a function of PC_k, also displayed wave-like patterns (Figure 3).

The PC1-PC2 plot exhibited a symmetric arch, whereas the PC2-PC3 plot exhibited a fish-like pattern. Higher PC plots exhibited patterns similar to Lissajous curves. The results for $n = 30$ were similar to those obtained with $n = 100$, and showed that the number of populations does not need to be very large for the asymptotic theory to apply (Figure S1 and Figure S2). The PC plots also looked like those obtained in simulations of one-dimensional stepping-stone models [34]. The similarity was not expected a priori, because the covariance matrix of the stepping-stone model, which has a stationary block-Toeplitz structure, differs from the non-stationary structure of the umbrella model. Whatever the value of n, the remarkable fact is that the sinusoidal patterns arise without any geographic structuring, but because a genetically homogeneous founder population gives birth to many small drifting subpopulations through its temporal evolution.

Unfiltered data set. For unfiltered data from the UM, the contribution of singletons to genetic variation modified the shapes of PCs, and thereby, the patterns of population structure changed significantly (Figure 4, Figures S3-S4). For the first five PCs, we found that the approximations of PC eigenvalues were highly accurate, and again very close to numerical eigenvalues obtained from solving the discrete system of linear equations (Figure 4 top left panels, and Figure S5 for $n = 30$). The PCs exhibited wave-like patterns as predicted by the Bessel functions obtained from theoretical approximation by the integral operator (Figure 4, Figure S3). The amplitude of oscillation of these functions varied with the age of the individual ancestor, corresponding to the PC ranking. The PC plots exhibited shapes that were not observed in simulations of one-dimensional stepping-stone models. The PC1-PC2 plot still exhibited an arch pattern, but the arch had an asymmetrical shape, with a more pronounced asymmetry for smaller n (Figure S3 and Figure S5). The first PCs remained close to those obtained for the filtered data set, showing that the first axes of variation are mainly influenced by genetic drift in the ancestral population.

However, the subsequent PCs exhibited patterns that were unseen in the PCA of the filtered data set. PCs associated with smaller variances were flat until the age of the ancestor, exhibited a large amplitude oscillation, and then returned back to zero quickly (Figure S4). Those shapes made PC plots difficult to display graphically, and only a few first PCs could be usefully interpreted, especially in the smaller n case.

Human data analysis. To evaluate the relevance of the umbrella model to biological data, we analyzed human genomic data for fifty-nine individuals originating from $n = 30$ human populations in Asia and in the Americas [31]. The individual identifiers and the geographic coordinates of the sampled populations were given in Table S1, and displayed in Figure 5. The study of those populations was motivated by the settlement of the Americas, for which the prevalent models outline Asian migration from the Beringia land bridge and subsequent dispersal of the ancestral population throughout the Americas [19, 42]. A subset of around 555k SNPs was drawn from the whole genome sequence data, after filtering out missing data and variants with minor allele frequency less than 5%. To keep the dimension of the data similar to those used in the simulation study, we checked that the results obtained with 10k SNPs were robust to subsampling and similar to those presented in Figure 5 (Figure S6).

The first two eigenvectors of the PCA displayed a horseshoe pattern for which the most extreme populations were the Dai population of China (21° N, 100° E) and the Karitiana population of Brazil (−13.5° N, 348° E). The observed arch agreed with the predictions of a UM for which the most ancient populations were Asian and the most recent ones were South American (Figure 5). Nevertheless, a constant ancestor population size in the UM did not allow a perfect fit to the data. For example, the number of present-day populations was overestimated by the UM ($n = 41$). In approximations of PC2 and PC3, the number of populations in the UM ($n = 31$) was closer to the number of present-day populations in the samples ($n = 30$), and

Figure 5. Human data analysis. *Top panel*: Geographic coordinates of 59 samples from the Simons Genome Diversity Project (Table S1). *Bottom panel*: PC plots for the genotype data (554,716 SNPs). The color gradient reflects the geographic distance to China. Blue: Dai (21° N, 100° E), Red: Karitiana (Brazil, −13.5° N, 348° E). The grey lines represent the predictions from the umbrella model with $n = 41$ populations (PC1-PC2 plot) and $n = 32$ populations (PC2-PC3 plot).

Figure 6. Comparison of PCA and STRUCTURE for human samples. For 59 Asian and American samples from the Simons Genome Diversity Project, PC1 is consistent with an umbrella model of evolution (*top panel*), whereas STRUCTURE wrongly suggests a fifty-fifty percent admixture model for Aleutian and Eskimo populations (*bottom panel*). The color gradient reflects the geographic distance to China. Blue: Dai (21° N, 100° E), Red: Karitiana (Brazil, −13.5° N, 348° E).

the second and the third eigenvectors of the PCA displayed a fish-like pattern that supported the predictions of the diffusion model.

To evaluate whether the sinusoidal patterns observed in PCA are more difficult to interpret than ancestry estimates obtained with other methods, we performed an analysis of population genetic structure with a program that reproduces the results of STRUCTURE [39, 18]. The STRUCTURE analysis identified two ancestral genetic groups corresponding to the Asian and South American populations in the PC plot (Figure 6). Asian genetic ancestry measured with STRUCTURE (blue color in Figure

6) exhibited a very high correlation with the first PC ($r = 99.8\%$). A direct interpretation of the STRUCTURE plot, exhibiting a geographic cline of admixture, could lead to the conclusion that Aleut and Eskimo populations resulted from secondary contact between Asian and American populations. This direct interpretation of the STRUCTURE estimates contradicts recent evidence that migrations connected to the peopling of the Aleutian islands are genetically linked to a single Siberian source related to Paleo-Eskimos [13]. The overall analysis showed that sinusoidal patterns observed in PC plots are not always artificial outcomes arising from spatially structured data. Those patterns could actually reveal the temporal nature of migration of peoples better than other types of population structure analysis.

4   Conclusion

This study provided a detailed mathematical analysis of the eigenvalues and eigenvectors obtained from the principal component analysis of a diffusion model of genetic variation. The model was based on hierarchy of population splits from a genetically homogeneous population without any geographical structure in the sample. Our results described continuous approximations of PC eigenvectors for large sample size. When singletons are removed from the genetic data, the PCA eigenvectors took simple mathematical expressions, defined as cosine functions of increasing periodicity. For unfiltered data sets, the first PC eigenvectors could be approximated by Bessel's functions that exhibit complex wave-like patterns.

Novembre and Stephens showed that sinusoidal patterns arise in principal components generally when PCA is applied to spatial data [34]. The waves were interpreted as mathematical artifacts that can be observed when covariance between locations decays with spatial distance. Their study suggested that the artificial patterns could be amplified by spatial interpolation methods used to display PC eigenvectors as geographic maps. Our results support the alternative evidence that wave-like patterns

could arise from genetic drift only, and may not be artificial outcomes of spatially structured data.

Several studies have argued that human genetic variation agrees with a model of isolation-by-distance, whereas others see a role for long-range migrations and bottlenecks [23]. Regarding the implication of the results for human data, we showed an example of complex arch and fish-like patterns observed PC plots when studying the migration of an Asian ancestral population to the Bering Straits and its dispersal throughout the Americas. Although the PC eigenvectors reproduced patterns observed in a one-dimensional equilibrium isolation-by-distance model, this multiple-island model represents an unlikely explanation for the genetic data [13]. In addition, an analysis of population structure based on a STRUCTURE-like algorithm led to misleading estimates of admixture coefficients in Beringian populations. The confounding effect of hierarchical population splits on admixture estimates was confirmed by coalescent simulations (Figure S7, see [27] for how to misinterpret STRUCTURE results). Thus STRUCTURE-like ancestry estimation algorithms did not lead to better explanations than the sinusoidal patterns obtained from the PCA of the human data. For the Asian and Amerindian samples, the patterns of variation observed in PC plots were consistent with a history of serial population splits from a unique ancestral homogeneous gene pool. This observation has an anthropological implication for the concept of human race: either a unique species or as many races as individuals are represented in the sample, making the concept of race inappropriate for the studied populations [28].

Appendices

Appendix A. Simulation of umbrella model. Simulations of the umbrella model corresponded to the following `ms` command [22]

```
./ms n L -s m
```

```
-I n 1 1 ... 1
-n 1 1 -n 2 1 ... -n n 1
-ej 1 n n-1 -ej 2 n-1 n-2 ... -ej n-1 2 1
```

Appendix B. Proof of the approximation result for filtered data sets. According to [32], a theoretical approximation for the mathematical expectation of the empirical covariance matrix $\mathbb{E}[\mathbf{C}]$ is defined by equation (4). For non-singleton variants, the average total branch length in the tree is equal to n and the average coalescence time for ancestor i and ancestor j is equal to the maximum between i and j. To find the eigenvectors and eigenvalues of $\mathbb{E}[\mathbf{C}]$, we studied the spectrum of matrix $\bar{T}\mathbb{E}[\mathbf{C}]/n^2$, which is equal to $\mathbb{E}[\mathbf{C}]/n$. We looked for an eigenvector $\mathbf{u}$ and his corresponding eigenvalue μ such that $\mathbb{E}[\mathbf{C}]\mathbf{u}/n = \mu\mathbf{u}$. We set $\lambda = n\mu$, so that we have $\mathbb{E}[\mathbf{C}]\mathbf{u} = \lambda\mathbf{u}$. To solve the last equation, we sought solutions of the form $\mathbf{u}(i) = u(i/n)$, $i = 1, \ldots, n$, where $u(x)$ is a continuous function determined by

$$\left(\frac{\bar{t}(i)}{n} - \frac{\bar{t}}{n}\right)\frac{1}{n}\sum_{j=1}^{n}u(j/n) + \frac{1}{n}\sum_{j=1}^{n}u(j/n)\frac{\bar{t}(j)}{n} - \frac{1}{n}\sum_{j=1}^{n}\max(i/n, j/n)u(j/n) = \mu u(i/n).$$

$$(16)$$

Approximating the sums by Riemann integrals, we obtained

$$\frac{\bar{t}(i)}{n} = \frac{1}{n}\sum_{k=1}^{n}\max(i/n, k/n) \approx \int_0^1 \max(x, y)dy = \frac{x^2 + 1}{2},$$

and

$$\frac{\bar{t}}{n} = \frac{1}{n^2}\sum_{k=1}^{n}\sum_{\ell=1}^{n}\max(k/n, \ell/n) \approx \int_0^1\int_0^1 \max(y, z)dydz = \frac{2}{3}.$$

With similar approximations for the other terms in equation (16), the discrete eigen-

value problem becomes a continuous eigenvalue problem

$$\left[\frac{x^2+1}{2} - \frac{2}{3}\right] \int_0^1 u(y)dy + \int_0^1 \int_0^1 \max(y,z)u(y)dydz - \int_0^1 \max(x,y)u(y)dy = \mu u(x).$$
(17)

Simplifying (17), we obtained

$$\frac{3x^2+2}{6} \int_0^1 u(y)dy + \frac{1}{2} \int_0^1 y^2 u(y)dy - \int_0^1 \max(x,y)u(y)dy = \mu u(x).$$
(18)

Differentiating (18) yielded

$$x \int_0^1 u(y)dy - \int_0^x u(y)dy = \mu u'(x).$$
(19)

A second differentiation led to the following differential equation

$$\mu u''(x) + u(x) = \int_0^1 u(y)dy.$$

The solutions of the differential equation have the following form

$$u(x) = A\cos\left(\frac{x}{\sqrt{\mu}}\right) + B\sin\left(\frac{x}{\sqrt{\mu}}\right) + C,$$
(20)

where

$$C = \int_0^1 u(y)dy,$$

and A and B are constant values determined by the boundary conditions described below. A first boundary condition was obtained by setting $x = 1$ into equation (17)

$$\mu u(1) = \int_0^1 \int_0^1 \max(y,z)u(y)dydz - \frac{2}{3}\int_0^1 u(y)dy.$$
(21)

Incorporating $u(x)$, as given by equation (20), into equation (21) led to

$$\mu u(1) = A\int_0^1 \int_0^1 \max(y,z)\cos\left(\frac{y}{\sqrt{\mu}}\right)dydz + B\int_0^1 \int_0^1 \max(y,z)\sin\left(\frac{y}{\sqrt{\mu}}\right)dydz.$$
(22)

Setting $x = 0$ in equation (19) gives $u'(0) = 0$ which leads to $B = 0$. Setting $x = 1$ gives $u'(1) = 0$, which implies that the eigenvalues are equal to $\mu_k = 1/k^2\pi^2$. As a consequence, equation (22) is equal to

$$
\begin{aligned}
\mu u_k(1) = A_k \frac{\cos(k\pi)}{k^2\pi^2} + \frac{C_k}{k^2\pi^2} &= A_k \int_0^1 \int_0^1 \max(y, z) \cos\left(k\pi y\right) dy dz \\
&= A_k \int_0^1 \frac{y^2 + 1}{2} \cos\left(k\pi y\right) dy \\
&= A_k \frac{\cos(k\pi)}{k^2\pi^2}.
\end{aligned}
$$

From this equations, we obtained that $C_k = 0$. This finished the proof of our main result: The normalized eigenvectors and their corresponding eigenvalues have the following form

$$
u_k(x) = \sqrt{2} \cos(xk\pi), \qquad \lambda_k = \frac{n}{k^2\pi^2}, \qquad k = 1, \ldots, n.
$$

Appendix C. Normalizing constant for the unfiltered data analysis. The normalizing constant for the unfiltered data analysis is defined by

$$
\begin{aligned}
\frac{1}{A_k^2} = {} & \left(J_1(\sqrt{2\lambda_k}n) - \rho_k Y_1(\sqrt{2\lambda_k}n) \right)^2 \left[\frac{4}{n^3\lambda_k^2} + \frac{2}{n^3\lambda_k^3} - 1 - \frac{2}{n^2\lambda_k} \right] \\
& + \left(J_1(\sqrt{2(\lambda_k n - 2)n}) - \rho_k Y_1(\sqrt{2(\lambda_k n - 2)n}) \right)^2 \left[1 + \frac{2}{n(\lambda_k n - 2)} \right] \\
& - \frac{4}{n(n\lambda_k)^{3/2}\sqrt{\lambda_k n - 2}} \left(J_1(\sqrt{2\lambda_k}n) - \rho_k Y_1(\sqrt{2\lambda_k}n) \right) \\
& \qquad\qquad \times \left(J_1(\sqrt{2(\lambda_k n - 2)n}) - \rho_k Y_1(\sqrt{2(\lambda_k n - 2)n}) \right),
\end{aligned}
$$

where $\rho_k = J_2(\sqrt{2\lambda_k}n)/Y_2(\sqrt{2\lambda_k}n)$.

References

[1] Arenas, M., François, O., Currat, M., Ray, N., Excoffier, L. (2013). Influence of admixture and paleolithic range contractions on current European diversity gradients. Mol. Biol. Evol., 30(1), 57-61.

[2] Austerlitz F., Jung-Muller B., Godelle B., Gouyon, P.-H. (1997). Evolution of coalescence times, genetic diversity and structure during colonization. Theor. Popul. Biol., 51, 148–164.

[3] Branco, C., Velasco, M., Benguigui, M., Currat, M., Ray, N., Arenas, M. (2018). Consequences of diverse evolutionary processes on American genetic gradients of modern humans. Heredity, 121(6), 548-556.

[4] Cavalli-Sforza, L. L. Edwards, A. W. F. (1967). Phylogenetic analysis: models and estimation procedures. Am. J. Hum. Genet., 19, 233–257.

[5] Cavalli-Sforza L.L., Edwards A.W.F., Geerts S. (1963). Analysis of human evolution. In:Genetics today: Proceedings of the 11th International Congress of Genetics, The Hague, The Netherlands. New York: Pergamon 3, 923-993.

[6] DeGiorgio, M., Jakobsson, M., Rosenberg, N. A. (2009). Explaining worldwide patterns of human genetic variation using a coalescent-based serial founder model of migration outward from Africa. Proc. Natl. Acad. Sci. U.S.A., 106, 16057–16062.

[7] DeGiorgio, M., Degnan, J. H. , Rosenberg, N. A. (2011). Coalescence-time distributions in a serial founder model of human evolutionary history. Genetics, 189, 579–593.

[8] Deshpande, O., Batzoglou, S., Feldman, M. W., Cavalli-Sforza, L. (2009). A serial founder effect model for human settlement out of Africa. Proc. Royal Soc. B., 276, 291-300.

[9] Diaconis, P., Goel, S., Holmes, S. (2008). Horseshoes in multidimensional scaling and local kernel methods. Ann. Appl. Stat., 2, 777-807.

[10] Duforet-Frebourg, N., Slatkin, M. (2016). Isolation-by-distance-and-time in a stepping-stone model. Theor. Pop. Biol., 108, 24-35.

[11] Excoffier,L., Foll, M., Petit R. J. (2009). Genetic consequences of range expansions. Annu. Rev. Ecol. Evol. Syst., 40, 481–501.

[12] Felsenstein, J. *Inferring Phylogenies*. Sunderland, MA: Sinauer associates, 2004.

[13] Flegontov, P., Altınışık, N. E., Changmai, P., Rohland, N., Mallick, S., Adamski, N. et al. (2019). Palaeo-Eskimo genetic ancestry and the peopling of Chukotka and North America. Nature, 570, 236-240.

[14] François, O., Currat, M., Ray, N., Han, E., Excoffier, L., Novembre, J. (2010). Principal component analysis under population genetic models of range expansion and admixture. Mol. Biol. Evol., 27(6), 1257-1268.

[15] François O, Jay F. (2020). Factor analysis of ancient population genomic samples. Nat. Commun. 11, 4661.

[16] François, O., Gain, C. (2021). A spectral theory for Wright's inbreeding coefficients and related quantities. PLoS Genet., 17(7), e1009665.

[17] Frichot, E., Schoville, S. D., Bouchard, G. François, O. (2012). Correcting principal component maps for effects of spatial autocorrelation in population genetic data. Front. Genet., 3, 254.

[18] Frichot, E., Mathieu, F., Trouillon, T., Bouchard, G., François, O. (2014). Fast and efficient estimation of individual ancestry coefficients. Genetics, 196(4), 973-983.

[19] Goebel, T., Waters, M. R., O'Rourke, D. H. (2008). The late Pleistocene dispersal of modern humans in the Americas. Science, 319(5869), 1497-1502.

[20] Hallatschek, O., Hersen, P., Ramanathan, S., Nelson, D. R. (2007). Genetic drift at expanding frontiers promotes gene segregation. Proc. Natl. Acad. Sci. U.S.A., 104, 19926–19930.

[21] Hotelling H. (1936). Relations between two sets of variates. Biometrika, 28, 321-377.

[22] Hudson, R. R. (2002). Generating samples under a Wright–Fisher neutral model of genetic variation. Bioinformatics, 18, 337-338.

[23] Hunley, K. L., Healy, M. E., Long, J. C. (2009). The global pattern of gene identity variation reveals a history of long-range migrations, bottlenecks, and local mate exchange: implications for biological race. Am. J. Phys. Anthropol., 139, 35-46.

[24] Jolliffe I. *Principal Component Analysis.* Springer New-York, 1986.

[25] Kimura, M. Diffusion models in population genetics. (1964). J. Appl. Probab. 1, 177–232.

[26] Kimura, M. *The Neutral Theory of Molecular Evolution.* Cambridge University Press, Cambridge, UK, 1983.

[27] Lawson, D. J., Van Dorp, L., Falush, D. (2018). A tutorial on how not to over-interpret STRUCTURE and ADMIXTURE bar plots. Nat. Commun., 9(1), 1-11.

[28] Lewontin, R. (1972). The apportionment of human genetic diversity. In: Dobzhansky T, Hecht M, Steere W, editors. Evolutionary biology, Vol. 6. New York: Appleton-Century-Crofts.

[29] Loève, M. Fonctions aléatoires du second ordre. (1948). In Levy, P. (ed.) *Processus Stochastiques et Mouvement Brownien*, Gauthier-Villars, Paris, France.

[30] Lynch, M., Hill, W. G. (1986). Phenotypic evolution by neutral mutation. Evolution, 40, 915-935.

[31] Mallick, S., Li, H., Lipson, M., Mathieson, I., Gymrek, M., Racimo, F., *et al.* (2016). The Simons genome diversity project: 300 genomes from 142 diverse populations. Nature, 538, 201-206.

[32] McVean G. (2009). A genealogical interpretation of principal components analysis. PLoS Genet., 5, e1000686.

[33] Menozzi P., Piazza A., Cavalli-Sforza L.L. (1978). Synthetic maps of human gene frequencies in Europeans. Science, 201, 786-792.

[34] Novembre, J., Stephens, M. (2008). Interpreting principal component analyses of spatial population genetic variation. Nat. Genet., 40, 646.

[35] Patterson N., Price A.L., Reich D. (2006). Population structure and eigenanalysis. PLoS Genet., 2, e0020190.

[36] Peter, B. M. (2021). Modelling complex population structure using F-statistics and Principal Component Analysis. bioRxiv.

[37] Phillips, L. B., Brown, G. P., Webb, J. K., Shine, R. (2006). Invasion and the evolution of speed in toads. Nature, 439, 803–803.

[38] Pierce, A. A., Zalucki, M. P., Bangura, M., Udawatta, M., Kronforst, M. R., Altizer, S., *et al.* (2014). Serial founder effects and genetic differentiation during worldwide range expansion of monarch butterflies. Proc. Royal Soc. B., 281, 2014-2230.

[39] Pritchard, J. K., Stephens, M., Donnelly, P. (2000). Inference of population structure using multilocus genotype data. Genetics, 155(2), 945-959.

[40] Ramachandran, S., Deshpande, O., Roseman, C. C., Rosenberg, N. A., Feldman, M. W., Cavalli-Sforza, L. L. (2005). Support from the relationship of genetic and geographic distance in human populations for a serial founder effect originating in Africa. Proc. Natl. Acad. Sci. U.S.A. 102, 15942-15947.

[41] Slatkin, M., Excoffier, L. (2012). Serial founder effects during range expansion: A spatial analog of genetic drift. Genetics, 191, 171–181.

[42] Waters, M. R. (2019). Late Pleistocene exploration and settlement of the Americas by modern humans. Science, 365(6449), eaat5447.

Supplementary materials

Figure S1. Approximation of PC eigenvalues and eigenvectors for filtered data from the UM ($n = 30$). Simulation of the umbrella model performed for $n = 30$ haploid individuals and $L = 18,002$ filtered single nucleotide polymorphism loci. *Top panels*: Approximations of PC eigenvalues by $L\lambda_k/n = L/\pi^2 k^2$ (left) and by the numerical values obtained from the discrete eigenvalue equations based on McVean's equation for $n = 30$ (right). Eigenvalues are displayed on a base 10 logarithm scale. *Bottom panels*: Approximations of PC eigenvectors by their continuous approximations, $\mathbf{u}_k(x) = \sqrt{2}\cos(k\pi x)$, for $x = i/n$, $k = 1$ to 4. PC eigenvectors are colored from red (most recent samples) to blue (most ancient samples) and the continuous approximations are represented by grey circles.

Figure S2. Wave patterns in PC plots for filtered data ($n = 30$). A simulation of the umbrella model was performed for $n = 30$ haploid individuals, and resulted in $L = 18,002$ unfiltered single nucleotide polymorphism loci. *Left column*: PC plots for the simulated data. *Right column*: Continuous plots derived from the asymptotic theory for $x = i/n$ and $k = 1$ to 4. Eigenvectors are represented in red (most recent samples) to blue (most ancient samples) colors.

Figure S3. Wave patterns in PC plots for unfiltered data ($n = 100$). Simulation of the umbrella model performed for $n = 100$ haploid individuals and $L = 300,000$ unfiltered single nucleotide polymorphism loci. *Left column*: PC plots for the simulated data. *Right column*: Continuous plots derived from the asymptotic theory (Bessel approximations). Eigenvectors are represented in red (most recent samples) to blue (most ancient samples) colors.

Figure S4. PC eigenvectors for unfiltered data ($n = 100$). Simulation of the umbrella model performed for $n = 100$ haploid individuals and $L = 300,000$ unfiltered single nucleotide polymorphism loci. Six PC eigenvectors for the simulated data and their prediction from the discrete system of equations. Eigenvectors are represented in red (most recent samples) to blue (most ancient samples) colors.

Figure S5. Approximation of PC eigenvalues and eigenvectors for unfiltered data ($n = 30$). Simulation of the umbrella model performed for $n = 30$ haploid individuals and $L = 300,000$ unfiltered single nucleotide polymorphism loci. *Top panels*: Approximations of the first PC eigenvalues by $L\lambda_k/n$ (continuous approximation) (left) and by the numerical values obtained from the discrete eigenvalue equations for $n = 30$ (right). Eigenvalues are displayed on a base 10 logarithm scale. *Bottom panels*: PC plots for the simulated data and continuous plots derived from the asymptotic theory. Eigenvectors are represented in red (most recent samples) to blue (most ancient samples) colors.

Figure S6. Human data analysis PC plots for 10k filtered genotypes from 59 individuals from the SGDP. The color gradient reflects the geographic distance to China. Blue: Dai (21° N, 100° E), Red: Karitiana (Brazil, −13.5° N, 348° E). The grey lines represent the predictions from the umbrella model with $n = 41$ populations (PC1-PC2 plot) and $n = 32$ populations (PC2-PC3 plot).

Figure S7. Ancestry coefficients under the UM. Simulation of the umbrella model performed for $n = 100$ haploid individuals and $L = 5,938$ filtered single nucleotide polymorphism loci. *Left*: First principal component. *Right*: Ancestry coefficient estimates from a STRUCTURE-like program run for $K = 2$.

Table S1. List of human samples used in the study

Instance.ID	Master.ID	Group.ID	Country	Lat.	Long.
A_Dai-5	A_Dai-5	Dai	China	21	100
S_Dai-2	HGDP01312	Dai	China	21	100
B_Dai-4	HGDP01308	Dai	China	21	100
S_Lahu-2	HGDP01320	China_Lahu	China	22	100
S_Lahu-1	HGDP01323	China_Lahu	China	22	100
S_Miao-2	HGDP01198	Miao	China	28	109
S_Miao-1	HGDP01191	Miao	China	28	109
S_Tujia-2	HGDP01098	Tujia	China	29	109
S_Tujia-1	HGDP01095	Tujia	China	29	109
A_Han-4	A_Han-4	Han	China	32.26	114
S_Han-1	HGDP00783	Han	China	32.26	114
B_Han-3	HGDP00775	Han	China	32.26	114
S_She-1	HGDP01335	She	China	27	119
S_She-2	HGDP01333	She	China	27	119
S_Mongola-2	HGDP01223	Mongola	China	45	111
S_Mongola-1	HGDP01228	Mongola	China	45	111
S_Korean-1	ND19394	Korean	South Korea	37.60	127
S_Korean-2	NA00726	Korean	South Korea	37.60	127
S_Daur-2	HGDP01215	Daur	China	48.49	124
S_Oroqen-2	HGDP01211	Oroqen	China	50.43	126.5
S_Oroqen-1	HGDP01203	Oroqen	China	50.43	126.5
S_Japanese-2	HGDP00773	Japanese	Japan	37.85	139
S_Japanese-1	HGDP00749	Japanese	Japan	37.85	139
S_Japanese-3	S_Japanese-3	Japanese	Japan	37.85	139.001
S_Hezhen-1	HGDP01240	Hezhen	China	47.49	133.5
S_Hezhen-2	HGDP01242	Hezhen	China	47.49	133.5
S_Even-3	Nlk3	Even	Russia	57.53	135.88
S_Even-1	Nlk1	Even	Russia	57.53	135.88
S_Even-2	Nlk18	Even	Russia	57.53	135.88
S_Yakut-1	HGDP00956	Yakut	Russia	62.98	129.5
S_Yakut-2	HGDP00951	Yakut	Russia	62.98	129.5
S_Itelman-1	S_Itelman-1	Itelmen	Russia	57	157

S_Aleut-2	Ale20	Aleut	Russia	55.18	166
S_Aleut-1	Ale22	Aleut	Russia	55.18	166
S_Eskimo_Naukan-2	Nesk_25	Eskimo_Naukan	Russia	66.02	169.71
S_Eskimo_Naukan-1	Nesk_22	Eskimo_Naukan	Russia	66.02	169.71
S_Chukchi-1	S_Chukchi-1	Chukchi	Russia	69	169
S_Eskimo_Chaplin-1	Esk29	Eskimo_Chaplin	Russia	64.48	172.86
S_Chipewyan-2	NAD59	Chipewyan	Canada	59.60	-107.3
S_Chipewyan-1	NAD93	Chipewyan	Canada	59.60	-107.3
S_Cree-2	Cree6	Cree	Canada	50.30	-102.5
S_Cree-1	Cree5	Cree	Canada	50.30	-102.5
S_Pima-2	HGDP01044	Pima	Mexico	29	-108
S_Pima-1	HGDP01047	Pima	Mexico	29	-108
S_Nahua-2	S_Nahua-2	Nahua	Mexico	19.01	-99.06
S_Mayan-1	HGDP00855	Mayan	Mexico	19	-91
S_Mayan-2	HGDP00857	Mayan	Mexico	19	-91
S_Zapotec-2	S_Zapotec-2	Mexico_Zapotec	Mexico	16.50	-97.2
S_Zapotec-1	S_Zapotec-1	Mexico_Zapotec	Mexico	16.50	-97.2
S_Piapoco-1	HGDP00702	Piapoco	Colombia	3	-68
S_Piapoco-2	HGDP00706	Piapoco	Colombia	3	-68
S_Surui-2	HGDP00852	Surui	Brazil	-11	-62
S_Surui-1	HGDP00846	Surui	Brazil	-11	-62
A_Karitiana-4	A_Karitiana-4	Karitiana	Brazil	-10	-63
B_Karitiana-3	HGDP01015	Karitiana	Brazil	-10	-63
S_Karitiana-1	HGDP01012	Karitiana	Brazil	-10	-63
S_Quechua-1	NA11201	Quechua	Peru	-13.5	-72
S_Quechua-3	S_Quechua-3	Quechua	Peru	-13.5	-72
S_Quechua-2	NA11200	Quechua	Peru	-13.5	-72