X chromosomes show relaxed selection and complete somatic dosage compensation across *Timema* stick insect species

Darren J. Parker[1,2,3], Kamil S. Jaron[1,2,4], Zoé Dumas[1], Marc Robinson-Rechavi[1,2], and Tanja Schwander[1]

1. Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland
2. Swiss Institute of Bioinformatics, Lausanne, Switzerland
3. School of Natural Sciences, Bangor University, United Kingdom
4. Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3FL

Correspondence should be addressed to d.parker@bangor.ac.uk or Tanja.Schwander@unil.ch

Abstract

Sex chromosomes have evolved repeatedly across the tree of life. As they are present in different copy numbers in males and females, they are expected to experience different selection pressures than the autosomes, with consequences including a faster rate of evolution, increased accumulation of sexually antagonistic alleles, and the evolution of dosage compensation. Whether these consequences are general or linked to idiosyncrasies of specific taxa is not clear as relatively few taxa have been studied thus far. Here we use whole-genome sequencing to identify and characterize the evolution of the X chromosome in five species of *Timema* stick insects with XX:X0 sex determination. The X chromosome had a similar size (approximately 11% of the genome) and gene content across all five species, suggesting that the X chromosome originated prior to the diversification of the genus. Genes on the X showed evidence of relaxed selection (elevated dN/dS) and a slower evolutionary rate (dN + dS) than genes on the autosomes, likely due to sex-biased mutation rates. Genes on the X also showed almost complete dosage compensation in somatic tissues (heads and legs), but dosage compensation was absent in the reproductive tracts. Contrary to prediction, sex-biased genes showed little enrichment on the X, suggesting that the advantage X-linkage provides to the accumulation of sexually antagonistic alleles is weak. Overall, we found the consequences of X-linkage on gene sequences and expression to be similar across *Timema* species, showing the characteristics of the X chromosome are surprisingly consistent over 30 million years of evolution.

Introduction

One of the most common forms of genomic variation between individuals within species stems from sex chromosomes. Sex chromosomes differ in copy number between males and females, which has a large effect on the evolutionary forces acting on genes located on them. Specifically, when males are the heterogametic sex (i.e., in XY or X0 systems), the genes on the X chromosome are present in only a single copy in males, while genes on the autosomes are present in two copies. This fundamental difference is expected to have a large effect on the evolutionary forces acting on genes located on the X chromosome, with consequences including a faster rate of sequence evolution, increased accumulation of sexually antagonistic alleles, and the evolution of dosage compensation mechanisms (Bachtrog *et al.*, 2011; Wright *et al.*, 2016; Lenormand & Roze, 2021). Such effects should apply across taxa, meaning we should observe common patterns of X chromosome evolution in diverse species. Despite this, studies of sex chromosome evolution have shown a great deal of variation among study systems, including differences in the influence selection and drift have on the content of the X and in the extent of dosage compensation (Bachtrog *et al.*, 2011; Gu & Walters, 2017). Currently, it is difficult to understand the factors that govern this variation as only few taxa (typically with only a single or few representative species) have been studied. To elucidate these factors, studies of X chromosome evolution are thus needed from multiple species from a wide range of taxa (Palmer *et al.*, 2019). Such studies will allow us to disentangling general sex chromosome-linked patterns from species-specific patterns, and allow us to develop a fuller understanding of sex chromosome evolution.

The rate of sequence evolution is expected to differ between the X chromosome and autosomes for two main reasons. Firstly, the X is hemizygous in males, meaning that recessive or partially recessive mutations in X-linked genes will be more exposed to selection than mutations in genes on the autosomes, allowing for beneficial mutations to be fixed and deleterious mutations to be purged more effectively (Charlesworth *et al.*, 1987). On the other hand, as the X is only present in a single copy in males, its effective population size is expected to be smaller than that of the autosomes. This means that the effects of drift will also be stronger for genes on the X chromosome than on the autosomes, which will tend to reduce the fixation rate for advantageous mutations but increase it for deleterious mutations (Wright, 1931; Vicoso & Charlesworth, 2009). Together, these effects have been used as an explanation for the overall faster evolution of the X chromosome (the faster-X effect) seen in many species (Mank *et al.*, 2010; Meisel & Connallon, 2013; Parsch & Ellegren, 2013; Charlesworth *et al.*, 2018). In addition, the difference in X chromosome copy number between males and females is expected to facilitate the fixation of sexually antagonistic alleles (Rice, 1984; Gibson *et al.*, 2002; Mullon *et al.*, 2012). This is because the X chromosome spends two-thirds of its time in females, giving an advantage to dominant female-beneficial alleles on the X. In addition, recessive alleles on the X will be exposed to selection on the X in males, giving an advantage to male-beneficial alleles. These complex forces have the potential to shape how sexually antagonistic variation is distributed across the genome, which can in turn influence broad evolutionary processes such as speciation (Coyne & Orr, 2004; Payseur *et al.*, 2018) and sexual conflict (Bachtrog *et al.*, 2011; Mank *et al.*, 2014; Wilkinson *et al.*, 2015).

The fact that genes on the X are present in different copy numbers in males and females can also create a problem for gene expression, as for many genes expression is proportional to their copy number (Birchler & Veitia, 2012; Birchler, 2016). As such, species with differentiated sex chromosomes should have evolved dosage compensation mechanisms to equalise expression of the X chromosome in males and females (Ohno, 1967; Charlesworth, 1978, 1996). Note that such dosage compensation mechanisms can evolve either in response to sex chromosomes differentiation or alongside it (Lenormand *et al.*, 2020). Dosage compensation has been demonstrated across a wide range of taxa (Disteche, 2012; Mank, 2013; Gu & Walters, 2017), including model species such as *Drosophila melanogaster* (Conrad & Akhtar, 2012) and *Caenorhabditis elegans* (Meyer, 2000) where this phenomenon has been studied in detail (Parkhurst & Meneely, 1994; Lucchesi, 1998; Meyer, 2000; Straub & Becker, 2011; Conrad & Akhtar, 2012). Despite this commonality, it has become increasingly clear that the extent to which genes on the X are dosage compensated varies among species. Several studied species show only partial or no dosage compensation (Mank, 2013). The extent of dosage compensation may also differ by tissue type, with reduced dosage compensation observed in the reproductive tracts of e.g. *C. elegans* (Kelly *et al.*, 2002; Pirrotta, 2002) or *D. melanogaster* (Oliver, 2002; Meiklejohn *et al.*, 2011; Mahadevaraju *et al.*, 2021). However, it is not clear how widespread tissue-specific dosage compensation is, as work in non-model species often use whole-body samples for examining expression (Gu & Walters, 2017)**.**

Here we expand our knowledge of the evolutionary characteristics of sex chromosomes by identifying and studying the X chromosome in *Timema* stick insects. Aspects of X chromosome evolution have been previously studied in several insect orders (Odonata (Chauhan *et al.*, 2021), Hemiptera (Pal & Vicoso, 2015; Richard *et al.*, 2017), Orthoptera (Rayner *et al.*, 2021), Strepsiptera (Mahajan & Bachtrog, 2015), Coleoptera (Prince *et al.*, 2010; Mahajan & Bachtrog, 2015), and Diptera (Bone & Kuroda, 1996; Marín *et al.*, 1996; Deng *et al.*, 2011; Nozawa *et al.*, 2014; Jiang *et al.*, 2015; Vicoso & Bachtrog, 2015; Rose *et al.*, 2016)). However, to date no studies have examined X chromosome evolution in stick insects (Phasmatodea), an order which originated approximately 130 mya (Simon *et al.*, 2019) and contains around 3100 extant species (Bradler & Buckley, 2018). First, we identified the X chromosome in five species of *Timema* that diverged approximately 30 mya (Riesch *et al.*, 2017). *Timema* have an XX/X0 sex determination system (Schwander & Crespi, 2009). To determine if genes on the X chromosome show the predicted faster rate of sequence evolution than genes on the autosomes, we examined sequence evolution rates in each species. In addition, we tested if the X chromosome is enriched for sexually antagonistic alleles. This was done by examining if genes with sex-biased expression are enriched on the X chromosome, as the evolution of sex-biased expression is thought to be driven primarily by sexually antagonistic selection (Ellegren & Parsch, 2007; Innocenti & Morrow, 2010; Griffin *et al.*, 2013). Finally, we examined if the genes on the X chromosome are dosage compensated by comparing male and female gene expression in three composite tissues (heads, legs, and reproductive tracts). Our study thus provides a detailed study of several key aspects of X chromosome evolution in a previously unstudied group, revealing that the characteristics of these X chromosomes are conserved over at least 30 million years of evolution.

Materials and methods

Sample collection and sequencing

We used a combination of available genomic data from females in addition to newly collected data for males for five sexually reproducing species of *Timema* (*T. bartmani, T. cristinae*, *T. californicum*, *T. podura*, and *T. poppensis*). Reads from five females per species were downloaded from NCBI (Bioproject accession number: PRJNA670663). We collected four males from each of the five species from natural populations in California, from the same (or a geographically very close) population as the available females (Table S8). DNA extractions were done on whole-body adult males using the Qiagen Mag Attract HMW DNA kit following the manufacturer instructions. Sequencing libraries were generated for each male using a TruSeq DNA nano prep kit (550bp insert size). Libraries were then sequenced using Illumina HiSeq 2500 at the Lausanne Genomic Technologies Facility.

Using coverage to identify X-linked scaffolds

Reads were trimmed before mapping using Trimmomatic (v. 0.36) (Bolger *et al.*, 2014) to remove adapter and low-quality sequences (options: ILLUMINACLIP:3:25:6 LEADING:9 TRAILING:9 SLIDINGWINDOW:4:15 MINLEN:90). Reads from each individual were mapped to their species' reference genome (Jaron *et al.*, 2022) (Bioproject accession number: PRJEB31411) using BWA-MEM (v. 0.7.15) (Li, 2013). Multi-mapping and poor quality alignments were filtered (removing reads with XA:Z or SA:Z tags or a mapq < 30). PCR duplicates were removed with Picard (v. 2.9.0) (http://broadinstitute.github.io/picard/). Coverage was then estimated for all scaffolds at least 1000 bp in length using BEDTools (v. 2.26.0) (Quinlan & Hall, 2010). Per base coverage distributions were inspected visually for each individual and libraries with extremely non-normal coverage distributions were excluded from further analysis (Fig. S1-S5) .

To compare coverage between males and females, coverage was first summed for all male and all female libraries per scaffold (for scaffolds at least 1000 bp in length). Male and female coverage was then normalised by modal coverage to adjust for differences in overall coverage. X-linked scaffolds were then identified using the $\log_2$ ratio of male to female coverage. Autosomal scaffolds should have equal coverage in males and females ($\log_2$ ratio of male to female coverage ≈ 0) and X-linked scaffolds should have half the coverage in males as in females ($\log_2$ ratio of male to female coverage ≈ -1). Each species showed frequency peaks near these values (Fig. S6). X linked scaffolds were classified in two ways: a 'liberal' classification (following (Vicoso & Bachtrog, 2015)) whereby scaffolds were classified as X-linked if the $\log_2$ ratio of male to female coverage < autosomal peak - 0.5, and a 'stringent' classification (following (Pal & Vicoso, 2015)) whereby scaffolds were classified as X-linked if the $\log_2$ ratio of male to female coverage was within 0.1 of the value of the X-linked peak.

To compare X-linked scaffolds between species, we examined the overlap of one-to-one orthologs (previously identified in (Jaron *et al.*, 2022)) on the X-linked scaffolds. To determine if the overlap of X-linked orthologs between species was greater than expected, we used the

171   SuperExactTest package (v. 0.99.4) (Wang *et al.*, 2015) in R (R Core Team, 2017).
172   Additionally, we used a tblastn approach to identify which regions of the *Bacillus rossius*
173   genome correspond to the 210 shared X linked orthologs in *Timema* (maximum e-value = 1 x
174   10^{-20}, minimum query coverage = 50%).
175

Heterozygosity, nucleotide diversity, effective population size

To calculate heterozygosity, reads were mapped to the reference genomes as described
above. We then additionally performed indel realignment with GATK (v. 3.7) (Van der Auwera
et al., 2013). A maximum-likelihood estimate of the number of heterozygous sites per scaffold
was then calculated using AngsD (v. 0.921) (Korneliussen *et al.*, 2014) (options: -doSaf 1 -gl
1 -minQ 20 -minMapQ 40 -fold 1 -doCounts 1, with a minimum depth of 5 and a maximum
depth of twice the median genome coverage once sites with 0 coverage were excluded) for
each sample (for all scaffolds at least 1000 bp in length). The proportion of heterozygous sites
was then calculated for each scaffold and the median proportion of heterozygous sites was
weighted by the number of covered sites on a scaffold. We used a weighted median for
scaffolds as shorter contigs are typically enriched for repeats and therefore less likely to give
reliable estimates of heterozygosity. A similar approach was used to calculate pairwise
nucleotide diversity for each species, but using only female samples (options: -doSaf 1 -gl 1 -
minQ 20 -minMapQ 40 -fold 1 -doCounts 1 with a minimum depth of 5 and a maximum depth
of twice the median genome coverage once sites with 0 coverage were excluded). The number
of individuals (-nind) and the minimum number of individuals a site must be present in (-minind)
were both set to the number of samples analysed. Effective population size (Ne) was
estimated for the X and autosomes using the estimates of nucleotide diversity, assuming that
nucleotide diversity is equal to $4Ne\mu$ (μ = mutation rate). Estimates of the mutation rate were
taken from *Drosophila melanogaster* (2.8×10^{-09}) (Keightley *et al.*, 2014) and *Heliconius*
melpomene (2.9×10^{-09}) (Keightley *et al.*, 2015).

Excluding male achiasmy

An absence of recombination in males could influence the genetic diversity of the X and
autosomes. Previous work in different phasmid species has shown that males do recombine
(White, 1976; Marescalchi *et al.*, 1986), however, this has not been shown in *Timema*. To
examine this, we re-surveyed images of Giemsa-stained chromosome preparations from
(Schwander & Crespi, 2009) which were originally generated for karyotyping purposes.
Specifically we looked for the presence and position of chiasmata on pre-metaphase plates
obtained from male testes.

Selection analyses

210   We took values of dN/dS (number of non-synonymous substitutions per non-synonymous site
211   / number of synonymous substitutions per synonymous site) for each species branch from
212   Jaron et al (2022). Briefly, branch-site models with rate variation at the DNA level (Davydov et

al., 2019) were run using the Godon software (https://bitbucket.org/Davydov/godon/, version 2020-02-17, option BSG --ncat 4) for each gene with an ortholog found in at least three of the species of *Timema* used here. Godon estimates the proportion of sites evolving under purifying selection (p0), neutrality (p1), and positive selection. We used only sites evolving under purifying or neutrality to calculate dN/dS. To test for differences in dN/dS between the X and autosomes, we used Wilcoxon tests with p-values adjusted for multiple comparisons using Benjamini and Hochberg's algorithm (Benjamini & Hochberg, 1995). GC was calculated using the seqinR package (v.4.2-8) (Charif & Lobry, 2007) in R. We examined the occurrence of positive selection with two approaches. First, as in Jaron et al (2022), we used the branch-site models as described above to identify branches with evidence for positively selected sites. However, such branch-site models may lack power to detect positive selection so in addition we also used GODON to perform site-based model comparisons. Specifically, we compared the fit of model M8 (which allows for positively selected sites) and M8a (which does not allow for any positively selected sites) (Yang *et al.*, 2000), also with rate variation at the DNA level (Davydov et al., 2019). The two models were compared using a log-likelihood ratio test for each gene. p-values were corrected for multiple tests using Benjamini and Hochberg's algorithm. We then used a Fisher's exact test to determine if positively selected genes were overrepresented on the X.

Gene expression analyses

RNA-seq reads from three composite tissues (heads, legs and reproductive tracts) for males and females (3 replicates per sex) for each of the five species are publically available (Parker *et al.*, 2019a; b) (Bioproject accession number: PRJNA392384). Adapter sequences were removed using Cutadapt (v. 2.3) (Martin, 2011) before quality trimming reads with Trimmomatic (v. 0.36) (Bolger *et al.*, 2014) (options: LEADING:10 TRAILING:10 SLIDINGWINDOW:4:20 MINLEN:80). Trimmed reads were then mapped to reference genomes using STAR (v. 2.6.0c, default options). HTSeq v.0.9.1 (Anders *et al.*, 2015) was used to count the number of reads uniquely mapped to the exons of each gene, with the following options (htseq-count --order=name --type=exon --idattr=gene_id --stranded=reverse). Expression analyses were performed using the Bioconductor package EdgeR (v 3.32.1) (Robinson *et al.*, 2010), and done separately for each species and tissue. Normalisation factors for each library were computed using the TMM method and were used to calculate normalised expression levels (either FPKM (Fragments Per Kilobase of transcript per Million mapped reads) or TPM (Transcripts Per Million mapped reads)). For the main analyses, genes with low expression (less than 2 FPKM (or TPM) in 2 or more libraries per sex) were excluded. This filtering step was used to exclude any sex-specifically expressed genes as our goal was to examine how the expression of genes differs in males and females. This decision could influence our results if sex-limited gene expression was extensive on the X chromosome. To investigate this we repeated our analysis with the inclusion of sex-limited genes and found similar results to the main analyses (Fig. S7 - S8). To examine dosage compensation, we used the $\log_2$ ratio of mean male expression level to female expression level and used Wilcoxon tests (adjusted for multiple testing using Benjamini and Hochberg's algorithm (Benjamini & Hochberg, 1995)) to determine if this ratio differed between genes on the autosomal and X-linked scaffolds.

To determine the significance of sex on gene expression, we fit a generalised linear model
(GLM) with a negative binomial distribution with sex as an explanatory variable and used a
GLM likelihood ratio test to determine the significance for each gene. P-values were then
corrected for multiple tests using Benjamini and Hochberg's algorithm (Benjamini & Hochberg,
1995). In the main analysis sex-biased genes were then defined as genes that showed
difference in expression between males and females with a false-discovery rate (FDR) < 0.05
(we also repeated analyses with the additional condition that genes must show a greater than
two fold difference in expression to ensure our results are robust to the effects of sex-biased
allometry (Montgomery & Mank, 2016)). Note that all genes not classified as sex-biased were
classified as unbiased genes. We then examined if male- or female- biased genes were under-
or over- represented on autosomal or X-linked scaffolds using Fisher's exact tests (adjusted
for multiple-testing using Benjamini and Hochberg's algorithm (Benjamini & Hochberg, 1995)).
To examine species by sex interactions in gene expression, we used a similar similar GLM
approach as above, with sex, species, and species by sex interaction as explanatory variables
for genes with an ortholog in each species. Genes with an FDR < 0.1 for the interaction term
were considered to have significant species by sex interaction. To determine if sex-biased
genes on the X were underrepresented for species by sex interactions, we used a Fisher's
exact test.
276
It is possible that dosage compensation may vary along the X chromosome. Unfortunately,
our current reference genomes are too fragmented to investigate this question. Previous work
by Nosil et al. (Nosil *et al.*, 2018) however produced a more contiguous genome assembly of
one of our study species, *T. cristinae,* which has been further constructed into a linkage map.
To use this synteny information for each of our species, we anchored the scaffolds from each
of our genome assemblies to the Nosil et al. (Nosil *et al.*, 2018) reference genome (BioProject
Accession PRJNA417530) using MUMmer (version 4.0.0beta2) (Marçais *et al.*, 2018) with
parameter --mum. The alignments were processed by other tools within the package: show-
coords with parameters -THrcl to generate tab-delimited alignment files and dnadiff to
generate 1-to-1 alignments. We used only uniquely anchored scaffolds for which we were able
to map at least 10k nucleotides to the Nosil et al. (2018) reference genome. Nosil et al (2018)
indicated that linkage group 13 was the X chromosome ((Nosil *et al.*, 2018)). To determine if
this was correct, we repeated our coverage analyses on the Nosil et al. (2018) assembly. From
this, we found that most scaffolds that make up linkage group 13 did not show reduced
coverage in males (Fig. S9). In addition, several scaffolds from other linkage groups did show
reduced coverage in males (Fig. S9). In order to use as much of the synteny information as
possible we "cleaned" the Nosil et al. (2018) assembly by removing X-linked scaffolds from
autosomal linkage groups 1-12. These scaffolds were then assigned to a new, unordered
collection of scaffolds from the X chromosome, together with X-linked scaffolds from linkage
group 13 and from those not assigned to any linkage group in Nosil et al. (2018). Scaffolds
from linkage group 13 that were not X-linked were assigned to linkage group NA. We classified
scaffolds in the Nosil et al. (2018) assembly as X-linked if most of a scaffold was covered by
aligned scaffolds from our assembly assigned to the X rather than autosomes (i.e. if aligned
scaffolds assigned as X covered more than twice as many bases as those assigned as
autosomal for a particular scaffold, it was classed as X-linked).

Results

Identifying X-linked scaffolds

We used a coverage approach to identify X-linked scaffolds in our previous genome assemblies (Jaron *et al.*, 2022). Genomic reads from four males and five females from each species were mapped onto the corresponding reference genome. After filtering low-quality alignments and non-uniquely mapping reads (see Methods) the median coverage per sample ranged from 11x to 31x (18.5x on average, see Table S1). Visual inspection of coverage distributions (Figs S1-S5) found that while most libraries had either one (in females) or two (in males) coverage peaks, three libraries (*T. podura* (H56, Fig. S4), *T. poppensis* (ReSeq Ps08 and Reseq_Ps12, Fig. S5)) did not show a clear coverage peak, and were excluded from all further analyses.

To identify X-linked scaffolds, we used the $\log_2$ ratio of (normalised) coverage of males to females. As males have only a single copy of the X and females have two, X linked scaffolds should have twice as much coverage in females as males ($\log_2$ male:female coverage ≈ -1). Autosomes are expected to have the same coverage in both sexes (log2 male:female coverage ≈ 0). Considering all scaffolds with a $\log_2$ ratio of male to female coverage < autosomal peak -0.5 to be X-linked, we classified between 12 and 14 % of each genome as X-linked (Table 1, Fig. S6), which fits well with the X chromosome size observed in karyotypes (Schwander & Crespi, 2009). This approach may mean some autosomal contigs may be misclassified as X-linked, thus we also repeated all our analyses with a more stringent classification scheme (scaffolds with a $\log_2$ ratio of male to female coverage within 0.1 of the X linked peak) (Table S2, Fig. S6). Using the more stringent classification scheme produced very similar results (not shown). Of note, most differences between the classification schemes are for short scaffolds (1000-4999 bp), which represent ~20% of the genome assemblies. When these are excluded the two classification schemes classify almost the same set of scaffolds as X-linked (Fig. S10, Table S3). Finally, we also examined the heterozygosity of the X in males. As expected, the heterozygosity of the X is close to zero and much lower than on the autosomes, corroborating our X-linked scaffold assignments (Fig. S11).

Table 1 | Number of genes and scaffolds classified as X-linked or autosomal

Species	Min scaffold length	% genome X-linked	N Autosomal genes	N X-linked genes	N genes not classified
T. bartmani	1000	12.11	12804	1244	18
T. cristinae	1000	12.21	12542	1316	24
T. californicum	1000	12.22	13207	1344	12
T. podura	1000	13.75	15069	1457	3
T. poppensis	1000	12.40	14115	1454	36

The X chromosome is conserved across *Timema*

Comparing orthologs across the five different species shows X chromosome gene content is conserved, with >90% of X-linked orthologs shared between the 5 species (Fig. 1, Fig. S12). This overlap is much greater than expected by chance (FDR < 6.863 x 10^{-316}). This suggests

that the X chromosome is homologous in all five species, which last shared a common ancestor approximately 30 million years ago (Riesch *et al.*, 2017). Additionally, we used species genome alignments to an independent *T. cristinae* genome assembly (Nosil *et al.*, 2018) to assign scaffolds to linkage groups based on a single reference. By applying coverage analyses we were able to identify and correct the X linked scaffolds in the Nosil et al. 2018 assembly (see Methods). Using this corrected reference, we found that contigs aligned to X-linked scaffolds showed reduced coverage in males but not females in each species (Figs S13-S17), again indicating the X chromosome is the same in all species. Finally, using BLAST we found that the majority (72%) of the shared X-linked genes in *Timema* (for which we were able to obtain a significant hit) were also present on the X chromosome of *Bacillus rossius* (Fig. S18). The split between *Timema* and all other extant phasmids (the Euphasmatodea, which includes *Bacillus*) occurred approximately 120 mya (Simon *et al.*, 2019), suggesting that the X chromosome in phasmids predates this split.

Fig. 1 | The X chromosome is conserved between *Timema* species. A. Venn-diagrams showing the number of shared X-linked orthologs between species. **B.** Number of shared orthologs (expected, maximum possible). The observed amount of overlap was much greater than expected in all comparisons (FDR < 6.863 x 10^{-316}). Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps = *T. poppensis*, and Tpa = *T. podura*.

The X chromosome has reduced genetic variation

We used female samples to examine two related measures of genetic variation: heterozygosity and nucleotide diversity (π). Only female samples were used, as males are hemizygous for the X, which would affect the estimation of heterozygosity on the X directly. It would also affect the estimation of nucleotide diversity indirectly, as X haplotypes containing recessive lethals on the X will be absent in male samples. We found that both heterozygosity and nucleotide diversity (π) were lower on X-linked than on autosomal scaffolds (Fig. S11, S19), as expected. In all species, the ratios of X to autosomes for both of these measures (heterozygosity = 0.16 to 0.62 (Fig. 2), π = 0.19 to 0.48 (Fig. 2)) were lower than the 0.75 that would be expected from the reduced effective population size of the X relative to the autosomes. This pattern was also seen when comparing the X to each of the autosomal linkage groups individually (Fig. S20, S21). Nucleotide diversity estimates were then used to estimate the effective population size of each species. From this, we estimated the autosomal effective population size in *Timema* to range from ~150,000 (*T. poppensis*) to ~2,000,000 (*T. podura*) (Table S4). Finally, we corroborated that male meiosis was chiasmate, which was revealed by the presence of at least one chiasma per autosome on pre-metaphase plates (Fig. S22).

Fig. 2 | Genetic variation on the X chromosome is lower than expected. A. Ratio of heterozygosity on the X and autosomes in 3 to 5 females per species. **B.** Ratio of pairwise nucleotide diversity (π) on the X and autosomes. Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps = *T. poppensis*, and Tpa = *T. podura*. The dotted lines indicate the neutral expectation in a population with a balanced sex ratio.

The X chromosome shows evidence for reduced purifying selection but no differences in positive selection

Genes on the X chromosome show an elevated dN/dS relative to genes on the autosomes in each species (Fig. 3A). This effect remains when accounting for differences in GC and mean expression level (permutation ANOVA $p < 0.05$ for all species). This increased ratio appears to be driven by reduced purifying selection, as X-linked genes are not enriched for positively selected genes (Fig. 3B, Table S9).

Although dN/dS is elevated for genes on the X, overall divergence (branch length, dN + dS) is lower for genes on the X than genes on the autosomes (Fig. S23A). More specifically, dS is lower on the X for all species (Fig. S23B) and values of dN are only higher on the X for two species (*T. podura* and *T. poppensis*, Fig. S23C). The reduction in dS on the X is likely due to an overall lower mutation rate in the female than male germline, resulting in fewer mutations on the X due to the X's more frequent transmission through females (Kirkpatrick & Hall, 2004; Ellegren, 2007). Taken together, this suggests that while the genes on the X are subject to reduced purifying selection, the overall divergence of X-linked genes is smaller.

Fig. 3 | Sequence evolution on the X and autosomes. A. Average dN/dS across genes**.**
Error bars indicate standard error. Asterisks indicate the significance (FDR) of Wilcoxon tests
(***<0.001, **<0.01, *<0.05). **B.** Proportion of positively selected genes. Positively selected
genes were not enriched on the X chromosome (Fisher's exact test p value = 0.50). Species
names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps =
T. poppensis, and Tpa = *T. podura*.

***Timema* show complete dosage compensation in heads and legs, but no dosage**
compensation in reproductive tracts

We examined if the X chromosome is dosage compensated in *Timema* by comparing gene
expression in males and females in three different composite tissues (heads, legs, and
reproductive tracts) for each of our five species. While for the main analysis we calculated
expression values as FPKM, we also repeated our analyses using TPM. Analyses based on
TPM showed very similar results and are provided as supplemental figures and tables. We
examined the $\log_2$ ratio of male to female expression on the X and the autosomes. For the
autosomes and a dosage compensated X, the $\log_2$ value should be approximately 0, whereas
a non-dosage compensated X would have a value of approximately -1. We found that in heads
and legs the ratio was close to 0 with only small differences between the X and the autosomes
(Figs. 4, S24, S25), indicating almost-complete complete dosage compensation in these
tissues. By contrast, in the reproductive tracts the ratio of male to female expression for genes
on the X is close to -1 (Figs. 4, S24, S25), indicating an absence of dosage compensation in

this tissue. This observation was also seen when comparing the X to each of the autosomal
linkage groups individually (Figs. S26, S27). An alternative, mutually non-exclusive possibility
is that the greatly reduced expression observed in genes on the X in male reproductive tracts
is due to a large enrichment of female-biased genes and depletion of male-biased genes.
Although we cannot formally exclude this possibility, three lines of evidence indicate that lack
of dosage compensation in reproductive tracts is the best explanation for our findings. Firstly,
a lack of dosage compensation is expected to result in a two-fold reduction of expression,
meaning X linked genes in males should show a major peak with a two-fold reduction in
expression (Mank & Ellegren, 2009; Vicoso *et al.*, 2013; Pal & Vicoso, 2015), which is what
we observe (Fig. 5). Secondly, if the X chromosomes facilitates the accumulation of sex-
biased genes, we should be able to observe this effect in all tissues (Jaquiéry *et al.*, 2021).
While we find a large enrichment of female-biased genes and a depletion of male-biased
genes on the X in the reproductive tracts, this is not found in the other tissues, with only a
slight enrichment of female-biased genes in the heads of two species and a depletion of male-
biased genes in one species (Fig. 6, Fig. S28, Table S5, Table S6). This suggests that
selection for the enrichment of female-biased genes and the depletion of male-biased genes
on the X is weak, and thus unlikely to generate the large effect sizes we see in the reproductive
tracts. Finally, a reduction in expression of the male X due to a lack of dosage compensation
is expected to be consistent across species, since they share the same X chromosome (see
above). By contrast, sex biased expression independently of dosage is more likely to be
species-specific given the very fast turnover of sex biased genes observed between closely
related species in several taxa (Zhang *et al.*, 2007; Harrison *et al.*, 2015) including *Timema*
(Parker *et al.*, 2019b). To distinguish between these patterns, we tested if sex-biased genes
on the X in the reproductive tracts are underrepresented for genes with species by sex
interactions. We found that genes on the X are underrepresented for genes showing species
by sex interactions in the reproductive tracts ($p = 0.010$), and that this is not the case for the
heads ($p = 0.938$) or legs ($p = 0.795$). This shows that sex-differences in expression on the X
are more consistent between species in the reproductive tracts than in the heads and legs
(also see Fig. S29), and further supports a lack of dosage compensation in the reproductive
tracts rather than a large enrichment of female-biased genes on the X.
457

458
Fig. 4 | Gene expression on the X and autosomes in heads, legs, and reproductive
tracts. A. Average expression levels in males and females on the X and autosomes **B.** Log2
of male to female expression ratio for the X and autosomes. Dashed lines represent a two-
fold reduction in expression in males (as expected if there was no dosage compensation).
Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*,
Tps = *T. poppensis*, and Tpa = *T. podura*.

By examining the expression of the X and autosomes in the different tissues, we can infer the
type of dosage compensation. In the dosage compensated tissues (heads and legs) genes on
the autosomes and X have similar overall expression levels in both males and females, and
the overall expression of the X is similar to that of the autosomes (Fig. 4, Fig. S24, Table S7).
By contrast, in the reproductive tracts, where dosage compensation is lacking, expression of
genes on the X is much lower in males than in females. This difference seems to be driven by
changes in male expression, as X-linked gene expression in females remains similar to the
autosomes (Fig. 4, Fig. S24, Table S7). This supports a mechanism of dosage compensation
by hyper-transcription of the X in males, a mechanism common among other insects (Gu &
Walters, 2017).
476
477 Although we find almost complete dosage compensation, it is possible that its extent could
478 vary along the X chromosome (Mullon *et al.*, 2015). To address this, we examined the extent
479 of dosage along the longest X-linked scaffolds in the genome, and found no evidence of
480 variation along the X chromosome (Fig. S30 - S34, see also Fig. S35 - S46 for expression
481 variation along the autosomes).
482

Fig. 5 | Ratio of male and female gene expression on the X (orange) and autosomes (black) in heads, legs, and reproductive tracts. Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps = *T. poppensis*, and Tpa = *T. podura*.

Sex-biased genes are not enriched on the X chromosome

Sexually antagonistic mutations are expected to fix more easily on the X chromosome than the autosomes. As the evolution of sex-biased gene expression is thought to be primarily driven by sexually antagonistic selection, it is expected that the X chromosome will be a hotspot for sex-biased gene expression (Rice, 1984; Gibson *et al.*, 2002; Ellegren & Parsch, 2007; Innocenti & Morrow, 2010; Griffin *et al.*, 2013). Contrary to this expectation, we find most sex-biased genes on the autosomes (% of sex-biased genes on the X ranges from 8.3 - 14.3, Table S5), and we find little evidence for enrichment of sex-biased genes on the X in the head and leg tissues (Fig. 6, Table S5, Table S6). Note that almost all genes on the X appear to be sex-biased in the reproductive tracts, but this effect is largely due to a lack of dosage compensation in this tissue (see above). In addition, while we do find an enrichment of female-biased genes in the heads of two species (*T. bartmani* and *T. podura*, Fig. 6) and a depletion of male-biased genes in *T. bartmani,* the effect sizes are small and the effect becomes insignificant when considering only sex-biased genes with at least a twofold difference in expression between males and females (Fig. S28). These findings suggest that the selective pressures to accumulate female-biased genes and reduce male-biased genes on the X are weak and/or that there are constraints to the gene content or expression levels on the *Timema* X chromosome.

Fig. 6 | Proportion of female- and male- biased genes on the X and autosomes in reproductive tract, head and leg samples. Note the scale changes between tissue-types. Asterisks indicate the significance level (FDR) of Fisher's exact tests (***<0.001, **<0.01, *<0.05). Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps = *T. poppensis*, and Tpa = *T. podura*.

Discussion

The difference in copy numbers of the X chromosome in males and females is expected to have profound effects on its evolution. In particular, the X chromosome is predicted to evolve at a faster rate due to a combination of hemizygous selection and increased drift (Wright, 1931; Charlesworth *et al.*, 1987; Vicoso & Charlesworth, 2009), to accumulate sexually antagonistic alleles (Rice, 1984; Gibson *et al.*, 2002), and to evolve dosage compensation mechanisms (Disteche, 2012; Mank, 2013; Gu & Walters, 2017; Lenormand *et al.*, 2020). Support for these predictions has been mixed from the taxa examined so far, yet the factors responsible for the variation among different taxa are poorly understood (Bachtrog *et al.*, 2011; Gu & Walters, 2017). In this study, we examine these predictions across five species of *Timema* stick insects. Overall, we find evidence for relaxed selection and complete dosage compensation in somatic tissues, but little evidence for the accumulation of sexually antagonistic alleles (in the form of sex-biased genes) on the X. Patterns of X chromosome evolution were generally consistent across *Timema* species, suggesting that the factors influencing sex chromosome evolution in this group are also largely the same.

Sex chromosome conservation is highly variable between taxa, with extensive turnover between species in some groups, e.g. beetles (Coleoptera) (Blackmon & Demuth, 2014), flies (Diptera) (Vicoso & Bachtrog, 2015), or frogs (Ranidae) (Jeffries *et al.*, 2018), and conservation for over a hundred million years in others, e.g. Eutherian mammals (Lahn & Page, 1999; Cortez *et al.*, 2014; Marshall Graves, 2015), moths and butterflies (Lepidoptera) (Fraïsse *et al.*, 2017), or birds (Shetty *et al.*, 1999; Xu & Zhou, 2020). The factors influencing turnover rate are complex and interacting (Vicoso, 2019), however, one key factor is the level of differentiation between the X and Y chromosomes, with more differentiated chromosomes less likely to turnover (Pokorná & Kratochvíl, 2009; Vicoso, 2019). Our finding that the X chromosome in *Timema* is old (likely conserved for over 120 million years) supports this idea. XX/X0 sex determination systems as found in *Timema* (Schwander & Crespi, 2009) are thought to derive from XX/XY systems with highly differentiated X and Y chromosomes and represent the end point of the gradual loss of gene content from the Y (Bergero & Charlesworth, 2009). As such, XX/X0 systems could be considered the most extreme example of sex chromosome differentiation possible, which may mean that XX/X0 systems are particularly unlikely to turnover (but see (Blackmon & Demuth, 2014)).

Gene sequence evolution on the X chromosome is expected to be faster than that on the autosomes due to a combination of increased drift and hemizygous selection (Charlesworth *et al.*, 1987; Vicoso & Charlesworth, 2009). While this prediction should apply universally, empirical support is mixed (Mank *et al.*, 2010; Meisel & Connallon, 2013; Charlesworth *et al.*, 2018; Pinharanda *et al.*, 2019; Whittle *et al.*, 2020). The cause of this variation is unclear, in particular, because typically only single lineages are examined. In single lineages it is difficult to disentangle the influence of X-linkage from lineage-specific effects such as recent bottlenecks, differences in operational sex ratio, or population size. By examining the influence of X-linkage in multiple *Timema* species covering a span of 30 million years of divergence, we can assess how consistent the effects of X linkage are, allowing for a comprehensive assessment of X chromosome evolution in this genus. Using this approach, we found consistent evidence for a relaxed selection (as measured by dN/dS), known as the "faster X

effect", in all species, and that this effect is primarily driven by reduced purifying selection on
the X. Interestingly, while dN/dS was higher for genes on the X, the overall amount of
divergence was lower. This result is likely driven by a lower net mutation rate on the X relative
to autosomes as a result of a lower mutation rate in the female than male germline (Kirkpatrick
& Hall, 2004; Ellegren, 2007).

The reduction of purifying selection on the X relative to the autosomes is expected to be largest
when the effective population size of the X (Ne_X) is much smaller than that of the autosomes
(Ne_A) (Vicoso & Charlesworth, 2009). The neutral expectation in a population with a balanced
sex ratio is that the ratio of Ne_X to Ne_A will be approximately 0.75 (Wright, 1969; Hartl & Clark,
2006), however, demographic and selective processes can have a large effect on this ratio
(Nunney, 1993; Caballero, 1995; Charlesworth, 2001). In *Timema*, Ne_X / Ne_A values are much
smaller than 0.75 (0.19 to 0.48, estimated from nucleotide diversity), meaning the effective
population size of the X is much smaller than that of the autosomes. Departures from the
expected 0.75 nucleotide diversity ratio are common across animals and are typically thought
to be a consequence of sex-biased demography (Mank *et al.*, 2010). One common cause for
such departures is that in many systems reproductive skew is stronger for males than females,
meaning fewer males contribute to the next generation than females. In *Timema*, females
mate multiply and skew offspring towards particular males (Arbuthnott *et al.*, 2015), suggesting
that in *Timema* reproductive skew is indeed greater in males than females. In male
heterogametic species (such as *Timema*), such a reproductive skew should disproportionately
reduce diversity on the autosomes, and result in an X to autosome nucleotide diversity ratio >
0.75. Here however, we observe the opposite pattern, a reduction in the X to autosome
nucleotide diversity ratio. As such, the reduced diversity on the X we observe is unlikely to be
due to sex-specific variation in reproductive success. There are many potential alternative
demographic and selective explanations for this finding, however the most likely (discussed
below) are that the X has a lower recombination rate than the autosomes, and/or that *Timema*
undergo frequent population bottlenecks. Since we show that crossing over occurs in both
sexes, the effective recombination rate for the X is likely lower than for the autosomes (as
recombination between different copies of the X can only occur in females). Low
recombination rates intensify the consequences of selective sweeps and background
selection on genetic diversity and could thus contribute to the reduced genetic diversity on the
X we observe (Betancourt *et al.*, 2004; Charlesworth, 2012, 2013; Wilson Sayres, 2018).
Population bottlenecks could also contribute to the reduced genetic diversity on the X.
Although bottlenecks will reduce genetic diversity for both the X and the autosomes, it is
expected that they will disproportionately reduce genetic diversity on the X (Pool & Nielsen,
2007). This factor may be particularly relevant as *Timema* live in fire-prone habitats where
fires may lead to frequent population bottlenecks.

Independently of the mechanisms responsible for the strongly reduced effective population
size of the X in *Timema*, it supports our interpretation that the faster-X effect is driven by less
effective purifying selection. Most previous studies of the faster-X effect have focused on
species with Ne_X / Ne_A values ≥ 0.75 (Mank *et al.*, 2010), making direct comparisons with our
study difficult. The exceptions to this are studies on birds and *Heliconius* butterflies which have
values of Ne_Z / Ne_A in a similar range as *Timema*. Note that both these taxa have
heterogametic females (ZW), where low Ne_Z / Ne_A ratios are believed to stem from strong
reproductive skew among males (Vicoso & Charlesworth, 2009; Mank *et al.*, 2010). In birds,

a faster-Z (faster-X) effect is commonly observed and appears to be driven primarily by less
effective purifying selection (Mank *et al.*, 2010). In contrast, in *Heliconius* evidence for a faster-
Z effect is weaker and is thought to be driven by increased levels of adaptive evolution on the
Z, with no evidence for reduced purifying selection on the Z (Pinharanda *et al.*, 2019). The
cause of this difference is unclear. However, it has been suggested that it may be due to the
overall higher effective population size in *Heliconius* (Ne ≈ 2,000,000 (Keightley *et al.*, 2015))
compared to birds (Ne = 200,000 - 600,000 (Primmer *et al.*, 2002; Axelsson *et al.*, 2004;
Jennings & Edwards, 2005; Backström *et al.*, 2008)), meaning that selection can be efficient
in *Heliconius* even with the relatively reduced effective population size of the sex chromosome
(Mank *et al.*, 2010; Pinharanda *et al.*, 2019). This interpretation conflicts with our findings in
Timema, as the (autosomal) effective population size varies greatly between species (from
~150,000 in *T. poppensis* to ~2,000,000 in *T. podura*), yet we observe a similar reduction in
purifying selection on the X in all species. The reason for this variation between study systems
is thus unclear, and highlights the need for future studies across diverse taxa. It should be
noted that the values of Ne we estimate for *Timema* should be considered as rough
approximations as our calculations of Ne assume the mutation rate in *Timema* is similar to
that in *D. melanogaster* and *H. melpomene*. Since these are distantly related species the
mutation rate in *Timema* could differ markedly, which would mean our estimates of Ne would
be systematically under- or over- estimated. While this complicates direct comparisons with
other taxa, it is unlikely to influence our comparisons among *Timema* species for which only
the relative differences in Ne are important.
629
The X chromosome has long been predicted to be a hotspot for sexually antagonistic variation
(Rice, 1984; Gibson *et al.*, 2002). The evolution of sex-biased gene expression is thought to
be driven by sexually antagonistic selection, and thus sex-biased gene expression should be
overrepresented on the X chromosome ((Ellegren & Parsch, 2007; Innocenti & Morrow, 2010;
Griffin *et al.*, 2013) but see (Hitchcock & Gardner, 2020; Ruzicka & Connallon, 2020)). In
Timema, we find very little support for this prediction with only a small enrichment of sex-
biased genes on the X in one of our five species. In combination with studies in *Drosophila
melanogaster* (Ruzicka *et al.*, 2019), *Callosobruchus maculatus* (Sayadi *et al.*, 2019), and
Ischnura elegans (Chauhan *et al.*, 2021) which also found little or no enrichment of sexually
antagonistic alleles or sex-biased gene expression on the sex chromosomes, our study
suggests that the advantage of accumulating sexually antagonistic alleles on the X be may be
smaller than often assumed. The reasons for this are unclear. However, it is likely that the
advantage X-linkage gives to sexually antagonistic alleles is balanced by other forces such as
epistatic interactions (Arnqvist *et al.*, 2014) or sex-specific dominance (Fry, 2010). Both forces
favour the accumulation of sexually antagonistic alleles on the autosomes. Additionally, it
should be noted that our approach to use sex-biased genes as a proxy for sexually
antagonistic variation may be underpowered as it will miss any genes that are evolving under
sexual conflict but have not (yet) evolved sex-biased gene expression. Despite this, several
studies do show the expected enrichment of sexually antagonistic alleles or sex-biased gene
expression on the X: in *Tribolium castaneum* (Whittle *et al.*, 2020), Diptera (Innocenti &
Morrow, 2010; Vicoso & Bachtrog, 2015), Hemiptera (Pal & Vicoso, 2015), and nematodes
(Albritton *et al.*, 2014). A key challenge for future work will thus be to integrate studies that
quantify multiple factors that influence the accumulation of sexually antagonistic alleles (e.g.
epistatic interactions, sex-specific dominance, reproductive skew, etc.) to understand how the
variation between studies and taxa is produced.

In many species dosage compensation is thought to be important for ameliorating the costs of misexpression of genes on the X chromosome (Marín *et al.*, 2000). Although common, there is a great deal of variation in the extent to which genes are dosage compensated. Here we find almost complete dosage compensation in the somatic tissues of all five *Timema* species, as reported for other species with X0 systems (e.g. Nematodes (Meyer, 2000) and crickets (Rayner *et al.*, 2021)). Dosage compensation is expected for X0 systems as they are thought to arise from XY systems where the Y chromosome has degraded to the point it can be lost without a large decrease in fitness. By this stage, most genes on the X should already be haplo-sufficient in males. In addition, the evolution of dosage compensation could itself hasten the loss of the Y chromosome, as genes that have functional copies on both the X and the Y will be misexpressed if chromosome-wide dosage compensation evolves (Vicoso & Bachtrog, 2009; Lenormand & Roze, 2021).

In contrast to the somatic tissues, male reproductive tracts displayed a lack of dosage compensation. Reduced expression of X chromosome in male reproductive tissue has been observed in a number of species including mammals (Khil *et al.*, 2004; Disteche, 2012; Sangrithi & Turner, 2018), *C. elegans* (Kelly *et al.*, 2002; Pirrotta, 2002) and insects such as *D. melanogaster* (Oliver, 2002; Meiklejohn *et al.*, 2011; Mahadevaraju *et al.*, 2021) or *Teleogryllus oceanicus* (Rayner *et al.*, 2021). Whether reduced X expression in male reproductive tracts generally stems from a lack of dosage compensation is, however, not clear. Indeed, reduced X expression can be caused by several non-mutually exclusive mechanisms, including the accumulation of female-biased genes on the X (Mank, 2009), the movement of male-beneficial genes to the autosomes (Vibranovski *et al.*, 2009), or the inactivation of X chromosomes in the germ line (Lee, 2005; Vibranovski, 2014). In *Timema*, we suggest an absence of dosage compensation is the most likely explanation as expression of X-linked genes in the male reproductive tracts show a major peak of genes with expression approximately half of that observed in females, similar to that observed in species that lack dosage compensation (Mank & Ellegren, 2009; Vicoso *et al.*, 2013). It should be noted, however, that the expression reduction we see on the male X in *Timema* is actually slightly less than the half we would expect from a lack of dosage compensation alone. This suggests that other factors, described above, may also have an influence. Disentangling the contribution of each of these factors will require further work. Our work, however, clearly shows that expression in reproductive tissues behaves differently than in somatic tissues, highlighting the importance of studying these tissues separately particularly when assessing the extent of dosage compensation (Gu & Walters, 2017).

Dosage compensation in *Timema* somatic tissues appears to be achieved by the upregulation of the X in males (type I, (Gu & Walters, 2017)) as indicated by similar expression levels of genes on the X and the autosomes in both sexes. This interpretation assumes that the ancestral state of X chromosome expression (before it was a sex chromosome) was similar to the other autosomes. This seems likely as expression across all *Timema* autosomes is similar. However, assessing the ancestral state of X expression is largely impossible as the *Timema* X chromosome appears to be very old (~120 million years). As a consequence, there are likely no species available for comparison with conserved genome organisation but with X chromosomes that are homologous to different *Timema* autosomes. Similar to *Timema*, upregulation of the X chromosome in males also appears to be the mechanism for dosage compensation in all other XX/XY or XX/X0 insect systems yet studied, even when dosage

compensation is incomplete (Gu & Walters, 2017). With our study, this amounts to information
from seven different insect orders (Odonata (Chauhan *et al.*, 2021), Phasmatodea (this study),
Hemiptera (Pal & Vicoso, 2015; Richard *et al.*, 2017), Orthoptera (Rayner *et al.*, 2021),
Strepsiptera (Mahajan & Bachtrog, 2015), Coleoptera (Prince *et al.*, 2010; Mahajan &
Bachtrog, 2015), and Diptera (Bone & Kuroda, 1996; Marín *et al.*, 1996; Deng *et al.*, 2011;
Nozawa *et al.*, 2014; Jiang *et al.*, 2015; Vicoso & Bachtrog, 2015; Rose *et al.*, 2016))
suggesting that this mechanism may be universal for insect species with heterogametic males.
710

Conclusions

How consistent the consequences of sex linkage are for gene sequence and expression
evolution across taxa remains an open question. Here we examine several key aspects of sex
chromosome evolution in a previously neglected group, phasmids. Overall, we find evidence
for several predicted consequences, including complete dosage compensation of the X (in
somatic tissues) and a faster rate of evolution of X-linked than autosomal genes. By contrast,
we find little evidence that sex linkage facilitates the accumulation of sexually antagonistic
alleles. While our results were consistent across different *Timema* species, they also show
key differences from studies in other taxa, highlighting the importance of studying sex
chromosome evolution across a diverse set of species to distinguish general patterns caused
by sex linkage from species-specific idiosyncrasies.

Data and code availability

Raw sequence reads have been deposited in NCBI's sequence read archive under the bioproject: PRJNA725673 (Table S8). Scripts for the analyses in this paper are available at: https://github.com/DarrenJParker/Timema_sex_chr_evol_code and will be archived at Zenodo upon acceptance. Data was processed to generate plots and statistics using R v4.0.3 and Python v.3.7.3 unless otherwise stated.

Acknowledgements

We would like to thank Chloe Larose, Kirsten Jalvingh, and Bart Zijlstra for their assistance in the field. We thank Sébastien Moretti for his assistance with running Godon. We also thank Matthew Hahn for useful comments. This project was supported by several grants from the Swiss Science Foundation: PP00P3_170627 and 31003A_182495 to TS as well as CRSII3_160723 to TS, MRR and Nicolas Galtier. We would like to thank the former Vital-IT platform of the SIB Swiss Institute of Bioinformatics (SIB) and the DCSR for maintaining the computer infrastructure at the University of Lausanne.

References

Albritton, S.E., Kranz, A.-L., Rao, P., Kramer, M., Dieterich, C. & Ercan, S. 2014. Sex-biased gene expression and evolution of the x chromosome in nematodes. *Genetics* **197**: 865–883.

Anders, S., Pyl, P.T. & Huber, W. 2015. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**: 166–169.

Arbuthnott, D., Crespi, B.J. & Schwander, T. 2015. Female stick insects mate multiply to find compatible mates. *Am. Nat.* **186**: 519–530.

Arnqvist, G., Vellnow, N. & Rowe, L. 2014. The effect of epistasis on sexually antagonistic genetic variation. *Proceedings of the Royal Society B: Biological Sciences* **281**: 20140489.

Axelsson, E., Smith, N.G.C., Sundström, H., Berlin, S. & Ellegren, H. 2004. Male-biased mutation rate and divergence in autosomal, Z-linked and W-linked introns of chicken and turkey. *Mol. Biol. Evol.* **21**: 1538–1547. academic.oup.com.

Bachtrog, D., Kirkpatrick, M., Mank, J.E., McDaniel, S.F., Pires, J.C., Rice, W., *et al.* 2011. Are all sex chromosomes created equal? *Trends Genet.* **27**: 350–357. Elsevier.

Backström, N., Fagerberg, S. & Ellegren, H. 2008. Genomics of natural bird populations: a gene-based set of reference markers evenly spread across the avian genome. *Mol. Ecol.* **17**: 964–980. Wiley.

Benjamini, Y. & Hochberg, Y. 1995. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. R. Stat. Soc. Series B Stat. Methodol.* **57**: 289–300. [Royal Statistical Society, Wiley].

Bergero, R. & Charlesworth, D. 2009. The evolution of restricted recombination in sex chromosomes. *Trends Ecol. Evol.* **24**: 94–102.

Betancourt, A.J., Kim, Y. & Orr, H.A. 2004. A pseudohitchhiking model of X vs. autosomal diversity. *Genetics* **168**: 2261–2269.

Birchler, J.A. 2016. Parallel universes for models of X chromosome dosage compensation in *Drosophila*: a review. *Cytogenet. Genome Res.* **148**: 52–67.

Birchler, J.A. & Veitia, R.A. 2012. Gene balance hypothesis: connecting issues of dosage sensitivity across biological disciplines. *Proc. Natl. Acad. Sci. U. S. A.* **109**: 14746–14753.

Blackmon, H. & Demuth, J.P. 2014. Estimating tempo and mode of Y chromosome turnover: explaining Y chromosome loss with the fragile Y hypothesis. *Genetics* **197**: 561–572.

Bolger, A.M., Lohse, M. & Usadel, B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.

Bone, J.R. & Kuroda, M.I. 1996. Dosage compensation regulatory proteins and the evolution of sex chromosomes in *Drosophila*. *Genetics* **144**: 705–713.

Bradler, S. & Buckley, T.R. 2018. Biodiversity of Phasmatodea. In: *Insect Biodiversity*, pp. 281–313. John Wiley & Sons, Ltd, Chichester, UK.

Caballero, A. 1995. On the effective size of populations with separate sexes, with particular
reference to sex-linked genes. *Genetics* **139**: 1007–1011.

Charif, D. & Lobry, J.R. 2007. SeqinR 1.0-2: a contributed package to the R project for
statistical computing devoted to biological sequences retrieval and analysis. Springer
Verlag, New York.

Charlesworth, B. 2013. Background selection 20 years on: the Wilhelmine E. Key 2012
invitational lecture. *J. Hered.* **104**: 161–171.

Charlesworth, B. 1978. Model for evolution of Y chromosomes and dosage compensation.
Proc. Natl. Acad. Sci. U. S. A. **75**: 5618–5622. National Acad Sciences.

Charlesworth, B. 2001. The effect of life-history and mode of inheritance on neutral genetic
variability. *Genet. Res.* **77**: 153–166.

Charlesworth, B. 1996. The evolution of chromosomal sex determination and dosage
compensation. *Curr. Biol.* **6**: 149–162.

Charlesworth, B. 2012. The role of background selection in shaping patterns of molecular
evolution and variation: evidence from variability on the *Drosophila* X chromosome.
Genetics **191**: 233–246.

Charlesworth, B., Campos, J.L. & Jackson, B.C. 2018. Faster-X evolution: Theory and
evidence from *Drosophila*. *Mol. Ecol.* **27**: 3753–3771.

Charlesworth, B., Coyne, J.A. & Barton, N.H. 1987. The relative rates of evolution of sex
chromosomes and autosomes. *Am. Nat.* **130**: 113–146. The University of Chicago
Press.

Chauhan, P., Swaegers, J., Guillen, R.A.S., Svensson, E.I., Wellenreuther, M. & Hansson,
B. 2021. Genome assembly, sex-biased gene expression and dosage compensation in
the damselfly *Ischnura elegans*. *Genomics* **113**: 1828–1837.

Conrad, T. & Akhtar, A. 2012. Dosage compensation in *Drosophila melanogaster*: epigenetic
fine-tuning of chromosome-wide transcription. *Nat. Rev. Genet.* **13**: 123–134.

Cortez, D., Marin, R., Toledo-Flores, D., Froidevaux, L., Liechti, A., Waters, P.D., *et al.* 2014.
Origins and functional evolution of Y chromosomes across mammals. *Nature* **508**: 488–
493. Nature Publishing Group.

Coyne, J.A. & Orr, H.A. 2004. *Speciation*. Sinauer Associates, Sunderland, USA.

Deng, X., Hiatt, J.B., Nguyen, D.K., Ercan, S., Sturgill, D., Hillier, L.W., *et al.* 2011. Evidence
for compensatory upregulation of expressed X-linked genes in mammals,
Caenorhabditis elegans and *Drosophila melanogaster*. *Nat. Genet.* **43**: 1179–1185.

Disteche, C.M. 2012. Dosage compensation of the sex chromosomes. *Annu. Rev. Genet.*
46: 537–560.

Ellegren, H. 2007. Characteristics, causes and evolutionary consequences of male-biased
mutation. *Proc. Biol. Sci.* **274**: 1–10.

Ellegren, H. & Parsch, J. 2007. The evolution of sex-biased genes and sex-biased gene
expression. *Nat. Rev. Genet.* **8**: 689–698.

Fraïsse, C., Picard, M.A.L. & Vicoso, B. 2017. The deep conservation of the Lepidoptera Z

chromosome suggests a non-canonical origin of the W. *Nat. Commun.* **8**: 1–9. Nature
Publishing Group.

Fry, J.D. 2010. The genomic location of sexually antagonistic variation: some cautionary
comments. *Evolution* **64**: 1510–1516.

Gibson, J.R., Chippindale, A.K. & Rice, W.R. 2002. The X chromosome is a hot spot for
sexually antagonistic fitness variation. *Proc. Biol. Sci.* **269**: 499–505.

Griffin, R.M., Dean, R., Grace, J.L., Rydén, P. & Friberg, U. 2013. The shared genome is a
pervasive constraint on the evolution of sex-biased gene expression. *Mol. Biol. Evol.* **30**:
2168–2176.

Gu, L. & Walters, J.R. 2017. Evolution of sex chromosome dosage compensation in animals:
a beautiful theory, undermined by facts and bedeviled by details. *Genome Biol. Evol.* **9**:
2461–2476.

Harrison, P.W., Wright, A.E., Zimmer, F., Dean, R., Montgomery, S.H., Pointer, M.A., *et al.*
2015. Sexual selection drives evolution and rapid turnover of male gene expression.
Proc. Natl. Acad. Sci. U. S. A. **112**: 4393–4398.

Hartl, D.L. & Clark, A.G. 2006. *Principles of Population Genetics*, 4th ed. Sinauer
Associates.

Hitchcock, T.J. & Gardner, A. 2020. A gene's-eye view of sexual antagonism. *Proc. Biol. Sci.*
287: 20201633.

Innocenti, P. & Morrow, E.H. 2010. The sexually antagonistic genes of *Drosophila*
melanogaster. *PLoS Biol.* **8**: e1000335.

Jaquiéry, J., Simon, J.-C., Robin, S., Richard, G., Peccoud, J., Boulain, H., *et al.* 2021.
Masculinization of the X-chromosome in aphid soma and gonads. *bioRxiv*
2021.08.13.453080.

Jaron, K.S., Parker, D.J., Anselmetti, Y., Van, P.T., Bast, J., Dumas, Z., et al. 2022.
Convergent consequences of parthenogenesis on stick insect genomes. *Science*
Advances 8: eabg3842.

Jeffries, D.L., Lavanchy, G., Sermier, R., Sredl, M.J., Miura, I., Borzée, A., *et al.* 2018. A
rapid rate of sex-chromosome turnover and non-random transitions in true frogs. *Nat.*
Commun. **9**: 4088.

Jennings, W.B. & Edwards, S.V. 2005. Speciational history of Australian grass finches
(Poephila) inferred from thirty gene trees. *Evolution* **59**: 2033–2047. Wiley.

Jiang, X., Biedler, J.K., Qi, Y., Hall, A.B. & Tu, Z. 2015. Complete dosage compensation in
Anopheles stephensi and the evolution of sex-biased genes in mosquitoes. *Genome*
Biol. Evol. **7**: 1914–1924.

Keightley, P.D., Ness, R.W., Halligan, D.L. & Haddrill, P.R. 2014. Estimation of the
spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib
family. *Genetics* **196**: 313–320.

Keightley, P.D., Pinharanda, A., Ness, R.W., Simpson, F., Dasmahapatra, K.K., Mallet, J., *et*
al. 2015. Estimation of the spontaneous mutation rate in *Heliconius melpomene*. *Mol.*
Biol. Evol. **32**: 239–243.

Kelly, W.G., Schaner, C.E., Dernburg, A.F., Lee, M.-H., Kim, S.K., Villeneuve, A.M., *et al.*
2002. X-chromosome silencing in the germline of *C. elegans*. *Development* **129**: 479–
492. journals.biologists.com.

Khil, P.P., Smirnova, N.A., Romanienko, P.J. & Camerini-Otero, R.D. 2004. The mouse X
chromosome is enriched for sex-biased genes not subject to selection by meiotic sex
chromosome inactivation. *Nat. Genet.* **36**: 642–646.

Kirkpatrick, M. & Hall, D.W. 2004. Male-biased mutation, sex linkage, and the rate of
adaptive evolution. *Evolution* **58**: 437–440. Wiley.

Korneliussen, T.S., Albrechtsen, A. & Nielsen, R. 2014. ANGSD: Analysis of next generation
sequencing data. *BMC Bioinformatics* **15**: 356. Springer.

Lahn, B.T. & Page, D.C. 1999. Four evolutionary strata on the human X chromosome.
Science **286**: 964–967.

Lee, J.T. 2005. Sex chromosome inactivation: the importance of pairing. *Curr. Biol.* **15**:
R249–52.

Lenormand, T., Fyon, F., Sun, E. & Roze, D. 2020. Sex Chromosome Degeneration by
Regulatory Evolution. *Curr. Biol.* **30**: 3001–3006.e5.

Lenormand, T. & Roze, D. 2021. Y recombination arrest and degeneration in the absence of
sexual dimorphism. *bioRxiv* 2021.05.18.444606.

Li, H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. *arXiv* 1303.3997. arxiv.org.

Lucchesi, J.C. 1998. Dosage compensation in flies and worms: the ups and downs of X-
chromosome regulation. *Curr. Opin. Genet. Dev.* **8**: 179–184. Elsevier.

Mahadevaraju, S., Fear, J.M., Akeju, M., Galletta, B.J., Pinheiro, M.M.L.S., Avelino, C.C., *et*
al. 2021. Dynamic sex chromosome expression in *Drosophila* male germ cells. *Nat.*
Commun. **12**: 892.

Mahajan, S. & Bachtrog, D. 2015. Partial dosage compensation in Strepsiptera, a sister
group of beetles. *Genome Biol. Evol.* **7**: 591–600.

Mank, J.E. 2013. Sex chromosome dosage compensation: definitely not for everyone.
Trends Genet. **29**: 677–683. Elsevier Ltd.

Mank, J.E. 2009. The W, X, Y and Z of sex-chromosome dosage compensation. *Trends*
Genet. **25**: 226–233. Elsevier BV.

Mank, J.E. & Ellegren, H. 2009. All dosage compensation is local: gene-by-gene regulation
of sex-biased expression on the chicken Z chromosome. *Heredity* **102**: 312–320.

Mank, J.E., Hosken, D.J. & Wedell, N. 2014. Conflict on the sex chromosomes: cause,
effect, and complexity. *Cold Spring Harb. Perspect. Biol.* **6**: a017715.

Mank, J.E., Vicoso, B., Berlin, S. & Charlesworth, B. 2010. Effective population size and the
Faster-X effect: empirical results and their interpretation. *Evolution* **64**: 663–674. Wiley.

Marçais, G., Delcher, A.L., Phillippy, A.M., Coston, R., Salzberg, S.L. & Zimin, A. 2018.
MUMmer4: A fast and versatile genome alignment system. *PLoS Comput. Biol.* **14**:
e1005944.

Marescalchi, O., Masetti, M. & Scali, V. 1986. C-banding karyotype and male meiosis of
Acrophylla wuelfingi (redtenbacher) (Phasmatodea: Phasmatidae). *Aust. J. Entomol.* **25**:
223–227. Wiley.

Marín, I., Franke, A., Bashaw, G.J. & Baker, B.S. 1996. The dosage compensation system of
Drosophila is co-opted by newly evolved X chromosomes. *Nature* **383**: 160–163. Nature
Publishing Group.

Marín, I., Siegal, M.L. & Baker, B.S. 2000. The evolution of dosage-compensation
mechanisms. *Bioessays* **22**: 1106–1114. Wiley.

Marshall Graves, J.A. 2015. Evolution of vertebrate sex chromosomes and dosage
compensation. *Nat. Rev. Genet.* **17**: 33–46. Nature Publishing Group.

Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet J* **17**: 10–12.

Meiklejohn, C.D., Landeen, E.L., Cook, J.M., Kingan, S.B. & Presgraves, D.C. 2011. Sex
chromosome-specific regulation in the *Drosophila* male germline but little evidence for
chromosomal dosage compensation or meiotic inactivation. *PLoS Biol.* **9**: e1001126.

Meisel, R.P. & Connallon, T. 2013. The faster-X effect: integrating theory and data. *Trends*
Genet. **29**: 537–544.

Meyer, B.J. 2000. Sex in the worm: counting and compensating X-chromosome dose.
Trends Genet. **16**: 247–253.

Montgomery, S.H. & Mank, J.E. 2016. Inferring regulatory change from gene expression: the
confounding effects of tissue scaling. *Mol. Ecol.* **25**: 5114–5128.

Mullon, C., Pomiankowski, A. & Reuter, M. 2012. The effects of selection and genetic drift on
the genomic distribution of sexually antagonistic alleles. *Evolution* **66**: 3743–3753.

Mullon, C., Wright, A.E., Reuter, M., Pomiankowski, A. & Mank, J.E. 2015. Evolution of
dosage compensation under sexual selection differs between X and Z chromosomes.
Nat. Commun. **6**: 7720. Nature Publishing Group.

Nosil, P., Villoutreix, R., de Carvalho, C.F., Farkas, T.E., Soria-Carrasco, V., Feder, J.L., *et*
al. 2018. Natural selection and the predictability of evolution in *Timema* stick insects.
Science **359**: 765–770. American Association for the Advancement of Science.

Nozawa, M., Fukuda, N., Ikeo, K. & Gojobori, T. 2014. Tissue- and stage-dependent dosage
compensation on the neo-X chromosome in *Drosophila pseudoobscura. Mol. Biol. Evol.*
31: 614–624.

Nunney, L. 1993. The influence of mating system and overlapping generations on effective
population size. *Evolution* **47**: 1329–1341. [Society for the Study of Evolution, Wiley].

Ohno, S. 1967. *Sex chromosomes and sex-linked genes*. Springer Verlag, Berlin.

Oliver, B. 2002. Genetic control of germline sexual dimorphism in *Drosophila. Int. Rev. Cytol.*
219: 1–60.

Pal, A. & Vicoso, B. 2015. The X Chromosome of Hemipteran insects: conservation, dosage
compensation and sex-biased expression. *Genome Biol. Evol.* **7**: 3259–3268.

Palmer, D.H., Rogers, T.F., Dean, R. & Wright, A.E. 2019. How to identify sex chromosomes

941 and their turnover. *Mol. Ecol.* **28**: 4709–4724.

942 Parker, D.J., Bast, J., Jalvingh, K., Dumas, Z., Robinson-Rechavi, M. & Schwander, T.
943 2019a. Repeated evolution of asexuality involves convergent gene expression changes.
944 *Mol. Biol. Evol.* **36** : 350–364.

945 Parker, D.J., Bast, J., Jalvingh, K., Dumas, Z., Robinson-Rechavi, M. & Schwander, T.
946 2019b. Sex-biased gene expression is repeatedly masculinized in asexual females. *Nat.*
947 *Commun.* **10**: 4638.

948 Parkhurst, S.M. & Meneely, P.M. 1994. Sex determination and dosage compensation:
949 lessons from flies and worms. *Science* **264**: 924–932. science.sciencemag.org.

950 Parsch, J. & Ellegren, H. 2013. The evolutionary causes and consequences of sex-biased
951 gene expression. *Nat. Rev. Genet.* **14**: 83–87.

952 Payseur, B.A., Presgraves, D.C. & Filatov, D.A. 2018. Introduction: Sex chromosomes and
953 speciation. *Mol. Ecol.* **27**: 3745–3748. Wiley.

954 Pinharanda, A., Rousselle, M., Martin, S.H., Hanly, J.J., Davey, J.W., Kumar, S., *et al.* 2019.
955 Sexually dimorphic gene expression and transcriptome evolution provide mixed
956 evidence for a fast-Z effect in *Heliconius*. *J. Evol. Biol.* **32**: 194–204.

957 Pirrotta, V. 2002. Silence in the germ. *Cell* **110**: 661–664.

958 Pokorná, M. & Kratochvíl, L. 2009. Phylogeny of sex-determining mechanisms in squamate
959 reptiles: are sex chromosomes an evolutionary trap? *Zool. J. Linn. Soc.* **156**: 168–183.
960 Oxford Academic.

961 Pool, J.E. & Nielsen, R. 2007. Population size changes reshape genomic patterns of
962 diversity. *Evolution* **61**: 3001–3006.

963 Primmer, C.R., Borge, T., Lindell, J. & Saetre, G.-P. 2002. Single-nucleotide polymorphism
964 characterization in species with limited available sequence information: high nucleotide
965 diversity revealed in the avian genome. *Mol. Ecol.* **11**: 603–612. Wiley.

966 Prince, E.G., Kirkland, D. & Demuth, J.P. 2010. Hyperexpression of the X chromosome in
967 both sexes results in extensive female bias of X-linked genes in the flour beetle.
968 *Genome Biol. Evol.* **2**: 336–346.

969 Quinlan, A.R. & Hall, I.M. 2010. BEDTools: a flexible suite of utilities for comparing genomic
970 features. *Bioinformatics* **26**: 841–842. academic.oup.com.

971 Rayner, J.G., Hitchcock, T.J. & Bailey, N.W. 2021. Variable dosage compensation is
972 associated with female consequences of an X-linked, male-beneficial mutation. *Proc.*
973 *Biol. Sci.* **288**: 20210355.

974 R Core Team. 2017. R: A Language and Environment for Statistical Computing. R
975 Foundation for Statistical Computing, Vienna, Austria.

976 Rice, W.R. 1984. Sex chromosomes and the evolution of sexual dimorphism. *Evolution* **38**:
977 735–742.

978 Richard, G., Legeai, F., Prunier-Leterme, N., Bretaudeau, A., Tagu, D., Jaquiéry, J., *et al.*
979 2017. Dosage compensation and sex-specific epigenetic landscape of the X
980 chromosome in the pea aphid. *Epigenetics Chromatin* **10**: 30.

Riesch, R., Muschick, M., Lindtke, D., Villoutreix, R., Comeault, A.A., Farkas, T.E., *et al.*
2017. Transitions between phases of genomic differentiation during stick-insect
speciation. *Nature Ecology & Evolution* **1**: 0082. Nature Publishing Group.

Robinson, M.D., McCarthy, D.J. & Smyth, G.K. 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–
140.

Rose, G., Krzywinska, E., Kim, J., Revuelta, L., Ferretti, L. & Krzywinski, J. 2016. Dosage
compensation in the African Malaria Mosquito *Anopheles gambiae*. *Genome Biol. Evol.*
8: 411–425.

Ruzicka, F. & Connallon, T. 2020. Is the X chromosome a hot spot for sexually antagonistic
polymorphisms? Biases in current empirical tests of classical theory. *Proceedings of the*
Royal Society B: Biological Sciences **287**: 20201869. Royal Society.

Ruzicka, F., Hill, M.S., Pennell, T.M., Flis, I., Ingleby, F.C., Mott, R., *et al.* 2019. Genome-
wide sexually antagonistic variants reveal long-standing constraints on sexual
dimorphism in fruit flies. *PLoS Biol.* **17**: e3000244.

Sangrithi, M.N. & Turner, J.M.A. 2018. Mammalian X chromosome dosage compensation:
perspectives from the germ line. *Bioessays* **40**: e1800024. Wiley Online Library.

Sayadi, A., Martinez Barrio, A., Immonen, E., Dainat, J., Berger, D., Tellgren-Roth, C., *et al.*
2019. The genomic footprint of sexual conflict. *Nat Ecol Evol* **3**: 1725–1730.

Schwander, T. & Crespi, B.J. 2009. Multiple direct transitions from sexual reproduction to
apomictic parthenogenesis in *Timema* stick insects. *Evolution* **63**: 84–103.

Shetty, S., Griffin, D.K. & Graves, J.A. 1999. Comparative painting reveals strong
chromosome homology over 80 million years of bird evolution. *Chromosome Res.* **7**:
289–295.

Simon, S., Letsch, H., Bank, S., Buckley, T.R., Donath, A., Liu, S., *et al.* 2019. Old World
and New World Phasmatodea: phylogenomics resolve the evolutionary history of stick
and leaf Insects. *Frontiers in Ecology and Evolution* **7**: 345.

Straub, T. & Becker, P.B. 2011. Transcription modulation chromosome-wide: universal
features and principles of dosage compensation in worms and flies. *Curr. Opin. Genet.*
Dev. **21**: 147–153. Elsevier.

Van der Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., Del Angel, G., Levy-Moonshine,
A., *et al.* 2013. From FastQ data to high confidence variant calls: the Genome Analysis
Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* **43**: 11.10.1–11.10.33. Wiley
Online Library.

Vibranovski, M.D. 2014. Meiotic sex chromosome inactivation in *Drosophila*. *J Genomics* **2**:
104–117. ncbi.nlm.nih.gov.

Vibranovski, M.D., Zhang, Y. & Long, M. 2009. General gene movement off the X
chromosome in the *Drosophila* genus. *Genome Res.* **19**: 897–903.

Vicoso, B. 2019. Molecular and evolutionary dynamics of animal sex-chromosome turnover.
Nat Ecol Evol **3**: 1632–1641. nature.com.

Vicoso, B. & Bachtrog, D. 2015. Numerous transitions of sex chromosomes in Diptera. *PLoS*
Biol. **13**: e1002078.

Vicoso, B. & Bachtrog, D. 2009. Progress and prospects toward our understanding of the
evolution of dosage compensation. *Chromosome Res.* **17**: 585–602.

Vicoso, B. & Charlesworth, B. 2009. Effective population size and the faster-X effect: an
extended model. *Evolution* **63**: 2413–2426.

Vicoso, B., Emerson, J.J., Zektser, Y., Mahajan, S. & Bachtrog, D. 2013. Comparative sex
chromosome genomics in snakes: differentiation, evolutionary strata, and lack of global
dosage compensation. *PLoS Biol.* **11**: e1001643. Public Library of Science.

Wang, M., Zhao, Y. & Zhang, B. 2015. Efficient test and visualization of multi-set
intersections. *Sci. Rep.* **5**: 16923.

White, M.J.D. 1976. *Blattodea, Mantodea, Isoptera, Grylloblattodea, Pasmatodea,*
Dermaptera and Embioptera. Animal Cytogenetics Vol. 3: Insecta 2. Gebruder
Borntraeger, Berlin.

Whittle, C.A., Kulkarni, A. & Extavour, C.G. 2020. Absence of a Faster-X effect in beetles
(Tribolium, Coleoptera). *G3: Genes, Genomes, Genetics* **10**: 1125–1136. G3: Genes,
Genomes, Genetics.

Wilkinson, G.S., Breden, F., Mank, J.E., Ritchie, M.G., Higginson, A.D., Radwan, J., *et al.*
2015. The locus of sexual selection: moving sexual selection studies into the post-
genomics era. *J. Evol. Biol.* **28**: 739–755.

Wilson Sayres, M.A. 2018. Genetic diversity on the sex chromosomes. *Genome Biol. Evol.*
10: 1064–1078.

Wright, A.E., Dean, R., Zimmer, F. & Mank, J.E. 2016. How to make a sex chromosome.
Nat. Commun. **7**: 12087. Nature Publishing Group.

Wright, S. 1969. *Evolution and the Genetics of Populations, Volume 2: Theory of Gene*
Frequencies. University of Chicago Press, Chicago, USA.

Wright, S. 1931. Evolution in Mendelian Populations. *Genetics* **16**: 97–159. Genetics.

Xu, L. & Zhou, Q. 2020. The female-specific W chromosomes of birds have conserved gene
contents but are not feminized. *Genes* **11**: 1126.

Yang, Z., Nielsen, R., Goldman, N. & Pedersen, A.M. 2000. Codon-substitution models for
heterogeneous selection pressure at amino acid sites. *Genetics* **155**: 431–449.

Zhang, Y., Sturgill, D., Parisi, M., Kumar, S. & Oliver, B. 2007. Constraint and turnover in
sex-biased gene expression in the genus *Drosophila*. *Nature* **450**: 233–237.