# X chromosomes show a faster evolutionary rate and complete somatic dosage compensation across *Timema* stick insect species

Darren J. Parker[1,2], Kamil S. Jaron[1,2,3], Zoé Dumas[1], Marc Robinson-Rechavi[1,2], and Tanja Schwander[1]

1. Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland
2. Swiss Institute of Bioinformatics, Lausanne, Switzerland
3. Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3FL

Correspondence should be addressed to DarrenJames.Parker@unil.ch or Tanja.Schwander@unil.ch

# Abstract

Sex chromosomes have evolved repeatedly across the tree of life. As they are present in different copy numbers in males and females, they are expected to experience different selection pressures than the autosomes, with consequences including a faster rate of evolution, increased accumulation of sexually antagonistic alleles, and the evolution of dosage compensation. Whether these consequences are general or linked to idiosyncrasies of specific taxa is not clear as relatively few taxa have been studied thus far. Here we use whole-genome sequencing to identify and characterize the evolution of the X chromosome in five species of *Timema* stick insects with XX:X0 sex determination. The X chromosome had a similar size (approximately 11% of the genome) and gene content across all five species, suggesting that the X chromosome originated prior to the diversification of the genus. Genes on the X showed evidence of a faster evolutionary rate than genes on the autosomes, likely due to less effective purifying selection. Genes on the X also showed almost complete dosage compensation in somatic tissues (heads and legs), but dosage compensation was absent in the reproductive tracts. Contrary to prediction, sex-biased genes showed little enrichment on the X, suggesting that the advantage X-linkage provides to the accumulation of sexually antagonistic alleles is weak. Overall, we found the consequences of X-linkage on gene sequences and expression to be similar across *Timema* species, showing the characteristics of the X chromosome are surprisingly consistent over 30 million years of evolution.

# Introduction

One of the most common forms of genomic variation between individuals within species stems from sex chromosomes. Sex chromosomes differ in copy number between males and females, which has a large effect on the evolutionary forces acting on genes located on them. Specifically, when males are the heterogametic sex (i.e., in XY or X0 systems), the genes on the X chromosome are present in only a single copy in males, while genes on the autosomes are present in two copies. This fundamental difference is expected to have a large effect on the evolutionary forces acting on genes located on the X chromosome, with consequences including a faster rate of sequence evolution, increased accumulation of sexually antagonistic alleles, and the evolution of dosage compensation mechanisms (Bachtrog et al. 2011; Wright et al. 2016; Lenormand and Roze 2021). Such effects should apply across taxa, meaning we should observe common patterns of X chromosome evolution in diverse species. Despite this, studies of sex chromosome evolution have shown a great deal of variation among study systems, including differences in the influence selection and drift have on the content of the X and in the extent of dosage compensation (Bachtrog et al. 2011; Gu and Walters 2017). Currently, it is difficult to understand the factors that govern this variation as only few taxa (typically with only a single or few representative species) have been studied. To elucidate these factors, studies of X chromosome evolution are thus needed from multiple species from a wide range of taxa (Palmer et al. 2019). Such studies will allow us to disentangling general sex chromosome-linked patterns from species-specific patterns, and allow us to develop a fuller understanding of sex chromosome evolution.

The rate of sequence evolution is expected to differ between the X chromosome and autosomes for two main reasons. Firstly, the X is hemizygous in males, meaning that recessive or partially recessive mutations in X-linked genes will be more exposed to selection than mutations in genes on the autosomes, allowing for beneficial mutations to be fixed and deleterious mutations to be purged more effectively (Charlesworth et al. 1987). On the other hand, as the X is only present in a single copy in males, its effective population size is expected to be smaller than that of the autosomes. This means that the effects of drift will also be stronger for genes on the X chromosome than on the autosomes, which will tend to reduce the fixation rate for advantageous mutations but increase it for deleterious mutations (Wright 1931; Vicoso and Charlesworth 2009). Together, these effects have been used as an explanation for the overall faster evolution of the X chromosome (the faster-X effect) seen in many species (Mank et al. 2010; Meisel and Connallon 2013; Parsch and Ellegren 2013; Charlesworth et al. 2018). In addition, the difference in X chromosome copy number between males and females is expected to facilitate the fixation of sexually antagonistic alleles (Rice 1984; Gibson et al. 2002; Mullon et al. 2012). This is because the X chromosome spends two-thirds of its time in females, giving an advantage to dominant female-beneficial alleles on the X. In addition, recessive alleles on the X will be exposed to selection on the X in males, giving an advantage to male-beneficial alleles. These complex forces have the potential to shape how sexually antagonistic variation is distributed across the genome, which can in turn influence broad evolutionary processes such as speciation (Coyne and Orr 2004; Payseur et al. 2018) and sexual conflict (Bachtrog et al. 2011; Mank et al. 2014; Wilkinson et al. 2015).

The fact that genes on the X are present in different copy numbers in males and females can
also create a problem for gene expression, as for many genes expression is proportional to
their copy number (Birchler and Veitia 2012; Birchler 2016). As such, species with
differentiated sex chromosomes should have evolved dosage compensation mechanisms to
equalise expression of the X chromosome in males and females (Ohno 1967; Charlesworth
1978; Charlesworth 1996). Note that such dosage compensation mechanisms can evolve
either in response to sex chromosomes differentiation or alongside it (Lenormand et al. 2020).
Dosage compensation has been demonstrated across a wide range of taxa (Disteche 2012;
Mank 2013; Gu and Walters 2017), including model species such as *Drosophila melanogaster*
(Conrad and Akhtar 2012) and *Caenorhabditis elegans* (Meyer 2000) where this phenomenon
has been studied in detail (Parkhurst and Meneely 1994; Lucchesi 1998; Meyer 2000; Straub
and Becker 2011; Conrad and Akhtar 2012). Despite this commonality, it has become
increasingly clear that the extent to which genes on the X are dosage compensated varies
among species. Several studied species show only partial or no dosage compensation (Mank
2013). The extent of dosage compensation may also differ by tissue type, with reduced dosage
compensation observed in the reproductive tracts of e.g. *C. elegans* (Kelly et al. 2002; Pirrotta
2002) or *D. melanogaster* (Oliver 2002; Meiklejohn et al. 2011; Mahadevaraju et al. 2021).
However, it is not clear how widespread tissue-specific dosage compensation is, as work in
non-model species often use whole-body samples for examining expression (Gu and Walters
2017)**.**
102
Here we expand our knowledge of the evolutionary characteristics of sex chromosomes by
identifying and studying the X chromosome in *Timema* stick insects. Aspects of X
chromosome evolution have been previously studied in several insect orders (Odonata
(Chauhan et al. 2021), Hemiptera (Pal and Vicoso 2015; Richard et al. 2017), Orthoptera
(Rayner et al. 2021), Strepsiptera (Mahajan and Bachtrog 2015), Coleoptera (Prince et al.
2010; Mahajan and Bachtrog 2015), and Diptera (Bone and Kuroda 1996; Marín et al. 1996;
Deng et al. 2011; Nozawa et al. 2014; Jiang et al. 2015; Vicoso and Bachtrog 2015; Rose et
al. 2016)). However, to date no studies have examined X chromosome evolution in stick
insects (Phasmatodea), an order which originated approximately 130 mya (Simon et al. 2019)
and contains around 3100 extant species (Bradler and Buckley 2018). First, we identified the
X chromosome in five species of *Timema* that diverged approximately 30 mya (Riesch et al.
2017). *Timema* have an XX/X0 sex determination system (Schwander and Crespi 2009). To
determine if genes on the X chromosome show the predicted faster rate of sequence evolution
than genes on the autosomes, we examined sequence evolution rates in each species. In
addition, we tested if the X chromosome is enriched for sexually antagonistic alleles. This was
done by examining if genes with sex-biased expression are enriched on the X chromosome,
as the evolution of sex-biased expression is thought to be driven primarily by sexually
antagonistic selection (Ellegren and Parsch 2007; Innocenti and Morrow 2010; Griffin et al.
2013). Finally, we examined if the genes on the X chromosome are dosage compensated by
comparing male and female gene expression in three composite tissues (heads, legs, and
reproductive tracts). Our study thus provides a detailed study of several key aspects of X
chromosome evolution in a previously unstudied group, revealing that the characteristics of
these X chromosomes are conserved over at least 30 million years of evolution.

# Results

**Identifying X-linked scaffolds**

We used a coverage approach to identify X-linked scaffolds in our previous genome assemblies (Jaron et al. 2021). Genomic reads from four males and five females from each species were mapped onto the corresponding reference genome. After filtering low-quality alignments and non-uniquely mapping reads (see Methods) the median coverage per sample ranged from 11x to 31x (18.5x on average, see Table S1). Visual inspection of coverage distributions (Figs S1-S5) found that while most libraries had either one (in females) or two (in males) coverage peaks, three libraries (*T. podura* (H56, Fig. S4), *T. poppensis* (ReSeq Ps08 and Reseq_Ps12, Fig. S5)) did not show a clear coverage peak, and were excluded from all further analyses.

To identify X-linked scaffolds, we used the $\log_2$ ratio of (normalised) coverage of males to females. As males have only a single copy of the X and females have two, X linked scaffolds should have twice as much coverage in females as males ($\log_2$ male:female coverage $\approx$ -1). Autosomes are expected to have the same coverage in both sexes ($\log_2$ male:female coverage $\approx$ 0). Considering all scaffolds with a $\log_2$ ratio of male to female coverage < autosomal peak -0.5 to be X-linked, we classified between 12 and 14 % of each genome as X-linked (Table 1, Fig. S6), which fits well with the X chromosome size observed in karyotypes (Schwander and Crespi 2009). This approach may mean some autosomal contigs may be misclassified as X-linked, thus we also repeated all our analyses with a more stringent classification scheme (scaffolds with a $\log_2$ ratio of male to female coverage within 0.1 of the X linked peak) (Table S2, Fig. S6). Using the more stringent classification scheme produced very similar results (not shown). Of note, most differences between the classification schemes are for short scaffolds (1000-4999 bp), which represent ~20% of the genome assemblies. When these are excluded the two classification schemes classify almost the same set of scaffolds as X-linked (Fig. S7, Table S3). Finally, we also examined the heterozygosity of the X in males. As expected, the heterozygosity of the X is close to zero and much lower than on the autosomes, corroborating our X-linked scaffold assignments (Fig. S8).

**Table 1 | Number of genes and scaffolds classified as X-linked or autosomal**

| Species | Min scaffold length | % genome X-linked | N Autosomal genes | N X-linked genes | N genes not classified |
|---|---|---|---|---|---|
| *T. bartmani* | 1000 | 12.11 | 12804 | 1244 | 18 |
| *T. cristinae* | 1000 | 12.21 | 12542 | 1316 | 24 |
| *T. californicum* | 1000 | 12.22 | 13207 | 1344 | 12 |
| *T. podura* | 1000 | 13.75 | 15069 | 1457 | 3 |
| *T. poppensis* | 1000 | 12.40 | 14115 | 1454 | 36 |

**The X chromosome is conserved across *Timema***

Comparing orthologs across the five different species shows X chromosome gene content is conserved, with >90% of X-linked orthologs shared between the 5 species (Fig. 1, Fig S9). This overlap is much greater than expected by chance (FDR < 6.863 x 10^{-316}). This suggests that the X chromosome is homologous in all five species, which last shared a common ancestor approximately 30 million years ago (Riesch et al. 2017). Additionally, we used

species genome alignments to an independent *T. cristinae* genome assembly (Nosil et al. 2018) to assign scaffolds to linkage groups based on a single reference. By applying coverage analyses we were able to identify and correct the X linked scaffolds in the Nosil et al. 2018 assembly (see Methods). Using this corrected reference, we found that contigs aligned to X-linked scaffolds showed reduced coverage in males but not females in each species (Figs S10-S14), again indicating the X chromosome is the same in all species. Finally, using BLAST we found that the majority (72%) of the shared X-linked genes in *Timema* (for which we were able to obtain a significant hit) were also present on the X chromosome of *Bacillus rossius* (Fig. S15). The split between *Timema* and all other extant phasmids (the Euphasmatodea, which includes *Bacillus*) occurred approximately 120 mya (Simon et al. 2019), suggesting that the X chromosome in phasmids predates this split.

**Fig. 1 | The X chromosome is conserved between *Timema* species. A.** Venn-diagrams showing the number of shared X-linked orthologs between species. **B.** Number of shared orthologs (expected, maximum possible). The observed amount of overlap was much greater than expected in all comparisons (FDR < 6.863 x 10$^{-316}$). Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps = *T. poppensis*, and Tpa = *T. podura*.

**The X chromosome has reduced genetic variation**

We used female samples to examine two related measures of genetic variation: heterozygosity and nucleotide diversity ($\pi$). Only female samples were used, as males are hemizygous for the X, which would affect the estimation of heterozygosity on the X directly. It would also affect the estimation of nucleotide diversity indirectly, as X haplotypes containing recessive lethals on the X will be absent in male samples. We found that both heterozygosity and nucleotide diversity ($\pi$) were lower on X-linked than on autosomal scaffolds (Fig. S8, S16), as expected. In all species, the ratios of X to autosomes for both of these measures (heterozygosity = 0.16 to 0.62 (Fig. 2), $\pi$ = 0.19 to 0.48 (Fig. 2)) were lower than the 0.75 that would be expected from the reduced effective population size of the X relative to the autosomes. This pattern was also seen when comparing the X to each of the autosomal linkage groups individually (Fig. S17, S18). Nucleotide diversity estimates were then used to

estimate the effective population size of each species. From this, we estimated the autosomal effective population size in *Timema* to range from ~150,000 (*T. poppensis*) to ~2,000,000 (*T. podura*) (Table S4).

**Fig. 2 | Genetic variation on the X chromosome is lower than expected. A.** Ratio of heterozygosity on the X and autosomes in 3 to 5 females per species. **B.** Ratio of pairwise nucleotide diversity (π) on the X and autosomes. Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps = *T. poppensis*, and Tpa = *T. podura*. The dotted lines indicate the neutral expectation in a population with a balanced sex ratio.

## The X chromosome shows evidence for reduced purifying selection but no differences in positive selection

The X chromosome shows a faster rate of sequence divergence than the autosomes as shown by an elevated dN/dS for X-linked genes (Fig. 3A). This increased rate appears to be driven by reduced purifying selection, as X-linked genes are not enriched for positively selected genes (Fig. 3B).

**Fig. 3 | Sequence evolution on the X and autosomes. A.** Average dN/dS across genes. Error bars indicate standard error. **B.** Proportion of positively selected genes. Positively selected genes were not enriched on the X chromosome (Fisher's exact test p value = 0.50). Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps = *T. poppensis*, and Tpa = *T. podura*.

## *Timema* show complete dosage compensation in heads and legs, but no dosage compensation in reproductive tracts

We examined if the X chromosome is dosage compensated in *Timema* by comparing gene expression in males and females in three different composite tissues (heads, legs, and reproductive tracts) for each of our five species. While for the main analysis we calculated expression values as FPKM, we also repeated our analyses using TPM. Analyses based on TPM showed very similar results and are provided as supplemental figures and tables. We examined the $\log_2$ ratio of male to female expression on the X and the autosomes. For the

autosomes and a dosage compensated X, the $\log_2$ value should be approximately 0, whereas a non-dosage compensated X would have a value of approximately -1. We found that in heads and legs the ratio was close to 0 with only small differences between the X and the autosomes (Figs. 4, S19, S20), indicating almost-complete complete dosage compensation in these tissues. By contrast, in the reproductive tracts the ratio of male to female expression for genes on the X is close to -1 (Figs. 4, S19, S20), indicating an absence of dosage compensation in this tissue. This observation was also seen when comparing the X to each of the autosomal linkage groups individually (Figs. S21-S22). An alternative, mutually non-exclusive possibility is that the greatly reduced expression observed in genes on the X in male reproductive tracts is due to a large enrichment of female-biased genes and depletion of male-biased genes. Although we cannot formally exclude this possibility, three lines of evidence indicate that lack of dosage compensation in reproductive tracts is the best explanation for our findings. Firstly, a lack of dosage compensation is expected to result in a two-fold reduction of expression, meaning X linked genes in males should show a major peak with a two-fold reduction in expression (Mank and Ellegren 2009; Vicoso et al. 2013; Pal and Vicoso 2015), which is what we observe (Fig. 5). Secondly, if the X chromosomes facilitates the accumulation of sex-biased genes, we should be able to observe this effect in all tissues (Jaquiéry et al. 2021). While we find a large enrichment of female-biased genes and a depletion of male-biased genes on the X in the reproductive tracts, this is not found in the other tissues, with only a slight enrichment of female-biased genes in the heads of two species and a depletion of male-biased genes in one species (Fig. 6, Fig. S23, Table S5, Table S6). This suggests that selection for the enrichment of female-biased genes and the depletion of male-biased genes on the X is weak, and thus unlikely to generate the large effect sizes we see in the reproductive tracts. Finally, a reduction in expression of the male X due to a lack of dosage compensation is expected to be consistent across species, since they share the same X chromosome (see above). By contrast, sex biased expression independently of dosage is more likely to be species-specific given the very fast turnover of sex biased genes observed between closely related species in several taxa (Zhang et al. 2007; Harrison et al. 2015) including *Timema* (Parker et al. 2019a). To distinguish between these patterns, we tested if sex-biased genes on the X in the reproductive tracts are underrepresented for genes with species by sex interactions. We found that genes on the X are underrepresented for genes showing species by sex interactions in the reproductive tracts ($p = 0.010$), and that this is not the case for the heads ($p = 0.938$) or legs ($p = 0.795$). This shows that sex-differences in expression on the X are more consistent between species in the reproductive tracts than in the heads and legs (also see Fig. S24), and further supports a lack of dosage compensation in the reproductive tracts rather than a large enrichment of female-biased genes on the X.

**Fig. 4 | Gene expression on the X and autosomes in heads, legs, and reproductive tracts. A.** Average expression levels in males and females on the X and autosomes **B.** Log2 of male to female expression ratio for the X and autosomes. Dashed lines represent a two-fold reduction in expression in males (as expected if there was no dosage compensation). Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps = *T. poppensis*, and Tpa = *T. podura*.

By examining the expression of the X and autosomes in the different tissues, we can infer the
type of dosage compensation. In the dosage compensated tissues (heads and legs) genes on
the autosomes and X have similar overall expression levels in both males and females, and
the overall expression of the X is similar to that of the autosomes (Fig 4, Fig S19, Table S7).
By contrast, in the reproductive tracts, where dosage compensation is lacking, expression of
genes on the X is much lower in males than in females. This difference seems to be driven by
changes in male expression, as X-linked gene expression in females remains similar to the
autosomes (Fig 4, Fig S19, Table S7). This supports a mechanism of dosage compensation
by hyper-transcription of the X in males, a mechanism common among other insects (Gu and
Walters 2017).

Although we find almost complete dosage compensation, it is possible that its extent could
vary along the X chromosome (Mullon et al. 2015). To address this, we examined the extent
of dosage along the longest X-linked scaffolds in the genome, and found no evidence of
variation along the X chromosome (Fig. S25 - S29, see also Fig. S30 - S41 for expression
variation along the autosomes).

**Fig. 5 | Ratio of male and female gene expression on the X (orange) and autosomes (black) in heads, legs, and reproductive tracts**. Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps = *T. poppensis*, and Tpa = *T. podura*.

**Sex-biased genes are not enriched on the X chromosome**

Sexually antagonistic mutations are expected to fix more easily on the X chromosome than the autosomes. As the evolution of sex-biased gene expression is thought to be primarily driven by sexually antagonistic selection, it is expected that the X chromosome will be a hotspot for sex-biased gene expression (Rice 1984; Gibson et al. 2002; Ellegren and Parsch 2007; Innocenti and Morrow 2010; Griffin et al. 2013). Contrary to this expectation, we find most sex-biased genes on the autosomes (% of sex-biased genes on the X ranges from 8.3 - 14.3, Table S5), and we find little evidence for enrichment of sex-biased genes on the X in the head and leg tissues (Fig. 6, Table S5, Table S6). Note that almost all genes on the X appear to be sex-biased in the reproductive tracts, but this effect is largely due to a lack of dosage compensation in this tissue (see above). In addition, while we do find an enrichment of female-biased genes in the heads of two species (*T. bartmani* and *T. podura*, Fig. 6) and a depletion of male-biased genes in *T. bartmani,* the effect sizes are small and the effect becomes insignificant when considering only sex-biased genes with at least a twofold difference in expression between males and females (Fig. S23). These findings suggest that the selective pressures to accumulate female-biased genes and reduce male-biased genes on the X are weak and/or that there are constraints to the gene content or expression levels on the *Timema* X chromosome.

**Fig. 6 | Proportion of female- and male- biased genes on the X and autosomes in reproductive tract, head and leg samples.** Note the scale changes between tissue-types. Asterisks indicate the significance level (FDR) of Fisher's exact tests (***<0.001, **<0.01, *<0.05). Species names are abbreviated as Tbi = *T. bartmani*, Tce = *T. cristinae*, Tcm = *T. californicum*, Tps = *T. poppensis*, and Tpa = *T. podura*.

# Discussion

The difference in copy numbers of the X chromosome in males and females is expected to have profound effects on its evolution. In particular, the X chromosome is predicted to evolve at a faster rate due to a combination of hemizygous selection and increased drift (Wright 1931; Charlesworth et al. 1987; Vicoso and Charlesworth 2009), to accumulate sexually antagonistic alleles (Rice 1984; Gibson et al. 2002), and to evolve dosage compensation mechanisms (Disteche 2012; Mank 2013; Gu and Walters 2017; Lenormand et al. 2020). Support for these predictions has been mixed from the taxa examined so far, yet the factors responsible for the variation among different taxa are poorly understood (Bachtrog et al. 2011; Gu and Walters 2017). In this study, we examine these predictions across five species of *Timema* stick insects. Overall, we find evidence for a faster-X effect and complete dosage compensation in somatic tissues, but little evidence for the accumulation of sexually antagonistic alleles on the X. Patterns of X chromosome evolution were generally consistent across *Timema* species, suggesting that the factors influencing sex chromosome evolution in this group are also largely the same.

Sex chromosome conservation is highly variable between taxa, with extensive turnover between species in some groups, e.g. beetles (Coleoptera) (Blackmon and Demuth 2014), flies (Diptera) (Vicoso and Bachtrog 2015), or frogs (Ranidae) (Jeffries et al. 2018), and conservation for over a hundred million years in others, e.g. Eutherian mammals (Lahn and Page 1999; Cortez et al. 2014; Marshall Graves 2015), moths and butterflies (Lepidoptera) (Fraïsse et al. 2017), or birds (Shetty et al. 1999; Xu and Zhou 2020). The factors influencing turnover rate are complex and interacting (Vicoso 2019), however, one key factor is the level of differentiation between the X and Y chromosomes, with more differentiated chromosomes less likely to turnover (Pokorná and Kratochvíl 2009; Vicoso 2019). Our finding that the X chromosome in *Timema* is old (likely conserved for over 120 million years) supports this idea. XX/X0 sex determination systems as found in *Timema* (Schwander and Crespi 2009) are thought to derive from XX/XY systems with highly differentiated X and Y chromosomes and represent the end point of the gradual loss of gene content from the Y (Bergero and Charlesworth 2009). As such, XX/X0 systems could be considered the most extreme example of sex chromosome differentiation possible, which may mean that XX/X0 systems are particularly unlikely to turnover (but see (Blackmon and Demuth 2014)).

Gene sequence evolution on the X chromosome is expected to be faster than that on the autosomes due to a combination of increased drift and hemizygous selection (Charlesworth et al. 1987; Vicoso and Charlesworth 2009). While this prediction should apply universally, support for a faster-X effect is mixed (Mank et al. 2010; Meisel and Connallon 2013; Charlesworth et al. 2018; Pinharanda et al. 2019; Whittle et al. 2020). The cause of this variation is unclear, in particular, because typically only single lineages are examined. In single lineages it is difficult to disentangle the influence of X-linkage from lineage-specific effects such as recent bottlenecks, differences in operational sex ratio, or population size. By examining the influence of X-linkage in multiple *Timema* species covering a span of 30 million years of divergence, we can assess how consistent the effects of X linkage are, allowing for a comprehensive assessment of X chromosome evolution in this genus. Using this approach,

368  we found consistent evidence for a faster X effect in all species, and that this effect is primarily
369  driven by reduced purifying selection on the X.
370
The reduction of purifying selection on the X relative to the autosomes is expected to be largest
when the effective population size of the X ($Ne_X$) is much smaller than that of the autosomes
($Ne_A$) (Vicoso and Charlesworth 2009). The neutral expectation in a population with a
balanced sex ratio is that the ratio of $Ne_X$ to $Ne_A$ will be approximately 0.75 (Wright 1969; Hartl
and Clark 2006), however, demographic and selective processes can have a large effect on
this ratio (Nunney 1993; Caballero 1995; Charlesworth 2001). In *Timema*, $Ne_X$ / $Ne_A$ values
are much smaller than 0.75 (0.19 to 0.48, estimated from nucleotide diversity), meaning the
effective population size of the X is much smaller than that of the autosomes. Departures from
the expected 0.75 nucleotide diversity ratio are common across animals and are typically
thought to be a consequence of sex-biased demography (Mank et al. 2010). In male
heterogametic species such as *Timema*, increased reproductive skew for males (for example
under harem-like mating systems) will disproportionately reduce diversity on the autosomes.
This would result in an X to autosome nucleotide diversity ratio > 0.75. Here we observe the
opposite pattern, a reduction in the autosome nucleotide diversity ratio. As such it is likely that
the reduced diversity on the X is not due to demography but perhaps due to reduced
recombination rates on the X relative to the autosomes, and/or to selective differences on the
X. In particular, the X chromosome is likely to be susceptible to an increased frequency of
selective sweeps / background selection which will disproportionately reduce genetic diversity
on the X (Betancourt et al. 2004; Charlesworth 2012; Charlesworth 2013; Wilson Sayres
2018).
391
Independently of the mechanisms responsible for the strongly reduced effective population
size of the X in *Timema*, it supports our interpretation that the faster-X effect is driven by less
effective purifying selection. Most previous studies of the faster-X effect have focused on
species with $Ne_X$ / $Ne_A$ values ≥ 0.75 (Mank et al. 2010), making direct comparisons with our
study difficult. The exceptions to this are studies on birds and *Heliconius* butterflies which have
values of $Ne_Z$ / $Ne_A$ in a similar range as *Timema*. Note that both these taxa have
heterogametic females (ZW), where low $Ne_Z$ / $Ne_A$ ratios are believed to stem from strong
reproductive skew among males (Vicoso and Charlesworth 2009; Mank et al. 2010). In birds,
a faster-Z (faster-X) effect is commonly observed and appears to be driven primarily by less
effective purifying selection (Mank et al. 2010). In contrast, in *Heliconius* evidence for a faster-
Z effect is weaker and is thought to be driven by increased levels of adaptive evolution on the
Z, with no evidence for reduced purifying selection on the Z (Pinharanda et al. 2019). The
cause of this difference is unclear. However, it has been suggested that it may be due to the
overall higher effective population size in *Heliconius* ($Ne \approx 2,000,000$ (Keightley et al. 2015))
compared to birds ($Ne = 200,000 - 600,000$ (Primmer et al. 2002; Axelsson et al. 2004;
Jennings and Edwards 2005; Backström et al. 2008)), meaning that selection can be efficient
in *Heliconius* even with the relatively reduced effective population size of the sex chromosome
(Mank et al. 2010; Pinharanda et al. 2019). This interpretation conflicts with our findings in
*Timema*, as the (autosomal) effective population size varies greatly between species (from
~150,000 in *T. poppensis* to ~2,000,000 in *T. podura*), yet we observe a similar reduction in
purifying selection on the X in all species. The reason for this variation between study systems
is thus unclear, and highlights the need for future studies across diverse taxa.
414

The X chromosome has long been predicted to be a hotspot for sexually antagonistic variation (Rice 1984; Gibson et al. 2002). The evolution of sex-biased gene expression is thought to be driven by sexually antagonistic selection, and thus sex-biased gene expression should be overrepresented on the X chromosome ((Ellegren and Parsch 2007; Innocenti and Morrow 2010; Griffin et al. 2013) but see (Hitchcock and Gardner 2020; Ruzicka and Connallon 2020)). In *Timema*, we find very little support for this prediction with only a small enrichment of sex-biased genes on the X in one of our five species. In combination with studies in *Drosophila melanogaster* (Ruzicka et al. 2019), *Callosobruchus maculatus* (Sayadi et al. 2019), and *Ischnura elegans* (Chauhan et al. 2021) which also found little or no enrichment of sexually antagonistic alleles or sex-biased gene expression on the sex chromosomes, our study suggests that the advantage of accumulating sexually antagonistic alleles on the X be may be smaller than often assumed. The reasons for this are unclear. However, it is likely that the advantage X-linkage gives to sexually antagonistic alleles is balanced by other forces such as epistatic interactions (Arnqvist et al. 2014) or sex-specific dominance (Fry 2010). Both forces favour the accumulation of sexually antagonistic alleles on the autosomes. Despite this, several studies do show the expected enrichment of sexually antagonistic alleles or sex-biased gene expression on the X: in *Tribolium castaneum* (Whittle et al. 2020), Diptera (Innocenti and Morrow 2010; Vicoso and Bachtrog 2015), Hemiptera (Pal and Vicoso 2015), and nematodes (Albritton et al. 2014). A key challenge for future work will thus be to integrate studies that quantify multiple factors that influence the accumulation of sexually antagonistic alleles (e.g. epistatic interactions, sex-specific dominance, reproductive skew, etc.) to understand how the variation between studies and taxa is produced.

In many species dosage compensation is thought to be important for ameliorating the costs of misexpression of genes on the X chromosome (Marín et al. 2000). Although common, there is a great deal of variation in the extent to which genes are dosage compensated. Here we find almost complete dosage compensation in the somatic tissues of all five *Timema* species, as reported for other species with X0 systems (e.g. Nematodes (Meyer 2000) and crickets (Rayner et al. 2021)). Dosage compensation is expected for X0 systems as they are thought to arise from XY systems where the Y chromosome has degraded to the point it can be lost without a large decrease in fitness. By this stage, most genes on the X should already be haplo-sufficient in males. In addition, the evolution of dosage compensation could itself hasten the loss of the Y chromosome, as genes that have functional copies on both the X and the Y will be misexpressed if chromosome-wide dosage compensation evolves (Vicoso and Bachtrog 2009; Lenormand and Roze 2021).

In contrast to the somatic tissues, male reproductive tracts displayed a lack of dosage compensation. Reduced expression of X chromosome in male reproductive tissue has been observed in a number of species including mammals (Khil et al. 2004; Disteche 2012; Sangrithi and Turner 2018), *C. elegans* (Kelly et al. 2002; Pirrotta 2002) and insects such as *D. melanogaster* (Oliver 2002; Meiklejohn et al. 2011; Mahadevaraju et al. 2021) or *Teleogryllus oceanicus* (Rayner et al. 2021). Whether reduced X expression in male reproductive tracts generally stems from a lack of dosage compensation is, however, not clear. Indeed, reduced X expression can be caused by several non-mutually exclusive mechanisms, including the accumulation of female-biased genes on the X (Mank 2009), the movement of male-beneficial genes to the autosomes (Vibranovski et al. 2009), or the inactivation of X chromosomes in the germ line (Lee 2005; Vibranovski 2014). In *Timema*, we suggest an absence of dosage compensation is the most likely explanation as expression of X-linked genes in the male

reproductive tracts show a major peak of genes with expression approximately half of that observed in females, similar to that observed in species that lack dosage compensation (Mank and Ellegren 2009; Vicoso et al. 2013). It should be noted, however, that the expression reduction we see on the male X in *Timema* is actually slightly less than the half we would expect from a lack of dosage compensation alone. This suggests that other factors, described above, may also have an influence. Disentangling the contribution of each of these factors will require further work. Our work, however, clearly shows that expression in reproductive tissues behaves differently than in somatic tissues, highlighting the importance of studying these tissues separately particularly when assessing the extent of dosage compensation (Gu and Walters 2017).

Dosage compensation in *Timema* somatic tissues appears to be achieved by the upregulation of the X in males (type I, (Gu and Walters 2017)) as indicated by similar expression levels of genes on the X and the autosomes in both sexes. This interpretation assumes that the ancestral state of X chromosome expression (before it was a sex chromosome) was similar to the other autosomes. This seems likely as expression across all *Timema* autosomes is similar. However, assessing the ancestral state of X expression is largely impossible as the *Timema* X chromosome appears to be very old (~120 million years). As a consequence, there are likely no species available for comparison with conserved genome organisation but with X chromosomes that are homologous to different *Timema* autosomes. Similar to *Timema*, upregulation of the X chromosome in males also appears to be the mechanism for dosage compensation in all other XX/XY or XX/X0 insect systems yet studied, even when dosage compensation is incomplete (Gu and Walters 2017). With our study, this amounts to information from seven different insect orders (Odonata (Chauhan et al. 2021), Phasmatodea (this study), Hemiptera (Pal and Vicoso 2015; Richard et al. 2017), Orthoptera (Rayner et al. 2021), Strepsiptera (Mahajan and Bachtrog 2015), Coleoptera (Prince et al. 2010; Mahajan and Bachtrog 2015), and Diptera (Bone and Kuroda 1996; Marín et al. 1996; Deng et al. 2011; Nozawa et al. 2014; Jiang et al. 2015; Vicoso and Bachtrog 2015; Rose et al. 2016)) suggesting that this mechanism may be universal for insect species with heterogametic males.

**Conclusions**

How consistent the consequences of sex linkage are for gene sequence and expression evolution across taxa remains an open question. Here we examine several key aspects of sex chromosome evolution in a previously neglected group, phasmids. Overall, we find evidence for several predicted consequences, including complete dosage compensation of the X (in somatic tissues) and a faster rate of evolution of X-linked than autosomal genes. By contrast, we find little evidence that sex linkage facilitates the accumulation of sexually antagonistic alleles. While our results were consistent across different *Timema* species, they also show key differences from studies in other taxa, highlighting the importance of studying sex chromosome evolution across a diverse set of species to distinguish general patterns caused by sex linkage from species-specific idiosyncrasies.

# Methods

## Sample collection and sequencing

We used a combination of available genomic data from females in addition to newly collected data for males. Reads from five females per species were downloaded from NCBI (Bioproject accession number: PRJNA670663). We collected four males from each of the five species from natural populations in California, from the same (or a geographically very close) population as the available females (Table S8). DNA extractions were done on whole-body adult males using the Qiagen Mag Attract HMW DNA kit following the manufacturer instructions. Sequencing libraries were generated for each male using a TruSeq DNA nano prep kit (550bp insert size). Libraries were then sequenced using Illumina HiSeq 2500 at the Lausanne Genomic Technologies Facility.

## Using coverage to identify X-linked scaffolds

Reads were trimmed before mapping using Trimmomatic (v. 0.36) (Bolger et al. 2014) to remove adapter and low-quality sequences (options: ILLUMINACLIP:3:25:6 LEADING:9 TRAILING:9 SLIDINGWINDOW:4:15 MINLEN:90). Reads from each individual were mapped to their species' reference genome (Jaron et al. 2021) (Bioproject accession number: PRJEB31411) using BWA-MEM (v. 0.7.15) (Li 2013). Multi-mapping and poor quality alignments were filtered (removing reads with XA:Z or SA:Z tags or a mapq < 30). PCR duplicates were removed with Picard (v. 2.9.0) (http://broadinstitute.github.io/picard/). Coverage was then estimated for all scaffolds at least 1000 bp in length using BEDTools (v. 2.26.0) (Quinlan and Hall 2010). Per base coverage distributions were inspected visually for each individual and libraries with extremely non-normal coverage distributions were excluded from further analysis (Fig S1-S5) .

To compare coverage between males and females, coverage was first summed for all male and all female libraries per scaffold (for scaffolds at least 1000 bp in length). Male and female coverage was then normalised by modal coverage to adjust for differences in overall coverage. X-linked scaffolds were then identified using the $\log_2$ ratio of male to female coverage. Autosomal scaffolds should have equal coverage in males and females ($\log_2$ ratio of male to female coverage $\approx$ 0) and X-linked scaffolds should have half the coverage in males as in females ($\log_2$ ratio of male to female coverage $\approx$ -1). Each species showed frequency peaks near these values (Fig S6). X linked scaffolds were classified in two ways: a 'liberal' classification (following (Vicoso and Bachtrog 2015)) whereby scaffolds were classified as X-linked if the $\log_2$ ratio of male to female coverage < autosomal peak - 0.5, and a 'stringent' classification (following (Pal and Vicoso 2015)) whereby scaffolds were classified as X-linked if the $\log_2$ ratio of male to female coverage was within 0.1 of the value of the X-linked peak.

To compare X-linked scaffolds between species, we examined the overlap of one-to-one orthologs (previously identified in (Jaron et al. 2021)) on the X-linked scaffolds. To determine if the overlap of X-linked orthologs between species was greater than expected, we used the SuperExactTest package (v. 0.99.4) (Wang et al. 2015) in R (R Core Team 2017). Additionally,

548  we used a tblastn approach to identify which regions of the *Bacillus rossius* genome
549  correspond to the 210 shared X linked orthologs in *Timema* (minimum e-value = 1 x 10$^{-20}$,
550  minimum query coverage = 50%).

## Heterozygosity, nucleotide diversity, effective population size

To calculate heterozygosity, reads were mapped to the reference genomes as described
above. We then additionally performed indel realignment with GATK (v. 3.7) (Van der Auwera
et al. 2013). A maximum-likelihood estimate of the number of heterozygous sites per scaffold
was then calculated using AngsD (v. 0.921) (Korneliussen et al. 2014) (options: -doSaf 1 -gl 1
-minQ 20 -minMapQ 40 -fold 1 -doCounts 1, with a minimum depth of 5 and a maximum depth
of twice the median genome coverage once sites with 0 coverage were excluded) for each
sample (for all scaffolds at least 1000 bp in length). The proportion of heterozygous sites was
then calculated for each scaffold and the median proportion of heterozygous sites was
weighted by the number of covered sites on a scaffold. A similar approach was used to
calculate pairwise nucleotide diversity for each species, but using only female samples
(options: -doSaf 1 -gl 1 -minQ 20 -minMapQ 40 -fold 1 -doCounts 1 with a minimum depth of
5 and a maximum depth of twice the median genome coverage once sites with 0 coverage
were excluded). The number of individuals (-nind) and the minimum number of individuals a
site must be present in (-minind) were both set to the number of samples analysed. Effective
population size was estimated for the X and autosomes using the estimates of nucleotide
diversity, assuming that nucleotide diversity is equal to $4Ne\mu$. Estimates of the mutation rate
($\mu$) were taken from *Drosophila melanogaster* (2.8 x 10$^{-09}$) (Keightley et al. 2014) and
*Heliconius melpomene* (2.9 x 10$^{-09}$) (Keightley et al. 2015).

## Selection analyses

We took values of dN/dS for each species branch from Jaron et al (2021). Briefly, branch-site
models with rate variation at the DNA level (Davydov et al., 2019) were run using the Godon
software (https://bitbucket.org/Davydov/godon/, version 2020-02-17, option BSG --ncat 4) for
each gene with an ortholog found in at least three of the species of *Timema* used here. Godon
estimates the proportion of sites evolving under purifying selection (p0), neutrality (p1), and
positive selection. We used only sites evolving under purifying or neutrality to calculate dN/dS.
To test for differences in sequence divergence rates (dN/dS) between the X and autosomes,
we used Wilcoxon tests with p-values adjusted for multiple comparisons using Benjamini and
Hochberg's algorithm (Benjamini and Hochberg 1995). Using the same method, Jaron et al
(2021) also identified branches with evidence for positively selected sites. We determined if
these were overrepresented on the X using a Fisher's exact test.

## Gene expression analyses

RNA-seq reads from three composite tissues (heads, legs and reproductive tracts) for males
and females (3 replicates per sex) for each of the five species are publically available (Parker
et al. 2019a; Parker et al. 2019b) (Bioproject accession number: PRJNA392384). Adapter
sequences were removed using Cutadapt (v. 2.3) (Martin 2011) before quality trimming reads

with Trimmomatic (v. 0.36) (Bolger et al. 2014) (options: LEADING:10 TRAILING:10
SLIDINGWINDOW:4:20 MINLEN:80). Trimmed reads were then mapped to reference
genomes using STAR (v. 2.6.0c, default options). HTSeq v.0.9.1 (Anders et al. 2015) was
used to count the number of reads uniquely mapped to the exons of each gene, with the
following options (htseq-count --order=name --type=exon --idattr=gene_id --
stranded=reverse). Expression analyses were performed using the Bioconductor package
EdgeR (v 3.32.1) (Robinson et al. 2010), and done separately for each species and tissue.
Normalisation factors for each library were computed using the TMM method and were used
to calculate normalised expression levels (either FPKM or TPM). For the main analyses, genes
with low expression (less than 2 FPKM (or TPM) in 2 or more libraries per sex) were excluded.
This filtering step was used to exclude any sex-specifically expressed genes as our goal was
to examine how the expression of genes differs in males and females. This decision could
influence our results if sex-limited gene expression was extensive on the X chromosome. To
investigate this we repeated our analysis with the inclusion of sex-limited genes and found
similar results to the main analyses (Fig. S42 - S43). To examine dosage compensation, we
used the $\log_2$ ratio of mean male expression level to female expression level and used
Wilcoxon tests (adjusted for multiple testing using Benjamini and Hochberg's algorithm
(Benjamini and Hochberg 1995)) to determine if this ratio differed between genes on the
autosomal and X-linked scaffolds.

To determine the significance of sex on gene expression, we fit a generalised linear model
(GLM) with a negative binomial distribution with sex as an explanatory variable and used a
GLM likelihood ratio test to determine the significance for each gene. P-values were then
corrected for multiple tests using Benjamini and Hochberg's algorithm (Benjamini and
Hochberg 1995). In the main analysis sex-biased genes were then defined as genes that
showed difference in expression between males and females with an FDR < 0.05 (we also
repeated analyses with the additional condition that genes must show a greater than two fold
difference in expression to ensure our results are robust to the effects of sex-biased allometry
(Montgomery and Mank 2016)). Note that all genes not classified as sex-biased were
classified as unbiased genes. We then examined if male- or female- biased genes were under-
or over- represented on autosomal or X-linked scaffolds using Fisher's exact tests (adjusted
for multiple-testing using Benjamini and Hochberg's algorithm (Benjamini and Hochberg
1995)). To examine species by sex interactions in gene expression, we used a similar similar
GLM approach as above, with sex, species, and species by sex interaction as explanatory
variables for genes with an ortholog in each species. Genes with an FDR < 0.1 for the
interaction term were considered to have significant species by sex interaction. To determine
if sex-biased genes on the X were underrepresented for species by sex interactions, we used
a Fisher's exact test.

It is possible that dosage compensation may vary along the X chromosome. Unfortunately,
our current reference genomes are too fragmented to investigate this question. Previous work
by Nosil et al. (Nosil et al. 2018) however produced a more contiguous genome assembly of
one of our study species, *T. cristinae,* which has been further constructed into a linkage map.
To use this synteny information for each of our species, we anchored the scaffolds from each
of our genome assemblies to the Nosil et al. (Nosil et al. 2018) reference genome (BioProject
Accession PRJNA417530) using MUMmer (version 4.0.0beta2) (Marçais et al. 2018) with
parameter --mum. The alignments were processed by other tools within the package: show-
coords with parameters -THrcl to generate tab-delimited alignment files and dnadiff to

generate 1-to-1 alignments. We used only uniquely anchored scaffolds for which we were able
to map at least 10k nucleotides to the Nosil et al. (2018) reference genome. Nosil et al (2018)
indicated that linkage group 13 was the X chromosome ((Nosil et al. 2018)). To determine if
this was correct, we repeated our coverage analyses on the Nosil et al. (2018) assembly. From
this, we found that most scaffolds that make up linkage group 13 did not show reduced
coverage in males (Fig. S44). In addition, several scaffolds from other linkage groups did show
reduced coverage in males (Fig. S44). In order to use as much of the synteny information as
possible we "cleaned" the Nosil et al. (2018) assembly by removing X-linked scaffolds from
autosomal linkage groups 1-12. These scaffolds were then assigned to a new, unordered
collection of scaffolds from the X chromosome, together with X-linked scaffolds from linkage
group 13 and from those not assigned to any linkage group in Nosil et al. (2018). Scaffolds
from linkage group 13 that were not X-linked were assigned to linkage group NA. We classified
scaffolds in the Nosil et al. (2018) assembly as X-linked if most of a scaffold was covered by
aligned scaffolds from our assembly assigned to the X rather than autosomes (i.e. if aligned
scaffolds assigned as X covered more than twice as many bases as those assigned as
autosomal for a particular scaffold, it was classed as X-linked).

# Acknowledgements

656
We would like to thank Chloe Larose, Kirsten Jalvingh, and Bart Zijlstra for their assistance in
the field. This project was supported by several grants from the Swiss Science Foundation:
PP00P3_170627 and 31003A_182495 to TS as well as CRSII3_160723 to TS, MRR and
Nicolas Galtier. We would like to thank the former Vital-IT platform of the SIB Swiss Institute
of Bioinformatics (SIB) and the DCSR for maintaining the computer infrastructure at the
University of Lausanne.

# Data and code availability

664  Raw sequence reads have been deposited in NCBI's sequence read archive under the
665  bioproject: PRJNA725673 (Table S8). Scripts for the analyses in this paper are available at:
666  https://github.com/DarrenJParker/Timema_sex_chr_evol_code and will be archived at
667  Zenodo upon acceptance. Data was processed to generate plots and statistics using R v4.0.3
668  and Python v.3.7.3 unless otherwise stated.

# References

Albritton SE, Kranz A-L, Rao P, Kramer M, Dieterich C, Ercan S. 2014. Sex-biased gene expression and evolution of the x chromosome in nematodes. *Genetics* 197:865–883.

Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31:166–169.

Arnqvist G, Vellnow N, Rowe L. 2014. The effect of epistasis on sexually antagonistic genetic variation. *Proceedings of the Royal Society B: Biological Sciences* 281:20140489.

Axelsson E, Smith NGC, Sundström H, Berlin S, Ellegren H. 2004. Male-biased mutation rate and divergence in autosomal, Z-linked and W-linked introns of chicken and turkey. *Mol. Biol. Evol.* 21:1538–1547.

Bachtrog D, Kirkpatrick M, Mank JE, McDaniel SF, Pires JC, Rice W, Valenzuela N. 2011. Are all sex chromosomes created equal? *Trends Genet.* 27:350–357.

Backström N, Fagerberg S, Ellegren H. 2008. Genomics of natural bird populations: a gene-based set of reference markers evenly spread across the avian genome. *Mol. Ecol.* 17:964–980.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. R. Stat. Soc. Series B Stat. Methodol.* 57:289–300.

Bergero R, Charlesworth D. 2009. The evolution of restricted recombination in sex chromosomes. *Trends Ecol. Evol.* 24:94–102.

Betancourt AJ, Kim Y, Orr HA. 2004. A pseudohitchhiking model of X vs. autosomal diversity. *Genetics* 168:2261–2269.

Birchler JA. 2016. Parallel universes for models of X chromosome dosage compensation in *Drosophila*: a review. *Cytogenet. Genome Res.* 148:52–67.

Birchler JA, Veitia RA. 2012. Gene balance hypothesis: connecting issues of dosage sensitivity across biological disciplines. *Proc. Natl. Acad. Sci. U. S. A.* 109:14746–14753.

Blackmon H, Demuth JP. 2014. Estimating tempo and mode of Y chromosome turnover: explaining Y chromosome loss with the fragile Y hypothesis. *Genetics* 197:561–572.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120.

Bone JR, Kuroda MI. 1996. Dosage compensation regulatory proteins and the evolution of sex chromosomes in *Drosophila*. *Genetics* 144:705–713.

Bradler S, Buckley TR. 2018. Biodiversity of Phasmatodea. In: Insect Biodiversity. Chichester, UK: John Wiley & Sons, Ltd. p. 281–313.

Caballero A. 1995. On the effective size of populations with separate sexes, with particular reference to sex-linked genes. *Genetics* 139:1007–1011.

Charlesworth B. 1978. Model for evolution of Y chromosomes and dosage compensation. *Proc. Natl. Acad. Sci. U. S. A.* 75:5618–5622.

Charlesworth B. 1996. The evolution of chromosomal sex determination and dosage

compensation. *Curr. Biol.* 6:149–162.

Charlesworth B. 2001. The effect of life-history and mode of inheritance on neutral genetic
variability. *Genet. Res.* 77:153–166.

Charlesworth B. 2012. The role of background selection in shaping patterns of molecular
evolution and variation: evidence from variability on the *Drosophila* X chromosome.
*Genetics* 191:233–246.

Charlesworth B. 2013. Background selection 20 years on: the Wilhelmine E. Key 2012
invitational lecture. *J. Hered.* 104:161–171.

Charlesworth B, Campos JL, Jackson BC. 2018. Faster-X evolution: Theory and evidence
from *Drosophila*. *Mol. Ecol.* 27:3753–3771.

Charlesworth B, Coyne JA, Barton NH. 1987. The relative rates of evolution of sex
chromosomes and autosomes. *Am. Nat.* 130:113–146.

Chauhan P, Swaegers J, Guillen RAS, Svensson EI, Wellenreuther M, Hansson B. 2021.
Genome assembly, sex-biased gene expression and dosage compensation in the
damselfly *Ischnura elegans*. *Genomics* 113:1828–1837.

Conrad T, Akhtar A. 2012. Dosage compensation in *Drosophila melanogaster*: epigenetic fine-
tuning of chromosome-wide transcription. *Nat. Rev. Genet.* 13:123–134.

Cortez D, Marin R, Toledo-Flores D, Froidevaux L, Liechti A, Waters PD, Grützner F,
Kaessmann H. 2014. Origins and functional evolution of Y chromosomes across
mammals. *Nature* 508:488–493.

Coyne JA, Orr HA. 2004. Speciation. Sunderland, USA: Sinauer Associates

Deng X, Hiatt JB, Nguyen DK, Ercan S, Sturgill D, Hillier LW, Schlesinger F, Davis CA, Reinke
VJ, Gingeras TR, et al. 2011. Evidence for compensatory upregulation of expressed X-
linked genes in mammals, *Caenorhabditis elegans* and *Drosophila melanogaster*. *Nat.*
*Genet.* 43:1179–1185.

Disteche CM. 2012. Dosage compensation of the sex chromosomes. *Annu. Rev. Genet.*
46:537–560.

Ellegren H, Parsch J. 2007. The evolution of sex-biased genes and sex-biased gene
expression. *Nat. Rev. Genet.* 8:689–698.

Fraïsse C, Picard MAL, Vicoso B. 2017. The deep conservation of the Lepidoptera Z
chromosome suggests a non-canonical origin of the W. *Nat. Commun.* 8:1–9.

Fry JD. 2010. The genomic location of sexually antagonistic variation: some cautionary
comments. *Evolution* 64:1510–1516.

Gibson JR, Chippindale AK, Rice WR. 2002. The X chromosome is a hot spot for sexually
antagonistic fitness variation. *Proc. Biol. Sci.* 269:499–505.

Griffin RM, Dean R, Grace JL, Rydén P, Friberg U. 2013. The shared genome is a pervasive
constraint on the evolution of sex-biased gene expression. *Mol. Biol. Evol.* 30:2168–2176.

Gu L, Walters JR. 2017. Evolution of sex chromosome dosage compensation in animals: a
beautiful theory, undermined by facts and bedeviled by details. *Genome Biol. Evol.*
9:2461–2476.

Harrison PW, Wright AE, Zimmer F, Dean R, Montgomery SH, Pointer MA, Mank JE. 2015.
Sexual selection drives evolution and rapid turnover of male gene expression. *Proc. Natl.*
*Acad. Sci. U. S. A.* 112:4393–4398.

Hartl DL, Clark AG. 2006. Principles of Population Genetics. 4th ed. Sinauer Associates

Hitchcock TJ, Gardner A. 2020. A gene's-eye view of sexual antagonism. *Proc. Biol. Sci.*
287:20201633.

Innocenti P, Morrow EH. 2010. The sexually antagonistic genes of *Drosophila melanogaster*.
*PLoS Biol.* 8:e1000335.

Jaquiéry J, Simon J-C, Robin S, Richard G, Peccoud J, Boulain H, Legeai F, Tanguy S,
Prunier-Leterme N, Le Trionnaire G. 2021. Masculinization of the X-chromosome in aphid
soma and gonads. *bioRxiv*:2021.08.13.453080.

Jaron KS, Parker DJ, Anselmetti Y, Van PT, Bast J, Dumas Z, Figuet E, François CM, Hayward
K, Rossier V, et al. 2021. Convergent consequences of parthenogenesis on stick insect
genomes. *bioRxiv*:2020.11.20.391540.

Jeffries DL, Lavanchy G, Sermier R, Sredl MJ, Miura I, Borzée A, Barrow LN, Canestrelli D,
Crochet P-A, Dufresnes C, et al. 2018. A rapid rate of sex-chromosome turnover and non-
random transitions in true frogs. *Nat. Commun.* 9:4088.

Jennings WB, Edwards SV. 2005. Speciational history of Australian grass finches (Poephila)
inferred from thirty gene trees. *Evolution* 59:2033–2047.

Jiang X, Biedler JK, Qi Y, Hall AB, Tu Z. 2015. Complete dosage compensation in *Anopheles*
*stephensi* and the evolution of sex-biased genes in mosquitoes. *Genome Biol. Evol.*
7:1914–1924.

Keightley PD, Ness RW, Halligan DL, Haddrill PR. 2014. Estimation of the spontaneous
mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib family. *Genetics*
196:313–320.

Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, Davey JW,
Jiggins CD. 2015. Estimation of the spontaneous mutation rate in *Heliconius melpomene*.
*Mol. Biol. Evol.* 32:239–243.

Kelly WG, Schaner CE, Dernburg AF, Lee M-H, Kim SK, Villeneuve AM, Reinke V. 2002. X-
chromosome silencing in the germline of *C. elegans*. *Development* 129:479–492.

Khil PP, Smirnova NA, Romanienko PJ, Camerini-Otero RD. 2004. The mouse X chromosome
is enriched for sex-biased genes not subject to selection by meiotic sex chromosome
inactivation. *Nat. Genet.* 36:642–646.

Korneliussen TS, Albrechtsen A, Nielsen R. 2014. ANGSD: Analysis of next generation
sequencing data. *BMC Bioinformatics* 15:356.

Lahn BT, Page DC. 1999. Four evolutionary strata on the human X chromosome. *Science*
286:964–967.

Lee JT. 2005. Sex chromosome inactivation: the importance of pairing. *Curr. Biol.* 15:R249–
R252.

Lenormand T, Fyon F, Sun E, Roze D. 2020. Sex Chromosome Degeneration by Regulatory
Evolution. *Curr. Biol.* 30:3001–3006.e5.

Lenormand T, Roze D. 2021. Y recombination arrest and degeneration in the absence of
sexual dimorphism. *bioRxiv*:2021.05.18.444606.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
*arXiv*:1303.3997.

Lucchesi JC. 1998. Dosage compensation in flies and worms: the ups and downs of X-
chromosome regulation. *Curr. Opin. Genet. Dev.* 8:179–184.

Mahadevaraju S, Fear JM, Akeju M, Galletta BJ, Pinheiro MMLS, Avelino CC, Cabral-de-Mello
DC, Conlon K, Dell'Orso S, Demere Z, et al. 2021. Dynamic sex chromosome expression
in *Drosophila* male germ cells. *Nat. Commun.* 12:892.

Mahajan S, Bachtrog D. 2015. Partial dosage compensation in Strepsiptera, a sister group of
beetles. *Genome Biol. Evol.* 7:591–600.

Mank JE. 2009. The W, X, Y and Z of sex-chromosome dosage compensation. *Trends Genet.*
25:226–233.

Mank JE. 2013. Sex chromosome dosage compensation: definitely not for everyone. *Trends*
*Genet.* 29:677–683.

Mank JE, Ellegren H. 2009. All dosage compensation is local: gene-by-gene regulation of sex-
biased expression on the chicken Z chromosome. *Heredity* 102:312–320.

Mank JE, Hosken DJ, Wedell N. 2014. Conflict on the sex chromosomes: cause, effect, and
complexity. *Cold Spring Harb. Perspect. Biol.* 6:a017715.

Mank JE, Vicoso B, Berlin S, Charlesworth B. 2010. Effective population size and the Faster-
X effect: empirical results and their interpretation. *Evolution* 64:663–674.

Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. 2018. MUMmer4: A
fast and versatile genome alignment system. *PLoS Comput. Biol.* 14:e1005944.

Marín I, Franke A, Bashaw GJ, Baker BS. 1996. The dosage compensation system of
Drosophila is co-opted by newly evolved X chromosomes. *Nature* 383:160–163.

Marín I, Siegal ML, Baker BS. 2000. The evolution of dosage-compensation mechanisms.
*Bioessays* 22:1106–1114.

Marshall Graves JA. 2015. Evolution of vertebrate sex chromosomes and dosage
compensation. *Nat. Rev. Genet.* 17:33–46.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet J* 17:10–12.

Meiklejohn CD, Landeen EL, Cook JM, Kingan SB, Presgraves DC. 2011. Sex chromosome-
specific regulation in the *Drosophila* male germline but little evidence for chromosomal
dosage compensation or meiotic inactivation. *PLoS Biol.* 9:e1001126.

Meisel RP, Connallon T. 2013. The faster-X effect: integrating theory and data. *Trends Genet.*
29:537–544.

Meyer BJ. 2000. Sex in the worm: counting and compensating X-chromosome dose. *Trends*
*Genet.* 16:247–253.

Montgomery SH, Mank JE. 2016. Inferring regulatory change from gene expression: the

confounding effects of tissue scaling. *Mol. Ecol.* 25:5114–5128.

Mullon C, Pomiankowski A, Reuter M. 2012. The effects of selection and genetic drift on the
genomic distribution of sexually antagonistic alleles. *Evolution* 66:3743–3753.

Mullon C, Wright AE, Reuter M, Pomiankowski A, Mank JE. 2015. Evolution of dosage
compensation under sexual selection differs between X and Z chromosomes. *Nat.*
*Commun.* 6:7720.

Nosil P, Villoutreix R, de Carvalho CF, Farkas TE, Soria-Carrasco V, Feder JL, Crespi BJ,
Gompert Z. 2018. Natural selection and the predictability of evolution in *Timema* stick
insects. *Science* 359:765–770.

Nozawa M, Fukuda N, Ikeo K, Gojobori T. 2014. Tissue- and stage-dependent dosage
compensation on the neo-X chromosome in *Drosophila pseudoobscura*. *Mol. Biol. Evol.*
31:614–624.

Nunney L. 1993. The influence of mating system and overlapping generations on effective
population size. *Evolution* 47:1329–1341.

Ohno S. 1967. Sex chromosomes and sex-linked genes. Berlin: Springer Verlag

Oliver B. 2002. Genetic control of germline sexual dimorphism in *Drosophila*. *Int. Rev. Cytol.*
219:1–60.

Pal A, Vicoso B. 2015. The X Chromosome of Hemipteran insects: conservation, dosage
compensation and sex-biased expression. *Genome Biol. Evol.* 7:3259–3268.

Palmer DH, Rogers TF, Dean R, Wright AE. 2019. How to identify sex chromosomes and their
turnover. *Mol. Ecol.* 28:4709–4724.

Parker DJ, Bast J, Jalvingh K, Dumas Z, Robinson-Rechavi M, Schwander T. 2019a. Sex-
biased gene expression is repeatedly masculinized in asexual females. *Nat. Commun.*
10:4638.

Parker DJ, Bast J, Jalvingh K, Dumas Z, Robinson-Rechavi M, Schwander T. 2019b.
Repeated evolution of asexuality involves convergent gene expression changes. *Mol.*
*Biol. Evol.* 36 :350–364.

Parkhurst SM, Meneely PM. 1994. Sex determination and dosage compensation: lessons
from flies and worms. *Science* 264:924–932.

Parsch J, Ellegren H. 2013. The evolutionary causes and consequences of sex-biased gene
expression. *Nat. Rev. Genet.* 14:83–87.

Payseur BA, Presgraves DC, Filatov DA. 2018. Introduction: Sex chromosomes and
speciation. *Mol. Ecol.* 27:3745–3748.

Pinharanda A, Rousselle M, Martin SH, Hanly JJ, Davey JW, Kumar S, Galtier N, Jiggins CD.
2019. Sexually dimorphic gene expression and transcriptome evolution provide mixed
evidence for a fast-Z effect in *Heliconius*. *J. Evol. Biol.* 32:194–204.

Pirrotta V. 2002. Silence in the germ. *Cell* 110:661–664.

Pokorná M, Kratochvíl L. 2009. Phylogeny of sex-determining mechanisms in squamate
reptiles: are sex chromosomes an evolutionary trap? *Zool. J. Linn. Soc.* 156:168–183.

Primmer CR, Borge T, Lindell J, Saetre G-P. 2002. Single-nucleotide polymorphism
characterization in species with limited available sequence information: high nucleotide
diversity revealed in the avian genome. *Mol. Ecol.* 11:603–612.

Prince EG, Kirkland D, Demuth JP. 2010. Hyperexpression of the X chromosome in both
sexes results in extensive female bias of X-linked genes in the flour beetle. *Genome Biol.*
*Evol.* 2:336–346.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* 26:841–842.

Rayner JG, Hitchcock TJ, Bailey NW. 2021. Variable dosage compensation is associated with
female consequences of an X-linked, male-beneficial mutation. *Proc. Biol. Sci.*
288:20210355.

R Core Team. 2017. R: A Language and Environment for Statistical Computing. Available
from: https://www.R-project.org/

Rice WR. 1984. Sex chromosomes and the evolution of sexual dimorphism. *Evolution* 38:735–
742.

Richard G, Legeai F, Prunier-Leterme N, Bretaudeau A, Tagu D, Jaquiéry J, Le Trionnaire G.
2017. Dosage compensation and sex-specific epigenetic landscape of the X chromosome
in the pea aphid. *Epigenetics Chromatin* 10:30.

Riesch R, Muschick M, Lindtke D, Villoutreix R, Comeault AA, Farkas TE, Lucek K, Hellen E,
Soria-Carrasco V, Dennis SR, et al. 2017. Transitions between phases of genomic
differentiation during stick-insect speciation. *Nature Ecology & Evolution* 1:0082.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* 26:139–140.

Rose G, Krzywinska E, Kim J, Revuelta L, Ferretti L, Krzywinski J. 2016. Dosage
compensation in the African Malaria Mosquito *Anopheles gambiae*. *Genome Biol. Evol.*
8:411–425.

Ruzicka F, Connallon T. 2020. Is the X chromosome a hot spot for sexually antagonistic
polymorphisms? Biases in current empirical tests of classical theory. *Proceedings of the*
*Royal Society B: Biological Sciences* 287:20201869.

Ruzicka F, Hill MS, Pennell TM, Flis I, Ingleby FC, Mott R, Fowler K, Morrow EH, Reuter M.
2019. Genome-wide sexually antagonistic variants reveal long-standing constraints on
sexual dimorphism in fruit flies. *PLoS Biol.* 17:e3000244.

Sangrithi MN, Turner JMA. 2018. Mammalian X chromosome dosage compensation:
perspectives from the germ line. *Bioessays* 40:e1800024.

Sayadi A, Martinez Barrio A, Immonen E, Dainat J, Berger D, Tellgren-Roth C, Nystedt B,
Arnqvist G. 2019. The genomic footprint of sexual conflict. *Nat Ecol Evol* 3:1725–1730.

Schwander T, Crespi BJ. 2009. Multiple direct transitions from sexual reproduction to
apomictic parthenogenesis in *Timema* stick insects. *Evolution* 63:84–103.

Shetty S, Griffin DK, Graves JA. 1999. Comparative painting reveals strong chromosome
homology over 80 million years of bird evolution. *Chromosome Res.* 7:289–295.

Simon S, Letsch H, Bank S, Buckley TR, Donath A, Liu S, Machida R, Meusemann K, Misof

B, Podsiadlowski L, et al. 2019. Old World and New World Phasmatodea: phylogenomics resolve the evolutionary history of stick and leaf Insects. *Frontiers in Ecology and Evolution* 7:345.

Straub T, Becker PB. 2011. Transcription modulation chromosome-wide: universal features and principles of dosage compensation in worms and flies. *Curr. Opin. Genet. Dev.* 21:147–153.

Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, Jordan T, Shakir K, Roazen D, Thibault J, et al. 2013. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* 43:11.10.1–11.10.33.

Vibranovski MD. 2014. Meiotic sex chromosome inactivation in *Drosophila*. *J Genomics* 2:104–117.

Vibranovski MD, Zhang Y, Long M. 2009. General gene movement off the X chromosome in the *Drosophila* genus. *Genome Res.* 19:897–903.

Vicoso B. 2019. Molecular and evolutionary dynamics of animal sex-chromosome turnover. *Nat Ecol Evol* 3:1632–1641.

Vicoso B, Bachtrog D. 2009. Progress and prospects toward our understanding of the evolution of dosage compensation. *Chromosome Res.* 17:585–602.

Vicoso B, Bachtrog D. 2015. Numerous transitions of sex chromosomes in Diptera. *PLoS Biol.* 13:e1002078.

Vicoso B, Charlesworth B. 2009. Effective population size and the faster-X effect: an extended model. *Evolution* 63:2413–2426.

Vicoso B, Emerson JJ, Zektser Y, Mahajan S, Bachtrog D. 2013. Comparative sex chromosome genomics in snakes: differentiation, evolutionary strata, and lack of global dosage compensation. *PLoS Biol.* 11:e1001643.

Wang M, Zhao Y, Zhang B. 2015. Efficient test and visualization of multi-set intersections. *Sci. Rep.* 5:16923.

Whittle CA, Kulkarni A, Extavour CG. 2020. Absence of a Faster-X effect in beetles (Tribolium, Coleoptera). *G3: Genes, Genomes, Genetics* 10:1125–1136.

Wilkinson GS, Breden F, Mank JE, Ritchie MG, Higginson AD, Radwan J, Jaquiery J, Salzburger W, Arriero E, Barribeau SM, et al. 2015. The locus of sexual selection: moving sexual selection studies into the post-genomics era. *J. Evol. Biol.* 28:739–755.

Wilson Sayres MA. 2018. Genetic diversity on the sex chromosomes. *Genome Biol. Evol.* 10:1064–1078.

Wright AE, Dean R, Zimmer F, Mank JE. 2016. How to make a sex chromosome. *Nat. Commun.* 7:12087.

Wright S. 1931. Evolution in Mendelian Populations. *Genetics* 16:97–159.

Wright S. 1969. Evolution and the Genetics of Populations, Volume 2: Theory of Gene Frequencies. Chicago, USA: University of Chicago Press

Xu L, Zhou Q. 2020. The female-specific W chromosomes of birds have conserved gene

947       contents but are not feminized. *Genes* 11:1126.

948   Zhang Y, Sturgill D, Parisi M, Kumar S, Oliver B. 2007. Constraint and turnover in sex-biased
949       gene expression in the genus *Drosophila*. *Nature* 450:233–237.

950
951