Similarities Between Bacterial GAD and Human GAD65:

Implications in Gut Mediated Autoimmune Type 1 Diabetes

Suhana Bedi[1*], Tiffany M. Richardson[2*], Baofeng Jia[3*], Hadeel Saab[4], Fiona S.L. Brinkman[3],

Monica Westley[5#]

[1]Department of Natural Sciences and Mathematics, the University of Texas at Dallas, USA

[2]Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, USA

[3]Department of Molecular Biology and Biochemistry, Simon Fraser University, Burnaby, CA

[4]Intern, the(sugar)science

[5]Founder, the(sugar)science

[*]authors contributed equally to this work

[#]corresponding author

Corresponding author: Monica Westley Ph.D., the(sugar)science, monica@thesugarscience.org

Keywords: GAD, GAD65, microbiome, beta cell, type 1 diabetes, in silico.

Abstract

A variety of islet autoantibodies (AAbs) can predict and possibly dictate eventual type 1 diabetes

(T1D) diagnosis. Upwards of 75% of those with T1D are positive for AAbs against glutamic acid

decarboxylase (GAD65), a producer of gamma-aminobutyric acid (GABA) in human pancreatic

beta cells. Interestingly, bacterial populations within the human gut also express GAD65 and

produce GABA. Evidence suggests that dysbiosis of the microbiome may correlate with T1D

pathogenesis and physiology. Therefore, autoimmune linkages between the gut microbiome and

islets susceptible to autoimmune attack need to be further elucidated. Utilizing *silico* analyses, we

show here that 25 GAD sequences from different human gut bacterial sources show sequence and

motif similarities to human beta cell GAD65. Our motif analyses determined that a majority of gut

GAD sequences contain the pyroxical dependent decarboxylase domain of human GAD65 which

is important for its enzymatic activity. Additionally, we showed overlap with known human

GAD65 T-cell receptor epitopes which may implicate the immune destruction of beta cells. Thus,

we propose a physiological hypothesis in which changes in the gut microbiome in those with T1D

result in a release of bacterial GAD, thus causing miseducation of the host immune system. Due

to the notable similarities, we found between humans and bacterial GAD, these deputized immune

cells may then go on to target human beta cells leading to the development of T1D.

Introduction

Glutamic acid decarboxylase (GAD65) is a prominent autoantibody in type 1 diabetes (T1D). For those diagnosed as a teen, in addition to having an HLA DR3/DR2 signature, it is likely to be the first autoantibody clinically detected in the blood. GAD65 antibodies are prevalent in the majority of T1D cases, occurring in over 70% of patients [1]. Autoantibodies directed against GAD65 recognize linear and conformational epitopes throughout the protein. The functional region of GAD65 contains a catalytic group involving several residues that are distantly located in the primary amino acid sequence of the protein. The enzymatic function of GAD65 and subsequent regulation of GABA production is dependent on a key residue within the catalytic loop, Lys396 [2]. This residue covalently binds pyridoxal 5'-phosphate (PLP), a co-factor that is essential for the production of GABA by GAD65. Thus, T1D-related immune recognition of GAD65 is highly dependent on the exposure of these regions to autoreactive antigen-presenting cells (APCs). These activated APCs can then go on to attack GAD65-expressing pancreatic beta cells leading to their destruction and eventual T1D diagnosis.

In addition to beta cells, a multitude of bacteria within the gut express GAD (bacterial nomenclature for human GAD65). While there is great diversity within the gut microbiota, more than 95% of gut bacterial species fall into four major microbial phyla: Firmicutes, Bacteroidetes, Actinobacteria, and Protecteobacteria [3]. Bifidobacterium adolescentis, in particular, is an important gut bacterial GABA producer and has the highest prevalence of gad genes in their genomes [4]. These species provide a homeostatic environment that fosters and facilitates gut epithelial integrity and immunity, along with providing energy. Varied literature supports both a decrease in microbial diversity and an increase in a "leaky gut" phenotype throughout T1D [5].

GABA producing Bifidobacterium bacterial counts drop as the Firmicutes/Bacteroides (F/B) ratios

decrease in T1D [6]. There is also a reduction in the abundance of Clostridium clusters IV and

XIVa and mucin-degrading bacteria such as Prevotella and Akkermansia. An important metabolite

produced by these symbiotic bacterial populations, butyrate, supports gut integrity and has beta

cell-specific anti-inflammatory effects [7, 8, 9, 10, 11]. The reduction of these microbes and their

byproducts may be a contributing factor to why children at risk for T1D have higher intestinal

permeability and T1D-associated gut dysbiosis.

One hypothesis for the interrelationship between dysbiosis and T1D is that as these bacterial

populations die, they release GAD65 mimetics that trigger the immune system. Presumably, as

GABA-producing bacteria like B. adolescentis die, either from antibiotic misuse or other

pathology, immune cells located in Peyer's patches and deep in the enterocyte border detect the

inappropriate presence of GAD. Thus, APCs can act to further inform CD8+ T cells in nearby

lymph nodes. The lymphatic pathways between the intestinal submucosa and pancreatic lymph

nodes may provide a route for these "miseducated" CD8+ T cells to gain access to GAD65+ beta

cells. GAD65 may also originate from beta cells leading to an increase in epitope spreading and a

misguided immune attack on these insulin-producing cells.

To initiate the exploration of this hypothesis, we utilized in silico methodologies to compare gut

microbial species that decline at the onset of T1D and express GAD, with GAD65 sequences in

pancreatic beta cells. We show that there is notable sequence similarity, including relevant motif

and T-cell epitope overlaps between several bacterial species and beta cells particularly in the

functionally important pyroxical dependent decarboxylase domain of GAD65. This in silico

exercise suggests that further investigation is warranted of the relationship between microbial

disappearance and appearance of GAD65 autoantibodies during the prodrome and presentation of

T1D.

Materials and Methods

GAD Sequence and T cell receptor epitope curation

Twenty-five bacterial GAD proteins from species of interest, alongside those of mouse,

chimpanzee, and human GAD65 sequences were curated from the NCBI genes database (Table

1). The bacterial species were chosen by considering their association with the gut microbiome,

their significance in the gut microbiome alterations seen in T1D patients, human-related pathogen

status, and non-human-related environmental species. GAD65 T cell receptor epitopes were

curated from a literature review.

All GAD sequences in bacteria and the three mammalian species under consideration were

obtained by performing a BLASTP search, with human GAD65 as the query sequence.

Furthermore, only those sequences with significant e and p values were well annotated on the

NCBI database and/or RefSeq and were identified as GAD proteins were considered. All other

hits, including unnamed protein products, were ignored.

Multiple Sequence Alignment and Motif Identification

Multiple sequence alignment was performed using ClustalW (Ver. 1.2.2) with a gap opening

penalty of 10 and a gap extension penalty of 0.2 [12]. The resulting alignment is visualized and

annotated with CLC Sequence Viewer (Ver. 8.0) to highlight conservation and amino acid

properties. Identification of ungapped motifs was done using the MEME tool (Ver. 5.4.1) under

the MEME suite tools for motif discovery and enrichment [13]. The output consisted of 10 motifs,

shared by various (not necessarily all) organisms into consideration. Subsequently, each motif was

run against protein families, using a Pfam (Ver 33.1) search. The Motifs identified by MEME were

mapped against the query sequences using MAST(Ver. 5.4.1), to obtain the coordinates of motifs,

in each gad sequence [13].

Logo plot & T Cell Receptor Epitopes

The multiple sequence alignment obtained was converted into a logo plot, using Weblogo (Ver

2.8.2) [14]. Here, each logo consists of a stack of symbols, and each symbol's height is indicative

of the amino acid conservation, at a particular location. All motifs and T-cell receptor epitope

coordinates, obtained in the previous steps, were marked on the logo plot. The plot was then

trimmed, with a special emphasis on regions with high overlap between motifs and T-cell receptor

epitopes.

Results

Multiple sequence alignment highlights several conserved residues between human and

bacterial GAD

A total of 25 GAD protein sequences, including 3 mammalian GAD65 (human, chimpanzee, and

mouse) and 22 bacterial GAD, were aligned (Table 1 & Fig. 1). The 3 mammalian GAD65

exhibited high conservation with each other, while 20 out of the 22 bacteria GAD exhibited high

similarity to one another. Two bacteria, *Bdellobibrio sp. Qaytius* and *Sediminibacterium spp.* did

not exhibit any protein conservation. This may be due to a misannotation within the RefSeq

genome. We also found interesting similarities and differences between the mammalian and

bacterial GAD sequences. There were two regions of interest: AA389-397 and AA418-425.

AA389-397 contained several charged residues interspaced with hydrophobic residues, potentially

a conserved alpha-helix structure. AA418-AA425 contains conserved HK residues. These two

residues represent a known pyridoxal phosphate-binding site in *E.coli* GAD (H275 and K276)*,*

that is likely conserved in mammalian GAD65. The length of mammalian GAD65 was ~120 amino

acids longer than bacterial GAD. Based on our alignment, this difference is mostly centered around

the N-terminus, which suggests adaptations and inclusions of signal peptides required for complex

cellular trafficking in mammalian cells. Also, we identified a few small gaps within the core

regions of GAD between species, these gaps are unlikely of any biological significance.

Table 1. GABA producing bacterial species and their respective GAD gene

Category	Species	Gene ID
Human	Human	2572
Primate	Chimpanzee	466026
Mouse	Mouse	14417
T1D	Streptococcus pneumoniae	Z49109.1
T1D	Bifidobacterium moukalabense	61040350
T1D	Bifidobacterium adolescentis	56675515
T1D	Bacteroides fragilis	61595025
T1D	Ruminococcus	Genome not found
T1D	Veillonella	Genome not found
T1D	Blautia	Genome not found
T1D	Lachnospira	Genome not found
T1D	Alistipes obesi	Genome not found
T1D	Clostridium spp.	29570578
T1D	Prevotella	Genome not found
T1D	Parabacteroides distasonis	CP040468
Gut	Escherichia coli	946058
Gut	Listeria monocytogenes	985123
Gut	Salmonella enterica	13913147
Gut	Streptomyces sp. NRRL B-24085	6211654
Gut	Carnobacterium maltaromaticum	56848084
Gut	Nocardia brasiliensis	13733698
Gut	Lactobacillaceae	56991876
Pathogen	Gammaproteobacteria	QNEL01000191.1
Pathogen	Klebsiella pneumoniae IS22	12545741
Environment	Planctomycetes bacterium	KAA3606119.1
Environment	Sediminibacterium sp.	JADMJG010000006.1
Environment	Moorea producens	CP017599.1
Environment	Magnetococcales bacterium	JADGBX010000054.1
Environment	Scytonema hofmannii	ANNX02000047.1
Environment	Candidatus Aminicenantes bacterium	WVZL01000014.1
Environment	Bdellovibrio sp. qaytius	CP025734.1
Environment	Verrucomicrobia	NZEQ01000004.1

Motif analysis uncovers regional similarities of human and bacterial GAD enzymatic regions

Our motif analysis found 10 ungapped motifs with each motif present in at least 10 of the sequences under consideration (Fig. 2). Even though multiple common motifs between human GAD65 and bacterial GAD can be observed, it is noteworthy that two motif blocks are conserved all across the organisms considered, which is indicative of the sequence similarity observed in the multiple sequence alignment. In addition, except for *Streptococcus pneumoniae,* one motif block was consistently present in all sequences. Furthermore, these motifs were queried on the Pfam and the CDD databases, and all 10 motifs were identified as matches to the pyridoxal-dependent decarboxylase (PDP) conserved domain, which is instrumental in the biosynthesis of GABA, from glutamate. This indicates that human GAD65 and bacterial GAD share sequence similarities in a functional region of GAD.

Overlap exists between T cell receptor epitopes of GAD65 and bacterial GAD

To compare the overlap between T cell receptor epitopes, human GAD65, and bacterial GAD, a logo plot of the multiple sequence alignment was made using three representative bacterial GAD and three animal GAD65 proteins. Motif positions for each sequence were manually annotated using the motif coordinates obtained in motif discovery and enrichment analysis. Thereafter, the logo plot was annotated with T cell receptor epitopes and the overlapping regions were observed. The fine lines indicate the motif coordinates, while the red blocks indicate the CD8+ epitopes and blue blocks indicate the CD4+ epitopes. Three regions overlapped with a motif, which as stated in the motif analysis result description, hit the PDP domain (Fig. 3). This was also confirmed by a CDD search. Of the amino acids in which both CD4 and CD8 receptor epitopes overlap, there is at least one amino acid residue that is conserved across bacterial GAD and GAD65 (Fig. 4). Such

overlaps emphasize a possible explanation for how T cells could recognize both bacterial and

human GAD65.

Discussion

In this study, we explored the sequence similarities between GAD in twenty GABA-producing

bacteria and GAD65 in humans. We found that human and bacterial GAD have similar motifs and

conserved residues. We identified notable conservation centered around the pyridoxal dependent

decarboxylase domain of GAD65 found in pancreatic beta cells, particularly around two key

substrate-binding residues. Moreover, some conserved sequences in bacterial GAD overlap with

known human GAD65 T cell receptor epitopes. As one example, GAD65-specific HLA-DR4

(DRB1*0401)–restricted murine T-cell hybridoma line T33.1 recognizes the GAD65 274-286

epitope, and this epitope was conserved across bacteria and human samples examined here (Figure

1). This *in silico* proof-of-concept suggests a possible relationship between disappearance of GAD

containing GABA-producing bacteria in the human gut microbiome and the appearance of GAD65

autoantibodies during the onset of T1D that needs to be further elucidated.

It is important to note a few limitations within our analyses. Many gut microbial genomes and

genome annotations were incomplete, particularly for some GABA-producing bacterial species of

importance that are significantly changed in T1D. We did not include several species due to

missing GAD genome annotations. Together, only 20 sequences were analyzed for their

homology, with many belonging to environmental organisms. A larger cohort would further

strengthen our sequence analyses and hypotheses. Furthermore, we hypothesized that the change

in anti-GAD65 is partly due to the changes in the gut microbiome. Additional studies are needed

to determine what insult(s) would cause such a disturbance and what cellular mechanisms are

involved in the miseducation of immune cells due to the release of GAD in the gut. The window

for the disappearance of the *gad* containing bacterial species should be identified as to whether it

precedes or follows the appearance of autoantibodies to GAD65 in humans. Lastly, the analysis

relied on known and curated T cell epitopes of GAD65 from the literature review, in which we

demonstrated similarity to bacterial GAD. Whether or not the bacterial peptides bind to human T

cell receptors needs to be further verified *in vitro.*

We postulate that bacterial GAD found in GABA producing bacteria may act as an antigen to

activate submucosal T cells as a result of gut microbiome bacterial destruction, either viral or

antibiotic induced. Further, we posit these primed bacterial GAD T cells bind GAD65 in the islet

environment due to the GAD and GAD65 epitope similarity. That leaves us to explore how

GAD65, an intracellular enzyme found in alpha and beta islet cells, might get recognized by

bacterial derived anti-GAD T cells.

In beta cells, GAD65 is intracellular and interacts with the cytosolic leaflet of Golgi membranes

and secretory vesicles due to its post-translational palmitoylation (lipidation) at cysteines 30 and

45 [15]. GAD65's shuttling to the Golgi for palmitoylation requires a signal located between

residues 1-27 of the protein [16, 17]. GAD65 may transiently interact with the cell's cytosolic

leaflet upon exocytosis of beta cell granules [18]. Pre-clinical studies demonstrated that GAD65

can associate with the plasma membrane and enter the extracellular space [19].

GAD65 is certainly present in alpha and beta cells and may interact with a GAD T cell from the

gut at the plasma membrane of both islet cell types, but there may be another route of contact. A

recent review highlights how beta cells can become "deputized" and act as antigen-presenting cells

(APCs) under certain conditions [20] . In this scenario, "deputized" beta cells that present GAD65

might have an opportunity to interact with a GAD sensitized T cell. Endothelial cells in the highly

vascularized islet environment have been named as additional "deputized" APCs. These

endothelial cells may be exposed to transient plasma membrane resident GAD65, or GAD65

containing EVs from stressed pancreatic beta cells [21, 22] and act to present them to anti-GAD

gut derived T cells.

A "two-hit" scenario is one where bacterial GAD primed T cells recognize alpha and beta cell

derived GAD65 to impact both intra islet cell interaction as well as simultaneously disrupting local

immune response. In islet α-cells, GABA induces membrane hyperpolarization and suppresses

glucagon secretion, while in islet β-cells GABA induces membrane depolarization and increases

insulin secretion [23]. One consequence of a decrease of GAD65 in the islet environment is the

resultant decrease in GABA production resulting in hyperglycemia which is toxic to beta cells.

GABA is also an immune cell modulator, shown to increase Tregs, decrease Th1 and CTLs

fostering a tolerant immune environment [23]. A GAD65 depletion leading to a GABA decrease

in the islet could pack a double punch of generating dysfunctional alpha and beta cells operating

in a more highly reactive immune micro-environment.

One can imagine a scenario where GAD primed T cells from the gut, become initially alerted to

the islet environment as GAD65 is presented via deputized alpha, beta, endothelial cells or some

combination of the three. Once alerted, GAD primed T cells destroy beta or alpha cells that display

GAD65 at the plasma membrane, reducing the overall GABA levels in the islets with resulting

consequences of hyperglycemia which destabilizes beta cells, in addition to altered immune

phenotype that favors attack of the islets.

Taken together, our *in silico* analyses of genome sequences and motifs from bacterial, human, and

T cell epitopes identified several homologies that may inform further experimentation and inquiry

into the relationship between T1D pathogenesis and microbiome dysbiosis.

References

[1] Petersen JS, Hejnaes KR, Moody A, et al. Detection of GAD65 antibodies in diabetes and other

autoimmune diseases using a simple radioligand assay. Diabetes. 1994;43(3):459-467.

doi:10.2337/diab.43.3.459

[2] Fenalti G, Law RH, Buckle AM, et al. GABA production by glutamic acid decarboxylase is

regulated by a dynamic catalytic loop. Nat Struct Mol Biol. 2007;14(4):280-286.

doi:10.1038/nsmb1228

[3] Sommer F, Bäckhed F. The gut microbiota--masters of host development and physiology. Nat

Rev Microbiol. 2013;11(4):227-238. doi:10.1038/nrmicro2974

[4] Duranti S, Ruiz L, Lugli GA, et al. Bifidobacterium adolescentis as a key member of the human

gut microbiota in the production of GABA. Sci Rep. 2020;10(1):14112. Published 2020 Aug 24.

doi:10.1038/s41598-020-70986-z

[5] Zhou H, Sun L, Zhang S, Zhao X, Gang X, Wang G. Evaluating the Causal Role of Gut

Microbiota in Type 1 Diabetes and Its Possible Pathogenic Mechanisms. Front Endocrinol

(Lausanne). 2020;11:125. Published 2020 Mar 24. doi:10.3389/fendo.2020.00125

[6] Murri M, Leiva I, Gomez-Zumaquero JM, et al. Gut microbiota in children with type 1 diabetes

differs from that in healthy children: a case-control study. BMC Med. 2013;11:46. Published 2013

Feb 21. doi:10.1186/1741-7015-11-46

[7] de Goffau MC, Fuentes S, van den Bogert B, et al. Aberrant gut microbiota composition at the

onset of type 1 diabetes in young children. Diabetologia. 2014;57(8):1569-1577.

doi:10.1007/s00125-014-3274-0

[8] Prause M, Pedersen SS, Tsonkova V, Qiao M, Billestrup N. Butyrate Protects Pancreatic Beta

Cells from Cytokine-Induced Dysfunction. Int J Mol Sci. 2021;22(19):10427. Published 2021 Sep

27. doi:10.3390/ijms221910427

[9] Leiva-Gea I, Sánchez-Alcoholado L, Martín-Tejedor B, et al. Gut Microbiota Differs in

Composition and Functionality Between Children With Type 1 Diabetes and MODY2 and Healthy

Control Subjects: A Case-Control Study. Diabetes Care. 2018;41(11):2385-2395.

doi:10.2337/dc18-0253

[10] Harbison JE, Roth-Schulze AJ, Giles LC, et al. Gut microbiome dysbiosis and increased

intestinal permeability in children with islet autoimmunity and type 1 diabetes: A prospective

cohort study. Pediatr Diabetes. 2019;20(5):574-583. doi:10.1111/pedi.12865

[11] Soyucen E, Gulcan A, Aktuglu-Zeybek AC, Onal H, Kiykim E, Aydin A. Differences in the

gut microbiota of healthy children and those with type 1 diabetes. Pediatr Int. 2014;56(3):336-343.

doi:10.1111/ped.12243

[12] Thompson, J D et al. "CLUSTAL W: improving the sensitivity of progressive multiple

sequence alignment through sequence weighting, position-specific gap penalties and weight

matrix choice." Nucleic acids research vol. 22,22 (1994): 4673-80. doi:10.1093/nar/22.22.4673

[13] Timothy L. Bailey, James Johnson, Charles E. Grant, William S. Noble, "The MEME Suite",

Nucleic Acids Research, 43(W1):W39-W49, 2015.

[14] Crooks, Gavin E et al. "WebLogo: a sequence logo generator." Genome research vol. 14,6

(2004): 1188-90. doi:10.1101/gr.849004

[15] Kanaani J, Diacovo MJ, El-Husseini Ael-D, Bredt DS, Baekkeskov S. Palmitoylation controls

trafficking of GAD65 from Golgi membranes to axon-specific endosomes and a Rab5a-dependent

pathway to presynaptic clusters. J Cell Sci. 2004;117(Pt 10):2001-2013. doi:10.1242/jcs.01030

[16] Solimena M, Dirkx R Jr, Radzynski M, Mundigl O, De Camilli P. A signal located within

amino acids 1-27 of GAD65 is required for its targeting to the Golgi complex region. J Cell Biol.

1994;126(2):331-341. doi:10.1083/jcb.126.2.331

[17] Christgau S, Aanstoot HJ, Schierbeck H, et al. Membrane anchoring of the autoantigen

GAD65 to microvesicles in pancreatic beta-cells by palmitoylation in the NH2-terminal domain.

J Cell Biol. 1992;118(2):309-320. doi:10.1083/jcb.118.2.309

[18] Kanaani J, el-Husseini Ael-D, Aguilera-Moreno A, Diacovo JM, Bredt DS, Baekkeskov S. A

combination of three distinct trafficking signals mediates axonal targeting and presynaptic

clustering of GAD65. J Cell Biol. 2002;158(7):1229-1238. doi:10.1083/jcb.200205053

[19] Jin H, Wu H, Osterhaus G, et al. Demonstration of functional coupling between gamma -

aminobutyric acid (GABA) synthesis and vesicular GABA transport into synaptic vesicles. Proc

Natl Acad Sci U S A. 2003;100(7):4293-4298. doi:10.1073/pnas.0730698100

[20] Li Y, Sun F, Yue TT, et al. Revisiting the Antigen-Presenting Function of β Cells in T1D

Pathogenesis. Front Immunol. 2021;12:690783. Published 2021 Jul 14.

doi:10.3389/fimmu.2021.690783

[21] Cianciaruso C, Phelps EA, Pasquier M, et al. Primary Human and Rat β-Cells Release the

Intracellular Autoantigens GAD65, IA-2, and Proinsulin in Exosomes Together With Cytokine-

Induced Enhancers of Immunity. Diabetes. 2017;66(2):460-473. doi:10.2337/db16-0671

[22] Greening JE, Tree TI, Kotowicz KT, et al. Processing and presentation of the islet autoantigen

GAD by vascular endothelial cells promotes transmigration of autoreactive T-cells. Diabetes.

2003;52(3):717-725. doi:10.2337/diabetes.52.3.717

[23] Wan Y, Wang Q, Prud'homme GJ. GABAergic system in the endocrine pancreas: a new target

for diabetes treatment. Diabetes Metab Syndr Obes. 2015;8:79-87. Published 2015 Feb 3.

doi:10.2147/DMSO.S50642

Acknowledgments

We wish to express our gratitude to the entire volunteer team at the(sugar)science and to our

donors and sponsors that help us continue our mission of helping scientists who study Type 1

diabetes connect, collaborate, and accelerate research.

Funding

the(sugar)science is an all-volunteer 501c3 non-profit organization. B.J. was supported by a

Canadian Institutes of Health Research (CIHR) doctoral scholarship. T.M.R. was supported by a

National Science Foundation Graduate Research Fellowship, and F.S.L.B. has a Simon Fraser

University distinguished professorship and support from Natural Sciences and Engineering

Research Council of Canada (NSERC).

Author Contributions

M.W. devised the original experimentation and acquired funding; M.W., S.B., and B.J.

contributed to the conceptualization; S.B. and B.J. performed the investigation and formal

analysis; M.W., F.S.L.B, S.B., B.J., and T.M.R. interpreted the results; M.W., T.M.R., B.J., S.B.,

and H.S. wrote the original draft; all authors reviewed and edited the manuscript.

Figure 1. Multiple sequence alignment of twenty bacterial GAD protein and three animal

GAD65 protein highlights conserved residues. Across a diverse number of bacterial species

and 3 animal species, 12 amino acids were found to be 100% conserved (shaded in blue). Amino

acids 1-110 are likely signal peptides required for eukaryotic protein translation and cellular

transport not needed in bacteria. H275 and K276 are key residues required for PDP binding and

are conserved in human GAD65 and bacterial GAD.

Figure 2. Motif Discovery Analysis of twenty bacterial GAD protein and three animal

GAD65 proteins to identify motifs conserved across various species and their functional

relevance with GAD protein. Across all 23 GAD sequences under consideration, 10 motifs

were each found in at least 10 of the sequences used for motif discovery. Each motif appears in a

separate color below with the individual sequences described as well. In addition, 2 of the 10

motifs were found in all of the sequences. Each motif was then queried for, using Pfam and CDD

searches and it was found that all 10 of the motifs, hit the PDP domain in the GAD65 protein.

All motifs match the Pyridoxal-dependent decarboxylase (PDP) conserved domain

Name	p-value	Motif Locations
Homo_sapiens	1.26e-189	
Pan_troglodytes	8.29e-190	
Mus_musculus	3.17e-189	
Streptococcus_pneumoniae	2.00e-36	
1_Bifidobacterium_moukalabense	1.37e-231	
2_Bifidobacterium_adolescentis	2.20e-228	
Escherichia_coli_str._K-12_substr._MG1655	2.35e-195	
Listeria_monocytogenes	2.59e-228	
Salmonella_enterica	6.08e-51	
Streptomyces	1.64e-205	
Carnobacterium_maltaromaticum	3.30e-224	
Nocardia_brasiliensis	2.75e-200	
Levilactobacillus_brevis	4.91e-182	
Bacteroides	2.47e-181	
Clostridium_perfringens	2.04e-233	
Gammaproteobacteria_bacterium	1.30e-163	
Klebsiella_pneumoniae	4.94e-107	
Planctomycetes_bacterium	3.74e-149	
Moorea_producens_PAL-8-15-08-1	1.64e-172	
Magnetococcales_bacterium	7.41e-163	
Scytonema_hofmannii_PCC_7110	5.53e-169	
Candidatus_Aminicenantes_bacterium	1.68e-165	
Verrucomicrobiales_bacterium	4.88e-144	

Motif	Symbol	Motif Consensus
1.		HVDAASGGMFAPFIEPDLVWDFRLKRVVSINTSGHKYGLVYPGLGWVVWR
2.		KYKJWLHVDAAWGGSLLFSRKHKHLLDGSELADSVTWBPHKMMGVPLICS
3.		VDENTIGVVATAGTTVTGAYDPIKAJADI
4.		GTSTTGGSEANMLAGLAARFR
5.		TFCTTWMDDEATKLMAEALDKNAIDKDEYPQTAELESRCVNMJADLWHAP
6.		LPEELIFKVSYLGGEMPTFAJNFSRPAAQIIGQYYNFLRLGFEGYRRVQQ
7.		DDEWTLYDLGDRLLQCGRRVDAYKLWLSW
8.		YLKYSVRTGHPQFFNQLFGGFDLVGLLAEWLTSTTNTSMYTYEIAPVATL
9.		QVCWEKFCRYWDVEMREVPMSEEHLSLDPDEVLDY
10.		RLVAFVSEQAHYSFKKAANLLGIGTDNVIKVKTDERGRMIPSDLEAAILZ

Figure 3. Conserved regions of GAD share common CD8+ and CD4+ T cell receptor

epitopes. A total of five regions of interest were identified by overlapping T cell receptor

epitopes to the sequence logo of three representative bacterial GAD and three animal GAD65

proteins. Three regions contained a motif match to Group II pyridoxal-dependent

decarboxylases. Of the amino acids in which both CD4+ and CD8+ receptor epitopes overlap,

there is at least 1 amino acid residue that is conserved across bacterial GAD and GAD65. Red is

CD8+ epitopes and Blue is CD4+ epitopes.

Figure 4. Miseducation of antigen-presenting cells (APCs) by bacterial GAD may

contribute to pancreatic beta cell immune-mediated destruction. The death of GABA-

producing bacteria at the onset of T1D may release bacterial GAD65 which could lead to the

miseducation of APCs. Therefore, this miseducation could provoke the immune system to attack

and produce antibodies targeting human GAD65 expressing beta cells because of the similarity

and conservation between bacterial GAD and human GAD65. The pancreatic lymphatic system

could provide a pathway for these miseducated APCs to traverse from the duodenum into

pancreatic islets.