Non-uniform distribution of dendritic nonlinearities differentially engages thalamostriatal and corticostriatal inputs onto cholinergic interneurons

Abbreviated title: Dendritic nonlinearities in striatal cholinergic interneurons

Osnat Oz[1], Lior Matityahu[1], Aviv Mizrahi-Kliger[1,2], Alexander Kaplan[1,2], Noa Berkowitz[1], Lior Tiroshi[1†], Hagai Bergman[1,2,3], Joshua A. Goldberg[1*]

[1]Department of Medical Neurobiology, Institute of Medical Research Israel – Canada, The Faculty of Medicine,
[2]Edmond and Lily Safra Center for Brain Sciences,
The Hebrew University of Jerusalem, 9112102 Jerusalem, Israel
[3]Department of Neurosurgery, Hadassah Medical Center, 9112102 Jerusalem, Israel

[†]Current address: Department of Neurobiology, Northwestern University, Evanston, IL, 60208, USA

[*]Corresponding author, joshua.goldberg2@mail.huji.ac.il

Abstract

The tonic activity of striatal cholinergic interneurons (CINs) is modified differentially by their afferent inputs. Although their unitary synaptic currents are identical, in most CINs cortical inputs onto distal dendrites only weakly entrain them, whereas proximal thalamic inputs trigger abrupt pauses in discharge in response to salient external stimuli. To test whether the dendritic expression of the active conductances that drive autonomous discharge contribute to the CINs' capacity to dissociate cortical from thalamic inputs, we used an optogenetics-based method to quantify dendritic excitability. We found that the persistent sodium (NaP) current gave rise to dendritic boosting, and that the hyperpolarization-activated cyclic nucleotide-gated (HCN) current gave rise to a subhertz membrane resonance. This resonance may underlie our novel finding of an association between CIN pauses and internally-generated slow wave events in sleeping non-human primates. Moreover, our method indicated that dendritic NaP and HCN currents were preferentially expressed in proximal dendrites. We validated the non-uniform distribution of NaP currents pharmacologically; with two-photon imaging of dendritic back-propagating action potentials; and by demonstrating boosting of thalamic, but not cortical, inputs by NaP currents. Thus, the localization of active dendritic conductances in CIN dendrites mirrors the spatial distribution of afferent terminals and may promote their differential responses to thalamic *vs.* cortical inputs.

Introduction

The striatal cholinergic interneuron (CIN) is a key modulator of the striatal microcircuitry, impacting the neuronal excitability, synaptic transmission and synaptic plasticity of spiny projection neurons (SPNs), as well as other striatal interneurons (Abudukeyoumu et al., 2019; Assous, 2021; Goldberg et al., 2012; Matityahu et al., 2022). There are two processes that drive the ongoing release of acetylcholine (ACh) by CINs. First, the two glutamatergic inputs arising from cerebral cortex and intralaminar nuclei of the thalamus can drive CINs to discharge (Bradfield et al., 2013; Ding et al., 2010; Doig et al., 2014; Kosillo et al., 2016; Lapper and Bolam, 1992; Mamaligas et al., 2019; Matsumoto et al., 2001; Sharott et al., 2012; Thomas et al., 2000). While the unitary synaptic currents generated by these two inputs in CINs are identical (Aceves Buendia et al., 2019), thalamic inputs to CINs dominate in the sense that they that give rise to larger excitatory post-synaptic potentials (EPSPs) in acute striatal slices (Johansson and Silberberg, 2020) and can trigger abrupt pause responses, often flanked by excitatory peaks, to external saliency-related cues (Apicella et al., 1991; Goldberg and Reynolds, 2011; Graybiel et al., 1994; Kimura et al., 1984; Matsumoto et al., 2001; Morris et al., 2004; Raz et al., 1996). In contrast, cortical inputs to CINs are weaker in that, in acute striatal slices, they give rise to smaller EPSPs in most CINs (Mamaligas et al., 2019) and cannot trigger the pause response (Ding et al., 2010), although, these differences are less pronounced in intact animals (Doig et al., 2014).

Second, even in the absence of afferent input, CINs exhibit multiple autonomously generated discharge patterns – including regular and irregular pacemaking, as well as burst firing (Bennett et al., 2000; Bennett and Wilson, 1999; Goldberg and Reynolds, 2011; Goldberg and Wilson, 2010). These firing patterns are generated by an interplay between various nonlinear ionic currents including voltage- and calcium-activated K^+ currents, as well as two voltage-dependent pacemaker currents: the hyperpolarization-activated cyclic nucleotide-gated (HCN) current; and the persistent Na^+ (NaP) current (Bennett et al., 2000; Deng et al., 2007; Goldberg et al., 2009; Goldberg and Wilson, 2010, 2005; McGuirt et al., 2021; Oswald et al., 2009; Song and Surmeier, 1996; Wilson, 2005; Wilson and Goldberg, 2006). The main purpose of these pacemaker currents presumably is to guarantee the ongoing release of ACh onto the striatal microcircuitry by sustaining the autonomous discharge of CINs. Nevertheless, these pacemaker (and other subthreshold) currents will also impact how cortical and thalamic inputs are integrated by the CINs.

Attaining a mechanistic (e.g., dynamical systems) understanding of how the repertoire

of CIN firing patterns is generated, requires a full characterization of the nonlinear properties

of the pacemaker (and other) currents (e.g., by determining their voltage dependence and

kinetics) – which is a daunting endeavor. In contrast, understanding how these currents

impact synaptic inputs is a simpler task. Because individual synaptic inputs are small, the

membrane nonlinearities can be *linearized* making the analysis of their impact simpler and

more general – a treatment called the quasi-linear membrane approximation. This analysis

dates back to Mauro (Koch, 1984; Mauro et al., 1970) and has shown that quasi-linear

membranes can give rise to two qualitatively different transformations of inputs:

amplification and *resonance* (Goldberg et al., 2007; Hutcheon and Yarom, 2000). Being a

linear approximation, the quasi-linear approximation is amenable to Fourier analysis, which

helps to better define these transformations as linear (time-invariant) filters on the input in

frequency space.

Amplification arises from regenerative ionic currents – that provide positive feedback

– including inward (depolarizing) currents activated by depolarization, such as the NaP

current. Here the main effect in frequency space is amplification of the amplitude response

(as compared to the response of a passive linear membrane). Resonance arises from

restorative currents – that provide negative feedback – including inward currents activated by

hyperpolarization, such the HCN current. In frequency space, the defining properties of

resonance is a peak (at a non-zero frequency) in the amplitude response, and a zero-crossing

of the phase delay (at a nearby frequency). On a practical level, the quasi-linear response

properties of a membrane can be measured by providing a small-amplitude, sinusoidally-

modulated voltage command to an intracellularly-recorded neuron and recording the resultant

sinusoidal output current. Calculating the ratio of the voltage amplitude to the current

response yields an estimate of the membrane's impedance (which is, loosely speaking, an

indication of the neuron's "input resistance" to a sinusoidal input as a function of its

frequency). Membrane impedances have been reported for various neuronal types in the brain

(Hutcheon et al., 1996; Ulrich, 2002), including in the striatum (Beatty et al., 2015). Beatty

and collaborators (2015) found that CINs exhibit a resonance in the vicinity of 1 Hz.

Moreover, they found that the shape of the impedance function depended on the holding

voltage, which is understandable given that the amplitude of the various subthreshold

currents is voltage-dependent. Finally, with use of tetrodotoxin (TTX), a selective antagonist

of voltage-activated Na^+ (Nav) channels, Beatty and collaborators (2015) demonstrated that

NaP currents contributed to these filtering properties of the CIN membrane.

Use of somatic voltage perturbations, however, fails to discriminate the role played by

the CINs' dendritic arbor *per se* in transforming synaptic inputs. The CINs' dendritic arbor

can span well over half a millimeter from the soma (Wilson, 2004), with cortical inputs

terminating on distal dendrites and thalamic inputs terminating perisomatically and on

proximal dendrites (Doig et al., 2014; Lapper and Bolam, 1992; Mamaligas et al., 2019;

Thomas et al., 2000). Both dendritic morphology (as taught by cable theory) and dendritic

nonlinearities will lead to distal cortical inputs being integrated differently from proximal

thalamic input. Thus, the impact of membrane nonlinearities on the quasi-linear

approximation will depend on *where* they are expressed throughout the dendritic arbor. To

address this question, we recently developed an optogenetics-based experimental method

(that relies on the use of quasi-linear cable theory and Fourier analysis) to determine the

duration of the delays introduced by dendrites, and how these delays impact the rapidity and

fidelity of a neuron's response to its input. We used our method to study GABAergic neurons

of the substantia nigra pars reticulata (SNr) that expressed channelrhodopsin-2 (ChR2). We

illuminated (with a 470 nm LED) either a small perisomatic region or the entire dendritic

arbor. Comparison of the two illumination regimes enabled us to demonstrate that dendrites

(that in the SNr can be >700 μm long) introduce a significant integration delay. The analysis

also yielded that SNr dendrites behaved like passive linear filters, without evidence for

amplification or resonances, or any dependence on holding voltage (Tiroshi and Goldberg,

2019).

In the present study, we apply our method to study nonlinearities of CIN dendrites.

We demonstrate that HCN and NaP currents shape the quasi-linear response properties of

CINs, and that dendrites contribute additional phase delays. Furthermore, we show that our

analysis can reveal information about dendritic location of membrane nonlinearities, and use

the analysis to deduce that both HCN and NaP currents are expressed primarily proximally.

Blocking NaP currents pharmacologically revealed that only perisomatic illumination triggers

a boosting response. The proximal distribution of NaP currents is further supported by

measuring: a) how far autonomously generated backpropagating action potentials (bAPs)

actively invade the dendritic arbor; and b) the differential boosting by NaP currents of

proximal thalamostriatal vs. distal corticostriatal EPSPs.

Figure 1: CIN membranes exhibit voltage-dependent quasi-linear properties. **A.** CINs exhibit a voltage sag due to HCN currents, and autonomous pacemaking due to NaP currents. **B.** Application of 2 mV sinusoidal voltage commands to the soma, of increasing frequencies, elicits a current response that is voltage dependent (black trace, –55 mV; red trace, –70 mV). **C.** Estimation of the impedance (left) and phase shift (right) show that at –55 mV, CINs exhibit an amplified impedance and that at –70 mV, CINs exhibit a resonance (non-monotonic impedance and negative phase delays). Solid lines are parameter fits for $(\alpha^2 + \beta^2)^{-1/2}$ up to a scale factor (left), and phase shift, ϕ_s (right, see Eq. 5).

Results

Ionic currents underlying amplification and resonance in cholinergic interneurons

The HCN and NaP currents depolarize CINs over largely non-overlapping voltage ranges. The HCN current is mostly active below –60 mV and is responsible for the voltage sag in response to a hyperpolarizing current pulse (Figure 1A, "HCN"), whereas the NaP current takes over at –60 mV and is necessary and sufficient (Bennett et al., 2000) to drive CINs to action potential threshold (Figure 1A, "NaP"). Therefore, because NaP is a regenerative current, while HCN is a restorative current, we would expect the current responses to an oscillating voltage command to depend strongly on whether the membrane voltage is clamped above or below –60 mV (Beatty et al., 2015). We therefore held mouse CINs in whole-cell voltage clamp (*n = 10* neurons, *N = 4* mice), first at –55 mV, and

subjected them to a voltage command that was composed of a continuous sequence of

sinusoidal cycles with an amplitude of 2 mV and a frequency that increases discretely from

0.2 to 20 Hz. The current amplitude was very small at low frequencies and increased

monotonically to the high frequency (Figure 1B, black), which is suggestive of an impedance

curve with amplified lower frequencies. Loosely speaking, this means that the CINs' "input

resistance" to a low frequency oscillatory input currents is boosted. As expected (Figure 1C,

black), the impedance curve, $|Z(f)|$, exhibited an amplifying structure, with the phase delay

being strictly positive. In order to quantify the degree of amplification, we fit the model of

the phase delay, ϕ_s, for an iso-potential cell with quasi-linear properties (see Materials and

Methods). In this fit, there is a (negative) amplifying parameter, μ_n – that is derived from

biophysical properties of amplifying current (e.g., the slope of the activation curve, reversal

potential, etc.) (Goldberg et al., 2007) – which was estimated to be $\mu_n = -3.9$.

When neurons were held at –70 mV ($n = 10$ neurons, $N = 7$ mice), the current

response was very different, it was much larger at subhertz frequencies as compared to the

experiment at –55 mV, and then exhibited what looks like a slightly decreasing amplitude for

frequencies near 1 Hz, followed by an amplitude increase at higher frequencies (Figure 1B,

red). Estimation of the amplitude response and phase delay (Figure 1C, red), revealed

significantly different curves (amplitude: $P = 6 \cdot 10^{-25}$, phase: $P = 1 \cdot 10^{-17}$, ANCOVA) with a

clear resonance peak at approximately 1 Hz, and a zero crossing at a slightly lower frequency

(Figure 1C, red). Fitting the somatic phase delay yielded a reasonable fit only when both an

amplifying and resonant component were included in the fit with the amplifying parameter

($\mu_n = -3.4$) being only slightly reduced relative to the –55 mV fit. In contrast, the additional

(positive) resonance parameter, μ_h – which is derived from the biophysical properties of the

restorative current – was estimated to be $\mu_h = 1.6$ (see Materials and Methods) based on the

fit to the negative lobe in the phase delay.

The previous experiments suggest that the amplifying effect at –55 mV occurs due to

prominence of the amplifying NaP current in that voltage range, whereas the resonance

visible at –70 mV is due to the HCN current that dominates that voltage range. This

conclusion was supported by the fact that application of 1 μM TTX ($n = 9$ neurons, $N = 4$

mice), that abolished autonomous spiking and slightly hyperpolarized the CIN (Figure 2A),

reduced the impedance ($P = 9 \cdot 10^{-9}$) and exhibited a trend toward a shortened phase delay (P

$= 0.137$, ANCOVA, Figure 2B). These changes are captured by the amplification parameter

being estimated as less negative ($\mu_n = -2.0$), which reduces ϕ_s (Eq. 5 in Materials and

Methods). Similarly, application of 10 µM ZD7288, the selective HCN antagonist ($n = 9$

neurons, $N = 7$ mice), which abolished the sag response (Figure 2C), abolished the resonance

peak in the impedance curve ($P = 4 \cdot 10^{-16}$) and significantly reduced the negative lobe in the

phase delay ($P = 0.018$, ANCOVA, Figure 2D), which was captured by the resonance

parameter being reduced to $\mu_h = 0.5$.

Figure 2: Amplification is caused by NaP currents, whereas the resonance is caused by HCN currents. A. TTX (blue) prevents autonomous spiking (black). **B.** TTX prevents amplification of the impedance (left), and reduces the phase shift (right). **C.** ZD7288 (red) abolishes the voltage sag (black). **D.** ZD7288 abolishes the resonance peak in the impedance, and the negative phase shifts in the subhertz range. Solid lines are parameter fits as in Figure 1.

<u>Optogenetic interrogation of the spatial distribution of NaP and HCN currents in the CIN</u>

<u>dendritic arbor</u>

While the above experiments demonstrate the NaP and HCN currents are capable of

transforming subthreshold voltage fluctuations, the question remains as to where within the

CIN's somatodendritic compartments these currents perform their amplifying and restorative

actions, respectively. One extreme scenario is that they are restricted to the axosomatic region

(where they are needed for sustaining the autonomous firing patterns of the CINs). In that

scenario, the dendrites could be entirely linear, passively transmitting the distal

depolarizations to the soma. Only at the soma are the inputs then transformed by these

currents. However, a more realistic scenario is that these currents are also expressed

dendritically and exert their nonlinear influence on synaptic inputs more distally. But in this

scenario, we would still want to know where along the dendrites the currents are expressed.

To determine the precise location of these channels would require advanced *in situ* molecular

and anatomical techniques and/or direct electrophysiological recordings from CIN dendrites

in conjunction with advanced imaging techniques. While these experiments may, in principle,

be done, we wondered whether the optogenetics-based technique that we recently developed

to interrogate the role of dendrites in synaptic integration (Tiroshi and Goldberg, 2019) could

help us address this question.

We previously showed that the impact of dendrites on synaptic integration can be

quantified by studying the response of neurons that express ChR2 post-synaptically to (470

nm) illumination of their dendritic arbor. In particular, we compared between two spatial

patterns of illumination of the somatodendritc arbor of SNr neurons: either a small

perisomatic region (approximately 100 μm in diameter) or the entire dendritic arbor. By

using sinusoidally-modulated blue LED illumination at various temporal frequencies, we

were able to calculate the phase delays produced by both spatial patterns, and found that

illumination of the entire dendritic arbor introduced larger phase delays. In order to quantify

the effect, we fit the data to a tractable theoretical model of a semi-infinite cable (Goldberg et

al., 2007; Tiroshi and Goldberg, 2019). As mentioned above, in the case of the SNr neuron,

we found that the dendrites were well-fit by a passive linear cable model, whose parameters

(i.e., time and space constants) we could estimate. The conclusion that SNr dendrites were

largely linear was further supported by the finding that these phase delays were voltage-

independent (Tiroshi and Goldberg, 2019).

Because CINs exhibit prominent amplifying and resonating currents that are strongly

voltage dependent (Figures 1&2), we posited that by using the same optogenetic technique

and semi-infinite cable model we would be able to quantify the contribution of CIN dendrites

to post-synaptic integration. In particular, we hypothesized that by fitting a *quasi-linear* cable

model, we would be able to quantify to what degree CIN dendrites *per se* possess amplifying

or resonating properties (See Materials and Methods). Finally, by comparing illumination of

the proximal *vs.* the entire dendritic arbor, we could learn something about the localization of

the nonlinearities along the dendritic arbor. To this end, we crossed mice that express Cre-

recombinase under a ChAT promoter with the Ai32 mouse that expresses ChR2 and EYFP in

a Cre-dependent manner (see Materials and Methods). The cholinergic neuropil and

individual CINs could be clearly visualized in the dorsal striatum of these ChAT-ChR2 mice

(Figure 3A). Individual CINs were patched and recorded in voltage clamp, while illuminating

either the proximal region with a 60X water-immersion objective (Figure 3A) or the entire

slice with a 5X air objective, with a continuous sequence of sinusoidally modulated

illumination waveforms at various frequencies (Figure 3B, blue).

Figure 3: Optogenetic interrogation of the quasi-linear properties of CIN dendrites indicates that dendritic nonlinearities are more prominent proximally. A. A CIN in a sagittal slice from ChAT-ChR2 mouse is patch-clamped in the whole-cell mode while either a small proximal region around the soma or the full-field are illuminated with a sinusoidally modulated 470 nm LED. **B.** The current response to the proximal (black) and full-field (red) illumination differ, with the phase of the full-field illumination right-shifted at the higher frequencies (20 Hz is highlighted in yellow). **C.** Phase shifts at –55 mV holding potential, calculated for proximal (black) and full-field illumination (red). A tendency towards negative phase shifts is present at 0.4 Hz (green arrow). Fitting the passive model at –55 mV demonstrated that the effective range of illumination (r) is larger for the full field fit (Eq. 2). **D.** Phase shifts at –70 mV holding potential, exhibit a negative phase shift, and the resonance parameter (μ_h) is smaller for the full field fit, as is the magnitude of the amplification parameter (μ_n, see main text). The elevation in these parameters' magnitudes when illuminating proximally relative to full-field suggest that the surface densities of NaP and HCN currents is higher proximally. **E.** Phase response for proximal (left) and full-field illumination (right) in TTX at –55 mV. **F.** Amplitude response for proximal (left) and full-field illumination (right) at –55 mV reveals an opposite effect of TTX.

Comparison of the somatic current traces in response to proximal (Figure 3B, black)

vs. full-field (red) illumination demonstrated that the phase of the full-field-generated current

is delayed relative to the proximally-generated current. This could be observed in the raw

data both for the low and high (yellow inset) frequencies. The effect at the high frequency

was very evident when plotting the phase delay curves both when the CINs were held at –55

mV (Figure 3C) and at –70 mV (Figure 3D). Estimation of these phase delays revealed

delays that were considerably larger than those observed with the electrical somatic

stimulation (Figure 1). The main contributor to the large delays are the kinetics of the ChR2,

ϕ_C (Eq. 6b, in Materials and Methods) with an additional dendritic delay ϕ_d (Eq. 2 in

Materials and Methods). So we used independent measurements of the ChR2 kinetics in

CINs (Figure 3–figure supplement 1), and previous literature about these kinetics (Nagel et

al., 2003; Tchumatchenko et al., 2013) to fit the phase and amplitude contribution of ChR2,

as explained in Appendix 1.

We found that fitting our model of ChR2 kinetics to the significantly-different phases

observed in the proximal and full-field illuminations at – 55 mV (*n = 5* neurons, *N = 4* mice,

P = 4·10⁻³, ANCOVA) yielded that the full-field illumination activated roughly twice the

electrotonic range (*r = 1.07*) that was activated by the proximal illumination (*r = 0.46* ,

Figure 3C). However, a closer look at these fits reveals that the phase delay at 0.4 Hz (for

both the proximal and full field illumination), is not captured by this model. This dip in phase

delay (Figure 3C, green arrow) at this low frequency is reminiscent of what a restorative

current is expected to do. In order to accentuate the effect of the restorative HCN currents, we

repeated the experiment at the –70 mV holding potential (Figure 3D). In this case (*n = 14*

neurons, *N = 10* mice), the phase delays at the lower frequencies (especially at 0.4 Hz) were

negative, which was reminiscent of the results from the electrical voltage stimulation experiments (Figure 1) at the –70 mV holding potential. Estimation of the amplitude responses for proximal and distal stimulation at both holding potentials revealed that they were less sensitive at revealing the resonance structure (Figure 3–figure supplement 2), that was more readily read off from the phase responses (Figure 3D) in the sense that phase estimates provided tighter error bars than the amplitude estimates.

When fitting a full ChR2 kinetics plus quasi-linear dendrite model (incorporating both amplifying and restorative parameters) to the –70 mV measurements (Figure 3D), we found that the curves were significantly different ($P = 8 \cdot 10^{-4}$, ANCOVA) and the estimates of the effective electrotonic range of illumination remained similar to those estimate with the passive model (Figure 3C): $r = 1.06$ for the full-field and $r = 0.58$ for the proximal illumination. Here too the restorative parameter was estimated to be larger for the proximal fit ($\mu_h = 3.4$) relative to the full field fit ($\mu_h = 1.7$), suggesting that the HCN currents are denser more proximally.

Table 1 summarizes the model parameters used to fit the curves in Figures 1-3. In Appendix 1, we discuss how the various parameters affect the model and provide a more detailed description of how the parameter space was searched to fit the model.

Figure, Curve	μ_n	μ_h	τ_n (ms)	τ_h (ms)	γ_R	τ (ms)	r	Amp (GΩ)
1C, -55 mV	-3.9189	-	28	-	5.4296	41.8	-	2.4728
1C, -70 mV	-3.3854	1.5863	11.6	1077.8	5.8722	26.1	-	1.4229
2B, TTX	-2.0178	-	31.7	-	3.4610	19.4	-	1.3809
2B, ZD7288	-1.8988	0.4844	13.4	9107.5	2.6111	17.2	-	0.8323
3C, proximal	-	-	-	-	-	35.8	0.4360	-
3C, distal	-	-	-	-	-	26.8	1.0651	-
3D, proximal	-1.4763	3.4400	60.0	551.2	3.3405	33.6	0.5788	-
3D, distal	-0.8885	1.6646	49.2	403.3	3.8850	45.3	1.0646	-
Table 1. Parameters fit to quasilinear model in Figures 1–3.								

The conclusion that the HCN current is concentrated proximally is buttressed by the fact that the negative phase delays in the subhertz frequency range tended to be less negative for the full-field illumination ($P = 0.037$, Wilcoxon rank sum test on phase delays at 0.4 Hz). In the framework of our quasi-linear model, the only mathematically possible way to recreate this finding while increasing the effective electrotonic range r being illuminated is by reducing the restorative parameter μ_h. Otherwise, if μ_h were constant then illuminating a larger region of the cable – by increasing r – would necessarily recruit more restorative current, thereby making the negative delays more negative (See Appendix 1).

Aside from the deviation at 0.4 Hz (Figure 3C, green arrow, which may be indicative

of a resonant component), the other phase measurements at –55 mV are consistent with a

simple model of ChR2 kinetics plus a passive dendritic delay, thereby raising the possibility

that optogenetic dendritic activation fails to engage NaP currents. This failure could suggest

that NaP currents are relatively absent from the dendrites. However, our somatic

measurements showed clear evidence that NaP currents affect the quasi-linear properties of

the soma (Figure 2B). Additionally, the fit to the phases at –70 mV, required the inclusion of

an amplification parameter, which was estimated to have a larger magnitude for the proximal

illumination ($\mu_n = -1.5$) than for the full-field illumination ($\mu_n = -0.9$). It therefore seems

more likely that the Nav channels that underlie the NaP currents are localized proximally and

taper off distally. To test this, we repeated the measurements at –55 mV before and after

application of TTX. While TTX application may have slightly increased the phase delays for

both proximal and full-field illumination (Figure 3E) – which could reflect a reduction in the

overall dendritic membrane conductance of the CINs, and therefore a lengthening of its space

constant – the amplitude responses (Figure 3F) unequivocally show that TTX exerts an

opposite affect when proximal *vs.* full-field illumination are used. While full-field

illumination in TTX increased the somatic current's amplitude response (control: $n = 9$

neurons, $N = 7$ mice; TTX: $n = 14$ neurons, $N = 8$ mice, $P = 7 \cdot 10^{-9}$, ANCOVA) presumably

by reducing the current escape via the dendritic membrane, proximal illumination in TTX

reduced the somatic current's amplitude response (control: $n = 9$ neurons, $N = 7$ mice; TTX:

$n = 12$ neurons, $N = 6$ mice, $P = 1.1 \cdot 10^{-4}$, ANCOVA) indicating that proximal NaP currents

indeed boost the somatic current, as concluded from the somatic experiments (Figure 2).

Thus, this pharmacological result provides *model-independent* evidence that the NaP currents

are expressed primarily proximally and less so distally. In the following section, we provide

independent evidence in support of this conclusion.

Figure 4: Autonomous action potentials actively back-propagate in CIN dendrites up to 70 μm from the soma. A. Autonomous discharge of a CIN that was patched and filled with fluo-4 and Alexa Fluor 568 for 2PLSM imaging (image). **B.** Line scans at various distances from the soma exhibit Ca^{2+} oscillations caused by bAPs. **C.** Calculating the spike-triggered average of these oscillations and fitting an alpha-function gives an estimate of the amplitude of these oscillations (in % $\Delta F/F_o$). **D.** The scatter plot of these amplitudes as a function of distance from the soma (11 CINs from 7 mice are pooled) exhibits a large degree of variability. However, a 35 μm moving average (black line) exhibits that the Ca^{2+} transients begin to decay approximately 70 μm from the soma, indicating that bAPs are supported by Nav channels up to that point (the initial increase up to 20 μm is due to the increase in the surface-to-volume ratio in the large proximal dendrites).

<u>Distance of dendritic bAP invasion indicates location of amplifying Nav channels</u>

The persistent and fast-inactivating Nav currents flow through the same Nav channels

(Alzheimer et al., 1993). Therefore, a method that is indicative of where these NaV channels

are located will indicate where the NaP current can be found. One such method involves

determining to what distance from the soma dendritic bAPs invade the dendritic arbor. To

this end, we used 2PLSM Ca^{2+} imaging to measure the size of the Ca^{2+} transients elicited by

bAPs in autonomously firing CINs at various distances from the soma (Figure 4A). We

conducted line scans (*n = 11* neurons, *N = 7* mice) to measure the Ca^{2+} signals at various

distances from the soma (Figure 4B). Next, we estimated the size of the spike triggered

average (STA) of the $\Delta F/F_0$ Ca^{2+} signal, by averaging around spontaneous APs measured at

the soma (Figure 4C). The amplitude of the STA was estimated by fitting it with an alpha-

function (see Materials and Methods). A scatter plot of STA amplitude *vs.* distance from

soma demonstrates a large degree of variability (Figure 4D). Nevertheless, applying a 35 μm

moving average to the scatter plot reveals a trapezoidal dependence of Ca^{2+} transient sizes

(Figure 4D, black). We previously demonstrated that somatic Ca^{2+} transients are smaller than

dendritic transients due to the difference in the surface-area-to-volume ratio (Goldberg et al.,

2009; Rehani et al., 2019). In the same vein, the initial dip in the size of the transients at short

distances from the soma result from the large size of the proximal dendrites (Figure 4D),

relative to the distal dendrites. Neglecting that effect, we found that the size of the bAP-

driven Ca^{2+} transients remains constant up to approximately 70 μm from the soma, and then

drops off. Additionally, in 7 CINs (*N = 5* mice) in which we had measurements of *ΔF/F₀* at

both proximal and distal (>70 μm) locations, we found that the median distal signal was

significantly lower than the proximal one by 29% (*P = 0.047*, Wilcoxon rank sum test). This

spatial dependence of the Ca^{2+} transient amplitude, suggests that the bAP maintains a

constant amplitude due to the presence of Nav channels that sustain their regenerative nature

out to some 70 μm from the soma. Farther out the Ca^{2+} transients decrease presumably due to

a drop off in Nav channel expression, which leads to a lower amplitude bAPs, and hence less

Ca^{2+} via voltage-activated Ca^{2+} channels. Thus, this measurement strengthens the conclusion

that NaP currents are present in CIN dendrites – primarily in proximal dendrites (up to

approximately 70 μm from the soma).

<u>NaP currents boost proximal thalamic inputs but not distal cortical inputs to CINs</u>

The above data indicate that Nav channels are preferentially expressed in proximal

dendrites. These channels should amplify synaptic inputs impinging on the dendrites. In the

case of linear dendrites – either passive or even quasi-linear dendrites – the relative boosting

will affect all inputs at all locations equally. However, as demonstrated in Appendix 2 – using

a simulated nonlinear dendrite that expresses the NaP current proximally – proximal inputs

are more strongly *nonlinearly* boosted than distal inputs, because the latter decay passively

before reaching the amplifying region, and are therefore boosted less.. Thus, because

thalamic inputs on to CINs terminate proximally, whereas cortical inputs terminate distally

(Doig et al., 2014; Lapper and Bolam, 1992; Mamaligas et al., 2019; Thomas et al., 2000), in

order to generate comparable EPSPs we presumably need to activate many more distal

cortical inputs than proximal thalamic inputs. However, because distal cortical inputs are

presumably amplified less their spatial summation at the soma will also be amplified less. In

contrast, because proximal thalamic inputs consist of individual inputs, each of which

engages the local nonlinear amplification, their summation at the soma will also exhibit more amplification. Thus, we expect only the thalamic – but not cortical – inputs to be significantly boosted by Nav current. To test this, we transcranially inoculated the parafascicular nucleus (PfN) of Vglut2-ires-Cre mice with adeno-associated viruses (AAVs) harboring Cre-dependent ChR2 and EYFP genes (Figure 5A) (Aceves Buendia et al., 2019; Rehani et al., 2019). Two weeks later, we tested the sensitivity of optogenetically-evoked monosynaptic EPSPs (Figure 5–figure supplement 1A) to ranolazine (30 µM), a selective antagonist of the NaP current, in CINs patched in acute striatal slices. The mean amplitude of the EPSP was reduced by a median of 16% ($n = 11$ neurons, $N = 4$ mice, $P = 0.04$, Wilcoxon signed-rank test, Figure 5B). In order to test whether the more distal cortical inputs are modulated by NaP, we used the Thy1-ChR2 mice (Aceves Buendia et al., 2019; Arenkiel et al., 2007) that expresses ChR2 in corticostriatal fibers (Figure 5C), but neither in the PfN (Figure 5–figure supplement 1B) nor in the pedunculopontine nucleus (Gradinaru et al., 2009) that also provides monosynaptic glutamatergic projection to CINs (Assous et al., 2019). The monosynaptic EPSPs (Figure 5–figure supplement 1A) elicited by cortical inputs to CINs in these mice were unaffected by ranolazine ($n = 11$ neurons, $N = 6$ mice, $P = 0.74$, Wilcoxon signed-rank test, Figure 5D). Moreover, ranolazine had no effect on paired-pulse ratios (Aceves Buendia et al., 2019) at either synapse (Figure 5–figure supplement 1C), ruling out a presynaptic mechanism-of-action. Thus, we conclude that the post-synaptic modulation of glutamatergic inputs to CINs by NaP currents corresponds to the spatial distribution of these currents as derived from the quasi-linear properties of CINs and from our 2PLSM imaging of bAPs.

Figure 5: Thalamic – but not cortical EPSPs – onto CINs are boosted by NaP currents in healthy mice. A. The parafascicular nucleus (PfN) of Vglut2-Cre mice was inoculated with AAVs harboring Cre-dependent ChR2, so that 470 nm LED illumination of striatal slices activated monosynaptic PfN terminals while CINs were patched in current clamp mode. **B.** Optogenetically evoked thalamic EPSPs in CINs (held between –50 mV and –60 mV) were reduced by 30 μM ranolazine. **C.** CINs were patched in current clamp in Thy1-ChR2 mice, so that 470 nm LED illumination of striatal slices activated nominally cortical terminals. **D.** Optogenetically evoked monosynaptic cortical EPSPs were unaffected by ranolazine.

Figure 6: TANs, but not SPNs, exhibit a pause-like response to slow-wave events occurring during natural non-REM sleep in non-human primates (NHPs). A. simultaneous recording of LFP and an SPN in an NHP during N2 and N3 stages of sleep. **B.** simultaneous recording of LFP and a TAN in an NHP during N2 and N3 stages of sleep. **C.** SPN firing rate conditioned on the occurrence of an LFP slow wave event (6,065 triggers). **D.** Average striatal LFP signal triggered on the occurrence of slow wave events (see Materials and Methods). **E.** TAN firing rate conditioned on the occurrence of slow wave event (28,603 triggers). **F.** simultaneous recording of LFP and a TAN in an NHP during a sleep spindle. **G.** Average striatal LFP signal triggered on the occurrence of a sleep spindle (see Materials and Methods). **H.** TAN firing rate conditioned on the occurrence of a sleep spindle (5,829 triggers). Gray box indicates the 99% confidence intervals.

Tonically active neurons in non-human primates exhibit a pause-like response to slow wave oscillations during sleep

Our data, alongside previous studies (Beatty et al., 2015), demonstrate that somatodendritic HCN currents give rise to resonances in the subhertz range in CIN surface membranes. In contrast, SPNs exhibit no resonance (Beatty et al., 2015). This suggests CINs in intact animals may exhibit increased sensitivity to oscillatory inputs in that frequency range, whereas SPNs should not. Delta waves in the electroencephalogram (EEG) and in cortical and sub-cortical local field potentials (LFP) are prominent during non-REM sleep across species (Brown et al., 2012; Liu and Dan, 2019), including in non-human primates (NHPs) (Mizrahi-Kliger et al., 2018). Because the LFP is widely thought to represent subthreshold cellular activity, affected by afferent and recurrent synaptic inputs (Buzsáki et al., 2012), we hypothesized that the tonically active neurons (TANs) in the NHP striatum – that are comprised primarily of CINs (Aosaki et al., 1995; Kawaguchi, 1993; Wilson et al., 1990) – will exhibit stronger entrainment to slow-wave LFP events during sleep, while SPNs under the same conditions will not. Indeed, triggering the spike trains of TANs and SPNs (from $N = 2$ NHPs) on slow wave events that occur during sleep stages N2 and N3 (Mizrahi-Kliger et al., 2018; Riedner et al., 2007), which represent relatively deep non-REM sleep, demonstrates that while the firing rate of SPNs was unaffected ($n = 83$), the firing rates of TANs ($n = 122$) was modulated by these slow-wave events (Figure 6A-E). In contrast, the TANs firing was not modulated during higher frequency sleep spindle events (Figure 6F-H), in agreement with the subhertz membrane resonance peak we observed in the acute slice experiments. Importantly, the biphasic response exhibited by TANs to slow-wave events bears a striking resemblance to the "classical" TAN response to external cues (Aosaki et al., 1995; Apicella et al., 1991; Joshua et al., 2008; Kimura et al., 1984). Because these responses in awake primates require an intact thalamic projection (Bradfield et al., 2013; Goldberg and Reynolds, 2011; Matsumoto et al., 2001; Schulz and Reynolds, 2013; Smith et al., 2004; Yamanaka et al., 2018) (and there is no reason to assume that this is altered in sleep), our findings (Figure 6) provide support to the hypothesis that the CIN membrane resonance contributes to a thalamically-driven biphasic CIN response during slow-wave sleep, possibly because this projection terminates proximally on CINs. Importantly, because the primates are asleep in a sound-proof room, this is, to the best of our knowledge, the first report of striatal TANs responding to an internally-generated brain event, and not an external saliency-related cue.

Discussion

Quasi-linear dendritic properties of CINs and the non-uniform distribution of their active conductances

In the current study, we applied a cable-theoretic and optogenetics-based formalism – that we developed previously to study dendritic properties of SNr neurons (Tiroshi and Goldberg, 2019) – to study the dendritic properties of CINs. Unlike SNr neurons that were found to act like passive linear cables exhibiting no voltage dependent properties, we find that CIN dendrites exhibit both amplification and resonances in a voltage dependent manner: amplification is more prominent at depolarized subthreshold potentials (approximately –55 mV), whereas resonances are more prominent at more hyperpolarized potentials (approximately –70 mV). Moreover, we found that our method is able to provide information about where in the dendritic arbor these quasi-linear properties are localized. We extracted boosting (μ_n) and resonance (μ_h) parameters for proximal vs. full-field illumination, and found that the magnitude of both parameters is smaller when the entire dendritic arbor is illuminated. In other words, activating more of the distal dendritic membrane dilutes the amplifying and resonating effects, meaning that the additional membrane illuminated in the full-field condition contributed less to amplification and resonances. The simplest interpretation of this finding is that the active conductances that give rise to amplification and resonance are more highly expressed proximally and apparently taper off farther out in the dendritic arbor. We identified the relevant currents by studying the quasi-linear properties of CIN somata with voltage perturbations. We found that amplification is TTX-sensitive and therefore arises from Nav channels, that give rise to the NaP current. Similarly, we found that resonances require HCN channels.

We used one-direct and two indirect methods to validate the preferentially proximal localization of NaP currents. First, TTX application revealed that boosting occurs only at the soma (Figure 2B) and in proximal (but not distal, Figure 3F) dendritic membranes. Importantly, the TTX results demonstrate that our conclusion that boosting is restricted to the soma and proximal dendrites is *independent* of the quasi-linear model fit. Second, we implemented a widely-used approach of estimating the distance of dendritic invasion of bAPs by imaging the Ca^{2+} influx that accompanies them (Carter and Sabatini, 2004; Day et al., 2008; Goldberg et al., 2009; Kerr, 2004; Rehani et al., 2019; Tanimura et al., 2016). We found – not unlike in SPN neurons – that bAPs maintain their amplitude out to 70 μm from

the soma and then begin to decay, suggesting that Nav currents that support bAP propagation

taper off from that point onwards. Because strictly speaking a change in Ca^{2+} transients could

result from a change in the concentration of voltage-activated Ca^{2+} channels, future work can

verify our results by directly testing where locally puffed TTX attenuates the bAP (perhaps

even with the use of voltage sensitive probes). Second, we demonstrated that only thalamic –

but not cortical – EPSPs exhibit sensitivity to the selective NaP blocker. Because the thalamic

terminals are located more proximally (Lapper and Bolam, 1992; Mamaligas et al., 2019;

Thomas et al., 2000), this provides further evidence that NaP currents are more prominent

locally. The preferential localization of HCN currents proximally in CIN dendrites requires

additional validation, particularly because other neurons exhibit an opposite pattern (Berger

et al., 2001; Harnett et al., 2015; Kole et al., 2006). However, we were less interested in

validating the proximal localization of HCN currents, because unlike NaP currents, activation

of HCNs is conditioned on CINs being actively (after-)hyperpolarized, during which bAPs

are unlikely to occur. We did however test a context in which HCN currents, and the

resonance they underlie, are likely to affect CIN collective dynamics. We found that TANs

respond biphasically to internally-generated slow wave activity during sleep, but not to sleep

spindles. This preferential entrainment is consistent with the fact that CIN membranes exhibit

a resonance in the subhertz range as found by us and others (Beatty et al., 2015). The

elevated impedance (or frequency-dependent access resistance) in this range means that

inputs fluctuating in this range will more efficiently depolarize the CINs, and therefore more

likely to trigger additional action potentials than inputs fluctuating at a frequency that is far

from the resonance frequency. Accordingly, SPNs that do not exhibit resonances (Beatty et

al., 2015) are not entrained by the slow wave oscillation. Our findings in sleeping NHPs are

at odds with various rodent studies that found that cortical slow-wave activity is weakly

associated with the discharge of TANs in anesthetized rodents, and more strongly associated

with SPN activity (Goldberg et al., 2003; Mahon, 2001; Reig and Silberberg, 2014; Schulz et

al., 2011; Sharott et al., 2012; Stern et al., 1998). These differences may be attributable to

species differences and/or differences between natural sleep and anesthesia, which may differ

in the degree to which thalamic *vs.* cortical inputs engage striatal neurons. Moreover, the

dendritic nonlinearities in SPNs (Plotkin et al., 2011) may be preferentially engaged during

anesthesia, thereby causing SPNs to respond more strongly under anesthesia than during

natural sleep.

Advantages and limitations of the quasi-linear formalism and the optogenetics-based approach

As mentioned in the Introduction, the quasi-linear approach provides a tool to characterize membrane excitability in general functional terms, without requiring the complete biophysical characterization of membrane currents. Unfortunately, the method is not invertible: even after estimating all the quasi-linear parameters one cannot deduce the full biophysical characterization of the underlying channels even if one has a precise mathematical model of these channels. Moreover, even though we extracted the values of μ_n and μ_h, we do not know how to translate them into a quantitative measure of channel density. Still, the ability to compare parameters for various spatial illumination patterns enables us to reach qualitative conclusions regarding relative channel density. Additionally, our attribution of μ_n to NaP and μ_h to HCN is only provisional. CINs possess other channels that have a major influence on their firing patterns such as A-type, inward rectifying, and various Ca^{2+}-activated K^+ channels, to mention a few (Bennett et al., 2000; Deng et al., 2007; Goldberg et al., 2009; Goldberg and Wilson, 2010, 2005; McGuirt et al., 2021; Oswald et al., 2009; Song and Surmeier, 1996; Wilson, 2005; Wilson and Goldberg, 2006), that were not included in our analysis. Future work should elaborate how these and other channels contribute to the quasi-linear properties of CINs and their dendrites.

Using optogenetics provides another level of practicality. The seminal studies that characterized the filtering properties of dendrites or pyramidal neurons (Berger et al., 2001; Goldberg et al., 2007; Hutcheon et al., 1996; Ulrich, 2002) required dual soma-dendrite patching which is not practical for all neuronal types whose dendrites taper off rapidly. Expressing opsins in the membrane being activated, as we did, means that a single somatic patch electrode can suffice to conduct the quasi-linear characterization. We currently use this approach to illuminate large regions of the dendritic arbor simultaneously, which means we can only derive large-scale dendritic properties. In the future, localized laser stimulation of dendritic regions (visualized with the help of a fluorescent marker in the patch pipette), or even two-photon laser activation of opsins (provided one can find stimulation parameters that do not harm the dendrites) could be used as an alternative approach. On the backdrop of optogenetics being utilized almost exclusively to study circuit mapping (Häusser, 2021; Kim et al., 2017; Petreanu et al., 2009) by expressing opsins presynaptically, we believe our study joins other studies (Higgs and Wilson, 2017; Tiroshi and Goldberg, 2019) to underscore the

value of using opsins expressed postsynaptically to study neuronal – even dendritic –

excitability.

<u>Somatic amplification and resonance in CINs</u>

We preceded our optogenetic characterization of dendritic nonlinearities with the

electrical characterization of quasi-linear somatic properties using a sinusoidal voltage

command applied through the patch pipette. As previously reported by Beatty and

collaborators (Beatty et al., 2015), CIN somata exhibit a membrane resonance in the subhertz

range, that depends on the holding potential. However, while Beatty and collaborators found

that an amplified resonance is present in the more depolarized (approximately – 55 mV)

range, we found that the resonance is only pronounced in the more hyperpolarized

(approximately –70 mV) range, where it depends on the HCN current. It is possible that the

resonances observed by Beatty and collaborators (Beatty et al., 2015) at –55 mV arose from

ineffective clamping of dendrites with the slightly higher resistance electrodes used in that

study. In support of this proposition, we also found evidence for the resonance arising from

unclamped (dendritic) membranes. When the CINs were clamped at –55 mV, while somatic

voltage perturbations did not produce a resonance, optogenetic stimulation (both proximal

and full field) at 0.4 Hz produced a more negative phase than at the neighboring 0.2 Hz and

0.6 Hz stimulation (Figure 3C, green arrow). This downward deflection in phase is

reminiscent of the fullblown negative subhertz region that occurred with optogenetic

stimulation when the CINs were clamped at –70 mV (Figure 3D). Thus, it is likely that

resonance originated in both studies from dendritic HCN currents. Because in the study of

Beatty and collaborators (Beatty et al., 2015) TTX lowered the impedance, without removing

the subhertz resonance peak, we conclude that the effect of TTX in both studies is to remove

the NaP-dependent boosting, without directly affecting resonance. It also seems that the

location of the peak amplitudes estimated by Beatty and collaborators (Beatty et al., 2015) is

perhaps 1 Hz larger than the location of the peaks in our study. However, this difference

could result from the fact that they used a chirp stimulation where the frequency increases

continuously, whereas we used several periods of perfect sinusoidal waveforms with a

discrete set of frequencies. Moreover, it can be shown mathematically that for the quasi-

linear model the zero crossing in the phase response occurs at a slightly smaller frequency

than the peak in the amplitude response, so this may also contribute to the impression that the

resonance frequency observed in our experiments is slightly different from that observed in

Beatty and collaborators (Beatty et al., 2015).

<u>Dendritic contribution to capacity of CINs to differentiate between their excitatory inputs</u>

In NHPs, TANs that are comprised primarily of CINs (Aosaki et al., 1995; Goldberg

and Wilson, 2010; Kawaguchi, 1993; Wilson et al., 1990), encode – through a brief pause in

their tonic firing – external stimuli that are salient (and often unexpected, even conveying a

stop or behavioral shift signal (Aoki et al., 2018; Thorn and Graybiel, 2010)) or stimuli that

are associated with reward (Apicella et al., 1991; Goldberg and Reynolds, 2011; Kimura et

al., 1984). The pause response requires an intact projection from the PfN of the thalamus

(Bradfield et al., 2013; Goldberg and Reynolds, 2011; Matsumoto et al., 2001; Schulz et al.,

2011; Schulz and Reynolds, 2013; Smith et al., 2004; Yamanaka et al., 2018), indicating that

TANs are attuned to thalamic input. While TANs do not respond to ongoing movement

(Aosaki et al., 1994; Apicella et al., 1991; Kimura et al., 1984; Raz et al., 1996), and hence

seem less attuned to sensorimotor cortical input (Sharott et al., 2012), CIN activity in awake

behaving mice is strongly modulated by self-initiated movements (Gritton et al., 2019; Howe

et al., 2019; Yarom and Cohen, 2011). Nevertheless, slice physiology studies in rodents

(Aceves Buendia et al., 2019; Assous, 2021; Assous et al., 2017; Ding et al., 2010; Johansson

and Silberberg, 2020; Kosillo et al., 2016; Threlfell et al., 2012) clearly show that thalamic

inputs to CINs are stronger in the sense that they give rise to larger EPSPs (Johansson and

Silberberg, 2020) and can trigger a pause-like responses (Ding et al., 2010), whereas cortical

inputs to most CINs (Mamaligas et al., 2019) are weaker in that they give rise, in acute slices,

to smaller EPSPs and cannot trigger the pause-like response – although this distinction is less

pronounced in anesthetized rodents (Doig et al., 2014). Thus, it is clear that CINs

differentiate between these two excitatory inputs and respond differently to them. A major

contributor to the CINs' capacity to dissociate thalamic and cortical input is the differential

distribution of their respective terminals on the dendritic arbor: thalamic input terminate

perisomatically and on proximal dendrites, whereas cortical input terminates on distal

dendrites (Doig et al., 2014; Lapper and Bolam, 1992; Mamaligas et al., 2019; Thomas et al.,

2000). Our finding that the higher expression of NaP currents in proximal dendrites

preferentially boosts thalamic over cortical inputs, suggests that active conductances are

expressed in CIN dendrites in a manner that corresponds and reinforces the effect of the

spatial separation between the terminals of the two inputs.

<u>Adaptive changes in dendritic excitability in movement disorders</u>

The capacity of the CIN dendritic excitability to mirror the distribution of afferent

terminals is also an adaptive process. We previously reported that CIN dendritic excitability

is elevated in the Q175 mouse model of Huntington's disease (HD) (Tanimura et al., 2016).

In contrast to control (Thy1-ChR2) mice in which cortical EPSPs in CINs are insensitive to

ranolazine (Figure 5), cortical excitatory postsynaptic currents (EPSCs) in CINs from Q175

mice (crossed with Thy1-ChR2 mice) are strongly reduced by ranolazine. The acquired

dependence of distal cortical inputs on NaP currents in Q175 mice results from an

upregulation of the NaP current, which was also evidenced by bAPs invading farther out into

the CINs' dendritic arbor (Tanimura et al., 2016). This elevated excitability in the Q175

mouse is probably a homeostatic response aimed at elevating the postsynaptic sensitivity to

the remaining synaptic contents after the loss of afferent cortical and thalamic inputs

observed in HD mouse models (Deng et al., 2013). Because thalamostriatal inputs are also

altered in models of Parkinson's disease (PD) (Aceves Buendia et al., 2019; Parker et al.,

2016; Tanimura et al., 2019), future work should determine whether the excitability of CIN

dendrites is altered in these models, as well. Importantly, dendrites are positioned at the

interface between synaptic inputs and the intrinsic properties of CIN. While changes in

synaptic transmission and intrinsic excitability of CINs have received attention in models of

PD, HD and other movement disorders (Abudukeyoumu et al., 2019; Aceves Buendia et al.,

2019; Choi et al., 2020; Ding et al., 2006, 2011; Eskow Jaunarajs et al., 2015; Mallet et al.,

2019; Paz et al., 2021; Pisani et al., 2007; Plotkin and Goldberg, 2019; Poppi et al., 2021;

Tanimura et al., 2019; Tubert and Murer, 2021), studying alterations in dendritic excitability

in these models is a complementary approach that remains largely unchartered territory.

The membrane channels that are expressed in CINs (e.g., Nav, HCN, Kv4, etc.) are

targets for neuromodulation under normal, healthy circumstances (Deng et al., 2007; Helseth

et al., 2021; Maurice, 2004; Song and Surmeier, 1996). Therefore, it is likely that the

dendritic excitability of CINs can be modulated under physiological conditions, as well, as a

way to adjust the tuning of CINs to their afferent inputs. Conversely, if boosting and

resonances are present primarily proximally, the selective tuning of CINs to subhertz slow-

wave oscillations is probably stronger for thalamic inputs (that terminate proximally) than for

cortical inputs. Future experiments can address this issue by testing how silencing the PfN

affects the observed entrainment of TANs to slow-waves (Figure 6).

In summary, we have used a new optogenetics-based approach, complemented by

other electrophysiological and imaging approaches, to demonstrate that the spatial

localization of active dendritic conductances that endow CIN dendrites with quasi-linear

filtering properties corresponds to the spatial distribution of their two main afferent excitatory

inputs. This matching up of presynaptic terminals with post-synaptic excitability, probably

contributes to the capacity of CINs to respond differentially to cortical and thalamic inputs.

Sensorimotor information of cortical origin seems to be integrated continuously in a moment-

by-moment fashion. In contrast, thalamic inputs lead to abrupt pauses in TAN firing, even in

response to internally-generated brain-states (e.g., during slow-wave activity). The possibility

that dendritic arbors adapt to the spatiotemporal patterns of afferent inputs is likely an

important principle of neural computation that deserves further attention, particularly in the

framework of autonomously active neurons such at CINs, SNr neurons and other basal

ganglia pacemakers.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Genetic reagent (M. musculus)	B6;129S6-Chattm2(cre)Lowl/J	The Jackson Laboratory	Strain #:006410 RRID:IMSR_JAX:006410	ChAt-ires-CRE
Genetic reagent (M. musculus)	B6;129S-Gt(ROSA)26Sortm32(CAG-COP4*H134R/EYFP)Hze/J	The Jackson Laboratory	Strain #:012569 RRID:IMSR_JAX:012569	Ai32-conditional ChR2
Genetic reagent (M. musculus)	STOCK Slc17a6tm2(cre)Lowl/J	The Jackson Laboratory	Strain #:016963 RRID:IMSR_JAX:016963	Vglut2-ires-cre
Chemical compound, drug	Mecamylamine hydrochloride	Sigma-Aldrich	Lot # 019M4108V CAS: 826-39-1	

Chemical compound, drug	Atropine sulfate salt monohydrate	Sigma-Aldrich	Lot # BCBH8339V CAS No.: 5908-99-6	
Chemical compound, drug	SR 95531 hydrobromide (Gabazine)	Hello Bio	CAS: 104104-50-9	
Chemical compound, drug	DNQX	TOCRIS	CAS: 2379-57-9	
Chemical compound, drug	D-AP5	Hello Bio	CAS:79055-68-8	
Chemical compound, drug	CGP 55845 hydrochloride	Hello Bio	CAS: 149184-22-5	
Chemical compound, drug	Dihydro-β-erythroidine hydrobromide	TOCRIS	CAS: 29734-68-7	
Chemical compound, drug	Ranolazine dihydrochloride	Sigma-Aldrich	CAS Number: 95635-56-6	Product Number R6152
Chemical compound, drug	ZD7288	MedChemExpress	CAS No. : 133059-99-1	Synonyms: ICI D7288
Chemical compound, drug	Tetrodotoxin citrate	Hello Bio	CAS: 18660-81-6	
Chemical compound, drug	4-Aminopyridine	Sigma-Aldrich	CAS Number: 504-24-5	
Chemical compound, drug	Phosphate buffered saline tablets	Sigma-Aldrich	MDL number: MFCD00131855	Product Number P4417
Chemical compound, drug	Paraformaldehyde	Sigma-Aldrich	CAS Number: 30525-89-4	

Chemical compound, drug	XYLAZINE AS HYDROCHLORIDE	EUROVET ANIMAL HEALTH B.V	CAS: 082-91-92341-00	
Chemical compound, drug	CLORKETAM	VETOQUINOL	CAS: 1867-66-9	
Chemical compound, drug	Meloxicam	Chanelle Pharmaceuticals Manufacturing ltd	CAS Number: 71125-38-7	
Chemical compound, drug	isoflurane	Primal Critical Care	CAS Number: 26675-46-7	
Software	MATLAB	MathWorks	RRID:SCR_001622	Data analysis
Software	XPPAUT	X-Windows Phase Plane plus Auto	RRID:SCR_001996	Modelling
Software	WinWCP	University of Strathclyde Glasgow	RRID:SCR_014713	Data acquisition
Software	FEMTOSmart Software: MESc	FEMTONICS	RRID:SCR_018309	2P Data acquisition
Software	NIS-Elements Basic Research	Nikon instruments	RRID:SCR_002776	Confocal images acquisition
Software	Signal 6	Cambridge Electronic Design	RRID:SCR_017081	
Software	AlphaLab SnR	Alpha-Omega Engineering	https://www.alphaomega-eng.com/	
Software	Electrode Positioning System	Alpha-Omega Engineering	https://www.alphaomega-eng.com/	

Others	Model 940 Small Animal Stereotaxic Instrument with Digital Display Console	Kopf Instruments	https://kopfinstruments.com/product/model-940-small-animal-stereotaxic-instrument-with-digital-display-console/	
Others	Sound-attenuating room	IAC acoustics	https://www.iacacoustics.com/	
Others	Glass-coated Tungsten electrodes	Alpha-Omega Engineering	https://www.alphamega-eng.com/	
Others	Pre-gelled surface electrodes	Rhythmlink International	https://rhythmlink.com/	
Others	subdermal needle electrodes	Rhythmlink International	https://rhythmlink.com/	
Others	Infra-red camera	Norpix Inc., Canada	https://www.norpix.com/	

Ethics statement: All experimental protocols were conducted in accordance with the National

Institutes of Health Guide for the Care and Use of Laboratory Animals, and with the Hebrew

University guidelines for the use and care of laboratory animals in research. The experiments

adhered to, received prior written approval from and were supervised by the Institutional

Animal Care and Use Committee of the Faculty of Medicine, under protocols: MD-16-13518-

4 (H.B.) and MD-18-15657-3 (J.A.G.).

Mice: Homozygous Ai32 (RCL-ChR2(H134R)/EYFP) mice (The Jackson laboratory [Jax]

stock: 012569) that express floxed ChR2 and an EYFP fusion protein under the CAG

promoter were crossed with homozygous ChAT-IRES-Cre (Δneo) mice the express Cre

recombinase under the choline acetyltransferase promoter (Jax stock: 031661). The ChAT-

ChR2 offspring (4-8 weeks old/both sexes) were used for the majority of experiments. To

investigate corticostriatal transmission, we used homozygous transgenic Thy1-ChR2 mice

(B6.Cg-Tg (Thy1-COP4/EYFP) 18Gfng/1, Jax stock: 007612), that express ChR2 under the

Thy1 promoter in cortical neurons (Arenkiel et al., 2007). To investigate thalamostriatal

transmission, we used Vglut2-ires-Cre mice (Jax stock: 016963).

Non-Human Primates (NHPs): Data were obtained from two healthy, young adult, female

vervet monkeys. The monkeys were habituated to sleeping in a primate chair, positioned in a

dark, double-walled sound-attenuating room. The primate chair restrained the monkeys' hand

626	and body movements but otherwise allowed them to be in a position similar to their natural
627	(sitting) sleeping posture. Detailed sleep habituation, surgery and sleep staging were reported
628	previously (Mizrahi-Kliger et al., 2018). For extracellular recordings, the monkeys' heads
629	were immobilized with a head holder and eight glass-coated tungsten microelectrodes were
630	advanced the dorsal striatum. Electrical signals were amplified with a gain of 20, filtered
631	using a 0.075 Hz (2 pole) to 10 kHz (3 pole) Butterworth filter and sampled at 44 kHz by a
632	16-bit analog/digital converter. Spiking activity was sorted online using a template matching
633	algorithm. The striatum was identified based on its stereotaxic coordinates according to MRI
634	imaging and primate atlas data (Martin and Bowden, 2000). Tonically active neurons (TANs)
635	and spiny projection neurons (SPNs) were identified using real-time assessment of their
636	electrophysiological features. Spiking and local field potentials (LFP) were recorded only for
637	identified recording sites with stable recording quality (i.e., where single-neuron spiking
638	yielded an average isolation score ≥0.85).

639	We followed established procedures for slow wave detection in the LFP (Riedner et al.,
640	2007). Briefly, the LFP signal was filtered at the 0.5-4 Hz range and putative slow wave events
641	whose duration was 0.25-2 seconds were kept for further analysis. Next, the slow wave peaks
642	were sorted according to their amplitude. Artifacts were removed by discarding all events
643	whose amplitude exceeded 5 standard deviations above the mean. Finally, conditional firing
644	rate analysis was only performed on 30% of the slow wave events with the highest amplitude.
645	Conventional procedures were also used for sleep spindle detection in the LFP (Sela et al., 2016,
646	p. 2016). Briefly, the detection algorithm was only used for striatal sites whose LFP showed
647	significant 10-17 Hz (spindle range) activity. LFP was filtered at the 10-17 Hz range and the
648	Hilbert transform was then used to extract the instantaneous amplitude. Events exceeding 3
649	standard deviations above the mean were deemed potential spindle events, and a threshold of
650	one half of a standard deviation above the mean was used to detect the start and end point of an
651	individual sleep spindle. A potential sleep spindle was defined as such only if it lasted 0.5-3
652	seconds, and provided it did not exhibit a relatively high (more than 4.5 standard deviations
653	above the mean) amplitude in a control 20-30 Hz range. The spindle data were obtained using
654	a 4-pole 4-25 Hz Butterworth filter.

655	Histology: An 8 weeks old male Thy1-ChR2 mouse was deeply anesthetized with a terminal
656	dose of ketamine-xylazine followed by perfusion through the heart of cold PBS and 4% PFA.
657	The removed brain was kept overnight at 4˚C in 4% PFA. The next day, the brain was
658	washed 3 X 15 minutes before 50 μm coronal slices of the PfN were cut with a vibratome

(Leica VT1000S). VECTASHIELD (Vector Laboratories) was applied onto mounted slices

to protect from bleaching. Coronal slices of the PfN were imaged using confocal microscope

(Nikon A1R) using 10x lens and a 20x oil immersion lens to visualize constitutive EYFP

expression.

Stereotaxic viral inoculation in Vglut2-ires-Cre mice: Mice were deeply anesthetized with

isoflurane in a non-rebreathing system (2.5% induction, 1-1.5% maintenance) and placed in a

stereotaxic frame (Kopf Instruments, Tujunga, CA). Temperature was maintained at 35°C

with a heating pad, artificial tears were applied to prevent corneal drying, and animals were

hydrated with a bolus of injectable saline (10 ml/kg) mixed with analgesic (2.5 mg/kg

Meloxicam). Stereotaxic injections into caudal intralaminar nuclei of thalamus were

performed under aseptic conditions. Adeno-associated viruses (AAV) serotype 9 carrying

double-floxed fusion genes for hChR2 (E123A) and EYFP under an EF1a promoter

(University of Pennsylvania Vector Core, Addgene #35507) were used to transfect PfN

neurons. Injection coordinates were from Bregma: lateral, 0.65 mm; posterior, 2.3 mm; and

3.35 mm depth from surface of brain (Rehani et al., 2019). A small hole was bored into the

skull with a micro drill bit and a glass pipette was slowly inserted at the PfN coordinates. To

minimize backflow, solution was slowly injected; a total volume of 250 nl (>2.5x1012

GC/ml) of the AAV constructs was injected over a period of approximately 1.5 min and the

pipette was left in place for 5 min before slowly retracting it. Slice physiology experiments

were conducted 2-3 weeks after surgery.

Slice preparation: Mice were deeply anesthetized with ketamine (200 mg/kg)–xylazine (23.32

mg/kg) and perfused transcardially with ice-cold-modified artificial cerebrospinal fluid

(ACSF) bubbled with 95% O_2–5% CO_2, and containing (in mM) 2.5 KCl, 26 $NaHCO_3$, 1.25

Na2HPO4, 0.5 $CaCl_2$, 10 $MgSO_4$, 0.4 ascorbic acid, 10 glucose and 210 sucrose. The brain

was removed, and 275 µm thick sagittal slices containing the striatum were cut in ice-cold-

modified ACSF. Slices were then submerged in ACSF, bubbled with 95% O_2–5% CO_2,

containing (in mM) 2.5 KCl, 126 NaCl, 26 $NaHCO_3$, 1.25 Na_2HPO_4, 2 $CaCl_2$, 2 $MgSO_4$ and

10 glucose, and stored at room temperature for at least 1 h prior to recording.

Electrophysiological recording: The slices were transferred to the recording chamber

mounted on a Zeiss Axioskop 60X, 0.9 NA fixed-stage microscope and perfused with

oxygenated ACSF at 31ºC. During the optogenetic stimulation experiments, in order to

guarantee that the effects we measured were generated post-synaptically, the ACSF solution

contained (in µM) 10 DNQX to block AMPA receptors, 50 D-APV to block NMDA

receptors, 10 gabazine (SR95531) to block GABA$_A$ receptors, - 2 CGP55845 to block

GABA$_B$ receptors, 10 atropine to block muscarinic ACh receptors, and 10 mecamylamine to

block nicotinic ACh receptors. In the experiments in which optogenetics were used to

stimulate cortical or thalamic input we used the same blockers, except for DNQX which was

left out. An Olympus 40X, 0.8 NA water-immersion objective with a 26.5 mm field number

(FN) was used to examine the slice using standard infrared differential interference contrast

video microscopy. Patch pipette resistance was typically 4-5 MΩ when filled with recording

solutions. The junction potential estimated at 7-8 mV was not corrected. In EPSC

measurements, the intracellular solution contained (in mM): 127.5 CsCH$_3$SO$_3$, 7.5 CsCl, 10

HEPES, 10 TEA-Cl, 4 phosphocreatine disodium, 0.2 EGTA, 0.21 Na$_2$GTP, and 2 Mg$_{1.5}$ATP

(pH=7.3 with CsOH, 280-290 mOsm/kg). In the Ca^{2+} imaging experiments (see below) the

internal solution contained (in mM) 135 K-gluconate, 5 KCl, 2.5 NaCl, 5 Na-

phosphocreatine, 10 HEPES, 0.1 fluo-4 (Molecular Probes), 0.1 Alexa Fluor 568 (for

morphological visualization, Molecular Probes), 0.21 Na$_2$GTP, and 2 Mg$_{1.5}$ATP, pH 7.3 with

KOH (280–290 mOsm/kg). In all other experiments, the intracellular solution contained (in

mM) 135.5 KCH$_3$SO$_3$, 5 KCl, 2.5 NaCl, 5 Na-phosphocreatine, 10 HEPES, 0.2 EGTA, 0.21

Na$_2$GTP, and 2 Mg$_{1.5}$ATP, pH 7.3 with KOH (280–290 mOsm/kg). Electrophysiological

recordings were obtained with a MultiClamp 700B amplifier (Molecular Devices, Sunnyvale,

CA). Signals were filtered at 10 kHz online, digitized at 10 or 20 kHz and logged onto a

personal computer with the Signal 6 software (Cambridge Electronic Design, Cambridge,

UK). All reagents, drugs and toxins were purchased from either Merck/Sigma-Aldrich

(Darmstadt, Germany), Tocris Biosciences (Bristol, UK), MedChemExpress (Monmouth

Junction, NJ, USA) or HelloBio (Bristol, UK).

<u>Voltage-perturbation experiments</u>: CINs were held at either –55 mV or –70 mV and were

given an 83 second-long voltage command structured as a concatenated sequence of

sinusoids from a discrete set of frequencies ranging from 0.2 to 20 Hz with an amplitude of 2

mV (3 or 5 seconds per frequency, such that each frequency was represented by an integer

multiple of its fundamental period). Phase shifts between the voltage sinusoidal and the

somatic current response were determined by the location of the peak in the cross-correlation

function (CCF) of the two traces (whose units are mV·pA), for each stimulation frequency

and for each illumination condition. The impedance at each frequency, $|Z(f)|$, was calculated

from the maximal amplitude of the CCF as $|Z(f)| = (2\text{ mV})^2/\text{max(CCF)}$ (so that its units are

$G\Omega$).

<u>Optogenetic stimulation</u>: Optogenetic stimulation was performed with blue-light (470 nm)

LED illumination via the objective (Mightex, Toronto, ON, Canada). We used two spatial

illumination regimes: a) *proximal illumination* wherein an opaque disk with a central pinhole

was placed in the back focal plane of the 60X water-immersion objective such that a ~130

μm diameter region around the soma was illuminated (Tiroshi and Goldberg, 2019), thereby

targeting the soma and proximal dendrites; and b) *full-field illumination* of the entire slice

with a 5X air objective which excites the soma and the entire dendritic field. In all

experiments, LED light intensity was chosen such that stimulation generated comparable

current responses for both regimes. We used the same sequence of sinusoids described above,

only this time the voltage driving the LED was modulated (the minimal voltage was the

LED's voltage threshold, 40 mV). The phase delays were again calculated according to the

latency of the peak of the CCF between the LED voltage command and somatic current.

Note, that the phases were corrected by 0.5 (i.e., by π in radians) due to the fact that the ChR2

inward current is in antiphase with the LED's voltage command. The amplitude response was

calculated from the peak value of the CCF, normalized by the amplitude of the 470 nm LED

command (i.e., 1V for proximal illumination and 0.1 V for the full-field illumination, so that

its units are picoamperes).

To activate the excitatory synaptic inputs in the Thy1-ChR2 and in the Vglut2-mice a

full-field 470 nm LED 1 ms-long pulses were used with GABA, ACh and NMDA receptor

blockers in the ACSF. For EPSPs we average 25 trials (3 s intervals, and trials with spikes

were omitted). Paired pulse ratio (PPR) measurements consisted of 64 trials of two pulses

(100 ms apart, 3 s interval). The mean EPSC amplitude was calculated as the difference

between the mean peak current and the mean baseline current that preceded the pulse. PPR

was the ratio of the second mean EPSC to the first mean EPSC. To demonstrate that the

EPSCs were monosynaptic they were recorded before and after application of 1 μM TTX and

100 μM 4-AP (Petreanu et al., 2009).

To estimate the kinetics of the ChR2 currents, brief 1ms-long 470 nM LED pulses

(1V for proximal and 0.1 V for the full-field illumination) were repeated 250 times and the

resulting average current response was measured, and fit with an alpha function

$$A\left(e^{-t/\tau_r} - e^{-t/\tau_d}\right)$$ (Eq. 1)

to estimate the τ_r and τ_d, the rise and decay times, respectively.

<u>Two-photon laser scanning microscopy (2PLSM)</u>: The two-photon excitation source was a

Chameleon Vision II tunable Ti:Sapphire pulsed laser (Coherent, Santa Clara, CA, USA)

tuned to 820 nm. The images were collected with the Femto2D system (Femtonics, Budapest,

Hungary) which includes two 3 mm galvo-scanners, one gated GaAsP and one multi-alkaline

non-descanned photomultiplier tube for imaging fluo-4 and Alexa Fluor, respectively. Z-

stacks of optical sections (spaced 2 μm apart) were collected using 0.2 μm pixels and 15 μs

dwell times. Optical and electrophysiological data were obtained using the software package

MES (Femtonics), which also integrates the control of all hardware units in the microscope.

The software automates and synchronizes the imaging signals and electrophysiological

protocols. Data was extracted from the MES package to personal computers using custom-

made code in MATLAB. We recorded spontaneously occurring bAPs with line scans at

various distances measured radially from the tip of the soma. Spike triggered averages of the

calcium measurements ($\Delta F/F_0$) were estimated and an alpha-function (Eq. 1) was fit to them.

The value of the peak of the fitted alpha-function was used as a measure of the size of the

spontaneous bAP at that location.

<u>Drugs and Reagents:</u> TTX was used to block voltage-activated Na^+ currents. Ranolazine was

used to block NaP. ZD7288 was used to block the HCN current. Gabazine (SR-95531) and

CGP 55845 were used to block $GABA_A$ and $GABA_B$ receptors, respectively. 4-

Aminopyridine (4-AP) was used to enable optogenetically-driven monosynaptic release in

the presence of TTX. All reagents, drugs and toxins were purchased from either

Merck/Sigma-Aldrich (Darmstadt, Germany), Tocris Biosciences (Bristol, UK) or HelloBio

(Bristol, UK).

<u>Data analysis and statistics</u>: Data were analyzed and curve fitting was performed using

custom code in Matlab (MathWorks, Natick, MA, USA). Simulations of the nonlinear

dendrite (Appendix 2) was were performed using XPPAUT (Ermentrout, 2002). The

nonparametric two-tailed Wilcoxon signed-rank test was used for matched samples and the

Wilcoxon rank-sum test was used for independent samples. The parametric ANCOVA test

was used to test significant changes in the amplitude and phase curves as a function of the

natural logarithm of the frequencies (an operation which spreads out this independent

parameter more uniformly). Null hypotheses were rejected if the P-value was below 0.05.

For the TAN and SPN locking to slow wave or spindle peak analysis, confidence

intervals (at P-value of 0.01) were calculated based on the distribution of conditional firing

rates 1 s before and after the slow wave peak.

Parameter fitting to the phase delay: In our previous study (Tiroshi and Goldberg, 2019), we

modeled the dendritic arbor as a semi-infinite cable with a homogeneous quasi-linear

membrane (i.e., the current density of each nonlinearity is constant along the dendrite). When

a segment of length r (measured in units of the dendrite's space constant) from the soma is

activated with a sinusoidal current injection, the dendritic phase delay is given by

$$\phi_d = \frac{1}{2\pi}\left(\tan^{-1}\frac{q}{p} - \tan^{-1}\frac{\sin qr}{e^{pr} - \cos qr}\right) \qquad \text{(Eq. 2)}$$

794   where

$$p = \sqrt{\frac{\sqrt{\alpha^2 + \beta^2} + \alpha}{2}} \quad q = \sqrt{\frac{\sqrt{\alpha^2 + \beta^2} - \alpha}{2}} \qquad \text{(Eq. 3)}$$

796   α and β are functions of frequency and are determined by the linearization of the dendritic

797   nonlinearities as explained in Goldberg et al. (2007) and Remme and Rinzel (2011), with a

798   negative amplifying parameter, μ_n, and a positive resonance parameter, μ_h (See Appendix 1).

$$\alpha(f) = \gamma_R + \frac{\mu_n}{1 + (2\pi f \tau_n)^2} + \frac{\mu_h}{1 + (2\pi f \tau_h)^2} \qquad \text{(Eq. 4a)}$$

$$\beta(f) = 2\pi f \left[\tau - \frac{\mu_n \tau_n}{1 + (2\pi f \tau_n)^2} - \frac{\mu_h \tau_h}{1 + (2\pi f \tau_h)^2}\right] \qquad \text{(Eq. 4b)}$$

Additional parameters include total dendritic conductance (relative to leak), γ_R, the

membrane time constant τ, and the time constants representing the kinetics of the nonlinear

dendritic conductances, as explained in Goldberg et al. (2007) and in Remme and Rinzel

(2011). In some cases we only used the amplifying parameter in the fit (e.g., Figure 1C), and

in Figure 3C, we used a Eq. 2 in the case of a passive dendrite for which $\alpha(f) = 1$ and

$\beta(f) = 2\pi f \tau$.

Using the same formalism it is easy to show that for the (isopotential) soma (Eq. 5) with a quasi-linear membrane the phase delay is given by $\phi_s = \frac{1}{2\pi}\tan^{-1}\frac{\beta}{\alpha}$

807 corresponding to an amplitude that is equal to $(\alpha^2 + \beta^2)^{-1/2}$ up to a scaling factor. In the

808 main text, we point out that when μ_n becomes less negative ϕ_s is reduced. This is because

809 when μ_n becomes less negative, α is increased (Eq. 4a) and β is decreased (Eq. 4b).

810 The amplitude attenuation and phase delays generated by the ChR2 kinetics are

811 calculated from the Fourier transform of the alpha function (Eq. 1) and are given by

$$A_C(f) \propto \{(1 - \tau_r\tau_d(2\pi f)^2)^2 + (2\pi f[\tau_r + \tau_d])^2\}^{-1/2} \qquad \text{(Eq. 6a)}$$

$$\phi_C(f) = \frac{1}{2\pi}\tan^{-1}\left(\frac{2\pi f[\tau_r + \tau_d]}{1 - \tau_r\tau_d(2\pi f)^2}\right) \qquad \text{(Eq. 6b)}$$

Acknowledgments

814 This work was funded by a European Research Council (ERC) Consolidator Grant (no.

815 646886) to J.A.G. and grants from the Israel Science Foundation (no. 2051/20), German

816 Research Foundation (CRC TRR295), Israel-China Binational Science Foundation and the

817 Silverstein Foundation to H.B. We would like to thank Ronit Cherki, Tamar Licht and

818 Anatoly Shapochnikov for excellent technical assistance.

Competing interests

820 The authors declare no competing interests.

References

Abudukeyoumu N, Hernandez-Flores T, Garcia-Munoz M, Arbuthnott GW. 2019.
Cholinergic modulation of striatal microcircuits. *Eur J Neurosci* **49**:604–622.
doi:10.1111/ejn.13949
Aceves Buendia J de J, Tiroshi L, Chiu W, Goldberg JA. 2019. Selective remodeling of
glutamatergic transmission to striatal cholinergic interneurons after dopamine
depletion. *Eur J Neurosci* **49**:824–833. doi:10.1111/ejn.13715
Alzheimer C, Schwindt P, Crill W. 1993. Modal gating of Na+ channels as a mechanism of
persistent Na+ current in pyramidal neurons from rat and cat sensorimotor cortex. *J*
Neurosci **13**:660–673. doi:10.1523/JNEUROSCI.13-02-00660.1993

Aoki S, Liu AW, Akamine Y, Zucca A, Zucca S, Wickens JR. 2018. Cholinergic
interneurons in the rat striatum modulate substitution of habits. *Eur J Neurosci*
47:1194–1205. doi:10.1111/ejn.13820
Aosaki T, Kimura M, Graybiel AM. 1995. Temporal and spatial characteristics of tonically
active neurons of the primate's striatum. *J Neurophysiol* **73**:1234–1252.
Aosaki T, Tsubokawa H, Ishida A, Watanabe K, Graybiel AM, Kimura M. 1994. Responses
of tonically active neurons in the primate's striatum undergo systematic changes
during behavioral sensorimotor conditioning. *J Neurosci* **14**:3969–3984.
Apicella P, Scarnati E, Schultz W. 1991. Tonically discharging neurons of monkey striatum
respond to preparatory and rewarding stimuli. *Exp Brain Res* **84**:672–675.
Arenkiel BR, Peca J, Davison IG, Feliciano C, Deisseroth K, Augustine GJ, Ehlers MD, Feng
G. 2007. In vivo light-induced activation of neural circuitry in transgenic mice
expressing channelrhodopsin-2. *Neuron* **54**:205–218.
doi:10.1016/j.neuron.2007.03.005
Assous M. 2021. Striatal cholinergic transmission. Focus on nicotinic receptors' influence in
striatal circuits. *Eur J Neurosci* **53**:2421–2442. doi:10.1111/ejn.15135
Assous M, Dautan D, Tepper JM, Mena-Segovia J. 2019. Pedunculopontine Glutamatergic
Neurons Provide a Novel Source of Feedforward Inhibition in the Striatum by
Selectively Targeting Interneurons. *J Neurosci* **39**:4727–4737.
doi:10.1523/JNEUROSCI.2913-18.2019
Assous M, Kaminer J, Shah F, Garg A, Koós T, Tepper JM. 2017. Differential processing of
thalamic information via distinct striatal interneuron circuits. *Nature Communications*
8:15860. doi:10.1038/ncomms15860
Beatty Joseph A., Song SC, Wilson CJ. 2015. Cell-type-specific resonances shape the
responses of striatal neurons to synaptic input. *Journal of Neurophysiology* **113**:688–
700. doi:10.1152/jn.00827.2014
Bennett BD, Callaway JC, Wilson CJ. 2000. Intrinsic membrane properties underlying
spontaneous tonic firing in neostriatal cholinergic interneurons. *J Neurosci* **20**:8493–
8503. doi:20/22/8493 [pii]
Bennett BD, Wilson CJ. 1999. Spontaneous activity of neostriatal cholinergic interneurons in
vitro. *J Neurosci* **19**:5586–5596.
Berger T, Larkum ME, Lüscher H-R. 2001. High I_h Channel Density in the Distal Apical
Dendrite of Layer V Pyramidal Cells Increases Bidirectional Attenuation of EPSPs.
Journal of Neurophysiology **85**:855–868. doi:10.1152/jn.2001.85.2.855
Bradfield LA, Bertran-Gonzalez J, Chieng B, Balleine BW. 2013. The Thalamostriatal
Pathway and Cholinergic Control of Goal-Directed Action: Interlacing New with
Existing Learning in the Striatum. *Neuron* **79**:153–166.
doi:10.1016/j.neuron.2013.04.039
Brown RE, Basheer R, McKenna JT, Strecker RE, McCarley RW. 2012. Control of Sleep
and Wakefulness. *Physiological Reviews* **92**:1087–1187.
doi:10.1152/physrev.00032.2011
Buzsáki G, Anastassiou CA, Koch C. 2012. The origin of extracellular fields and currents —
EEG, ECoG, LFP and spikes. *Nat Rev Neurosci* **13**:407–420. doi:10.1038/nrn3241
Carter AG, Sabatini BL. 2004. State-Dependent Calcium Signaling in Dendritic Spines of
Striatal Medium Spiny Neurons. *Neuron* **44**:483–493.
doi:10.1016/j.neuron.2004.10.013
Choi SJ, Ma TC, Ding Y, Cheung T, Joshi N, Sulzer D, Mosharov EV, Kang UJ. 2020.
Alterations in the intrinsic properties of striatal cholinergic interneurons after
dopamine lesion and chronic L-DOPA. *eLife* **9**:e56920. doi:10.7554/eLife.56920

Day M, Wokosin D, Plotkin JL, Tian X, Surmeier DJ. 2008. Differential Excitability and
Modulation of Striatal Medium Spiny Neuron Dendrites. *Journal of Neuroscience*
28:11603–11614. doi:10.1523/JNEUROSCI.1840-08.2008
Deng P, Zhang Y, Xu ZC. 2007. Involvement of I(h) in dopamine modulation of tonic firing
in striatal cholinergic interneurons. *J Neurosci* **27**:3148–3156. doi:27/12/3148 [pii]
10.1523/JNEUROSCI.5535-06.2007
Deng YP, Wong T, Bricker-Anthony C, Deng B, Reiner A. 2013. Loss of corticostriatal and
thalamostriatal synaptic terminals precedes striatal projection neuron pathology in
heterozygous Q140 Huntington's disease mice. *Neurobiol Dis* **60**:89–107.
doi:10.1016/j.nbd.2013.08.009
Ding J, Guzman JN, Tkatch T, Chen S, Goldberg JA, Ebert PJ, Levitt P, Wilson CJ, Hamm
HE, Surmeier DJ. 2006. RGS4-dependent attenuation of M4 autoreceptor function in
striatal cholinergic interneurons following dopamine depletion. *Nat Neurosci* **9**:832–
842. doi:10.1038/nn1700
Ding JB, Guzman JN, Peterson JD, Goldberg JA, Surmeier DJ. 2010. Thalamic Gating of
Corticostriatal Signaling by Cholinergic Interneurons. *Neuron* **67**:294–307.
doi:10.1016/j.neuron.2010.06.017
Ding Y, Won L, Britt JP, Lim SAO, McGehee DS, Kang UJ. 2011. Enhanced striatal
cholinergic neuronal activity mediates L-DOPA-induced dyskinesia in parkinsonian
mice. *Proceedings of the National Academy of Sciences* **108**:840–845.
doi:10.1073/pnas.1006511108
Doig NM, Magill PJ, Apicella P, Bolam JP, Sharott A. 2014. Cortical and Thalamic
Excitation Mediate the Multiphasic Responses of Striatal Cholinergic Interneurons to
Motivationally Salient Stimuli. *Journal of Neuroscience* **34**:3101–3117.
doi:10.1523/JNEUROSCI.4627-13.2014
Ermentrout B. 2002. Simulating, Analyzing, and Animating Dynamical Systems: A Guide to
XPPAUT for Researchers and Students. Philadelphia: Society for Industrial and
Applied Mathematics.
Eskow Jaunarajs KL, Bonsi P, Chesselet MF, Standaert DG, Pisani A. 2015. Striatal
cholinergic dysfunction as a unifying theme in the pathophysiology of dystonia.
Progress in Neurobiology **127–128**:91–107. doi:10.1016/j.pneurobio.2015.02.002
Goldberg JA, Deister CA, Wilson CJ. 2007. Response Properties and Synchronization of
Rhythmically Firing Dendritic Neurons. *Journal of Neurophysiology* **97**:208–219.
doi:10.1152/jn.00810.2006
Goldberg JA, Ding JB, Surmeier DJ. 2012. Muscarinic Modulation of Striatal Function and
Circuitry In: Fryer AD, Christopoulos A, Nathanson NM, editors. Muscarinic
Receptors, Handbook of Experimental Pharmacology. Berlin, Heidelberg: Springer
Berlin Heidelberg. pp. 223–241. doi:10.1007/978-3-642-23274-9_10
Goldberg JA, Kats SS, Jaeger D. 2003. Globus Pallidus Discharge Is Coincident with Striatal
Activity during Global Slow Wave Activity in the Rat. *J Neurosci* **23**:10058–10063.
doi:10.1523/JNEUROSCI.23-31-10058.2003
Goldberg JA, Reynolds JN. 2011. Spontaneous firing and evoked pauses in the tonically
active cholinergic interneurons of the striatum. *Neuroscience* **198**:27–43.
doi:10.1016/j.neuroscience.2011.08.067
Goldberg JA, Teagarden MA, Foehring RC, Wilson CJ. 2009. Nonequilibrium Calcium
Dynamics Regulate the Autonomous Firing Pattern of Rat Striatal Cholinergic
Interneurons. *Journal of Neuroscience* **29**:8396–8407.
doi:10.1523/JNEUROSCI.5582-08.2009
Goldberg JA, Wilson CJ. 2010. The Cholinergic Interneurons of the Striatum: Intrinsic
Properties Underlie Multiple Discharge Patterns. In: Steiner H, Tseng KY, editors.

Handbook of Basal Ganglia Structure and Function. London: Academic Press. pp.
133–149.

Goldberg JA, Wilson CJ. 2005. Control of Spontaneous Firing Patterns by the Selective
Coupling of Calcium Currents to Calcium-Activated Potassium Currents in Striatal
Cholinergic Interneurons. *Journal of Neuroscience* **25**:10230–10238.
doi:10.1523/JNEUROSCI.2734-05.2005

Gradinaru V, Mogri M, Thompson KR, Henderson JM, Deisseroth K. 2009. Optical
deconstruction of parkinsonian neural circuitry. *Science* **324**:354–359.
doi:10.1126/science.1167093

Graybiel A, Aosaki T, Flaherty A, Kimura M. 1994. The basal ganglia and adaptive motor
control. *Science* **265**:1826–1831. doi:10.1126/science.8091209

Gritton HJ, Howe WM, Romano MF, DiFeliceantonio AG, Kramer MA, Saligrama V,
Bucklin ME, Zemel D, Han X. 2019. Unique contributions of parvalbumin and
cholinergic interneurons in organizing striatal networks during movement. *Nat
Neurosci* **22**:586–597. doi:10.1038/s41593-019-0341-3

Harnett MT, Magee JC, Williams SR. 2015. Distribution and Function of HCN Channels in
the Apical Dendritic Tuft of Neocortical Pyramidal Neurons. *Journal of Neuroscience*
35:1024–1037. doi:10.1523/JNEUROSCI.2813-14.2015

Häusser M. 2021. Optogenetics — The Might of Light. *N Engl J Med* **385**:1623–1626.
doi:10.1056/NEJMcibr2111915

Helseth AR, Hernandez-Martinez R, Hall VL, Oliver ML, Turner BD, Caffall ZF, Rittiner
JE, Shipman MK, King CS, Gradinaru V, Gerfen C, Costa-Mattioli M, Calakos N.
2021. Cholinergic neurons constitutively engage the ISR for dopamine modulation
and skill learning in mice. *Science* **372**:eabe1931. doi:10.1126/science.abe1931

Higgs MH, Wilson CJ. 2017. Measurement of phase resetting curves using optogenetic
barrage stimuli. *Journal of Neuroscience Methods* **289**:23–30.
doi:10.1016/j.jneumeth.2017.06.018

Howe M, Ridouh I, Allegra Mascaro AL, Larios A, Azcorra M, Dombeck DA. 2019.
Coordination of rapid cholinergic and dopaminergic signaling in striatum during
spontaneous movement. *eLife* **8**:e44903. doi:10.7554/eLife.44903

Hutcheon B, Miura RM, Puil E. 1996. Subthreshold membrane resonance in neocortical
neurons. *Journal of Neurophysiology* **76**:683–697. doi:10.1152/jn.1996.76.2.683

Hutcheon B, Yarom Y. 2000. Resonance, oscillation and the intrinsic frequency preferences
of neurons. *Trends in Neurosciences* **23**:216–222. doi:10.1016/S0166-
2236(00)01547-2

Johansson Y, Silberberg G. 2020. The Functional Organization of Cortical and Thalamic
Inputs onto Five Types of Striatal Neurons Is Determined by Source and Target Cell
Identities. *Cell Reports* **30**:1178-1194.e3. doi:10.1016/j.celrep.2019.12.095

Joshua M, Adler A, Mitelman R, Vaadia E, Bergman H. 2008. Midbrain Dopaminergic
Neurons and Striatal Cholinergic Interneurons Encode the Difference between
Reward and Aversive Events at Different Epochs of Probabilistic Classical
Conditioning Trials. *Journal of Neuroscience* **28**:11673–11684.
doi:10.1523/JNEUROSCI.3839-08.2008

Kawaguchi Y. 1993. Physiological, morphological, and histochemical characterization of
three classes of interneurons in rat neostriatum. *J Neurosci* **13**:4908–4923.
doi:10.1523/JNEUROSCI.13-11-04908.1993

Kerr JND. 2004. Action Potential Timing Determines Dendritic Calcium during Striatal Up-
States. *Journal of Neuroscience* **24**:877–885. doi:10.1523/JNEUROSCI.4475-03.2004

Kim CK, Adhikari A, Deisseroth K. 2017. Integration of optogenetics with complementary
methodologies in systems neuroscience. *Nat Rev Neurosci* **18**:222–235.
doi:10.1038/nrn.2017.15
Kimura M, Rajkowski J, Evarts E. 1984. Tonically discharging putamen neurons exhibit set-
dependent responses. *Proceedings of the National Academy of Sciences* **81**:4998–
5001. doi:10.1073/pnas.81.15.4998
Koch C. 1984. Cable theory in neurons with active, linearized membranes. *Biol Cybern*
50:15–33. doi:10.1007/BF00317936
Kole MHP, Hallermann S, Stuart GJ. 2006. Single I_h Channels in Pyramidal Neuron
Dendrites: Properties, Distribution, and Impact on Action Potential Output. *J*
Neurosci **26**:1677–1687. doi:10.1523/JNEUROSCI.3664-05.2006
Kosillo P, Zhang Y-F, Threlfell S, Cragg SJ. 2016. Cortical Control of Striatal Dopamine
Transmission via Striatal Cholinergic Interneurons. *Cereb Cortex* **26**:4160–4169.
doi:10.1093/cercor/bhw252
Lapper SR, Bolam JP. 1992. Input From the Frontal Cortex and the Nucleus To Cholinergic
Interneurons in the Dorsal of the Rat **51**:533–545.
Liu D, Dan Y. 2019. A Motor Theory of Sleep-Wake Control: Arousal-Action Circuit. *Annu*
Rev Neurosci **42**:27–46. doi:10.1146/annurev-neuro-080317-061813
Mahon S. 2001. Relationship between EEG Potentials and Intracellular Activity of Striatal
and Cortico-striatal Neurons: an In Vivo Study under Different Anesthetics. *Cerebral*
Cortex **11**:360–373. doi:10.1093/cercor/11.4.360
Mallet N, Leblois A, Maurice N, Beurrier C. 2019. Striatal Cholinergic Interneurons: How to
Elucidate Their Function in Health and Disease. *Front Pharmacol* **10**:1488.
doi:10.3389/fphar.2019.01488
Mamaligas AA, Barcomb K, Ford CP. 2019. Cholinergic Transmission at Muscarinic
Synapses in the Striatum Is Driven Equally by Cortical and Thalamic Inputs. *Cell*
Reports **28**:1003-1014.e3. doi:10.1016/j.celrep.2019.06.077
Martin RF, Bowden DM. 2000. Primate brain maps: structure of the macaque brain. Elsevier.
Matityahu L, Malgady JM, Schirelman M, Johansson Y, Wilking JA, Silberberg G, Goldberg
JA, Plotkin JL. 2022. A tonic nicotinic brake controls spike timing in striatal spiny
projection neurons. *eLife* **11**:e75829. doi:10.7554/eLife.75829
Matsumoto N, Minamimoto T, Graybiel AM, Kimura M. 2001. Neurons in the Thalamic
CM-Pf Complex Supply Striatal Neurons With Information About Behaviorally
Significant Sensory Events. *Journal of Neurophysiology* **85**:960–976.
doi:10.1152/jn.2001.85.2.960
Maurice N. 2004. D2 Dopamine Receptor-Mediated Modulation of Voltage-Dependent Na+
Channels Reduces Autonomous Activity in Striatal Cholinergic Interneurons. *Journal*
of Neuroscience **24**:10289–10301. doi:10.1523/JNEUROSCI.2155-04.2004
Mauro A, Conti F, Dodge F, Schor R. 1970. Subthreshold Behavior and Phenomenological
Impedance of the Squid Giant Axon. *Journal of General Physiology* **55**:497–523.
doi:10.1085/jgp.55.4.497
McGuirt AF, Post MR, Pigulevskiy I, Sulzer D, Lieberman OJ. 2021. Coordinated Postnatal
Maturation of Striatal Cholinergic Interneurons and Dopamine Release Dynamics in
Mice. *J Neurosci* **41**:3597–3609. doi:10.1523/JNEUROSCI.0755-20.2021
Mizrahi-Kliger AD, Kaplan A, Israel Z, Bergman H. 2018. Desynchronization of slow
oscillations in the basal ganglia during natural sleep. *Proc Natl Acad Sci USA*
115:E4274–E4283. doi:10.1073/pnas.1720795115
Morris G, Arkadir D, Nevet A, Vaadia E, Bergman H. 2004. Coincident but distinct
messages of midbrain dopamine and striatal tonically active neurons. *Neuron* **43**:133–
143. doi:10.1016/j.neuron.2004.06.012 S0896627304003551 [pii]

Nagel G, Szellas T, Huhn W, Kateriya S, Adeishvili N, Berthold P, Ollig D, Hegemann P,
Bamberg E. 2003. Channelrhodopsin-2, a directly light-gated cation-selective
membrane channel. *Proc Natl Acad Sci USA* **100**:13940–13945.
doi:10.1073/pnas.1936192100
Oswald MJ, Oorschot DE, Schulz JM, Lipski J, Reynolds JN. 2009. IH current generates the
afterhyperpolarisation following activation of subthreshold cortical synaptic inputs to
striatal cholinergic interneurons. *J Physiol* **587**:5879–5897. doi:jphysiol.2009.177600
[pii] 10.1113/jphysiol.2009.177600
Parker PRL, Lalive AL, Kreitzer AC. 2016. Pathway-Specific Remodeling of Thalamostriatal
Synapses in Parkinsonian Mice. *Neuron* **89**:734–740.
doi:10.1016/j.neuron.2015.12.038
Paz RM, Tubert C, Stahl AM, Amarillo Y, Rela L, Murer MG. 2021. Levodopa Causes
Striatal Cholinergic Interneuron Burst-Pause Activity in Parkinsonian Mice. *Mov*
Disord **36**:1578–1591. doi:10.1002/mds.28516
Petreanu L, Mao T, Sternson SM, Svoboda K. 2009. The subcellular organization of
neocortical excitatory connections. *Nature* **457**:1142–5. doi:10.1038/nature07709
Pisani A, Bernardi G, Ding J, Surmeier DJ. 2007. Re-emergence of striatal cholinergic
interneurons in movement disorders. *Trends in Neurosciences* **30**:545–553.
doi:10.1016/j.tins.2007.07.008
Plotkin JL, Day M, Surmeier DJ. 2011. Synaptically driven state transitions in distal
dendrites of striatal spiny neurons. *Nat Neurosci* **14**:881–888. doi:10.1038/nn.2848
Plotkin JL, Goldberg JA. 2019. Thinking Outside the Box (and Arrow): Current Themes in
Striatal Dysfunction in Movement Disorders. *Neuroscientist* **25**:359–379.
doi:10.1177/1073858418807887
Poppi LA, Ho-Nguyen KT, Shi A, Daut CT, Tischfield MA. 2021. Recurrent Implication of
Striatal Cholinergic Interneurons in a Range of Neurodevelopmental,
Neurodegenerative, and Neuropsychiatric Disorders. *Cells* **10**:907.
doi:10.3390/cells10040907
Raz A, Feingold A, Zelanskaya V, Vaadia E, Bergman H. 1996. Neuronal synchronization of
tonically active neurons in the striatum of normal and parkinsonian primates. *J*
Neurophysiol **76**:2083–2088.
Rehani R, Atamna Y, Tiroshi L, Chiu W-H, de Jesús Aceves Buendía J, Martins GJ,
Jacobson GA, Goldberg JA. 2019. Activity Patterns in the Neuropil of Striatal
Cholinergic Interneurons in Freely Moving Mice Represent Their Collective Spiking
Dynamics. *eNeuro* **6**:ENEURO.0351-18.2018. doi:10.1523/ENEURO.0351-18.2018
Reig R, Silberberg G. 2014. Multisensory Integration in the Mouse Striatum. *Neuron*
83:1200–1212. doi:10.1016/j.neuron.2014.07.033
Remme MWH, Rinzel J. 2011. Role of active dendritic conductances in subthreshold input
integration. *J Comput Neurosci* **31**:13–30. doi:10.1007/s10827-010-0295-7
Riedner BA, Vyazovskiy VV, Huber R, Massimini M, Esser S, Murphy M, Tononi G. 2007.
Sleep Homeostasis and Cortical Synchronization: III. A High-Density EEG Study of
Sleep Slow Waves in Humans. *Sleep* **30**:1643–1657. doi:10.1093/sleep/30.12.1643
Schulz JM, Oswald MJ, Reynolds JNJ. 2011. Visual-Induced Excitation Leads to Firing
Pauses in Striatal Cholinergic Interneurons. *Journal of Neuroscience* **31**:11133–
11143. doi:10.1523/JNEUROSCI.0661-11.2011
Schulz JM, Reynolds JNJ. 2013. Pause and rebound: sensory control of cholinergic signaling
in the striatum. *Trends in Neurosciences* **36**:41–50. doi:10.1016/j.tins.2012.09.006
Sela Y, Vyazovskiy VV, Cirelli C, Tononi G, Nir Y. 2016. Responses in Rat Core Auditory
Cortex are Preserved during Sleep Spindle Oscillations. *Sleep* **39**:1069–1082.
doi:10.5665/sleep.5758

Sharott A, Doig NM, Mallet N, Magill PJ. 2012. Relationships between the Firing of Identified Striatal Interneurons and Spontaneous and Driven Cortical Activities In Vivo. *Journal of Neuroscience* **32**:13221–13236. doi:10.1523/JNEUROSCI.2440-12.2012

Smith Y, Raju DV, Pare JF, Sidibe M. 2004. The thalamostriatal system: a highly specific network of the basal ganglia circuitry. *Trends Neurosci* **27**:520–527. doi:10.1016/j.tins.2004.07.004 S0166-2236(04)00232-2 [pii]

Song WJ, Surmeier DJ. 1996. Voltage-dependent facilitation of calcium channels in rat neostriatal neurons. *J Neurophysiol* **76**:2290–2306.

Stern EA, Jaeger D, Wilson CJ. 1998. Membrane potential synchrony of simultaneously recorded striatal spiny neurons in vivo. *Nature* **394**:475–478. doi:10.1038/28848

Tanimura A, Du Y, Kondapalli J, Wokosin DL, Surmeier DJ. 2019. Cholinergic Interneurons Amplify Thalamostriatal Excitation of Striatal Indirect Pathway Neurons in Parkinson's Disease Models. *Neuron* **101**:444-458.e6. doi:10.1016/j.neuron.2018.12.004

Tanimura A, Lim SAO, Aceves Buendia J de J, Goldberg JA, Surmeier DJ. 2016. Cholinergic Interneurons Amplify Corticostriatal Synaptic Responses in the Q175 Model of Huntington's Disease. *Front Syst Neurosci* **10**. doi:10.3389/fnsys.2016.00102

Tchumatchenko T, Newman JP, Fong M, Potter SM. 2013. Delivery of continuously-varying stimuli using channelrhodopsin-2. *Front Neural Circuits* **7**. doi:10.3389/fncir.2013.00184

Thomas TM, Smith Y, Levey AI, Hersch SM. 2000. Cortical Inputs to m2-Immunoreactive Striatal Interneurons in Rat and Monkey. *Synapse* **37**:252–261.

Thorn CA, Graybiel AM. 2010. Pausing to Regroup: Thalamic Gating of Cortico-Basal Ganglia Networks. *Neuron* **67**:175–178. doi:10.1016/j.neuron.2010.07.010

Threlfell S, Lalic T, Platt NJ, Jennings KA, Deisseroth K, Cragg SJ. 2012. Striatal Dopamine Release Is Triggered by Synchronized Activity in Cholinergic Interneurons. *Neuron* **75**:58–64. doi:10.1016/j.neuron.2012.04.038

Tiroshi L, Goldberg JA. 2019. Population dynamics and entrainment of basal ganglia pacemakers are shaped by their dendritic arbors. *PLoS Comput Biol* **15**:e1006782. doi:10.1371/journal.pcbi.1006782

Tubert C, Murer MG. 2021. What's wrong with the striatal cholinergic interneurons in Parkinson's disease? Focus on intrinsic excitability. *Eur J Neurosci* **53**:2100–2116. doi:10.1111/ejn.14742

Ulrich D. 2002. Dendritic Resonance in Rat Neocortical Pyramidal Cells. *Journal of Neurophysiology* **87**:2753–2759. doi:10.1152/jn.2002.87.6.2753

Wilson C, Chang H, Kitai S. 1990. Firing patterns and synaptic potentials of identified giant aspiny interneurons in the rat neostriatum. *J Neurosci* **10**:508–519. doi:10.1523/JNEUROSCI.10-02-00508.1990

Wilson CJ. 2005. The mechanism of intrinsic amplification of hyperpolarizations and spontaneous bursting in striatal cholinergic interneurons. *Neuron* **45**:575–585.

Wilson CJ. 2004. Basal Ganglia In: Shepherd GM, editor. The Synaptic Organization of the Brain. New York: Oxford University Press. pp. 361–413.

Wilson CJ, Goldberg JA. 2006. Origin of the slow afterhyperpolarization and slow rhythmic bursting in striatal cholinergic interneurons. *J Neurophysiol* **95**:196–204. doi:00630.2005 [pii] 10.1152/jn.00630.2005

Yamanaka K, Hori Y, Minamimoto T, Yamada H, Matsumoto N, Enomoto K, Aosaki T, Graybiel AM, Kimura M. 2018. Roles of centromedian parafascicular nuclei of thalamus and cholinergic interneurons in the dorsal striatum in associative learning of

environmental events. *J Neural Transm* **125**:501–513. doi:10.1007/s00702-017-1713-z

Yarom O, Cohen D. 2011. Putative Cholinergic Interneurons in the Ventral and Dorsal Regions of the Striatum Have Distinct Roles in a Two Choice Alternative Association Task. *Front Syst Neurosci* **5**. doi:10.3389/fnsys.2011.00036