1 **Mapping nationally and globally at-risk species to identify hotspots for (and**
2 **gaps in) conservation**
3
4

Marie E Hardouin[1] (https://orcid.org/0000-0003-1524-3278, marie.hardouin@mail.mcgill.ca)

Anna L Hargreaves[1]* (https://orcid.org/0000-0001-9680-9696, anna.hargreaves@mcgill.ca)

1. Department of Biology, McGill University, 1205 Dr. Penfield Ave, Montreal Quebec,

Canada, H3A 1B1

* = corresponding author. +1 514 398 7401

ABSTRACT

Protecting habitat of species-at-risk is critical to their recovery, but can be contentious. For

example, protecting species that are locally imperilled but globally common (e.g. species that

only occur in a jurisdiction at the edge of their geographic range) is often thought to distract from

protecting globally-imperilled species. However, such perceived trade-offs are based on the

assumption that threatened groups have little spatial overlap, which is rarely quantified. Here, we

compile range maps of terrestrial species-at-risk in Canada to assess the geographic overlap of

nationally and globally at-risk species with each other, among taxonomic groups, and with

protected areas. While many nationally-at-risk taxa only occurred in Canada at their northern

range edge (median=4% of range in Canada), nationally-at-risk species were not significantly

more peripheral in Canada than globally-at-risk species. Further, 56% of hotspots of nationally-

at-risk taxa were also hotspots of globally-at-risk taxa in Canada, undercutting the perceived

trade-off in their protection. Hotspots of nationally-at-risk taxa also strongly overlapped with

hotspots of individual taxonomic groups, though less so for mammals. While strong spatial

overlap across threat levels and taxa should facilitate efficient habitat protection, <7% of the area

in Canada's at-risk hotspots is protected, and more than 70% of nationally and globally-at-risk

species in Canada have <10% of their Canadian range protected. Our results counter the

perception that protecting nationally vs. globally at-risk species are at odds, and identify critical

areas to target as Canada strives to increase its protected areas and promote species-at-risk

recovery.

Keywords: Canada, protected areas, species at risk, conservation hotspots, peripheral species,

1. INTRODUCTION

Habitat destruction is one of the primary threats to global biodiversity (Baillie et al., 2004). Indeed, the loss and degradation of habitat is the main threat to 85% of species listed in the International Union for the Conservation of Nature's Red List of Threatened Species (IUCN, 2015) and a leading barrier to the recovery of imperilled species (Kerr & Deguise, 2004; Vellend, 2003). Protecting habitat through the establishment of protected areas is thus a crucial component of species-at-risk conservation. At the local scale, protected areas can alleviate land-conversion pressures on existing populations of species at risk, and at the landscape scale they can provide crucial habitat corridors that could facilitate species range shifts in response to climate change (Littlefield et al., 2019; Thomas et al., 2012).

While protected areas are a key conservation tool, it can be challenging to establish them in the areas most important for mitigating biodiversity loss. Habitat of species-at-risk often coincides with areas of high human population density or land use (Allan et al., 2019; Caissy et al., 2020), potentially generating mismatches between economic and conservation priorities. Combined with a lack of political will or funding (Watson et al., 2014), this can result in imperiled taxa being poorly represented within protected area networks (Venter et al., 2014). Given these challenges, optimizing which species-at-risk habitat is proposed for protection is crucial. A promising prioritization strategy is to identify 'hotspots' with particularly high taxonomic richness. This approach has been applied to specific taxonomic groups (Cameron & Hargreaves, 2020; Ceballos & Ehrlich, 2006; Fattorini et al., 2012; Orme et al., 2005), but it is unclear how well hotspots overlap among taxa. For instance, there is limited overlap in hotspots of vertebrates and plants in China (Xu et al., 2018) and of birds, mammals, and invertebrates in the UK

(Prendergast et al., 1993). Poor hotspot overlap among taxa complicates prioritizing areas for

conservation.

Even when hotspots align among taxa, prioritizing hotspots of at-risk species may be

controversial when species are rare (and therefore deemed at-risk) in one jurisdiction but

widespread elsewhere. Species can be rare in a jurisdiction simply because they only occur there

at the edge of their geographic range (henceforth "peripheral" taxa; Bunnell et al., 2004). This

can focus regional conservation efforts on range-edge populations of species that are globally

secure, particularly in high-latitude countries that often contain the poleward range-edge of

species more common toward the equator (Gibson et al., 2009). For instance, >75% of Canada's

nationally at-risk plants and Finland's rare beetles are largely distributed outside these countries

(Caissy et al., 2020; Komonen, 2007), and most US-state lists of threatened birds are dominated

by taxa with low global risk (Wells et al., 2010). Protecting locally imperilled species has

therefore been accused of being 'parochial', and coming at the expense of protecting more

endemic or globally imperilled species (Hunter & Hutchinson, 1994; Rodrigues & Gaston,

2002). However, whether such a trade-off really exists if often unknown. Hotspots of locally at-

risk species can co-occur with hotspots of overall diversity (Cameron & Hargreaves, 2020), and

the extent to which hotspots of locally-at-risk and globally-at-risk species co-occur has rarely if

ever been formally assessed.

An excellent case study for exploring how well hotspots of at-risk taxa co-occur, particularly

between nationally and globally at-risk species, is Canada. As the world's second largest country

and the northernmost country in the Americas, habitat protection in Canada is critical for the

species already there and the many predicted to shift northward under climate change. Because

many nationally-at-risk taxa are peripheral in Canada (though this has only been quantified for

plants and mammals), critics have argued that prioritizing them for conservation detracts from

conserving globally-at-risk taxa (Kraus et al., 2021; Raymond et al., 2018). However, the

geographic overlap between nationally and globally at-risk species has not been quantified.

Further, while we know that Canada's nationally at-risk taxa cluster in southern Canada

(Coristine et al., 2018; Gibson et al., 2009), it is unclear whether this is driven by species-rich

taxonomic groups (e.g. vascular plants, which cluster south; Caissy et al., 2020), and therefore

how well overall hotspots of at-risk species would reflect hotspots of less species-rich groups

(e.g. mammals, which have northwestern clusters; Cameron & Hargreaves, 2020). Identifying

conservation hotspots in Canada is also particularly timely. Habitat loss threatens more than

three-quarters of Canada's nationally at-risk species (Woo-Durand et al., 2020), many of which

have very little range overlap with protected areas (Caissy et al., 2020; Kraus & Hebb, 2020),

and Canada has recently committed to protecting 25% of its land by 2025, up from the 12.5%

conserved in 2020 (ECCC, 2021a; ECCC, 2021b).

Using range maps of Canada's nationally and globally at-risk terrestrial taxa, we run three sets of

analyses related to their conservation. *1) Hotspots*. We identify where species-at-risk are

concentrated in Canada, and ask (Q1): How well do hotspots overlap among taxonomic groups,

and between nationally and globally at-risk species? We predicted that most hotspots of

nationally at-risk species would occur in southern Canada, where Canada's biodiversity, human

population, and human land use all cluster (Coristine et al., 2018; Kraus & Hebb, 2020). We did

not have specific predictions about overlap given the distinct distributions of nationally at-risk

plants and mammals (Caissy et al., 2020; Cameron & Hargreaves, 2020) and general sense that

Canada's globally-at-risk species are often in the far north. *2) Peripherality.* We quantify the

proportion of each at-risk taxon's range in Canada, and ask (Q2): Does peripherality (the extent

to which taxa only occur in Canada at their range edge) vary among threat statuses and

taxonomic groups, and between nationally and globally at-risk species? We expected that

nationally-at-risk taxa would be mostly peripheral, and significantly more peripheral than

globally-at-risk taxa, since national conservation assessment focuses on Canadian range area

whereas the IUCN assesses global range area (COSEWIC, 2019; IUCN, 2012). *3) Habitat*

protection. (Q3) How well is Canada protecting at-risk taxa? As many nationally at-risk species

occur in highly human-modified southern Canada (Coristine et al., 2018; Coristine & Kerr,

2011), we were not optimistic about overlap with protected areas.

2. METHODS

Eligibility for protection under Canada's Species at Risk Act (SARA) is determined by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC). COSEWIC has seven subcommittees dedicated to assessing nine terrestrial taxonomic groups (Table 1) using quantitative criteria established by the IUCN. Taxa can be species, subspecies or populations. COSEWIC first prioritizes which taxa to assess, and at this stage prioritizes taxa likely to become globally extinct (SARA 15.1b). For prioritized taxa, COSEWIC commissions, reviews and approves an assessment report based on the 'best biological information' available (SARA, 2002), then uses the report to recommend a threat status. COSEWIC initially considers only Canadian populations, then reviews whether adjustments are warranted given the likelihood of demographic rescue from populations outside Canada (Environment and Climate Change Canada, 2017). Here we consider species assessed as nationally 'at-risk', i.e. Special Concern (may become threatened or endangered); Threatened (likely to become endangered if threats are not mitigated); or Endangered (facing imminent extirpation or extinction), and ignore taxa deemed Extirpated, Extinct, or Data deficient (i.e. no current Canadian range or reliable range map) or Not at risk. The final decision to list and protect taxa under SARA rests with the federal government after considering the socioeconomic implications (SARA, 2002). We use COSEWIC rather than SARA designations as they more closely reflect biology.

131 **Table 1.** Sample sizes for at-risk terrestrial taxa in Canada. Nationally at-risk taxa are those with a COSEWIC status of Special
132 Concern, Threatened or Endangered. Globally at-risk taxa are those with an IUCN status of Vulnerable, Endangered or Critically
133 Endangered. Range maps in C and D were obtained from COSEWIC for all taxa except birds, whose maps were obtained from
134 BirdLife International and so are independent of sample sizes in A and B. C, D and G indicate the final sample sizes for Questions 1,
135 2, and 3. Missing Canadian range maps for globally at-risk species in F were obtained by digitizing COSEWIC maps when available.

	Nationally at-risk Taxa				Globally at-risk Taxa		
	A. Populations with a COSEWIC status	B. Taxa in A with a COSEWIC range map[*]	C. Taxa with a digitized Canadian range map (Q1, Q3[**])	D. Taxa in C with a digitized global range map (Q2)[***]	E. Taxa in Canada with an IUCN status	F. Taxa with an IUCN range polygon	G. Taxa with a digitized range map (Q2)
Birds	76	NA	56	54	10	10	10
Mammals	42	41	37	36	7	7	7
Reptiles	41	34	32	26	3	1	3
Amphibians	27	24	21	17	0	0	0
Arthropods	70	66	60	54	12	6	9
Molluscs	40	39	34	32	17	14	15
Vascular Plants	207	200	184	171	21	15	17
Mosses	20	20	17	10	0	0	0
Lichens	23	23	21	14	6	2	3
Total	546	NA	462	414	76	55	64

[*] If populations in A) were from the same species and only one range map was provided for that species, they were merged in B).
[**] Q3: analyses of hotspot overlap with protected areas uses n in C; analyses of range overlap with protected areas by threat status has
slightly lower n, as populations of the same species with different threat statuses had to be excluded.
[***] If the populations merged in B) did not have the same COSEWIC status, they were excluded.

2.1 Data Collection

<u>2.1.1 COSEWIC Range Maps</u>

COSEWIC assessment reports are publicly available as PDFs and generally include a range map

of the assessed species (Government of Canada, 2021). For all taxa except birds (see 2.1.2) we

updated a database developed by Pascale Caissy (see Caissy et al., 2020) by digitizing range

maps in COSEWIC reports using the Quantum GIS 3.16 geographic information software (QGIS

Development Team, 2021). We digitized the global range maps when possible, but 34 taxa had

global range maps that were not properly digitizable (maps were incomplete, imprecise, or in a

projection that could not be digitized), but Canadian range maps that were. For these taxa we

digitized Canadian ranges only. These taxa were excluded from peripherality analyses (Q2) but

included in the hotspot and protected area analyses (Q1, Q3) for which only the Canadian ranges

were needed (Table 1).

To digitize COSEWIC maps, we first georeferenced maps by associating ≥15 (generally 40-60)

specific points on the PDF maps with their corresponding coordinates on a base map of the

world; these points were clear landmarks such as sharp land mass corners, waterbody edges,

roads, or jurisdictional boundaries. Maps then underwent a thin plane spline transformation to

project them in the World Geodetic System 1984 projection (WGS 84, EPSG:4326), a common

global latitude and longitude-based coordinate reference system (Caissy et al., 2020; Kennedy &

Kopp, 2000).

When ranges were already represented as polygons, we created a polygon shapefile by tracing

the outline by hand; for ranges represented as point occurrences, we created a point shapefile by

adding each point. If there were at least 4 points in the Americas, we generated a convex polygon

(the smallest polygon shape encompassing all the points), which generally matched the taxon's

extent of occupancy as described in the text of COSEWIC assessment reports. If there were <4

point occurrences and we were unable to generate a convex polygon (12 taxa), we generated a

buffer around each point instead. We set the buffer distance for each taxon to ensure that the

taxon's total buffered area would be roughly equivalent to the extent of occupancy reported in its

COSEWIC report. We cut out the Great Lakes area from all range maps except for mollusc taxa

that were present in these lakes.

2.1.2 BirdLife Range Maps

Most bird species with a COSEWIC assessment are migratory, so their range maps had to be

divided between their resident, breeding, non-breeding, and passage (e.g. migratory stopovers)

ranges to meaningfully compare their Canadian and global distributions. We obtained range

maps with this information from BirdLife International, who provided us with an ESRI File

Geodatabase containing a separate distribution polygon shapefile for each seasonal range of each

species (BirdLife International and Handbook of the Birds of the World, 2019). From these, we

retained only ranges that overlapped a polygon shapefile of Canada (in QGIS3.16; QGIS

Development Team, 2021). For species that spend multiple seasons in Canada (e.g. both

breeding and passage) we merged the seasonal polygons and used this merged shapefile for

analyses. Thus, we compared each species' Canadian range with their corresponding seasonal

global range. For instance, if a species only uses land in Canada during its breeding and passage

season, we compared its Canadian breeding + passage range to its global breeding + passage

range.

<u>2.1.3 IUCN Range Maps</u>

To map globally at-risk species in Canada, we searched the IUCN Red List of Threatened

Species database (IUCN, 2021) for species in the categories Vulnerable, Endangered or

Critically Endangered (i.e. 'at-risk'), whose 'land region' included Canada, and assessment

scope = "global". IUCN assessment reports are publicly available as PDFs, and most have

corresponding polygon shapefiles of the species' range. We downloaded these polygons,

excluding 9 species whose IUCN reports were flagged as needing updating and 4 species whose

Canadian ranges were defined as "vagrant" and did not appear in their global range maps. For

each species we removed polygon areas defined by IUCN as "possibly extinct". For species

without a distribution polygon provided, we digitized the species' range from its COSEWIC

report if available (9 non-bird species; Table 1).

2.2 Data Extraction

Data extraction and analyses were done in R (version 4.1.0, R Core Team, 2021). To identify

hotspots with unusually high concentrations of at-risk taxa, we projected each at-risk taxon's

digitized range map in the Albers equal area conic projection, which is commonly used for

Canada (Kennedy & Kopp, 2000). We then overlaid these maps on a map of Canada divided into

100 x 100 km grid cells (1276 cells total) and counted the number of nationally and globally at-

risk species in each grid cell. For each taxonomic grouping (the nine taxonomic groups of

nationally-at-risk taxa, nationally-at-risk taxa overall, and globally-at-risk taxa overall) we

identified cells with the highest species richness ("hotspots"), up to a maximum of 5% of the

1276 grid cells (max. 63 cells) (Cameron and Hargreaves 2020; adapted from Prendergast et al.

(1993) Reid (1998)). E.g. cells containing 5 nationally-at-risk mammal taxa were not counted as

mammal hotspots, as doing so would have resulted in >63 hotspot cells, leaving us with 55

mammal hotspot cells each with 6 or more at-risk mammal taxa. Thus the number of hotspot

cells differed among taxonomic groups (42 to 62 hotspot cells per group).

For peripherality analyses, we projected digitized range maps in the World Mollweide equal area

projection (WGS 84, ESRI:54009) to accommodate taxa with worldwide ranges. To assess

peripherality, we quantified the proportion of each taxon's western hemisphere range that occurs

in Canada (i.e. the inverse of peripherality). We deemed western-hemisphere range to be a more

biologically relevant denominator than global range, since the difficulty of moving between

hemispheres makes it unlikely that populations on other continents would substantially impact

Canadian populations. To calculate Canadian and western hemisphere range areas (km^2), we

intersected each digitized map with a Canada or western hemisphere (North, Central and South

America) boundary map respectively (Natural Earth, 2020), thus cropping all maps to land only.

We calculated the proportion of each taxon's Canadian range and the proportion of each hotspot

that overlapped officially protected areas using the Canadian Protected and Conserved Areas

Database (ECCC, 2021b). This publicly available database includes polygon shapefiles of

"terrestrial protected areas" and "terrestrial areas with other effective area-based conservation

measures" (ECCC, 2021b). We projected these protected area shapefiles in the Albers equal area

conic projection. We then intersected the protected area shapefiles with 1) each at-risk taxon's

range polygon to calculate the km^2 of protected habitat within each Canadian range; and 2) the

map of Canada divided into 100 x 100 km grid cells from Q1 to calculate the km^2 of protected

habitat within each cell. We then divided the area of protected habitat within each grid cell by the

grid cell's total area, which was not always 100 x 100 km as some grid cells were cut by

Canada's borders.

2.3 Data Analyses

Analyses assessed differences among taxonomic groups for nationally-at-risk species, and

compared nationally and globally at-risk species. We did not split analyses of globally-at-risk

species by taxonomic group, due to small sample sizes in several groups (Table 1). Data used in

analyses will be publicly archived on acceptance.

2.3.1 Hotspot analyses

To assess hotspot overlap among pairs of groups (e.g. birds and mosses, nationally-at-risk and

globally-at-risk), we counted the number of cells that were hotspots for both groups, then

calculated the percentage of shared hotspots. As different taxonomic groups had different

numbers of hotspots (see 2.2 above), we calculated overlap twice, one with each group as a

denominator, then used their average. For example, birds (48 hotspots) and mosses (42 hotspots)

shared 6 hotspot cells, resulting in a mean % overlap of 13% (i.e. the mean of 6/48 x 100 and

6/42 x 100).

2.3.2 Peripherality analyses

We tested whether the proportion of a taxon's western hemisphere range in Canada (proportional

response) varied among taxonomic groups and threat statuses using generalized linear models

(GLMs) with a beta-regression error structure (*glmmTMB* package version 1.1.2; Brooks et al.,

2017). Beta-regressions deal with proportional data that does not result from a binomial process

(such as percent areas), but cannot handle 0s and 1s. To deal with endemic species (i.e. range

proportion in Canada = 1), we transformed all 'range proportion in Canada' values according to

the equation (Douma & Weedon, 2019):

$$p^* = \frac{p(n-1) + \frac{1}{c}}{n}$$

where p^* is the proportion ('range proportion in Canada'), n is the total number of observations

in the dataset (number of taxa) and c is the number of categories (2, i.e. range proportion in

Canada and range proportion outside Canada).

First, we tested whether the 'range proportion in Canada' differed among COSEWIC statuses

and/or taxonomic groups of nationally-at-risk taxa. Each nationally-at-risk taxon contributed one

data point (sample sizes in Table 1D) so we used a generalized linear model (*Model 1*: prop

range in Canada ~ COSEWIC status x taxonomic group). We grouped vascular plants and

mosses together ('plants') due to the small number of at-risk mosses (10 taxa; Table 1). Here and

for all beta-regression models (see below) we tested the significance of fixed effects using

likelihood ratio tests, which compare models with and without the fixed effect, compared to a χ^2

distribution (*anova* function in R). As the 2-way interaction was not significant ($\chi^2_{df14} = 16.8$, $P =$

0.26), we dropped it from the final model to facilitate interpretation of main effects. For

significant main effects, we used least-squared mean contrasts to determine which taxonomic

groups or threat statuses differed from each other (*lsmeans* package version 2.30.0; Lenth, 2016).

Second, we tested the widely-held assumption that nationally-at-risk species are more peripheral

in Canada than globally-at-risk species. We ran two models, each with one fixed effect, 'threat

status'. In *Model 2*, threat status had 6 categories (3 from COSEWIC and 3 from IUCN), to test

whether peripherality varied with level of endangerment. In *Model 3,* we collapsed these into 2

categories 'nationally-at-risk' or 'globally-at-risk'. In both models, each nationally-at-risk taxon

and each globally-at-risk taxon contributed one data point (sample sizes in Table 1D&G).

Because 42 taxa are considered at-risk by both COSEWIC and IUCN, and therefore occur in the

data twice, we used a generalized linear mixed model with a random intercept for 'species' (prop

range in Canada ~ threat status + (1| species)). Significance of 'threat status' and differences

among statuses were determined as in Model 1.

<u>2.3.3 Protected areas analyses</u>

We tested how well hotspots of at-risk species were protected compared to less species-rich cells

using beta regression GLMs. The response (proportion of a cell protected) was transformed as in

2.3.2 to deal with 0s and 1s (none vs. all of a cell protected). We tested whether the proportion of

a grid cell that was protected (response) differed depending on whether or not the cell was a

hotspot (binary predictor; model structure = prop cell protected ~ hotspot status), with one model

for nationally-at-risk hotspots (*Model 4*) and one for globally-at-risk hotspots (*Model 5*). Second,

we tested whether the proportion of the cell that was protected varied with the number of at-risk

taxa in the grid cell. Visual examination of the data suggested that this relationship might vary

between hotspot and non-hotspot cells, so we included an interaction with hotspot status (prop

cell protected ~ # at-risk taxa x hotspot status). While this model also assessed the effect of

hotspot status, we derive that effect from Models 4 and 5 as hotspot status is related to # at-risk

taxa. We again ran one model for nationally-at-risk taxa (*Model 6)* and one for globally-at-risk

taxa (*Model 7)*. We obtained backtransformed fit lines and 95% confidence intervals using the

visreg package (version 2.7.0, Breheny & Burchett, 2017).

Finally, we tested how well ranges of at-risk species were being protected and whether this

varied with threat status. We ran one beta regression model for nationally-at-risk taxa (*Model 8:*

prop Canadian range protected ~ COSEWIC status) and one for globally-at-risk taxa (*Model 9:*

prop Canadian range protected ~ IUCN status).

3. RESULTS

3.1 Hotspots of at-risk taxa

In Canada, hotspots of nationally-at-risk taxa generally clustered along the southern border

(Figure 1, top nine maps). This was true for most taxonomic groups, with the notable exceptions

of at-risk mammals, whose diversity hotspots followed the western mountains, and at-risk

lichens, whose hotspots were mostly coastal (Figure 1). When all of Canada's nationally at-risk

taxa in our data were combined (bottom left map in Figure 1), their hotspots clustered along

Canada's southern borders from western Canada and the prairies to the Great Lakes and St-

Lawrence river regions.

Hotspots showed substantial taxonomic overlap. Taxonomic groups shared 31.5% of their

hotspot cells on average (Figure 1). Overlap between hotspots of individual taxonomic groups

and hotspots of nationally-at-risk taxa overall was even higher (mean = 50.1%). More than three

quarters of the hotspots of at-risk reptiles, arthropods and vascular plants overlapped with

nationally-at-risk hotspots (Figure 1). Hotspots of nationally-at-risk taxa contained up to 132 at-

risk species, and 40% contained at-risk species from all 9 taxonomic groups. Together, the 4.4%

of land in Canada included in nationally-at-risk hotspots was home to 80% of the 462 nationally-

at-risk taxa in our data, including: 97% of the reptile taxa, 95% of amphibians, 92% of

arthropods, 89% of birds, 85% of molluscs, 76% of plants, 65% of mosses, 65% of mammals,

and 62% of lichens. While groups with the most at-risk species tended to have high overlap with

national hotspots, overlap was not solely driven by richness. For example, there are twice as

many at-risk vascular plant taxa in our data (184) than the next highest group (arthropods, 60

taxa), but arthropod hotspots had higher overlap with nationally-at-risk hotspots (84% vs. 70%;

Figure 1).

Globally at-risk taxa in Canada showed a remarkably similar distribution to nationally-at-risk

taxa (Figure 1, bottom two maps). Contrary to our expectation that globally-at-risk taxa might be

distributed farther north, hotspots of globally-at-risk taxa also clustered along Canada's southern

border, but were almost entirely in eastern Canada from the Great Lakes to the southern

Maritimes. More than half (56%) of globally at-risk hotspots were also hotspots of nationally at-

risk taxa. Together, these 34 overlapping hotspots, representing only 2.4% of Canada's land area,

were home to 64% of the globally-at-risk taxa and 43% of the nationally-at-risk taxa in our data.

Figure 1. Hotspots of at-risk taxa in Canada. Maps are divided into 1276 100 x 100km grid cells,

which are shaded in a greyscale according to the number of at-risk taxa they contain (greyscales

differ among maps; legend below each map shows the maximum number of at-risk taxa per grid

cell). Top three rows of maps show nationally-at-risk taxa separated by taxonomic group (yellow

icons = vertebrates; orange = invertebrates; green = plants; blue = lichens). Bottom two maps

show all nationally-at-risk (left) and globally-at-risk (right) taxa in Canada across taxonomic

groups. Hotspot cells (i.e. those with the most at-risk taxa up to a maximum of 5% of the total

grid cells) are outlined in red. The matrix of overlap (bottom right) indicates the percentage of

hotspot cells that coincide for each pairwise combination.

3.2 Peripherality of at-risk taxa in Canada

Most nationally-at-risk taxa only occur in Canada at the northern edge of their range (Figure 2A).

While most taxonomic groups contained some endemic species (47 nationally-at-risk species had

100% of their western-hemisphere range in Canada, e.g. Vancouver Island Marmot, Banff

Springs Snail, Magdalen Islands Grasshopper), 71% of nationally-at-risk taxa had less than 20%

of their western-hemisphere range in Canada (mean = 23.8% of range in Canada, median =

4.4%). The extent to which taxa were peripheral in Canada varied among COSEWIC threat

statuses (χ^2_{df2} = 15.1, $P < 0.001$), and taxonomic groups (χ^2_{df7} = 21.7, $P = 0.003$; *Model 1*). At-

risk mammals had significantly more of their range in Canada (i.e. were less peripheral) than at-

risk reptiles, amphibians, or plants (Figure 2C). The most imperilled taxa had significantly less

of their range in Canada (i.e. were more peripheral) than the least imperilled at-risk taxa (Figure

2D).

Contrary to our predictions, many globally at-risk taxa also occurred in Canada only at their

northern range edge. 54% of globally-at-risk taxa in Canada had less than 20% of their western

hemisphere range in Canada (mean = 31.2% of range in Canada, median = 17.4%). All IUCN

and COSEWIC threat categories contained both taxa with <5% of their range in Canada, and

taxa that are endemic or almost endemic (IUCN Vulnerable taxa had up to 86.4% (for reindeer)

of their range) in Canada. The extent to which taxa were peripheral in Canada varied

significantly among the 6 threat statuses (*Model 2*: χ^2_{df5} = 22.0, $P = < 0.001$; Figure 2D coloured

bars), but only because nationally Endangered taxa had less of their range in Canada than Special

Concern taxa, as found in *Model 1*. No category of nationally-at-risk taxa was more peripheral

than any category of globally-at-risk taxa, nor did peripherality differ between nationally- and

globally-at-risk taxa overall (*Model 3*; χ^2_{df1} = 2.8, P = 0.09; Figure 2D grey bars).

Figure 2. Geographic distribution and peripherality of at-risk taxa in Canada (that had digitizable

Canadian and global range maps; see Table 1). Maps show Canadian ranges of **(A)** nationally-at-

risk taxa, separated by their COSEWIC threat status, and **(B)** globally-at-risk taxa, combined

across IUCN at-risk threat statuses. Coloured dots correspond to threat status, from least (purple)

to most imperilled (red). **C)** Nationally-at-risk taxonomic groups differed in how peripheral they

were; bars without shared letters are significantly different (*Model 1*). **D)** Nationally-at-risk taxa

were not significantly more peripheral in Canada than globally-at-risk taxa when compared

among the 6 threat statuses (*Model 2*; coloured bars) or between nationally and globally-at-risk

taxa overall (*Model 3*; grey bars). Box plots show the median (thick middle line), 25th and 75th

percentiles (box), 1.5x the interquartile range (whiskers), and raw data (horizontally jittered

points).

3.3 Habitat protection of at-risk taxa

Areas with particularly high concentrations of at-risk species (at-risk hotspots) are not

particularly well protected. Whereas hotspots of nationally and globally at-risk taxa cluster in

southern Canada, Canada's largest expanses of protected habitat are further north (Figure 3). The

percentage of a grid cell protected did not differ between cells that were hotspots of nationally-

at-risk taxa (mean = 8.0 ± 12.3% of cell protected, median = 3.5%) and cells that were not (mean

= 12.9 ± 22.6%, median = 2.4%; *Model 4*: χ^2_{dfl} = 1.2, *P* = 0.27), nor between cells that were

hotspots of globally-at-risk taxa (mean = 4.7 ± 5.9%, median of 2.7%) and cells that were not

(mean = 13.1 ± 22.6%, median = 2.5%; *Model 5*: χ^2_{dfl} = .51, *P* = 0.48). For both nationally and

globally at-risk taxa, the percentage of a grid cell protected increased as the number of at-risk

taxa in the cell increased, but only for non-hotspot cells (significant interaction # at-risk taxa x

hotspot status: *Model 6*: χ^2_{dfl} = 32.7, *P* < 0.001; *Model 7:* χ^2_{dfl} = 18.9, *P* < 0.001; Figure 3C&D).

Of the 1276 grid cells across Canada, 34% contained no protected areas, while 0.5% were fully

protected. Only 6.8% of the land in nationally-at-risk hotspots was protected. Protection is even

lower for land in hotspots of globally-at-risk taxa (5.1% protected), and lower still for areas that

were hotspots for both nationally and globally-at-risk taxa (3.8% protected).

The Canadian ranges of individual at-risk taxa were also poorly protected on average. Nationally

and globally at-risk taxa in Canada had only 5.5% and 5.7% (median) of their Canadian range

protected, respectively. Most nationally-at-risk taxa (74%) and globally at-risk taxa (83%) had

less than 10% of their Canadian range protected. This is despite the fact that many at-risk taxa

have small range areas in Canada (median range area in Canada = 12,157 km^2 for nationally-at-

risk taxa and 105,634 km^2 for globally-at-risk taxa). Habitat protection differed among

COSEWIC statuses, with the most imperilled taxa having *less* of their Canadian range protected

than the least imperilled at-risk taxa (*Model 8*: $\chi^2_{df2} = 15.3$, $P < 0.001$; Figure 3E). Habitat

protection was equally low among IUCN statuses (*Model 9*: $\chi^2_{df2} = 0.39$, $P = 0.82$; Figure 3F).

Figure 3. Habitat protection of nationally-at-risk taxa (left) and globally-at-risk taxa (right) in

Canada. **A-B)** location of Canada's publicly protected and conserved areas (green), and hotspots

of at-risk taxa (from Figure 1) outlined in red. **C-D)** protection increased significantly with the

number of at-risk taxa in a grid cell, but only for cells that were not hotspots of at-risk taxa

(black). Each point represents one of the 1276 grid cells in Canada (raw data), trend lines and

95% confidence intervals are extracted from *Model 6* (C) and *Model 7* (D). **E-F)** Percentage of

at-risk taxa's Canadian range included in protected areas. Differing letters represent significant

differences among COSEWIC threat statuses (*Model 8*; E) or IUCN threat statuses (*Model 9;* F).

Sample sizes are in Table 1. Box plot formats are as in Figure 2.

4. DISCUSSION

Our analyses of Canada's at-risk terrestrial taxa support some preconceptions about their spatial distribution, but counter others. First, our results confirm that most species considered at-risk in Canada only occur in Canada at the northern edge of their range (Figure 2). This aligns with previous qualitative assessments of species-at-risk in Canada (Gibson et al. 2009), but had not been quantified across taxa before. However, our results do not support the perception that nationally-at-risk taxa are more peripheral than globally-at-risk taxa in Canada. While globally-at-risk species did have slightly more of their range in Canada on average, they also clustered toward the southern border and 54% had less than 20% of their range in Canada (Figure 2). These results underscore that conservation of at-risk species and biodiversity in Canada, as in many high latitude countries, is fundamentally linked to the conservation of range-edge populations.

Our results also challenge the perception that protecting nationally-at-risk species inherently involves a trade-off with protecting globally-at-risk species. As so many of Canada's nationally-at-risk taxa clusters in the south due to populations at the edge of their range (Kraus et al., 2020), there has been significant debate about how to prioritize their conservation (Fraser 2000), and sustained criticisms that doing so comes at the cost of protecting globally-imperilled taxa (Jones & Fredricksen, 1999; Raymond et al., 2018). Our results suggest that at least in terms of habitat protection, this is a false dichotomy. More than half (56%) of the areas with the highest concentration of nationally-at-risk taxa in Canada were also hotspots of globally-at-risk species in Canada (Figure 1). Hotspots of nationally-at-risk taxa encompassed 64% of the globally at-risk taxa in our data, showing that protecting one threat group can significantly benefit the other.

Not only do hotspots of nationally-at-risk taxa contain many globally-at-risk taxa, they are also

taxonomically diverse. A single nationally-at-risk hotspot contained up to 132 at-risk species

(Figure 1), and all nationally-at-risk hotspots contained at-risk species from at least 6 taxonomic

groups. However, nationally-at-risk hotspots did not represent at-risk mammals as well as they

did other taxa. This is important as non-human mammals often garner disproportionate

conservation interest from human mammals (Davies et al., 2019; Donaldson et al., 2017; Titley

et al., 2017). Hotspots of nationally-at-risk mammals do contain disproportionately high overall

mammal diversity (Cameron & Hargreaves, 2020), so conservation efforts in these areas are

certainly worthwhile. However, our results caution that using at-risk mammals as charismatic

flagship species will not necessarily focus conservation efforts toward the habitat most needed

by other at-risk taxa. Protecting habitat in nationally-at-risk hotspots, on the other hand, can have

a significant conservation benefit for at-risk mammals. Hotspots of nationally-at-risk species

together contained >50% of the at-risk taxa in each taxonomic group, which is highly promising

for planning diverse protected areas.

Strong overlap in hotspots of different taxonomic groups and threat levels should facilitate

habitat protection to promote species-at-risk recovery, but we see little evidence that protected

areas have so far been planned with species-at-risk in mind. The amount of land protected did

increase with the number of at-risk species in non-hotspots cells (Figure 3), which is an

improvement over a complete lack of relationship between protected areas and density of species

at-risk 15 years ago (Deguise & Kerr 2006). However, this effect was slight, and most nationally

and globally at-risk species have only a tiny fraction of their Canadian range protected (Figure

3). Areas that are hotspots for both nationally-at-risk and globally-at-risk taxa should be of the

highest conservation priority, but only 3.8% of these areas are publicly protected. This shows a

sharp disconnect between species at-risk policy and protected area designations.

Why is Canada not doing better at protecting species-at-risk within its borders? The disconnect

could reflect delays and biases in identifying critical habitat or implementing recovery plans

(Bird & Hodges, 2017; Findlay et al., 2009; Mooers et al., 2010), or lack of co-ordination

between the disparate government departments responsible for species-at-risk recovery planning

and protected area designations. It strongly suggests that protected area selection has so far been

driven by objectives other than species-at-risk protection. While protected area objectives can

include historical, scenic, or cultural significance (Benidickson, 2013), designation may also be

influenced by minimizing resource exploitation conflicts, rather than optimizing species-at-risk

conservation (Bolliger et al., 2020; Venter et al., 2017) as is the case in many protected area

networks worldwide (Clancy et al., 2020; Venter et al., 2014). The difficulty of protecting habitat

for species-at-risk species underscores the importance of identifying priority areas that can

promote as many conservation objectives as possible. Our findings of high spatial overlap among

at-risk taxa is great news: it means Canada's recent commitment to protect more natural area

could easily be used to promote species-at-risk recovery. Further, protecting natural landscapes

in densely populated areas like southern Canada can also benefit the many people living there

(Jiricka-Pürrer et al., 2019; White et al., 2019), who are increasingly seeking natural areas and

greenspace (Bain, 2021; Parks Canada, 2019).

Of course, habitat protection generally happens at a smaller scale that the 100 x 100 km grid cells

used in our hotspot analysis. This would be especially true for new protected areas developed in

southern Canada where private land ownership is high. Since species rarely occupy an entire

hotspot grid cell or even all the area within their distribution polygons (Rotenberry &

Balasubramaniam, 2020), protecting a smaller subsection of habitat might not encompass a

hotspot's full suite of overlapping taxa. Even so, coarsely mapping Canada's nationally and

globally at-risk hotspots provides a useful starting point for future, more local-scale analyses to

pinpoint priority areas. Further, patchy species distributions mean that small parks and protected

areas can have a disproportionate benefit for individual species-at-risk.

Another reason to protect habitat in hotspots of species at risk, even if not all species in the

region occur in the local protected area, is that species range shifts driven by climate change are

already changing taxonomic overlap (Yong et al., 2016). Protecting land within the

taxonomically diverse, at-risk hotspots we identified could offer habitat refuges to surrounding

species whose ranges do not yet overlap that area. Northward range shifts are especially

important given our finding that 71% and 54% of the nationally and globally at-risk taxa in

Canada occur at their northern range edge. Protecting habitat in at-risk hotspots would not only

protect a taxonomically diverse set of at-risk populations, but could also provide essential

corridors for edge populations to shift toward higher latitudes and track a changing climate

(Chen et al., 2011; Gibson et al., 2009). Protecting habitat within Canada's at-risk hotspots is

therefore crucial for Canada's current and future biodiversity.

5. ACKNOWLEDGMENTS

We thank Pascale Caissy for spearheading and developing methods for digitizing COSEWIC

range maps, and the Liber Ero Chair in Biodiversity Conservation at McGill University for

financial support in developing this database. We also thank Pascale and Victor Cameron for

helping MEH with further map digitization and quantifying spatial distribution hotspots,

respectively, and the McGill Geographic Information Centre for help with geographic data.

Funding for this project was provided by the Natural Sciences and Engineering Research Council

of Canada (NSERC) through an Undergraduate Science Research Award to MEH and Discovery

Grant to ALH.

6. REFERENCES

Allan, J. R., Watson, J. E. M., Di Marco, M., O'Bryan, C. J., Possingham, H. P., Atkinson, S. C., & Venter, O. (2019). Hotspots of human impact on threatened terrestrial vertebrates. *PLoS Biology*, *17*(3), e3000158. https://doi.org/10.1371/journal.pbio.3000158

Baillie, J. E. M., Hilton-Taylor, C., & Stuart, S. N. (2004). 2004 IUCN Red List of Threatened Species. A global species assessment. IUCN, Gland, Switzerland and Cambridge, UK. xxiv + 191 pp

Bain, J. (2021). Visitor stats reveal how Canada's parks and historic sites fared in 2020. *National Parks Traveler.* Available from https://www.nationalparkstraveler.org/2021/03/visitor-stats-reveal-how-canadas-parks-and-historic-sites-fared-2020

Benidickson, J. (2013). Legal framework for protected areas: Canada. *IUCN-EPLP*, *81*, 1–35. https://doi.org/10.2139/ssrn.2296254

Bird, S. C., & Hodges, K. E. (2017). Environmental science & policy critical habitat designation for Canadian listed species : slow, biased, and incomplete. *Environmental Science and Policy*, *71*, 1–8. https://doi.org/10.1016/j.envsci.2017.01.007

BirdLife International and Handbook of the Birds of the World. (2019). Bird species distribution maps of the world. Version 2019.1. Available from http://datazone.birdlife.org/species/requestdis

Bolliger, C. S., Raymond, C. V., Schuster, R., & Bennett, J. R. (2020). Spatial coverage of protection for terrestrial species under the Canadian Species at Risk Act. *Ecoscience*, *27*(2), 141–147. https://doi.org/10.1080/11956860.2020.1741497

Breheny, P., & Burchett, W. (2017). Visualization of regression models using visreg. *The R Journal*, *9*(2), 56–71.

Brooks, M. E., Kristensen, K., Benthem, K. J. Van, Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R*

Journal, *9*(2), 378–400.

Bunnell, F. L., Campbell, R. W., & Squires, K. A. (2004). Conservation priorities for peripheral

species : the example of British Columbia. *Canadian Journal of Forestry Research*, *34*,

2240–2247. https://doi.org/10.1139/X04-102

Caissy, P., Klemet-N'Guessan, S., Jackiw, R., Eckert, C. G., & Hargreaves, A. L. (2020). High

conservation priority of range-edge plant populations not matched by habitat protection or

research effort. *Biological Conservation*, *249*, 108732.

https://doi.org/10.1016/j.biocon.2020.108732

Cameron, V., & Hargreaves, A. (2020). Spatial distribution and hotspots of mammals in Canada.

Facets, *5*, 1–12. https://doi.org/10.1101/2020.01.29.925461

Ceballos, G., & Ehrlich, P. R. (2006). Global mammal distributions, biodiversity hotspots, and

conservation. *Proceedings of the National Academy of Sciences of the United States of*

America, *103*(51), 19374–19379. https://doi.org/10.1073/pnas.0609334103

Chen, I. C., Hill, J. K., Ohlemüller, R., Roy, D. B., & Thomas, C. D. (2011). Rapid range shifts

of species associated with high levels of climate warming. *Science*, *333*(6045), 1024–1026.

https://doi.org/10.1126/science.1206432

Clancy, N. G., Draper, J. P., Wolf, J. M., Abdulwahab, U. A., Pendleton, M. C., Brothers, S.,

Brahney, J., Weathered, J., & Hammill, E. (2020). Protecting endangered species in the

USA requires both public and private land conservation. *Scientific Reports*, 1–8.

https://doi.org/10.1038/s41598-020-68780-y

Coristine, L. E., Jacob, A. L., Schuster, R., Otto, S. P., Baron, N. E., Bennett, N. J., Bittick, S. J.,

Dey, C., Favaro, B., Ford, A., Nowlan, L., Orihel, D., Palen, W. J., Polfus, J. L., Shiffman,

D. S., Venter, O., & Woodley, S. (2018). Informing Canada's commitment to biodiversity

conservation: a science-based framework to help guide protected areas designation through

Target 1 and beyond. *Facets*, *3*(1), 531–562. https://doi.org/10.1139/facets-2017-0102

Coristine, L. E., & Kerr, J. T. (2011). Habitat loss, climate change, and emerging conservation

challenges in Canada 1. *Canadian Journal of Zoology*, *89*(5), 435–451.

https://doi.org/10.1139/Z11-023

COSEWIC. (2019). COSEWIC Assessment Process, Categories, and Guidelines.

Davies, T., Cowley, A., Bennie, J., Leyshon, C., Inger, R., Carter, H., Robinson, B., Duffy, J. P.,

Casalegno, S., Lambert, G., & Gaston, K. (2019). Correction: Popular interest in vertebrates

does not reflect extinction risk and is associated with bias in conservation investment (PLoS

ONE (2018) 13: 9 (e0203694) DOI: 10.1371/journal.pone.0203694). *PLoS ONE*, *14*(2), 1–

13. https://doi.org/10.1371/journal.pone.0212101

Deguise, I. E., & Kerr, J. T. (2006). Protected Areas and Prospects for Endangered Species

Conservation in Canada. *Conservation Biology, 20*(1), 48-55.

https://doi.org/10.1111/j.1523-1739.2005.00274.x

Donaldson, M. R., Burnett, N. J., Braun, D. C., Suski, C. D., Hinch, S. G., Cooke, S. J., & Kerr,

J. T. (2017). Taxonomic bias and international biodiversity conservation research. *Facets*,

1(1), 105–113. https://doi.org/10.1139/facets-2016-0011

Douma, J. C., & Weedon, J. T. (2019). Analysing continuous proportions in ecology and

evolution: a practical introduction to beta and Dirichlet regression. *Methods in Ecology and*

Evolution, *10*(9), 1412–1430. https://doi.org/10.1111/2041-210X.13234

Environment and Climate Change Canada. (2021a). Canada Target 1 challenge. Available from

https://www.canada.ca/en/environment-climate-change/services/nature-legacy/canada-

target-one-challenge.html

Environment and Climate Change Canada. (2021b). Canadian protected and conserved areas

database. Available from https://www.canada.ca/en/environment-climate-

change/services/national-wildlife-areas/protected-conserved-areas-database.html

Fattorini, S., Dennis, R. L. H., & Cook, L. M. (2012). Use of Cross-Taxon congruence for

hotspot identification at a regional scale. *PLoS ONE*, *7*(6), 1–6.

https://doi.org/10.1371/journal.pone.0040018

Findlay, C. S., Elgie, S., Giles, B., & Burr, L. (2009). Species listing under Canada's Species at

Risk Act. *Conservation Biology*, *23*(6), 1609–1617. https://doi.org/10.1111/j.1523-

1739.2009.01255.x

Gibson, S. Y., Van Der Marel, R. C., & Starzomski, B. M. (2009). Climate change and

conservation of leading-edge peripheral populations. *Conservation Biology*, *23*(6), 1369–

1373. https://doi.org/10.1111/j.1523-1739.2009.01375.x

Government of Canada. (2021). Species at risk public registry. Available from

https://species-registry.canada.ca/index-en.html

Hunter, M. L., & Hutchinson, A. (1994). The virtues and shortcomings of parochialism:

conserving species that are locally rare, but globally common. *Conservation Biology*, *8*(4),

1163–1165. https://doi.org/10.1046/j.1523-1739.1994.08041163.x

IUCN. (2012). IUCN Red List categories and criteria: Version 3.1. Second edition. Gland,

Switzerland and Cambridge, UK: IUCN. iv + 32pp.

IUCN. (2015). Conservation successes overshadowed by more species declines – IUCN Red List

update | IUCN. Available from https://www.iucn.org/content/conservation-successes-

overshadowed-more-species-declines-–-iucn-red-list-update

IUCN. (2021). The IUCN Red List of Threatened Species. Version 2021-1. Available from

https://www.iucnredlist.org. Downloaded on 2021-05-08.

Jiricka-Pürrer, A., Tadini, V., Salak, B., Taczanowska, K., Tucki, A., & Senes, G. (2019). Do

protected areas contribute to health and well-being? A cross-cultural comparison.

International Journal of Environmental Research and Public Health, *16*(7).

https://doi.org/10.3390/ijerph16071172

Jones, L., & Fredricksen, L. (1999). Crying wolf? Public policy on endangered species in

Canada. *Critical Issues Bulletin, The Fraser Institute*, 1–23.

Kearney, S. G., Adams, V. M., Fuller, R. A., Possingham, H. P., & Watson, J. E. M. (2020).

Estimating the benefit of well-managed protected areas for threatened species conservation.

Oryx, *54*(2), 276–284. https://doi.org/10.1017/S0030605317001739

Kennedy, M., & Kopp, S. (2000). Understanding Map Projections: GIS by ESRI. *ESRI Press.*

https://gis.icao.int/icaoetod/map_projections[1].pdf

Kerr, J. T., & Deguise, I. (2004). Habitat loss and the limits to endangered species recovery.

Ecology Letters, *7*(12), 1163–1169. https://doi.org/10.1111/j.1461-0248.2004.00676.x

Komonen, A. (2007). Are we conserving peripheral populations? An analysis of range structure
of longhorn beetles (Coleoptera: Cerambycidae) in Finland. *Journal of Insect Conservation*,
11(3), 281–285. https://doi.org/10.1007/s10841-006-9043-8

Kraus, D., & Hebb, A. (2020). Southern Canada's crisis ecoregions: identifying the most
significant and threatened places for biodiversity conservation. *Biodiversity and*
Conservation, *29*(13), 3573–3590. https://doi.org/10.1007/s10531-020-02038-x

Kraus, D., Murphy, S., & Armitage, D. (2021). Ten bridges on the road to recovering Canada's
endangered species. *Facets*, *6*, 1088–1127. https://doi.org/10.1139/facets-2020-0084

Lenth, R. V. (2016). Least-squares means: The R package lsmeans. *Journal of Statistical*
Software, *69*(1). https://doi.org/10.18637/jss.v069.i01

Littlefield, C. E., Krosby, M., Michalak, J. L., & Lawler, J. J. (2019). Connectivity for species on
the move: supporting climate-driven range shifts. *Frontiers in Ecology and the*
Environment, *17*(5), 270–278. https://doi.org/10.1002/fee.2043

Maxwell, S. L., Cazalis, V., Dudley, N., Hoffmann, M., Rodrigues, A. S. L., Stolton, S.,
Visconti, P., Woodley, S., Kingston, N., Lewis, E., Maron, M., Strassburg, B. B. N.,
Wenger, A., Jonas, H. D., Venter, O., & Watson, J. E. M. (2020). Area-based conservation
in the twenty-first century. *Nature*, *586*(7828), 217–227. https://doi.org/10.1038/s41586-
020-2773-z

Mooers, A. O., Doak, D. F., Scott Findlay, C., Green, D. M., Grouios, C., Manne, L. L.,
Rashvand, A., Rudd, M. A., & Whitton, J. (2010). Science, policy, and species at risk in
Canada. *BioScience*, *60*(10), 843–849. https://doi.org/10.1525/bio.2010.60.10.11

Natural Earth. (2020). Natural earth vector countries. Version 4.1.0 [online]: Available
from https://naturalearthdata.com

Orme, C. D. L., Davies, R. G., Burgess, M., Eigenbrod, F., Pickup, N., Olson, V. A., Webster, A.
J., Ding, T. S., Rasmussen, P. C., Ridgely, R. S., Stattersfield, A. J., Bennett, P. M.,
Blackburn, T. M., Gaston, K. J., & Owens, I. P. F. (2005). Global hotspots of species
richness are not congruent with endemism or threat. *Nature*, *436*(7053), 1016–1019.
https://doi.org/10.1038/nature03850

Parks Canada. (2019). *Parks Canada attendance 2019-20*. Parks Canada attendance.
https://publications.gc.ca/site/eng/9.839688/publication.html

Prendergast, J. R., Quinn, R. M., Lawton, J. H., Eversham, B. C., & Gibbons, D. W. (1993). Rare
species, the coincidence of diversity hotspots and conservation strategies. *Nature*,
365(6444), 335–337. https://doi.org/10.1038/365335a0

QGIS Development Team. (2021). QGIS Geographic Information System. Available from
http://qgis.osgeo.org.

R Core Team. (2021). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. Available from https://www.R-project.org/.

Raymond, C. V., Wen, L., Cooke, S. J., & Bennett, J. R. (2018). National attention to
endangered wildlife is not affected by global endangerment: a case study of Canada's
species at risk program. *Environmental Science and Policy*, *84*(August 2017), 74–79.
https://doi.org/10.1016/j.envsci.2018.03.001

Reid, W. V. (1998). Biodiversity hotspots. *Trends in Ecology and Evolution*, *13*(7), 275–280.
https://doi.org/10.1016/S0169-5347(98)01363-9

Rodrigues, A. S. L., & Gaston, K. J. (2002). Rarity and conservation planning across geopolitical
units. *Conservation Biology*, *16*(3), 674–682. https://doi.org/10.1046/j.1523-
1739.2002.00455.x

Rotenberry, J. T., & Balasubramaniam, P. (2020). Connecting species' geographical distributions
to environmental variables: range maps versus observed points of occurrence. *Ecography*,
43(6), 897–913. https://doi.org/10.1111/ecog.04871

Thomas, C. D., Gillingham, P. K., Bradbury, R. B., Roy, D. B., Anderson, B. J., Baxter, J. M.,
Bourne, N. A. D., Crick, H. Q. P., Findon, R. A., Fox, R., Hodgson, J. A., Holt, A. R.,
Morecroft, M. D., O'Hanlon, N. J., Oliver, T. H., Pearce-Higgins, J. W., Procter, D. A.,
Thomas, J. A., Walker, K. J., … Hill, J. K. (2012). Protected areas facilitate species' range
expansions. *Proceedings of the National Academy of Sciences of the United States of
America*, *109*(35), 14063–14068. https://doi.org/10.1073/pnas.1210251109

Titley, M. A., Snaddon, J. L., & Turner, E. C. (2017). Scientific research on animal biodiversity

is systematically biased towards vertebrates and temperate regions. *PLoS ONE*, *12*(12), 1–

14. https://doi.org/10.1371/journal.pone.0189577

Vellend, M. (2003). Habitat loss inhibits recovery of plant diversity as forests regrow. *Ecology*,

84(5), 1158–1164. https://doi.org/10.1890/0012-9658(2003)084[1158:HLIROP]2.0.CO;2

Venter, O., Fuller, R. A., Segan, D. B., Carwardine, J., Brooks, T., Butchart, S. H. M., Di Marco,

M., Iwamura, T., Joseph, L., O'Grady, D., Possingham, H. P., Rondinini, C., Smith, R. J.,

Venter, M., & Watson, J. E. M. (2014). Targeting global protected area expansion for

imperiled biodiversity. *PLoS Biology*, *12*(6). https://doi.org/10.1371/journal.pbio.1001891

Venter, O., Magrach, A., Outram, N., Klein, C. J., Possingham, H. P., Marco, M. Di, & Watson,

J. E. M. (2017). Bias in protected-area location and its effects on long-term aspirations of

biodiversity conventions. *Conservation Biology*, *32*(1), 127–134.

https://doi.org/10.1111/cobi.12970

Watson, J. E. M., Dudley, N., Segan, D. B., & Hockings, M. (2014). The performance and

potential of protected areas. *Nature*, *515*(7525), 67–73. https://doi.org/10.1038/nature13947

Wells, J. V., Robertson, B., Rosenberg, K. V., & Mehlman, D. W. (2010). Global versus local

conservation focus of U.S. state agency endangered bird species lists. *PLoS ONE*, *5*(1), 3–7.

https://doi.org/10.1371/journal.pone.0008608

White, M. P., Alcock, I., Grellier, J., Wheeler, B. W., Hartig, T., Warber, S. L., Bone, A.,

Depledge, M. H., & Fleming, L. E. (2019). Spending at least 120 minutes a week in nature

is associated with good health and wellbeing. *Scientific Reports*, *9*(1), 1–11.

https://doi.org/10.1038/s41598-019-44097-3

Woo-Durand, C., Matte, J. M., Cuddihy, G., McGourdji, C. L., Venter, O., & Grant, J. W. A.

(2020). Increasing importance of climate change and other threats to at-risk species in

Canada. *Environmental Reviews*, *28*(4), 449–456. https://doi.org/10.1139/er-2020-0032

Yong, D. L., Barton, P. S., Okada, S., Crane, M., & Lindenmayer, D. B. (2016). Birds as

surrogates for mammals and reptiles: are patterns of cross-taxonomic associations stable

over time in a human-modified landscape? *Ecological Indicators*, *69*, 152–164.

https://doi.org/10.1016/j.ecolind.2016.04.013