Synapse-associated astrocyte mitochondria stabilize motor circuits to prevent

excitotoxicity

Sonja A. Zolnoski[1], Emily L. Heckman[1], Chris Q. Doe[1], and Sarah D. Ackerman[1,2]*

[1]Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon, Eugene,

OR 97403

[2]Current address: Center for Brain Immunology and Glia, Department of Pathology and

Immunology, Washington University School of Medicine, Saint Louis, MO 63110

* Author for correspondence at sarah.ackerman@wustl.edu (SDA)

Orcid IDs: 0000-0002-0012-3364 (ELH), 0000-0001-5980-8029 (CQD), 0000-0001-8752-8972
(SDA)

ABSTRACT

Early stages of the devastating neurodegenerative disease amyotrophic lateral sclerosis (ALS) are characterized by motor neuron hyperexcitability. During this phase, peri-synaptic astrocytes are neuroprotective. When reactive, loss of wild-type astrocyte functions results in excitotoxicity. How astrocytes stabilize motor circuit function in early-stage ALS is poorly understood. Here, we used *Drosophila* motor neurons to define the role of astrocyte-motor neuron metabolic coupling in a model of ALS: astrocyte knockdown of the ALS-causing gene *tbph/TARDBP*. In wild-type, astrocyte mitochondria were dynamically trafficked towards active motor dendrites/synapses to meet local metabolic demand. Knockdown of *tbph* in astrocytes resulted in motor neuron hyperexcitability, reminiscent of early-stage ALS, which was met with a compensatory accumulation of astrocyte mitochondria near motor dendrites/synapses. Finally, we blocked mitochondria-synapse association in *tbph* knockdown animals and observed locomotor deficits and synapse loss. Thus, synapse-associated astrocyte mitochondria stabilize motor circuits to prevent the transition from hyperexcitability to excitotoxicity.

MAIN

Amyotrophic lateral sclerosis (ALS) is devastating neurodegenerative disorder in which

motor neurons are selectively lost from the central nervous system (CNS), resulting in

paralysis and eventual death. One major hurdle to understanding the molecular and cellular

basis of ALS is the genetic complexity of the disease, which can be caused or modified by

hundreds of mutations in greater than 50 genes[1–4]. Furthermore, less than ten percent of cases

with a known genetic basis are hereditary (familial ALS); thus, each mutation only represents

a very small percentage of the overall population suffering from ALS[1]. For example,

mutations in TAR DNA-binding protein 43 (TDP-43, encoded by the gene *TARDBP*)

represent 1.5-4.2% of familial ALS, but only .2-.8 % of all ALS cases[5]. Despite the low

representation of *TARDBP* mutations in the overall ALS population, changes in TDP-43

localization are apparent in ninety-five percent of ALS and forty-five percent of overlapping

frontotemporal dementia (FTD) cases[6]. Thus, TDP-43 represents a promising therapeutic

target to tackle the pathophysiological heterogeneity of ALS/FTD.

TDP-43 is a nuclear DNA/RNA binding protein that broadly regulates RNA

metabolism and subsequent gene expression. In ALS/FTD neurons and associated glia, TDP-

43 is translocated from the nucleus to form dense cytoplasmic inclusions. Accordingly, it is

thought that TDP-43-associated neuronal death results from both loss-of-function and toxic

gain-of-function phenotypes[7,8]. Although ALS is considered a motor neuron disease, recent

reports suggest that glial dysfunction can strongly modulate disease progression[9–12].

Astrocytes −the most abundant glial subtype− extend finely branched processes to intimately

ensheath central synapses, including those at motor neuron dendrites. This unique positioning

allows astrocytes to play several key roles that ensure neuronal health: provision of metabolic

support, ion buffering, neurotransmitter recycling, and regulation of synaptic

structure/function[13]. In ALS, healthy astrocytes undergo dramatic changes in cell morphology

and gene expression profile to transform from a functional homeostatic to dysfunctional

reactive state[1,14,15]. The early phases of ALS are characterized by motor neuron

hyperexcitability[16], and transformation of astrocytes into a reactive state is thought to tip the

balance from motor neuron hyperexcitability to excitotoxicity[16,17]. In contrast, the presence of

wild-type astrocytes in an otherwise mutant background is sufficient to slow disease onset and

progression in murine models of ALS[18]. Thus, discovering which wild-type function(s) of

astrocytes are most neuroprotective in pre-symptomatic ALS is of great clinical importance.

We took advantage of *Drosophila* as a system with well-established ALS models and

precise genetic access to motor neurons and astrocytes[19,20]. Specifically, we wanted to test the

hypothesis that reduction in astrocyte-motor neuron metabolic coupling transforms motor

circuit activity from hyperactive to excitotoxic in an ALS model. In the healthy nervous

system, we found that astrocytes position mitochondria near motor dendrites/synapses in an

activity-dependent manner, presumably to match metabolic demand. Following astrocyte-

specific knockdown of *tbph* (human *TARDBP*), we observed increased motor neuron activity

and increased localization of astrocyte mitochondria near motor dendrites/synapses, but no

excitotoxicity or behavioral changes. These data suggest that astrocyte mitochondria are tuned

to neighboring neuronal activity, and that proximity to synapses may be neuroprotective in

conditions where motor activity is elevated (e.g. astrocyte knockdown of *tbph*). To test this

hypothesis, we blocked mitochondrial trafficking in *tbph* knockdown animals and assessed

the effect on motor circuit structure and function. When mitochondria cannot form stable

associations with synapses in *tbph* knockdown animals, we observed significant increases in

crawling speed, consistent with motor neuron hyperexcitability, which preceded synapse loss

at the neuromuscular junction, consistent with excitotoxicity. We conclude that astrocytes

dynamically position their mitochondria to match local metabolic demand and facilitate long-

term stability of motor circuits.

RESULTS

Activity-dependent positioning of astrocyte mitochondria

Astrocytes are an essential energy source for neurons, facilitating both neuronal longevity and

long-term memory formation[13,21,22]. Not surprisingly, mitochondria are frequently visualized

inside fine astrocytic peri-synaptic processes by transmission electron microscopy[23]. We

sought to test whether ALS-causing mutations resulted in defective astrocyte-neuron

metabolic coupling and excitotoxicity. As diseased motor neurons exhibit elevated activity in

mammals[1], we first needed test if astrocyte mitochondria localization correlates with motor

neuron activity. We focused our work on the well-characterized aCC/RP2 motor neurons of

the *Drosophila* larval motor circuit[24]. We developed an image analysis pipeline in Imaris to

define presynapses that were co-localized with aCC/RP2 dendrites, similar to one previously

published[20,25]. We then determined the percentage of these synapses that were associated with

astrocyte mitochondria (Figure 1, see legend and methods for full details). Using these model

neurons, we recently discovered a critical period of motor circuit plasticity during

development when motor neurons robustly change their dendrite structure in response to

neuronal activity[20]. To assess activity-dependent mitochondria-synapse association without

the confounds of structural plasticity, we limited our studies to larval stages ≥ 8 hours after

larval hatching (h ALH), when the critical period is closed and dendrite morphology is not

dramatically affected by motor neuron activity. Following optogenetic activation of motor

neurons for one hour (*RN2-lexA, lexAop-Chrimson::tdTomato*), we observed a redistribution

of astrocyte mitochondria (*R25H07-gal4, UAS-mito::GFP*) from the periphery to the center of

the dendritic domain (Figure 2a-b"). This resulted in a significant increase in total astrocyte

mitochondrial volume within the dendritic domain, and a significant increase in the

percentage of motor synapses within close proximity (≤ 120 nm) to astrocyte mitochondria

(Figure 2c-d). Importantly, we did not observe any activity-dependent changes in overall

dendrite volume that could account for these phenotypes, as the critical period was closed

(Figure 2e).

Previous mammalian slice culture assays demonstrated that acute, pharmacological

activation of neurons results in pausing of astrocyte mitochondria near synapses in a calcium-

dependent manner[26,27]. To determine if *Drosophila* motor neuron activity causes astrocyte

mitochondrial stalling at motor dendrites/synapses, we used live imaging to visualize real-

time changes in mitochondrial behavior during optogenetic activation of neighboring motor

neurons. In contrast to simple stalling of mitochondria at synapses, we observed rapid, active

transport of astrocyte mitochondria from the periphery to the center of the dendritic domain

within one minute of motor neuron activation, resulting in a significant increase in

mitochondrial volume within the activated region. Interestingly, this volume plateaued after

five minutes of stimulation, suggesting that *in vivo*, mitochondria are initially actively

transported to dendrites/synapses, followed by pausing once a certain threshold of activity has

passed (Figure 2f-g, Supplemental Movies 1-2).

We next tested whether optogenetic silencing of motor neurons (*RN2-gal4, UAS-GtACR2::EYFP*) alters astrocyte mitochondrial distribution (*alrm-lexA, lexAop-*

mcherry::mito.OMM, Figure 3). Interestingly, while motor neuron activation was sufficient to recruit astrocyte mitochondria to synapses, synapse-associated mitochondria were largely stable throughout motor neuron silencing (Figure 3a-c). We did, however, observe a significant decrease in overall mitochondrial volume within the dendritic domain after silencing for one hour (Figure 3d), but not shorter times (Figure 3f-g, Supplemental Movies 3-4). Dendrite volume was stable throughout these manipulations, as the critical period was closed (Figure 3e). In sum, astrocyte mitochondria traffic to areas of elevated neuronal activity and away from areas of low neuronal activity to meet local metabolic demand.

Astrocyte-knockdown of *tbph/TARDBP* enhances mitochondria-synapse association

The early stages of ALS are characterized by motor neuron hyperexcitability, but not excitotoxicity[16]. During this phase, astrocytes maintain their homeostatic functions and are thought to be neuroprotective[14]. To test whether loss of activity-dependent astrocyte mitochondrial trafficking contributes to motor neuron loss in the *tbph* ALS model, we visualized astrocyte mitochondria (*R25H07-gal4, UAS-mito::GFP*) relative to motor neuron dendrites (*RN2-lexA, lexAop-Chrimson::tdTomato*) and synapses (Brp+) following astrocyte-specific knockdown of *tbph*. To achieve efficient knockdown, we used a well-established and previously published RNA interference line (*UAS-tbph*[RNAi])[28–30]. Astrocyte-specific knockdown of *tbph* resulted in a significant increase in mitochondria-synapse association and increased volume of astrocyte mitochondrial volume within the motor neuropil; optogenetic activation further enhanced these phenotypes (Figure 4a-d", quantified in e,f). Importantly, neither genotype nor manipulation impacted overall dendrite volume (Figure 4g). Together,

these data demonstrate that loss of astrocyte Tbph does not impair mitochondrial recruitment, but instead enhances mitochondrial trafficking to the dendritic domain.

Astrocyte-knockdown of *tbph*/*TDP-43* causes ectopic motor neuron activity

Thus far, we have shown that motor neuron activation can recruit astrocyte mitochondria to motor synapses (Figure 2), and this phenotype is enhanced following astrocyte knockdown of *tbph* (Figure 4). Motor neuron hyperexcitability is common in ALS models, including mouse and human-patient derived *TARDBP* mutant motor neurons[31,32]. Given that astrocyte knockdown of *tbph* resulted in accumulation of astrocyte mitochondria near motor synapses even without optogenetic stimulation of motor neurons (Figure 4e), we tested whether *tbph* knockdown in astrocytes alone is sufficient to cause motor neuron hyperexcitability. To test this hypothesis, we assessed calcium signaling within motor neuron dendrites (*RN2-lexA, lexAop-GCaMP7s*; Figure 5a) in astrocyte *tbph* knockdown animals (*R25H07-gal4, UAS-tbph*[RNAi]). During a 15-minute imaging period, control motor dendrites exhibited a baseline level of spontaneous calcium activity (Figure 5b-d, Supplemental Movie 5). In contrast, knock down of *tbph* in astrocytes led to a significant increase in the frequency of spontaneous calcium events, as well as an increase in the average amplitude of each event (Figure 5b-e, Supplemental Movie 6).

These data indicate that astrocyte knockdown of *tbph* is sufficient to cause motor neuron hyperexcitability. In murine ALS models, hyperexcitability is caused by a reduction in the astrocyte excitatory amino acid transporter EAAT2[33,34], which clears 90% of peri-synaptic glutamate in the adult brain[35]. Note that synapse-associated *Drosophila* astrocytes primarily express EAAT1, a related glutamate receptor that is the major EAAT subtype expressed

during mammalian CNS development[35] and is the only high-affinity glutamate transporter in

Drosophila[36]. Previous studies in *Drosophila* found that *eaat1* expression levels are

dysregulated in *tbph* mutants[37] and glia-specific overexpression of *eaat1* is sufficient to rescue

locomotor defects in *tbph* mutant larvae[38]. Similarly, we found a significant reduction in

EAAT1 levels on astrocytes following *tbph* knockdown (Figure 5f-h). These data support the

model that loss of *tbph* from astrocytes results in a reduction in EAAT1, accumulation of

excitatory neurotransmitters within the neuropil, and ultimately hyperexcitability.

Identification of cellular machinery for astrocyte mitochondrial trafficking

If astrocyte mitochondria are trafficked to motor synapses to meet energetic demand, perhaps

enhanced astrocyte mitochondrial trafficking in *tbph* knockdown conditions prevents

hyperexcitability (Figure 5) from turning into excitotoxicity. To test this hypothesis, we first

needed to determine how mitochondria are trafficked within astrocytes, which is vastly

understudied relative to neurons. We performed astrocyte-specific RNAi knockdown of

molecular motor and mitochondrial adaptor proteins using previously published RNAi lines[30],

and visualized astrocyte mitochondria (*R25H07-gal4, UAS-mito::GFP, UAS-RNAi*) by *in vivo*

time-lapse imaging and in fixed preparations (Supplemental Movies 7-11, Figure 6). We

screened eight lines targeting the adaptor protein *milton* (N=3 lines), the mitochondrial

membrane GTPase *miro* (N=1 line, previously linked to astrocyte mitochondrial

trafficking[39]), as well as the molecular motors *kinesin heavy chain* (N=2 lines) and *dynein*

heavy chain (N=2 lines). We found that astrocyte-specific knockdown of *milton* completely

blocked mitochondrial trafficking into the neuropil, without any observable defects in the

peri-synaptic localization of astrocyte processes (Supplemental Movie 8, Figure 6a-b",

quantified in f-g). We also observed significant reductions in mitochondrial trafficking

following *miro* or *khc* knockdown (Supplemental Movies 9-10, Figure 6c-d", quantified in f-

g). In neurons, Milton is a mitochondrial adaptor protein that serves as a linker between

mitochondria-bound Miro and microtubule-associated Kinesin heavy chain[40,41]. Together,

this complex facilitates long-distance, (+) end directed axonal trafficking of mitochondria

along microtubules. Conversely, astrocyte knockdown of the (-) end directed microtubule

motor protein Dynein heavy chain did not have any impact on mitochondrial trafficking into

the neuropil (Supplemental Movie 11, Figure 6e-h). At this developmental stage, astrocyte

membranes are rapidly growing into the neuropil to tile in between neuronal synapses[20,42].

Our data suggest that during this developmental stage of rapid astrocyte ingrowth,

microtubules are oriented (+) end out to facilitate mitochondrial trafficking into distal peri-

synaptic processes to meet the metabolic demands of neurons.

Astrocyte mitochondrial trafficking deficits modify locomotor behavior

Having defined the molecular machinery required for astrocyte mitochondrial trafficking, we

next wanted to determine whether increased mitochondria-synapse proximity following

astrocyte *tbph* knockdown is neuroprotective. To that end, we used a high-throughput

behavioral rig[43] to assay a series of locomotor behaviors (Figure 7a) in freely crawling

tbph[RNAi], *milton*[RNAi], and double knockdown animals at 72 h ALH (*R25H07-gal4, UAS-*

RNAi). We found that single knockdown of *tbph* in astrocytes did not have a significant

impact on locomotor behavior (Figure 7b-c, quantified in f-j). These data suggest that despite

elevated motor neuron activity at 8 h ALH (Figure 5), homeostatic mechanisms stabilize

overall motor circuit activity when astrocyte mitochondria can properly translocate to the

synapse (Figure 4). Astrocyte knockdown of *milton* alone caused significant reductions in larval distance traveled due to a combined decrease in crawling speed and duration. We also observed increased bending in *milton* knockdown animals, a measure of uncoordinated movement (Figure 7d, quantified in f-j). In contrast, when we simultaneously knocked down *tbph* and *milton*, we saw a significant increase in crawling speed, duration, and overall traversed distance. In double knockdowns, these phenotypes were still accompanied by increased bending, which occurs in *milton,* but not *tbph,* single knockdowns. These data suggest that *milton* knockdown is the primary driver of the uncoordinated phenotype (Figure 7e, quantified in f-j). Thus, in *tbph* single knockdown animals where mitochondria-synapse association is preserved, homeostatic mechanisms rebalance overall network activity to prevent developmental hyperexcitability (Figure 5) from manifesting in mature locomotor phenotypes (Figure 7b-c, quantified in f-j). When we blocked mitochondrial trafficking via *milton*[RNAi] in *tbph* knockdown animals, homeostatic mechanisms were not activated. Accordingly, ectopic motor signaling in the CNS (Figure 5) corresponded with hyperactive crawling (Figure 7).

Astrocyte mitochondrial trafficking deficits impair synapse maintenance

Blocking mitochondrial trafficking following *tbph* knockdown resulted in hyperactivity; we wanted to know if this was sufficient to shift hyperexcitability to excitotoxicity. To that end, we genetically labeled all motor neurons (*vglut-lexA, lexAop-myr::GFP*) while knocking down *tbph*, *milton*, or both in astrocytes (*R25H07-gal4, UAS-RNAi*). We then performed larval fillet preps to visualize the structure of the neuromuscular junction at 96 h ALH. We focused our analyses on the dorsal-most muscles where our model motor neurons

(aCC>muscle DA1; RP2>muscle DA2) and others (U1>muscle DO1; U2>muscle DO2)

elaborate their synaptic boutons[44]. Similar to our behavioral analyses, we observed no

significant difference in bouton number between control and *tbph* knockdown animals (Figure

8a-b, e-f), suggesting that homeostatic mechanisms that stabilize the motor circuit (Figure 7b-

c, quantified in f-j) may act at the level of presynaptic vesicle release[45]. Knockdown of *milton*

alone resulted in a significant increase in synaptic bouton number (Figure 8c, quantified in f).

Because knockdown of *milton* resulted in decreased crawling speed and distance (Figure 7d,

quantified in f-j), these data suggest that astrocyte knockdown of *milton* results in reduced

motor neuron signaling and a compensatory increase in bouton number, but does not achieve

homeostasis. Finally, we observed a significant decrease in total bouton number when we

blocked mitochondrial trafficking in *tbph* knockdown animals (*R25H07-gal4, UAS-tbph*[RNAi],

UAS-milton[RNAi], Figure 8d, quantified in f). Thus, mitochondria-synapse association is

necessary for synapse maintenance following astrocyte knockdown of *tbph*. In sum, our data

support the hypothesis that peri-synaptic localization of astrocyte mitochondria stabilizes

motor circuit structure and function in a *Drosophila* model of ALS.

DISCUSSION

Astrocyte neurotransmitter turnover and ALS

In pathological conditions (disease or injury), inflammation drives the transformation of healthy

astrocytes into reactive astrocytes (astrogliosis), which are characterized by morphological,

transcriptional, and functional changes[46]. Reactive astrocytes were first described over a century

ago and are considered a hallmark of neurodegenerative disease[47]. Accordingly, most research on

the role of astrocytes in neurodegeneration has focused on the putative toxic effects of reactive

astrocytes on compromised neuronal populations (*e.g.* motor neurons in ALS). One of the best-

studied ALS models featuring astrogliosis is the G93A variant of superoxide dismutase 1 ($SOD1^{G93A}$). SOD1 is an enzyme that reduces cellular levels of the free radical superoxide[48]. Mutations in SOD1 are associated with oxidative stress, mitochondrial dysfunction, inflammation, and motor neuron excitotoxicity due to misregulation of peri-synaptic neurotransmitter concentrations[49]. In murine models, astrocyte-specific overexpression of $SOD1^{G93A}$ causes astrocyte activation, triggers release of proinflammatory cytokines, and is sufficient to induce motor neuron death *in vitro* and *in vivo*[11,50,51]. Compound mutations (*IL-1α*$^{-/-}$ *TNFα*$^{-/-}$ *C1qa*$^{-/-}$) that prevent astrocyte reactivity can significantly slow disease progression (but not onset) and extend the life of $SOD1^{G93A}$ animals[17]. In contrast, transplantation of wild-type astrocytes into the spinal cord of $SOD1^{G93A}$ rats can delay both disease onset and progression[18]. These data implicate astrocyte-induced inflammation as a key determinant of ALS progression, but also point to early neuroprotective roles of wild-type astrocytes.

Here, we wanted to determine which astrocyte functions support motor circuit activity and maintenance in a *Drosophila* model of ALS. We focused on *tbph/TARDBP* as a common pathological indicator in ALS/FTD[6]. In mouse, depletion of astrocyte *Tardbp* is associated with astrocyte reactivity in mature motor circuits where symptoms have already manifested[31]. Although *Drosophila* astrocytes are not known to become reactive, glia-specific knockdown of *tbph* is linked to reduced adult lifespan and decreased motor coordination[37] and glia can modify the severity of ALS phenotypes in *tbph* null flies[52]. We focused our initial experiments on pre-symptomatic stages to determine what homeostatic function(s) of astrocytes are lost following *tbph* knockdown, and how that in turn impacts motor circuit maintenance. We found that loss of *tbph* in astrocytes alone enhances motor neuron activity, likely due to reductions in EAAT1 expression on astrocyte membranes and accumulation of extracellular glutamate. The *Drosophila*

neuromuscular junction (NMJ) exhibits striking homeostatic plasticity to stabilize motor output in experimental conditions where motor neuron activity is perturbed[45,53]. In accordance with these previous findings, we did not observe overt behavioral defects in astrocyte *tbph* knockdown animals. In addition, we did not observe any significant changes in synaptic bouton number following astrocyte *tbph* knockdown. Similarly, depletion of astrocyte *Tardbp* in mouse does not impact NMJ morphology[31]. These data support the hypothesis that when astrocyte mitochondria-synapse association is intact, homeostatic plasticity mechanisms at the NMJ (*e.g.* presynaptic release) can compensate for ectopic motor neuron activation in the CNS to stabilize motor output and in turn, locomotor behavior.

Astrocyte-neuron metabolic coupling ensures motor circuit homeostasis

A primary function of healthy astrocytes is to meet the metabolic demand of associated neurons[13,21]. We found that astrocyte mitochondria sense and respond to local changes in neuronal activity, as stimulation of motor neurons resulted in a rapid mobilization and then stabilization of astrocyte mitochondria near motor synapses. These findings are consistent with previous characterizations of activity-dependent mitochondrial trafficking in hippocampal slice cultures[27]. We questioned whether knockdown of *tbph* would impair activity-dependent mitochondrial trafficking. Instead, we observed enhanced localization of peri-synaptic astrocyte mitochondria following knockdown of *tbph*, theoretically to meet the energetic demand of hyperactive motor neurons. Indeed, when we blocked mitochondrial trafficking in *tbph* knockdown animals, we observed significant increases in crawling speeds that preceded synapse loss at the NMJ. Thus, in the absence of peri-synaptic astrocyte mitochondria, homeostatic mechanisms that stabilize motor circuit activity failed to engage, resulting in excitotoxicity.

Do astrocytes dampen neuronal activity?

Hyperexcitability is a major cause of motor neuron death in ALS[16]. We found that knockdown of

astrocyte *tbph* causes enhanced motor neuron activity due to decreased astrocyte EAAT1

expression. Similarly, overexpression of SOD1^{G93A} in astrocytes results in motor neuron

hyperexcitability due to decreased expression of EAAT2[54–56]. These data indicate that wild-type

astrocytes generally serve to restrict motor neuron activity to a given physiological range.

Interestingly, we recently identified a critical period of activity-dependent homeostatic plasticity

in the *Drosophila* motor circuit that is closed by astrocyte-motor neuron contact. Within this

window, motor neurons scale dendrite length, complexity, and connectivity in response to their

own activity; failure to close the critical period results in sustained changes to locomotor

behavior[20]. In the present study, we found that in a *Drosophila* model of ALS, blocking astrocyte

mitochondrial transport impaired homeostatic mechanisms that rebalance motor circuit activity

during development, resulting in mature synapse loss. These findings open up the intriguing

possibility that astrocytes buffer developmental activity to facilitate long-term circuit

maintenance and function, and that ultimately, neurodegenerative diseases reflect failed checks on

developmental activity.

Acknowledgements

We thank Marc Freeman for providing fly stocks. We thank members of the Doe lab for comments on the manuscript. Stocks obtained from the Bloomington Drosophila Stock Center were used in this study. Funding was provided by HHMI (CQD), R01 HD27056 (CQD), NIH K99NS121137 (SDA). SDA is a Milton Safenowitz Post-doctoral fellow of the ALSA.

Author Contributions

SDA conceived of the project; SDA performed and analyzed all experiments in Figures 1-4 and 6; SDA and SAZ performed and analyzed experiments in Figures 5-8; ELH analyzed GCaMP7s data in Figure 5; CQD provided feedback during the project; SDA, SAZ, and ELH wrote the paper and prepared the figures. All authors commented and approved of the manuscript.

Competing Interest Statement

The authors declare no competing financial or non-financial interests.

Data Availability Statement

This study did not generate/analyze datasets/code. Raw data for any main or supplemental figure (.lsm, .czi, or .avi files) can be supplied upon request.

REFERENCES

1. *Amyotrophic Lateral Sclerosis*. (Exon Publications, 2021).
    doi:10.36255/exonpublications.amyotrophiclateralsclerosis.2021.

2. Rosen, D. R. *et al.* Mutations in Cu/Zn superoxide dismutase gene are associated with
    familial amyotrophic lateral sclerosis. *Nature* **362**, 59–62 (1993).

3. Rowland, L. P. & Shneider, N. A. Amyotrophic Lateral Sclerosis. *N. Engl. J. Med.* **344**,
    1688–1700 (2001).

4. Lattante, S., Rouleau, G. A. & Kabashi, E. TARDBP and FUS Mutations Associated with
    Amyotrophic Lateral Sclerosis: Summary and Update. *Hum. Mutat.* **34**, 812–826 (2013).

5. Mejzini, R. *et al.* ALS Genetics, Mechanisms, and Therapeutics: Where Are We Now? *Front.*
    Neurosci. **13**, 1310 (2019).

6. Neumann, M. *et al.* Ubiquitinated TDP-43 in frontotemporal lobar degeneration and
    amyotrophic lateral sclerosis. *Science* **314**, 130–133 (2006).

7. Janssens, J. & Van Broeckhoven, C. Pathological mechanisms underlying TDP-43 driven
    neurodegeneration in FTLD-ALS spectrum disorders. *Hum. Mol. Genet.* **22**, R77–R87
    (2013).

8. Wegorzewska, I. & Baloh, R. H. TDP-43-based animal models of neurodegeneration: new
    insights into ALS pathology and pathophysiology. *Neurodegener. Dis.* **8**, 262–274 (2011).

9. Vahsen, B. F. *et al.* Non-neuronal cells in amyotrophic lateral sclerosis - from pathogenesis to
    biomarkers. *Nat. Rev. Neurol.* **17**, 333–348 (2021).

10. Lee, Y. *et al.* Oligodendroglia metabolically support axons and contribute to
    neurodegeneration. *Nature* **487**, 443–448 (2012).

11. Papadeas, S. T., Kraig, S. E., O'Banion, C., Lepore, A. C. & Maragakis, N. J. Astrocytes carrying the superoxide dismutase 1 (SOD1G93A) mutation induce wild-type motor neuron degeneration in vivo. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 17803–17808 (2011).

12. Joshi, A. U. *et al.* Fragmented mitochondria released from microglia trigger A1 astrocytic response and propagate inflammatory neurodegeneration. *Nat. Neurosci.* **22**, 1635–1648 (2019).

13. Lago-Baldaia, I., Fernandes, V. M. & Ackerman, S. D. More Than Mortar: Glia as Architects of Nervous System Development and Disease. *Front. Cell Dev. Biol.* **8**, 611269 (2020).

14. Izrael, M., Slutsky, S. G. & Revel, M. Rising Stars: Astrocytes as a Therapeutic Target for ALS Disease. *Front. Neurosci.* **14**, 824 (2020).

15. Lee, J. *et al.* Astrocytes and Microglia as Non-cell Autonomous Players in the Pathogenesis of ALS. *Exp. Neurobiol.* **25**, 233–240 (2016).

16. Do-Ha, D., Buskila, Y. & Ooi, L. Impairments in Motor Neurons, Interneurons and Astrocytes Contribute to Hyperexcitability in ALS: Underlying Mechanisms and Paths to Therapy. *Mol. Neurobiol.* **55**, 1410–1418 (2018).

17. Guttenplan, K. A. *et al.* Knockout of reactive astrocyte activating factors slows disease progression in an ALS mouse model. *Nat. Commun.* **11**, 3753 (2020).

18. Lepore, A. C. *et al.* Focal transplantation-based astrocyte replacement is neuroprotective in a model of motor neuron disease. *Nat. Neurosci.* **11**, 1294–1301 (2008).

19. Casci, I. & Pandey, U. B. A fruitful endeavor: modeling ALS in the fruit fly. *Brain Res.* **1607**, 47–74 (2015).

20. Ackerman, S. D., Perez-Catalan, N. A., Freeman, M. R. & Doe, C. Q. Astrocytes close a motor circuit critical period. *Nature* **592**, 414–420 (2021).

21. Tsacopoulos, M. & Magistretti, P. J. Metabolic coupling between glia and neurons. *J. Neurosci. Off. J. Soc. Neurosci.* **16**, 877–885 (1996).

22. Suzuki, A. *et al.* Astrocyte-neuron lactate transport is required for long-term memory formation. *Cell* **144**, 810–823 (2011).

23. Benjamin Kacerovsky, J. & Murai, K. K. Stargazing: Monitoring subcellular dynamics of brain astrocytes. *Neuroscience* **323**, 84–95 (2016).

24. Landgraf, M., Jeffrey, V., Fujioka, M., Jaynes, J. B. & Bate, M. Embryonic origins of a motor system: motor dendrites form a myotopic map in Drosophila. *PLoS Biol.* **1**, E41 (2003).

25. Sales, E. C., Heckman, E. L., Warren, T. L. & Doe, C. Q. Regulation of subcellular dendritic synapse specificity by axon guidance cues. *eLife* **8**, e43478 (2019).

26. Jackson, J. G. & Robinson, M. B. Reciprocal Regulation of Mitochondrial Dynamics and Calcium Signaling in Astrocyte Processes. *J. Neurosci. Off. J. Soc. Neurosci.* **35**, 15199–15213 (2015).

27. Jackson, J. G., O'Donnell, J. C., Takano, H., Coulter, D. A. & Robinson, M. B. Neuronal activity and glutamate uptake decrease mitochondrial mobility in astrocytes and position mitochondria near glutamate transporters. *J. Neurosci. Off. J. Soc. Neurosci.* **34**, 1613–1624 (2014).

28. Maccioni, R. *et al.* Standardized phytotherapic extracts rescue anomalous locomotion and electrophysiological responses of TDP-43 Drosophila melanogaster model of ALS. *Sci. Rep.* **8**, 16002 (2018).

29. Kushimura, Y. *et al.* Overexpression of ter94, Drosophila VCP, improves motor neuron degeneration induced by knockdown of TBPH, Drosophila TDP-43. *Am. J. Neurodegener. Dis.* **7**, 11–31 (2018).

30. Perkins, L. A. *et al.* The Transgenic RNAi Project at Harvard Medical School: Resources and Validation. *Genetics* **201**, 843–852 (2015).

31. Yang, C. *et al.* Partial loss of TDP-43 function causes phenotypes of amyotrophic lateral sclerosis. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E1121-1129 (2014).

32. Devlin, A.-C. *et al.* Human iPSC-derived motoneurons harbouring TARDBP or C9ORF72 ALS mutations are dysfunctional despite maintaining viability. *Nat. Commun.* **6**, 5999 (2015).

33. Sasaki, S., Komori, T. & Iwata, M. Excitatory amino acid transporter 1 and 2 immunoreactivity in the spinal cord in amyotrophic lateral sclerosis. *Acta Neuropathol. (Berl.)* **100**, 138–144 (2000).

34. Lapucci, A. *et al.* Effect of Class II HDAC inhibition on glutamate transporter expression and survival in SOD1-ALS mice. *Neurosci. Lett.* **656**, 120–125 (2017).

35. Kim, K. *et al.* Role of excitatory amino acid transporter-2 (EAAT2) and glutamate in neurodegeneration: opportunities for developing novel therapeutics. *J. Cell. Physiol.* **226**, 2484–2493 (2011).

36. Besson, M. T., Soustelle, L. & Birman, S. Identification and structural characterization of two genes encoding glutamate transporter homologues differently expressed in the nervous system of Drosophila melanogaster. *FEBS Lett.* **443**, 97–104 (1999).

37. Diaper, D. C. *et al.* Drosophila TDP-43 dysfunction in glia and muscle cells cause cytological and behavioural phenotypes that characterize ALS and FTLD. *Hum. Mol. Genet.* **22**, 3883–3893 (2013).

38. Romano, G. *et al.* Glial TDP-43 regulates axon wrapping, GluRIIA clustering and fly motility by autonomous and non-autonomous mechanisms. *Hum. Mol. Genet.* **24**, 6134–6145 (2015).

39. Stephen, T.-L. *et al.* Miro1 Regulates Activity-Driven Positioning of Mitochondria within Astrocytic Processes Apposed to Synapses to Regulate Intracellular Calcium Signaling. *J. Neurosci. Off. J. Soc. Neurosci.* **35**, 15996–16011 (2015).

40. Glater, E. E., Megeath, L. J., Stowers, R. S. & Schwarz, T. L. Axonal transport of mitochondria requires milton to recruit kinesin heavy chain and is light chain independent. *J. Cell Biol.* **173**, 545–557 (2006).

41. Schwarz, T. L. Mitochondrial trafficking in neurons. *Cold Spring Harb. Perspect. Biol.* **5**, a011304 (2013).

42. Stork, T., Sheehan, A., Tasdemir-Yilmaz, O. E. & Freeman, M. R. Neuron-glia interactions through the Heartless FGF receptor signaling pathway mediate morphogenesis of Drosophila astrocytes. *Neuron* **83**, 388–403 (2014).

43. Risse, B. *et al.* FIM, a novel FTIR-based imaging method for high throughput locomotion analysis. *PloS One* **8**, e53963 (2013).

44. Landgraf, M., Bossing, T., Technau, G. M. & Bate, M. The origin, location, and projections of the embryonic abdominal motorneurons of Drosophila. *J. Neurosci. Off. J. Soc. Neurosci.* **17**, 9642–9655 (1997).

45. Goel, P. & Dickman, D. Synaptic homeostats: latent plasticity revealed at the Drosophila neuromuscular junction. *Cell. Mol. Life Sci. CMLS* **78**, 3159–3179 (2021).

46. Escartin, C. *et al.* Reactive astrocyte nomenclature, definitions, and future directions. *Nat. Neurosci.* **24**, 312–325 (2021).

47. Andriezen, W. L. The Neuroglia Elements in the Human Brain. *Br. Med. J.* **2**, 227–230 (1893).

48. Fridovich, I. Superoxide dismutases. *Adv. Enzymol. Relat. Areas Mol. Biol.* **58**, 61–97 (1986).

49. Barber, S. C., Mead, R. J. & Shaw, P. J. Oxidative stress in ALS: a mechanism of neurodegeneration and a therapeutic target. *Biochim. Biophys. Acta* **1762**, 1051–1067 (2006).

50. Cassina, P. *et al.* Mitochondrial dysfunction in SOD1G93A-bearing astrocytes promotes motor neuron degeneration: prevention by mitochondrial-targeted antioxidants. *J. Neurosci. Off. J. Soc. Neurosci.* **28**, 4115–4122 (2008).

51. Nagai, M. *et al.* Astrocytes expressing ALS-linked mutated SOD1 release factors selectively toxic to motor neurons. *Nat. Neurosci.* **10**, 615–622 (2007).

52. Romano, G. *et al.* Downregulation of glutamic acid decarboxylase in Drosophila TDP-43-null brains provokes paralysis by affecting the organization of the neuromuscular synapses. *Sci. Rep.* **8**, 1809 (2018).

53. Paradis, S., Sweeney, S. T. & Davis, G. W. Homeostatic control of presynaptic release is triggered by postsynaptic membrane depolarization. *Neuron* **30**, 737–749 (2001).

54. Howland, D. S. *et al.* Focal loss of the glutamate transporter EAAT2 in a transgenic rat model of SOD1 mutant-mediated amyotrophic lateral sclerosis (ALS). *Proc. Natl. Acad. Sci. U. S. A.* **99**, 1604–1609 (2002).

55. Guo, H. *et al.* Increased expression of the glial glutamate transporter EAAT2 modulates excitotoxicity and delays the onset but not the outcome of ALS in mice. *Hum. Mol. Genet.* **12**, 2519–2532 (2003).

56. Rothstein, J. D. *et al.* Beta-lactam antibiotics offer neuroprotection by increasing glutamate transporter expression. *Nature* **433**, 73–77 (2005).

 METHODS

Lead Contact and Materials Availability

Additional information and inquiries regarding resources and reagents should be directed to

and will be fulfilled upon request by Lead Contact Sarah D Ackerman

(sarah.ackerman@wustl.edu).

Experimental Model and Subject Details

Fly husbandry

We raised all fly stocks at 25°C on standard cornmeal fly food. Animals were staged relative

to a 25°C standard. At 25°C, embryos take 21 hours to hatch into larvae[57]; larvae hatch and

develop 1.25x faster at 30°C, and 2x slower at 18°C.

Transgenes (in order of appearance)

1. *RN2-lexA*[20]

2. *13lexAop-CsChrimson::tdTomato*[20]

3. *R25H07-gal4* (BDSC# 49145)

4. *UAS-mito-HA-GFP/cyo* (BDSC# 8442)

5. *13xLexAop2-CD4-tdTom/T6b* (BDSC# 77139)

6. *RN2-gal4*[24]

7. *UAS-GtACR2::EYFP*[58]

8. *alrm-lexA::GAD* (Chromosome 3)[59]

9. *lexAop-mCherry::mito.OMM* (BDSC# 66530)

10. *10XUAS-myr::GFP* (BDSC# 32198)

11. *UAS-tbph*[RNAi] (BDSC# 39014)

12. *UAS-mito-HA-GFP* (chromosome 3, BDSC# 8443)

13. *UAS-luciferase*[RNAi] (BDSC# 31603)

14. *13XLexAop-IVS-jGCaMP7s* (BDSC# 80913)

15. *alrm-gal4* (Chromosome 2)[59]

16. *10XUAS-mcd8::GFP* (BDSC# 5137)

17. *UAS-milton*[RNAi] (BDSC#s 28385, 43173, 44477)

18. *UAS-khc*[RNAi] (BDSC#s 35409, 35770)

19. *UAS-miro*[RNAi] (BDSC# 43973)

20. *UAS-Dhc64C* (BDSC# 36583, 36698)

21. *Mi{Trojan-lexA:QFAD.2}VGlut[MI04979-TlexA:QFAD.2]/CyO* (BDSC# 60314)

22. *13XLexAop2-IVS-myr::GFP* (X Chromosome, BDSC# 32211)

Animal collections

For imaging of wild-type brains without optogenetics (Figures 2-3)*:* We reared crosses at 25°C in collection bottles fitted with 3.0% agar apple juice caps seeded with plain yeast paste (Baker's year + water). We then collected embryos on fresh 3.0% agar apple juice caps seeded with plain yeast paste for 1.5 hours (h) and aged at 25°C until hatching. At 0 h after larval hatching (ALH), larvae were collected and transferred to fresh apple caps (N=20 larvae per cap) containing yeast paste and aged until 8 h ALH, when they were dissected and prepared for imaging.

For imaging of RNAi brains without optogenetics (Figures 4-6, 8)*:* We reared crosses at 25°C in collection bottles fitted with 3.0% agar apple juice caps seeded with plain yeast paste (Baker's year + water). We then collected embryos on fresh 3.0% agar apple juice caps seeded with plain yeast paste for 1.5 hours (h) and aged at 30°C until hatching to maximize RNAi expression. For assessment of mitochondrial trafficking, larvae were collected at 0 h ALH and transferred to apple caps (N=20 larvae per cap) containing fresh year paste, aged 6.5 h (~ 8 h ALH at 25°C standard), when they were dissected and prepared for imaging. For assessment of synapse number at the neuromuscular junction, apple caps were transferred to fresh food bottles (corn meal) when animals reached 0 h ALH. Bottles were maintained at 30°C for 77 h (~ 96 h ALH at 25°C standard), when larvae were collected, dissected, and prepared for imaging.

For optogenetics on control animals (Figures 2-3)*:* We reared crosses at 25°C in collection bottles fitted with 3.0% agar apple juice caps seeded with fresh yeast paste that was supplemented with 0.5mM all-*trans* retinal (+ATR) (Sigma-Aldrich, R2500-100MG). Crosses were supplied fresh yeast paste daily (+ATR) for a minimum of 72 h prior to embryo collection to ensure maternal transfer of ATR into embryos. We then collected embryos on 3.0% agar apple juice caps with yeast paste (+ATR) for 1.5 h and aged at 25°C. To prevent premature optogenetic activation, we kept all crosses, embryos, and larvae in dark conditions until the appropriate developmental stage. At 25° C, *Drosophila* embryos hatch at 21 h after egg laying (AEL). Larvae were collected at 0 h ALH and transferred to fresh apple caps (N=10 larvae per cap) containing yeast paste (+ATR). For fixed preparations, larvae were then transferred to activating/silencing conditions for 1 hour and dissected at 8 h ALH. For

live imaging experiments, larvae were dissected at 8 h ALH and transferred to the confocal

microscope for manipulation/recording (read more below).

For optogenetics with RNAi (Figure 4)*:* We reared crosses at 25°C in collection bottles fitted

with 3.0% agar apple juice caps seeded with fresh yeast paste that was supplemented with

0.5mM all-*trans* retinal (+ATR) (Sigma-Aldrich, R2500-100MG). Crosses were supplied

fresh yeast paste daily (+ATR) for a minimum of 72 h prior to embryo collection to ensure

maternal transfer of ATR into embryos. We then collected embryos on 3.0% agar apple juice

caps with yeast paste (+ATR) for 1.5 h and aged at 30°C. To prevent premature optogenetic

activation, we kept all crosses, embryos, and larvae in dark conditions until the appropriate

developmental stage. At 0 h ALH, we transferred larvae to fresh apple caps (N=10 larvae per

cap) containing yeast paste (+ATR) and aged 6.5 h (~ 8 h ALH at 25°C standard) prior to

dissection for dark-reared controls, or aged 5.5 h followed by 1 h light activation. All animals

were dissected immediately following activity manipulation.

For behavioral analyses: We reared crosses at 25°C in collection bottles fitted with 3.0% agar

apple juice caps containing plain yeast paste. Embryos were then collected on 3.0% agar

apple juice caps with plain yeast paste for 1.5 h and aged at 30°C. Apple caps were

transferred to fresh food bottles (corn meal) when animals reached 0 h ALH. Bottles were

maintained at 30°C for 60 h (~ 72 h ALH at 25°C standard) until larvae reached wandering 3[rd]

instar stage, when they were transferred to the behavioral rig for testing.

METHOD DETAILS

<u>Sustained optogenetic activation/silencing</u>

We used the following published strategy[20] for optogenetic activation/silencing in all fixed

preparation experiments (Figures 2-4). At the designated stage, dark-reared larvae

(maintained on apple caps with yeast + ATR) were placed beneath a full spectrum light bulb,

shining with an average intensity of 10550 lx (determined with two independent photometer

software programs developed for Android: Light Meter© and Photometer©). We then

dissected larvae in low-light conditions (<100 lx) immediately following activity

manipulations. Yeast paste was placed in the middle of the apple cap to motivate larvae to

remain on the top of the agar plate throughout the duration of the manipulation. Any larvae

found outside of this area were not processed further.

<u>Immunohistochemistry</u>

Larval brains were dissected in sterile-filtered, ice-cold 1X PBS and mounted on 12mm #1

thickness poly-D-lysine coated round coverslips (Corning® BioCoat™, 354085 or German

Glass Cover Slips, Fisher Scientific, 72196-12). We fixed brains for 12 minutes in fresh 4%

paraformaldehyde (Electron Microscopy Sciences, 15710) in .3% PBSTriton, and then

washed in .3% PBSTriton to remove fixative. For larval fillet preps, we dissected larvae in

sterile-filtered, ice-cold HL3.1, fixed for 45 minutes in fresh 4% paraformaldehyde in

detergent-free 1X PBS (Electron Microscopy Sciences, 15710), and then washed in .3%

PBSTriton to remove fixative. At this point, all samples were treated the same way through

post-secondary washes. Samples were blocked overnight at 4°C in .3% PBSTriton

supplemented with 1% BSA (Fisher, BP1600-100), 1% normal donkey serum and 1% normal

goat serum (Jackson ImmunoResearch Laboratories, Inc., 017-000-121 and 005-000-121).

Samples were then incubated in primary antibody for one-two days at 4°C. The primary was

removed, and brains were washed overnight at 4°C with 0.3% PBSTriton. Samples were then

incubated in secondary antibodies overnight at 4°C. The secondary antibodies were removed,

and samples transferred to .3% PBSTriton overnight. Larval brains were then prepared for

DPX mounting. Brains were dehydrated with an ethanol series: 30%, 50%, 70%, 90%, each

for 5 minutes, then twice in 100% ethanol for 10 minutes each (Decon Labs, Inc., 2716GEA).

Finally, samples were incubated in xylenes (Fisher Chemical, X5-1) for 2 x 10 minutes, were

mounted onto slides containing DPX mountant (Millipore Sigma, 06552), and cured for 1-2

days before imaging. After post-secondary .3% PBSTriton washes, larval fillet preps were

washed into 1X PBS overnight. Fillet preps were then incubated in Phalloidin-405 (Alexa

Fluor™ Plus 405 Phalloidin, a30104) in 1X PBS for 5 minutes, washed back into 1X PBS,

and mounted in ProLong Glass Antifade Mountant (Thermo Fisher Scientific, P36982).

The following primary and secondary antibodies were used:

Primary Antibody (concentration)	Source	Figures
Chicken anti-GFP (1:1000)	Aves Cat. GFP-1010	1-5, 8
Rabbit anti-dsred (1:500)	Takara Bio Cat. 632496	1-2,4
Rabbit anti-cherry (1:500)	Novus Biologicals, Cat. NBP2-25157	1,3

Ms anti-Brp/Nc82 (1:100)	DSHB nc82	1-4, 6
Rabbit anti-EAAT1 (1:5000)	D. Van Meyel lab[60]	5
Rabbit anti-Gat (1:4000)	M. Freeman lab[42]	6
Ms anti-Dlg (1:100)	DSHB 4F3	8
Phalloidin-405 (1:40)	Alexa Fluor™, a30104	8

Secondary Antibodies:

All secondary antibodies were purchased from Jackson ImmunoResearch and used at a

working concentration of 1:400. The following antibodies were used: Alexa Fluor®

Rhodamine RedTM-X Donkey-Anti Mouse (715-295-151), Alexa Fluor® 647 AffiniPure

Donkey Anti-Mouse IgG (715-605-151), Alexa Fluor® 488 Donkey anti-Chicken (703-545-

155), Alexa Fluor® 647 Donkey-Anti Rabbit (711-605-152); Alexa Fluor® 488 Donkey anti-

Rabbit (711-545-152), Alexa Fluor® Rhodamine RedTM-X Donkey Anti-Rabbit (711-295-

152).

<u>Light Microscopy</u>

Fixed larval preparations for synapse quantifications (central and NMJ, Figures 1-4, 8) and all

live imaging experiments (Figures 2-5, Supplemental Movies 1-11) were imaged on a Zeiss

LSM 800 laser scanning confocal fitted with a 63x/1.40 NA Oil Plan-Apochromat DIC m27

objective lens and GaAsP photomultiplier tubes. Fixed larval preparations for astrocyte

morphology analyses (Figure 6) were imaged with a Zeiss LSM 700 laser scanning confocal

using a 63x/1.4 NA Oil Plan-Apochromat DIC m27 objective lens.

 Image processing and analyses

 Quantitative analyses were performed either in Imaris 9.2.0 or higher (Bitplane 9.2.0, Figures

 1-4, 6) or in FIJI (ImageJ 1.50d, https://imagej.net/Fiji, Figure 5, 8).

 Time-lapse imaging of fictive preparations

 The following assay was used for live imaging of aCC/RP2 dendrites and/or astrocyte

 mitochondria in isolated CNS (Figures 2-3, 5, Supplemental Movies 1-11). Larvae were

 dissected at 8 h ALH in a hemolymph-like solution (HL3.1); both lobes and the ventral nerve

 cord were kept intact. Isolated brains were mounted dorsal-side down on a 12mm #1 thickness

 poly-D-lysine coated round coverslips (Corning® BioCoat™, 354085 or German Glass Cover

 Slips, Fisher Scientific, 72196-12). The cover slip was then inverted onto a slide with vacuum

 grease as a bridge and pressed down until brain lobes were sandwiched between the slide and

 cover glass. Samples that were damaged during mounted were not imaged. Samples were

 imaged on a Zeiss LSM 800 laser scanning confocal fitted with a 63x/1.40 NA Oil Plan-

 Apochromat DIC m27 objective lens and GaAsP photomultiplier tubes, and imaged using a

 488 nm laser and/or 561 nm laser when appropriate. For optogenetic experiments (Figures 2-

 3), both dissections and mounting were performed under low light conditions (<100 lx) to

 delay optogenetic activation. Continuous scans were obtained every 30 seconds for 15

 minutes, with two hemisegments in the field of view. A z-stack of 12 μm (allowing for drift in

 Z) with .5 μm step size was performed. Finally, we imaged abdominal segments 5-6, which

 were less prone to drift over the 15-minute acquisition. For GCaMP7s assays (Figure 5), a z-

 stack of 19 μm (allowing for drift in Z) with 1 μm step size was imaged continuously with

 stacks acquired every 15 seconds for 15 minutes, with abdominal segments 3-7 in view. For

our mitochondria machinery screen (Supplemental Movies 7-11), we obtained 50 μm stacks

(step size of 1 μm) once a minute for 15 minutes, with abdominal segments 5-8 in view.

Behavioral analyses

We transferred larvae at 72 h ALH from food bottles (standard corn meal) to 1.2% agarose

plates for half an hour prior to locomotion assays to allow the larvae to acclimate to the new

crawling surface. Larvae were then transferred to a FIM behavior table[43] fitted with a fresh

1.2% agarose gel and allowed to further acclimate for two minutes prior to imaging. Two or

more independent cohorts of larvae (N=30 larvae per cohort) were tested per genotype. Larval

crawling was imaged at 4 Hz, 91 pixels/cm for one minute using a Basler acA2040-25gm

camera in the Pylon5 Camera Software Suite (Basler). Data were then analyzed using

FIMTrack software using standard settings[43]. Data files were blinded to genotype until after

analysis.

Figure preparation

Images in figures were prepared in Imaris (Bitplane AG, version 9.2.0 or higher) or in Fiji

(ImageJ 1.50d). Both software packages were used for rendering 3D, max, or orthogonal

projections of the data, which were then assembled using Adobe Illustrator. Schematics were

drawn in Microsoft Powerpoint.

QUANTIFICATION AND STATISTIC ANALYSIS

Synapse quantification: association of astrocyte mitochondria with presynapses

aCC/RP2-associated presynapses

For quantification of astrocyte mitochondria in close proximity to aCC/RP2-associated presynapses (Figures 1-4), data were acquired with a voxel size of .099 x .099 x .23 μm^3 and de-convoluted in Imaris. First, aCC/RP2 dendrites within a single hemisegment were reconstructed using the Imaris "Surface" module (no smoothing, thresholds varied with fluorescence intensity). A standard ROI spanned 100 x 100 pixels in XY, and 7 μm in Z with the top of the ROI beginning dorsally at the aCC/RP2 axons. The dendrite "Surface" was also used to determine dendrite volume. A distance transformation was then performed on the dendrite "Surface". Second, Brp+ presynapses within the ROI were annotated using the Imaris "Spots" functions and then classified as verified post-synapses if they fell within 90 nm of the "Surface" (accounts for chromatic aberration and the size of the synaptic cleft) based on previously validated criteria[20,25], and were used to build a new channel. Third, astrocyte mitochondria within the same ROI were reconstructed using the Imaris "Surface" module (.1 μm smoothing, thresholds varied with fluorescence intensity). A distance transformation was then performed on the astrocyte mitochondria "Surface". Finally, verified presynapse "Spots" were classified as mitochondria-associated if they fell within 120 nm of the astrocyte mitochondria "Surface" (accounts for chromatic aberration[25], the thickness of the astrocyte plasma membrane[61], and the thickness of both mitochondrial membranes[62]).

Astrocyte association with all presynapses

For quantification of astrocyte membrane and mitochondria-associated presynapses (Figure 6), data were acquired with a voxel size of .2 x .2 x .39 μm^3. Imaris reconstructions were applied to the entire larval brain. First, we reconstructed astrocyte membranes using the Imaris "Surface" module (no smoothing, thresholds varied with fluorescence intensity). A

distance transformation was performed on the membrane "Surface." Then, we constructed

astrocyte mitochondria using the Imaris "Surface" module (no smoothing, thresholds varied

with fluorescence intensity). A second distance transformation was performed on the

mitochondria "Surface." Finally, Brp+ presynapses were annotated using the Imaris "Spots"

functions and then categorized by proximity ($\leq$ 1 µm) to the astrocyte mitochondria or

membrane "Surface."

Analysis of time-lapse imaging samples (optogenetics)

For analysis of mitochondrial dynamics over time (Figures 2-3), data were collected with a

voxel size of .085 x .085 x .5 µm^3. Any samples that drifted out of the frame of view during

the acquisition period were not processed further. Data were then imported to Imaris and

analyzed using the "Surface" module (.1 µm smoothing, thresholding varied by fluorescence

intensity of sample). A standard ROI spanned 200 x 200 pixels in XY, and 12.5 µm in Z.

First, we built a "Surface" to mask the motor dendrites. Then, we built a "Surface" to mask

the astrocyte mitochondria. We then used the "Surface-Surface" overlap function (Imaris

9.7.0 or later) to calculate the volume of all mitochondria within the dendritic domain ($\leq$ 120

nm from motor dendrites), and to determine the total volume of "Surface" overlap (*aka*

volume of membrane overlap).

Analysis of GCaMP imaging data

Image files were blinded to genotype. Raw time-lapse image stacks were imported into Fiji and

converted to time-lapse maximum intensity projections. These converted files were loaded into a

custom MATLAB script[25]. The script first performs rigid registration to correct for movement

artifacts during recording, and then allows for ROI selection. ROIs were drawn around aCC/RP2

dendrites in 6 individual hemisegments, 3 per side. ROI size was constant across all analyzed files

(Figure 5). The script then quantifies and plots the frame-by-frame average dF/F within each ROI.

To identify peaks of calcium activity in an unbiased manner, our dF/F traces were next

loaded into the MATLAB program PeakCaller[63]. In PeakCaller, the following parameters were

specified for the program to identify peaks in each trace: Required Rise = 20%, Required Fall =

30%, Max Lookback and Lookahead pts = 10 frames. Peaks < dF/F=0.1 were excluded from

analysis. We used PeakCaller to identify the number of calcium peaks in each dF/F trace. We

logged the amplitude of each peak (dF/F) within a trace, and averaged these amplitudes together

to obtain a single average peak fluorescence value for each ROI.

After quantifying the frequency and amplitude of peaks for each ROI, sample files were

unblinded, and statistical comparisons were performed between Control and *tbph* RNAi groups

using a two-tailed unpaired t-test with unequal variance.

Analysis of EAAT1 levels on astrocyte membranes

Data were acquired with a voxel size of .099 x .099 x .23 μm^3 and imported into Fiji. For each

sample, raw integrated density (sum of fluorescence intensity unites) was measured from a

standard ROI covering 1313 μm of the neuropil in the anterior ventral nerve cord. We measured

raw integrated density for both EAAT1 and mcd8::GFP (astrocyte membrane marker) from four

independent z planes per sample. Finally, we normalized EAAT1 expression to mcd8::GFP

expression to account for any potential impacts of *tbph* knockdown of astrocyte membrane

elaboration (Figure 5).

 Analysis of synaptic boutons in larval fillet preparations

 Data were acquired from dorsal muscles only (DO1-DO2, DA1-DA2) with a voxel size of

 .198 x .198 x .3 μm^3. Samples were blinded and imported into Fiji, where we performed a

 max intensity projection of the channel corresponding to *vglut-lexA, lexAop-myr::GFP* (labels

 all motor neuron projections and synaptic boutons). Synaptic boutons were quantified

 manually using the Fiji "Multipoint" tool. After quantifying bouton numbers for an entire

 experimental set, sample files were unblinded and statistical comparisons were performed

 using a two-tailed unpaired t-test with unequal variance.

 Statistical analyses

 Statistics were performed using a combination of Microsoft Excel, MATLAB (MathWorks), and

 Prism (GraphPad) software. We used a combination of one-way ANOVA, Student's T-Tests

 (two-tailed, unequal variance), and Mann-Whitney Tests where appropriate (noted throughout).

 Error bars, Standard Error of the Mean unless otherwise noted. A 95% confidence interval was

 used to define the level of significance. Significance: *, $p<.05$; **, $p<.01$; ***, $p<.001$; ****,

 $p<.0001$, NS= not significant. All other pertinent information, including sample size, statistical

 test employed, and variance can be found in the figure legends or labeled within the figure.

METHOD REFERENCES

57. Crisp, S., Evers, J. F., Fiala, A. & Bate, M. The development of motor coordination in Drosophila embryos. *Dev. Camb. Engl.* **135**, 3707–3717 (2008).

58. Mauss, A. S., Busch, C. & Borst, A. Optogenetic Neuronal Silencing in Drosophila during Visual Processing. *Sci. Rep.* **7**, 13823 (2017).

59. Doherty, J., Logan, M. A., Taşdemir, O. E. & Freeman, M. R. Ensheathing glia function as phagocytes in the adult Drosophila brain. *J. Neurosci. Off. J. Soc. Neurosci.* **29**, 4768–4781 (2009).

60. Peco, E. *et al.* Drosophila astrocytes cover specific territories of the CNS neuropil and are instructed to differentiate by Prospero, a key effector of Notch. *Dev. Camb. Engl.* **143**, 1170–1181 (2016).

61. Ingólfsson, H. I. *et al.* Computational Lipidomics of the Neuronal Plasma Membrane. *Biophys. J.* **113**, 2271–2280 (2017).

62. Perkins, G. *et al.* Electron tomography of neuronal mitochondria: three-dimensional structure and organization of cristae and membrane contacts. *J. Struct. Biol.* **119**, 260–272 (1997).

63. Artimovich, E., Jackson, R. K., Kilander, M. B. C., Lin, Y.-C. & Nestor, M. W. PeakCaller: an automated graphical interface for the quantification of intracellular calcium obtained by high-content screening. *BMC Neurosci.* **18**, 72 (2017).

 FIGURES AND LEGENDS

Figure 1. Mapping astrocyte-motor neuron, mitochondria-synapse associations in

***Drosophila*.**

(**a**) Schematic for reader orientation. A, anterior. P, posterior. Astrocyte mitochondria (orange)

are distributed throughout the synapse-rich neuropil (cyan) near motor dendrites (magenta).

(**b**) Pipeline for identifying motor synapses in close proximity to astrocyte mitochondria at 8

hours after larval hatching (h ALH). (i-ii) Acquire confocal z-stack through neuropil with

fluorescently labeled astrocyte mitochondria (*R25H07-gal4, UAS-mito::GFP* or *alrm-lexA,*

799   *lexAop-mcherry::mito.OMM*), aCC/RP2 motor neurons (MN; *RN2-lexA, lexAop-TdTomato* or

800   *RN2-gal4, UAS-myr::GFP*), and presynapses (Brp+). Scale bar, 10 μm. (iii-iv) Mask the motor

801   dendrites with an Imaris "Surface." (v-vi) Identify presynapses colocalized with motor dendrites

802   (≤90 nm) and generate Imaris "Spots". (vii-viii) Mask astrocyte mitochondria with an Imaris

803   "Surface." (ix-x) Identify Spots within 120 nm of masked mitochondria. (iii-ix) Scale bar, 5 μm.

804   For more details, see Methods.

805

MN activation recruits astrocyte mitochondria to dendrites & synapses

Figure 2. Activation of MNs recruits astrocyte mitochondria to motor dendrites and synapses.

(**a-b"**) Representative images of dark-reared control (**a-a"**) and 1-hour activated (**b-b"**) motor dendrites (magenta, *RN2-lexA, lexAop-Chrimson::TdTomato)* with neighboring astrocyte synapses (orange, *R25H07-gal4, UAS-mito::GFP*) and presynapses (cyan, Brp+) in fixed preparations at 8 h ALH. Prime panels show synapse "Spots" co-localized ($\leq$ 120 nm) with astrocyte mitochondria. Double prime panels show orthogonal views of motor dendrites (top) and astrocyte mitochondria (bottom) within the dendritic domain (yellow circle). (**a-a'**, **b-b'**) Scale bar, 2 µm. (**a"**,**b"**) Scale bar, 10 µm.

(**c**) Quantification of the percent of synapses associated with astrocyte mitochondria ($p<.0001$, Mann-Whitney test). Control: n=16 animals. Activated: n=9 animals. Error bars = SEM.

(**d**) Quantification of the mitochondrial volume in the dendritic domain ($p=.044$, Mann-Whitney test). Control: n=19 animals. Activated: n=20 animals. Error bars = SEM.

(**e**) Quantification of dendrite volume ($p=.14$, Mann-Whitney test). Control: n=11 animals. Activated: n=9 animals. Error bars = SEM. NS, not significant.

(**c-e**) N values reflect animals from 3 technical replicates.

(**f-g**) Quantification of (**f**) astrocyte mitochondrial volume (*R25H07-gal4, UAS-mito::GFP*) within the dendritic domain and (**g**) overlapping mitochondria-dendrite volume in live samples (8 h ALH) imaged every 30 seconds for a 15-minute window of activation (pink, *RN2-lexA, lexAop-Chrimson::TdTomato*) or control (grey, *RN2-lexA, lexAop-myr::TdTomato*). Control: n=14 animals. Activation: n=15 animals. Green areas denote timepoints when $p<.05$ by one-way ANOVA. Error bars = SEM. N values reflect animals from 2 technical replicates.

MN silencing repels astrocyte mitochondria from dendrites

Figure 3. Silencing of MNs repels astrocyte mitochondria from motor dendrites.

(**a-b"**) Representative images of dark-reared control (**a-a"**) and 1-hour silenced (**b-b"**) motor

dendrites (magenta, *RN2-gal4, UAS-GtACR2::EYFP)* with neighboring astrocyte synapses

(orange, *alrm-lexA, lexAop-mcherry::mito.OMM*) and presynapses (cyan, Brp+) in fixed

preparations at 8 h ALH. Prime panels show synapse "Spots" co-localized ($\leq$ 120 nm) with

astrocyte mitochondria. Double prime panels show orthogonal views of motor dendrites (top) and

astrocyte mitochondria (bottom) within the dendritic domain (yellow circle). (**a-a'**, **b-b'**) Scale

bar, 2 μm. (**a"**,**b"**) Scale bar, 10 μm.

(**c**) Quantification of the percent of synapses associated with astrocyte mitochondria (p=.26,

Mann-Whitney test). Control: n=17 animals. Silenced: n=19 animals. Error bars = SEM. NS, not

significant.

(**d**) Quantification of the mitochondrial volume in the dendritic domain (p=.003, Mann-Whitney

test). Control: n=17 animals. Silenced: n=19 animals. Error bars = SEM.

(**e**) Quantification of dendrite volume (p=.1, Mann-Whitney test). Control: n=17 animals.

Silenced: n=19 animals. Error bars = SEM.

(**c-e**) N values reflect animals from 3 technical replicates.

(**f-g**) Quantification of (**f**) astrocyte mitochondrial volume (*alrm-lexA, lexAop-*

mcherry::mito.OMM) within the dendritic domain and (**g**) overlapping mitochondria-dendrite

volume in live samples (8 h ALH) imaged every 30 seconds for a 15-minute window of silencing

(green, *RN2-gal4, UAS-GtACR2::EYFP*) or control (grey, *RN2-gal4, UAS-myr::GFP*). Control:

n=11 animals. Silencing: n=9 animals. Error bars = SEM. N values reflect animals from 2

technical replicates.

Astrocyte knockdown of tbph enhances mitochondrial trafficking

Figure 4. Astrocyte knockdown of *tbph* enhances activity-dependent mitochondria-synapse association.

(**a-b"**) Representative images of dark-reared control (**a-a"**) and 1-hour activated (**b-b"**) motor dendrites (magenta, *RN2-lexA, lexAop-Chrimson::TdTomato)* with neighboring astrocyte synapses (orange, *R25H07-gal4, UAS-mito::GFP*) and presynapses (cyan, Brp+) in fixed preparations of control (*UAS-luciferase*[RNAi]) animals at 8 h ALH. Prime panels show synapse "Spots" co-localized ($\leq$ 120 nm) with astrocyte mitochondria. Double prime panels show orthogonal views of motor dendrites (top) and astrocyte mitochondria (bottom) within the dendritic domain (yellow circle). (**a-a'**, **b-b'**) Scale bar, 2 µm. (**a"**,**b"**) Scale bar, 10 µm.

(**c-d"**) Representative images of dark-reared control (**a-a"**) and 1-hour activated (**b-b"**) motor dendrites (magenta, *RN2-lexA, lexAop-Chrimson::TdTomato)* with neighboring astrocyte synapses (orange, *R25H07-gal4, UAS-mito::GFP*) and presynapses (cyan, Brp+) in fixed preparations of astrocyte *tbph* knockdown (*UAS-tbph*[RNAi]) animals at 8 h ALH. Prime panels show synapse "Spots" co-localized ($\leq$ 120 nm) with astrocyte mitochondria. Double prime panels show orthogonal views of motor dendrites (top) and astrocyte mitochondria (bottom) within the dendritic domain (yellow circle). (**c-c'**, **d-d'**) Scale bar, 2 µm. (**c"**,**d"**) Scale bar, 10 µm.

(**e**) Quantification of the percent of synapses associated with astrocyte mitochondria. Statistical significance quantified by one-way ANOVA: dark-reared *versus* activated control ($p<.0001$); dark-reared *versus* activated *tbph*[RNAi] ($p<.03$); dark-reared comparisons ($p<.03$); activated comparisons ($p<.03$).

(**f**) Quantification of mitochondrial volume in the dendritic domain. Statistical significance quantified by one-way ANOVA: dark-reared *versus* activated control ($p<.005$); dark-reared

versus activated *tbph*^RNAi (p<.13); dark-reared comparisons (p<.008); activated comparisons

(p<.04). NS, not significant.

(**g**) Quantification of dendrite volume. Statistical significance quantified by one-way ANOVA:

dark-reared *versus* activated control (p<.55); dark-reared *versus* activated *tbph*^RNAi (p<.84); dark-

reared comparisons (p<.3); activated comparisons (p<.15).

(**e-g**) Dark-reared control: n=16 animals. Activated control: n=15 animals. Dark-reared *tbph*^RNAi:

n=8 animals. Activated *tbph*^RNAi: n=6 animals. Error bars = SEM. N values reflect animals from 3

technical replicates.

Astrocyte *tbph*^RNAi increases frequency and amplitude of MN activity

a

Basal Activity | Peak Activity

GCaMP6s ROIs

i ii iii

8 h ALH

b

# of Calcium Spikes

Control | *tbph*^RNAi

c

Average Amplitude (dF/F)

Control | *tbph*^RNAi

luciferase^RNAi Control

d

Left_i
Left_ii
Left_iii

Right_i
Right_ii
Right_iii

dF/F_0

0 Time (s) 200

astro>*tbph*^RNAi

e

Left_i
Left_ii
Left_iii

Right_i
Right_ii
Right_iii

dF/F_0

0 Time (s) 200

Knockdown of tbph causes significant reduction in astrocyte EAAT1

luciferase^RNAi Control | astro>*tbph*^RNAi

f **g**

alrm>GFP EAAT1

8 h ALH

f' **g'**

EAAT1

h

Normalized EAAT1 expression

control | *tbph*^RNAi

888

Figure 5. Astrocyte *tbph* knockdown increases frequency and amplitude of motor neuron activity.

(**a**) Maximum intensity projection of GCaMP7s activity in aCC/RP2 (*RN2-lexA, lexAop-GCaMP7s*) in control at basal (left) and peak (right) activity at 8 h ALH. Anterior to the left, dorsal is up. Pink boxes are ROIs used for dF/F quantification.

(**b**) Frequency of calcium peak events for control (gray, *R25H07-gal4, UAS-luciferase*^{RNAi}) and *tbph* knockdown (pink, *R25H07-gal4, UAS-tbph*^{RNAi}) animals. *tbph*^{RNAi} animals exhibited a higher frequency of calcium peaks (Student's t-test, p=0.023). Each point represents an individual hemisegment. Control: n=39 hemisegments from 7 animals, *tbph*^{RNAi} : n=42 hemisegments from 7 animals. Error bars = SD.

(**c**) Average dF/F amplitude of calcium peaks for control (gray) and *tbph*^{RNAi} (pink) animals. *tbph*^{RNAi} animals exhibited an increase in calcium peak amplitude (Student's t-test, p=0.013). Each point represents an individual hemisegment. Control: n=39 hemisegments from 7 animals, *tbph*^{RNAi} : n=42 hemisegments from 7 animals. Error bars = SD.

(**b-c**) N values reflect animals from 3 technical replicates.

(**d-e**) Representative dF/F traces from a single control (**d**) and *tbph*^{RNAi} (**e**) animal. Each trace is from an individual hemisegment. Hemisegment position (Left/Right, Anterior(i)-Posterior(iii)) indicated to the right of each trace. Peaks of calcium activity identified by PeakCaller (see Methods) are shown as blue dots in each trace.

(**f-g**) Orthogonal view showing EAAT1 expression in (**f**) control astrocytes (*alrm-gal4, UAS-mcd8GFP*) or (**g**) astrocytes following *tbph* RNAi knockdown (*UAS-tbph*^{RNAi}) at 8 h ALH. Prime panels show EAAT1 staining alone. Control: n=40 sections from 10 animals, *tbph*^{RNAi}: n=52 segments from 13 animals. N values reflect animals from 2 technical replicates.

912   (**h**) Quantification of EAAT1 expression relative to membrane bound GFP on astrocyte

913   membranes (p=.018, Mann-Whitney test). Error bars = SEM.

Figure 6. Astrocyte mitochondria traffic along (+) ended microtubules.

(**a-e"**) Representative orthogonal views of astrocyte membranes (Gat+, green), mitochondria

(*R25H07-gal4, UAS-mito::GFP*, orange), and neighboring synapses (Brp+, cyan) in (**a-a"**)

luciferase[RNAi] (luc) control brains at 8 h ALH, or (**b-e"**) 8 h ALH brains following astrocyte-

specific knockdown of molecular machinery required for mitochondrial transport in axons.

Prime panels show astrocyte membranes and mitochondria alone. Double prime panels show

mitochondria alone. Scale bar, 5 µm for all panels.

(**f**) Quantification of the percent of all synapses within 1 µm of astrocyte mitochondria.

Statistical significance compared to control determined by one-way ANOVA: *milton*[RNAi 1]

(p<.0001), *milton*[RNAi 2] (p<.0001), *milton*[RNAi 3] (p<.002), *khc*[RNAi 1] (p<.04), *khc*[RNAi 2] (p<.007),

miro[RNAi] (p<.0001), *Dhc64C*[RNAi 1] (p<.14), *Dhc64C*[RNAi 2] (p<.43).

(**g**) Quantification of the percent of all synapses within 1 µm of astrocyte membranes.

Statistical significance compared to control determined by one-way ANOVA: *milton*[RNAi 1]

($p<.84$), *milton*[RNAi 2] ($p<.85$), *milton*[RNAi 3] ($p<.24$), *khc*[RNAi 1] ($p<.91$), *khc*[RNAi 2] ($p<.97$),

miro[RNAi] ($p<.0001$), *Dhc64C*[RNAi 1] ($p<.37$), *Dhc64C*[RNAi 2] ($p<.88$).

(**f-g**) Control: n=9. *Milton*[RNAi 1]: n=6. *Milton*[RNAi 2]: n=5. *Milton*[RNAi 3]: n=6. *Khc*[RNAi 1]: n=5.

Khc[RNAi 2]: n=5 animals. *Miro*[RNAi]: n=5 animals. *Dhc64C*[RNAi 1]: n=5 animals. *Dhc64C*[RNAi 1]:

n=4 animals. Error bars = SEM. N values reflect animals from 2 technical replicates.

(**h**) Graphical summary of screen results.

Figure 7. Astrocyte mitochondrial trafficking modifies larval crawling behavior.

(**a**) Schematic of behavioral paradigms tested.

(**b-e**) FIMTrack traces from individual, free-crawling larvae recorded for 1 minute at 4 Hz.

Luciferase[RNAi] (luc) Control: n=222 animals. *Tbph*[RNAi]: n=126 animals. *Milton*[RNAi 1]: n=139

animals. *Milton*$^{RNAi\ 2}$: n=85 animals. *Tbph*^{RNAi} + *milton*$^{RNAi\ 1}$: n=79 animals. *Tbph*^{RNAi} +

milton$^{RNAi\ 2}$: n=82 animals. N values reflect animals from 3 technical replicates.

(**f-j**) Quantification of behavioral metrics relative to control. Statistical significance

determined by Student's T-Test for (**f**) velocity (*tbph*^{RNAi}, p<.57; *milton*$^{RNAi\ 1}$, p<.0001;

milton$^{RNAi\ 2}$, p<.4; *tbph*^{RNAi} + *milton*$^{RNAi\ 1}$, p<.0001; *tbph*^{RNAi} + *milton*$^{RNAi\ 2}$, p<.14), (**g**)

distance to origin (*tbph*^{RNAi}, p<.26; *milton*$^{RNAi\ 1}$, p<.003; *milton*$^{RNAi\ 2}$, p<.21; *tbph*^{RNAi} +

milton$^{RNAi\ 1}$, p<.002; *tbph*^{RNAi} + *milton*$^{RNAi\ 2}$, p<.03), (**h**) accumulated distance (*tbph*^{RNAi},

p<.26; *milton*$^{RNAi\ 1}$, p<.0001; *milton*$^{RNAi\ 2}$, p<.03; *tbph*^{RNAi} + *milton*$^{RNAi\ 1}$, p<.0001; *tbph*^{RNAi} +

milton$^{RNAi\ 2}$, p<.06), (**i**) cumulative bending angle (*tbph*^{RNAi}, p<.68; *milton*$^{RNAi\ 1}$, p<.03;

milton$^{RNAi\ 2}$, p<.05; *tbph*^{RNAi} + *milton*$^{RNAi\ 1}$, p<.05; *tbph*^{RNAi} + *milton*$^{RNAi\ 2}$, p=.06), and (**j**)

crawling (*tbph*^{RNAi}, p<.33; *milton*$^{RNAi\ 1}$, p<.0001; *milton*$^{RNAi\ 2}$, p=.05; *tbph*^{RNAi} + *milton*$^{RNAi\ 1}$,

p<.02; *tbph*^{RNAi} + *milton*$^{RNAi\ 2}$, p<.22). Error bars = SEM.

Figure 8. Peri-synaptic astrocyte mitochondria are necessary for synapse maintenance.

(**a-d'**) Representative images of 96 h ALH neuromuscular junctions (NMJs, Green: *vglut-lexA, lexAop-myr::GFP*; Magenta: Dlg+; White: Phalloidin) from luciferase[RNAi] (luc) control (**a**) or following astrocyte-specific knockdown of (**b**) *tbph*, (**c**) *milton*, or (**d**) both *tbph* and *milton* together (*R25H07-gal4, UAS-RNAi*). Prime panels show GFP channel used for quantification of synaptic boutons (arrowheads). Control: n=98 NMJs from n=16 animals. *Tbph*[RNAi]: n=81 NMJs from n=14 animals. *Milton*[RNAi 1]: n=61 NMJs from n=12 animals. *Milton*[RNAi 2]: n=25 NMJs from n=4 animals. *Tbph*[RNAi] + *milton*[RNAi 1]: n=64 NMJs from n=12 animals. *Tbph*[RNAi] + *milton*[RNAi 2]: n=26 NMJs from n=4 animals. N values reflect animals from 2 technical replicates.

964   (**e**) Schematic for reader orientation.

965   (**f**) Quantification of bouton number (type 1b only). Statistical significance determined by

966   Student's T-Test relative to control: *tbph*^RNAi^ (p<.19), *milton*^RNAi 1^ (p<.02), *milton*^RNAi 2^

967   (p<.03), *tbph*^RNAi^ + *milton*^RNAi 1^ (p<.002), *tbph*^RNAi^ + *milton*^RNAi 2^ (p<.04).

SUPPLEMENTAL MOVIE LEGENDS

Supplementary Movies 1-11: All movies are .mov files and can be visualized using standard

media players.

Supplementary Movie 1. Time-lapse imaging of astrocyte mitochondria in myr::TdTomato

control for Chrimson-activation.

Representative time-lapse movie showing astrocyte mitochondrial dynamics (*R25H07-gal4,*

UAS-mito::GFP, green) around aCC/RP2 dendrites (*RN2-lexA, lexAop-myr::TdTomato*) in a

fictive brain preparation at 8 h ALH. Animals were supplied with ATR and reared in the dark to

mimic conditions for Chrimson-activation. A single z-stack was acquired every 30 seconds (each

stack taking 25" to acquire) and brains were imaged for a total of 15 minutes. Astrocyte

mitochondria were highly motile, though largely stable in volume over the 15-minute period. Left

hemisegment shows Imaris "Surface" reconstruction of astrocyte mitochondria, right shows raw

data in neighboring hemisegment.

Supplementary Movie 2. Time-lapse imaging of astrocyte mitochondria during motor

neuron Chrimson-activation.

Representative time-lapse movie showing astrocyte mitochondrial dynamics (*R25H07-gal4,*

UAS-mito::GFP, green) around aCC/RP2 dendrites (*RN2-lexA, lexAop-Chrimson::TdTomato*) in

a fictive brain preparation at 8 h ALH. Animals were supplied with ATR and dark-reared. A

single z-stack was acquired every 30 seconds (each stack taking 25" to acquire) and brains were

imaged for a total of 15 minutes. Astrocyte mitochondria were rapidly trafficked towards motor

dendrites in the first several minutes, followed by relative stability. Left hemisegment shows

Imaris "Surface" reconstruction of astrocyte mitochondria, right shows raw data in neighboring

hemisegment.

Supplementary Movie 3. Time-lapse imaging of astrocyte mitochondria in myr::GFP

control for GtACR2 silencing.

Representative time-lapse movie showing astrocyte mitochondrial dynamics (*alrm-lexA, lexAop-*

mcherry::mito.OMM, magenta) around aCC/RP2 dendrites (*RN2-gal4, UAS-GtACR2::EYFP*) in a

fictive brain preparation at 8 h ALH. Animals were supplied with ATR and reared in the dark to

mimic conditions for GtACR2-silencing. A single z-stack was acquired every 30 seconds (each

stack taking 25" to acquire) and brains were imaged for a total of 15 minutes. Astrocyte

mitochondria were highly motile, though largely stable in volume over the 15-minute period. Left

hemisegment shows Imaris "Surface" reconstruction of astrocyte mitochondria, right shows raw

data in neighboring hemisegment.

Supplementary Movie 4. Time-lapse imaging of astrocyte mitochondria during motor

neuron GtACR2-silencing.

Representative time-lapse movie showing astrocyte mitochondrial dynamics (*alrm-lexA, lexAop-*

mcherry::mito.OMM, magenta) around aCC/RP2 dendrites (*RN2-gal4, UAS-GtACR2::EYFP*) in a

fictive brain preparation at 8 h ALH. Animals were supplied with ATR and reared in the dark. A

single z-stack was acquired every 30 seconds (each stack taking 25" to acquire) and brains were

imaged for a total of 15 minutes. Astrocyte mitochondria were highly motile, though largely

1017 stable in volume over the 15-minute period. Left hemisegment shows Imaris "Surface"

1018 reconstruction of astrocyte mitochondria, right shows raw data in neighboring hemisegment.

1019

1020 **Supplementary Movie 5. Time-lapse imaging of wild-type aCC/RP2 activity.**

1021 Representative time-lapse movie showing control aCC/RP2 activity in a fictive brain preparation

1022 at 8 h ALH (*RN2-lexA, lexAop-GCaMP7s* ; *R25H07-gal4, UAS-luciferase*^RNAi^). A single z-stack

1023 was acquired every 15 seconds (each stack taking 15" to acquire) and brains were imaged for a

1024 total of 15 minutes.

1025

Supplementary Movie 6. Time-lapse imaging of aCC/RP2 activity following astrocyte

knockdown of *tbph*.

Representative time-lapse movie showing aCC/RP2 activity in a fictive brain preparation at 8 h

ALH (*RN2-lexA, lexAop-GCaMP7s*) following astrocyte knockdown of *tbph* (*R25H07-gal4,*

UAS-tbph^RNAi^). A single z-stack was acquired every 15 seconds (each stack taking 15" to acquire)

and brains were imaged for a total of 15 minutes. Activity was elevated relative to control.

1032

Supplementary Movie 7. Time lapse imaging showing control astrocyte mitochondrial

dynamics.

Representative time-lapse movie showing astrocyte mitochondria (*R25H07-gal4, UAS-*

mito::GFP) dynamics in a control brain (*UAS-luciferase*^RNAi^) at 8 h ALH. A single z-stack was

acquired one a minute (each stack taking 15" to acquire) and brains were imaged for a total of 10

minutes. Mitochondria were distributed throughout the neuropil.

1039

Supplementary Movie 8. Time lapse imaging showing astrocyte mitochondrial dynamics following astrocyte knockdown of *milton*.

Representative time-lapse movie showing astrocyte mitochondria (*R25H07-gal4, UAS-mito::GFP*) dynamics in a *milton* knockdown brain (*UAS-milton*[RNAi]) at 8 h ALH. A single z-stack was acquired one a minute (each stack taking 15" to acquire) and brains were imaged for a total of 10 minutes. Mitochondria were absent from the neuropil.

Supplementary Movie 9. Time lapse imaging showing astrocyte mitochondrial dynamics following astrocyte knockdown of *khc*.

Representative time-lapse movie showing astrocyte mitochondria (*R25H07-gal4, UAS-mito::GFP*) dynamics in a *khc* knockdown brain (*UAS-khc*[RNAi]) at 8 h ALH. A single z-stack was acquired one a minute (each stack taking 15" to acquire) and brains were imaged for a total of 10 minutes. Mitochondria were reduced in the neuropil.

Supplementary Movie 10. Time lapse imaging showing astrocyte mitochondrial dynamics following astrocyte knockdown of *miro*.

Representative time-lapse movie showing astrocyte mitochondria (*R25H07-gal4, UAS-mito::GFP*) dynamics in a *miro* knockdown brain (*UAS-miro*[RNAi]) at 8 h ALH. A single z-stack was acquired one a minute (each stack taking 15" to acquire) and brains were imaged for a total of 10 minutes. Mitochondria were reduced in the neuropil.

Supplementary Movie 11. Time lapse imaging showing astrocyte mitochondrial dynamics following astrocyte knockdown of *Dhc64C*.

1063   Representative time-lapse movie showing astrocyte mitochondria (*R25H07-gal4, UAS-*

1064   *mito::GFP*) dynamics in a *Dhc64C* knockdown brain (*UAS-Dhc64C*[RNAi]) at 8 h ALH. A single z-

1065   stack was acquired one a minute (each stack taking 15" to acquire) and brains were imaged for a

1066   total of 10 minutes. Mitochondria distributed normally in the neuropil.