Title:

Mapping consistent, reproducible, and transcriptionally relevant functional connectome hubs of the human brain

Authors:

Zhilei Xu[1,2,3], Mingrui Xia[1,2,3], Xindi Wang[1,2,3], Xuhong Liao[4], Tengda Zhao[1,2,3], Yong He[1,2,3,5] *

[1]State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal University, Beijing, China

[2]Beijing Key Laboratory of Brain Imaging and Connectomics, Beijing Normal University, Beijing, China

[3]IDG/McGovern Institute for Brain Research, Beijing Normal University, Beijing, China

[4]School of Systems Science, Beijing Normal University, Beijing, China

[5]Chinese Institute for Brain Research, Beijing, China

***Corresponding author:**

Yong He, Ph.D., E-mail: yong.he@bnu.edu.cn

Manuscript information: 30 text pages, 7 figures, 1 table.

(Supplementary Information: 20 text pages, 9 figures, 10 tables.)

Abstract

Human brain connectomes include sets of densely connected regions, known as connectome hubs, which play a vital role in understanding global brain communication, cognitive processing, and brain disorders. However, the consistency and reproducibility of functional connectome hubs' anatomical localization have not been established to date and the genetic signatures underlying robust connectome hubs remain unknown. Here, we conduct the first worldwide, harmonized meta-connectomic analysis by pooling resting-state functional MRI data of 5,212 healthy young adults across 61 independent cohorts. We identify highly consistent and reproducible functional connectome hubs in heteromodal and unimodal regions both across cohorts and across individuals. These connectome hubs show heterogeneous connectivity profiles and are critical for both intra- and inter-network communications. Using post-mortem gene expression data, we show that these connectome hubs have a spatiotemporally distinctive transcriptomic pattern dominated by genes involved in the neuropeptide signaling pathway, neurodevelopmental processes, and metabolic processes. These results highlight the robustness of macroscopic connectome hubs and their potential cellular and molecular underpinnings.

Introduction

Functional connectome mapping studies have identified sets of densely connected regions in large-scale human brain networks, which are known as hubs[1]. Connectome hubs play a crucial role in global brain communication[1, 2] and support a broad range of cognitive processing, such as working memory[3] and semantic processing[4]. Growing evidence suggests that these highly connected brain hubs are preferentially targeted by many neuropsychiatric disorders[5-8], which provides critical clues for understanding the biological mechanisms of disorders and establishing biomarkers for disease diagnosis[8, 9] and treatment evaluation[10] ([1, 2, 11, 12] for reviews).

Despite such importance, there is considerable inconsistency in anatomical locations of functional connectome hubs among existing studies. For example, components of the default-mode network (DMN) have been frequently reported as connectome hubs, yet the spatial pattern is highly variable across studies. In particular, several studies have shown highly connected hubs in the lateral parietal regions of the DMN[7, 8, 13, 14], whereas others have reported midline structures of the DMN[15-19]. Several works have identified primary sensorimotor and visual regions as connectome hubs[13, 14, 16-19], yet others did not replicate these findings[7, 8, 15]. Subcortical regions, such as the thalamus and amygdala, have also been inconsistently reported as hubs[8, 15, 16, 18] and non-hubs[7, 13, 14, 17, 19]. Thus, the consistency and reproducibility of functional connectome hubs have been difficult to establish to date, which can be attributed to inadequate sample size and differences in imaging scanner, imaging protocol, data processing, and connectome analysis strategies. Here, we aimed to establish a harmonized meta-analysis model to identify robust functional connectome hubs in healthy young adults by combining multiple cohorts with uniform protocols for data quality assurance, image processing, and connectome analyses.

Once the robust connectome hubs are identified, we will further examine their genetic signatures. It has been well demonstrated that the connectome architecture of the human brain is inheritable, such as functional connectivity of the DMN[20] and the cost-efficiency optimization[21]. Moreover, the functional connectomes can be regulated by genotypic variation both during rest[22] and in cognitive tasks[23], especially involving the DMN[22, 23] and frontoparietal network (FPN)[23]. Growing evidence also suggests spatial correspondence between transcriptomic profiles and connectome architectures[24-26] ([27] for review). Thus, we reasoned that the robust macroscopic connectome hubs could be associated with microscopic genetic signatures. Elucidating these genetic signatures will substantially benefit our understanding of how connectome hubs emerge in development, function in complex cognition, and are involved in disease.

To address these issues, we provide the first worldwide harmonized meta-connectomic analysis of functional brain hubs by pooling a large-sample resting-state functional MRI (rsfMRI) dataset of 5,212 healthy young adults (aged 18–36 years) across 61 cohorts. Fig 1 illustrates the sample size and age ranges of each cohort. To uncover the genetic signatures underlying these connectome hubs, we conducted machine learning approaches to distinguish connectome hubs from non-hubs using transcriptomic data from the Allen Human Brain Atlas (AHBA) and explored their developmental evolutions using the BrainSpan Atlas.

Results

Identifying consistent connectome hubs using a harmonized meta-analysis model.

Prior to the meta-analysis, we constructed a voxelwise functional connectome matrix for each individual by computing the Pearson's correlation coefficient between preprocessed rsfMRI time series of all pairs of gray matter voxels (47,619 voxels). Then, the functional connectivity strength (FCS) of each voxel was computed as the sum of connection weights between the given voxel and all the other voxels. This resultant FCS map was further normalized with respect to its mean and standard deviation across voxels[7]. For each cohort, we performed a general linear model on these normalized FCS maps to reduce age and gender effects. As a result, we obtained a mean FCS map and its corresponding variance map for each cohort that were used for subsequent meta-analyses.

To identify the most consistent connectome hubs, we conducted a voxelwise random-effects meta-analysis on the mean and variance FCS maps of the 61 cohorts. Such an analysis addressed the across-cohort heterogeneity of functional connectomes, resulting in a robust FCS pattern and its corresponding standard error (SE) map (Fig 2A). Then, we identified consistent connectome hubs whose FCS values were significantly ($p < 0.001$, cluster size > 200 mm^3) higher than the global mean (i.e., zero) using a voxelwise Z value map computed by dividing the FCS map by the SE map. To determine the significance levels of these observed Z values, a nonparametric permutation test[28] with 10,000 iterations was performed. Finally, we estimated voxelwise effect sizes using Cohen's d metric computed by dividing the Z value map by the square root of the cohort number (Fig 2B, left). According to prior brain network parcellations[29, 30], these identified hub voxels (15,461 voxels) were spatially distributed in multiple brain networks, including the DMN (27.5%), dorsal attention network (DAN) (16.5%), FPN (15.9%), ventral attention network (VAN) (15.6%), somatomotor network (SMN) (14.4%), and visual network (VIS) (9.9%) (Fig 2B, right). Using a local maxima localization procedure, we identified 35 robust brain hubs across 61 cohorts (Fig 2B, left; Table 1), involving various heteromodal and unimodal areas. Specifically, the most robust findings resided in several lateral parietal regions, including the bilateral ventral postcentral gyrus, supramarginal gyrus, and angular gyrus.

The identified connectome hubs are reproducible across cohorts and individuals.

During identifying the above highly consistent connectome hubs, the random-effects meta-analysis revealed high heterogeneity of FCS across cohorts (Fig 2C, left). The cumulative distribution function plot shows more than 95% voxels with I^2 (heterogeneity score) exceeding 50% (Fig 2C, right), indicating high heterogeneity across cohorts in almost all brain areas (see also Fig S1). To determine whether the connectome hubs identified here are dominated by certain cohorts or are reproduced across-subject/cohort, we performed a leave-one-cohort-out validation analysis and an across-subject/cohort conjunction analysis.

Leave-one-cohort-out validation analysis. We repeated the above harmonized meta-analysis hub identification procedure after leaving one cohort out at a time. Comparing the identified hubs using all cohorts (Fig 2B) with those after leaving one cohort out obtained extremely high Dice's coefficients (*mean±sd*: 0.990±0.006; range: 0.966-0.997). For hub peaks, leaving one cohort out

resulted in very few displacements (mostly fewer than 6 mm, Fig 2D). Thus, connectome hubs
identified using the 61 cohorts were not dominated by specific cohorts.

Across-subject/cohort conjunction analysis. We defined the top N (N = 15,461, which is the
voxel number of hubs in Fig 2B) voxels with the highest FCS values of a subject or a cohort as
connectome hubs for that subject or that cohort. Then, for each voxel, we assessed hub
occurrence probability values across subjects and cohorts. The identified hubs using all cohorts
were highly overlapped with the top N voxels with the highest hub occurrence probability values
both across all subjects and across all cohorts, indicated by a high Dice's coefficient (*Dice* =
0.867, Fig 2E, left; *Dice* = 0.924, Fig 2E, right). When the identified hubs using all cohorts were
compared with the top N voxels with the highest hub occurrence probability values across
randomly selected subjects or across randomly selected cohorts, the Dice's coefficient
approached 99% of its maximum value after exceeding 510 subjects (Fig 2F, left) and 35 cohorts
(Fig 2F, right), respectively. This indicated that the identified connectome hubs were highly
reproducible both across cohorts and across individuals.

Validation analysis demonstrated that the above results did not depend on analysis parameters,
such as the connection threshold (Fig S2 and S3), and were not driven by the size of the brain
network to which they belong[31] (Fig S4), suggesting the robustness of our main findings.

Connectome hubs have heterogeneous functional connectivity profiles.

Next, we further examined whether these robust brain hubs (Fig 2B and Table 1) have distinctive
functional connectivity profiles that represent their unique roles in network communication. To
gain detailed and robust functional connectivity profiles of each hub region, we conducted a
seed-to-whole-brain connectivity meta-analysis in a harmonized protocol again. For each of the
35 hub regions, we obtained an estimated Cohen's d effect size map that characterizes the robust
whole-brain connectivity pattern relevant to the seed region across the 61 cohorts (Fig 3A). We
then divided the connectivity map of each hub into eight brain networks according to prior
parcellations[29, 30], resulting in an 8×35 connectivity matrix with each column representing the
voxel percentage of each of the eight networks connected with a hub.

Hierarchical clustering analysis on the connectivity matrix clearly divided the 35 hubs into three
clusters (Fig 3B). Cluster I consists of 21 hubs that are primarily connected with extensive areas
in the DAN, VAN, FPN, and SMN (orange, Fig 3B). Cluster II consists of four hubs that are
densely connected with VIS (green, Fig 3B). Cluster III consists of 10 hubs that have robust
connections with the DMN and LIMB (blue, Fig 3B). Of particular interest is that within Cluster
III, a left posterior middle frontal hub called ventral area 8A (8Av) shows a distinctive
connectivity profile in contrast to the other nine hubs, manifested as having robust connections
with bilateral frontal FPN regions (Fig 3A and Fig S5). This imply that the left 8Av hub is a key
connector between the DMN and FPN, which can be supported by the recent finding of a
control-default connector located in the posterior middle frontal gyrus[32]. Although both Cluster I
and III hubs are connected with subcortical structure, they are connected with different
subcortical nuclei (Fig 3B and Fig S6). Finally, whereas all hubs possess dense intranetwork
connections, most also retain significant internetwork connections (Fig S7), which preserves
efficient communication across the whole brain network feasible.

Transcriptomic data distinguishes connectome hubs from non-hubs.

A supervised machine learning classifier based on XGBoost[33] and 10,027 genes' transcriptomic data from the AHBA[34] was trained to distinguish connectome hubs from non-hubs (Fig 4A). The sensitivity, specificity, and accuracy rate of the XGBoost classifier were stably estimated by repeating the training and testing procedure 1,000 times. This classifier performed better than chance in all 1,000 repetitions and achieved an overall accuracy rate of 65.3% (Fig 4B). In cross-validation, connectome hubs and non-hubs were classified with a sensitivity of 71.1% and specificity of 63.4%, respectively. The testing procedure yielded a comparable sensitivity of 69.7% and specificity of 62.0%. After training the classifier, each gene's contribution to the optimal prediction model was determined. We noted that some key genes contributed two or three orders of magnitude more than other genes (Fig 4C). The contributions of the top 300 mostly contributed key genes were consistent between the first 500 repetitions and the second 500 repetitions (Pearson's $r = 0.958$, $p < 10^{-6}$, Fig 4D), suggesting a high reproducibility.

To exclude the XGBoost model's potential bias relating to the mostly contributed key genes, we replicated the above classification results using another machine learning model based on the support vector machine (SVM) that was trained using only the top N key genes with the greatest contributions to the XGBoost classifier (Fig 4E). Because no data were available to determine how many key genes were sufficient to train an SVM classifier, we examined the count N from 100 to 300. The SVM classifier achieved a very high peak accuracy rate of 91.8% with approximately the top 150 key genes in the easiest classification task (Fig 4F) and also achieved a reasonable peak accuracy rate of 67.8% with approximately the top 150 key genes even in the most difficult classification task (Fig 4G). By contrast, SVM classifiers trained using 150 randomly selected genes performed worse than that using the top 150 key genes in all 1,000 repetitions (Fig 4H). Thus, these robust connectome hubs were significantly associated with a transcriptomic pattern dominated by approximately 150 key genes (Table S1).

Connectome hubs have a spatiotemporally distinctive transcriptomic pattern.

Gene Ontology (GO) enrichment analysis using GOrilla[35] demonstrated that the above 150 key genes were mostly enriched in the neuropeptide signaling pathway (*fold enrichment* (*FE*) = 8.9, uncorrected $p = 1.2{\times}10^{-5}$, Table S2). GO enrichment analysis using the ranked 10,027 genes according to their contributions to the XGBoost classifier also confirmed the most enriched GO term of the neuropeptide signaling pathway (*FE* = 5.7, uncorrected $p < 10^{-6}$, Table S3). The ranked 10,027 genes were also associated with the developmental process (*FE* = 1.2), cellular developmental process (*FE* = 1.3), anatomical structure development (*FE* = 1.3), and neuron projection arborization (*FE* = 13.7) (uncorrected $ps < 5.5{\times}10^{-4}$, Table S3). We speculated that connectome hubs have a distinctive transcriptomic pattern of neurodevelopmental processes in contrast to non-hubs.

We repeated the GO enrichment analysis of the above 150 key genes using DAVID[36, 37]. In addition to the mostly enriched GO term of the neuropeptide signaling pathway (*FE* = 8.7, uncorrected $p = 5.8{\times}10^{-4}$), there were 10 GO terms associated with metabolic process, such as the positive regulation of cellular metabolic process (*FE* = 1.4, uncorrected $p = 0.031$, Table S4). Disease association analysis demonstrated metabolic disease associated with the greatest number of key genes (60 genes, *FE* = 1.2, uncorrected $p = 0.094$, Table S5). Accordingly, it is rational to

speculate that connectome hubs have a distinctive transcriptomic pattern of metabolic processes in contrast to non-hubs.

To confirm the above two speculations of GO enrichment analysis results, we examined transcription level differences between hub and non-hub regions for genes previously implicated in key neurodevelopmental processes[38] (Table S6) and main neuronal metabolic pathways[39] (oxidative phosphorylation[40] and aerobic glycolysis[41], Table S7). Permutation tests revealed hub regions with significantly higher transcription levels for genes associated with dendrite development, synapse development, and aerobic glycolysis than non-hub regions (one-sided Wilcoxon rank-sum tests, Bonferroni-corrected $ps \leq 0.032$, Fig 5A). In addition, hub regions had a weak trend of lower transcription levels for genes associated with axon development, myelination, neuron migration, and oxidative phosphorylation (Fig 5A). These transcription level differences were consistent with our speculations of GO enrichment analysis results.

These above transcriptomic results were derived from the AHBA, an adult transcriptomic dataset. To explore their developmental evolutions, we inspected transcriptomic trajectory differences between hub and non-hub regions using the BrainSpan Atlas[42]. We observed pronounced diverging transcriptomic trajectories between hub and non-hub regions in these key neurodevelopmental processes and main neuronal metabolic pathways (Fig 5B and Fig S8). For neuron migration, the transcription level in hub regions is higher than that in non-hub regions from the mid-fetal period to early infancy. For dendrite, synapse, axon development, and myelination, transcriptomic trajectories of hub regions apparently diverge from those of non-hubs during childhood and adolescence, during which hub regions have higher transcription levels for dendrite and synapse development but lower transcription levels for axon development and myelination. These results are in agreement with the observation of primary somatosensory, auditory, and visual (V1/V2) cortices with lower synapse density but higher myelination than the prefrontal area[43, 44]. Mmoreover, hub regions have higher transcription levels than non-hub regions for aerobic glycolysis since the early childhood period and for oxidative phosphorylation during childhood and adolescence. These transcriptome analyses achieved convergent results between the AHBA and BrainSpan Atlas.

Together, functional connectome hubs have a spatiotemporally distinctive transcriptomic pattern in contrast to non-hubs, which is dominated by genes involved in the neuropeptide signaling pathway, neurodevelopmental processes, and metabolic processes.

Connectome hubs have more intricate fiber configuration and higher metabolic rate.

Growing evidence has suggested a striking spatial correspondence between transcriptomic profile and structural connectivity in the human brain[27]. We speculated that the above microscale transcriptomic differences between hub and non-hub regions in key neurodevelopmental processes may result in macroscale structural connectivity profile differences. Using a fiber length profiling dataset[45], we observed that hub regions possess more fibers with a length exceeding 40 mm but less fibers with a length shoter than 40 mm (one-sided Wilcoxon rank-sum tests, Bonferroni-corrected $ps \leq 0.008$, Fig 6). That is, hub regions have more short, medium, and long fibers, whereas non-hub regions have more very short (< 40 mm) fibers, suggesting a more intricate fiber configuration in hub regions.

The above transcriptome analyses have shown a higher transcription level of oxidative
phosphorylation and aerobic glycolysis in hub regions than in non-hubs. We validated these
observations using a metabolism dataset derived from positron emission tomography[46] and found
that hub regions not only have a higher metabolic rate than non-hubs in oxidative
phosphorylation (indicated by the cerebral metabolic rate for oxygen) and aerobic glycolysis
(indicated by the glycolytic index), but also have more blood supply (indicated by the cerebral
blood flow) (one-sided Wilcoxon rank-sum tests, Bonferroni-corrected $ps < 0.001$, Fig 7). This is
in agreement with prior observations of a tight coupling between FCS and blood supply[1, 47].

Discussion

Using a worldwide harmonized meta-connectomic analysis of 5,212 healthy young adults across
61 cohorts, we provided the first description of highly consistent and reproducible functional
connectome hubs in the resting human brain. Using transcriptomic data from the AHBA and
BrainSpan Atlas, we reported that these robust connectome hubs have a spatiotemporally
distinctive transcriptomic pattern in contrast to non-hub regions. These results advanced our
understanding of the robustness of macroscopic functional connectome hubs and their potential
cellular and molecular substrates.

Extant reports have shown largely inconsistent and less reproducible hub localizations[7, 8, 13-19],
which may arise from high heterogeneity in the included subjects, data acquisition, and analysis
strategies across studies. To diminish these potential confounding factors, we employed stringent
participant inclusion criteria that included only healthy young adults aged 18 to 36 years and
adopted harmonized data preprocessing and connectome analysis protocols across cohorts.
Nevertheless, the random-effects meta-analysis revealed high heterogeneity among cohorts in
almost all brain areas, which implied that heterogeneity of imaging scanners and/or imaging
protocols could be an important cause for inconsistent and less reproducible results across prior
studies. Thus, our study was indispensable by conducting a harmonized random-effects meta-
analysis model in which both intracohort variation (i.e., sampling errors) and intercohort
heterogeneity were considered[48]. In addition, our validation results showed that the spatial
distribution of functional connectome hubs was relatively stable when using more than 510
subjects and 35 cohorts, demonstrating that 5,212 subjects from 61 cohorts were adequate to
minimize both sampling errors and heterogeneity among cohorts. Considering only dozens of
subjects in most prior studies[7, 8, 13-15, 17, 19], the low statistical power attributed to inadequate
subjects could be another cause for prior inconsistent and less reproducible hub localizations.
Finally, we used harmonized image processing and connectome analysis protocols across cohort,
which avoided methodological variation and reduced potential methodological defects that have
not been resolved in prior studies. See an extension discussion in Supplementary Text II.

The present results demonstrated that the 35 highly consistent and reproducible connectome hubs
show heterogeneous functional connectivity profiles, forming three clusters. Twenty-one hubs
(Cluster I) are connected with extensive areas in the DAN, VAN, FPN, and SMN. Previous
investigations indicated that they are core regions of the DAN (left AIP, right 7PC, left 7Am,
bilateral PFt, left FEF, bilateral 6a, right 6v, and right FST)[29, 49], VAN (left 43, left FOP4, right
46, right 6r, right PF, left PFop, left SCEF, right 5mv)[29, 49], and FPN (left p9-46v and right

IFSa)[29, 50]. In addition, hub regions involved in the sensorimotor pathway (right VIP, right FST,
left 7Am, and left FEF)[51] are also connected with the visual association cortex, acting as
connectors between the VIS and the SMN, DAN, and VAN. Information flow along the primary
visual, visual association, and higher-level sensorimotor cortices is undertaken by the four
occipital hubs (Cluster II) left VMV1, right V4, and bilateral V3A that are all densely connected
with the VIS and portions of the SMN, DAN, and VAN. This aligns with the role of their
homologous regions in the non-human primate cerebral cortex[51]. The remaining 10 hubs (Cluster
III) are all located in canonical DMN regions[52]. One of them, the left 8Av hub, is robustly
connected with both DMN and lateral prefrontal FPN regions, acting as a connector between the
DMN and FPN. This can be supported by the recent finding of a control-default connector
located in the posterior middle frontal gyrus[32] and may also be a case of the hypothesis of
parallel interdigitated subnetworks[53]. This observation offers a significant complementary
interpretation to the conventional assumption that the DMN is anticorrelated with other
networks[52]. Considering that communication between the DMN and other networks is of
significant relevance to neuropsychiatric disorders[54], such as autism spectrum disorders[55], we
speculated that the left 8Av hub may be a promising target region for therapeutic interventions.

We demonstrated that these robust brain hubs have a spatiotemporally distinctive transcriptomic
pattern dominated by genes with the highest enrichment for the neuropeptide signaling pathway.
Because neuropeptides are a main type of synaptic transmitter that is widely distributed in the
human central nervous system[56], robust neuropeptide signaling pathways are indispensable for
efficient synaptic signal transduction that sustains dense and flexible functional connections of
hub regions. In addition, hub regions have higher transcription levels for main neuronal
metabolic pathways in contrast to non-hubs. This is reasonable because massive synaptic
activities in hub regions demand high material and metabolic costs, which is in accordance with
our observation of more blood supply and higher oxidative phosphorylation and aerobic
glycolysis levels in hub regions. This is also in agreement with prior observations of a tight
coupling between FCS and blood supply[1, 47].

We found that connectome hubs possess a spatiotemporally distinctive transcriptomic pattern of
key neurodevelopmental processes in contrast to non-hubs. Specifically, connectome hubs have
higher transcription levels for dendrite and synapse development and lower transcription levels
for axon development and myelination during childhood, adolescence, and adulthood. These
findings are compatible with previous observations of the prefrontal area having higher synapse
density but lower myelination than primary somatosensory, auditory, and visual (V1/V2)
cortices[43, 44]. Higher transcription levels for dendrite and synapse development in hub regions are
necessary for the overproduction of synapses that will be selectively eliminated based on the
demand of the environment and gradually stabilized before full maturation[57], which is "the major
mechanism for generating diversity of neuronal connections beyond their genetic
determination"[58]. Lower transcription levels for axon development and myelination will prolong
the myelination period in hub regions, which characterizes a delayed maturation phase[59]. Marked
delay of anatomical maturation in human prefrontal and lateral parietal cortices has been
frequently observed both in human development[58, 60] and in primate evolution[59], which provides
more opportunities for social learning to establish diverse neuronal circuits that contribute to our
complex[58] and species-specific[59] cognitive capabilities. We also observed higher transcription
levels for neuron migration in hub regions from mid-fetal period to early infancy. This is in
agreement with the report of extensive migration of young neurons persisting for several months

after birth in the human frontal cortex[61]. Meanwhile, the migration and final laminar positioning
of postmitotic neurons are regulated by common transcription factors[62], which suggests that a
higher transcription level for neuron migration in hub regions facilitates the construction of more
intricate interlaminar connectivity. These microscale divergences of key neurodevelopmental
processes may result in a more intricate macroscale structural connectivity proflie in hub regions.

Human neurodevelopment is an intricate and protracted process, during which the transcriptome
of the human brain requires precise spatiotemporal regulation[38]. Thus, in addition to contributing
to our complex cognitive capabilities, the distinctive transcriptomic pattern of neurodevelopment
in hub regions may also increase connectome hubs' susceptibility to neuropsychiatric disorders[58,
59], which means small disturbance in the magnitude or the timing of this transcriptomic pattern
may have long-term consequences on brain anatomical topography or functional activation. This
is in line with our observation of psychiatric disorders being the most significant disease
associated with the top 150 key genes (Table S5). This implies that uncovering the intricate
transcriptomic pattern, diverse neuronal circuits, anatomical topography, and functional
activation of connectome hubs provide crucial and promising routes for understanding the
pathophysiological mechanisms underlying neurodevelopmental disorders, such as autism
spectrum disorders[38, 55] and schizophrenia[5, 38, 58, 59].

Of note, we conducted transcriptome-connectome association analysis using machine learning
approaches in which non-linear mathematical operations were implemented rather than linear
operations, such as linear correlation[24], linear regression[25], or partial least squares[26]. It has been
argued that observations of transcriptome-connectome spatial association have a high false-
positive rate through linear regression[63] and linear correlation[64] and may be largely shifted
toward the first principal component axis of the dataset through partial least squares[65]. These
investigations imply that prior transcriptome-connectome association results by linear
mathematical operations may include high false-positive observations that are independent of
connectome measurements, such as genes enriched for ion channels[24-26]. By contrast, high
reproducibility across different machine learning models and across different GO enrichment
analysis tools and convergent results from the AHBA and BrainSpan Atlas made it very unlikely
that our findings were false-positive observations.

Some results of the present study should be interpreted cautiously because of methodological
issues. First, we identified the robust connectome hubs using preprocessed rsfMRI data with
global signal regression because of its great promise in minimizing physiological artifacts on
functional connectomes[66]. Validation analysis demonstrated that hub distribution identified
without global signal regression was more likely derived from physiological artifacts rather than
by ongoing neuronal activity (Supplementary Text III and Fig S9). Second, the AHBA dataset
only includes partial human genes, of which approximately half were excluded in data
preprocessing[34], which may have induced incomplete observations in our data-driven analysis.
Finally, our transcriptomic signature results addressed only the association between connectome
hubs and transcriptomic patterns and did not explore causation between them. Exploring more
detailed mechanisms underlying this association is attractive and may be practicable for non-
human primate brains in future studies.

Methods

Dataset.

We included a large-sample rsfMRI dataset of 5,212 healthy young adults (aged 18–36 years, 2,377 males) across 61 cohorts from Asia, Europe, North America, and Australia. Data of each cohort were collected with participants' written informed consent and with approval by the respective local institutional review boards. All data passed strict quality controls and were routinely preprocessed with a uniform pipeline. For details, see Supplementary Text I.

Identifying robust functional connectome hubs using a harmonized meta-analysis.

For each individual, we constructed a voxelwise functional connectome matrix by computing the Pearson's correlation coefficient between preprocessed rsfMRI time series of all pairs of voxels within a predefined gray matter mask (47,619 voxels). The gray matter mask was divided into seven large-scale cortical networks[29] and a subcortical network[30]. The cerebellum was not included due to largely incomplete coverage during rsfMRI scanning in most cohorts. Negative functional connections were excluded from our analysis due to neurobiologically ambiguous interpretations[67]. To further reduce signal noise and simultaneously avoid potential sharing signals between nearby voxels, both weak connections (Pearson's $r < 0.1$) and connections terminating within 20 mm were set to zero[68]. We validated the threshold of weak connections using 0.05 and 0.2.

For each voxel, we computed the FCS as the sum of connection weights between the given voxel and all the other voxels. We further normalized this resultant FCS map with respect to its mean and standard deviation across voxels[7]. For each cohort, we performed a general linear model on these normalized FCS maps to reduce age and gender effects. For each voxel, we constructed the general linear model as:

$$FCS_i = \beta_0 + \beta_{Age} * (Age_i - MeanAge) + \beta_{Gender} * Gender_i + \varepsilon_i \tag{1}$$

FCS_i, Age_i, $Gender_i$, and ε_i indicate the FCS, age, gender, and residual of the ith individual, respectively. $MeanAge$ indicates the mean age of that cohort. The general linear model exported a mean FCS map and its corresponding variance map for each cohort.

The mean and variance FCS maps of the 61 cohorts were submitted to a random-effects meta-analysis model[48] to address across-cohort heterogeneity of functional connectomes. The detailed computational procedures are described in the book[48]. A short summary of these procedures was provided in Supplementary Text I. This resulted in a consistent FCS pattern and its corresponding SE map. We compared the FCS of each voxel with the average of the whole brain (i.e., zero) using a Z value[48]:

$$Z = \frac{FCS - 0}{SE} \tag{2}$$

In line with previous neuroimaging meta-analysis study[69], we performed 10,000 one-sided nonparametric permutation tests[28] to assign a p value to the observed Z value. For each iteration, after randomizing the spatial correspondence among cohorts' mean FCS maps (the spatial

correspondence between a cohort's mean FCS map and its variance map was not changed), we
repeated the computation procedure of the random-effects meta-analysis for each voxel and
extracted the maximum Z value of all voxels to construct a null distribution. A p value was
assigned to each voxel by comparing the observed Z value to the null distribution. For a
significance level below 0.05, this p value closely tracks the Bonferroni threshold[28]. Finally, we
defined functional connectome hubs as brain regions with a p value less than 0.001 and cluster
size greater than 200 mm^3. The thresholds of p value and cluster size were similar with the
activation likelihood estimation algorithm[69]. We extracted MNI coordinates for each local peak Z
value terminating beyond 15 mm within each brain cluster using the *wb_command -volume-*
extrema command (https://humanconnectome.org/software/workbench-command/-volume-
extrema) in Connectome Workbench v1.4.2. Effect size was estimated using Cohen's d metric [48]:

$$d = \frac{Z}{\sqrt{k}} \qquad\qquad (3)$$

419 k is the number of cohorts in the meta-analysis.

Mapping seed-to-whole-brain connectivity profiles of functional connectome hubs.

We modeled each hub seed region as a sphere with a 6-mm radius centered on the hub peak and
computed Pearson's correlation coefficients between the seed region's preprocessed rsfMRI time
series and the time series of all gray matter voxels. The time series of the seed region was
computed by averaging the time series of all gray matter voxels in the seed sphere. These
correlation coefficients were further transformed to Fisher's z for normality. In line with above,
we constructed a general linear model on these Fisher's z value maps within each cohort to
reduce age and gender effects and performed a random-effects meta-analysis on these Fisher's z
value maps across cohorts to address the across-cohort heterogeneity, resulting in a robust
Fisher's z pattern and its corresponding SE map. Then, We compared the Fisher's z value of each
voxel with zero using a Z value[48] and estimated effect size using Cohen's d metric[48] as described
in equations (2) and (3). We performed 10,000 one-sided nonparametric permutation tests[28] to
identify the most consistent functional connection Z value map with a p value less than 0.001 and
cluster size greater than 200 mm^3. Finally, we divided the connectivity map of each hub into
eight brain networks mentioned above and represented the functional connectivity profile of a
hub as the voxel percentage of each of the eight networks connected with it to address the effect
of network size. To illustrate the left 8Av hub's connectivity profile, we also mapped its
homologous region the right 8Av region's connectivity profile (Fig S5).

Identifying transcriptomic signatures underlying functional connectome hubs.

We trained classifiers based on XGBoost and SVM to distinguish connectome hubs from non-
hubs using transcriptomic features from the preprocessed AHBA dataset[34]. The top 150 genes
(Table S1) mostly contributed to the classification results were submitted to GO enrichment
analyses using GOrilla[35] (http://cbl-gorilla.cs.technion.ac.il) and DAVID[36, 37] v6.8
(https://david.ncifcrf.gov). For details, see Supplementary Text I. Based on GO enrichment
analysis results, we tested transcription level differences of gene sets involved in key
neurodevelopmental processes[38] (Table S6) and main neuronal metabolic pathways[39] (oxidative
phosphorylation[40] and aerobic glycolysis[41], Table S7) between connectome hubs and non-hubs

through one-sided Wilcoxon rank-sum test. In line with prior studies[38, 41], we used the first
principal component of each gene set's transcription level to plot and to perform the statistical
analysis (Fig 5A). For illustration purposes, we normalized the first principal component of each
gene set's transcription level respect to its minimum and maximum values across all brain
samples to range from 0 to 1.

To explore developmental details, we inspected transcriptomic trajectory differences between
connectome hubs and non-hubs in the above gene sets using the BrainSpan Atlas[42]. In line with
prior studies[38, 41], we used the first principal component of each gene set's transcription level to
plot transcriptomic trajectories and visually inspected transcriptomic trajectory differences
between connectome hubs and non-hubs (Fig 5B). Transcriptomic trajectories were plotted using
locally weighted regression by smoothing the first principal component of each gene set's
transcription level against log2[post-conceptional days] as in prior study[38]. Of note, considering
apparent transcriptomic differences compared to the neocortex[38], we excluded the striatum,
mediodorsal nucleus of the thalamus, and cerebellar cortex in the transcriptomic trajectory
analysis but not the amygdala and hippocampus whose transcriptomic trajectories are more
similar to those of the neocortex than to those of other subcortical structures[38]. Analysis using
only neocortical regions revealed almost unchanged results (Fig S8).

To validate above results derived from transcriptome datasets, we tested fiber number
differences between connectome hubs and non-hubs through one-sided Wilcoxon rank-sum test
(Fig 6). Fiber number data across different length bins was derived from a fiber length profiling
dataset[45]. For each fiber length bin, fiber number of each vertex was normalized with respect to
its mean and standard deviation across vertices. We further examined differences between
connectome hubs and non-hubs in metabolic measurements of blood supply (the cerebral blood
flow), oxidative phosphorylation (the cerebral metabolic rate for oxygen), and aerobic glycolysis
(the glycolytic index) through one-sided Wilcoxon rank-sum test (Fig 7). These measurements
were derived from a positron emission tomography study[46] and assigned to 82 Brodmann areas
and seven subcortical structures. Brodmann areas with more than 50% vertices or subcortical
structures with more than 50% voxels identified as hubs were regarded as hub regions.

Statistical analysis.

We performed statistical analyses using MATLAB R2013a. Statistical significance of brain
clusters in Fig 2B and 3A and Fig S2B, S3B, S5, and S9A were determined by comparing the
observed Z values in equation (2) with their corresponding null distributions constructed by
above mentioned 10,000 one-sided nonparametric permutation tests[28]. To determine the
statistical significance of one-sided Wilcoxon rank-sum tests in Fig 5A, 6, and 7 and Fig S9D,
we constructed 1,000 surrogate hub identification maps with the spatial autocorrelations being
corrected using a generative model[70] and repeated calculating rank-sum statistics using these
surrogate hub identification maps to construct a null distribution. Then, p values derived by
comparing the observed rank-sum statistics with their corresponding null distributions were
Bonferroni-corrected. Surrogate hub identification maps in Fig 5A, 6, and 7 were constructed
based on the hub identification map in Fig 2B. Surrogate hub identification maps in Fig S9D
were constructed based on the hub identification map in Fig S9A.

Data availability

The MRI data of the first 60 cohorts listed in Table S8 are available at the International Neuroimaging Data-sharing Initiative (http://fcon_1000.projects.nitrc.org), Brain Genomics Superstruct Project (https://doi.org/10.7910/DVN/25833), Human Connectome Project (https://www.humanconnectome.org), MPI-Leipzig Mind-Brain-Body Project (https://openneuro.org/datasets/ds000221), and Age-ility Project (https://www.nitrc.org/projects/age-ility). The MRI data of the PKU cohort are under active use by the reporting laboratory and will be available upon reasonable request. The preprocessed AHBA dataset is available at https://doi.org/10.6084/m9.figshare.6852911. The normalized BrainSpan Atlas dataset is available at http://brainspan.org/static/download.html. The fiber length profiling dataset[45] is available at https://balsa.wustl.edu/study/1K3l.

Code availability

The code to reproduce the results and visualizations of this manuscript is available at https://github.com/zhileixu/FunctionalConnectomeHubs.

Author Contributions

Conceptualization: Z.X., Y.H.; Methodology: Z.X., Y.H., M.X., X.W., X.L., T.Z.; Investigation: Z.X.; Visualization: Z.X.; Supervision: Y.H.; Writing—original draft: Z.X., Y.H.; Writing—review & editing: Y.H., Z.X., M.X., X.L., T.Z., X.W.

Acknowledgments

We thank Drs. Huali Wang and Xiaodan Chen for data acquisition of the PKU cohort and Drs. Qiushi Wang and Nan Zhang for valuable advice on MRI data quality controls. This work was supported by the National Natural Science Foundation of China [82021004, 31830034, and 81620108016 to Y.H., 82071998 and 81671767 to M.X., 81971690 to X.L., 81801783 to T.Z.], Changjiang Scholar Professorship Award [T2015027 to Y.H.], National Key Research and Development Project [2018YFA0701402 to Y.H.], Beijing Nova Program [Z191100001119023 to M.X.], and Fundamental Research Funds for the Central Universities [2020NTST29 to M.X.].

Conflict of Interest

The authors declare no competing interests.

References

1. van den Heuvel MP, Sporns O. Network hubs in the human brain. *Trends in Cognitive Sciences* **17**, 683-696 (2013).

2. Bullmore E, Sporns O. The economy of brain network organization. *Nature Reviews Neuroscience* **13**, 336-349 (2012).

3. Liu J*, et al.* Intrinsic Brain Hub Connectivity Underlies Individual Differences in Spatial Working Memory. *Cerebral Cortex* **27**, 5496-5508 (2016).

4. Xu Y, Lin Q, Han Z, He Y, Bi Y. Intrinsic functional network architecture of human semantic processing: Modules and hubs. *Neuroimage* **132**, 542-555 (2016).

5. van den Heuvel MP*, et al.* Abnormal Rich Club Organization and Functional Brain Dynamics in Schizophrenia. *JAMA Psychiatry* **70**, 783-792 (2013).

6. Crossley NA*, et al.* The hubs of the human connectome are generally implicated in the anatomy of brain disorders. *Brain* **137**, 2382-2395 (2014).

7. Buckner RL*, et al.* Cortical Hubs Revealed by Intrinsic Functional Connectivity: Mapping, Assessment of Stability, and Relation to Alzheimer's Disease. *The Journal of Neuroscience* **29**, 1860 (2009).

8. Dai Z*, et al.* Identifying and Mapping Connectivity Patterns of Brain Network Hubs in Alzheimer's Disease. *Cerebral Cortex* **25**, 3723-3742 (2014).

9. Wang J*, et al.* Disrupted Functional Brain Connectome in Individuals at Risk for Alzheimer's Disease. *Biological Psychiatry* **73**, 472-481 (2013).

10. Wang L*, et al.* The Effects of Antidepressant Treatment on Resting-State Functional Brain Networks in Patients With Major Depressive Disorder. *Human Brain Mapping* **36**, 768-778 (2015).

11. Fornito A, Zalesky A, Breakspear M. The connectomics of brain disorders. *Nature Reviews Neuroscience* **16**, 159-172 (2015).

12. Gong Q, He Y. Depression, Neuroimaging and Connectomics: A Selective Overview. *Biological Psychiatry* **77**, 223-235 (2015).

13.   Achard S, Salvador R, Whitcher B, Suckling J, Bullmore E. A Resilient, Low-Frequency, Small-World Human Brain Functional Network with Highly Connected Association Cortical Hubs. *The Journal of Neuroscience* **26**, 63-72 (2006).

14.   Liao X-H*, et al.* Functional brain hubs and their test–retest reliability: A multiband resting-state functional MRI study. *NeuroImage* **83**, 969-982 (2013).

15.   Cole MW, Pathak S, Schneider W. Identifying the brain's most globally connected regions. *Neuroimage* **49**, 3132-3148 (2010).

16.   Tomasi D, Volkow ND. Functional connectivity density mapping. *Proc Natl Acad Sci USA* **107**, 9885-9890 (2010).

17.   Fransson P, Aden U, Blennow M, Lagercrantz H. The Functional Architecture of the Infant Brain as Revealed by Resting-State fMRI. *Cerebral Cortex* **21**, 145-154 (2011).

18.   Tomasi D, Volkow ND. Association between Functional Connectivity Hubs and Brain Networks. *Cerebral Cortex* **21**, 2003-2013 (2011).

19.   de Pasquale F*, et al.* The connectivity of functional cores reveals different degrees of segregation and integration in the brain at rest. *Neuroimage* **69**, 51-61 (2013).

20.   Glahn DC*, et al.* Genetic control over the resting brain. *Proc Natl Acad Sci USA* **107**, 1223-1228 (2010).

21.   Fornito A*, et al.* Genetic Influences on Cost-Efficient Organization of Human Cortical Functional Networks. *The Journal of Neuroscience* **31**, 3261-3270 (2011).

22.   Wiggins JL*, et al.* The impact of serotonin transporter (5-HTTLPR) genotype on the development of resting-state functional connectivity in children and adolescents: a preliminary report. *Neuroimage* **59**, 2760-2770 (2012).

23.   Gordon EM, Stollstorff M, Devaney JM, Bean S, Vaidya CJ. Effect of Dopamine Transporter Genotype on Intrinsic Functional Connectivity Depends on Cognitive State. *Cerebral Cortex* **22**, 2182-2196 (2012).

24.   Richiardi J*, et al.* Correlated gene expression supports synchronous activity in brain networks. *Science* **348**, 1241-1244 (2015).

599 25. Diez I, Sepulcre J. Neurogenetic profiles delineate large-scale connectivity dynamics of
600   the human brain. *Nature Communications* **9**, 3876 (2018).

601
602 26. Vertes PE*, et al.* Gene transcription profiles associated with inter-modular hubs and
603   connection distance in human functional magnetic resonance imaging networks. *Phil Trans*
604   *R Soc B* **371**, (2016).

605
606 27. Fornito A, Arnatkevičiūtė A, Fulcher BD. Bridging the Gap between Connectome and
607   Transcriptome. *Trends in Cognitive Sciences* **23**, 34-50 (2019).

608
609 28. Nichols TE, Holmes AP. Nonparametric Permutation Tests For Functional Neuroimaging:
610   A Primer with Examples. *Human Brain Mapping* **15**, 1-25 (2002).

611
612 29. Yeo BT*, et al.* The organization of the human cerebral cortex estimated by intrinsic
613   functional connectivity. *Journal of Neurophysiology* **106**, 1125-1165 (2011).

614
615 30. Tzourio-Mazoyer N*, et al.* Automated Anatomical Labeling of Activations in SPM Using
616   a Macroscopic Anatomical Parcellation of the MNI MRI Single-Subject Brain.
617   *Neuroimage* **15**, 273-289 (2002).

618
619 31. Power JD, Schlaggar BL, Lessov-Schlaggar CN, Petersen SE. Evidence for Hubs in
620   Human Functional Brain Networks. *Neuron* **79**, 798-813 (2013).

621
622 32. Gordon EM*, et al.* Three Distinct Sets of Connector Hubs Integrate Human Brain Function.
623   *Cell reports* **24**, 1687-1695 e1684 (2018).

624
625 33. Chen T, Guestrin C. XGBoost: A Scalable Tree Boosting System. *arXiv*, (2016).

626
627 34. Arnatkevic Iute A, Fulcher BD, Fornito A. A practical guide to linking brain-wide gene
628   expression and neuroimaging data. *Neuroimage* **189**, 353-367 (2019).

629
630 35. Eden E, Navon R, Steinfeld I, Lipson D, Yakhini Z. GOrilla: a tool for discovery and
631   visualization of enriched GO terms in ranked gene lists. *BMC bioinformatics* **10**, 48 (2009).

632
633 36. Huang da W, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward
634   the comprehensive functional analysis of large gene lists. *Nucleic Acids Research* **37**, 1-13
635   (2009).

636
637 37. Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene
638   lists using DAVID bioinformatics resources. *Nature protocols* **4**, 44-57 (2009).

640   38.   Kang HJ, *et al.* Spatio-temporal transcriptome of the human brain. *Nature* **478**, 483-489
641   (2011).

643   39.   Harris Julia J, Jolivet R, Attwell D. Synaptic Energy Use and Supply. *Neuron* **75**, 762-777
644   (2012).

646   40.   Arroyo JD*, et al.* A Genome-wide CRISPR Death Screen Identifies Genes Essential for
647   Oxidative Phosphorylation. *Cell Metabolism* **24**, 875-885 (2016).

649   41.   Goyal MS, Hawrylycz M, Miller JA, Snyder AZ, Raichle ME. Aerobic Glycolysis in the
650   Human Brain Is Associated with Development and Neotenous Gene Expression. *Cell*
651   *Metabolism* **19**, 49-57 (2014).

653   42.   Miller JA*, et al.* Transcriptional landscape of the prenatal human brain. *Nature* **508**, 199-
654   206 (2014).

656   43.   Sherwood CC, Gómez-Robles A. Brain Plasticity and Human Evolution. *Annual Review*
657   *of Anthropology* **46**, 399-419 (2017).

659   44.   Huttenlocher PR, Dabholkar AS. Regional Differences in Synaptogenesis in Human
660   Cerebral Cortex. *The Journal of comparative neurology* **387**, 167-178 (1997).

662   45.   Bajada CJ, Schreiber J, Caspers S. Fiber length profiling: A novel approach to structural
663   brain organization. *Neuroimage* **186**, 164-173 (2019).

665   46.   Vaishnavi SN, Vlassenko AG, Rundle MM, Snyder AZ, Mintun MA, Raichle ME.
666   Regional aerobic glycolysis in the human brain. *Proc Natl Acad Sci USA* **107**, 17757-17762
667   (2010).

669   47.   Liang X, Zou Q, He Y, Yang Y. Coupling of functional connectivity and regional cerebral
670   blood flow reveals a physiological basis for network hubs of the human brain. *Proc Natl*
671   *Acad Sci USA* **110**, 1929-1934 (2013).

673   48.   Cumming G. *Understanding the new statistics: Effect sizes, confidence intervals, and*
674   *meta-analysis*. Routledge (2013).

676   49.   Fox MD, Corbetta M, Snyder AZ, Vincent JL, Raichle ME. Spontaneous neuronal activity
677   distinguishes human dorsal and ventral attention systems. *Proc Natl Acad Sci USA* **103**,
678   10046-10051 (2006).

50.   Marek S, Dosenbach NUF. The frontoparietal network: function, electrophysiology, and
importance of individual precision mapping. *Dialogues in clinical neuroscience* **20**, 133-
140 (2018).

51.   Felleman DJ, Van Essen DC. Distributed Hierarchical Processing in the Primate Cerebral
Cortex. *Cerebral Cortex* **1**, 1-47 (1991).

52.   Buckner RL, Andrews-Hanna JR, Schacter DL. The Brain's Default Network Anatomy,
Function, and Relevance to Disease. *Annals of the New York Academy of Sciences* **1124**,
1-38 (2008).

53.   Braga RM, Buckner RL. Parallel Interdigitated Distributed Networks within the Individual
Estimated by Intrinsic Functional Connectivity. *Neuron* **95**, 457-471 e455 (2017).

54.   Menon V. Large-scale brain networks and psychopathology: a unifying triple network
model. *Trends in Cognitive Sciences* **15**, 483-506 (2011).

55.   Xie Y, *et al.* Alterations in Connectome Dynamics in Autism Spectrum Disorder: A
Harmonized Mega- and Meta-Analysis Study Using the ABIDE Dataset. *Biological
Psychiatry*, (2021).

56.   Nicholls JG, Martin AR, Fuchs PA, Brown DA, Diamond ME, Weisblat DA. *From Neuron
to Brain, Fifth Edition*. Sinauer Associates Sunderland, MA (2012).

57.   Changeux J-P, Danchin A. Selective stabilisation of developing synapses as a mechanism
for the specification of neuronal networks. *Nature* **264**, 705-712 (1976).

58.   Petanjek Z, *et al.* Extraordinary neoteny of synaptic spines in the human prefrontal cortex.
Proc Natl Acad Sci USA **108**, 13281-13286 (2011).

59.   Miller DJ, *et al.* Prolonged myelination in human neocortical evolution. *Proc Natl Acad
Sci USA* **109**, 16480–16485 (2012).

60.   Lebel C, Gee M, Camicioli R, Wieler M, Martin W, Beaulieu C. Diffusion tensor imaging
of white matter tract evolution over the lifespan. *Neuroimage* **60**, 340-352 (2012).

61.   Paredes MF, *et al.* Extensive migration of young neurons into the infant human frontal lobe.
Science **354**, (2016).

62.   Kwan KY, Sestan N, Anton ES. Transcriptional co-regulation of neuronal migration and
720   laminar identity in the neocortex. *Development* **139**, 1535-1546 (2012).

721
722   63.   Wei Y*, et al.* Statistical testing in transcriptomic-neuroimaging studies: A how-to and
723   evaluation of methods assessing spatial and gene specificity. *Human Brain Mapping* **43**,
724   885-901 (2022).

725
726   64.   Fulcher BD, Arnatkeviciute A, Fornito A. Overcoming false-positive gene-category
727   enrichment in the analysis of spatially resolved transcriptomic brain atlas data. *Nature
728   Communications* **12**, 2669 (2021).

729
730   65.   Helmer M*, et al.* On stability of Canonical Correlation Analysis and Partial Least Squares
731   with application to brain-behavior associations. *bioRxiv*, (2020).

732
733   66.   Ciric R*, et al.* Benchmarking of participant-level confound regression strategies for the
734   control of motion artifact in studies of functional connectivity. *Neuroimage* **154**, 174-187
735   (2017).

736
737   67.   Schwarz AJ, McGonigle J. Negative edges and soft thresholding in complex network
738   analysis of resting state functional connectivity data. *Neuroimage* **55**, 1132-1146 (2011).

739
740   68.   Power Jonathan D*, et al.* Functional Network Organization of the Human Brain. *Neuron*
741   **72**, 665-678 (2011).

742
743   69.   Eickhoff SB, Laird AR, Grefkes C, Wang LE, Zilles K, Fox PT. Coordinate-Based
744   Activation Likelihood Estimation Meta-Analysis of Neuroimaging Data: A Random-
745   Effects Approach Based on Empirical Estimates of Spatial Uncertainty. *Human Brain
746   Mapping* **30**, 2907-2926 (2009).

747
748   70.   Burt JB, Helmer M, Shinn M, Anticevic A, Murray JD. Generative modeling of brain maps
749   with spatial autocorrelation. *Neuroimage* **220**, 117038 (2020).

Fig 1. Enhanced box plot of the age ranges of each cohort. M/F, males/femals.

Fig 2. Highly consistent and reproducible functional connectome hubs. A Robust FCS
pattern and its corresponding variance (standard error, SE) map estimated using a harmonized
voxelwise random-effects meta-analysis across 61 cohorts. **B** Left: The most consistent
functional connectome hubs ($p < 0.001$, cluster size > 200 mm^3); white spheres represent hub
peaks. Right: Hub voxel distribution in eight large-scale brain networks; insets, the seven large-
scale cortical networks[29] were rendered on the left hemisphere. SUB, subcortical network;
LIMB, limbic network. **C** Left: Heterogeneity measurement I^2 estimated through the random-
effects meta-analysis. Right: Cumulative distribution function plot of I^2. **D** Left: Heatmap of
displacements of the 35 hub peaks after leaving one cohort out. Right: Bar plot of the probability
across the 35 hub peaks whose displacement was less than 6 mm after leaving one cohort out. **E**
Hub occurrence probability map across all subjects (left) and all cohorts (right). White lines
delineate boundaries of the identified hubs in **B**. **F** Dice's coefficient of the identified hubs in **B**
compared with the top N (voxel number of the identified hubs in **B**) voxels with the highest hub
occurrence probability values across randomly selected subjects (left) and randomly selected
cohorts (right). Blue shading represents the standard deviation across 2,000 random selections.

Fig 3. Functional connectivity profiles of connectome hubs. A Functional connectivity profiles of the 35 hubs. White spheres represent hub seeds. Blue lines delineate boundaries of the seven cortical networks shown in Fig 2B. **B** Top: Dendrogram derived by hierarchical clustering of the connectivity percentage matrix. Middle: The 35 hubs were rendered using three different colors according to the hierarchical clustering solution. Bottom: Radar charts showing heterogeneous connectivity profiles of the three hub clusters.

Fig 4. Transcriptomic data distinguishes connectome hubs from non-hubs. **A** Schematic
diagram of using the XGBoost model to classify brain samples as a hub or non-hub. **B**
Performance of the XGBoost classifier. Each dot represents one repetition in **A**. The horizontal
gray dashed line represents the chance level accuracy rate (50%). The horizontal green dashed
line represents the average accuracy rate of the XGBoost classifier across 1,000 repetitions. **C**

Density plot of 10,027 genes' logarithmic average contributions across 1,000 repetitions to the XGBoost classifier. Genes with the greatest contributions were regarded as key genes. **D** Regression plot of the logarithmic average contributions of the top 300 key genes across the first 500 repetitions versus those across the second 500 repetitions. Each dot represents one gene. **E** Schematic diagram of using the SVM model to classify brain samples as a hub or non-hub. **F and G** Accuracy rate of the SVM classifier versus the count of key genes used to distinguish 382 hub samples from 382 non-hub samples with the highest rate (**F**) or lowest rate (**G**) to be correctly classified by the XGBoost classifier. Each dot represents one SVM classifier. Black curves were estimated by locally weighted regression. **H** Performance of the SVM classifier. Horizontal lines correspond to the SVM classifier trained using top 150 key genes in **G**. Each dot represents one repetition using 150 randomly selected genes in **E**. The horizontal gray dashed line represents the chance level accuracy rate (50%).

Fig 5. Connectome hubs have a spatiotemporally distinctive transcriptomic pattern. A
Transcription level differences between hub samples (n=382) and non-hub samples (n=776) for
genes associated with key neurodevelopmental processes[38] and main neuronal metabolic
pathways[39]. Boxplot edges, gray lines, and whiskers and dots depict the 25th and 75th
percentiles, median, and extreme nonoutlier and outlier values, respectively. Significance of one-
sided Wilcoxon rank-sum tests were determined by 1,000 permutation tests and were labeled
with Bonferroni-corrected *p* values. **B** Transcriptomic trajectory differences between hub and
non-hub regions for genes involved in key neurodevelopmental processe[38] and main neuronal
metabolic pathways[39]. w, post-conceptional week; y, postnatal year; a.u., arbitrary unit.

Fig 6. Connectome hubs have more intricate fiber configuration. Left: Fiber number for different fiber length bins was derived from a fiber length profiling dataset[45]. To save space, we only presented data of the left hemisphere and used data of both hemispheres in boxplots and statistics. White lines delineate boundaries of the identified hubs in Fig 2B. Right: Fiber number difference between connectome hubs (red, n=25,944) and non-hubs (blue, n=33,195). Boxplot edges, gray lines, and whiskers and dots depict the 25th and 75th percentiles, median, and extreme nonoutlier and outlier fiber number values, respectively. Significance of one-sided Wilcoxon rank-sum tests were determined by 1,000 permutation tests and were labeled with Bonferroni-corrected *p* values. a.u., arbitrary unit.

Fig 7. Connectome hubs have higher metabolic rate. The cerebral metabolic rate for oxygen, glycolytic index, and cerebral blood flow of 82 Brodmann areas and seven subcortical structures were provided by a prior study[46]. White lines delineate boundaries of the identified hubs in Fig 2B. Boxplot edges, gray lines, and whiskers and dots depict the 25th and 75th percentiles, median, and extreme nonoutlier and outlier metabolic measurement values, respectively. Brodmann areas with more than 50% vertices or subcortical structures with more than 50% voxels identified as hubs were regarded as hub regions (n=29), vice versa as non-hub regions (n=60). Significance of one-sided Wilcoxon rank-sum tests was determined by 1,000 permutation tests and were labeled with Bonferroni-corrected *p* values. a.u., arbitrary unit.

     Table 1. Highly consistent functional connectome hubs.

No.	Hub	Location	MNI coordinates			Cohen's d	FCS	SE
			x	y	z			
1	Right PFt	PFt (superoanterior BA* 40)	60	-21	45	6.267	1.072	0.022
2	Left PFt	PFt (superoanterior BA 40)	-60	-24	36	6.151	0.949	0.020
3	Right PF	PF (posterior BA 40)	60	-27	24	5.785	1.239	0.027
4	Left SCEF	Supplementary and cingulate eye field	0	0	51	5.635	1.000	0.023
5	Left PGi	PGi (inferior BA 39)	-51	-66	30	5.168	1.075	0.027
6	Left PFop	PF opercular (inferoanterior BA 40)	-63	-27	18	5.160	1.095	0.027
7	Left 43	Area 43	-57	3	3	4.927	1.114	0.029
8	Right 6r	Rostral area 6	57	6	0	4.916	1.184	0.031
9	Right PGi	PGi (inferior BA 39)	54	-60	30	4.739	1.007	0.027
10	Right 8BL	Area 8B lateral	21	36	51	4.655	0.713	0.020
11	Right 7PC	Area 7PC	36	-45	54	4.414	0.712	0.021
12	Left 9p	Area 9 posterior	-15	45	45	4.199	0.639	0.019
13	Right 6v	Ventral area 6	54	9	33	4.037	0.766	0.024
14	Left 8Av	Ventral area 8A	-39	18	48	3.990	0.561	0.018
15	Left AIP	Anterior intra-parietal area	-33	-45	45	3.474	0.567	0.021
16	Right FST	Fundus of the superior temporal area	54	-60	0	3.156	0.729	0.030
17	Right 9m	Area 9 middle	3	54	24	3.128	0.609	0.025
18	Left 31pv	Area 31p ventral	-3	-51	33	3.049	0.784	0.033
19	Right VIP	Ventral intra-parietal complex	18	-63	57	2.984	0.572	0.025
20	Right 6a	Area 6 anterior	33	3	63	2.975	0.454	0.020
21	Left FOP4	Frontal opercular area 4	-33	21	6	2.858	0.828	0.037
22	Right 5mv	Area 5m ventral	12	-30	45	2.822	0.701	0.032
23	Right 46	Area 46	36	42	30	2.779	0.656	0.030
24	Left 10v	Area 10v	0	57	-9	2.769	0.731	0.034
25	Left p9-46v	Area posterior 9-46v	-42	36	27	2.591	0.561	0.028
26	Left V3A	Area V3A	-15	-90	33	2.575	0.684	0.034
27	Left TE1a	Area TE1 anterior	-63	-15	-15	2.527	0.595	0.030
28	Right TE1a	Area TE1 anterior	60	-9	-21	2.494	0.580	0.030
29	Right IFSa	Anterior inferior frontal suleus	48	39	12	2.468	0.480	0.025
30	Left 7Am	Medial area 7A	-12	-60	60	2.461	0.475	0.025
31	Right V3A	Area V3A	18	-87	36	2.442	0.645	0.034
32	Right V4	Fourth visual area	24	-63	-9	2.339	0.446	0.024
33	Left 6a	Area 6 anterior	-24	3	63	2.317	0.331	0.018
34	Left VMV1	Ventromedial visual area 1	-18	-60	-6	1.937	0.397	0.026
35	Left FEF	Frontal eye fields	-45	-9	57	1.412	0.640	0.058

     * BA, Brodmann area.