Mutation, selection, and the prevalence of the *C. elegans* mortal germline phenotype

Sayran Saber[1,*], Michael Snyder[1,*], Moein Rajaei[1], and Charles F. Baer[1,2]

1- Department of Biology, University of Florida, Gainesville, FL USA

2- University of Florida Genetics Institute, Gainesville, FL USA

* These authors contributed equally

Communication to: cbaer@ufl.edu

Abstract

C. elegans strains with the mortal germline (Mrt) phenotype become progressively sterile over the course of a few tens of generations. Mrt is proximately controlled epigenetically, and is typically temperature-dependent, being penetrant at temperatures near the upper range of *C. elegans*' tolerance. Previous studies have suggested that Mrt presents a relatively large mutational target, and that Mrt is not uncommon in natural populations of *C. elegans*. The Mrt phenotype is not monolithic. Some strains exhibit a strong Mrt phenotype, in which individuals invariably become sterile over a few generations, whereas other strains show a weaker (less penetrant) phenotype in which the onset of sterility is slower and more stochastic. We present results in which we (1) quantify the rate of mutation to the Mrt phenotype, and (2) quantify the frequency of Mrt in a collection of 95 wild isolates. Over the course of ~16,000 meioses, we detected one mutation to a strong Mrt phenotype, resulting in a point estimate of the mutation rate $U_{Mrt} \approx 6 \times 10^{-5}$/genome/generation. We detected no mutations to a weak Mrt phenotype. 5/95 wild isolates had a strong Mrt phenotype, and although quantification of the weak Mrt phenotype is inexact, the weak Mrt phenotype is not rare in nature. We estimate a strength of selection against mutations conferring the strong Mrt phenotype $\bar{s} \approx 0.1\%$, similar to selection against mutations affecting competitive fitness. The appreciable frequency of weak Mrt variants in nature combined with the low mutation rate suggests that Mrt may be maintained by balancing selection.

Introduction

The *C. elegans* "mortal germline" (Mrt) phenotype is an epigenetically controlled, transgenerationally heritable trait in which Mrt lineages become progressively sterile over the course of a few to a few tens of generations (Ahmed and Hodgkin 2000; Smelick and Ahmed 2005). However, like any trait governed epigenetically, it has an ultimate, underlying genetic basis. The Mrt phenotype was first discovered in a mutant strain defective in germline telomere replication and DNA repair (Ahmed and Hodgkin 2000). Subsequent studies have identified numerous Mrt mutants, many of which are associated with defects in nuclear RNAi (Katz et al. 2009; Buckley et al. 2012; Spracklin et al. 2017). Many, but not all, Mrt mutants are temperature-sensitive, with continued exposure to high temperature leading to onset of sterility. Based on the frequency of appearance of Mrt mutants in forward genetic screens, it was suggested that many genes are capable of producing the Mrt phenotype if mutated (Smelick and Ahmed 2005), or in other words, the Mrt phenotype presents a large mutational target (Houle 1998). Note that "large mutational target" in this context is not synonymous with "polygenic" in the usual sense, because even if many genes potentially affect the trait, a mutation in any one gene is sufficient to produce the Mrt phenotype. On the other hand, Mrt probably is polygenic, with subtle phenotypic variation resulting from segregating variants at many loci. However, it would be challenging to discern whether a given genotype becomes sterile after (say) 13 generations vs. 14 generations, on average.

Over the past two decades, the realm of *C. elegans* biology has expanded beyond its initial role as a model system par excellence for functional biology to include studies of natural variation (Dirksen et al. 2016; Felix and Duveau 2012; Schulenburg and Felix 2017; Cook et al. 2017). It soon became apparent that some wild isolates could not be maintained in culture at 25° C (near the upper range of *C. elegans* thermal tolerance), and further, that most such strains had the temperature-sensitive Mrt phenotype (Frezal et al. 2018). Temperature-sensitive Mrt strains can typically be rescued by exposure to cool temperature (15° C) for a

generation, and it is unclear if long-term (multi-generational) exposure to temperatures

sufficiently high to induce the Mrt phenotype are common in *C. elegans'* natural environment.

At first glance, the Mrt phenotype would seem to be the manifestation of context-dependent

mutations, which are neutral in the wild, and only become deleterious in the lab environment.

However, that scenario requires bidirectional mutation, such that Mrt alleles mutate into wild-

type alleles as well as the reverse; if not, the population would eventually mutate its way to

fixation for the Mrt phenotype. Bidirectional mutation is possible, of course, but the evidence at

hand suggests it is not common, because the Mrt phenotype is associated with loss-of-function

mutations.

A straightforward alternative to context-dependent neutrality is that Mrt alleles are

deleterious in nature, in which case genetic variation is maintained by mutation-(purifying)

selection balance (MSB). That possibility is intuitively attractive because, all else equal, sterility

will never be favored by natural selection. All else may not be equal, however; the Mrt

phenotype may be a pleiotropic correlate of some other trait(s) for which variation is maintained

by some form of balancing selection (BS).

To begin to sort out the various possibilities by which variation for the Mrt phenotype is

maintained, we need to know (1) the rate of input of new genetic variation by mutation, and (2)

the frequency of the Mrt phenotype in nature. At any one locus, the equilibrium frequency of a

deleterious allele at MSB, $\hat{q} \approx \frac{\mu}{s}$, where μ is mutation rate from wild-type to the deleterious allele

and s is the strength of selection against the mutant allele (the homozygous effect in an

organism with near-complete self-fertilization, such as *C. elegans*; (Haldane 1927)). In a

(nearly) completely inbred population, by extrapolation over the entire genome, the probability

that an individual is not Mrt $\approx 1 - \sum_i \hat{q}_i$, summed over all *i* loci capable of yielding the Mrt

phenotype when mutated. In a population at MSB, the expected frequency of the Mrt

phenotype is approximately $\frac{U}{\bar{s}}$, where U is the genome-wide rate of mutation to Mrt alleles and $\bar{s}$

is the average strength of selection against an Mrt allele.

We estimated the rate of mutation to the Mrt phenotype from two sets of *C. elegans*

laboratory mutation accumulation (MA) lines, which evolved in the near-absence of natural

selection for approximately 250 generations. On average, each MA line carries about 65 unique

spontaneous base-substitution and small indel mutations (Rajaei et al. 2021), and probably a

few larger structural variants (A. S. Saxena and Baer, unpublished results). In addition, we

estimated the frequency of the Mrt phenotype in a worldwide collection of ~100 wild isolates.

From these data, we infer the approximate strength of purifying selection acting on new Mrt

mutations. Although we refer to "the mortal germline" as if it was a discrete, presence/absence

trait, in reality, the mortal germline exists along a continuum (Frezal et al. 2018), and our

analysis takes that fact into account.

Materials and Methods

Mutation Accumulation Experiment. Details of the mutation accumulation protocol are given in

Baer et al. (2005). N2 is the standard laboratory strain of *C. elegans*; PB306 is a wild isolate

generously provided by Scott Baird. The basic protocol follows that of Vassilieva and Lynch

(1999) and is outlined in **Supplemental Figure S1**. Briefly, 100 replicate populations (MA lines)

were initiated from a cryopreserved stock of a highly-inbred ancestor ("G0") at mutation-drift

equilibrium and propagated by transferring a single immature hermaphrodite at one-generation

(four-day) intervals. Lines were maintained on 60mm NGM agar plates, spotted with 100 *μl* of

an overnight culture of the OP50 strain of *E. coli* B, at a constant 20°C. The lines were

propagated for 250 transfers (Gmax=250), beginning in March, 2001 and culminating with a

final cryopreservation in 2005.

Wild Isolates. A collection of wild isolates of *C. elegans* was obtained from Erik Andersen

(Northwestern University) in 2015 and cryopreserved in the Baer lab. A list of the wild isolates

is given in **Supplemental Table S3**. The genome sequences of the wild isolates along with

collection information are available at https://www.elegansvariation.org/.

Mortal Germline Assay. The assay is based on that of Frezal et al. (2018), and is schematically

depicted in **Figure 1**. Cryopreserved samples of the G0 ancestor were thawed onto 35 mm

Figure 1. Schematic diagram of the Mrt assay. See Methods for details of the experiment.

plates seeded with OP50 and incubated at 20° for three days, at which time 18 L4-stage

hermaphrodites were picked individually to seeded 35 mm plates and incubated at 15°. The 18

replicates of the G0 ancestor were subsequently treated identically to the MA lines; we refer to

these as "pseudolines" (PS). The following day, 35 randomly selected N2 (block 1) or PB306

(block 2) MA lines were thawed from cryopreserved samples and incubated at 15° C. PS and

MA lines were allowed to reproduce for one generation (g.-3) at 15° C, at which point ten

replicates of each line (MA and PS) were initiated by transferring a single L4 hermaphrodite to a

seeded 35 mm plate. Each replicate was allowed to reproduce at 15°C for two more generations

generation (g.-2, g.-1), at which time three L4s from each replicate were transferred to a seeded

35 mm plate (g.0) and incubated at 25° C. Subsequently, three L4s were transferred at three-

day intervals for the duration of the assay. The N2 assay (block 1) was propagated for 21

generations; we terminated the PB306 assay (block 2) after 14 generations because it was

evident that there were no lines with a Mrt phenotype.

The assay of the wild isolates was identical to that of the N2 MA lines (21 generations)

except there were only three replicates per line rather than ten.

Data Access

Mrt assay data are included in supplemental tables S1 and S3 and in Dryad ###. Genome

sequence data of MA line 578 is deposited in the NCBI Sequence Read Archive under

Accession number PRJNA665851, sample SAMN16272702.

Results

Mutation

Raw survival data are given in **Supplemental Table S1**. Discriminating the Mrt from non-Mrt

phenotype in the MA lines turns out to be straightforward. In the N2 G0 progenitor, only ten of

the 180 replicates (18 PS lines, 10 replicates/line) became sterile before termination of the

assay at generation 21, and only one of the 18 PS lines had more than one replicate fail to finish

the assay. Of the 34 N2 MA lines assayed, one (line 540) incurred a heat-sensitive sterile

mutation, identified as such because all ten replicates of the line were sterile after the first

generation at 25°. Temperature-sensitive sterile and lethal mutations are well-documented in

many organisms, and are distinct from Mrt. One line (line 578) had an obvious strong Mrt

phenotype; all ten replicates were sterile by generation ten (median time to sterility = six

generations). In contrast, no replicate of any other N2 MA line became sterile before generation

7, and only one line (516) had more than one replicate become sterile before generation 11.

However, the other seven replicates of line 516 made it to the finish, and in the three replicates

that went extinct, the worms did not appear to have the classical sterile phenotype (see Figure

1D of Frezal et al. 2018); rather, they simply looked sickly.

Of the 180 replicates of the PB306 progenitor, only one became sterile prior to

completion of the assay at generation 14. Of the 33 PB306 MA lines, one (line 471) had 6/10

replicates fail before generation seven. However, line 471 has low fitness even at 20°, and the

remaining four replicates survived to the end. Nor did the replicates that failed have an obvious

sterile phenotype; rather, they were characterized by slow growth and dead worms.

Accordingly, we do not classify line 471 as Mrt. No other PB306 MA line had more than one

replicate fail to complete the assay.

From these data, we conclude that 1/34 N2 lines and 0/33 PB306 lines accumulated a

Mrt mutation. We can calculate the point estimate of the genome-wide rate of mutation to Mrt

as $U_{Mrt}=k/Lt$, where k is the number of Mrt mutations observed (one in N2 and zero in PB306,

assuming that the one observed Mrt phenotype is the result of a single mutation, which it

appears to be), L is the number of MA lines included, and t is the number of generations of MA.

Note that this is the haploid rate, but that mutations accumulated in diploids; double the number

of genomes (for diploidy) is cancelled by the probability of loss of a new neutral mutation in an

MA line, which is 1/2. Pooling over the two sets of lines, the point estimate of $U_{Mrt}=$ 1/(67×250)≈

6×10^{-5}/genome/generation. The 95% confidence interval on the value of U_{Mrt} consistent with

the observed number of mutations can be calculated from the Poisson probability of observing k

mutations $P(k)=e^{-\lambda}(\lambda^k/k!)$, where λ is the expected number of mutations and $k=1$ (Schug et al.

1997). The smallest and largest values of λ for which $P(1)=0.025$ establish the lower and upper

95% confidence limit on λ; dividing that value by the product of the number of MA lines (L=67)

and the number of generations of MA (t≈250) gives the confidence interval on U_{Mrt}. The

approximate 95% CI on U_{Mrt} is (1.5×10^{-6} – 3.2×10^{-4}/genome/generation). The per-nucleotide

mutation rate in these lines is approximately 2.8×10^{-9}/generation (Saxena et al. 2019; Rajaei et

al. 2021) and the *C. elegans* genome is approximately 10^{8} bp, which implies that the mutational

target of the Mrt phenotype is about 0.02%, and possibly as much as 0.1% of the *C. elegans*

genome.

Given that one, and only one, MA line has a clear Mrt phenotype, we scrutinized its

genome for candidate mutations. Line 578 carries 37 unique base-substitutions, 13 deletions,

and four insertions relative to the genome of the progenitor of the N2 MA lines (**Supplemental**

Table S2). There is one obvious candidate, an 11-base frameshift insertion in an exon of the

nrde-2 gene. *nrde-2* is so-named for its <u>N</u>uclear <u>R</u>NAi <u>D</u>efective phenotype (Guang et al. 2008),

is involved in heterochromatin assembly by small RNA as well as nuclear RNAi, and has been

shown to be involved in temperature-dependent transgenerational nuclear silencing (Sakaguchi

et al. 2014).

Standing genetic variation

95 wild isolates ("strains") were chosen haphazardly, based on a collection by E. C. Andersen.

Assay data are given in **Supplemental Table S3**. Unfortunately, unlike that of the MA lines, the

phenotypic distribution of the wild isolates cannot be unambiguously categorized into Mrt and

Not Mrt. The difficulty has (at least) two sources. First, and most importantly, the sample size

per strain is much smaller (three replicates per strain, as opposed to ten per MA line), and

second, there appear to be small-effect QTL segregating in the population that contribute a non-

trivial fraction of the heritable variation (Frezal et al. 2018).

Five of the 95 strains had an unambiguous strong Mrt phenotype, based on the criterion

that no replicate in any assay survived past 12 generations (**Figure 2**). Progressively loosening

the Mrt criteria, 18 strains had no replicate remaining by generation 21, and another 13 strains

had only a single replicate remaining by generation 21. 42 strains remained fertile at 21

generations in all three replicates. Depending on the stringency of the criteria, the frequency of

the Mrt phenotype in the wild isolates is as low as 5/95 (~5%) and as high as 53/95 (~56%).

Figure 2. Average time to sterility of wild isolates (n=3 reps/isolate). Strain CB4858 is

temperature-dependent sterile and not Mrt.

Mutation, selection, and the maintenance of genetic variation

We begin with the most stringently defined wild isolates, of which there are five. These strains

clearly have the same strong Mrt phenotype as MA line 578. Solving the equation $\hat{q} \approx \frac{U}{s}$ for $\hat{q} \approx$

0.05 and U=6×10^{-5}/generation, we infer a strength of selection s≈0.001. That strength of

purifying selection is on the order of that inferred to be acting against deleterious alleles that

affect competitive fitness (Yeh et al. 2017). The frequency of the strong Mrt phenotype is

entirely consistent with genetic variation being maintained by MSB.

The MA data are not so clear with respect to weaker Mrt phenotypes. The distribution of

mean time-to-sterility among MA lines is no different from that of the G0 PS lines in either MA

background (**Figure 3**). An upper bound on the genome-wide mutation rate to weak Mrt alleles

that is consistent with observing no MA line with that phenotype can be calculated from the

Poisson probability of observing k mutations $P(k)=e^{-\lambda}(\lambda^{k}/k!)$, where λ is the expected number of

mutations and $k=0$ (Schug et al. 1997). $P(k)=0.025$ establishes the upper 95% confidence limit

of the expected number of mutations consistent with observing no mutations; dividing that value

by the product of the number of MA lines (L) and the number of generations of MA (t) gives the

upper 95% confidence limit on the genomic mutation rate U consistent with observing no

mutations. U as high as 2.2×10^{-4}/generation is consistent with the observed absence of weak

Mrt phenotypes among the MA lines.

Figure 3. Frequency distribution of time to sterility of individual replicates in the MA assay.

MA lines above the mid-line, G0 pseudolines (PS) below. (**A**) N2. (**B**) PB306.

Discussion

The high frequency of weak Mrt phenotypes in the wild isolates seems incongruous with the

failure to observe even a single MA line with a weak Mrt phenotype. Long-term maintenance of

neutral variation requires bidirectional mutation. It is evident that the strong Mrt phenotype is

deleterious in the lab environment. Fortuitously, we have a strain (XZ1516) in long-term mass

culture at 20° that has a strong temperature-dependent Mrt phenotype that is weakly penetrant

at 20°. As an *ad hoc* test for back-mutation of a strong Mrt phenotype in the presence of

selection, we initiated ten replicates of our Mrt assay at 25° with XZ1516 worms that had been

cryopreserved after ~80 generations in mass culture at 20°. All ten replicates were sterile by

seven generations. Obviously that would have been a more meaningful test had we kept the

strain in mass culture at 25° rather than 20° (and used more than a single strain), but it at least

suggests that back-mutations from a strong Mrt phenotype are infrequent. Of course, the

genetic basis underlying weak Mrt phenotypes is likely to be different from that of the strong Mrt

phenotype, which has been shown to typically result from loss-of-function mutations at protein-

coding loci (e.g., MA line 578). Based on what is known about quantitative traits in general

(Manolio et al. 2009; Boyle et al. 2017), it seems likely that much variation in weak Mrt is the

result of variation in the magnitude and/or timing of expression of genes that confer a strong Mrt

phenotype when silenced. On the other hand, "typical" quantitative traits accumulate abundant

mutational variance (Houle et al. 1996; Davies et al. 2016), which is not the case for the weak

Mrt phenotype in these lines.

Taken together, abundant genetic variation in nature coupled with a low rate of input of

variation by mutation points toward variation being maintained by some type of balancing

selection. We do not yet know enough about the natural history of the Mrt phenotype to identify

candidate mechanisms, except to note the close correspondence between genes that produce a

Mrt phenotype and the RNAi mechanism. Natural targets of RNAi include transposable

elements and viruses (Robert et al. 2005; Fischer et al. 2013), each of which could plausibly

constitute an agent of balancing selection.

Acknowledgments

We thank Marie-Anne Félix for her helpful comments, and for piquing our interest in the mortal

germline phenotype. Support was provided by NIH award GM127433 to CFB and V. Katju.

Literature Cited

Ahmed, S., and J. Hodgkin, 2000 MRT-2 checkpoint protein is required for germline immortality and telomere replication in *C. elegans*. *Nature* 403 (6766):159-164.

Baer, C.F., F. Shaw, C. Steding, M. Baumgartner, A. Hawkins *et al.*, 2005 Comparative evolutionary genetics of spontaneous mutations affecting fitness in rhabditid nematodes. *Proceedings of the National Academy of Sciences of the United States of America* 102 (16):5785-5790.

Boyle, E.A., Y.I. Li, and J.K. Pritchard, 2017 An Expanded View of Complex Traits: From Polygenic to Omnigenic. *Cell* 169 (7):1177-1186.

Buckley, B.A., K.B. Burkhart, S.G. Gu, G. Spracklin, A. Kershner *et al.*, 2012 A nuclear Argonaute promotes multigenerational epigenetic inheritance and germline immortality. *Nature* 489 (7416):447-451.

Cook, D.E., S. Zdraljevic, J.P. Roberts, and E.C. Andersen, 2017 CeNDR, the *Caenorhabditis elegans* natural diversity resource. *Nucleic Acids Research* 45 (D1):D650-D657.

Davies, S.K., A.M. Leroi, A. Burt, J. Bundy, and C.F. Baer, 2016 The mutational structure of metabolism in *Caenorhabditis elegans*. *Evolution* 70:2239-2246.

Dirksen, P., S.A. Marsh, I. Braker, N. Heitland, S. Wagner *et al.*, 2016 The native microbiome of the nematode *Caenorhabditis elegans*: gateway to a new host-microbiome model. *Bmc Biology* 14.

Felix, M.A., and F. Duveau, 2012 Population dynamics and habitat sharing of natural populations of *Caenorhabditis elegans* and *C. briggsae*. *Bmc Biology* 10.

Fischer, S.E.J., Q. Pan, P.C. Breen, Y. Qi, Z. Shi *et al.*, 2013 Multiple small RNA pathways regulate the silencing of repeated and foreign genes in *C. elegans*. *Genes & Development* 27 (24):2678-2695.

Frezal, L., E. Demoinet, C. Braendle, E. Miska, and M.A. Felix, 2018 Natural genetic variation in a multigenerational phenotype in *C. elegans*. *Current Biology* 28 (16):2588-+.

Guang, S., A.F. Bochner, D.M. Pavelec, K.B. Burkhart, S. Harding *et al.*, 2008 An Argonaute transports siRNAs from the cytoplasm to the nucleus. *Science* 321 (5888):537-541.

Haldane, J.B.S., 1927 A mathematical theory of natural and artificial selection, Part V: Selection and mutation. *Proceedings of the Cambridge Philosophical Society* 23:838-844.

Houle, D., 1998 How should we explain variation in the genetic variance of traits? *Genetica* 103:241-253.

Houle, D., B. Morikawa, and M. Lynch, 1996 Comparing mutational variabilities. *Genetics* 143 (3):1467-1483.

Katz, D.J., T.M. Edwards, V. Reinke, and W.G. Kelly, 2009 A *C. elegans* LSD1 demethylase contributes to germline Immortality by reprogramming epigenetic memory. *Cell* 137 (2):308-320.

Manolio, T.A., F.S. Collins, N.J. Cox, D.B. Goldstein, L.A. Hindorff *et al.*, 2009 Finding the missing heritability of complex diseases. *Nature* 461 (7265):747-753.

Rajaei, M., A. Shekhar Saxena, L.M. Johnson, M.C. Snyder, T.A. Crombie *et al.*, 2021 Mutability of mononucleotide repeats, not oxidative stress, explains the discrepancy between laboratory-accumulated mutations and the natural allele-frequency spectrum in *C. elegans*. *bioRxiv*:2021.2002.2009.430480.

Robert, V.J.P., T. Sijen, J. van Wolfswinkel, and R.H.A. Plasterk, 2005 Chromatin and RNAi factors protect the *C. elegans* germline against repetitive sequences. *Genes & Development* 19 (7):782-787.

Sakaguchi, A., P. Sarkies, M. Simon, A.-L. Doebley, L.D. Goldstein *et al.*, 2014 *Caenorhabditis elegans* RSD-2 and RSD-6 promote germ cell immortality by maintaining small interfering RNA populations. *Proceedings of the National Academy of Sciences* 111 (41):E4323-E4331.

Saxena, A.S., M.P. Salomon, C. Matsuba, S.D. Yeh, and C.F. Baer, 2019 Evolution of the mutational process under relaxed selection in *Caenorhabditis elegans*. *Molecular Biology and Evolution* 36 (2):239-251.

Schug, M.D., T.F.C. Mackay, and C.F. Aquadro, 1997 Low mutation rates of microsatellite loci in Drosophila melanogaster. *Nature Genetics* 15 (1):99-102.

Schulenburg, H., and M.A. Felix, 2017 The Natural Biotic Environment of *Caenorhabditis elegans*.
Genetics 206 (1):55-86.
Smelick, C., and S. Ahmed, 2005 Achieving immortality in the *C. elegans* germline. *Ageing Research*
Reviews 4 (1):67-82.
Spracklin, G., B. Fields, G. Wan, D. Becker, A. Wallig *et al.*, 2017 The RNAi inheritance machinery of
Caenorhabditis elegans. *Genetics* 206 (3):1403-1416.
Vassilieva, L.L., and M. Lynch, 1999 The rate of spontaneous mutation for life-history traits in
Caenorhabditis elegans. *Genetics* 151 (1):119-129.