Microbial cysteine degradation is a source of hydrogen sulfide in oxic freshwater lakes

Patricia Q. Tran[1,2]*, Samantha C. Bachand[1]*, Jacob C. Hotvedt[1]*, Kristopher Kieft[1,3], Elizabeth A. McDaniel[1,3], Katherine D. McMahon[1,4], Karthik Anantharaman[1+]

1. Department of Bacteriology, University of Wisconsin-Madison, WI, USA
2. Department of Integrative Biology, University of Wisconsin-Madison, WI, USA
3. Microbiology Doctoral Training Program, University of Wisconsin-Madison, WI, USA
4. Department of Civil and Environmental Engineering, University of Wisconsin-Madison, WI, USA

* Patricia Q. Tran, Samantha C. Bachand and Jacob C. Hotvedt are co-first authors.

+ Corresponding author: Karthik Anantharaman (karthik@bact.wisc.edu)

Abstract

The sulfur-containing amino acid cysteine is abundant in the environment including in freshwater lakes. Biological cysteine degradation can result in hydrogen sulfide (H_2S), a toxic and ecologically relevant compound that is a central player in biogeochemical cycling in aquatic environments. Here, we investigated the ecological significance of cysteine in oxic freshwater lake environments, using isolated cultures, controlled growth experiments, and multi-omics. We screened bacterial isolates enriched from natural lake water for their ability to produce H_2S when provided cysteine. In total, we identified 29 isolates that produced H_2S and belonged to the phyla *Bacteroidetes, Proteobacteria,* and *Actinobacteria*. To understand the genomic and genetic basis for cysteine degradation and H_2S production, we further characterized 3 freshwater isolates using whole-genome sequencing (using a combination of short-read and long-read sequencing), and quantitatively tracked cysteine and H_2S levels over their growth ranges: *Stenotrophomonas maltophilia* (Gammaproteobacteria), *Stenotrophomonas bentonitica* (Gammaproteobacteria) and *Chryseobacterium piscium* (Bacteroidetes). We observed a decrease in cysteine and increase in H_2S, and identified genes involved in cysteine degradation in all 3 genomes. Finally, to assess the presence of these organisms and genes in the environment, we surveyed a five-year time series of metagenomic data from the same isolation source (freshwater Lake Mendota, WI, USA) and identified their presence throughout the time series. Overall, our study shows that sulfur-containing amino acids can drive microbial H_2S production in oxic environments. Future considerations of sulfur cycling and biogeochemistry in oxic environments should account for H_2S accumulation from degradation of organosulfur compounds.

Importance

Hydrogen sulfide (H_2S), a naturally occurring gas with biological origins, can be toxic to living organisms. In aquatic environments, H_2S production typically originates from anoxic (lacking oxygen) environments such as sediments, or the bottom layers of thermally stratified lakes. However, the degradation of sulfur-containing amino acids such as cysteine, which all cells and life forms rely on, can be a source of ammonia and H_2S in the environment. Unlike other approaches for biological H_2S production such as dissimilatory sulfate reduction, cysteine degradation can occur in the presence of oxygen. Yet, little is known about how cysteine degradation influences sulfur availability and cycling in freshwater lakes. In our study, we identified diverse bacteria from a freshwater lake that can produce H_2S in the presence of O_2. Our study highlights the ecological importance of oxic H_2S production in natural ecosystems and necessitates a change in our outlook of sulfur biogeochemistry.

Introduction

In most natural environments, hydrogen sulfide gas (H_2S) production is usually attributed to defined groups of bacteria and archaea (1, 2) , and occurs primarily in anoxic environments. During the process of dissimilatory sulfate reduction, sulfate acts as a terminal electron acceptor, and is converted to hydrogen sulfide. However, other pathways for H_2S production exist, namely assimilatory sulfate reduction, in which H_2S contributes to cell growth and increased biomass,

65   and the desulfurylation (desulfurization) of sulfur-containing amino acids such as cysteine which
66   can lead to production of pyruvate, ammonia, and H_2S (3). It is believed that assimilatory sulfate
67   reduction contributes to growth but does not release H_2S from the cell, while dissimilatory
68   sulfate reduction and cysteine degradation can both contribute to growth and release ecologically
69   relevant nitrogen and sulfur compounds into the ecosystem.
70
The sulfur cycle is composed of several assimilatory and dissimilatory pathways, which interact
in complex ways through biotic and abiotic factors. Sulfur cycling in freshwater ecosystems can
be ecologically significant, especially in places where strong redox gradients exist (4). For
example, in high arctic lakes, sulfur-compounds are suggested to serve as biogeochemical hubs
(5). Cysteine, a sulfur containing amino acid, is proposed to be an overlooked source of both
carbon (6) and sulfur. Additionally, seston (particles in water comprised of both living and non-
living organisms) contains organosulfur containing lipids which settle into the sediments, and
contributes to the sulfur pool, even in highly oligotrophic lakes such as Lake Superior (7).
79
In seasonally stratified lakes consisting of oxygenated warm water (epilimnion) floating atop
colder anoxic waters (hypolimnion), H_2S is often abundant in the hypolimnion (8, 9), due to
oxygen demand driving terminal electron acceptor depletion. However, an overlooked player in
the pool of available H_2S are the use of organosulfur compounds, such as cysteine, by microbes.
Cysteine is required to produce many proteins and is also important for protein structure. It is
one of the two amino acids (methionine being the other) that contains a sulfur group; however,
the sulfhydryl group on cysteine is more reactive and can lead to H_2S formation. Like all amino
acids, cysteine also contains an amine group that will form ammonia once the molecule is
degraded. As such, cysteine degradation (desulfurylation) by microbes leads to H_2S production.
H_2S is ecologically relevant because it can be toxic to plants and animals. During periods of
anoxia, H_2S can accumulate to levels beyond the threshold for living organisms, and can cause
massive fish kills (10). Unlike other H_2S sources, cysteine desulfurylation can occur under oxic
conditions, thereby expanding the environmental scope of this sulfur pool. Indeed, cysteine can
be desulfurylated under oxic conditions in the laboratory, but little information exists on the
natural prevalence of this process in lakes. We expect that H_2S production in oxic environments
(during the mixed water column periods of the year, and throughout the stratified period in the
mixed epilimnion) could result from cysteine breakdown by microbes.
97
In this study, we investigated the prevalence of organosulfur degradation (desulfurylation) in a
freshwater lake, using both laboratory and genomic evidence, to advance our understanding of
oxic sulfur cycling in aquatic ecosystems (**Figure 1**). First, we grew bacterial isolates enriched
from Lake Mendota's oxic epilimnion to quantify H_2S and ammonia production, which informs
the potential for organosulfur degradation in an oxygenated aquatic environment. We found 18
isolates producing H_2S under oxygenic conditions. We selected three H_2S-producing isolates for
detailed characterization using full-genome sequencing and chemical analyses to track cysteine
concentrations and H_2S accumulation during their growth: *Stenotrophomonas maltophilia*
(Gammaproteobacteria), *Stenotrophomonas S. bentonitica* (Gammaproteobacteria) and
Chryseobacterium piscium (Bacteroidetes). In all three isolates, cysteine decreased and H_2S
increased over their exponential growth curve under oxic conditions. Finally, we contextualized
our laboratory results using a time-series of metagenomic data from the same isolation source
(Lake Mendota, WI, USA), to study the temporal importance of organosulfur degradation. We

found that genes for cysteine desulfurylation were present and abundant throughout the time-series suggesting that the ability to degrade cysteine is well represented in Lake Mendota.

Figure 1. Overview of methods used in this study. Isolates were enriched from oxygenated lake water. Isolates were screened for H_2S and ammonia production, and those that produced H_2S were selected for 16S rRNA sequencing. Then based on the results, three isolates were selected for whole-genome sequencing using a combination of short and long-reads. Genome characterization of functional potential and taxonomic classification was conducted on the isolates. Screening of genes involved in cysteine degradation was conducted in the isolates, and a five-year metagenomic time series of Lake Mendota (2008-2012).

Results

Isolates capable of H$_2$S production in oxic conditions

To answer the question of whether bacteria could produce H$_2$S in the presence of oxygen, we grew pure culture isolates originally recovered from the water column of temperate eutrophic Lake Mendota. We grew the isolates in moderately rich media (R2A, see methods) under control and treatment conditions (cysteine addition) and tracked H$_2$S production after 24 hours (**Figure S1, Table S1**). Using qualitative H$_2$S measurements, we found that 18 isolates produced H$_2$S and ammonia when grown in the presence of amended cysteine. We performed 16S rRNA gene sequencing on the 29 isolates that produced H$_2$S, regardless of whether they produced ammonia or not when amended with cysteine. Isolates that produced both H$_2$S and ammonia were identified as *Stenotrophomonas rhizophila* (Betaproteobacteria), *Stenotrophomonas maltophilia* (Betaproteobacteria), *Citrobacteria gillenii* (Gammaproteobacteria), and *Chryseobacterium sp.* (Bacteroidetes), whereas those producing H$_2$S but not ammonia were identified as *Pseudomonas arsenicoxydans, Pseudomonas mandelii, Pseudomonas migulae, Pseudomonas thivervalensis,* and *Microbacterium flavescens*.

Detailed microbiological, chemical and genomic characterization in selected isolates

Next, we selected 3 isolates (#43, #13 and #66) representing distinct species based on 16S rRNA sequence taxonomy (97% identity), that produced H$_2$S for further characterization. These detailed characterizations include OD$_{600}$-based growth rates and paired quantitative measurements of cysteine and H$_2$S concentrations in the spent medium. Cysteine addition resulted in concommital H$_2$S production over time (**Figure 2A, Table S2**). All organisms used L-cysteine preferably to D-cysteine (**Figure 2B, Table S3**).

150
Figure 2. Further characterization of the three isolates and demonstration of cysteine
degradation and H₂S production. A. Higher amounts of H$_2$S was produced by the three isolates
over 50 hours compared to negative controls. B, C, D. Identification of different forms of
cysteine that can be degraded. L-cysteine decreased in all isolates compared to the control (B,
D). D-cysteine also decreased over time in all samples except the negative control, however, the
net amount decreased was less compared to L-cysteine. Cysteine concentrations were measured
as cystine as described in the methods. Because isolates #13 and #43 were assessed in a different
experimental run than #66, but using the exact same instruments and methods, plots B, C and D
are separated by HPLC runs. Due to large sample volume it was not possible to test all isolates
and conditions in one HPLC run.

161
Next, we performed whole-genome sequencing using combined short-read and long-read
sequencing on these 3 isolates (**Table 1**). We performed whole-genome sequencing because
functional information such as gene content cannot be predicted reliably from 16S rRNA
sequencing alone. The full genome of Isolate 43 was assembled into a single circular genome,
and taxonomically assigned to *Stenotrophomonas maltophilia*. Unlike the 16S rRNA sequence
which assigned it to *S. rhizophila*, the full-genome was actually closer to *S. maltophilia*. The full
genome of isolate #13, could be assembled into 2 long contigs, and was taxonomically assigned
to *S. bentonitica*. The *Chryseobacter* genome was assembled into one circular genome, and

assigned to *Chryseobacterium piscium*. All three genomes were estimated to be 100% complete based on CheckM. Overall, the 16S rRNA gene amplicon sequencing performed prior agreed with full-genome sequencing assignment in some cases, and in others, the whole-genome sequencing assignment allowed finer taxonomic resolution (such as in the case of Isolate 13). Overall, whole-genome sequencing provided more information about the isolates' metabolic potential. The genomic content was then used to inform how or why H_2S might be produced in oxic environments, as shown in the laboratory experiments.

Table 1. Genome characteristics of 3 selected isolates that produce H_2S in the presence of oxygen and cysteine.

Name	Genome Size (bp)	Completeness[1]	GC content	Taxonomy[2]	Number of contigs
13-LM-B-02-08 (referred to as #13)	4,188,104	100%	66.8%	d__Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__Xanthomonadales;f__Xanthomonadaceae;g__Stenotrophomonas;**s__Stenotrophomonas maltophilia_O**	1
43-LM-B-01-03 (referred to as #43)	4,325,715	100%	66.5%	d__Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__Xanthomonadales;f__Xanthomonadaceae;g__Stenotrophomonas;**s__Stenotrophomonas bentonitica**	2
LM_BA_5.2 (referred to as #66)	1,375,102	100%	33.7%	d__Bacteria;p__Bacteroidota;c__Bacteroidia;o__Flavobacteriales;f__Weeksellaceae;g__Chryseobacterium;**s__Chryseobacterium piscium**	7

[1] Using CheckM (See Methods)   [2] Using GTDB-tk (See Methods)

We used gene annotations of the 3 isolates to infer the presence of genes involved in cysteine metabolism. We identified genes involved in cysteine degradation to ammonia, pyruvate, and H_2S: *metC*, *malY*, *tnaA*, *cysM*, *cysK* (which involve the use of L-cysteine as the substrate), and *dcyD* (which involve the use of D-cysteine as the substrate) (**Table S4**). However, we note that these genes may have other enzymatic activities, such as cysteine biosynthesis instead of degradation (**Table S5**). (**Figure 3**)

Figure 3. Diagram showing pathways for sulfur and organosulfur metabolism. Several
pathways for hydrogen sulfide and cysteine production exist in microorganisms. The
presence/absence of key genes (*cysK*, *cysM*, *malY*, *metC*, *tnaA*, *sseA*, *aspB*, and *dcyD* along the
blue arrows) in the three isolate genomes are shown by filled (present) circles, and unfilled (not
present) circles.

Leveraging the full-genomic content of the 3 isolates (**Table S6**), we proposed a joint cellular
map based on identified metabolic functions and pathways in the genomes (**Figure 4A**). All
three isolates contained pathways associated with central carbon metabolism: including the TCA
cycle, glycolysis, gluconeogenesis, the pentose phosphate pathway and the glyoxylate cycle.
They could all generate fatty acids using fatty acid biosynthesis, and catabolize fatty acids
through the beta-oxidation pathway. As expected, they had genes for cysteine metabolism
including *metC*, *malY*, *tnaA*, *cysM*, *cysK*, and *dcyD*, cysteine biosynthesis pathways from
homocysteine and serine, as well pathways for degradation of other amino acids including
methionine.

Despite these similarities, the three isolates also have distinguishing characteristics amongst
them (**Table S6, Figure 4B**). For example, while all isolates encoded genes for sulfur oxidation

208 (sulfur dioxygenase), genes for thiosulfate oxidation were present in the two *Stenotrophomonas*
209 isolates but not *Chryseobacterium*. The *Chryseobacterium* isolate encoded for a urease
210 suggesting the use of organic nitrogen in the form of urea but this was absent in the two
211 *Stenotrophomonas* isolates. Finally, genes for sugar utilization were identified in the two
212 *Stenotrophomonas* isolates but not in *Chryseobacterium*.
213

214
215 **Figure 4. A. Cellular map showing important metabolic pathways and transporters which**
216 **were common to all three isolates.** A complete list is found in **Table S6**. The KEGG module
217 identifiers are listed in purple whenever relevant. **B. Heat map showing selected metabolic**
218 **functions and pathways in the three isolates.** A complete list is found in **Table S6**.

Presence of cysteine-degrading organisms and genes in a five-year metagenomic environmental time-series

To put these laboratory results and lab-grown organisms into a natural environment context, we leveraged a previously published metagenomic time-series collected from the oxygenated upper mixed layer of Lake Mendota spanning 2008-2012 (97 time points) to search for the presence of cysteine degradation genes in the microbial communities.

First, we searched the time-series to see if organisms in our study were also present in the time series. To do this, we linked the 16S rRNA gene sequences of the isolated organisms to the assembled metagenomes (i.e. contigs) from the time series. We found that while the 16S rRNA sequences were also present in the time series (**Table S7, S8, S9**), and broadly distributed over time, these scaffolds were not part of binned genomes. Therefore, little information about these isolates would be gathered from metagenomic data only. As such, the full-genome sequencing we performed was particularly helpful in understanding the full genomic structure of the H_2S-producing organisms.

Second, we searched for the 6 genes associated with cysteine degradation and H_2S production (**Table S5**) (in binned and unbinned contigs) using KEGG HMMs with hmmsearch. In total, we searched over 22 million amino acid sequences and identified 1882 hits to the 5 marker genes found in the isolate metagenomes, *dcyD* was not found (**Figure 5, Table S10**). *cysK* and *malY* were the genes with the most corresponding matches at any time point, followed by *metC* and *cysM*. Only 2 scaffolds contained *tnaA*. Overall, after correcting for genome size, there was no obvious temporal trend of the genes, although genes were found throughout the 5-year time series.

Among these cysteine-degrading gene sequences, several were identified in binned metagenome-assembled genomes (MAGs) (**Figure 5B, Table S11**), which allowed for the assignment of taxonomy. Overall, we identified 139 genes to be distributed in genomes of organisms from Actinobacteria, Bacteroidetes, Chloroflexota, Cyanobacteria, Planctomycetes, Proteobacteria, and Verrucomicrobia, representing common freshwater lineages (11). The *tnaA* gene was only present in Bacteroidetes, but other genes were more broadly distributed among taxonomic groups.

253 **Figure 5. A. Counts of cysteine desulfurylation genes (5 genes) in a 5-year time-series of Lake**
254 **Mendota.** We also searched for dcyD (6th gene) but did not identify it in any sample. **B. Presence of**
255 **these 5 genes in metagenome-assembled metagenomes from the same time series.**

Discussion

Different types of H$_2$S production: the fate of cysteine and the origin of the H$_2$S

The H$_2$S-producing isolates fell into two groups when grown in media with cysteine: ammonia-
producing and ammonia-consuming. We hypothesize that those able to produce ammonia and
H$_2$S in the presence of cysteine, but not under "control" conditions were those that were
potentially contributing to the H$_2$S pool in the lake. However, it is possible that the isolates that
consume ammonia also conduct the cysteine degradation pathway, but do not excrete ammonia
from the cell. Instead, since ammonia is an important metabolic precursor, it could be used by
the organism instead of being released into the media, making it undetectable with our analytical
technique.

We do not believe that this occurs with H$_2$S, and that most, if not all, of the isolates that produce
H$_2$S in the presence of oxygen will excrete it from the cell. H$_2$S is toxic to organisms that
undergo aerobic cellular respiration (12). One potential reason to keep H$_2$S in the cell is that
some bacteria appear to use H$_2$S as a protective compound against antibiotics, but H$_2$S
accumulation also creates a large amount of stress on the cell making it useful only in extreme
situations (13). While it is possible that organisms may use H$_2$S internally as a sulfur source, we
did not identify any genes encoding sulfide quinone oxidoreductases, flavocytochrome c

dehydrogenases, or other genes for the oxidation or transformation of H_2S. All three isolates
were obligate aerobes based on laboratory assays.
276
Of the 20 amino acids, little is known about cysteine in the environment. One of the difficulties
in studying the fate of cysteine in oxic environments is that it can be abiotically and
spontaneously oxidized into cystine (14), which *Escherichia coli* has been shown to uptake (15).
In a study of *E.coli* K-12 that lacked a cysteine transporter, cysteine could enter the cell through
transporters dedicated to other amino acids, when no amino acids alternatives other than cysteine
were present in the medium (16). It is likely that the majority of H_2S produced by the cell in the
three isolates originates from cysteine given the demonstrated reduction in cystine concentrations
when the isolates are grown with added cysteine (**Figure 2**), coupled with the accumulation of
H_2S and ammonia.

Genomic structure of the H_2S-producing isolates

287
288 Overall, the three isolates selected for whole genome sequencing revealed genes for cysteine
289 degradation into H_2S. Based on laboratory studies, they were able to produce H_2S in the presence
290 of oxygen. The isolates were obligate aerobes, which presents interesting questions about the life
291 history of these organisms.
292
Little information is known about the ecology of *Stenotrophomonas maltophilia*,
Stenotrophomonas bentonitica, and *Chryseobacterium piscium* in the natural environment. *S.*
maltophilia is a cosmopolitan bacterium in nature, and found in a range of natural environments,
particularly in association with plants(17). *S. bentonitica* was originally characterized in
bentonite formations, was predicted to have high tolerance to heavy metals (18), and has been
observed in arctic seawater (19). *C. piscium* was isolated from a fish in the arctic ocean (20), but
its ecological significance in the oceans remains unknown. This previously described *C. piscium*
strain LMG 23089 was not reported to produce H_2S yet our genetic analyses suggest that it has
the enzymatic machinery to degrade cysteine.
302
303 One possible explanation for this discrepancy is that LMG 23089 was previously grown on SIM
304 medium to test H_2S, which is lower resolution than the modern H_2S probes which measure μM
305 concentrations, and because SIM medium uses thiosulfate as a sulfur source. As a side test on
306 isolate #66, H_2S was not produced when thiosulfate was provided, but H_2S was produced when
307 cysteine was provided.
308
One particular finding of this study was that none of the 6 genes searched for cysteine
degradation into H_2S and ammonia was common to *all* three isolates, despite all 3 isolates
showing the same cysteine-decrease, ammonia-increase, and H_2S-increase over time. This could
be explained by alternative, perhaps less straightforward pathways for H_2S production. One
pathway is led by a gene named cystathionine gamma-lyase ("CTH" or "CSE"). In some bacteria
and mammals, this enzyme is involved in H_2S production (21). An HMM search for this enzyme
showed that it was present in Isolate #13, 43 and #66. While it was not initially included in the
initial methods and study, this could hint to another commonality among oxic H_2S producing
organisms.

Challenges associated with measuring oxic H₂S production from organosulfur in the environment

Extrapolating these laboratory results to widespread distribution of organosulfur degradation in the natural environment necessitates several steps, namely because of the major knowledge gaps that exist concerning the sulfur cycle in freshwater lakes, and because bridging the gap between cultivation-based, omics-based (22), and field-based experiments is needed. Foremost, the identity, distribution, and availability of organosulfur compounds broadly across lakes globally is currently mostly unknown. Cysteine is notoriously difficult to measure, and many previous studies characterizing the amino acid composition of the water column only measure the sulfur-containing organosulfur compound taurine (23, 24).

Organic sulfur in the form of cysteine is an important organosulfur amino acid, and is important in protein folding and function (25). As such, there is a difference in the fates of cysteine when it exists bound in cell walls, versus when cysteine is free in the water column and available for degradation by bacteria. While cysteine has been shown to contribute to the carbon pool and carbon flow in lakes (6), more quantitative field measurements are necessary to support whether cysteine also serves as a sulfur pool. Yet, other forms of organosulfur have important significance in aquatic environments. In marine environments, for example, DMSP (dimethylsulfoxonium propionate) is a critical component of the marine organosulfur cycle (26).

Additionally, current differences between computational gene similarity searches versus *in vivo* enzymatic functions are challenging to assess for the genes responsible for the cysteine degradation into pyruvate, ammonia, and H₂S. One reason is that the enzymatic activity of the gene has mostly been described in model organisms such as *E. coli*, and it has been shown that gene expression can be induced by genetic factors, or environmental factors such as metals (27). At least 6 genes have been proposed to have this enzymatic activity, yet each gene may serve different functions *in situ*, and it is difficult to assert directionality of enzymatic function based on metagenomic or genomic analyses only. To this end, the isolated bacterial strains from this study, which are non-model organisms, and originate from the natural freshwater lake environment, may be used for further detailed biochemical, physiological, and microbiological studies. Further characterization of these bacterial isolates using gene knockout, gene induction, or heterologous gene expression studies may inform the functional activity of these genes in nature.

Implications of oxic H₂S production by microbes in freshwaters

This study demonstrates that H₂S production by microbes in lake ecosystems occurs in the presence of oxygen, using evidence from pure culture bacterial isolates, and screening of long-term metagenomic time-series. By combining lake-to-laboratory experiments, we show that multiple bacterial strains spanning Gammaproteobacteria, Betaproteobacteria, Actinobacteria, and Bacteroidetes are all able to produce H₂S under oxic conditions, and at temperatures that would be ecologically relevant for surface lake water during the summer. Surface water temperatures in Lake Mendota can reach up to 27°C, and the top few meters of water surface are

362 saturated in oxygen. Worldwide, maximum lake surface temperature can range between 23 to
363 31°C (28).
364
Unlike dissimilatory sulfate reduction, cysteine use by bacteria to generate ammonia, pyruvate,
and H_2S, is not dependent on sulfate as an initial reaction substrate. Increased sulfate
concentrations are shown to lead to higher sulfate reduction rates in shallow eutrophic
freshwater, the sulfur originating from algal decay for example (29). While Lake Mendota is a
low-iron and high-sulfate lake (30), not all lakes have elevated sulfate levels, and therefore, H_2S
production might previously not have been thought of as relevant to study. However, sulfur-
containing amino-acids can have many origins. In lakes, concentrations of amino acids (free
dissolved and combined) often reflect the input and outputs of the lake (31, 32). For example,
amino acids contributed a detectable amount to the nitrogen cycle, and bacterial utilization of
amino acids contributes to nitrogen pool and cycling in natural ecosystems, (32) although
cysteine amino acids were not measured in that study.
376
Freshwater lakes that are dimictic can become stratified in temperature and oxygen during the
summer, and oxygen concentrations vary throughout the year. In the fall and spring, oxygen is
abundant, and cysteine degradation into H_2S could be a relevant process for the sulfur pool, and
H_2S fluxes to the atmosphere could be significant since wind is prevalent. Under ice during the
winter, where oxygen is plentiful, H_2S could be produced but could be consumed or oxidized. On
the other hand, gases would be trapped under ice. During summer, the anoxic hypolimnion and
sediments are known H_2S sources due to dissimilatory sulfate reduction, but density gradients
would prevent H_2S from reaching the atmosphere. However, the oxygenated mixed epilimnion
could be an H_2S source through organosulfur degradation. If we consider the importance of oxic
hydrogen sulfide production, which could occur year-round, the H_2S pool and the scope of sulfur
transformations may be greater than anticipated, if we focus solely on the anoxic hypolimnion
(**Figure 6**). Future work aiming to understand the broader distribution of sulfur-containing amino
acids and other organosulfur compounds in freshwaters, their fates and transformations, as well
as their contribution to H_2S production, will inform global sulfur biogeochemical cycling.
391

392
Figure 6. Conceptual figure showing the potential relevance of cysteine degradation in a
freshwater lake environment, with respect to oxygen availability and seasonality.
Oxygenated seasons and part of the lake water columns are shown with an asterisk. Significant
research gaps include cysteine concentrations in the natural environment over time, hypothesized
H_2S fluxes across different layers in the lake water column, and contribution of different H_2S
sources in the hypolimnion. In all season, portions of the water column can be oxygenated.

Methods

Enrichment cultures of isolates from a temperate freshwater lake

Lake Mendota (43°06′24″ N 89°25′29″ W) is a temperate eutrophic lake in South-Central
Wisconsin, in Madison, WI, USA. Lake Mendota is part of the Long-Term Ecological Research
Network North Temperate Lakes (NTL-LTER, https://lter.limnology.wisc.edu/about/lakes).
Lake Mendota encounters annual stratification and annual seasonal anoxia in the hypolimnion.
Lake water was collected on September 14, 2018 from an integrated water sample (0m to 12m)
from Lake Mendota at the "Deep Hole" station (43°05'54", 89°24'28"), where the maximum
depth is 23.5m. The water samples were collected during stratification from the oxygenated
epilimnion. The lake water was collected in pre-acid washed 2 L sampling bottles using a
flexible PVC tube, and brought back on shore within hours for immediate processing. Serial
dilution was performed and lake bacteria were grown on PCB (plate count media broth) agar
media, at room temperature (~21°C), in the lab in the light. The PCB media was made of: 1 L

water, 5g/L of yeast extract, 10g/L of tryptone and 2g/L of dextrose/D-glucose. If grown on solid
media, 10g of agar per 1L media was added. Enrichment resulted in about 60 isolates.
414

Screening for cysteine degradation into H₂S and ammonia

416 Isolates were able to grow on PCB and R2A media. R2A media is a culture medium for bacteria
417 that typically grow in freshwater. It is less "nutrient rich" than PCB media, and therefore slightly
418 closer to natural lake water. For the screening of the isolates for H_2S production, we grew them
419 on R2A media. Each isolate had two treatments: grown in R2A media without cysteine for the
420 control, and grown in R2A media with amended cysteine as the treatment.
421
422 R2A media consisted of (1 L of water): 0.5g of casein, 0.5g of dextrose, 0.5g of starch, 0.5g of
423 yeast extract, 0.3g of K_2HPO_4, 0.3g of sodium pyruvate, 0.25g of peptone, 0.25g of beef extract,
424 0.024g of $MgSO_4$, and autoclave. To make the same media for plates, we added 15g of agar
425 before autoclaving. For controls, isolates were grown in the media without cysteine amendments.
426 For "treatments", 2mM cysteine was added.
427
To assess the amount of cysteine degradation into H_2S and ammonia, we screened each of the 60
isolates for H_2S and/or ammonia accumulation. To test H_2S production, we grew the strains
individually in liquid media, and used lead acetate test strips (Fisher Scientific, USA) to
qualitatively assess H_2S accumulation in the headspace. A darkening of the strip shows that H_2S
was produced. To test ammonia concentrations after 24 hours, we measured samples at time zero
and 24 hours using Ammonia Salicylate Reagent Powder Pillows and Ammonia Cyanurate
Reagent Powder Pillows (Hatch Reagents) and used spectrophotometry at the 655nm
wavelength.
436

Identification of H₂S producing bacteria using 16S rRNA sequencing

Colony PCR and DNA extractions were conducted using the EtNa Crude DNA Extraction and
ExoSAP-ITTM PCR Product Cleanup protocols on the isolates that tested positive for producing
H_2S (10). Full length 16S rRNA gene products were generated for sequencing using universal
16S rRNA primers (27f, 1492r)(33).DNA concentration yields were measured using the QuBit
dsDNA HS assay kit (QuBit). DNA was sequenced at the University of Wisconsin-Madison
Biotechnology Center (Madison, WI, USA) The program 4Peaks (34) was used to clean the base
pairs by quality-checking followed by homology search using BLASTn against the NCBI
Genbank database (accessed December 2019) (35) to identify the sequences.

Detailed characterization of 3 hydrogen-sulfide producing isolates

We selected 3 isolates that could aerobically produce H_2S for further detailed characterization.
We selected these isolates because some of the 18 isolates that produced H_2S when grown with
cysteine had identical 16S rRNA sequences, therefore we chose isolates that had distinct 16S
rRNA sequences for full-genome sequencing. Additionally, using 16S rRNA gene sequencing of
the isolates, one was only assigned to *Stenotrophomonas sp.*, and we believed that whole-

genome sequencing would enable us to get a higher taxonomic confirmation and more complete
information.
454
We performed DNA extraction using the PowerSoil Powerlyzer kit (Qiagen) without protocol
modifications, and sent the genomic DNA for full genome sequencing at the Microbial Genome
Sequencing Center (MIGS) (Pittsburg, PA) with combined short read Illumina and long read
nanopore sequencing. The data was processed by MIGS to assemble the short-reads (Illumina
Next Seq 2000) and long-reads (Oxford Nanopore Technologies) into full-genomes. Quality
control and adapter trimming was performed with bcl2fastq (Illumina) and porechop
(https://github.com/rrwick/Porechop) for Illumina and Oxford Nanopore Technologies (ONT)
MinION sequencing respectively. Hybrid assembly with Illumina and ONT reads was performed
with Unicycler (36). Genome annotation of the 3 isolates was done with Prokka v.1.14.5(37),
using the --rfam setting.
465
Genome completeness and contamination were estimated using CheckM v.1.1.3 (38)
lineage_wf. Taxonomic classification was conducted using GTDB-tk v.0.3.2 (39) with the
database release r95. The full-genome taxonomic classification agreed with the prior 16S rRNA
sequencing results, but we were further able to identify Isolate 43 as *Stenotrophomonas*
bentonitica. We ran METABOLIC-G v.4.0 (40) to identify genes associated with cysteine
degradation and other metabolic pathways.
472
473 Growth measurements of the three isolates were measured using OD_{600} with a
474 spectrophotometer, with measurements every 1 hour. The isolates were grown in R2A broth
475 media, shaken in an incubator at 27°C. Aliquots were collected over the growth range for
476 cysteine measurements across the growth curves, as described below. A H_2S microsensor
477 (Unisense) was used to measure H_2S over time.
478

479 **Methods to measure Cysteine**

480
Cysteine concentrations were measured as cystine, as described in (41) (https://osf.io/9k8a6/).
One of the reasons for measuring cystine instead of cysteine is that in oxic environments,
cysteine is oxidized rapidly into cystine (14, 15). Additionally, unless LC-MS is used, cysteine
can be difficult to measure directly. Samples were diluted 5:4:1 Sample:DI H_2O:DMSO and left
at room temperature for at least 24 hours. Chromatographic analysis was performed on an
Agilent 1260 Infinity II with an Agilent Zorbax Eclipse Plus C18 RRHT 4.6x50mm, 1.8µm, with
Guard column. Column temperature was maintained at 40°C using an Agilent 1260 TCC
(G1316A).
489
Gradient elution was performed using Mobile phase A (MPA) consisting of 10mM Na_2HPO_4,
10mM Sodium tetraborate decahydrate, in DI H_2O, adjusted to pH 8.2 with HCl, filtered to
0.45µm. Mobile phase B (MPB) consisted of 45:45:10 Acetonitrile:Methanol:DI H_2O. Gradient
used for elution was as follows: 0 minutes 98% MPA, 2% MPB; 0.2 minutes 98% MPA, 2%
MPB; 6.67 minutes 46% MPA, 54% MPB; 6.77 minutes 0% MPA, 100% MPB; 7.3 minutes 0%
MPA, 100% MPB; 7.4 minutes 98% MPA, 2% MPB; 8 minutes 98% MPA, 2% MPB. Flow rate
was 2.0mL/min. The pump used was an Agilent Infinity Series G1311B quat pump. Pre-column

derivatization was performed using an Agilent 1260 ALS (G1329B) with an injector program.
Detection was performed using an Agilent 1260 Infinity II MWD (G7165A) at 338nm with
10nm bandwidth. Reference was 390nm with 20nm bandwidth. Recovery was tested during
method development. Recoveries of cystine ranged from 87.2-101.5%, with an average of
92.1%.

Methods to measure H$_2$S using a microsensor

503 Aliquots of at least 1 mL were taken from cultures at desired times after inoculation. H$_2$S
504 concentrations were measured by suspending the H$_2$S probe (Unisense) in the aliquot and leaving
505 it in place until the measurement stabilized. Measurements were manually edited to exclude data
506 gathered while the probe was stabilizing in the sample.
507

Generation of Metagenome-assembled genomes (MAGs)

Sequencing of the Lake Mendota time series for 2008-2012 was previously conducted at the
Joint Genome Institute (42), containing 97 time points (and therefore 97 metagenomic
datasets)(43). In summary, raw reads were quality filtered using fastp (44), and individually
assembled using metaSPAdes(45). Each metagenome was reciprocally mapped to each
individual assembly using BBMap v38.07 (46) with 95% sequence identity cutoff. Differential
coverage mapping to all samples was used to bin contigs into metagenome-assembled genomes
(MAGs) using Metabat2 v.2.12.1 (47). Bins were quality assessed with CheckM v.1.1.2 (38),
dereplicated with dRep v.2.4.2 (48), and classified with GTDB-tk v.0.3.2 (49) with default
settings. This resulted in a total of 116 MAGs from Lake Mendota (**Table S12**) which are
available for download at the Open Science Framework (https://osf.io/qkt9m/).

Searching for cysteine genes and isolates presence in metagenomic time-series and MAGs
520

Genes for cysteine degradation were identified using HMMsearch v3.1b2 (50). HMMs were
downloaded from KoFam (51), accessed May 2020). The KO numbers for the six cysteine
degradation genes are: *metC* (K01760), *cysK* (K01738), *cysM* (K12339), *malY* (K14155), *tnaA*
(K01667), and *dcyD* (K05396) (**Table S4**). Our HMM files are available in **Supplementary File
1,** but are the same as published by KOfam, with the modification of manual addition of the TC
thresholds. HMM-based homology searches were conducted on the 97 Lake Mendota
metagenomes assemblies as described above.
528

Data availability

The 16S rRNA sequences for the 29 H$_2$S producing isolates, the full genome sequences
(nucleotides and amino acids) for isolates #13, #43 and #66 are available on OSF:
https://osf.io/g25eq/ during the peer-review process. The sequences will be deposited in NCBI
prior to publication. The 97 metagenomes were previously published in (43) and are available
through JGI's IMG/M and Genome Portal. The MAGs generated in this study can be found on
https://osf.io/qkt9m/

Acknowledgements

We are thankful to Anna Schmidt for collecting the original lake water from Lake Mendota in 2018, and to Adam Breister and Elizabeth Zanetakos who enriched the bacterial isolates during summer 2018. We thank Trina McMahon's Lab and the Long-Term Ecological Research Network, and the Center for Limnology for their field support and prior work on Lake Mendota. We are thankful for the University of Wisconsin's Water Science and Engineering Laboratory for the use of their HPLC instrumentation, and James Lazarcik for training and assistance with the instruments. We would like to also thank MIGS (the Microbial Genome Sequencing) center (PA, USA) for their assistance with sequencing and long-read data processing.

This work was supported by the USDA National Institute of Food and Agriculture (NIFA) under grant: Hatch project 1025641. Patricia Tran and Kristopher Kieft received the support from the Anna Grant Birge Memorial Award from the Center for Limnology for support for the project in 2019. Patricia Tran is supported by the Natural Science and Engineering Research Council (NSERC) of Canada Doctoral Fellowship. Samantha Bachand was supported by the National Science Foundation (NSF) Research Experience Undergraduate (REU) Award, and the University of Wisconsin-Madison' Holstrom Environmental Research Fellowship. Kristopher Kieft was supported by a Wisconsin Distinguished Graduate Fellowship Award from the University of Wisconsin-Madison, and a William H. Peterson Fellowship Award from the Department of Bacteriology, University of Wisconsin-Madison. Elizabeth McDaniel was supported by a fellowship through the Department of Bacteriology at the University of Wisconsin - Madison. This research was performed in part using the Wisconsin Energy Institute computing cluster, which is supported by the Great Lakes Bioenergy Research Center as part of the U.S. Department of Energy Office of Science. We thank the U.S. Department of Energy Joint Genome Institute for sequencing and assembly (CSP 394) of the Lake Mendota metagenomes.

Contributions

P.Q.T, S.C.B., J.C.H, K.K, and K.A contributed to study design and conceptualization. P.Q.T, S.C.B, J.C.H and K.K conducted experiments on the isolates. J.C.H performed the chemical analyses of the isolates. P.Q.T, S.C.B and J.C.H conducted genomic analyses. P.Q.T, S.C.B and E.A.M conducted metagenomic analyses. P.Q.T, S.C.B and J.C.H analyzed the data and generated figures. P.Q.T, S.C.B, J.C.H and K.A drafted and edited the manuscript. All authors provided feedback and suggestions.

References

1. Foti M, Sorokin DY, Lomans B, Mussman M, Zacharova EE, Pimenov NV, Kuenen JG, Muyzer G. 2007. Diversity, Activity, and Abundance of Sulfate-Reducing Bacteria in Saline and Hypersaline Soda Lakes. Appl Environ Microbiol 73:2093–2100.

2. INGVORSEN K, BROCK T. 1982. ELECTRON FLOW VIA SULFATE REDUCTION AND METHANOGENESIS IN THE ANAEROBIC HYPOLIMNION OF LAKE MENDOTA. LIMNOLOGY AND OCEANOGRAPHY 27:559–564.

3. Tchong S-I, Xu H, White RH. 2005. L-cysteine desulfidase: an [4Fe-4S] enzyme isolated from Methanocaldococcus jannaschii that catalyzes the breakdown of L-cysteine into pyruvate, ammonia, and sulfide. Biochemistry 44:1659–1670.

4. Berg JS, Jézéquel D, Duverger A, Lamy D, Laberty-Robert C, Miot J. 2019. Microbial diversity involved in iron and cryptic sulfur cycling in the ferruginous, low-sulfate waters of Lake Pavin. PLOS ONE 14:e0212787.

5. Vigneron A, Cruaud P, Culley AI, Couture R-M, Lovejoy C, Vincent WF. 2021. Genomic evidence for sulfur intermediates as new biogeochemical hubs in a model aquatic microbial ecosystem. Microbiome 9:46.

6. Göbbels L, Poehlein A, Dumnitch A, Egelkamp R, Kröger C, Haerdter J, Hackl T, Feld A, Weller H, Daniel R, Streit WR, Schoelmerich MC. 2021. Cysteine: an overlooked energy and carbon source. Sci Rep 11:2139.

7. Petersen M, Schreiner KM, Hyde E, Sheik C, Katsev S. 2019. Characterization of Organic Sulfur in Lake Superior, and its Contribution to Sulfur Biogeochemical Cycling 2019:B11L-2247.

8.   Eckert W, Conrad R. 2007. Sulfide and Methane Evolution in the Hypolimnion of a
Subtropical Lake: A Three-Year Study. Biogeochemistry 82:67–76.

9.   Peterson BD, McDaniel EA, Schmidt AG, Lepak RF, Janssen SE, Tran PQ, Marick RA,
Ogorek JM, DeWild JF, Krabbenhoft DP, McMahon KD. 2020. Mercury Methylation Genes
Identified across Diverse Anaerobic Microbial Guilds in a Eutrophic Sulfate-Enriched Lake.
Environ Sci Technol 54:15840–15851.

10. Reese BK, Anderson MA, Amrhein C. 2008. Hydrogen sulfide production and volatilization
in a polymictic eutrophic saline lake, Salton Sea, California. Sci Total Environ 406:205–
218.

11. Newton RJ, Jones SE, Eiler A, McMahon KD, Bertilsson S. 2011. A Guide to the Natural
History of Freshwater Lake Bacteria. Microbiol Mol Biol Rev 75:14.

12. Kabil O, Banerjee R. 2010. Redox Biochemistry of Hydrogen Sulfide. J Biol Chem
285:21903–21907.

13. Pal VK, Bandyopadhyay P, Singh A. 2018. Hydrogen Sulfide in Physiology and
Pathogenesis of Bacteria and Viruses. IUBMB Life 70:393–410.

14. Bagiyan GA, Koroleva IK, Soroka NV, Ufimtsev AV. 2003. Oxidation of thiol compounds by
molecular oxygen in aqueous solutions. Russian Chemical Bulletin 52:1135–1141.

15. Korshunov S, Imlay KRC, Imlay JA. 2020. Cystine import is a valuable but risky process
whose hazards Escherichia coli minimizes by inducing a cysteine exporter. Molecular
Microbiology 113:22–39.

16. Zhou Y, Imlay JA. Escherichia coli K-12 Lacks a High-Affinity Assimilatory Cysteine
Importer. mBio 11:e01073-20.

17.   An S, Berg G. 2018. Stenotrophomonas maltophilia. Trends in Microbiology 26:637–638.

18.   Chen M-X, Li H-Y, Ye X-S, He X-Y. Draft Genome Sequence of an Extracellular Protease-

617   Producing Bacterium, Stenotrophomonas bentonitica VV6, Isolated from Arctic Seawater.

618   Genome Announcements 6:e01610-17.

619   19.   Sánchez-Castro I, Bakkali M, Merroun ML. Draft Genome Sequence of Stenotrophomonas

620   bentonitica BII-R7T, a Selenite-Reducing Bacterium Isolated from Spanish Bentonites.

621   Genome Announcements 5:e00719-17.

622   20.   de Beer H, Hugo CJ, Jooste PJ, Vancanneyt M, Coenye T, Vandamme P 2006.

623   Chryseobacterium piscium sp. nov., isolated from fish of the South Atlantic Ocean off

624   South Africa. International Journal of Systematic and Evolutionary Microbiology 56:1317–

625   1322.

626   21.   Sun Q, Collins R, Huang S, Holmberg-Schiavone L, Anand GS, Tan C-H, van-den-Berg S,

627   Deng L-W, Moore PK, Karlberg T, Sivaraman J. 2009. Structural Basis for the Inhibition

628   Mechanism of Human Cystathionine γ-Lyase, an Enzyme Responsible for the Production

629   of H2S *. Journal of Biological Chemistry 284:3076–3085.

630   22.   Hatzenpichler R, Krukenberg V, Spietz RL, Jay ZJ. 2020. Next-generation physiology

631   approaches to study microbiome function at single cell level. Nat Rev Microbiol 18:241–

632   256.

633   23.   Hubberten U, Lara RJ, Kattner G. 1994. Amino acid composition of seawater and

634   dissolved humic substances in the Greenland Sea. Marine Chemistry 45:121–128.

635   24.   Cindy, Lee. 2005. Amino Acid and Amine Biogeochemistry in Marine Particulate Material

636   and Sediments. https://www.semanticscholar.org/paper/Amino-Acid-and-Amine-

Biogeochemistry-in-Marine-and-Cindy.-

Lee/8f4bba4498ee7eeebd720492a938fe792ecb91cf. Retrieved 12 November 2021.

25. Brosnan JT, Brosnan ME. 2006. The Sulfur-Containing Amino Acids: An Overview. The

Journal of Nutrition 136:1636S-1640S.

26. Thume K, Gebser B, Chen L, Meyer N, Kieber DJ, Pohnert G. 2018. The metabolite

dimethylsulfoxonium propionate extends the marine organosulfur cycle. Nature 563:412–

415.

27. Yamamoto K, Oshima T, Nonaka G, Ito H, Ishihama A. 2011. Induction of the Escherichia

coli cysK gene by genetic and environmental factors. FEMS Microbiol Lett 323:88–95.

28. Sharma S, Gray DK, Read JS, O'Reilly CM, Schneider P, Qudrat A, Gries C, Stefanoff S,

Hampton SE, Hook S, Lenters JD, Livingstone DM, McIntyre PB, Adrian R, Allan MG,

Anneville O, Arvola L, Austin J, Bailey J, Baron JS, Brookes J, Chen Y, Daly R, Dokulil M,

Dong B, Ewing K, de Eyto E, Hamilton D, Havens K, Haydon S, Hetzenauer H, Heneberry

J, Hetherington AL, Higgins SN, Hixson E, Izmest'eva LR, Jones BM, Kangur K, Kasprzak

P, Köster O, Kraemer BM, Kumagai M, Kuusisto E, Leshkevich G, May L, MacIntyre S,

Müller-Navarra D, Naumenko M, Noges P, Noges T, Niederhauser P, North RP, Paterson

AM, Plisnier P-D, Rigosi A, Rimmer A, Rogora M, Rudstam L, Rusak JA, Salmaso N,

Samal NR, Schindler DE, Schladow G, Schmidt SR, Schultz T, Silow EA, Straile D,

Teubner K, Verburg P, Voutilainen A, Watkinson A, Weyhenmeyer GA, Williamson CE,

Woo KH. 2015. A global database of lake surface temperatures collected by in situ and

satellite methods from 1985–2009. Sci Data 2:150008.

29.   Chen M, Li X-H, He Y-H, Song N, Cai H-Y, Wang C, Li Y-T, Chu H-Y, Krumholz LR, Jiang

H-L. 2016. Increasing sulfate concentrations result in higher sulfide production and

phosphorous mobilization in a shallow eutrophic freshwater lake. Water Res 96:94–104.

30.   Brock TD, Brock TD, Brock WA. 1985. A Eutrophic Lake: Lake Mendota, Wisconsin.

Springer-Verlag.

31.   Jørgensen NOG. 1987. Free amino acids in lakes: Concentrations and assimilation rates in

relation to phytoplankton and bacterial production1. Limnology and Oceanography 32:97–

111.

32.   Gardner W, Lee G. 1975. The role of amino acids in the nitrogen cycle of Lake Mendota.

Limnology and Oceanography - LIMNOL OCEANOGR 20:379–388.

33.   Lane DJ, Pace B, Olsen GJ, STAHLt DA, SOGINt ML, Pace NR. 1985. Rapid

determination of 16S ribosomal RNA sequences for phylogenetic analyses. Proc Natl Acad

Sci USA 5.

34.   Nucleobytes. 4Peaks: For peaks, four peaks. The DNA sequence trace viewer for OS X.

https://nucleobytes.com/4peaks/index.html. Retrieved 16 October 2020.

35.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.

BLAST+: architecture and applications. BMC Bioinformatics 10:421.

36.   Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial genome

assemblies from short and long sequencing reads. PLOS Computational Biology

13:e1005595.

37.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–

2069.

38.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM:

681   assessing the quality of microbial genomes recovered from isolates, single cells, and

682   metagenomes. Genome research 25:1043–55.

683   39.   Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz

684   P. 2018. A standardized bacterial taxonomy based on genome phylogeny substantially

685   revises the tree of life. Nature Biotechnology 36:996–1004.

686   40.   Zhou Z, Tran P, Liu Y, Kieft K, Anantharaman K. 2019. METABOLIC: A scalable high-

687   throughput metabolic and biogeochemical functional trait profiler based on microbial

688   genomes. bioRxiv 761643.

689   41.   Hotvedt, Jacob C, Anantharaman, Karthik. 2021. Rapid and cheap quantification of

690   Cysteine viaoxidationto Cystine by DMSO and Ortho-phthalaldehyde pre-column

691   derivatization.

692   42.   Bendall ML, Stevens SL, Chan L-K, Malfatti S, Schwientek P, Tremblay J, Schackwitz W,

693   Martin J, Pati A, Bushnell B, Froula J, Kang D, Tringe SG, Bertilsson S, Moran MA, Shade

694   A, Newton RJ, McMahon KD, Malmstrom RR. 2016. Genome-wide selective sweeps and

695   gene-specific sweeps in natural bacterial populations. The ISME Journal 10:1589–1601.

696   43.   Roux S, Chan LK, Egan R, Malmstrom RR, McMahon KD, Sullivan MB. 2017.

697   Ecogenomics of virophages and their giant virus hosts assessed through time series

698   metagenomics. Nature Communications 8.

699   44.   Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

700   Bioinformatics 34:i884–i890.

45.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new versatile

metagenomic assembler. Genome Res 27:824–834.

46.   Bushnell B. 2014. BBMap: A Fast, Accurate, Splice-Aware Aligner. LBNL-7065E.

Lawrence Berkeley National Lab. (LBNL), Berkeley, CA (United States).

47.   Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: an

adaptive binning algorithm for robust and efficient genome reconstruction from

metagenome assemblies. PeerJ 7:e7359.

48.   Olm MR, Brown CT, Brooks B, Banfield JF. 2017. dRep: a tool for fast and accurate

genomic comparisons that enables improved genome recovery from metagenomes

through de-replication. The ISME Journal 1–5.

49.   Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2020. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. Bioinformatics 36:1925–1927.

50.   Eddy SR. 2011. Accelerated profile HMM searches. PLoS Computational Biology 7.

51.   Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata H. 2019.

KofamKOALA: KEGG ortholog assignment based on profile HMM and adaptive score

threshold. bioRxiv https://doi.org/10.1101/602110.

List of Supplementary Items

List of Supplementary Tables
Table S1. Description of all initial isolates used in screening
Table S2. Quantitative measurement of H_2S over time in the 3 isolates
Table S3. Quantitative measurement of cystine over time in the 3 isolates.
Table S4. Description of the 6 genes involved in the degradation of L-cysteine or D-cysteine into pyruvate, ammonia and hydrogen sulfide
Table S5. Detailed list of all potential enzymatic activities of genes in the cysteine degradation and biosynthesis pathways.
Table S6. METABOLIC table: Full genomic content of the 3 isolates
Table S7. HMMsearch table output of 16S rRNA amplicons of the 28 isolates that produce H2S from Figure 1 against the Lake Mendota Time Series assembly.
Table S8. HMMsearch table output of 16S rRNA amplicons of the 28 isolates that produce H2S from Figure 1 against the metagenome-assembled-genomes from the LMTS.
Table S9. HMMsearch table output of 16S rRNA amplicons of the 28 isolates that produce H2S from Figure 1 against the SSU (16S small ribosomal subunit) of the metagenome-assembled-genomes from the LMTS.
Table S10. Cysteine hits the Assembly.
Table S11. Cysteine hits to MAGS
Table S12. MAG characteristics binned from LMTS.

List of Supplementary Figures
Figure S1. H_2S and ammonia production screening of the isolates

List of Supplementary Files
File S1. HMM profiles.zip

 Supplementary Figures

749
750

Figure S1. Qualitative accumulation of hydrogen sulfide among microbial isolates enriched from a freshwater lake water column. Filled circles represent isolates grown with cysteine, and open circles represent isolates grown without cysteine. The vertical line represents values corrected for the control (natural ammonia production/consumption in the negative control). All points to the right of the vertical lines indicate an accumulation of ammonia, and all points to the left of the vertical lines refer to those that consumed ammonia after 24 hours. The isolates #43, #13 and #66 (bolded) were selected for further analysis.

A.

B.