1

Oxytocin promotes social grooming in bonobos

James Brooks[1,2*], Fumihiro Kano[2,3,4*], Hanling Yeow[1,2], Naruki Morimura[1,2], Shinya Yamamoto[1,5]

[1]Wildlife Research Center, Kyoto University

[2]Kumamoto Sanctuary, Kyoto University

[3]Center for the Advanced Study of Collective Behaviour, University of Konstanz

[4]Max-Planck Institute of Animal Behavior

[5]Kyoto University Institute for Advanced Study, Kyoto University

*Co-first authors

Correspondence to: James Brooks

Email: jamesgerardbrooks@gmail.com (JB)

Address: 990, Otao, Misumi, Uki City, Kumamoto Prefecture, 8693201, Japan

Abstract:

Oxytocin has attracted research attention due to its role in promoting social bonding. One notable

recent hypothesis is the biobehavioral feedback loop, which posits that the oxytocin system has

evolved to support the formation and maintenance of social bonds through a positive feedback

loop, where oxytocin promotes social behaviours which then cause oxytocin release themselves.

In the two *Pan* species, humans' closest relatives, oxytocin is known to be released following key

behaviours related to social bonding, such as social grooming in chimpanzees and female-female

sexual behaviour in bonobos. However, no experimental evidence has demonstrated that oxytocin

promotes such socio-positive behaviours. To test this, we administered nebulized oxytocin or

saline placebo to a group of female bonobos and subsequently observed the change in their gross

behavior during free interaction. We found that bonobos groomed other group members

significantly more frequently in the oxytocin compared to placebo condition. Other behavioural

measures did not largely differ between conditions, except for a nonsignificant trend for reduction

in abnormal regurgitation/reingestion behaviour. Overall, we found that oxytocin promoted socio-

positive interaction in bonobos, providing support for the biobehavioural feedback loop

hypothesis of oxytocin in bonobo social evolution.

Keywords: Oxytocin, bonobos, social bonding, biobehavioural feedback loop, social grooming, *pan*

paniscus

Introduction:

Oxytocin is a hormone neuropeptide conserved through mammalian evolution and plays diverse roles in regulating social behaviors across species. Evidence has accumulated for a biobehavioural feedback loop in mammalian social behaviours such as maternal care [1–3], pair bonding [4], and even the dog-human bond [5,6], where oxytocin both promotes and is released by key behaviours related to social bonding. In the great ape species most closely related to humans the presence of such a positive feedback loop has also been suggested [7], though the importance of it in human and other great ape species' evolution remains unclear. Currently, experimental evidence is lacking as to whether oxytocin promotes socio-positive interaction in these species, a key piece of evidence that would support the presence of such a positive feedback loop in the *Pan* species.

Among non-human great apes, the majority of studies have been conducted through measurement of urinary oxytocin following key social behaviours. Crockford et al. [7] showed that urinary oxytocin levels in wild chimpanzees increase following social grooming, a key socio-positive behavior widely present in nonhuman primates, and proposed that a positive feedback loop may have evolved to support social bonding in this species. Relatedly, Moscovice et al. [18] found that urinary oxytocin levels in wild female bonobos increased following same-sex sexual behaviour, genito-genital (GG) rubbing. Bonobos also increased proximity and coalitionary support among females after GG-rubbing; though it remains unclear if oxytocin played a direct role in these behavioural changes. Other studies have additionally demonstrated that urinary oxytocin rises after food sharing [19], reconciliation [20], border patrols [21,22], and group hunting [21,23], further suggesting its importance to social bonds and coordination.

In several primate species, studies have demonstrated exogenous oxytocin can impact a

wide range of social behaviours (reviewed in [8]). In macaques, several studies have demonstrated

that oxytocin alters social gaze, such as increased attention to eyes [9], reduced attention to

negative and fearful facial expressions [10] as well as social threats [11], and more gaze following

[12]. In one of the first to test the effect of oxytocin in spontaneous social behaviour among

multiple macaques, although still confined to primate chairs in a laboratory setting, Jiang and Platt

[13] found evidence that oxytocin flattened the dominance hierarchy and enhanced synchrony of

mutual gaze. Marmosets similarly showed an increase in attention to eyes [14] following oxytocin

administration and an increase in anxiety and vigilance following administration of an oxytocin

antagonist [15]. Another study found that oxytocin promoted huddling in marmosets, while an

oxytocin antagonist reduced social proximity and huddling [16]. On the other hand, in capuchin

monkeys oxytocin was found to reduce food sharing through increasing interindividual distance

[17]; the authors interpreted these results as derived from oxytocin's anxiolytic effect, which

increased social distance and thereby decreased opportunities for food sharing [17].

Tthe results of three studies measuring behaviour following oxytocin administration in

non-human great apes were mixed. Proctor et al. [24] administered oxytocin to eight chimpanzees

individually for one trial each in both saline and oxytocin conditions then observed them in their

regular social groups. Although they did not find significant effects for any behaviours measured,

the authors note that it may be due to methodological issues, such as failures to find effective

dose of oxytocin for chimpanzees and effective time window to test the oxytocin effect, or due to

influence from groupmates who did not receive oxytocin before social interaction. Hall et al. [25]

similarly found no effect of oxytocin when chimpanzee dyads were administered oxytocin or saline

placebo and subsequently tested in a token exchange task. Each participant chose one of two

tokens to exchange and received rewards based on the choice of both participants in distributions

based on games such as the prisoner's dilemma and hawk-dove. However, although this study

administered oxytocin to a dyad, the authors reported the same methodological concerns for the

oxytocin administration procedure as well as a confound between experimental condition and

order. No clear patterns emerged in either the placebo or oxytocin conditions, limiting

interpretation of oxytocin's possible effect. On the contrary to these studies reporting null results,

Brooks et al. [26] found that oxytocin enhanced species-typical social gaze, increasing eye contact

in bonobos but not chimpanzees, indicating that oxytocin can modulate gaze behaviour. While the

species difference in the latter study cannot be attributed to differences in oxytocin administration

procedure, it remains unclear whether the lack of effect in the prior studies is due to methodology

of oxytocin administration or that exogenous oxytocin fails to significantly affect chimpanzee

social behaviour.

Critically, although it is central to the biobehavioural feedback loop hypothesis that both socio-

positive interactions cause oxytocin release and that oxytocin can lead to socio-positive

interactions, there is no direct evidence showing that oxytocin promotes any socio-positive

interaction in *Pan*. Given recent progresses in this line of research, it is worthwhile to test whether

oxytocin promotes key social behaviours related to social bonding in *Pan*, particularly in bonobos

using the updated methods of oxytocin administration. Therefore, this study administered

nebulized oxytocin or saline placebo to a whole group of female bonobos following the methods

employed in Brooks et al. [26] and subsequently observed the change in their gross interactive

behavior, including grooming and GG-rubbing, as well as other noninteractive behaviours during

their free interaction. Based on previous studies, we developed two hypotheses regarding how

oxytocin would impact bonobos' social behaviour. As mentioned, the biobehavioural feedback

loop hypothesis suggests that an oxytocin positive feedback loop has evolved to support *pan* social

bonding, and in particular we predicted that oxytocin would promote social grooming, which is

known to be released through social grooming in wild chimpanzees [7]. This would suggest that

oxytocin acts in a self-reinforcing loop through grooming as proposed by Crockford et al. [7],

forming a part of the endocrine basis of *pan* bond formation and maintenance. Alternately, the

social satiation hypothesis predicted that the oxytocin released through grooming instead evolved

to serve as a socially mediated anxiolytic. This hypothesis predicted that oxytocin, through its

anxiolytic effect, would reduce the need for grooming due to lower social tension, similar to the

reduced food sharing observed in Brosnan et al. [17] .

Methods:

Ethics statement:

All ape participants received regular feedings, daily enrichment, and had ad libitum access to

water. No change was made to their daily care routine for the purpose of this study. Apes were

never restrained at any point. Ethical approval number was WRC-2020-KS014A.

We carefully considered the safety of the OT administration as in previous studies. Again, we

based this decision on the fact that 1) OT is often administered to human children and adults, 2)

OT is active for only a short period of time following administration with no known side effects in

humans (MacDonald et al., 2011), 3) OT is naturally produced in bonobos and chimpanzees

following relevant behaviors (Crockford et al., 2013; Moscovice et al., 2019), and 4) no previous

studies administering OT intranasally to chimpanzees or bonobos reported any agonistic

interaction (Brooks et al., 2021; Hall et al., 2019; Proctor et al., 2016).

Participants:

Four adult female bonobos at Kumamoto Sanctuary participated in this research. Details about

participant ages and rearing histories can be found in supplementary material (Table S1). Animals

were not food or water deprived at any time and were given both physical and social

environmental enrichment in their daily life. The bonobos live in a dynamic grouping structure

where three of the four females are together on any given day, and the fourth is with two male

bonobos. These two males were not involved in this study because one male refused to participate

in any oxytocin experiments, and our aim was to test whole groups at a time. Three of the females

join the male bonobos with varying frequency, while the fourth (Lenore) is always with other

female bonobos. Individuals thus had a varying number of trials, with Lenore having the most due

to never joining the male group (24 trials), followed by Lolita (20 trials), followed by Louise and

Ikela (14 trials each) who are most often with the males.

Administration procedure:

Oxytocin administration procedures followed Brooks et al. [26]. Briefly, oxytocin was dissolved in

saline at a concentration of 40IU/mL. The oxytocin solution or placebo control was nebulized into

a box using a portable nebulizer (Omron NE-U100) at a minimum rate of 0.25mL/minute, for a

cumulative 4 minutes while apes drank juice (thus a total of 40IU or more was nebulized during

the administration period). Timing was stopped while apes' noses were outside the box.

Participation was voluntary and apes were never restrained.

Observation procedure:

Observation began 30 minutes after completion of administration criteria of all individuals (30

minutes from the completion of the last individual), in line with previous studies [8], and lasted for

one hour. The last individual to complete administration procedures was always within 30 minutes

of the first individual to finish, and thus all participants were observed for one hour, starting and

finishing between 30 minutes and 2 hours following completion of administration procedures on

any given day. Observation methods combined scan and event sampling. Specifically, every 2

minutes, interindividual proximity was estimated for each dyad into one of four categories (in

contact, within arm's reach-one individual could extend their arm to touch the other, < 3 meters,

and > 3 meters) as well as each individual's behaviour (grooming including direction, resting, self-

directed behaviour, moving, eating). All occurrences of play, GG-rubbing, abnormal behaviour, and

aggression towards groupmates (including displays) were additionally recorded (see Brooks et al.

[27] for more details about these observation methods).

Analysis

All analyses were conducted in R [28]. Behavioural scan data was analyzed with binomial GLMMs

(Generalized Linear Mixed Models) with package lme4 [29], where each individual at each point in

time was characterized as either engaged in (1) or not engaged in (0) a given behaviour. The model

included a test fixed effect of condition in addition to control fixed effects of day (counting

upwards from first experimental day) and time, with random effects of dyad and grouping

structure (a unique value was given for each possible combination of individuals) and random

slopes of each fixed effect for each random effect. Numeric effects were scaled to a mean of 0 and

standard deviation of 1. Random slope structure was kept maximal, except that the interaction

between random slopes and intercepts was removed due to issues with convergence. For

grooming data, we analyzed rates of active grooming (giving or mutual grooming), and receiving

grooming was valued as 0 for not actively grooming another individual.

171   Proximity data was analyzed using a CLMM (cumulative link mixed model) on ordinal data

using the package "ordinal" [30]. Fixed and random effect structures were the same as those in

the behavioural scan data analysis, except for the individual participant variable was replaced by

dyad (a unique value for each dyad), and the addition of two random effects to represent the two

individuals within a dyad (randomly distributed as individual variable 1 and 2). Random slope

structure was kept maximal and the interaction between random slopes and intercepts was

retained in this model.

All occurrence data was analyzed with a binomial GLMM, where each individual for each

day was characterized as having engaged in (1) or not engaged in (0) a given behaviour. The fixed

and random effect structure was the same as in the scan behaviour models, except for the time

variable removed due to data being summarized across a given observation day. Model syntax for

all model types can be found in supplemental material.

Results

Bonobos engaged in active social grooming significantly more in the oxytocin condition than the

control condition (= 0.48, SE = 0.16, χ^2 = 3.94, p = 0.047; Figure 1). Plots of this effect by

participant, group, and time can be found in supplementary material (figures S1, S2, and S3).

Figure 1. Mean active grooming rates in the oxytocin compared to saline control condition (giving

or mutual grooming). Error bars represent 95% confidence intervals.

We did not find significant differences in other behaviors between conditions. However, there was

a trend for closer interindividual proximity in the oxytocin compared to control condition (β = -

0.22, SE = 0.12, χ^2 = 2.85, p = 0.091) and for a reduction in the abnormal behaviour regurgitation

and reingestion in the oxytocin compared to control condition (β = -0.73, SE = 0.36, χ^2 = 3.08, p =

0.079). There were no significant changes in self-directed behaviour (β = -0.33, SE = 0.18, χ^2 = 2.04,

p =0.15) or rest (β =-0.13, SE = 0.18, χ^2 = 0.47, p =0.49). Bonobos engaged in GG-rubbing only once

(oxytocin condition) and displayed no aggression toward groupmates or any bouts of play during

the observation period. See supplementary material Table S2 for full details of all models.

Discussion:

Female bonobos groomed groupmates significantly more in the oxytocin condition than in the

saline condition. In combination with the previous studies showing oxytocin is released in bonobos

and chimpanzees following socio-positive interaction [7,18], our results provide experimental

support for the biobehavioural feedback loop hypothesis in bonobos. Other observed behaviours

did not largely differ between the oxytocin and saline conditions (self-directed behavior and rest)

or were rarely or never observed during our 1-hour observation window (GG-rubbing, play, and

aggression). There were trends for closer inter-individual proximity and reduced rates of

regurgitation and reingestion. Brosnan et al. [17] found that oxytocin increased interindividual

distance in capuchin monkeys (though individuals were separated by a mesh partition), possibly

through oxytocin's known anxiolytic effect, and this increase in interindividual distance then

reduced food-sharing. Our results did not demonstrate this pattern in freely interacting bonobos

and thus did not support the social satiation hypothesis. Potential reduction of regurgitation and

reingestion may be explained by oxytocin's effect on anxiety or digestion [31–33].

Although we present the first observation that oxytocin promotes socio-positive

interaction in female bonobos, there are several important limitations in this study. First, due to

limited possibility of testing, enclosures suitable for detailed observation, and some apes'

willingness to join experiments, the sample was limited to just four adult female bonobos. This

limits firm conclusions about a biobehavioural feedback loop, as Crockford et al. [7] measured

urinary oxytocin after grooming only in chimpanzees, while Moscovice et al. [18] focused on sexual

interactions rather than grooming in bonobos. While these previous studies characterized the

changes similarly as socio-positive behaviours strengthening bond formation, conclusive proof of a

biobehavioural feedback loop in one or both species would require evidence of increased urinary

oxytocin following grooming in bonobos, administration of oxytocin promoting GG-rubbing in

bonobos, or administration of oxytocin promoting grooming among bond partners in chimpanzees.

It remains possible that exogenous oxytocin promotes grooming in bonobos but not chimpanzees,

or that oxytocin is released following grooming in chimpanzees but not bonobos. Previous work

has also indicated sex-specific effects of oxytocin [34–36], and thus it remains unclear whether our

results can be generalized to different sex pairs. However, it should be noted that Crockford et al.

[7] did not find significant differences between female-female, female-male, and male-male dyads

in urinary oxytocin level following grooming in wild chimpanzees. Second, it has not been

demonstrated whether urinary oxytocin shows a similar increase following grooming in wild

bonobos, though Moscovice et al. [18] found that urinary oxytocin rose following GG-rubbing

which was interpreted as a similar socio-positive behaviour. GG-rubbing was very infrequent in our

study, possibly due to low overall social tension, precluding formal analysis. Relatedly, Proctor et al.

[24] did not find any behavioural change after administration of oxytocin on captive chimpanzees.

While several methodological limitations preclude interpretation of their results (and unlike our

study, only one individual, instead of a whole group, was tested and observed at a time), currently

it has not been demonstrated whether oxytocin promotes grooming in captive chimpanzees.

Finally, the small number of participants did not enable us to test the effect of closeness with and

selectivity to a certain groupmate, which may interact with the observed increase in grooming.

In conclusion our finding offered experimental evidence that oxytocin promotes socio-

positive interaction in bonobos. The oxytocin system thus may have been coopted through

evolution to the formation and maintenance of social bonds though a positive feedback loop.

Moreover, we suggest that oxytocin administration can be an effective tool in great ape research

involving free interactions among groupmates. Future work should further test potential

differences in oxytocin's effect between species, should examine inter-individual variation with

respect to social closeness and centrality, and should study how social contexts such as feeding

tension interact with this effect. We demonstrate that exogenous oxytocin can affect great ape

behaviour in naturalistic, spontaneous social interactions and provide experimental support for

the biobehavioural feedback loop hypothesis of oxytocin in the evolution of bonobo social bonding.

Acknowledgements:

We thank the bonobos at Kumamoto Sanctuary for participating in this study. We also thank

Etsuko Nogami and the other caretakers at Kumamoto Sanctuary for their support throughout this

project, Dr. Satoshi Hirata for his advice and support, and Yutaro Sato for helping in data collection.

This study was funded by Japan Society for the Promotion of Science (KAKENHI #21J21123 to J.B.,

#19H01772 and #20H05000 to F.K., and #19H00629 and #19H05736 to S.Y.)

References:

1. Nagasawa M, Okabe S, Mogi K, Kikusui T. 2012 Oxytocin and mutual communication in
mother-infant bonding. *Front. Hum. Neurosci.* **6**, 31. (doi:10.3389/fnhum.2012.00031)

2. Rilling JK, Young LJ. 2014 The biology of mammalian parenting and its effect on offspring
social development. *Science (80-.).* **345**, 771–776. (doi:10.1126/SCIENCE.1252723)

3. Da Costa APC, Guevara-Guzman RG, Ohkura S, Goode JA, Kendrick KM. 1996 The Role of
Oxytocin Release in the Paraventricular Nucleus in the Control of Maternal Behaviour in the
Sheep. *J. Neuroendocrinol.* **8**, 163–177. (doi:10.1046/j.1365-2826.1996.04411.x)

4. Insel TR, Young LJ. 2001 The neurobiology of attachment. *Nat. Rev. Neurosci.* **2**, 129–136.
(doi:10.1038/35053579)

5. Nagasawa M, Ogawa M, Mogi K, Kikusui T. 2017 Intranasal Oxytocin Treatment Increases
Eye-Gaze Behavior toward the Owner in Ancient Japanese Dog Breeds. *Front. Psychol.* **8**,
1624. (doi:10.3389/fpsyg.2017.01624)

6. Nagasawa M, Mitsui S, En S, Ohtani N, Ohta M, Sakuma Y, Onaka T, Mogi K, Kikusui T. 2015
Oxytocin-gaze positive loop and the coevolution of human-dog bonds. *Science (80-.).* **348**,
333–336. (doi:10.1126/science.1261022)

7. Crockford C, Wittig RM, Langergraber K, Ziegler TE, Zuberbühler K, Deschner T. 2013
Urinary oxytocin and social bonding in related and unrelated wild chimpanzees. *Proc. R. Soc.
B Biol. Sci.* **280**, 20122765. (doi:10.1098/rspb.2012.2765)

8. Bauman MD, Murai T, Hogrefe CE, Platt ML. 2018 Opportunities and challenges for
intranasal oxytocin treatment studies in nonhuman primates. *Am. J. Primatol.* **80**, 1–8.
(doi:10.1002/ajp.22913)

9. Dal Monte O, Noble PL, Costa VD, Averbeck BB. 2014 Oxytocin enhances attention to the
eye region in rhesus monkeys. *Front Neurosci* **8**, 41. (doi:10.3389/fnins.2014.00041)

10. Parr LA, Modi M, Siebert E, Young LJ, Rd G. 2013 Intranasal oxytocin selectively attenuates
rhesus monkeys' attention to negative facial expressions. *Psychoneuroendocrinology*
(doi:10.1016/j.psyneuen.2013.02.011)

11. Ebitz RB, Watson KK, Platt ML. 2013 Oxytocin blunts social vigilance in the rhesus macaque.
Proc. Natl. Acad. Sci. **110**, 11630–11635. (doi:10.1073/pnas.1305230110)

12. Putnam PT, Roman JM, Zimmerman PE, Gothard KM. 2016 Oxytocin enhances gaze-
following responses to videos of natural social behavior in adult male rhesus monkeys.
Psychoneuroendocrinology **72**, 47–53. (doi:10.1016/j.psyneuen.2016.05.016)

13. Jiang Y, Platt ML. 2018 Oxytocin and vasopressin flatten dominance hierarchy and enhance
behavioral synchrony in part via anterior cingulate cortex. *Sci. Rep.* **8**, 1–14.
(doi:10.1038/s41598-018-25607-1)

14. Kotani M, Shimono K, Yoneyama T, Nakako T, Matsumoto K, Ogi Y, Konoike N, Nakamura K,

Ikeda K. 2017 An eye tracking system for monitoring face scanning patterns reveals the
enhancing effect of oxytocin on eye contact in common marmosets.
Psychoneuroendocrinology **83**, 42–48. (doi:10.1016/J.PSYNEUEN.2017.05.009)

15. Cavanaugh J, Carp SB, Rock CM, French JA. 2016 Oxytocin modulates behavioral and
physiological responses to a stressor in marmoset monkeys. *Psychoneuroendocrinology* **66**,
22–30. (doi:10.1016/J.PSYNEUEN.2015.12.027)

16. Smith AS, Ågmo A, Birnie AK, French JA. 2010 Manipulation of the oxytocin system alters
social behavior and attraction in pair-bonding primates, Callithrix penicillata. *Horm. Behav.*
57, 255–262. (doi:10.1016/j.yhbeh.2009.12.004)

17. Brosnan SF, Talbot CF, Essler JL, Leverett K, Flemming T, Dougall P, Heyler C, Zak PJ. 2015
Oxytocin reduces food sharing in capuchin monkeys by modulating social distance.
Behaviour **152**, 941–961. (doi:10.1163/1568539X-00003268)

18. Moscovice LR, Surbeck M, Fruth B, Hohmann G, Jaeggi A V., Deschner T. 2019 The
cooperative sex: Sexual interactions among female bonobos are linked to increases in
oxytocin, proximity and coalitions. *Horm. Behav.* **116**, 104581.
(doi:10.1016/j.yhbeh.2019.104581)

19. Wittig RM, Crockford C, Deschner T, Langergraber KE, Ziegler TE, Zuberbühler K. 2014 Food
sharing is linked to urinary oxytocin levels and bonding in related and unrelated wild
chimpanzees. *Proc. R. Soc. B Biol. Sci.* **281**, 20133096. (doi:10.1098/rspb.2013.3096)

20. Preis A, Samuni L, Mielke A, Deschner T, Crockford C, Wittig RM. 2018 Urinary oxytocin
levels in relation to post-conflict affiliations in wild male chimpanzees (Pan troglodytes
verus). *Horm. Behav.* **105**, 28–40. (doi:10.1016/j.yhbeh.2018.07.009)

21. Samuni L, Preis A, Mundry R, Deschner T, Crockford C, Wittig RM. 2017 Oxytocin reactivity
during intergroup conflict in wild chimpanzees. *Proc. Natl. Acad. Sci.* **114**, 268–273.
(doi:10.1073/pnas.1616812114)

22. Samuni L, Preis A, Deschner T, Wittig RM, Crockford C. 2019 Cortisol and oxytocin show
independent activity during chimpanzee intergroup conflict. *Psychoneuroendocrinology* **104**,
165–173. (doi:10.1016/j.psyneuen.2019.02.007)

23. Samuni L, Preis A, Deschner T, Crockford C, Wittig RM. 2018 Reward of labor coordination
and hunting success in wild chimpanzees. *Commun. Biol.* **1**, 138. (doi:10.1038/s42003-018-
0142-3)

24. Proctor D, Calcutt SESE, Burke K, de Waal FBMFBM. 2016 Intranasal Oxytocin Failed to
Affect Chimpanzee (Pan troglodytes) Social Behavior. *Anim. Behav. Cogn.* **3**, 150–158.
(doi:doi:10.12966/abc.04.08.2016.)

25. Hall K, Smith M, Russell JL, Lambeth SP, Schapiro SJ, Brosnan SF. 2019 Chimpanzees Rarely
Settle on Consistent Patterns of Play in the Hawk Dove, Assurance, and Prisoner's Dilemma
Games, in a Token Exchange Task. *Anim. Behav. Cogn.* **6**, 48–70.
(doi:10.26451/abc.06.01.04.2019)

26.   Brooks J, Kano F, Sato Y, Yeow H, Morimura N, Nagasawa M, Kikusui T, Yamamoto S. 2021
Divergent effects of oxytocin on eye contact in bonobos and chimpanzees.
Psychoneuroendocrinology **125**, 105119. (doi:10.1016/j.psyneuen.2020.105119)

27.   Brooks J, Onishi E, Clark IR, Bohn M, Yamamoto S. 2021 Uniting against a common enemy:
Perceived outgroup threat elicits ingroup cohesion in chimpanzees. *PLoS One* **16**, e0246869.
(doi:10.1371/journal.pone.0246869)

28.   R Core Team. 2019 R: A Language and Environment for Statistical Computing. *R Found. Stat.*
Comput.

29.   Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using
{lme4}. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

30.   Christensen RHB. 2019 ordinal: Regression Models for Ordinal Data.

31.   Dobrovol'skaia ZA, Kosenko AF. 1989 [The role of oxytocin in realizing hypothalamic effects
on the motor function of the digestive tract]. *Fiziol. Zh. SSSR Im. I. M. Sechenova* **75**, 734–7.

32.   Richard P, Moos F, Freund-Mercier MJ. 1991 Central effects of oxytocin. *Physiol. Rev.* **71**,
331–70. (doi:10.1152/physrev.1991.71.2.331)

33.   Uvnäs-Moberg K. 1998 Antistress Pattern Induced by Oxytocin. *Physiology* **13**, 22–25.
(doi:10.1152/physiologyonline.1998.13.1.22)

34.   Gao S *et al.* 2016 Oxytocin, the peptide that bonds the sexes also divides them. *Proc. Natl.*
Acad. Sci. U. S. A. **113**, 7650–4. (doi:10.1073/pnas.1602620113)

35.   Caldwell HK. 2018 Oxytocin and sex differences in behavior. *Curr. Opin. Behav. Sci.* **23**, 13–
20. (doi:10.1016/J.COBEHA.2018.02.002)

36.   Dumais KM, Bredewold R, Mayer TE, Veenema AH. 2013 Sex differences in oxytocin
receptor binding in forebrain regions: Correlations with social interest in brain region- and
sex- specific ways. *Horm. Behav.* **64**, 693–701. (doi:10.1016/J.YHBEH.2013.08.012)

357