1

Testing the effect of oxytocin on social grooming in bonobos

James Brooks[1,2*], Fumihiro Kano[2,3,4*], Hanling Yeow[1,2], Naruki Morimura[1,2], Shinya Yamamoto[1,5]

[1]Wildlife Research Center, Kyoto University

[2]Kumamoto Sanctuary, Kyoto University

[3]Center for the Advanced Study of Collective Behaviour, University of Konstanz

[4]Max-Planck Institute of Animal Behavior

[5]Kyoto University Institute for Advanced Study, Kyoto University

*Co-first authors

Correspondence to: James Brooks

Email: jamesgerardbrooks@gmail.com (JB)

Address: 990, Otao, Misumi, Uki City, Kumamoto Prefecture, 8693201, Japan

Abstract:

Oxytocin has attracted research attention due to its role in promoting social bonding. One notable

hypothesis is the biobehavioral feedback loop, which posits that some functions of the oxytocin

system support the formation and maintenance of social bonds through a positive feedback loop,

where oxytocin promotes social behaviours which then cause oxytocin release themselves. In the

two *Pan* species, humans' closest relatives, oxytocin is known to be released following key

behaviours related to social bonding, such as social grooming in chimpanzees and female-female

sexual behaviour in bonobos. However, no experimental evidence has demonstrated that oxytocin

promotes such socio-positive behaviours in *Pan*. To test this, we administered nebulized oxytocin

or saline placebo to a group of female bonobos and subsequently observed the change in their

gross behavior during free interaction. We found an overall effect of more frequent grooming in

the oxytocin condition. However, on the individual level this effect remained significant for only

one participant in our follow-up models, suggesting future work should explore inter-individual

variation. These results provide some experimental support for the biobehavioural feedback loop

hypothesis of oxytocin in bonobo social bonding and encourage more attention towards the effect

of oxytocin on social behaviour in more naturalistic settings involving free interaction between

participants.

Keywords: Oxytocin, bonobos, social bonding, biobehavioural feedback loop, social grooming, *Pan*

paniscus

Introduction:

Oxytocin is a hormone neuropeptide conserved through mammalian evolution and plays diverse

roles in regulating social behaviors across species. Evidence has accumulated for an oxytocin

biobehavioural feedback loop in mammalian social behaviours such as maternal care [1–3], pair

bonding [4], and even the dog-human bond [5,6], where oxytocin both promotes and is released

by key behaviours related to social bonding. In great apes the presence of such a positive feedback

loop has also been suggested [7], though the importance of it in human and other great ape

species' evolution remains unclear. Currently, experimental evidence is lacking as to whether

oxytocin promotes socio-positive interaction in these species, a key piece of evidence that would

support the presence of such a positive feedback loop.

Among non-human great apes, the majority of studies have been conducted through

measurement of urinary oxytocin following key social behaviours. Crockford et al. [7] showed that

urinary oxytocin levels in wild chimpanzees increase following social grooming, a key socio-positive

behavior widely present in nonhuman primates, and proposed that a positive feedback loop

through oxytocin may have evolved to support social bonding in this species. Relatedly, Moscovice

et al. [8] found that urinary oxytocin levels in wild female bonobos increased following same-sex

sexual behaviour, genito-genital (GG) rubbing. Bonobos also increased proximity and coalitionary

support among females after GG-rubbing; though it remains unclear if oxytocin played a direct

role in these behavioural changes. Other studies have additionally demonstrated that urinary

oxytocin in chimpanzees rises after food sharing [9], reconciliation [10], border patrols [11,12], and

group hunting [11,13], further suggesting its importance to social bonds and coordination.

In several primate species, studies have demonstrated exogenous oxytocin can impact a

wide range of social behaviours (reviewed in [14]). In macaques, several studies have

demonstrated that oxytocin alters social gaze, such as increased attention to eyes [15], reduced

attention to negative and fearful facial expressions [16] as well as social threats [17], and more

gaze following [18]. In one of the first to test the effect of oxytocin in spontaneous social

behaviour among multiple macaques, although still confined to primate chairs in a laboratory

setting, Jiang and Platt [19] found evidence that oxytocin flattened the dominance hierarchy and

enhanced synchrony of mutual gaze. Marmosets similarly showed an increase in attention to eyes

[20] following oxytocin administration and an increase in anxiety and vigilance following

administration of an oxytocin antagonist [21]. Another study found that oxytocin promoted

huddling in marmosets, while an oxytocin antagonist reduced social proximity and huddling [22].

On the other hand, in capuchin monkeys oxytocin was found to reduce food sharing through

increasing interindividual distance [23]; the authors interpreted these results as derived from

oxytocin's anxiolytic effect, which increased social distance and thereby decreased opportunities

for food sharing [23].

The results of three studies measuring behaviour following oxytocin administration in non-

human great apes were mixed. Proctor et al. [24] administered oxytocin to eight chimpanzees

individually for one trial each in both saline and oxytocin conditions then observed them in their

regular social groups. Although they did not find significant effects for any behaviours measured,

the authors note that it may be due to methodological issues, such as in establishing an effective

dose of oxytocin for chimpanzees or influence from groupmates who did not receive oxytocin

before social interaction. Hall et al. [25] similarly found no effect of oxytocin when chimpanzee

dyads were administered oxytocin or saline placebo and subsequently tested in a token exchange

task. Each participant chose one of two tokens to exchange and received rewards based on the

choice of both participants in distributions based on games such as the prisoner's dilemma and

hawk-dove. However, although this study administered oxytocin to a dyad, the authors reported

the same methodological concerns for the oxytocin administration procedure as well as a

confound between experimental condition and order. No clear patterns emerged in either the

placebo or oxytocin conditions, limiting interpretation of oxytocin's possible effect. On the

contrary to these studies reporting null results, Brooks et al. [26] found that oxytocin enhanced

species-typical social gaze, increasing eye contact in bonobos but not chimpanzees, indicating that

oxytocin can modulate gaze behaviour. While the species difference in Brooks et al. cannot be

attributed to differences in oxytocin administration procedure, it remains unclear whether the lack

of effect in Proctor et al. and Hall et al. is due to methodology of oxytocin administration or that

exogenous oxytocin fails to significantly affect chimpanzee social behaviour.

Critically, although it is central to the biobehavioural feedback loop hypothesis that both socio-

positive interactions cause oxytocin release and that oxytocin can lead to socio-positive

interactions, there is no direct evidence showing that oxytocin promotes any socio-positive

interaction in *Pan*. Given recent progresses in this line of research, it is worthwhile to test whether

oxytocin promotes key social behaviours related to social bonding in *Pan*, particularly in bonobos

using the updated methods of oxytocin administration. Therefore, this study administered

nebulized oxytocin or saline placebo to a whole group of female bonobos following the methods

employed in Brooks et al. [26] and subsequently observed the change in their gross interactive

behavior, including grooming and GG-rubbing, as well as other noninteractive behaviours during

their free interaction. As mentioned, the biobehavioural feedback loop hypothesis suggests that

an oxytocin positive feedback loop has evolved to support *Pan* social bonding, and in particular we

predicted that oxytocin would promote social grooming, which is known to be released through

social grooming in wild chimpanzees [7]. For practical reasons, we tested bonobos instead of

chimpanzees. In bonobos a similar argument about oxytocin and socio-positive behaviour was

made in Moscovice et al. [8], who found that GG-rubbing led to higher urinary oxytocin in wild

female bonobos. This study aimed to observe all socio-positive behaviors including play, GG

rubbing, and grooming, but focused on social grooming in particular. This choice was made

because our bonobos are often in close proximity and engage readily in grooming after

experiments (oxytocin administration with rewards), while GG-rubbing is not often observed after

these events (possibly due to low overall tension).

Methods:

Ethics statement:

All bonobo participants received regular feedings, daily enrichment, and had ad libitum access to

water. No change was made to their daily care routine for the purpose of this study. Apes were

never restrained at any point. We carefully considered the safety of the oxytocin administration as

in previous studies. Again, we based this decision on the fact that 1) oxytocin is often administered

to human children and adults, 2) oxytocin is active for only a short period of time following

administration, 3) oxytocin is naturally produced in bonobos and chimpanzees following relevant

behaviors [7,8] , and 4) no previous studies administering oxytocin intranasally to chimpanzees or

bonobos reported any agonistic interaction [24–26]. All female bonobos were taking birth control

and thus no bonobos were pregnant at any time during the course of this experiment. Ethical

approval number was WRC-2020-KS014A.

Participants:

Four adult female bonobos at Kumamoto Sanctuary participated in this research. Details about

participant ages and rearing histories can be found in supplementary material (Table S1). Animals

were not food or water deprived at any time and were given both physical and social

environmental enrichment in their daily life. The bonobos live in a dynamic grouping structure

where three of the four females are together on any given day, and the fourth is with two male

bonobos. These two males were not involved in this study because one male refuses to participate

in any oxytocin experiments, and our aim was to test whole groups at a time with the same

condition. Three of the females join the male bonobos with varying frequency, while the fourth

(Lenore) is always with other female bonobos. Individuals thus had a varying number of trials, with

Lenore having the most due to never joining the male group (24 trials), followed by Lolita (20

trials), followed by Louise and Ikela (14 trials each) who are most often with the males. Transfers

between groups typically occur in the evening and are kept consistent for at least one day and up

to one week. The bonobos therefore had been in the same grouping structure for the whole day

prior to the start of experiments.

Administration procedure:

Oxytocin administration procedures followed Brooks et al. [26]. Briefly, oxytocin was dissolved in

saline at a concentration of 40IU/mL. The oxytocin solution or placebo control was nebulized into

a box using a portable nebulizer (Omron NE-U100) at a minimum rate of 0.25mL/minute, for a

cumulative 4 minutes while apes drank juice (thus a total of 40IU or more was nebulized during

the administration period). We paused counting the time while apes' noses were outside the box.

Participation to this administration was voluntary. Three of the bonobos could simultaneously

participate in oxytocin administration in their typical enclosure, while the fourth (Ikela) preferred

to move to another room to participate in the administration procedure. Thus, on days when Ikela

was in the group, we first completed the administration procedure with Ikela (accompanied by

other participant bonobos), and then returned her to the home enclosure with the other

participant bonobos; the other participant bonobos were then administered oxytocin (or saline

placebo control). On days when Ikela was not in the group, all participants were administered

oxytocin (or saline placebo) in their home enclosure. All group members received the same

condition (saline placebo or oxytocin) on any given day of experiments and finished administration

procedures within 30 minutes of one another.

One trial was performed in an experimental day. The order of conditions was pseudorandomized

such that the same condition (placebo and oxytocin) never occurred more than twice in

consecutive trials (experimental days) and that the same number of trials were conducted for

placebo and oxytocin condition for each participant and for each grouping structure. We had a

minimum of 2 days between trials to avoid any possible carryover effect of oxytocin. On each

experimental day, the experiment was performed between 11:00 and 12:15, and the observation

window therefore started between 11:30 and 12:45. Experimental days followed the same feeding

schedule; bonobos were fed breakfast around 9:00, and additional greenery is available for

foraging throughout the day. Experiments took place over five calendar months across two

calendar years.

Observation procedure:

Observation began 30 minutes after completion of administration procedures to the last individual,

and lasted for one hour. This window was chosen based on previous studies [14], where oxytocin's

effect is typically measured in the window between 30 minutes and 2 hours after completion of

administration procedures. In our experiment, the last individual to complete administration

procedures was always within 30 minutes of the first individual to finish, and thus all participants

were observed for one hour, starting and finishing between 30 minutes and 2 hours following

completion of administration procedures on any given day. We chose this window in order to

maximize data collection within the active window of oxytocin while ensuring consistency

between trials and participants, where observation windows were kept constant at 1 hour per trial

and all data points for all individuals were within the 30 minutes to 2 hour window employed in

previous studies.

Observation methods combined scan and event sampling. Specifically, every 2 minutes,

interindividual proximity was estimated for each dyad into one of four categories; in contact,

within arm's reach (one individual could extend their arm to touch the other), < 3 meters, and > 3

meters. In addition, at the same 2 minute intervals, we coded each individual's behaviour

(grooming - including direction and partner(s), resting, self-directed behaviour, moving, eating).

Finally, we recorded all occurrences of play, GG-rubbing, abnormal behaviour (in this group

primarily regurgitation and reingestion), and aggression towards groupmates (including displays).

We additionally recorded any agonistic or socio-sexual behaviour during the administration

procedures and during the 30 minutes before the formal observation window to check if the

presence of such interactions could account for our results. No agonistic behaviour was observed,

and GG-rubbing occurred on 4 trials during administration procedure, on 2 trials before oxytocin

condition and 2 trials before saline placebo condition.

Analysis

All analyses were conducted in R [29]. Behavioural scan data was analyzed with binomial GLMMs

(Generalized Linear Mixed Models) with package lme4 [30], where each individual at each

sampling point was characterized as either engaged in (1) or not engaged in (0) a given behaviour.

The model included a fixed effect of condition as a test effect and also fixed effects of time, the

square of time (time^2) and grouping structure (where a unique value was given for each possible

combination of individuals) as control effects. The model also included random effects of

participant and day (a factor with a unique value for each experimental day), as well as random

slopes of each fixed effect for each random effect. Numeric effects were z-transformed to have a

mean of 0 and the standard deviation of 1. Random slope structure was kept maximal, except that

the interaction between random slopes and intercepts was removed due to issues with

convergence [31]. For all GLMMs we checked model stability by comparing our models to those

which excluded levels of the random effects one at a time. We additionally calculated odds ratio

(OR) estimates and 95% confidence intervals for all significant effects. For the grooming data, we

analyzed rates of active grooming (giving or mutual grooming), and receiving grooming was valued

as 0 for not actively grooming another individual. Given considerable inter-individual variation

observed by visual inspection of the data, we additionally tested each individual with the same

model structure for significant effects (without the random effect of participant). We here used a

significance threshold of 0.0125 (0.05/4) to correct for multiple testing at the individual level.

Finally, we ran as models excluding each level of participant to check the overall model stability.

Proximity data was analyzed using a CLMM (cumulative link mixed model) on ordinal data

using the package "ordinal" [32]. Fixed and random effect structures were the same as those in

the behavioural scan data analysis, except for the individual participant variable was replaced by

dyad (a unique value for each dyad), and the addition of two random effects to represent the two

individuals within a dyad (randomly distributed as individual variable 1 and 2).

All occurrence data was analyzed with a binomial GLMM, where each individual for each

day was characterized as having engaged in (1) or not engaged in (0) a given behaviour. The fixed

and random effect structure was the same as in the scan behaviour models, except for the time

variables were removed due to data being summarized across a given observation day. Model

syntax for all model types can be found in supplemental material.

Results

Our model with active grooming as response revealed a significant effect of condition (oxytocin,

saline placebo; β = 1.16, SE = 0.49, χ^2 = 5.47, p = 0.019, OR = 3.18 (95% CI: 1.23, 8.23); Table 1,

Figure 1). In our stability check analysis excluding each participant one-by-one in the model, the

effect of oxytocin on grooming was significant in all models except for that excluding Louise

(excluding Ikela: β = 0.79, SE = 0.31 , χ^2 = 5.27, p = 0.022; excluding Lenore: β = 1.28, SE = 0.64, χ^2 =

8.97, p = 0.0027; excluding Lolita: β = 0.88, SE = 0.36, χ^2 = 5.06, p = 0.025; excluding Louise: β =

0.59, SE = 0.37, χ^2 = 2.34, p = 0.13). In our individual-level analysis of grooming, we found a

significant effect of condition only for Louise using a significance threshold of 0.0125 (Ikela: β =

0.86, SE = 1.51, χ^2 = 0.24, p = 0.63; Lenore: β = 0.79, SE = 0.35, χ^2 = 4.61, p = 0.032; Lolita: model

could not run (as Lolita never actively groomed in our dataset); Louise: β = 1.15, SE = 0.40, χ^2 =

9.09, p = 0.0026).

Figure 1: Rates of active (giving) grooming in the oxytocin and saline placebo conditions over the
observation window. A) Absolute rates of grooming across all trials and participants B) Time
course of grooming by condition, 1 represents giving grooming and 0 represents not giving
grooming C) Grooming rates by participant and condition D) Grooming rates by group structure by
condition.

Table 1; Details of grooming and self-directed behaviour models with significant terms in bold.

Response	Term	Estimate ± SE	χ^2	df	p
Active grooming	(Intercept)	-15.57±6.29			
	Test predictors:				
	Condition	**1.156±0.49**	**5.47**	**1**	**0.019***
	Control predictors:				
	Group		*4.73*	*2*	*0.094.*
	Ikela-Lenore-Louise	*12.61±6.29*			
	Lenore-Lolita-Louise	*8.14±6.70*			
	Time (time + time^2)	**1.59±0.44**	**9.15**	**1**	**0.0025***

Self-directed behaviour	(Intercept)	-2.81±0.60			
	Test predictors:				
	Condition	**-0.33±0.14**	**5.16**	**1**	**0.018***
	Control predictors:				
	Group		3.09	2	0.21
	Ikela/Lenore/Louise	-0.094±0.53			
	Lenore/Lolita/Louise	0.75±0.53			
	Time	**-0.31±0.13**	**4.42**	**1**	**0.035***

There was also a significant reduction in self-directed behaviour in the oxytocin compared to

placebo condition (β = -0.33, SE = 0.14, χ^2 = 5.16, p = 0.018, OR = 0.72 (95% CI: 0.54, 0.94); Table 1).

In our stability check analysis excluding each participant one-by-one in the model, the effect of

oxytocin on self-directed behaviour was significant in the model excluding Lenore and the model

excluding Lolita (excluding Ikela: β = -0.26, SE = 0.13, χ^2 = 3.52, p = 0.061; excluding Lenore: β = -

0.41, SE = 0.14, χ^2 = 8.89, p = 0.0029; excluding Lolita: β = -0.51, SE = 0.16, χ^2 = 9.81, p = 0.0017;

excluding Louise: β = -0.34, SE = 0.17, χ^2 = 3.65, p = 0.056). In our individual-level analysis, we

found a significant effect only for Ikela using a significance threshold of 0.0125 (Ikela: β = -0.71, SE

= 0.21, χ^2 = 12.19, p = 0.00048; Lenore: β = -0.15, SE = 0.28, χ^2 = 0.27, p = 0.60; Lolita: β = -0.59, SE

= 0.24, χ^2 = 5.58, p = 0.018; Louise: β = -0.58, SE = 0.41, χ^2 = 2.48, p = 0.11).

There were no significant differences in interindividual proximity (β = -0.15, χ^2 = 0.32, p = 0.57),

frequency of the abnormal behaviour regurgitation and reingestion (β = -1.05, SE = 0.66, χ^2 = 2.65,

p = 0.10), or rate of rest (β =-0.10, SE = 0.18, χ^2 = 0.34, p =0.56) between the oxytocin and placebo

condition. Bonobos engaged in GG-rubbing only once (oxytocin condition) and displayed no

aggression toward groupmates or any bouts of play during the observation period. See

supplementary material Table S2 for full details of all models.

Discussion:

We found that on the whole grooming rates of captive female bonobos were higher in the

oxytocin compared to saline placebo condition, though when conducting individual-level analyses

this was only significant after correcting for multiple testing in one participant (Louise). Still, in

combination with the previous studies showing oxytocin is released in bonobos and chimpanzees

following socio-positive interaction [7,8], our results provide some support for the biobehavioural

feedback loop hypothesis of oxytocin in social bonding for bonobos. The bonobos also engaged in

self-directed behaviour less in the oxytocin compared to placebo condition (though again this was

significant on an individual level in just one participant - Ikela), which is likely related to its

anxiolytic effect, though it should be noted we did not distinguish between kinds of self-directed

behaviours such as self-scratching or self-grooming which may have different relations to stress.

The proportion of rest and frequency of regurgitation and reingestion did not differ between

conditions, while GG-rubbing, play, and aggression were rarely or never observed during our 1-

hour observation window, likely due to low overall tension, precluding formal analysis.

Despite our overall model showing an increase in grooming in the oxytocin condition, our

individual-level analyses revealed only one individual, Louise, a 48-year-old female bonobo

relatively dominant to other groupmates. Lenore, a 38-year-old, also groomed groupmates more

in the oxytocin compared to the saline condition, but was not significant after correcting for

multiple testing. Ikela, a 29-year-old, groomed slightly more, but this effect was also not significant.

Lolita was never observed actively grooming throughout our experiment, and thus her rate of

grooming was completely unchanged by oxytocin. Future studies are needed to examine what

factors drive these potential individual differences. It is also important to note that given social

grooming necessarily requires a grooming partner, and given our experimental design

administering the same condition to all group members simultaneously, oxytocin's effect on the

group may not be entirely reducible to the individual level. Oxytocin may as a whole promote a

group dynamic more conducive to grooming, which is measurable in the behaviour of certain key

individuals. This hypothesis can be directly tested by administering oxytocin compared to saline

placebo only to Louise, and always saline placebo to others, and examining if the same effect is

found.

Although we present the first observation that oxytocin can promote socio-positive

interactions in female bonobos, there are several important limitations in this study. Due to

limited possibility of testing, enclosures suitable for detailed observation, and some apes'

willingness to join experiments, the sample was limited to just four adult female bonobos. This

limits firm conclusions about a biobehavioural feedback loop, as Crockford et al. [7] measured

urinary oxytocin after grooming only in chimpanzees, while Moscovice et al. [8] focused on sexual

interactions rather than grooming in bonobos. While these previous studies characterized the

changes similarly as socio-positive behaviours strengthening bond formation, conclusive proof of a

biobehavioural feedback loop in one or both species would require evidence of increased urinary

oxytocin following grooming in bonobos, administration of oxytocin promoting GG-rubbing in

bonobos, or administration of oxytocin promoting grooming among bond partners in chimpanzees.

It remains possible that exogenous oxytocin promotes grooming in bonobos but not chimpanzees,

or that oxytocin is released following grooming in chimpanzees but not bonobos. Of relevance,

Proctor et al. [24] did not find any behavioural change after administration of oxytocin on captive

chimpanzees, though they focused on oxytocin administration and observation to only one

individual at a time, instead of the group level as in the current study, limiting our ability to

directly compare the results. In addition, previous work has indicated sex-specific effects of

oxytocin [33–35], and thus it remains unclear whether our results can be generalized to different

sex pairs, though it should be noted that Crockford et al. [7] did not find significant differences

between female-female, female-male, and male-male dyads in urinary oxytocin level following

grooming in wild chimpanzees. We also could not investigate the possible effect of different

genetic background contributing to differences in the amount of change between conditions

across individuals, which should be directly explored in the future. Finally, the small number of

participants did not enable us to test the effect of existing social bond strength and relatedness

among groupmates, which may interact with the observed increase in grooming.

In conclusion, although limited, our finding offers some experimental evidence that

oxytocin can promote socio-positive interaction in bonobos. The oxytocin system may have been

coopted through evolution to the formation and maintenance of social bonds though a positive

feedback loop. Moreover, we suggest that oxytocin administration can be an effective tool in great

ape research involving free interactions among groupmates. Future work should further test

potential differences in oxytocin's effect between species, should examine inter-individual

variation with respect to social closeness and centrality, and should study how social contexts such

as feeding tension interact with this effect. We demonstrate that exogenous oxytocin can affect

great ape behaviour in naturalistic, spontaneous social interactions and provide experimental

support for the biobehavioural feedback loop hypothesis of oxytocin in the evolution of bonobo

social bonding.

Acknowledgements:

We thank the bonobos at Kumamoto Sanctuary for participating in this study. We also thank

Etsuko Nogami and the other caretakers at Kumamoto Sanctuary for their support throughout this

project, Dr. Satoshi Hirata for his advice and support, and Dr. Yutaro Sato for helping in data

collection. This study was funded by Japan Society for the Promotion of Science (KAKENHI

#21J21123 to J.B., #19H01772 and #20H05000 to F.K., and #19H00629 and #19H05736 to S.Y.)

References:

1. Nagasawa M, Okabe S, Mogi K, Kikusui T: **Oxytocin and mutual communication in mother-infant bonding**. *Frontiers in Human Neuroscience* 2012, **6**:31.

2. Rilling JK, Young LJ: **The biology of mammalian parenting and its effect on offspring social development**. *Science* 2014, **345**:771–776.

3. Da Costa APC, Guevara-Guzman RG, Ohkura S, Goode JA, Kendrick KM: **The Role of Oxytocin Release in the Paraventricular Nucleus in the Control of Maternal Behaviour in the Sheep**. *Journal of Neuroendocrinology* 1996, **8**:163–177.

4. Insel TR, Young LJ: **The neurobiology of attachment**. *Nature Reviews Neuroscience* 2001, **2**:129–136.

5. Nagasawa M, Ogawa M, Mogi K, Kikusui T: **Intranasal Oxytocin Treatment Increases Eye-Gaze Behavior toward the Owner in Ancient Japanese Dog Breeds**. *Frontiers in Psychology* 2017, **8**:1624.

6. Nagasawa M, Mitsui S, En S, Ohtani N, Ohta M, Sakuma Y, Onaka T, Mogi K, Kikusui T: **Oxytocin-gaze positive loop and the coevolution of human-dog bonds**. *Science* 2015, **348**:333–336.

7. Crockford C, Wittig RM, Langergraber K, Ziegler TE, Zuberbühler K, Deschner T: **Urinary oxytocin and social bonding in related and unrelated wild chimpanzees**. *Proceedings of the Royal Society B: Biological Sciences* 2013, **280**:20122765.

8. Moscovice LR, Surbeck M, Fruth B, Hohmann G, Jaeggi A V., Deschner T: **The cooperative sex: Sexual interactions among female bonobos are linked to increases in oxytocin, proximity and coalitions**. *Hormones and Behavior* 2019, **116**:104581.

9. Wittig RM, Crockford C, Deschner T, Langergraber KE, Ziegler TE, Zuberbühler K: **Food sharing is linked to urinary oxytocin levels and bonding in related and unrelated wild chimpanzees**. *Proceedings of the Royal Society B: Biological Sciences* 2014, **281**:20133096.

10. Preis A, Samuni L, Mielke A, Deschner T, Crockford C, Wittig RM: **Urinary oxytocin levels in relation to post-conflict affiliations in wild male chimpanzees (Pan troglodytes verus)**. *Hormones and Behavior* 2018, **105**:28–40.

11. Samuni L, Preis A, Mundry R, Deschner T, Crockford C, Wittig RM: **Oxytocin reactivity during intergroup conflict in wild chimpanzees**. *Proceedings of the National Academy of Sciences* 2017, **114**:268–273.

12. Samuni L, Preis A, Deschner T, Wittig RM, Crockford C: **Cortisol and oxytocin show independent activity during chimpanzee intergroup conflict**. *Psychoneuroendocrinology* 2019, **104**:165–173.

13.   Samuni L, Preis A, Deschner T, Crockford C, Wittig RM: **Reward of labor coordination and**
hunting success in wild chimpanzees. *Communications Biology* 2018, **1**:138.

14.   Bauman MD, Murai T, Hogrefe CE, Platt ML: **Opportunities and challenges for intranasal**
oxytocin treatment studies in nonhuman primates. *American Journal of Primatology* 2018,
80:1–8.

15.   Dal Monte O, Noble PL, Costa VD, Averbeck BB: **Oxytocin enhances attention to the eye**
region in rhesus monkeys. *Front Neurosci* 2014, **8**:41.

16.   Parr LA, Modi M, Siebert E, Young LJ, Rd G: **Intranasal oxytocin selectively attenuates rhesus**
monkeys' attention to negative facial expressions. *Psychoneuroendocrinology* 2013,
doi:10.1016/j.psyneuen.2013.02.011.

17.   Ebitz RB, Watson KK, Platt ML: **Oxytocin blunts social vigilance in the rhesus macaque**.
Proceedings of the National Academy of Sciences 2013, **110**:11630–11635.

18.   Putnam PT, Roman JM, Zimmerman PE, Gothard KM: **Oxytocin enhances gaze-following**
responses to videos of natural social behavior in adult male rhesus monkeys.
Psychoneuroendocrinology 2016, **72**:47–53.

19.   Jiang Y, Platt ML: **Oxytocin and vasopressin flatten dominance hierarchy and enhance**
behavioral synchrony in part via anterior cingulate cortex. *Scientific Reports* 2018, **8**:1–14.

20.   Kotani M, Shimono K, Yoneyama T, Nakako T, Matsumoto K, Ogi Y, Konoike N, Nakamura K,
Ikeda K: **An eye tracking system for monitoring face scanning patterns reveals the**
enhancing effect of oxytocin on eye contact in common marmosets.
Psychoneuroendocrinology 2017, **83**:42–48.

21.   Cavanaugh J, Carp SB, Rock CM, French JA: **Oxytocin modulates behavioral and**
physiological responses to a stressor in marmoset monkeys. *Psychoneuroendocrinology*
2016, **66**:22–30.

22.   Smith AS, Ågmo A, Birnie AK, French JA: **Manipulation of the oxytocin system alters social**
behavior and attraction in pair-bonding primates, Callithrix penicillata. *Hormones and*
Behavior 2010, **57**:255–262.

23.   Brosnan SF, Talbot CF, Essler JL, Leverett K, Flemming T, Dougall P, Heyler C, Zak PJ: **Oxytocin**
reduces food sharing in capuchin monkeys by modulating social distance. *Behaviour* 2015,
152:941–961.

24.   Proctor D, Calcutt SESE, Burke K, de Waal FBMFBM: **Intranasal Oxytocin Failed to Affect**
Chimpanzee (Pan troglodytes) Social Behavior. *Animal behaviour and cognition* 2016,
3:150–158.

25.   Hall K, Smith M, Russell JL, Lambeth SP, Schapiro SJ, Brosnan SF: **Chimpanzees Rarely Settle**
405   **on Consistent Patterns of Play in the Hawk Dove, Assurance, and Prisoner's Dilemma**
406   **Games, in a Token Exchange Task.** *Animal behavior and cognition* 2019, **6**:48–70.

407   26.   Brooks J, Kano F, Sato Y, Yeow H, Morimura N, Nagasawa M, Kikusui T, Yamamoto S:
408   **Divergent effects of oxytocin on eye contact in bonobos and chimpanzees.**
409   *Psychoneuroendocrinology* 2021, **125**:105119.

410   27.   MacDonald E, Dadds MR, Brennan JL, Williams K, Levy F, Cauchi AJ: **A review of safety, side-**
411   **effects and subjective reactions to intranasal oxytocin in human research.**
412   *Psychoneuroendocrinology* 2011, **36**:1114–1126.

413   28.   Brooks J, Onishi E, Clark IR, Bohn M, Yamamoto S: **Uniting against a common enemy:**
414   **Perceived outgroup threat elicits ingroup cohesion in chimpanzees.** *PLOS ONE* 2021,
415   **16**:e0246869.

416   29.   R Core Team: **R: A Language and Environment for Statistical Computing.** *R Foundation for*
417   *Statistical Computing* 2019,

418   30.   Bates D, Mächler M, Bolker B, Walker S: **Fitting Linear Mixed-Effects Models Using {lme4}.**
419   *Journal of Statistical Software* 2015, **67**:1–48.

420   31.   Barr DJ, Levy R, Scheepers C, Tily HJ: **Random effects structure for confirmatory hypothesis**
421   **testing: Keep it maximal.** *Journal of Memory and Language* 2013, **68**:255–278.

422   32.   Christensen RHB: **ordinal: Regression Models for Ordinal Data.** 2019,

423   33.   Gao S, Becker B, Luo L, Geng Y, Zhao W, Yin Y, Hu J, Gao Z, Gong Q, Hurlemann R, et al.:
424   **Oxytocin, the peptide that bonds the sexes also divides them.** *Proceedings of the National*
425   *Academy of Sciences of the United States of America* 2016, **113**:7650–4.

426   34.   Caldwell HK: **Oxytocin and sex differences in behavior.** *Current Opinion in Behavioral*
427   *Sciences* 2018, **23**:13–20.

428   35.   Dumais KM, Bredewold R, Mayer TE, Veenema AH: **Sex differences in oxytocin receptor**
429   **binding in forebrain regions: Correlations with social interest in brain region- and sex-**
430   **specific ways.** *Hormones and Behavior* 2013, **64**:693–701.

431