1 **Chronic adolescent exposure to cannabis in mice leads to sex-biased changes in**

2 **gene expression networks across brain regions**

Yanning Zuo[1,2], Attilio Iemolo[5], Patricia Montilla-Perez[5], Hai-Ri Li[5], Xia Yang[1,3,4],

Francesca Telese[5]

[1] Department of Integrative Biology and Physiology, University of California, Los

Angeles, CA

[2] Neuroscience Interdepartmental Program, University of California Los Angeles, CA

[3] Institute for Quantitative and Computational Biosciences, University of California, Los

Angeles, CA

[4] Brain Research Institute, University of California, Los Angeles, CA

[5] Department of Medicine, University of California, San Diego, CA

Corresponding author: Francesca Telese, Ph.D.

George Palade Laboratories, Room 124

University of California, San Diego

9500 Gilman Dr, Mail code 0640

La Jolla, California 92093

Phone: 858 -246-2607

Email: ftelese@ucsd.edu

Short title: Transcriptional changes after THC adolescence exposure

23
24
25

Abstract

During adolescence, frequent and heavy cannabis use can lead to serious adverse

health effects and cannabis use disorders (CUD). Rodent models of adolescent

exposure to the main psychoactive component of cannabis, delta-9-

tetrahydrocannabinol (THC), mimic the behavioral alterations observed in adolescent

users. However, the underlying molecular mechanisms remain largely unknown. Here,

we treated female and male mice with high doses of THC during early adolescence and

assessed their memory and social behaviors in late adolescence. We then profiled the

transcriptome of five brain regions involved in cognitive and addiction-related

processes. We applied gene coexpression network analysis and identified gene

coexpression modules, termed cognitive modules, that simultaneously correlated with

THC treatment and memory traits reduced by THC. The cognitive modules were related

to endocannabinoid signaling in the female dorsal medial striatum, inflammation in the

female ventral tegmental area, and synaptic transmission in the male nucleus

accumbens. Moreover, cross-brain region module-module interaction networks

uncovered intra- and inter-region molecular circuitries influenced by THC. Lastly, we

identified key driver genes of gene networks associated with THC in mice and genetic

susceptibility to CUD in humans. This analysis revealed a common regulatory

mechanism linked to CUD vulnerability in the nucleus accumbens of females and

males, which shared four key drivers (*Hapln4, Kcnc1, Elavl2, Zcchc12*). These genes

regulate transcriptional subnetworks implicated in addiction processes, synaptic

transmission, brain development, and lipid metabolism. Our study provides novel

insights into disease mechanisms regulated by adolescent exposure to THC in a sex-

and brain region-specific manner.

Introduction

Cannabis remains the most widely used psychoactive drug worldwide, particularly

among adolescents and young adults[1]. Recent data showed that more than one-third of

12th graders in the US used cannabis in the past year, reflecting an overall decline in

the perceived risk of regular cannabis use among adolescents[2]. The high prevalence

rates of cannabis use among adolescents pose a significant concern as cannabis

misuse by adolescents can lead to persistent cognitive impairments in learning,

attention, and memory[3–8]. Moreover, the early onset of cannabis use before 16 years of

age increases the risk of developing psychiatric disorders, including cannabis use

disorders (CUD) [9,10]. CUD has a strong genetic component (50-70% heritability). Recent

large-scale genome-wide association studies (GWAS) began to identify genetic variants

associated with CUD[11–15]. They also revealed a genetic correlation of CUD with other

mental health traits, including substance abuse, schizophrenia, and risk-taking[11–15].

However, in line with the nature of complex disease traits, common genetic variants

associated with lifetime cannabis use can explain only 11% of the phenotypic variance,

as revealed by one meta-analysis of eight GWAS[15]. It is possible that other

environmental factors, including cannabis exposure during critical developmental

periods, might affect molecular networks in critical brain regions and, in turn, influence

the development and severity of CUD.

The primary psychoactive component of cannabis is delta-9-tetrahydrocannabinol

(THC). The biological effects of THC are mediated mainly by members of the G protein-

coupled receptors (GPCR) family, such as cannabinoid receptors 1 (CB1R) and 2

(CB2R)[16]. The cannabinoid receptors, together with endogenous cannabinoids and the

enzymes responsible for their biosynthesis and metabolism, constitute the

endocannabinoid (eCB) system[17–19]. The eCB system plays a critical role in the

maturation of brain circuits during adolescence by regulating excitatory and inhibitory

neurotransmission[20]. Moreover, the fluctuations of eCB signaling during adolescence

influence the pubertal changes in gonadal hormone secretion[21]. This interaction

between eCB signaling and gonadal functions contributes to the emergence of sex-

biased behaviors during adolescence, including social, cognitive, and emotional

behaviors[22]. Substantial evidence from human or animal model studies has led to the

hypothesis that excessive exposure to THC during adolescence may disrupt the

physiological function of the eCB system, ultimately leading to sex-specific behavioral

abnormalities and increased risk for psychopathology later in life [5,22–25].

Despite this knowledge, there is limited data on genes and pathways affected by

adolescent exposure to THC in different brain regions of the female and male brains.

Two recent studies analyzed gene expression changes following chronic adolescent

exposure to THC in rats[26,27]. These studies demonstrated that chronic adolescent

exposure to THC alters gene expression networks that are associated with the

structural maturation of cortical cells in the prefrontal cortex (PFC) and with reward and

stress reactivity in the basolateral amygdala (Amy). However, these analyses were

limited to one brain region in male rats [26,27]. Therefore, more research is needed to

dissect brain region-specificity and cross-brain networks perturbed by chronic

adolescent exposure to THC in the female and male brains.

In this study, we treated female and male mice with THC during early adolescence and

assessed their behavior in late adolescence. We performed RNA sequencing (RNA-

seq) on five brain regions involved in cognitive and addiction-related processes,

including the prefrontal cortex (PFC), nucleus accumbens (NAc), dorsal medial striatum

(DMS), amygdala (Amy), and ventral tegmental area (VTA). We conducted gene

coexpression network analysis for each sex, within and between brain regions. Lastly,

we performed integrative genomic analyses of coexpression networks altered by THC in

mice and human genetic data from GWAS of CUD. This analysis identified genes and

pathways involved in THC-mediated behavioral aberrations in mice and linked them to

CUD in humans.

Methods and Materials

Detailed descriptions of experimental design and methods are included in the

Supplemental Methods. All methods used in this study have been published recently[28–

31]. RNA-seq datasets were deposited on the GEO database (accession GSE189821).

Results

THC exposure during adolescence impairs cognitive behaviors in a sex-specific

manner

To assess the behavioral effects of adolescent exposure to THC, we chronically treated

female and male mice with 10mg/kg of THC in early adolescence (5-7 weeks). We

assessed object recognition memory, social interaction, and anxiety-like behaviors in

late adolescence, two weeks after the last THC administration (Fig 1A).

By using the six different objects test (6-DOT)[28,32], we measured the effect of THC on

object recognition memory. THC decreased the absolute (D1) and relative (D2)

discrimination indexes by 36% (treatment effect $F_{(1,41)}=7.12$, $p=0.011$) and 30%

(treatment effect $F_{(1,42)}=5.3$, $p=0.026$), respectively, compared to the vehicle control

group (Fig 1B-C). There were no significant sex x treatment interactions for D1

($F_{(1,41)}=3.86$, $p=0.056$) or D2 ($F_{(1,42)}=1.89$, $p>0.1$) indexes. However, exploratory

posthoc analysis showed that, compared to the vehicle, THC decreased the D1 index

by 44% ($t=3.35$, $p=0.009$) and the D2 by 39% ($t=2.6$, $p=0.055$) in females but not in

males ($p>0.1$, Fig. 1B-C). Moreover, we observed a significant sex x treatment

interaction ($F_{(1,42)}=4.6$, $p=0.038$) in novel object exploration. A posthoc analysis

revealed that THC-treated females, but not males, tended to reduce novel object

exploration ($t=2.44$, $p=0.08$). The impact of THC was specific for the cognitive

components of this assay, as THC had no detectable effect on the distance traveled

during habituation (Fig. S1A) nor on the total time mice spent exploring the objects (all

main effects and interactions $p>0.1$, Fig. S1B-C).

We also tested the effect of THC on social behaviors using the three-chamber

interaction test. Adolescent exposure to THC significantly decreased social preference

by 12% ($F_{(1,41)}=6.3$, $p=0.016$) compared to the vehicle group (Fig. 1E). In contrast, sex

had no detectable effect on treatment (sex x treatment $F_{(1,41.5)}=0.1$, $p>0.1$).

Lastly, we examined anxiety-like behaviors with the elevated plus-maze. We did not

observe significant differences in the willingness of mice to explore open environments

(all main effects and interactions $p>0.1$, Fig. S 1D).

We excluded potential confounding effects on the exploratory activity by showing that

THC treatment was not associated with changes in body weight at the time of

behavioral testing (main effects and interactions p>0.1, Fig. S1E).

Overall, the behavioral analysis showed that exposure to THC during adolescence

impaired memory and social interaction in late adolescence, with more robust memory

deficits in female mice.

Identification of DEGs associated with chronic adolescent exposure to THC

To gain insights into the neurobiological mechanisms underlying the behavioral

alterations induced by THC, we profiled the transcriptome of PFC, DMS, NAc, Amy, and

VTA from the vehicle and THC-treated mice (n=6/tissue/treatment/sex). Principal

component analysis (PCA) validated the accuracy of brain dissection, showing that the

transcriptomes of each brain region differed from one other (Fig. 2A). Furthermore, PCA

plots within each brain region revealed substantial sex differences (Fig. S2A). In

contrast, the effect of THC treatment was subtle (Fig. S2A). Next, we identified

differentially expressed genes (DEGs) from each brain region and sex using different

significance cutoffs (Fig. 2B, Table S1). We found the largest number of DEGs in the

female Amy (n=743) and, in general, more DEGs in females in most brain regions

except for NAc, which had the largest number of DEGs in males (n=77) at a false

discovery rate (FDR)<0.1 and log fold change (logFC)>0.4 (Fig. 2B). Pathway analysis

showed overlapping pathways altered by THC in female Amy and DMS, including opioid

signaling and GPCR ligand binding (Fig. 2C). Moreover, DEGs in female Amy were also

related to long-term potentiation, axon guidance, addiction, retrograde cannabinoid

signaling, and translation (Fig. 2C). In contrast, DEGs in male NAc were involved in

interferon signaling and ubiquitin-mediated proteolysis (Fig. 2C).

To further explore the influence of sex on transcriptional responses to THC, we

compared gene expression changes between males and females. At a statistical cutoff

of FDR <0.1, this analysis yielded significant sex x treatment interactions for DEGs in

Amy (n=106), DMS (n=7), and NAc (n=13), but not for PFC and VTA (Fig. 2D, Fig. S2B-

D). Pathway analysis revealed that the genes upregulated by THC in the female Amy

but not in males (Fig. S2B) were linked to presynaptic SNARE complex formation and

several signal transduction pathways (e.g., mTOR, Wnt, oxytocin signaling (Fig. 2E). In

contrast, other DEGs groups with sex x treatment interactions were not enriched in

specific pathways (Fig. S2C-D).

In agreement with DEG analysis, threshold-free RRHO analysis showed minimal

overlap in DEGs when we compared gene expression changes between most pairs of

brain regions or between sexes (Suppl. Fig. 3).

These results indicate that females and males responded differently to THC in a brain

region-specific manner.

Identification of gene coexpression networks correlated with THC treatment and

cognitive traits.

Genes usually do not act alone but work as a network to achieve a biological function by

interacting in a signal transduction or metabolic pathway [33]. To better understand how

THC impacts biological networks, we applied WGCNA[34], a gene network modeling

approach, to identify groups of genes (modules) highly coexpressed or coregulated in

response to THC treatment within each brain region in each sex (Table S3). We then

performed trait-module correlation analysis using cognitive behavioral traits measured in

each mouse (Table S4). This analysis identified 27 modules across different brain

regions significantly correlated ($p<0.05$) with THC treatment that we referred to as

"THC-correlated modules" (Fig. 3A). While we did not observe any overlap between

THC-correlated modules and those correlated with social preference, we identified five

modules that simultaneously correlated with THC treatment and memory traits (Fig. 3B,

S4A-B). Therefore, we will refer to these modules as "cognitive modules". The cognitive

modules included female DMS *saddlebrown (*Fig. 3C), female VTA *bisque4* (Fig. 3D)

and *lightsteelblue1 (*Fig. 3E), and male NAc *orange* (Fig. 3F) and *darkgrey* (Fig. S4C).

Pathway enrichment analysis showed that DMS *saddlebrown* module was related to the

metabolism of the endogenous cannabinoid anandamide and cognitive disorders, such

as Alzheimer's disease (Fig. 3C). The VTA *bisque4* module was enriched in interferon

signaling and purinergic receptor genes (Fig. 3D), and the VTA *lightsteelblue1* module

was involved in non-neuronal differentiation processes (Fig. 3E). In contrast, genes

related to synaptic transmission were enriched in the male NAc orange module (Fig.

3F), but no pathway enrichment was identified for the male NAc *darkgrey* module.

Notably, only female-specific modules showed a positive correlation with memory traits

but a negative correlation with THC treatment (Fig. 3B). This observation suggests that

the female cognitive modules regulate memory formation but are disrupted by THC

treatment, reflecting the behavioral deficits observed in female mice. In contrast, the

male NAc *darkgrey* showed positive correlations with both THC and memory traits but

orange showed a negative correlation with both THC and memory traits (Fig. 3B),

suggesting that the relationship between THC treatment and memory is more complex

in males.

To identify the cell types that might contribute to the formation of cognitive modules, we

performed cell-type marker enrichment analysis. In support of the pathway analysis,

neuronal markers were enriched in the female DMS *saddlebrown* module and male NAc

orange module, while markers of astrocytes and microglia were enriched in the female

VTA *bisque4* and *lightsteelblue1* modules (Fig. 3G).

We found that THC-correlated modules and DEGs only partially overlapped (Fig. S5),

indicating that network analysis captures additional information about THC

transcriptional responses that go beyond changes in individual DEGs.

Overall, these results revealed that the effect of THC on memory is correlated with the

regulation of sex- and brain region-specific gene coexpression modules.

Cross-brain region module-module interactions affected by THC.

During adolescence, dynamic changes in the eCBs coincide with the remodeling of

circuit connectivity within and between brain regions, including corticolimbic

structures[35–37]. To better understand the impact of THC on cross-brain gene

coexpression networks in the female and male brains, we analyzed the correlations

between modules within and between brain regions. This analysis identified numerous

"THC-interconnected modules" that we define as those significantly correlated with

THC-correlated modules in the same region or across brain regions with correlation

coefficient $|r| > 0.5$ and $p < 0.05$ in females (Fig. 4A) and males (Fig. 4B). These modules

likely reflect gene networks indirectly influenced by THC. Many THC-interconnected

modules were also correlated with memory traits (colored nodes in Fig. 4A-B). As

shown by the Sankey diagrams in Figure 4C-D, positive and negative correlations were

relatively balanced across brain regions in both sexes. In females, higher levels of

connectivity were observed between Amy-Amy modules, followed by Amy-VTA, VTA-

VTA, DMS-DMS, and Amy-DMS (Fig. 4E). In males, higher levels of connectivity were

observed between NAc-NAc, followed by NAc-VTA, NAc-PFC, VTA-NAc, and VTA-VTA

(Fig. 4F).

These findings suggest that adolescent exposure to THC leads to changes in molecular

circuitries across different brain regions in a sex-specific manner.

Associations between coexpression modules altered by THC and human

cannabis use disorders

Recent GWAS have started to identify genetic variants associated with CUD[38]. To gain

further insights into the genes and pathways associated with CUD, we applied the

Mergeomic pipeline[31,39] to integrate the human CUD GWAS signals with THC-

correlated gene coexpression networks for each brain region and sex in mice (Fig. 5A).

We defined "CUD-associated modules" as those enriched in CUD-associated genes

informed by human GWAS (Table S6). There was no overlap between CUD-associated

modules and THC-correlated modules in females or minimal overlap (11%, 2 modules)

in males (Fig. 5B). In contrast, the overlap increased to 81.8% (9 modules) in females

and 55.7% (11 modules) in males when we included THC-interconnected modules in

the analysis (Fig. 5C). These results suggest that CUD-associated modules are likely

indirectly affected by THC.

To predict potential key regulators of CUD-associated networks, we performed key

driver (KD) analysis using tissue-specific Bayesian networks that infer causal

relationships between genes and that were constructed using independent human and

mouse data[39]. We identified top KD regulators of gene coexpression networks

associated with THC and CUD (overlapping modules in Fig. 5C) in males and females

(Table S7). While most brain regions showed sex-specific KD genes, the NAc showed

an overlap of four KD genes between males and females (*Hapln4, Kcnc1, Elavl2,*

Zcchc12, Fig. 5D*),* suggesting a common regulatory mechanism linked to CUD. Among

these genes, two encode for membrane proteins involved in the modulation of synaptic

plasticity. KNCN1 is a voltage-gated potassium channel [40], and HAPLN4 is a

component of the perineuronal net[41]. The other two KDs encode proteins involved in

transcriptional regulation. ELAVL2 is an RNA-binding protein involved in splicing in

neuronal development[42], and ZCCHC12 is a neuronal transcriptional coactivator[43]. Of

note, two KD genes, *Hapln4* and *Elavl2,* were also identified as CUD-associated genes.

The pathways analysis of the four KD-associated subnetworks shared between males

and females revealed that they regulate genes implicated in addiction processes,

neurotransmission, brain development, and lipid metabolism (Fig. 5D).

This integrative genomic analysis identified a connection between genes and pathways

altered by THC and associated with CUD vulnerability.

Discussion

Our work provides the first comprehensive, tissue- and sex-specific view of molecular

processes perturbed by adolescent THC treatment in mice and linked to CUD in

humans (Fig. 6). We identified gene coexpression networks disrupted by THC in

specific brain regions and correlated to memory deficits induced by THC in a sex-

specific manner. In addition, we identified key regulators that orchestrate brain region-

specific transcriptional subnetworks linked to adolescent exposure to THC and CUD

vulnerability.

In line with previous reports in rodent models[28,44–50], our behavioral analysis

demonstrated that adolescent exposure to THC in mice led to long-term impairments in

object recognition memory and social interaction, but not in anxiety-like behaviors. Our

study also showed sex differences in the effects of THC on recognition memory, which

was impaired more in females compared to male mice. Although previous preclinical

studies have not examined the influence of sex on the effects of THC on memory,

female rats have been reported to be more susceptible than males to the effects of THC

on locomotor activity, nociception, and reward processes[51–55]. In addition, clinical

studies showed that females are more sensitive to the harmful effects of THC on spatial

memory[56–60].

In agreement with the sexual dimorphism observed for the behavioral effects of THC,

we also reported extensive sex differences in gene expression patterns in response to

THC. First, female mice showed a larger number of DEGs across different brain regions

compared to males. Secondly, when we conducted a statistical analysis that explicitly

tested for sex differences in DEGs, we found significant sex x treatment interaction in

Amy, NAc, and DMS, suggesting that females and males respond differently to THC.

These results are new, as prior research on the transcriptional effects of adolescent

exposure to THC has focused only on male rats[26,27]. Miller et al reported that

adolescent exposure to THC in male rats was associated with gene expression changes

related to cytoskeleton and chromatin regulation in the PFC[26]. In contrast, we did not

identify any DEGs or THC-correlated modules in the male PFC in mice. Between-

species differences our different statistical thresholds in DEG analysis may explain the

discrepancy.

Which brain region drives distinct behavioral abnormalities induced by adolescent

exposure to THC is not entirely known. Our work, for the first time, simultaneously

investigated five brain regions. Our results indicated an extensive brain region

specificity in the genes and networks altered by THC. Amy and NAc may be sites of

particular importance as they were associated with the largest number of DEGs in

females and males, respectively. In line with these findings, brain morphological studies

of human cannabis users have shown that marijuana use may be associated with

disrupting the neural organization of the Amy and NAc[61]. In particular, previous studies

have documented that female teenagers who use marijuana are more susceptible than

males to structural abnormalities of the Amy, which were correlated with worse

internalizing symptoms[62]. Consistent with these observations, animal studies also

reported perturbation of synaptic transmission in the Amy and NAc following

administration of exogenous cannabinoids[63,64].

Moreover, our gene co-expression analysis also identified specific brain regions linked

to memory traits in mice. Specifically, the disruption of female cognitive modules in DMS

(*saddlebrown,* enriched for eCB related pathways) and VTA (*bisque4* and

lightsteelblue1, enriched for immune and non-neuronal differentiation pathways) was

correlated with the deficits in recognition memory observed in female mice. In

agreement with the pathway annotation of these modules, it is well known that the

endocannabinoid system plays an essential role in learning and memory processes,

with the engagement of the dorsal striatum specifically in encoding habit-related

memories[65–67]. Moreover, repetitive exposure to synthetic cannabinoids led to

inflammatory phenotypes, including astrogliosis in the VTA[68].

Our multiple brain region studies also allowed us to uniquely infer network connections

within and between brain regions. The THC-interconnected modules are likely indirectly

influenced by THC, as inferred from the module-module interaction network. We

speculate that the inter-region interaction network could predict how THC directly affects

one brain region that then cascades down to other brain regions. For example, our

analysis suggests that adolescent exposure to THC alters neural circuits that connect

Amy with VTA and DMS in females and neural circuits that connect NAc with VTA and

PFC in males. Future experiments perturbing the THC modules using animal models

will be necessary to validate these predictions.

Cannabis use disorder has a strong genetic component and is influenced by other

environmental factors, including social and developmental vulnerability. For example,

early initiation age in adolescence and a high frequency of cannabis use increase the

risk of CUD[10]. Our integrative genomic analysis identified CUD-THC subnetworks and

potential causal regulators, including four KD genes (*Hapln4, Kcnc1, Elavl2, Zcchc12)*

shared between male and female NAc and implicated in addiction processes, synaptic

transmission, brain development, and lipid metabolism. However, the involvement of

these genes in CUD has not been explored before, and follow-up studies will be needed

to confirm the role of the key driver genes in mediating THC effects on CUD in vivo.

Our results should be considered in light of certain limitations. First, we focus on

correlating transcriptomic changes that occur in late adolescence with behaviors

measured at the same time. The advantage of this approach is that it can capture gene

expression changes associated with a history of early adolescent exposure to THC.

However, it cannot directly assess the transcriptional and behavioral changes occurring

while the drug is onboard. The second limitation of our study is that we focus on

cognitive behaviors, such as recognition memory and social interaction. Still, other

behaviors are likely to be influenced by THC, including addiction-like phenotypes.

However, it is important to note that the addictive properties of THC are not well

modeled in mice. Thirdly, our study is limited to 5 brain regions and can miss other

additional gene regulatory networks. Thus, it would be important to expand this study to

other brain regions, such as the hippocampus, given its critical role in recognition

memory. Lastly, we cannot exclude that mechanisms other than gene expression

changes contribute to sex-specific THC-related behaviors. For example, sex-specific

hormonal changes[69] or pharmacokinetic factors[70] during adolescence and the

differential density of cannabinoid receptors in the female and male brain[71] could

contribute to the sex differences observed in our study. Our findings open numerous

new hypotheses that warrant future experimental validation.

In summary, our study is the first to integrate gene expression profiles, GWAS, and

network modeling to reveal comprehensive sex- and brain region-specific view of

biological processes and regulators influenced by cannabis use and linked to CUD

vulnerability.

Acknowledgments

This work was supported by the National Institute on Drug Use, USA [DP1DA042232,

U01DA050239 to FT]. XY is supported by the National Center for Advancing

Translational Sciences UCLA CTSI Grant UL1TR001881.

This publication includes data generated at the UC San Diego IGM Genomics Center

utilizing an Illumina NovaSeq 6000 that was purchased with funding from a National

Institutes of Health SIG grant (S10OD026929)

We would like to acknowledge H. Taylor and J. Hightower for technical assistance.

Conflict of Interest

The authors have no conflicts of interest to disclose.

Supplementary information is available at MP's website

References

1. United Nations. *World Drug Report 2020.* (UN, 2021).

2. Johnston, L. D. *et al.* Monitoring the Future national survey results on drug use, 1975-2018: Overview, key findings on adolescent drug use. *Mahidol Popul. Q. Gaz.* (2019).

3. Curran, H. V. *et al.* Keep off the grass? Cannabis, cognition and addiction. *Nat. Rev. Neurosci.* **17**, 293–306 (2016).

4. Silins, E. *et al.* Young adult sequelae of adolescent cannabis use: an integrative analysis. *Lancet Psychiatry* **1**, 286–293 (2014).

5. Volkow, N. D. Effects of Cannabis Use on Human Behavior-Reply. *JAMA Psychiatry* **73**, 996 (2016).

6. Castellanos-Ryan, N. *et al.* Adolescent cannabis use, change in neurocognitive function,

and high-school graduation: A longitudinal study from early adolescence to young

adulthood. *Dev. Psychopathol.* **29**, 1253–1266 (2017).

7. Volkow, N. D. *et al.* Effects of Cannabis Use on Human Behavior, Including Cognition,

Motivation, and Psychosis: A Review. *JAMA Psychiatry* vol. 73 292 (2016).

8. Crean, R. D., Crane, N. A. & Mason, B. J. An evidence based review of acute and long-

term effects of cannabis use on executive cognitive functions. *J. Addict. Med.* **5**, 1 (2011).

9. Simpson, A. K. & Magid, V. Cannabis Use Disorder in Adolescence. *Child Adolesc.*

Psychiatr. Clin. N. Am. **25**, 431–443 (2016).

10. Connor, J. P. *et al.* Cannabis use and cannabis use disorder. *Nat Rev Dis Primers* **7**, 16

(2021).

11. Johnson, E. C. *et al.* A large-scale genome-wide association study meta-analysis of

cannabis use disorder. *Lancet Psychiatry* **7**, 1032–1045 (2020).

12. Sherva, R. *et al.* Genome-wide Association Study of Cannabis Dependence Severity, Novel

Risk Variants, and Shared Genetic Risks. *JAMA Psychiatry* **73**, 472–480 (2016).

13. Demontis, D. *et al.* Genome-wide association study implicates CHRNA2 in cannabis use

disorder. *Nat. Neurosci.* **22**, 1066–1074 (2019).

14. Agrawal, A. *et al.* A genome-wide association study of DSM-IV cannabis dependence.

Addict. Biol. **16**, 514–518 (2011).

15. Pasman, J. A. *et al.* GWAS of lifetime cannabis use reveals new risk loci, genetic overlap

with psychiatric traits, and a causal influence of schizophrenia. *Nat. Neurosci.* **21**, 1161–

1170 (2018).

16. Devane, W. A., Dysarz, F. A., 3rd, Johnson, M. R., Melvin, L. S. & Howlett, A. C.

Determination and characterization of a cannabinoid receptor in rat brain. *Mol. Pharmacol.*

34, 605–613 (1988).

17. Mechoulam, R. & Parker, L. A. The endocannabinoid system and the brain. *Annu. Rev.*

Psychol. **64**, 21–47 (2013).

18. Lu, H.-C. & Mackie, K. An Introduction to the Endogenous Cannabinoid System. *Biol.*

Psychiatry **79**, 516–525 (2016).

19. Di Marzo, V., Melck, D., Bisogno, T. & De Petrocellis, L. Endocannabinoids: endogenous

cannabinoid receptor ligands with neuromodulatory action. *Trends Neurosci.* **21**, 521–528

(1998).

20. Viveros, M. P. *et al.* The endocannabinoid system in critical neurodevelopmental periods:

sex differences and neuropsychiatric implications. *J. Psychopharmacol.* **26**, 164–176

(2012).

21. Meyer, H. C., Lee, F. S. & Gee, D. G. The Role of the Endocannabinoid System and

Genetic Variation in Adolescent Brain Development. *Neuropsychopharmacology* **43**, 21–33

(2018).

22. Sturman, D. A. & Moghaddam, B. The neurobiology of adolescence: changes in brain

architecture, functional dynamics, and behavioral tendencies. *Neurosci. Biobehav. Rev.* **35**,

1704–1712 (2011).

23. Hurd, Y. L. *et al.* Cannabis and the Developing Brain: Insights into Its Long-Lasting Effects.

The Journal of Neuroscience vol. 39 8250–8258 (2019).

24. Maccarrone, M., Guzman, M., Mackie, K., Doherty, P. & Harkany, T. Programming of

neural cells by (endo)cannabinoids: from physiological rules to emerging therapies. *Nat.*

Rev. Neurosci. **15**, 786–801 (2014).

25. Hurd, Y. L., Michaelides, M., Miller, M. L. & Jutras-Aswad, D. Trajectory of adolescent

cannabis use on addiction vulnerability. *Neuropharmacology* **76 Pt B**, 416–424 (2014).

26. Miller, M. L. *et al.* Adolescent exposure to Delta(9)-tetrahydrocannabinol alters the

transcriptional trajectory and dendritic architecture of prefrontal pyramidal neurons. *Mol.*

Psychiatry **24**, 588–600 (2019).

27. Ferland, J.-M. N. *et al.* Dose mediates the protracted effects of adolescent THC exposure

on reward and stress reactivity in males relevant to perturbation of the basolateral

amygdala transcriptome. *Mol. Psychiatry* (2022) doi:10.1038/s41380-022-01467-0.

28. Iemolo, A. *et al.* Reelin deficiency contributes to long-term behavioral abnormalities induced

by chronic adolescent exposure to Delta9-tetrahydrocannabinol in mice.

Neuropharmacology **187**, 108495 (2021).

29. Zhou, X., Barkley-Levenson, A. M., Montilla-Perez, P., Telese, F. & Palmer, A. A.

Functional validation of a finding from a mouse genome-wide association study shows that

Azi2 influences the acute locomotor simulant response to methamphetamine. *Genes Brain*

Behav. **20**, e12760 (2021).

30. Shu, L. *et al.* Shared genetic regulatory networks for cardiovascular disease and type 2

diabetes in multiple populations of diverse ethnicities in the United States. *PLoS Genet.* **13**,

e1007040 (2017).

31. Shu, L. *et al.* Mergeomics: multidimensional data integration to identify pathogenic

perturbations to biological systems. *BMC Genomics* **17**, 874 (2016).

32. Sannino, S. *et al.* Role of the dorsal hippocampus in object memory load. *Learn. Mem.* **19**,

211–218 (2012).

33. Fionda, V. Networks in Biology. in *Encyclopedia of Bioinformatics and Computational*

Biology (eds. Ranganathan, S., Gribskov, M., Nakai, K. & Schönbach, C.) 915–921

(Academic Press, 2019).

34. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

35. Lee, T. T.-Y., Hill, M. N. & Lee, F. S. Developmental regulation of fear learning and anxiety

behavior by endocannabinoids. *Genes Brain Behav.* **15**, 108–124 (2016).

36. Katona, I. & Freund, T. F. Endocannabinoid signaling as a synaptic circuit breaker in

neurological disease. *Nat. Med.* **14**, 923–930 (2008).

37. Peters, K. Z., Cheer, J. F. & Tonini, R. Modulating the Neuromodulators: Dopamine,

Serotonin, and the Endocannabinoid System. *Trends Neurosci.* **44**, 464–477 (2021).

38. Hillmer, A. *et al.* Genetic determinants of cannabis use: a systematic review protocol. *Syst.*

Rev. **9**, 190 (2020).

39. Ding, J. *et al.* Mergeomics 2.0: a web server for multi-omics data integration to elucidate

disease networks and predict therapeutics. *Nucleic Acids Res.* **49**, W375–W387 (2021).

40. Bee, S., Ringland, A. & Coutellier, L. Social impairments in mice lacking the voltage-gated

potassium channel Kv3.1. *Behav. Brain Res.* **413**, 113468 (2021).

41. Edamatsu, M. *et al.* Hapln4/Bral2 is a selective regulator for formation and transmission of

GABAergic synapses between Purkinje and deep cerebellar nuclei neurons. *J. Neurochem.*

147, 748–763 (2018).

42. Berto, S., Usui, N., Konopka, G. & Fogel, B. L. ELAVL2-regulated transcriptional and

splicing networks in human neurons link neurodevelopment and autism. *Hum. Mol. Genet.*

25, 2451–2464 (2016).

43. Li, H. *et al.* Human ZCCHC12 activates AP-1 and CREB signaling as a transcriptional co-

activator. *Acta Biochim. Biophys. Sin.* **41**, 535–544 (2009).

44. Sabran-Cohen, T., Bright, U., Mizrachi Zer-Aviv, T. & Akirav, I. Rapamycin prevents the

long-term impairing effects of adolescence Δ-9-tetrahydrocannabinol on memory and

plasticity in male rats. *Eur. J. Neurosci.* **54**, 6104–6122 (2021).

45. Jouroukhin, Y. *et al.* Adolescent Δ9-Tetrahydrocannabinol Exposure and Astrocyte-Specific

Genetic Vulnerability Converge on Nuclear Factor-κB-Cyclooxygenase-2 Signaling to

Impair Memory in Adulthood. *Biol. Psychiatry* **85**, 891–903 (2019).

46. Saravia, R. *et al.* Concomitant THC and stress adolescent exposure induces impaired fear

extinction and related neurobiological changes in adulthood. *Neuropharmacology* **144**,

345–357 (2019).

47. Murphy, M. *et al.* Chronic Adolescent Δ9-Tetrahydrocannabinol Treatment of Male Mice

Leads to Long-Term Cognitive and Behavioral Dysfunction, Which Are Prevented by

Concurrent Cannabidiol Treatment. *Cannabis Cannabinoid Res* **2**, 235–246 (2017).

48. Zamberletti, E., Gabaglio, M., Prini, P., Rubino, T. & Parolaro, D. Cortical

neuroinflammation contributes to long-term cognitive dysfunctions following adolescent

delta-9-tetrahydrocannabinol treatment in female rats. *Eur. Neuropsychopharmacol.* **25**,

2404–2415 (2015).

49. Swartzwelder, N. A. *et al.* Effects of ethanol, Δ(9)-tetrahydrocannabinol, or their

combination on object recognition memory and object preference in adolescent and adult

male rats. *Neurosci. Lett.* **527**, 11–15 (2012).

50. O'Tuathaigh, C. M. P. *et al.* Chronic adolescent exposure to Δ-9-tetrahydrocannabinol in

COMT mutant mice: impact on psychosis-related and other phenotypes.

Neuropsychopharmacology **35**, 2262–2273 (2010).

51. Fattore, L. *et al.* Cannabinoid self-administration in rats: sex differences and the influence

of ovarian function. *Br. J. Pharmacol.* **152**, 795–804 (2007).

52. Craft, R. M., Marusich, J. A. & Wiley, J. L. Sex differences in cannabinoid pharmacology: a

reflection of differences in the endocannabinoid system? *Life Sci.* **92**, 476–481 (2013).

53. Craft, R. M., Wakley, A. A., Tsutsui, K. T. & Laggart, J. D. Sex Differences in Cannabinoid 1

vs. Cannabinoid 2 Receptor-Selective Antagonism of Antinociception Produced by Δ9-

Tetrahydrocannabinol and CP55,940 in the Rat. *J. Pharmacol. Exp. Ther.* **340**, 787–800

(2012).

54. Romero, E. M. *et al.* Antinociceptive, behavioural and neuroendocrine effects of CP 55,940

in young rats. *Brain Res. Dev. Brain Res.* **136**, 85–92 (2002).

55. Tseng, A. H. & Craft, R. M. Sex differences in antinociceptive and motoric effects of

cannabinoids. *Eur. J. Pharmacol.* **430**, 41–47 (2001).

56. Hernandez-Avila, C. A., Rounsaville, B. J. & Kranzler, H. R. Opioid-, cannabis- and alcohol-

dependent women show more rapid progression to substance abuse treatment. *Drug*

Alcohol Depend. **74**, 265–272 (2004).

57. Schepis, T. S. *et al.* Gender differences in adolescent marijuana use and associated

psychosocial characteristics. *J. Addict. Med.* **5**, 65–73 (2011).

58. Bassir Nia, A. *et al.* The relevance of sex in the association of synthetic cannabinoid use

with psychosis and agitation in an inpatient population. *J. Clin. Psychiatry* **80**, 0–0 (2019).

59. Makela, P. *et al.* Low doses of delta-9 tetrahydrocannabinol (THC) have divergent effects

on short-term spatial memory in young, healthy adults. *Neuropsychopharmacology* **31**,

462–470 (2006).

60. Pope, H. G., Jr, Jacobs, A., Mialet, J. P., Yurgelun-Todd, D. & Gruber, S. Evidence for a

sex-specific residual effect of cannabis on visuospatial memory. *Psychother. Psychosom.*

66, 179–184 (1997).

61. Gilman, J. M. *et al.* Cannabis use is quantitatively associated with nucleus accumbens and

amygdala abnormalities in young adult recreational users. *J. Neurosci.* **34**, 5529–5538

(2014).

62. McQueeny, T. *et al.* Gender effects on amygdala morphometry in adolescent marijuana

users. *Behav. Brain Res.* **224**, 128–134 (2011).

63. Azad, S. C. *et al.* Circuitry for associative plasticity in the amygdala involves

endocannabinoid signaling. *J. Neurosci.* **24**, 9953–9961 (2004).

64. Kolb, B., Gorny, G., Limebeer, C. L. & Parker, L. A. Chronic treatment with Delta-9-

tetrahydrocannabinol alters the structure of neurons in the nucleus accumbens shell and

medial prefrontal cortex of rats. *Synapse* **60**, 429–436 (2006).

65. Marsicano, G. & Lafenêtre, P. Roles of the endocannabinoid system in learning and

memory. *Curr. Top. Behav. Neurosci.* **1**, 201–230 (2009).

66. Moreira, F. A. & Lutz, B. The endocannabinoid system: emotion, learning and addiction.

Addict. Biol. **13**, 196–212 (2008).

67. Goodman, J. & Packard, M. G. The influence of cannabinoids on learning and memory

processes of the dorsal striatum. *Neurobiol. Learn. Mem.* **125**, 1–14 (2015).

68. Pintori, N. *et al.* Repeated exposure to JWH-018 induces adaptive changes in the

mesolimbic and mesocortical dopaminergic pathways, glial cells alterations, and

behavioural correlates. *Br. J. Pharmacol.* **178**, 3476–3497 (2021).

69. Cooper, Z. D. & Craft, R. M. Sex-Dependent Effects of Cannabis and Cannabinoids: A

Translational Perspective. *Neuropsychopharmacology* **43**, 34–51 (2018).

70. Tseng, A. H., Harding, J. W. & Craft, R. M. Pharmacokinetic factors in sex differences in

Delta 9-tetrahydrocannabinol-induced behavioral effects in rats. *Behav. Brain Res.* **154**,

77–83 (2004).

71. López-Gallardo, M. *et al.* Maternal deprivation and adolescent cannabinoid exposure

impact hippocampal astrocytes, CB1 receptors and brain-derived neurotrophic factor in a

sexually dimorphic fashion. *Neuroscience* vol. 204 90–103 (2012).

Figure Legends

Figure 1: Adolescent exposure to THC reduced recognition memory and social

interaction in a sex-specific manner. (A) Timeline of the study design. (B) absolute

discrimination index D1 and (C) relative discrimination index D2 are shown as mean ±

95% confidence intervals showing decreased recognition memory in THC-treated

groups compared to vehicle controls. (D) Exploration time(s) of the novel object is

expressed as mean ± SEM showing a decrease only in female mice. (E) Social

preference (%) is expressed as mean ± SEM and is reduced in both females and males.

Main effects and interactions found using LMM analysis and posthoc comparisons p

values using Tukey HSD test.

Figure 2: Adolescent THC exposure induced sex-specific transcriptional changes.

(A) PCA visualization of male and female RNAseq samples across brain regions and

treatment conditions. (B) Number of DEGs across brain regions and sexes using

different significance cutoffs in analyses within each sex and brain region. (C) Pathway

enrichment for DEGs in female Amy, DMS, and male NAc. Dot color depicts the

direction of regulation and dot size illustrates the significance. (D) Number of DEGs for

treatment by sex interaction across brain regions in analyses including both sexes for

each brain region. (E) Pathway enrichment for genes upregulated by THC only in

female Amy (sex x interaction). Dot size illustrates the significance.

Figure 3: Characterization of cognitive modules correlated with THC treatment

and mouse recognition memory. (A) Number of coexpression modules significantly

correlated with THC treatment (p<0.05). (B) Heatmap of cognitive modules correlated

with THC treatment and recognition memory. Color depicts the correlation coefficient

with THC or memory traits. *p<0.05; **p<0.01; ***p<0.001. (C-F) Visualization of

cognitive module networks and pathway annotations. The edges denote positive

correlations between pairs of genes. Only the top 100 edges based on topological

overlap weight were visualized due to the large size of the male NAc orange module.

(G) Cell type marker gene enrichment of cognitive modules.

Figure 4: Construction of THC-interconnected module map reveals potential intra-

or inter-region molecular circuitries disrupted by THC. (A-B) Visualization of female

(A) and male (B) THC-interconnected modules, which are correlated with THC-

correlated modules, with correlation coefficient |r|>0.5 and p<0.05. Nodes with filled

colors denote modules correlated with THC or memory traits. Nodes with orange

borderline depict THC-interconnected modules. (C-D) Sankey plots of female (C) and

male (D) THC-correlated module interactions. Link colors denote the direction of

correlation, with red indicating positive correlation and blue denoting negative

correlation. (E-F) Number of intermodular interactions in THC-interconnected modules

in females (E) and males (F). The color indicates the direction of correlation, with red

indicating positive correlation and blue denoting negative correlation.

Figure 5: Association of THC-related modules with human CUD. (A) Schematic of

Mergeomics pipeline. MDF: marker dependency filtering; MSEA: marker set enrichment

analysis; KDA: key driver analysis. (B) Overlap (%) between CUD-associated modules

and THC-correlated modules (C) or between CUD-associated modules and both THC-

correlated and interconnected modules (D). The overlap (%) is calculated as the

number of overlapping modules divided by the total number of CUD-associated

modules. (D) Visualization of Bayesian network shared by CUD-associated modules in

female and male NAc. Key driver genes are represented by large size diamond nodes.

Orange, blue, and grey nodes denote genes (male NAc orange module, genes in

female NAc blue module, and genes in the BN but not in the two aforementioned

modules, respectively. CUD-associated genes identified by Mergeomics using loci with

p<0.001 from the Johnson et al.[11] CUD GWAS are labeled with red borderline. Bar plot

depicts pathway enrichment of the genes in the CUD subnetwork. BN, Bayesian

network

Figure 6: Integrative genomic analysis identified sex- and brain region-specific

biological processes and regulators affected by THC and linked to CUD. (Top

panel) Behavioral analysis showed that adolescent exposure to THC leads to cognitive

impairments, with stronger memory deficits in female mice. (Middle panel) Gene

coexpression analysis identified sex- and brain region-specific cognitive modules

correlated with THC treatment and memory impairments. (Bottom panel) The integrative

genomic analysis identified potential causal regulators of gene networks associated with

THC treatment and CUD vulnerability.

Supplementary Figure 1: Behavioral characterization of female and male mice following

adolescent exposure to THC. **(A)** Locomotor activity is expressed as mean distance traveled

(m) ± SEM. Exploration (s) in familiarization phase T2 **(B)** and test phase T3 **(C)** are expressed

as mean ± SEM. **(D)** Bodyweight (g) is expressed as mean ± SEM for the THC administration

course (day 1 to 21) and day 36, 43, and 60 when the three behavioral assays started. Main

effects and interactions found using LMM analysis, and posthoc comparisons p values using

Tukey HSD test.

Supplementary Figure 2: Adolescent THC administration induced sex-specific

transcriptional changes. (A) PCA visualization of all samples in each brain region. Samples

are colored with sex and treatment conditions. (B) Heatmaps of treatment by sex interaction

DEGs in Amy. The left panel depicts DEGs upregulated in females with THC treatment, while

the right panel depicts DEGs upregulated in males with THC treatment. (C) Heatmaps of

treatment by sex interaction DEGs in NAc. (D) Heatmaps of treatment by sex interaction DEGs

in DMS. UP, upregulated; DN, downregulated.

Supplementary Figure 3. Minimal transcriptional overlap was detected in most

comparison pairs across sex and brain regions based on rank-rank hypergeometric

overlap (RRHO) analysis. (A) Schematic of RRHO heatmap interpretation. (B-D) RRHO

heatmaps show minimal overlap in transcription between pairs of brain regions within females

(B), males (C), and between sexes (D). (E-F) RRHO heatmaps show concordance (E) and

discordance (F) in transcription between pairs of brain regions. We define concordant as

positive correlation and discordant as negative correlation in DEG overlap patterns.

Supplementary Figure 4: **DEG enrichment in WGCNA coexpression modules was**

detected in a few brain regions. (A-B) Heatmaps of DEG enrichment in WGCNA coexpression

modules in individual brain regions. Select modules from female Amy, DMS, PFC, VTA and

male VTA, and NAc showed significant DEG enrichment. Female Amy had the largest number

of modules that showed DEG enrichment. Color depicts -log10 (P value) of the enrichment by

Fisher's exact test. *p<0.05; **p<0.01; ***p<0.001. DEG.all, DEG.up, and DEG.down denotes

all, up-regulated, and down-regulated DEGs of a certain brain region, respectively.

Supplementary Figure 5: Correlation of WGCNA coexpression modules with chronic THC

administration and mouse cognitive behavioral traits. (A-B) Heatmaps of module correlation

with THC treatment and cognitive traits in males (A) and females (B), respectively. Only brain

regions with significant correlation are shown. Color depicts correlation coefficient relative to

THC or memory traits. *p<0.05; **p<0.01; ***p<0.001. (C) Visualization of male NAc darkgrey

module. The edges denote positive correlations between pairs of genes. No pathway annotation

under the significance of FDR < 0.05 was found.

683
684
685

A RNA-seq

PC2: 32% variance / PC1: 51% variance

Brain region
- Amy
- NAc
- PFC
- DMS
- VTA

Condition
- ● THC
- ▲ Veh

B DEGs THC vs Veh

Number of DEGs

- FDR < 0.1 logFC > 0.4
- p < 0.01 logFC > 0.4

Female: Amy 743, 762; NAc 0, 32; PFC 2, 45; DMS 24, 84; VTA 12, 125
Male: Amy 0, 46; NAc 77, 273; PFC 0, 39; DMS 0, 14; VTA 6, 66

C DEGs THC vs Veh (FDR< 0.1 lgFC >0.4)

- Long-term potentiation
- Axon guidance
- Neurotransmitter receptor binding & downstream transmission
- Translation
- Opioid signaling
- Retrograde endocannabinoid signaling
- Cocaine/amphetamine addiction
- Signaling by GPCR
- GPCR ligand binding
- Neuroactive ligand-receptor interaction
- Interferon signaling
- Ubiquitin mediated proteolysis

Female Amy / Female DMS / Male NAc

-log10.FDR
- 2
- 3
- 4
- 5
- 6

Direction
- ● Up-regulated
- ● Down-regulated
- ● Both up and down-regulated

D Sex x Treatment (FDR < 0.1)

Number of DEGs: Amy 106, NAc 13, DMS 7

E DEG UP in Female Amy Sex x Treatment (FDR < 0.1)

- Morphine Addiction
- Wnt Signaling Pathway
- Oxytocin Signaling Pathway
- mTOR Signaling Pathway
- Disinhibition of SNARE Formation

-log10.FDR
1.4 1.5 1.6 1.7 1.8

Zuo et al. 2022, Sup Fig 5