Neural basis for anxiety and anxiety-related physiological responses during a driving situation:

An fMRI study

Takafumi Sasaoka[1], Tokiko Harada[1], Daichi Sato[2], Nanae Michida[2], Hironobu Yonezawa[2], Masatoshi

Takayama[2], Takahide Nouzawa[3], Shigeto Yamawaki[1]

1. Center for Brain, Mind, and KANSEI Sciences Research, Hiroshima University, Japan.

2. Mazda Motor Corporation, Hiroshima, Japan.

3. Office of Academic Research and Industry-Academia-Government and Community Collaboration,

Hiroshima University, Japan.

Corresponding author: Takafumi Sasaoka, PhD

Center for Brain, Mind, and KANSEI Sciences Research, Hiroshima University, 1-2-3 Kasumi,

Minami-ku, Hiroshima, 734-8551, Japan.

Phone: +81-82-257-1722; Fax: +81-82-257-1723

E-mail: tsasaoka@hiroshima-u.ac.jp

Running Title: An fMRI study on the neural basis of anxiety

Abstract

While the exteroceptive and interoceptive prediction of a negative event increases a person's

anxiety in daily life situations, the relationship between the brain mechanism of anxiety and the

anxiety-related autonomic response has not been fully understood. In this fMRI study, we examined

the neural basis of anxiety and anxiety-related autonomic responses in a daily driving situation.

Participants viewed a driving video clip in the first-person perspective. During the video clip,

participants were presented with a cue to indicate whether a subsequent crash could occur (attention

condition) or not (safe condition). Enhanced activities in the anterior insula, bed nucleus of the stria

terminalis, thalamus, and periaqueductal gray, and higher sympathetic nerve responses (pupil dilation

and peripheral arterial stiffness) were triggered by the attention condition but not with the safe

condition. Autonomic response-related functional connectivity was detected in the visual cortex,

cerebellum, brainstem, and MCC/PCC with the right anterior insula and its adjacent regions as seed

regions. Thus, the right anterior insula and adjacent regions, in collaboration with other regions play a

role in eliciting anxiety based on the prediction of negative events, by mediating anxiety-related

autonomic responses according to interoceptive information.

Keywords: anxiety, insula, peripheral arterial stiffness, pupillometry, sympathetic nerve response

Introduction

It has been emphasized that the fundamental function of the brain, as a "prediction machine," is prediction and prediction error processing (e.g., Clark 2013). The predictive coding theory (Friston 2005; Rao and Ballard 1999) has been proposed as a computational framework for brain prediction and prediction error processing. According to this theory, the brain works on the computational principle to minimize the prediction error (difference between the prediction of the cause of the sensory input and the actual sensory input). Thus, the cause of the sensory input is inferred: to minimize prediction errors, predictions are updated and actions are performed in order to change the sensory input and to confirm the predictions (active inference). Based on this theory, perception could be regarded as an inference of the cause of external sensation (i.e., exteroception) of the outer world by prediction and prediction-error processing of exteroception (Rao and Ballard 1999). Likewise, subjective feelings could be regarded as an inference of the cause of the bodily sensation (i.e., interoception) by prediction and prediction-error processing of interoception (e.g., Seth 2013, Barrett and Simmons 2015; Barrett et al. 2016; Barrett 2017).

Anxiety, a typical example of brain prediction processing, can be defined by anticipatory affective, cognitive, and behavioral changes in response to uncertainty regarding potential threats (Grupe and Nitshke 2013). When the brain predicts a threatening event that can disturb homeostasis, negative emotions are elicited and attention is directed to the incoming information; thus, actions are undertaken to avoid the event. This corresponds to a state of anxiety. The interoception processing

model based on the predictive coding theory (Seth 2013) assumes that the prediction of upcoming

threatening events results in updating the desired state of the organism, accompanied by autonomic

reflexes. Based on this assumption, we can predict that the elicitation of anxiety will trigger a

corresponding autonomic response. Understanding the neural basis for eliciting anxiety and

accompanying autonomic responses in a daily situation will contribute to understanding the

mechanisms of anxiety and anxiety disorders, as well as to implementing a system to appropriately

reduce anxiety by reading out anxiety-related autonomic responses.

Previous neuroimaging studies on anxiety disorder and phobia have suggested that the anterior

insula and amygdala play a key role in these conditions. A meta-analysis of neuroimaging studies on

anxiety disorder (Etkin and Wager 2007) showed that hyperactivation in the amygdala and insula was

more frequently related to social anxiety disorder and specific phobia than to posttraumatic stress

disorder. In line with this, Baur et al. (2013) demonstrated that state and trait anxiety were associated

with the functional and structural connectivity between the anterior insula and the amygdala. Specific

phobias, which are prevalent anxiety disorders, are associated with activation in the brain network

("fear network"; Sehlmeyer et al. 2009), including the medial prefrontal cortex, anterior cingulate

cortex (ACC), insula, and amygdala. Indeed, Schweckendiek et al. (2011) investigated fear learning in

arachnophobia using a fear conditioning paradigm. Their study showed that arachnophobia was linked

to enhanced activation in the fear network including the medial prefrontal cortex, ACC, insula,

thalamus, and amygdala. Moreover, they showed increased amygdala activation in response to the

phobia-related conditioned stimulus (CS), but not in response to the non-phobia-related CS. A meta-

analysis of fear conditioning neuroimaging studies (Fullana et al. 2016) showed that the fear CS

consistently activated the "central autonomic-interoceptive network," which includes the anterior

insula, dorsal ACC (dACC), and the subcortical viscero-sensory regions such as the midbrain

(periaqueductal gray [PAG] and parabrachial nucleus), ventral thalamus, hypothalamus, and the

pontomedullary junction. This activation reflects the involvement of the conscious experience of fear

and anxiety and the autonomic responses to threat. The amygdala involvement in the fear conditioning

was not apparent in the meta-analysis study. On the other hand, it has been reported that the extended

amygdala, including the bed nucleus of the stria terminalis (BNST), plays a crucial role in anxiety.

Indeed, Somerville et al. (2013) conducted an fMRI study showing that the BNST is associated with

the sustained response anticipating the aversive stimuli, whereas the amygdala is involved with the

transient response to such stimuli. Therefore, it is possible that the amygdala and the extended

amygdala have different roles in anxiety elicitation: the amygdala activity is related to fear itself,

whereas the BNST is more related to the anticipation of fearful events. In contrast, the default-mode

network (DMN), including the ventral medial prefrontal cortex and posterior cingulate cortex (PCC),

was deactivated when CS was presented relative to non-CS ("safe-signal") (Fullana et al. 2016). This

could yield various interpretations, such as fear-response inhibition, encoding episodic memory traces

of the conditioned-unconditioned stimuli (CS/US) association, and relief related to omission of the US.

However, the role of the safe-signal-related brain network remains unclear.

In the theory of anxiety disorders, some researchers have emphasized the deficit of the

interoceptive processing with the ability to mitigate anxiety. Paulus and Stein (2006) proposed that

individuals who are prone to anxiety show an altered interoceptive prediction in the anterior insula.

For the brain, excessive interoceptive prediction could need additional resources to reduce the

prediction error, resulting in increased anxiety in anticipation of aversive events. Interoceptive

information is often first processed in the brainstem, such as the medial nucleus of the solitary tract

and the parabrachial nucleus, and relayed to the insula by the thalamus (Chen et al. 2021). The

anterior insula has been suggested to be involved in the integration of interoceptive and exteroceptive

representations, as well as in their updating based on the incoming information (Craig 2009). Based

on these observations, we can speculate that perception of a cue associated with threatening events

triggers a prediction of negative body states generated by the anterior insula, and that accompanying

interoceptive information is processed by the subcortical regions, such as the thalamus and brainstem.

It is well known that autonomic responses, such as increasing heart rate (e.g., Deane 1961),

skin conductance (e.g., Epstein and Roupenian 1970), and pupil dilation (e.g., Bitsios et al. 2004) are

observed when people experience anxiety. However, there are few studies examining the brain

mechanisms involved in the autonomic responses related to anxiety. For instance, Wager et al. (2009a,

2009b) reported that the dorsal pregenual cingulate region was active in relation to the increase in the

heart rate while facing a socially threatening situation in which participants were waiting to deliver a

speech to others. However, many other indices reflect sympathetic nerve activity rather than the heart

rate. Peripheral arterial stiffness (β_{art}) has been proposed (Matsubara et al. 2018; Tsuji et al. 2021) as an index reflecting sympathetic activity. β_{art} can be estimated using electrocardiography, continuous sphygmomanometry, and photoplethysmography (PPG). Tsuji et al. (2021) demonstrated that β_{art} correlated with subjective pain ratings and brain activity in regions including the salience network (e.g., dACC), suggesting that β_{art} is a potential candidate for a sympathetic nerve index reflecting anxiety. Cardiovascular autonomic signals are often used as autonomic indices during driving, especially as a measure of mental workload (Paxion et al. 2014). For example, heart rate and a specific component of heart rate variability (0.10 Hz) reflect the mental workload during driving (e.g., Brookhuis and De Waard 1991; Liu and Lee 2006).

In the current study, we examined the brain mechanisms involved in eliciting anxiety and related to induced autonomic responses by using a situation of driving, one of the possible situations that would frequently cause anxiety in daily life. We performed simultaneous measurements via functional magnetic resonance imaging (fMRI) and other physiological indices, including electrocardiography (ECG), PPG, blood pressure monitoring, and pupillometry. In particular, we focused on pupil dilation and β_{art} as putative indications of sympathetic nerve activity. We hypothesized that the anterior insula and subcortical regions involved in threat and interoceptive processing (BNST, thalamus, and midbrain), are active in anticipation of the crash. We anticipated that if the amygdala was involved in the actual threatening event, its activity would increase in response to the crash, and that enhanced BNST activity would be detected in anticipation of the crash.

1 Moreover, we predicted significant physiological responses indicating sympathetic nerve activity (β_{art}

2 and pupillary dilation) in anticipation of the crash. These would hypothetically activate the network,

3 including the anterior insula, and/or modulate functional connectivity (FC) between the anterior insula

4 and other brain regions.

Materials and Methods

Experimental design

8 Study participants performed a simple reaction task while watching a driving video clip during an

9 MRI scan. Figure 1 shows a schematic of the experimental trial.

11 **Figure 1.** Schematic timeline of an experimental trial. In each trial, participants were presented with a

12 first-person perspective driving video clip. At 17 s following the video clip onset, participants were

13 presented with a word, either "Attention" or "Safe," for 2 s as a cue to indicate whether a subsequent

crash may occur (attention/crash condition) or not (safe condition). In the crash condition, a scooter

approached and hit the vehicle at 25 s following the video clip onset. At the end of the video clip, the

word "STOP" was presented for 1 s, and participants were asked to press a button as quickly as

possible. They received feedback as to whether their response was performed within 1 s following the

presentation of the word "STOP," via presentation of a word, either "GOOD" or "BAD.". We defined

the 10-s period from the video clip onset as "baseline," and the 8-s period from the cue onset as the

"anticipation period." See "Materials and Method" for details.

In the experiment, participants viewed a first-person perspective driving video clip generated by a

driving simulator (D3sim, Mitsubishi Precision Co., Ltd., Japan). The video clip displayed a vehicle

advancing through a narrow street for 26–28 s. At 17 s following the onset of the video clip, either the

word "Attention" or "Safe" was superimposed for 2 s on the video clip. Among the 16 trials in which

the word "Attention" appeared, six revealed a scooter approaching and hitting the vehicle, with the

presentation of a broken windshield and a collision sound (crash condition). In contrast, no scooter

approached the vehicle in the remaining 10 trials (attention condition) and in the 16 trials in which the

word "Safe" appeared (safe condition). Before the experiment, the participants were informed of the

potential scenarios after the presentation of each type of cue. At the end of each trial, the word "STOP"

was displayed for 1 s, upon which the participants had to press a button with their right index finger,

similar to a hard-braking event. The timing of the presentation of the word "STOP" was randomized

within 26–28 s following the video clip onset. Participants were presented with the word "GOOD" or "BAD" as feedback to indicate whether they responded within 1 s from the presentation of the word "STOP." We defined the 10-s period from the video clip onset as the "baseline" period and the 8-s period from the cue onset as the "anticipation" period.

After each trial, participants rated four items (anxiety, pleasantness, unpleasantness, and arousal), assessing their affective states during the anticipation period using a visual analog scale (VAS; Aitken 1969). For each item, the participants moved a cursor to indicate their rating (from 0 to 100) by pressing buttons with their middle and ring fingers. The experiment consisted of four sessions of eight trials each. Prior to the main trials, the participants performed three practice trials (one trial for each condition). Trait and state anxiety scores were obtained using the State-Trait Anxiety Inventory (STAI). Trait anxiety scores were obtained prior to the experiment. State anxiety scores were obtained before and after the experiment.

Visual stimuli were presented on an MRI-compatible, 32-inch liquid crystal display (NordicNeuroLab, Bergen, Norway) with a resolution of 1,920 × 1,080 pixels that subtended 30.4° × 17.4° visual angles. The participants perceived the visual stimuli through a mirror attached to the head coil. Auditory stimuli were presented with MRI-compatible, noise-canceling headphones (OptoActive, Optoacoustics Ltd., Or Yehuda, Israel). The volume of the auditory stimuli was adjusted to ensure that it was not too loud based on the participants' self-reports during the practice trials.

Participants

Thirty-four healthy adult participants completed this experiment (four women, all right-handed, aged 23.0 ± 2.12 [mean $\pm$ SD]). All participants had a driver's license and drove a car at least once a week. According to self-report data, none of the participants had a history of mental disorders. All participants provided written informed consent prior to participating in the study. This study was approved by the Research Ethics Committee of Hiroshima University (approval number E-965-3).

Analysis of behavioral data

The subjective rating data for each item were z-score normalized for each participant. To examine the differences in ratings among conditions and the occurrence of habituation, we performed a two-way repeated-measures analysis of variance (ANOVA) with factors of task condition and session for each item.

The reaction time (RT) data for each participant were averaged for each condition in the four sessions. To examine whether there were any differences in the mean RT, we performed a one-way repeated measures ANOVA with a factor of task condition. If any main effects or interactions were significant, we performed a post-hoc test using the modified Shaffer method.

Acquisition and analyses of autonomic responses

To examine autonomic responses related to the participants' anxiety, we measured the pupil area and blood pressure, and we performed ECG and PPG.

Pupil area

The pupil area of the participants' right eye was monitored using an MRI-compatible eye tracker (EyeLink 1000 Plus, SR Research, Osgoode, ON, Canada) at a sampling rate of 500 Hz. The pupil data were analyzed within the period from -2 s to 25 s of the onset (0 s) of the video clip. Data for one participant and those of six out of the remaining 132 sessions (33 participants × 4 sessions) were excluded from the statistical analysis owing to excessive eye blinks and measurement artifacts based on visual inspection. To avoid misrepresentation of a dataset, the data 200 ms before and after the eye blinks were replaced with "Not a Number" (NaN) since the data within this period were possibly distorted by eye blinks (Choe et al. 2016). To examine when a difference was statistically significant in the trial, we performed a paired t-test on the mean pupil areas between the conditions at each time point throughout the trial. Since we expected a difference in the autonomic responses between the attention and safe conditions during the anticipation period, and since the presentation of a cue itself could affect the autonomic responses right after the cue onset, we also calculated the mean pupil areas over the 5-s period from 3 s to 8 s following cue onset (20–25 s following video clip onset), and compared values between conditions using a paired t-test. We focused on the period before the crash (25 s following video clip onset); therefore, the crash and attention conditions were

collapsed (crash/attention condition) for this analysis.

Peripheral arterial stiffness

ECG signals were acquired from a three-lead ECG placed on the participant's chest. A PPG

was attached to the participant's left index finger. Systolic and diastolic blood pressure values during a

single heartbeat were measured using the CareTaker system (Biopac Systems, Goleta, CA, USA). A

blood pressure cuff was attached to the participant's left thumb. These physiological signals were

recorded using a Biopac MP 150 system at a sampling rate of 1,000 Hz. Since data could not be

obtained from one participant owing to a problem in the measurement device, the data of 33

participants were used for the statistical analysis. β_{art} was calculated for each heartbeat based on the

following equation (Tsuji et al. 2021):

$$\beta_{art} = \frac{\ln\left(\frac{P_{SYS}}{P_{DIA}}\right)}{P_{lmax} - P_{lmin}} \qquad (1)$$

where P_{lmax} and P_{lmin} are the maximum and minimum values of the PPG during a heartbeat,

respectively. P_{SYS} and P_{DIA} are systolic and diastolic blood pressures during a heartbeat. The ECG was

used to determine the R–R interval for extracting the P_{SYS}, P_{DIA}, P_{lmax}, and P_{lmin} for each heartbeat.

However, we could not robustly detect the R peaks in the ECG data obtained from most participants

owing to MRI artifacts. Therefore, we used the inverted PPG waveforms for each participant and

segmented the waveform between two adjacent peaks to determine each heartbeat.

1   To compare the β_{art} between the conditions, we calculated the ratio of the mean β_{art} averaged

2   over the 5-s period from 3 to 8 s following cue onset (20 to 25 s following video clip onset) to the

3   mean β_{art} averaged over the baseline, and subsequently performed a paired *t*-test.

MRI data acquisition

MRI data were acquired using a 3.0T MRI scanner (Siemens Magnetom Verio, Siemens AG,

Munich, Germany). The functional images were acquired using an echo-planar, T2*-weighted,

multiband gradient echo sequence with the following parameters: repetition time (TR), 1,000 ms;

echo time (TE), 30.00 ms; voxel size, $3.0 \times 3.0 \times 3.2$ mm; 42 slices; slice thickness, 3.2 mm; field of

view (FOV), 192 mm; flip angle, 80°; and acceleration factor, 3. The structural image of each

participant was acquired using T1-weighted, three-dimensional magnetization-prepared rapid

gradient-echo imaging (MPRAGE) with the following parameters: TR, 2,300 ms; TE, 2.98 ms; voxel

size, $1.0 \times 1.0 \times 1.0$ mm; flip angle, 9°; and FOV, 256 mm.

MRI data analysis

16   MRI data preprocessing and statistical analyses were performed using SPM12 software

17   (Wellcome Department of Cognitive Neurology, London, UK; www.fil.ion.ucl.ac.uk/spm). The first

18   10 echo-planar images (EPIs) were discarded to permit T1 equilibration effects. In order to account

for the correction of head movement, the remaining volumes were spatially realigned to the first of the

volumes and realigned to the mean of all the images. T1-weighted structural images were co-

registered with the first EPI of the remaining EPIs after discarding the first 10 EPIs. The co-registered

structural images were spatially normalized to the Montreal Neurological Institute template. The

parameters derived from this normalization process were subsequently applied to each EPI. The

normalized EPIs were spatially smoothed using an 8-mm full width at half maximum Gaussian kernel.

Voxel-based statistical analysis of the pre-processed EPIs was performed using a general linear

model (GLM). The blood oxygenation level-dependent (BOLD) response was related to the baseline

and anticipation periods, and the crash events were modeled as a boxcar function for the onset and

duration of each event (10 s for the baseline, 8 s for the anticipation period, and 0 s for the crash

event). Each resulting time series for each event was convolved with a canonical hemodynamic

response function, which was subsequently used as a regressor. Six head motion parameters, derived

from the realignment process, were also used as regressors to reduce motion-related artifacts. To

eliminate low-frequency drifts, a high-pass filter with a 128-s cut-off period was applied to the fMRI

time series. Serial autocorrelations between scans were corrected using a first-order autoregressive

model.

Regression coefficients for each event were computed for each participant using a fixed-effects

model. We created contrast images for the anticipation periods relative to the baseline in each of the

crash, attention, and safe conditions (anticipation_crash > baseline; anticipation_attention > baseline;

anticipation_safe > baseline) for the brain activity for each event. In addition, we created contrast

images to compare the brain activity during the anticipation period between the attention and safe

conditions (anticipation_attention > anticipation_safe; anticiaption_safe > anticipation_attention). We

also created contrast images for the crash events relative to the baseline. These contrast images were

subjected to group analysis using a random-effects model with a one-sample t-test. In the group

analysis, we set the statistical threshold at an uncorrected $p < 0.001$ at the voxel level and at a family-

wise error (FWE)-corrected $p < 0.05$ at the cluster level. We excluded the crash condition (i.e., the

condition in which the cue, "Attention," was followed by a car crash) from the main analyses (for the

common brain activity during the anticipation period, the difference in brain activity during the

anticipation period between the conditions, and the parametric modulation analyses), based on the

following reasons: (1) the crash condition contained the presentation of a broken windshield and a

collision sound, unlike the attention and safe conditions; (2) since subjective ratings were obtained

after each trial, it is possible that the additional aversive auditory-visual information affected the

subjective ratings in the crash condition.

We conducted parametric modulation analyses to examine subjective anxiety-related brain

activity. For parametric modulation of subjective anxiety, we used regressors for the anticipation

period in the trials under both attention and safe conditions, and z-score normalized scores of anxiety

in each corresponding trial as the parametric modulator.

For autonomic response-related brain activity, we conducted parametric modulation analyses

using β_{art} and the pupil area. In these analyses, trial-by-trial values of the ratio of mean β_{art} and pupil

area were used. These were calculated by averaging data over the 5-s period (from 3 to 8 s) following

the cue onset to those averaged over the baseline. We created parametric modulation regressors using

β_{art} and the pupil area for the anticipation period in the attention and safe conditions.

Autonomic response-related functional connectivity analysis

To examine the pupil- and β_{art}-related brain network, we examined FC between the anterior

insula and its adjacent regions (as seed regions) and the rest of the brain regions by conducting a

generalized form of context-dependent psychophysiological interactions (gPPI) using the CONN

toolbox (Whitfield-Gabrieli and Nieto-Castanon 2012). For gPPI analyses, we created GLMs,

including a regressor for the anticipation period in trials under both attention and safe conditions. The

processed GLM for each participant was imported to the gPPI model in the CONN toolbox. For each

participant, we created a separate gPPI model for pupil area and β_{art} parametric modulators. As first-

level covariates, we created gPPI regressors using parametric-modulation time-series based on trial-

by-trial values of the ratio of the mean β_{art} and pupil area averaged over the 5-s period from 3 to 8 s

following the cue onset to those averaged over the baseline. Based on our hypothesis, we defined seed

regions as the anterior insula (dorsal and ventral agranular insula) and the anatomical regions in the

cluster, including the anterior insula activated in the contrast condition (attention > safe). We used the

human Brainnetome Atlas (Fan et al. 2016) to define these anatomical regions of interest. In the

second-level analysis, seed-based connectivity maps of Fisher-transformed correlation coefficients

between BOLD time-series of each seed region and each individual voxel were entered into a one-

sample t-test to examine the brain regions showing significant pupil- and β_{art}-related FC with each

seed region. We applied a statistical threshold at an uncorrected $p < 0.001$ at the voxel level and a

false discovery rate (FDR)-corrected $p < 0.05$ at the cluster level.

Results

Behavioral data

Subjective ratings

Figure 2 shows the results of subjective ratings. Regarding subjective rating of anxiety, a

repeated-measures two-way ANOVA with factors of task conditions (crash, attention, and safe) and

session (sessions 1, 2, 3, and 4) revealed a significant main effect of task conditions (F [1.84, 60.83] =

316.596, partial $\eta^2 = 0.906$, $p_{corrected} < 0.001$; Chi-Muller correction for non-sphericity was applied).

Post-hoc analyses using the modified Shaffer method revealed that the participants rated subjective

anxiety highest in the crash condition, followed by the attention condition and the safe condition (all

ts [33] > 4.281, all ps < 0.001). In contrast, we observed no main effect of session (F [2.40, 79.08] =

0.423, partial $\eta^2 = 0.013$, $p_{corrected} = 0.693$) or interaction (F [5.35, 176.41] = 0.786, partial $\eta^2 = 0.023$,

$p_{corrected} = 0.569$). Considering the unpleasantness and arousal, repeated-measures two-way ANOVA

also revealed a significant main effect of condition (unpleasantness: F [2, 66] = 304.478, partial η^2 =

0.902, $p_{corrected}$ < 0.001; arousal: F [2, 66] = 156.673, partial η^2 = 0.826, $p_{corrected}$ < 0.001); however, no

main effect of session (unpleasantness: F [2.64, 86.96] = 1.004, partial η^2 = 0.03, $p_{corrected}$ = 0.388;

arousal: F [2.54, 83.78] = 0.696, partial η^2 = 0.021, $p_{corrected}$ = 0.534) or interaction (unpleasantness: F

[4.93, 162.75] = 0.898, partial η^2 = 0.026, $p_{corrected}$ = 0.483; arousal: F [4.99, 164.57] = 1.316, partial η^2

= 0.038, $p_{corrected}$ = 0.260) was observed. Post-hoc analyses again replicated the result for anxiety

(unpleasantness: all ts [33] > 10.207, p < 0.001; arousal: all ts [33] > 3.002, all ps < 0.01).

Considering pleasantness, a repeated-measures two-way ANOVA again revealed a significant main

effect of condition (F [1.94, 63.91] = 213.201, partial η^2 = 0.866, $p_{corrected}$ < 0.001). Post-hoc analyses

revealed that pleasantness was rated highest in the safe condition, followed by the attention condition

and the crash condition (all ts [33] > 9.249, all ps < 0.001). We observed no main effect of session (F

[3, 99] = 1.005, partial η^2 = 0.03, $p_{corrected}$ = 0.394), and a marginally significant interaction (F [4.89,

161.29] = 2.098, partial η^2 = 0.06, $p_{corrected}$ = 0.07).

The mean scores of STAI-trait, and STAI-state before (STAI-state pre) and after the

experiment (STAI-state post) were as follows: STAI-trait, 42.5 ± 7.50; STAI-state pre, 35.29 ± 7.50;

and STAI-state post, 34.82 ± 6.80. We conducted correlation analyses between all pairs of the mean

subjective ratings for each participant for each item and the STAI score (STAI-trait, STAI-state pre,

and STAI-state post). No significant correlation was observed between any of the pairs (all ps > 0.10).

Fig. 2. The results of subjective ratings (N = 34). SEM, standard error of the mean.

Reaction times

Figure 3 shows the mean RT for each condition. A one-way ANOVA on the mean RTs with a

factor of task condition revealed a significant main effect (F [1.73, 57.07] = 17.793, partial η^2 = 0.350,

$p_{corrected}$ < 0.001). Post-hoc analyses using the modified Shaffer method revealed that the mean RT for

the crash condition was faster than that for the attention and safe conditions (ts [33] < -4.060, ps <

0.001). However, we found no significant difference between the attention and safe conditions (t [33]

$= -1.070$, $p = 0.292$).

Fig. 3. Mean reaction time (N = 34). SEM, standard error of the mean.

Autonomic responses

Pupil dilation

Figure 4 shows the mean time series of all the participants' pupil responses during the task for

each condition. For data at each time point, we performed an FDR-corrected paired *t*-test. This

revealed significantly more pupil dilation in the crash/attention conditions than in the safe condition at

a threshold of $p_{FDR} < 0.05$, mainly in the period from approximately 1 s following the cue onset (i.e.,

18 s following the video clip onset) to the end of the trial. The gray areas in Figure 4 represent the

time points showing significant differences between the conditions. This indicates that significant

differences were robustly observed during the period from 3 to 8 s following cue onset. To examine

the statistical difference between conditions during this period, we performed a paired t-test between

the mean pupil areas in the crash/attention and safe conditions in the period from 3 to 8 s following

the cue onset. This again revealed significantly more pupil dilation in the crash/attention condition

than in the safe condition (t [33] = 4.4023, $p < 0.001$).

To examine the relationship between the pupil dilation and the subjective ratings of anxiety,

we conducted a correlation analysis using the difference between the mean rating scores of anxiety in

the attention and safe conditions for each participant, and the difference between the mean pupil areas

in the crash/attention and safe conditions for the period from 3 to 8 s following the cue onset. The

analysis revealed that the pupil dilation and the anxiety ratings showed a marginally significant

positive correlation ($r = 0.313$, t [31] = 1.834, $p = 0.076$, Supplementary Figure S1). For the other

items, we found a marginal positive correlation for arousal ($r = 0.318$, t [31] = 1.865, $p = 0.072$), and

no significant correlation for pleasantness or unpleasantness (pleasantness: $r = 0.270$, t [31] = 1.564, p

= 0.128; unpleasantness: $r = 0.149$, t [31] = 0.836, $p = 0.409$).

Fig. 4. The mean time-series of pupil areas for attention/crash and safe conditions. The gray areas

represent the time points showing significant differences between conditions (see text).

Peripheral arterial stiffness

Figure 5 shows the mean time series of all the participants' β_{art} during the task for each

condition. To examine the significance of the increase in β_{art} following cue onset, we compared the

ratio of the mean value of β_{art} during the 5-s period from 3 to 8 s following the cue onset to the mean

value over the baseline between the attention and safe conditions using a paired t-test. This revealed a

tendency for β_{art} in the attention condition to be greater than in the safe condition (t [32] = 1.80, p =

0.08).

Fig. 5. The mean time-series of peripheral arterial stiffness (β_{art}) for attention/crash and safe

conditions.

fMRI data

Brain activity during the anticipation period

To examine the common brain activity during the anticipation period relative to the baseline

in both the attention and safe conditions, we conducted a conjunction analysis. This revealed

significant activations in the bilateral front-parietal cortices, dorsal anterior cingulate cortex, and

bilateral anterior insulae (Figure 6, Table 1).

Fig. 6. Brain regions commonly activated in the attention and safe conditions revealed by a

conjunction analysis. SPL, superior parietal lobule; MFG, middle frontal gyrus; SFG, superior frontal

gyrus; opIFG, inferior frontal gyrus, opercular part; IPL, inferior parietal lobule; SMA, supplementary

motor area; dPM, dorsal premotor cortex; dACC, dorsal anterior cingulate cortex; aIns, anterior insula

Table 1. Common brain regions activated in both the attention and safe conditions during the

anticipation period based on the result of a conjunction analysis.

Anatomical region	Cluster size	MNI coordinates (mm)			*T*-value
		x	y	Z	
Left lingual gyrus	4,299	-12	-88	-9	18.67
Right calcarine gyrus		12	-88	-3	17.87
Right lingual gyrus		21	-76	-12	15.84
Right superior frontal gyrus	1201	24	-1	61	9.34

Left superior frontal gyrus		-24	-4	68	8.41
Left supplementary motor area		-6	11	52	7.39
Right inferior frontal gyrus, triangular part	116	60	23	16	5.35
Right inferior frontal gyrus, triangular part/anterior insula		39	23	13	4.45
Left inferior frontal gyrus, opercular part/left anterior insula	100	-45	14	10	5.15
Left inferior frontal gyrus, opercular part/left anterior insula		-36	17	10	4.70

Uncorrected $p < 0.001$ at peak level, family-wise error-corrected $p < 0.05$ at the cluster level. MNI,

Montreal Neurological Institute

To examine the brain regions related to each condition, we compared brain activation between

the attention and safe conditions during the anticipation period. In the attention condition, the right

anterior insula and the subcortical regions, including the PAG, thalamus, and BNST were more active

than in the safe condition (Figure 7, Table 2). We confirmed that the activated cluster had overlaps

with the atlas of the PAG (Keuken et al. 2017) and BNST (Torrisi et al. 2015, Figure 7B). In contrast,

the areas from the parietal operculum, including the secondary somatosensory area (SII) to the

posterior insula, and the precuneus were less active in the attention condition than in the safe

condition. Although these areas were deactivated under all conditions, deactivation was more evident

in the attention condition than in the safe condition (Figure 8, Table 3).

Parametric modulation using anxiety rating scores, including attention and safe conditions,

revealed that activations in the thalamus and PAG were positively correlated with the anxiety ratings.

In contrast, we found no brain regions showing a significant negative correlation with the anxiety

ratings at a voxel-level threshold of uncorrected $p < 0.001$ or a cluster-level FWE-corrected $p < 0.05$;

however, at a liberal threshold, the SII and posterior insula showed a negative correlation at a voxel-

level uncorrected $p < 0.005$ (cluster-corrected FWE-corrected $p = 0.066$; Figure 9, Table 4).

On the other hand, parametric modulation using physiological signals did not reveal any

significant activation for either pupil areas or β_{art}.

Fig. 7. (A) Brain regions showing more activation in the attention condition than in the safe condition.

(B) The overlap between the activated cluster (red) and the anatomical atlas of the BNST (Torrisi et al.

2015) and PAG (Keuken et al. 2017) (green). (C) Contrast estimates of the peak voxel in the activated

cluster in the right anterior insula for each condition. (D) The time course for BOLD signal change of

the peak voxel in the right anterior insula for each condition during the anticipation period. Each line

represents the mean event-related BOLD response over all participants for each condition with the

standard error of the mean as represented by the area with the corresponding color. The plot was

created by using the rfxplot Toolbox (Gläscher 2009).

aIns, anterior insula; PAG, periaqueductal gray; BNST, bed nucleus of the stria terminalis; BOLD,

1   blood oxygenation level-dependent.

3   **Table 2.** Brain regions significantly more activated in the attention condition than in the safe

4   condition

Anatomical region	Cluster size	MNI coordinates (mm)			*T*-value
		x	y	Z	
Left red nucleus	516	-6	-25	-9	7.53
Right caudate head		12	5	4	6.31
Left medial dorsal thalamus		-9	-19	13	5.97
Right inferior frontal gyrus, triangular part/anterior insula	105	36	26	7	5.96
Right inferior frontal gyrus, triangular part		48	29	16	3.76

5   Uncorrected $p < 0.001$ at peak level, family-wise error-corrected $p < 0.05$ at the cluster level. MNI,

6   Montreal Neurological Institute

Fig. 8. (A) Brain regions showing more activation in the safe condition than in the attention condition.

(B) Contrast estimates of the peak voxel in the activated cluster in the right SII/posterior insula for

each condition. (C) The time course for blood oxygenation level-dependent (BOLD) signal change of

the peak voxel in the right SII/posterior insula for each condition during the anticipation period. Each

line represents the mean event-related BOLD response over all participants for each condition with

the standard error of the mean as represented by the area with the corresponding color. The plot was

created by using the rfxplot Toolbox (Gläscher 2009).

1 vPM, ventral premotor cortex; IPL, inferior parietal lobule; SMA, supplementary motor area; dPM,

2 dorsal premotor cortex; SII, secondary somatosensory area; pIns, posterior insula.

4 **Table 3.** Brain regions significantly more activated in the safe condition than in the attention

5 condition

Anatomical region	Cluster size	MNI coordinates (mm)			*T*-value
		x	y	z	
Right Rolandic operculum	303	48	-13	23	5.70
Right Rolandic operculum		60	2	13	5.26
Right posterior insula		39	-10	16	4.99
Left posterior cingulate cortex	110	-3	-52	32	5.10
Left postcentral gyrus	307	-48	-19	26	5.06
Left posterior insula		-39	-13	13	4.95
Left postcentral gyrus		-57	-7	20	4.54
Left inferior parietal cortex	500	-54	-22	48	4.92
Left postcentral gyrus		-42	-34	61	4.74
Right paracentral lobule		6	-31	64	4.74
Left angular gyrus	99	-57	-67	29	4.85

Left angular gyrus		-45	-55	29	4.28
Left angular gyrus	303	-51	-67	42	3.90

Uncorrected $p < 0.001$ at peak level, family-wise error-corrected $p < 0.05$ at the cluster level. MNI,

Montreal Neurological Institute

Fig. 9. Brain activation correlated with the anxiety rating scores. (A) Brain regions positively

correlated with subjective anxiety (B) The overlap between the activated cluster (red) and the

anatomical atlas of the PAG (Keuken et al. 2017) (green). (C) Brain regions negatively correlated with

subjective anxiety at a liberal threshold (voxel-level uncorrected $p < 0.005$ and cluster-level FWE-

corrected $p = 0.066$). PAG, periaqueductal gray; BNST, bed nucleus of the stria terminalis; SII,

secondary somatosensory area; pIns, posterior insula.

Table 4.

(A) Brain regions showing activation positively correlated with subjective ratings of anxiety

Anatomical region	Cluster size	MNI coordinates (mm)			T-value
		x	y	z	
Left red nucleus	199	-6	-25	-6	6.43
Left medial dorsal thalamus		-3	-19	4	4.49
Left medial dorsal thalamus		-9	-19	10	4.46

1   Uncorrected $p < 0.001$ at peak level, family-wise error-corrected $p < 0.05$ at the cluster level. MNI,

2   Montreal Neurological Institute

4   (B) Brain regions showing activation negatively correlated with subjective ratings of anxiety

Anatomical region	Cluster size	MNI coordinates (mm)			T-value
		x	y	z	
Right Rolandic operculum	194	48	-4	20	4.74
Right Rolandic operculum		51	-13	23	3.82
Right posterior insula		42	-4	7	3.56

5   Uncorrected $p < 0.005$ at peak level. MNI, Montreal Neurological Institute

7   *Brain activity related to the crash*

The contrast in the crash event versus baseline showed extensive brain activation; therefore,

we reported the brain regions with a voxel-level FWE-corrected $p < 0.05$. The activated regions can

be classified into approximately three large clusters (Figure 10A, Table 5): (1) a posterior midline

cluster including the precuneus, cerebellum, and thalamus; (2) a frontal midline cluster including the

supplementary motor area and ACC, and (3) a fronto-temporoparietal cluster including the anterior

insula, inferior frontal cortex, superior temporal cortices, parietal operculum, and posterior insula.

This fronto-temporoparietal cluster included the amygdala (Figure 10B).

Fig. 10. (A) Brain regions significantly activated in the contrast of crash > baseline. (B) The amygdala

activation observed in the fronto-temporoparietal cluster.

Table 5. Brain regions significantly activated in the contrast crash > baseline

Anatomical region	Cluster size	MNI coordinates (mm)			T-value
		x	y	z	
Left thalamus (medial pulvinar)	11,381	-9	-25	-3	17.48
Right thalamus (medial geniculate nucleus)		12	-25	-6	17.02
Right lingual gyrus		15	-49	-3	15.97
Right pregenual ACC	1,561	9	44	20	11.21
Left superior medial frontal gyrus		3	17	42	10.67
Left supragenual ACC		-6	26	29	10.53
Cerebellum vermis	110	3	-58	-41	8.42
Left inferior parietal lobule	43	-33	-52	45	8.04
Right paracentral lobule	37	12	-37	55	6.39
Right precuneus		12	-46	55	6.20
Right precuneus		6	-52	55	5.98
Left cerebellum (VIII)	35	-12	-70	-48	6.27
Cerebellum vermis		6	-73	-41	6.16
Left cerebellum crus		-9	-76	-35	5.92
Left precuneus	5	-12	-49	45	6.19
Right inferior parietal lobule	8	30	-46	52	6.14
Right inferior parietal lobule		36	-49	45	6.05

Left middle frontal gyrus	23	-39	38	26	6.11
Left middle frontal gyrus		-36	29	36	5.92
Left middle frontal gyrus		-36	41	36	5.68
Right superior parietal lobule	1	54	-43	58	5.74
Left cerebellum (VIII)	1	-30	-64	-54	5.68
Right inferior parietal lobule	1	45	-52	42	5.66

Family-wise error-corrected $p < 0.05$ at the peak level. MNI, Montreal Neurological Institute; ACC,

anterior cingulate cortex

Autonomic response-related functional connectivity

Based on the brain activity in the contrast condition (attention > safe), we conducted gPPI

analyses using the right anterior insula (dorsal and ventral agranular insula) and the brain regions

including the anterior insula cluster, such as the right middle frontal gyrus (MFG) ventral area

(Brodmann area [BA] 9/46), right inferior frontal sulcus, right caudal inferior frontal gyrus (IFG)

(BA45), right rostral IFG (BA45), and right opercular IFG (BA44) as seed regions.

Regarding the pupil area, although we observed no brain region showing significant FC for the

right anterior insula as a seed region, we found brain regions showing significant pupil-related FCs

with the brain areas adjacent to the anterior insula: the right MFG and the right rostral IFG (BA45).

Moreover, the left cerebellum/brainstem showed a significant pupil-related FC with the right MFG,

and the left visual cortex showed a significant pupil-related FC with the right rostral IFG (Figure

11A–B, Table 6).

Regarding the β_{art}, there were significant FCs between the right dorsal agranular insula and the

middle/posterior cingulate cortex (MCC/PCC), and between the right ventral agranular insula and the

right visual cortex (Figure 11C–D, Table 6). In contrast, we found no brain region showing significant

FC with the regions adjacent to the anterior insula.

Fig. 11. (A–B) Brain regions showing pupil-related functional connectivity. (A) Brain regions

showing pupil-related functional connectivity with the right middle frontal gyrus as seed region. (B)

Brain regions showing pupil-related functional connectivity with the right rostral inferior frontal gyrus

as seed region. (C–D) Brain regions showing peripheral arterial stiffness (β_{art})-related functional

connectivity. (C) Brain regions showing β_{art}-related functional connectivity with the right dorsal

agranular insula as seed region. (D) Brain regions showing β_{art}-related functional connectivity with the

1 right ventral agranular insula as seed region. Vis, visual cortex; MCC, middle cingulate cortex; PCC,

2 posterior cingulate cortex.

4 **Table 6.** Brain regions showing pupil-related functional connectivity

5 (A) Brain regions showing pupil-related functional connectivity with the right MFG as seed region

Anatomical region	Cluster size	MNI coordinates (mm)			*T*-value
		x	y	z	
Left cerebellum	176	-20	-44	-32	5.26
Left cerebellum		-26	-34	-32	5.07
Brainstem		-14	-34	-32	3.50

6 Uncorrected $p < 0.001$ at peak level, false discovery rate-corrected $p < 0.05$ at the cluster level. MFG,

7 middle frontal gyrus; MNI, Montreal Neurological Institute

9 (B) Brain regions showing pupil-related functional connectivity with the right rostral IFG as seed

10 region

Anatomical region	Cluster size	MNI coordinates (mm)			*T*-value

		x	y	z	
Left cuneus	307	-14	-82	16	4.74
Left superior occipital gyrus		-18	-98	16	4.44
Left superior occipital gyrus		-14	-92	12	3.99

1 Uncorrected $p < 0.001$ at peak level, false discovery rate-corrected $p < 0.05$ at the cluster level. IFG,

2 inferior frontal gyrus; MNI, Montreal Neurological Institute

4 (C) Brain regions showing β_{art}-related functional connectivity with the right dorsal agranular insula as

5 seed region

Anatomical region	Cluster size	MNI coordinates (mm)			T-value
		x	y	z	
White matter	153	18	-22	36	4.94
Middle cingulate gyrus		14	-16	40	4.08
Posterior cingulate gyrus		14	-26	40	3.63

6 Uncorrected $p < 0.001$ at peak level, false discovery rate-corrected $p < 0.05$ at the cluster level. MNI,

1   Montreal Neurological Institute

3   (D) Brain regions showing β_{art}-related functional connectivity with the right ventral agranular insula

4   as seed region

Anatomical region	Cluster size	MNI coordinates (mm)			T-value
		x	y	z	
Right calcarine sulcus	155	10	-70	6	5.24
Right calcarine sulcus		10	-80	10	4.17
Right lingual gyrus		2	-70	4	3.97

Uncorrected $p < 0.001$ at peak level, false discovery rate-corrected $p < 0.05$ at the cluster level. MNI,

Montreal Neurological Institute

Discussion

In this study, we examined the neural basis of anxiety and anxiety-related autonomic

responses during a daily driving situation. We hypothesized that the brain network related to the

elicitation of anxiety and corresponding autonomic responses would be observed in the anterior

insula and subcortical regions. As predicted, during the period in which the participants

anticipated a car crash, we observed cortico-subcortical activations in the anterior insula, PAG,

thalamus, and BNST. The amygdala activity observed in response to a crash (but not detected in

anticipation of a crash) could extend our understanding of the role of the amygdala in the

elicitation of anxiety. Moreover, we observed deactivation in the posterior insula and SII, which

could reflect an allostatic response against the upcoming crash. This could offer new insights for

the safe-signal-related brain network that has been unclear from previous fear-conditioning

studies. We also observed significant pupil dilation, although the increase in β_{art} related to the

anticipation of a crash was not statistically significant. The FC analyses using these autonomic

responses revealed that β_{art}-related FC was found between the right dorsal anterior insula and

cingulate cortex, as well as between the right ventral anterior insula and visual cortex, and that

pupil-related FC was found between the right MFG and the cerebellum/brainstem, as well as

between the right rostral IFG and the visual cortex. Our results could extend the knowledge

regarding the neural basis of anxiety and anxiety-related physiological responses.

Brain regions involved in anxiety elicitation

     In the attention condition, the brain regions including the anterior insula, PAG, thalamus,

and BNST were more active than those in the safe condition. The anterior insula is reportedly

involved in the integration and prediction of interoceptive information (Craig 2009) and in the

allostatic process based on prediction (Barrett et al. 2017). The peak voxel of the anterior insula

cluster is located at the dorsal part of the anterior insula and extends to the operculum and

triangular parts of the IFG. These areas are consistent with those related to sustained fear

reported by Somerville et al. (2013). The dorsal anterior insula has been suggested to be

involved in cognitive and attentional functions rather than in emotional function (e.g., Chang et

al. 2013). This could suggest that the anterior insular activity during the attention condition

could reflect only the attentional process; however, based on the results of this study, such a

suggestion can be ruled out. The results of anxiety ratings indicated that participants felt more

anxious during the attention condition than during the safe condition. In contrast, RTs showed

no significant difference between conditions (see Figure 3), suggesting that attentional resources

allocated to the task were similar among the conditions. Furthermore, we found no significant

difference in activation in the dorsal attentional network, such as the dorsolateral prefrontal

cortex, frontal eye field, and posterior parietal cortex, between the attention and safe conditions.

Thus, our results suggest that the brain regions including the anterior insula were related to

anxiety, that is, negative emotion caused by prediction of an aversive event rather than just

attention.

The PAG receives projections from the prefrontal cortex, insula, and amygdala (Mantyh

1982). The PAG is suggested to play a role in homeostatic defense by integrating afferent

information from the periphery and information from higher centers (Linnman et al. 2012). For

instance, the PAG is known to be involved in pain modulation. It has been reported that when

pre-stimulus FC between the anterior insula and PAG is stronger, participants perceive a pain

stimulus as less painful (Ploner et al. 2010).

The thalamus is also reportedly involved in the modulation of ascending nociceptive

information (Tang et al. 2009). Moreover, it is known that the thalamus relays nociceptive

information to the insula, anterior cingulate cortex, and somatosensory cortices (Kummer et al.

2020). In the current study, we observed thalamic activation mainly in the dorsomedial region.

Considering that the dorsomedial thalamus is a part of the medial spinothalamocortical pathway

involved in nociceptive-specific responses (e.g., Gingold et al. 1991), the thalamic activity

observed in this study presumably reflects the anticipatory modulation of responses to

unpleasant stimuli accompanied by collision.

The BNST is known to be involved in emotion, threat, and autonomic processing (Davis

et al. 2010; Crestani et al. 2013). While animal studies have reported the structural connection

between the insula and BNST (e.g., Centanni et al. 2019), the BNST-insula connection in human

brains has also been revealed in recent MRI studies (e.g., Avery et al. 2014; Flook et al. 2020).

This connection is suggested to be involved in the translation of emotional states into behavioral

responses, including "fight or flight" (Flook et al. 2020). Based on these previous studies,

activation in the BNST and anterior insula detected in the attention condition might be involved

in the elicitation of autonomic and behavioral responses accompanied with anticipation of a

crash. Thus, our results suggest that a cortico-subcortical network comprising the anterior insula,

PAG, BNST, and thalamus is involved in the prediction of negative events and elicitation of

anxiety-related responses based on interoceptive information in the attention condition.

In contrast, the SII/posterior insula and precuneus were more activated in the safe

condition than in the attention condition. In the attention condition, where a crash was predicted,

the activation in these regions decreased as the timing of the crash approached (see Figure 8C).

Therefore, this deactivation could reflect the allostatic process preparing for the predicted

response to a possible upcoming aversive event. The SII/posterior insula has been reported to be

involved in processing aversive stimuli (e.g., Apkarian et al. 2005). In particular, the SII has

been suggested to represent somatosensory prediction error (Blakemore et al. 1998). In the

study by Blakemore et al., participants underwent two experimental conditions: the

experimenter tickled the participants' palms, and the participants tickled themselves. In the

condition of self-tickling, the SII was more deactivated than in the condition in which the

experimenter tickled the participants, suggesting that this deactivation reflects the process of

canceling out the sensory consequences, that is, a tickly feeling, generated by prediction. In our

study, crash-related activation overlapped in the SII/posterior insula cluster, as observed in the

contrast of safe > attention. This suggests that SII/posterior insula activation could reflect a part

of the aversive sensation elicited by the crash. Considering these observations, the deactivation

in the SII/posterior insula observed in the attention condition could be interpreted as an

allostatic attenuation triggered by the prediction of sensory input followed by an upcoming

aversive event.

The precuneus is part of the DMN. It has been proposed that the DMN is switched to the

central executive network according to the task demand by the salience network (Menon and

Uddin 2010). Therefore, deactivation in the precuneus could reflect inhibition of the DMN to

prepare and pay attention to the upcoming event in the attention condition.

Brain activity related to subjective anxiety

The results of the parametric modulation analyses showed that subjective anxiety was

significantly correlated with activation in the thalamus and PAG, and not with the anterior insula.

This could be explained by the relationship between anterior insula-PAG pre-stimulus FC and

subjective pain sensation (Ploner et al. 2010). Considering that the anterior insula-PAG

connection during the anticipation period determines subjective anxiety, when anticipatory

modulation of PAG by the anterior insula was insufficient, the participants could have felt more

anxious. In contrast, regardless of subjective anxiety, the anterior insula was active when an

upcoming crash could be predicted, which could contribute to the result that the anterior insula

was less related to subjective anxiety.

While many previous studies have suggested that the amygdala is involved in anxious

emotion, we did not find significant amygdala activity during the anticipation period in the

contrast of attention > safe or in the parametric modulation of subjective anxiety. However, the

amygdala was significantly active in response to a crash relative to the baseline (Figure 10B)[1].

This is consistent with the study of Somerville et al. (2013), who reported that the amygdala

---

[1] We must note that the crash condition was quite different from other conditions, as indicated in

the behavioral data. For example, the RTs to the word "STOP" in the crash condition were faster

than in other conditions. This could be attributed to the auditory-visual stimuli accompanied

with a crash that captured the participants' attention, thereby enhancing their responses to the

word "STOP." Furthermore, the subjective ratings for the crash condition were significantly

different from those for other conditions, indicating that the subjective ratings performed after

each trial were affected by the crash, as predicted.

responds to transient fear rather than sustained anxious emotion (anxiety). Thus, in the current

study, the absence of amygdala activity during the anticipation phase is reasonable.

Brain regions involved in anxiety-related autonomic responses

The autonomic response-related FC analyses revealed that the right anterior insula and

its adjacent regions showed significant FC with the visual cortex, cerebellum, brainstem, and

MCC/PCC.

Related to β_{art}, the right ventral and dorsal anterior insula showed FC with the visual

cortex and MCC/PCC, respectively. The anterior insula controls the autonomic nervous system

by regulating the subcortical brain areas, such as the PAG (Thayer and Lane 2009; Critchley and

Harrison 2013). Previous research has demonstrated that autonomic control of cardiac activity is

lateralized and mediated by the right insular cortex (Colivicchi et al. 2004; Craig 2009). For

instance, evidence from stroke studies has suggested that the right insula plays a major role in

cardiac autonomic control (e.g., Colivicchi et al. 2004). Craig (2009) proposed a hypothesis that

the right and left insular cortices are involved in the control of the sympathetic and

parasympathetic nervous systems, respectively. Based on these findings, the observation of β_{art}-

related FC for the right anterior insula as seed region is reasonable. The MCC/PCC cluster

showing FC with the right dorsal anterior insula was located at the posterior part of the MCC

and PCC. Particularly, the PCC is suggested to be involved in controlling the balance between

an internal and external attentional focus in collaboration with the right anterior insula (Leech

and Sharp 2014). The ventral anterior insula, which is involved in affective processing, showed

FC with the visual cortex. This might reflect the allostatic processing of upcoming visual

information based on anxiety elicitation. These observations suggest that the allostatic affective

processing was associated with the β_{art} change. However, Tsuji et al. (2021) demonstrated that

the β_{art} during pain stimulation was correlated with brain activity in the salience network,

including the dACC but not the anterior insula. This inconsistency could suggest that the

anterior insula is related to the β_{art} attributed to anticipation, whereas the dACC is related to that

caused by stimulation.

Regarding the pupil-related FC, the right ventral MFG and the rostral IFG showed FC

with the cerebellum/brainstem and the visual cortex, respectively. The brain regions including

the anterior insula/IFG and MFG have also been reported to be related to pupil change during

reward anticipation (Schneider et al. 2018). The ventral MFG and rostral IFG, which are

adjacent to the inferior frontal sulcus, are a part of the ventral attention network (Corbetta et al.

2008). The ventral attention network is known to be a part of the salience network and has been

suggested to be a "circuit breaker" that reorients focused attention by detecting salient stimuli

such as oddballs. The ventral MFG showed an FC with the cluster including the pons located

near (however, not including) the locus coeruleus (LC), which is involved in pupil control

(Schneider et al. 2016). Furthermore, FC between the IFG and visual cortex was observed. This

FC could reflect allostatic attentional control to visual information. Considering that no pupil-

related FC was observed with the anterior insula as seed, the pupil-related FC in the current

study could reflect a relatively cognitive aspect of allostatic processing, such as attentional

control to visual information accompanied by an upcoming collision.

Thus, our results suggest that the anterior insula and its adjacent regions, in collaboration

with other regions, could differentially realize an allostatic attention control and adaptive

physiological response to environmental information.

However, we did not observe FC with the brain regions involved in general direct

control of the pupil and artery, such as the LC for pupil dilation and the hypothalamus for the

artery. One possible interpretation of this result is that the regions observed in this study could

be involved in task-evoked modulation of the pupil and artery as part of the allostatic process.

During the resting state and the continuous attentive task, for instance, the pupil diameter is

known to be correlated with BOLD signals in the LC (Murphy et al. 2014). However, certain

studies have reported that LC activation is not related to task-evoked pupil dilation (Schneider

et al. 2018). Nevertheless, the relationship between the task-evoked pupil- and artery-related

regions and those involved in the general control of the pupil and artery should be further

examined in future research.

Regarding pupil dilation, we found a robust difference between the attention/crash and

safe conditions. This result suggests the possibility that the measurement of pupil dilation is

useful in detecting anxiety. However, we must note that the correlation between the pupil

dilation and subjective ratings of anxiety were not significant and that we also observed a weak

(but not significant) correlation between pupil dilation and subjective arousal. Therefore, future

studies controlling the arousal level should assess whether pupil dilation merely reflects the

arousal.

In contrast, the increase of β_{art} in the attention/crash conditions relative to the safe

condition was not significant. This result could have been caused by the difference in the

temporal properties of pupil dilation and β_{art}. Arousal-related pupil dilation reportedly appears at

nearly 200 ms (Sirois et al. 2014). In the current study, owing to the fast latency of pupil dilation,

the fluctuation in the participants' subjective emotional state could be reflected more accurately

than β_{art}. In contrast to pupil dilation, β_{art} is a relatively newly proposed index. Information on

the temporal properties and mechanism of β_{art} should be investigated in future studies.

Limitations and future directions

In the current study, we demonstrated that the network including the anterior insula plays a pivotal role for eliciting anxiety and for accompanying autonomic responses in a daily situation. However, our results do not elucidate clinical conditions, such as anxiety disorder. For instance, we found no significant correlation between the trait and state anxiety and the subjective ratings. Furthermore, there was no significant brain activation related to the individual difference in the trait anxiety nor the state anxiety before and after the experiment, as assessed with the STAI. This could be attributed to the fact that we only recruited healthy participants with a relatively narrow range of scores. To apply our findings to the understanding of anxiety disorder, one should compare brain activity and physiological responses between patients and normal controls. One of the aims of future studies should be to assess whether anxiety-prone individuals exhibit an increase in the pupil area and β_{art}, which will deepen our understanding of the mechanism of anxiety. In addition, we should note an imbalance in the male-female ratio in this study. Regarding gender differences in emotion, there has been a debate in the field of emotion studies. A review from Kret and De Gelder (2012) pointed out that men tend to show strong effects of threatening stimuli, such as enhanced physiological arousal and brain activity. The gender difference of the anxiety-related brain and physiological responses should be examined in future studies.

There is also a limitation to the localization of the small subcortical regions, such as the BNST and PAG. Although we confirmed that the activated cluster included the BNST and PAG using anatomical atlases, more precise localization is necessary for a better understanding of the roles of the nuclei in anxiety-related responses in future. Such may be achieved with specific scan parameters to target the small nuclei or MRI with a higher magnetic field (e.g., 7-T MRI).

Our results also showed that the increase in pupil area and peripheral arterial stiffness were commonly related to the network including the right anterior insula and its adjacent regions in a daily situation of driving a vehicle. This suggests that these are potential indices to detect the driver's anxiety during driving, underpinned by neuroscientific evidence. If our results are applied to daily driving situations, however, there are some limitations. First, the task was a simple fear-conditioning task, which did not contain actual control of a handle, accelerator, or brake, as in an actual driving situation. Moreover, we did not perform a comparison to a "non-driving" condition. Second, the experimental environment in the MRI scanner differs much from an actual driving situation. For instance, the participants did not hold a steering wheel or place their foot on a brake pedal. Furthermore, they were lying in a supine position in the MRI scanner. For industrial application, an experiment should be conducted with a realistic steering device (e.g., Okamoto et al. 2020) that can be used in an MRI scanner, enabling a more

realistic driving situation, and a non-driving condition should be included for comparison. It is

also necessary to perform brain imaging and physiological recordings using wearable devices

(e.g., Protzak and Gramann 2018) while driving an actual vehicle.

Conclusions

Our results suggest that cortico-subcortical regions, including the right anterior insula,

PAG, thalamus, and BNST, play a core role in anticipatory responses to upcoming threats in

daily driving situations. We also observed that multiple autonomic responses, such as pupil

dilation and β_{art}, were evident when an upcoming threat was predicted, and the right anterior

insula and its adjacent regions play a key role in eliciting autonomic responses. These results

suggest that pupil dilation and β_{art} reflect anxiety-related salience network activity, and that they

are potential candidate indices for detecting task-evoked anxious emotion based on

neuroscientific evidence.

Conflicts of Interest

DS, NM, HY, and MT are employed by Mazda Motor Corporation. TN was also

employed by Mazda Motor Corporation during the period when the current study was being

conducted. The other authors declare no conflict of interest.

Funding

This work was supported by JST COI (Grant No. JPMJCE1311).

Acknowledgments

We are grateful to Dr. Norihiro Sadato, Dr. Alan Fermin, Dr. Kentaro Ono, Dr. Ayumu

Matani, Dr. Toshio Tsuji, and Dr. Maro G. Machizawa for helpful discussions concerning this

study, and Ms. Noriko Miura and Ms. Tamami Tomita for assistance with the implementation of

the experiment. We would like to thank Editage (www.editage.com) for English language

editing.

References

Aitken RCB. 1969. Measurement of feelings using visual analogue scales. Proc R Soc Med.

62(10):989–993.

Apkarian AV, Bushnell MC, Treede RD, Zubieta JK. 2005. Human brain mechanisms of pain

perception and regulation in health and disease. Eur J Pain. 9(4):463–484.

Avery SN, Clauss JA, Winder DG, Woodward N, Heckers S, Blackford JU. 2014. BNST

neurocircuitry in humans. NeuroImage. 91:311–323.

Barrett LF. 2017. The theory of constructed emotion: an active inference account of

interoception and categorization. Soc Cogn Affect Neurosci. 12(1):1–23.

Barrett LF, Quigley KS, Hamilton P. 2016. An active inference theory of allostasis and

interoception in depression. Phil Trans R Soc B. 371(1708):20160011.

Barrett LF, Simmons WK. 2015. Interoceptive predictions in the brain. Nat Rev Neurosci.

16(7): 419–429.

Baur V, Hänggi J, Langer N, Jäncke L. 2013. Resting-state functional and structural

connectivity within an insula-amygdala route specifically index state and trait anxiety.

Biol Psychiatry. 73(1):85–92.

Bitsios, P, Szabadi E, Bradshaw CM. 2004. The fear-inhibited light reflex: Importance of the

anticipation of an aversive event. Int J Psychophysiol. 52(1):87–95.

Blakemore SJ, Wolpert DM, Frith CD. 1998. Central cancellation of self-produced tickle

sensation. Nat Neurosci. 1(7):635–640.

Brookhuis KA, de Vries G, De Waard D. 1991. The effects of mobile telephoning on driving

performance. Accid Anal Prev. 23(4):309–316.

Centanni SW, Morris BD, Luchsinger JR, Bedse G, Fetterly TL, Patel S, Winder DG. 2019.

Endocannabinoid control of the insular-bed nucleus of the stria terminalis circuit

regulates negative affective behavior associated with alcohol abstinence.

Neuropsychopharmacology. 44(3):526–537.

Chang LJ, Yarkoni T, Khaw MW, Sanfey G. 2013. Decoding the role of the insula in human

cognition: Functional parcellation and large-scale reverse inference. Cereb Cortex.

23(3):739–749.

Chen WG, Schloesser D, Arensdorf AM, Simmons JM, Cui C, Valentino R, Gnadt JW, Nielsen

L, St. Hillaire-Clarke C, Spruance VK, et al. 2021. The emerging science of

interoception: sensing, integrating, interpreting, and regulating signals within the self.

Trends Neurosci. 44(1):3–16.

Choe KW, Blake R, Lee SH. 2016. Pupil size dynamics during fixation impact the accuracy and

precision of video-based gaze estimation. Vis Res. 118:48–59.

Clark A. 2013. Whatever next? Predictive brains, situated agents, and the future of cognitive

science. Behav Brain Sci. 36(3):181–204.

Colivicchi F, Bassi A, Santini M, Caltagirone C. 2004. Cardiac autonomic derangement and

arrhythmias in right-sided stroke with insular involvement. Stroke. 35(9):2094–2098.

Corbetta M, Patel G, Shulman GL. 2008. The reorienting system of the human brain: from

environment to theory of mind. Neuron. 58(3):306–324.

Craig AD. 2009. How do you feel—now? The anterior insula and human awareness. Nat Rev

Neurosci. 10(1):59–70.

Crestani C, Alves F, Gomes F, Resstel L, Correa F, Herman J. 2013. Mechanisms in the bed

nucleus of the stria terminalis involved in control of autonomic and neuroendocrine

functions: a review. Curr Neuropharmacol. 11(2):141–159.

Critchley HD, Harrison NA. 2013. Visceral influences on brain and behavior. Neuron.

77(4):624–638.

Davis M, Walker DL, Miles L, Grillon C. 2010. Phasic vs sustained fear in rats and humans:

Role of the extended amygdala in fear vs anxiety. Neuropsychopharmacology.

35(1):105–135.

Deane GE. 1961. Human heart rate responses during experimentally induced anxiety. J Exp

Psychol. 61:489–493.

Epstein S, Roupenian A. 1970. Heart rate and skin conductance during experimentally induced

anxiety: The effect of uncertainty about receiving a noxious stimulus. J Pers Soc

Psychol. 16(1):20–28.

Etkin A, Wager TD. 2007. Functional neuroimaging of anxiety: A meta-analysis of emotional

processing in PTSD, social anxiety disorder, and specific phobia. Am J Psychiatry.

164(10):1476–1488.

Fan L, Li H, Zhuo J, Zhang Y, Wang J, Chen L, Yang Z, Chu C, Xie S, Laird AR, Fox PT,

Eickhoff SB, Yu C, Jiang T. 2016. The Human Brainnetome Atlas: A New Brain Atlas

Based on Connectional Architecture. Cereb Cortex. 26(8): 3508–3526.

Flook EA, Feola B, Avery SN, Winder DG, Woodward ND, Heckers S, Blackford JU. 2020.

BNST-insula structural connectivity in humans. NeuroImage. 210:116555.

Friston K. 2005. A theory of cortical responses. Phil Trans R Soc B. 360(1456):815–836.

Fullana MA, Harrison BJ, Soriano-Mas C, Vervliet B, Cardoner N, Àvila-Parcet A, Radua, J.

2016. Neural signatures of human fear conditioning: An updated and extended meta-

analysis of fMRI studies. Mol Psychiatry. 21(4):500–508.

Gingold SI, Greenspan JD, Apkarian AV. 1991. Anatomic evidence of nociceptive inputs to

primary somatosensory cortex: relationship between spinothalamic terminals and

thalamocortical cells in squirrel monkeys. J Comp Neurol. 308(3):467–490.

Gläscher J. 2009. Visualization of group inference data in functional neuroimaging. Neuroinformatics. 7(1):73–82.

Grupe DW. Nitschke JB. 2013. Uncertainty and anticipation in anxiety: An integrated neurobiological and psychological perspective. Nat Rev Neurosci. 14(7):488–501.

Keuken MC, Bazin PL, Backhouse K, Beekhuizen S, Himmer L, Kandola A, Lafeber JJ, Prochazkova L, Trutti A, Schäfer A, Turner R, Forstmann BU. 2017. Effects of aging on T1, T2*, and QSM MRI values in the subcortex. Brain Struct Funct. 222(6):2487–2505.

Kret ME, De Gelder B. 2012. A review on sex differences in processing emotional signals. Neuropsychologia. 50(7):1211–1221.

Kummer KK, Mitrić M, Kalpachidou T, Kress M. 2020. The medial prefrontal cortex as a central hub for mental comorbidities associated with chronic pain. Int J Mol Sci. 21(10):3440.

Leech R, Sharp DJ. 2014. The role of the posterior cingulate cortex in cognition and disease. Brain. 137(P7 1):12–32.

Linnman C, Moulton EA, Barmettler G, Becerra L, Borsook D. 2012. Neuroimaging of the periaqueductal gray: State of the field. NeuroImage. 60(1):505–522.

Liu BS, Lee YH. 2006. In-vehicle workload assessment: effects of traffic situations and cellular

telephone use. J Saf Res. 37(1):99–105.

Mantyh PW. 1982. Forebrain projections to the periaqueductral gray in the monkey, with

observations in the cat and rat. J Comp Neurol. 206(2):146–158.

Matsubara H, Hirano H, Hirano H, Soh Z, Nakamura R, Saeki N, Kawamoto M, Yoshizumi M,

Yoshino A, Sasaoka T, Yamawaki S, Tsuji T. 2018. Quantitative evaluation of pain

during electrocutaneous stimulation using a log-linearized peripheral arterial

viscoelastic model. Sci Rep. 8(1):3091.

Menon V, Uddin LQ. 2010. Saliency, switching, attention and control: a network model of

insula function. Brain Struct Funct. 214(5-6):655–667.

Murphy PR, O'Connell RG, O'Sullivan M, Robertson IH, Balsters JH. 2014. Pupil diameter

covaries with BOLD activity in human locus coeruleus. Hum Brain Mapp, 35(8):4140–

4154.

Okamoto Y, Sasaoka T, Sadato N, Fukunaga M, Yamamoto T, Soh Z, Nouzawa T, Yamawaki, S,

Tsuji T. 2020. Is human brain activity during driving operations modulated by the

viscoelastic characteristics of a steering wheel?: An fMRI study. IEEE Access.

8:215073–215090.

Paulus MP, Stein MB. 2006. An insular view of anxiety. Biol Psychiatry. 60(4):383–387.

Paxion J, Galy E, Berthelon C. 2014. Mental workload and driving. Front Psychol. 5:1344.

Ploner M, Lee MC, Wiech K, Bingel U, Tracey I. 2010. Prestimulus functional connectivity

determines pain perception in humans. Proc Natl Acad Sci U S A. 107(1):355–360.

Protzak J, Gramann K. 2018. Investigating established EEG parameter during real-world

driving. Front. Psychol. 9:2289.

Rao RPN, Ballard DH. 1999. Predictive coding in the visual cortex- a functional interpretation

of some extra-classical receptive-field effects. Nat Neurosci. 2(1):79–87.

Schneider M, Leuchs L, Czisch M, Sämann PG, Spoormaker VI. 2018. Disentangling reward

anticipation with simultaneous pupillometry / fMRI. NeuroImage. 178:11–22.

Schweckendiek J, Klucken T, Merz CJ, Tabbert K, Walter B, Ambach W, Vaitl, D, Stark R. 2011.

Weaving the (neuronal) web: Fear learning in spider phobia. NeuroImage. 54(1):681–

688.

Sehlmeyer C, Schöning S, Zwitserlood P, Pfleiderer B, Kircher T, Arolt V, Konrad C. 2009.

Human fear conditioning and extinction in neuroimaging: A systematic review. PLoS

ONE. 4(6):e5865.

Seth AK. 2013. Interoceptive inference, emotion, and the embodied self. Trends Cogn Sci.

17(11):565–573.

Sirois S, Brisson J. 2014. Pupillometry. Wiley Interdisciplinary Reviews: Cognitive Science. 5(6):679–92.

Somerville LH, Wagner DD, Wig GS, Moran JM, Whalen PJ, Kelley WM. 2013. Interactions between transient and sustained neural signals support the generation and regulation of anxious emotion. Cereb Cortex. 23(1):49–60.

Tang JS, Qu CL, Huo FQ. 2009. The thalamic nucleus submedius and ventrolateral orbital cortex are involved in nociceptive modulation: A novel pain modulation pathway. Prog Neurobiol. 89(4):383–389.

Thayer JF, Lane RD. 2009. Neuroscience and Biobehavioral Reviews Claude Bernard and the heart–brain connection: Further elaboration of a model of neurovisceral integration. Neurosci Biobehav Rev. 33(2):81–88.

Torrisi S, O'Connell K, Davis A, Reynolds R, Balderston N, Fudge JL, Grillon C, Ernst M. 2015. Resting state connectivity of the bed nucleus of the stria terminalis at ultra-high field. Hum Brain Mapp. 36(10):4076–4088.

Tsuji T, Arikuni F, Sasaoka T, Suyama S, Akiyoshi T, Soh Z, Hirano H, Nakamura R, Saeki N, Kawamoto M, et al. 2021. Peripheral arterial stiffness during electrocutaneous stimulation is positively correlated with pain related brain activity and subjective pain

intensity: an fMRI study. Sci Rep. 11(1):4425.

Wager TD, Waugh CE, Lindquist M, Noll DC, Fredrickson BL, Taylor SF. 2009a. Brain

mediators of cardiovascular responses to social threat. Part I: Reciprocal dorsal and

ventral sub-regions of the medial prefrontal cortex and heart-rate reactivity. NeuroImage.

47(3):821–835.

Wager TD, van Ast VA, Hughes BL, Davidson ML, Lindquist MA, Ochsner KN. 2009b. Brain

mediators of cardiovascular responses to social threat, Part II: Prefrontal-subcortical

pathways and relationship with anxiety. NeuroImage. 47(3):836–851.

Whitfield-Gabrieli S, Nieto-Castanon A. 2012. Conn: A functional connectivity toolbox for

correlated and anticorrelated brain networks. Brain Connect. 2(3): 125–141.