E-cadherin mediated AMIS localisation

Xuan Liang[1], Antonia Weberling[1], Chun Yuan Hii[1], Magdalena Zernicka-Goetz[1,2], Clare Buckley[1#]

[#] Corresponding author

[1] Department of Physiology, Development and Neuroscience

University of Cambridge

Downing Street

Cambridge

CB2 3BY

ceb85@cam.ac.uk

[2] Division of Biology and Biological Engineering

Caltech

Pasadena

Abstract

Individual cells within *de novo* polarising tubes and cavities must integrate their forming apical domains into a centralised apical membrane initiation site (AMIS). This is necessary to enable organised lumen formation within multi-cellular tissue. Despite the well documented importance of cell division in localising the AMIS, we have found a division-independent mechanism of AMIS localisation that relies instead on Cadherin-mediated cell-cell adhesion. Our study of *de novo* polarising mouse embryonic stem cells (mESCs) cultured in 3D suggest that cell-cell adhesion directs the localisation of apical proteins such as PAR-6 to a centralised AMIS. Unexpectedly, we also found that mESC cell clusters lacking functional E-Cadherin were still able to form a lumen-like cavity in the absence of AMIS localisation and did so at a later stage of development via a 'closure' mechanism, instead of via hollowing. This work suggests that there are two, interrelated mechanisms of apical polarity localisation: cell adhesion and cell division. Alignment of these mechanisms in space allows for redundancy in the system and ensures the development of a coherent epithelial structure within a growing organ.

Introduction

Most organs in the body arise from tubes or cavities made from polarised epithelial cells. These cells have a strict apico-basal orientation; they align their apical ends along a centrally located lumen. Some tubes, such as the anterior neural tube in amniotes, arise via folding and closure of an already polarised epithelial tissue, through mechanisms such as actomyosin-mediated apical constriction (Nikolopoulou et al., 2017). However, many tubes and cavities, such as the posterior neural tube, mammary acini, kidney tubules and mammalian epiblast arise via apical-basal polarisation within the centre of an initially solid tissue. The mechanisms by which such '*de novo'* polarisation is coordinated within dynamically growing tissue has been the focus of a significant body of research from several different models and have relevance both for understanding polarity-associated diseases and for directing organ bioengineering approaches.

Although the exact mechanisms are still under debate and may differ in different epithelia, Laminin, Integrin β1

and RAC1 signalling from the extra cellular matrix (ECM) is now well established to be necessary for directing

the overall apico-basal axis of polarisation of internally polarising tubes (Akhtar and Streuli, 2013; Bedzhov

and Zernicka-Goetz, 2014; Bryant et al., 2014; Buckley et al., 2013; Molè et al., 2021; Yu et al., 2005). What

is less clear is how the precise localisation of the apical membrane initiation site (AMIS) is directed at the single

cell level and how this is coordinated between neighbouring cells. The AMIS is a transient structure, marked

by the scaffolding protein Partitioning-defective-3 (PAR-3) and tight junctional components such as Zonula

occludens-1 (ZO-1), that defines where apically targeted proteins will fuse with the membrane, therefore

determining where the lumen will arise (Blasky et al., 2015; Bryant et al., 2010). It is important that the

subcellular localisation of the AMIS is coordinated between cells during morphogenesis to enable organised

lumen formation.

The current literature suggests that cell division plays an important role in AMIS localisation. In particular, the

post-mitotic midbody has been shown to anchor apically directed proteins (Li et al., 2014; Luján et al., 2016;

Rathbun et al., 2020; Schlüter et al., 2009; Wang et al., 2014). However, studies within the zebrafish neural

rod showed that, whilst misorientation of cell division results in disruption of the apical plane at a tissue level,

these phenotypes can be rescued by inhibiting cell division (Ciruna et al., 2006; Quesada-Hernandez et al.,

2010; Tawk et al., 2007; Zigman et al., 2011). We also previously demonstrated that individual zebrafish

neuroepithelial cells were able to recognise the future midline of the neural primordium and organise their

intracellular structure around this location in advance and independently of cell division. This resulted in the

initiation of an apical surface at whichever point the cells intersect the middle of the developing tissue, even if

this is part way along a cell length (Buckley et al., 2013). This suggests that, while cell division is undoubtably

a dominant mechanism in driving AMIS localisation, there must be an overlying symmetry-breaking event

during *de novo* polarisation. The earliest indication of midline positioning in the zebrafish neural rod was the

central accumulation of the junctional scaffolding protein Pard3 (PAR-3) and the adhesion protein N-Cadherin

(Buckley et al., 2013; Symonds et al., 2020). This led us to hypothesise that cell-cell adhesions could direct

the site for AMIS localisation during *de novo* polarisation. In line with this hypothesis, β-catenin mediated

maturation of N-cadherin was found to be necessary for the recruitment of the PAR apical complex protein

atypical protein kinase C (aPKC) in the chick neural tube (Herrera et al., 2021). Opposing localisations of ECM

and Cadherin proteins were also found to be sufficient to specify the apical-basal axis of hepatocytes in culture

(Zhang et al., 2020).

To test the role of cell-cell adhesions in AMIS localisation, we turned to mouse embryo stem cell (mESC)

culture in Matrigel, which has been used as an *in vitro* model for the *de novo* polarisation of the mouse epiblast

(Bedzhov and Zernicka-Goetz, 2014; Kim et al., 2021; Molè et al., 2021; Shahbazi et al., 2017). This allowed

us to study the initiation of apico-basal polarity of embryonic cells alongside the first cell-cell contacts between

isolated cells and small cell clusters. It also allowed us to determine within a mammalian model whether

division-independent polarisation is a conserved feature of *de novo* polarising structures. Unlike vertebrate

epithelial cell culture models such as Madin-Darby canine kidney (MDCK) cells, which initiate lumenogenesis

as early as the 2-cell stage when cultured in Matrigel (Blasky et al., 2015; Bryant et al., 2010), mESC cells in

Matrigel only form lumens at the multicellular stage after 48-72 hours in culture, coinciding with an exit in

pluripotency (Bedzhov and Zernicka-Goetz, 2014; Shahbazi et al., 2017). This results in a relatively clear

separation of the stages of *de novo* polarisation (figure 1A). Previous literature suggests that the AMIS is formed at the 2-cell stage, around 24-36 hours after culture in Matrigel, as denoted by membrane-localised PAR-3 and ZO-1 and sub-apical localisation of apical proteins such as Podocalyxin (PODYXL) (Shahbazi et al., 2017). The pre-apical patch (PAP) stage is formed after 36-48 hours in culture, as denoted by the fusion of apical proteins such as PODXYL, PAR-6 and aPKC to the apical membrane and the displacement of junctional proteins PAR-3, ZO-1 and E-Cadherin to the apico-lateral junctions (Kim et al., 2021; Shahbazi et al., 2017), following which lumenogenesis is initiated after 48-72 hours in culture.

To determine the role of cell division and of cell adhesion in mESC AMIS localisation, we analysed mESC cells cultured in Matrigel at the AMIS stage with and without cell division in wild type and E-Cadherin knock out cell lines. We then further analysed polarisation and lumenogenesis in the absence of E-Cadherin. Our results suggest that there is a division-independent mechanism of AMIS localisation that relies instead on E-Cadherin mediated cell-cell adhesions.

Results

Cell division is dispensable for AMIS localisation

First, we tested whether cell-division was necessary for AMIS localisation in mESC rosettes. We cultured naïve, unpolarized mESCs (ES-E14 cells) in 2D on gelatin with 2i/LIF and then treated them with mitomycin C to block cell division. We then isolated single cells and seeded them into Matrigel without 2i/LIF, in N2B27 differentiation medium (Figure 1B). Cell divisions were efficiently blocked during the first 24 hours post seeding, during which time individual cells contacted each other and formed cell clusters in the absence of cell division (Supplementary Movie 1, Figure S1 A,B).

To assess AMIS localisation, we carried out immunohistochemistry (IHC) for PAR-3 and ZO-1 at 24hrs post seeding. As previously published (Shahbazi et al., 2017), in addition to several puncta at the cell peripheries, both PAR-3 and ZO-1 localised to the membrane at the centre of cell-cell contacts, marking the AMIS in the majority of cell clusters (Figure 1Ci,Gi and S1E,F). Interestingly, division-blocked cells also localized PAR-3 and ZO-1 to the central membrane (Figure 1Cii,Gii, and S1E,F, quantified in 1D,I). In both control and division-blocked cells, there was a small proportion that had not yet localised the AMIS at the 24-hour stage (Figure 1D,I & S1D). E-Cadherin was upregulated along the whole length of the cell-cell interfaces in both dividing and non-dividing cell clusters, with a higher level of E-Cadherin at the cell-cell interface relative to the cell-matrix interface (Figure 1C and S1C,E). To quantify the subcellular localization of PAR-3, we carried out intensity profiles across the cell-cell interface of doublets (Figure 1F). This confirmed that PAR-3 localised to a small central area at the cell-cell interface in both control and division-blocked cells (Figure 1E and 3C). Golgi apparatus and centrosomes were also localised to the centre of cell-cell contacts both in dividing and non-dividing conditions (Figure 1G-J & S1F), confirming that mESCs were polarised in the absence of cell-division. Not only by mitomycin C treatment, but when we used aphidicolin to block cell divisions (Figure S1G), the division-blocked mESC clusters were also able to polarise PAR-3 (Figure S1H), ZO-1 and the Golgi apparatus (Figure S1I) upon making cell-cell contacts.

125 Together, these results demonstrate that cell division is dispensable for *de novo* AMIS localisation in polarizing
126 mESCs.

Cell-cell contact directs PAR-6 localisation

To understand the dynamics of apical protein polarisation in the absence of cell division, we generated a mESC
stable cell line expressing mCherry-PAR-6B and imaged cells live. In line with previous characterisation of
PAR-6 by IHC (Kim et al., 2021; Shahbazi et al., 2017), in control dividing cells, mCherry-PAR-6B localized to
the apical membrane by the PAP stage at 48 hours and to the luminal apical membrane from 72h (Figure 2A).
In addition, the transgene allowed us to better visualize PAR-6B puncta at earlier 24h AMIS stages of
development, before they had associated with the apical membrane. At this stage, PAR-6B was localized sub-
apically, polarized to towards the central region of cell-cell contact (Figure 2A). A similar polarised distribution
of PAR-6B was observed in both control and division-blocked cells (Figure 2B,C), demonstrating that cell
division is dispensable for apical protein polarisation.

We next assessed the dynamics of PAR-6B polarisation. In both control and division-blocked cells, non-cortical
mCherry-PAR-6B puncta were visible at the single cell stage. In control cells, these puncta then relocalised to
the abscission plane, following cell division (Figure 2D, supplementary movie 2). In division-blocked cells,
PAR-6B puncta dynamically relocalised to newly forming cell-cell contacts, eventually forming cell-cell clusters
with centrally localised PAR-6B (Figure 2D, supplementary movie 2).

These results suggest that cell-cell contact directs PAR-6B localisation at the central AMIS, independent of
cell division.

E-Cadherin adhesions are necessary for AMIS localisation

The above results suggest that there is a division-independent mechanism of AMIS localisation that relies
instead on cell-cell adhesions. Since E-Cadherin is the predominant adhesion molecule in non-neural epithelia,
we hypothesised that it might be important in AMIS localisation. To achieve a full removal of E-cadherin, we
employed an *E-Cadherin* knock-out (*Cdh1* KO) mESC line (Larue et al., 1996). Whilst E-Cadherin expression
was lost in *Cdh1* KO cells, they maintained P-Cadherin expression and were still able to form cell clusters
when cultured in Matrigel (Figure S2A, B).

To assess AMIS localisation, we again carried out IHC for PAR-3 and ZO-1 at 24hrs post seeding. As seen
earlier (Figure 1), both PAR-3 and ZO-1 localised to the central region of cell-cell contact within wild type (ES-
E14) doublets/clusters with and without division. However, PAR-3 and ZO-1 localisation was strongly inhibited
in the absence of E-Cadherin (Figure 3A-C). To confirm that these differences were not to do with the different
cell line backgrounds of WT and *Cdh1* KO cells, we also used RNAi to knock-down E-cadherin in ES-E14
mESCs. In contrast to the control two-cell clusters that formed the centralised apical PAR-3 domain after cell
divisions, PAR-3 at the central region of E-cadherin knock-down two-cell clusters was significantly reduced,
though E-cadherin was still sightly detectable upon the knock-down (Figure S2C, D).

To investigate AMIS localisation at a single cell level, we co-cultured division-blocked wild type (ES-E14) and
Cdh1 KO cells and analysed division-blocked chimeric mESC doublets, comprising one control and one *Cdh1*

KO cell. Whilst homogenous control doublets localised PAR-3 to the central region of the cell-cell interface, heterogeneous chimeric doublets did not localise PAR-3 centrally (Figure 3D,E). Golgi and centrosome localisation towards the cell-cell interface suggested that the overall axis of polarity was maintained, even in the absence of both cell division and E-Cadherin (Figure 3F-H & S2E).

We then analysed E-Cadherin at the centre-most regions and side regions flanking the centre-most regions in two-cell mESC clusters. E-Cadherin immunofluorescence signals (Figure S3A, B), as well as E-cadherin-GFP in the mESC line that expresses E-cadherin-GFP (Molè et al., 2021), were the same at these two regions (Figure S3C, D). However, fluorescence recovery after photobleaching (FRAP) of E-Cadherin-GFP at the two regions showed that the mobile fraction of E-cadherin-GFP was lower in the centre-most region than the side regions (Figure 3 I, J). Therefore, E-cadherin junctions are more stable at the centre-most region of the cell-cell interface, which may provide at least a partial explanation for why AMIS localisation occurs precisely at this region.

It has previously been demonstrated that a reduction in E-Cadherin can slow pluripotency exit (Soncin et al., 2009). However, pluripotency exit was previously shown not to alter AMIS formation (Shahbazi et al., 2017). In support of these results, we also found that cells maintained in the pluripotent state when cultured in Feeder cell medium provided with 2i/LIF still localised the AMIS, with and without cell division (Figure S4A-C). However, in line with our results showing lack of AMIS localisation in *Cdh1* KO cells cultured in the absence of 2i/LIF (Figure 3), cells cultured in the presence of 2i/LIF also could not localise an AMIS in the absence of E-Cadherin (Figure S4A-C). Despite this result, we wanted to check whether the stage of pluripotency exit differed between WT and *Cdh1* KO cells in our experiments since this might indicate a different speed of maturation. We therefore carried out IHC for Orthodenticle Homeobox 2 (OTX2) protein, which is necessary for pluripotency exit, and the pluripotency marker protein Nanog. Although, as expected, the overall level of nuclear OTX2 increased and Nanog decreased over the 24-hour course of development, we found no significant difference in post-mitotic levels of OTX2 and Nanog between WT and *Cdh1* KO cells (Figure S4D). This result suggests that there was no difference in the stage of pluripotency exit in the cell clusters that we analysed during this study and this is therefore unlikely to play a role in the lack of AMIS localisation seen in *Cdh1* KO cells.

These results demonstrate that E-Cadherin adhesions between cells are necessary for AMIS localisation but not for the overall axis of polarity. They also demonstrate that ECM in the absence of E-Cadherin is insufficient for AMIS localisation.

E-Cadherin adhesions are sufficient to initiate AMIS localisation, independent of ECM signalling and cell division

As discussed, ECM-mediated signalling plays an important role in orienting the axis of polarisation within *de novo* polarising systems. Recently, the apico-basal axis of cultured mature hepatocytes was established by a combination of ECM signalling and immobilised E-Cadherin (Zhang et al., 2020). However, PAR-3 has also recently been shown to polarise in mESCs lacking functional Integrin-β1 or cultured in agarose in the absence of ECM proteins (Molè et al., 2021). Our current study shows that the AMIS can localise in the absence of cell division but not in the absence of E-Cadherin. We therefore wanted to explore the relative roles of ECM, cell division and E-Cadherin in AMIS localisation.

To test this, we first eliminated the influence of ECM by culturing division-blocked mESCs (ES-E14) in 0.5%

Agarose and carried out IHC for PAR-3 after 30 hours in culture. These cells were still able to polarise PAR-3,

even in the absence of both cell division and ECM proteins (Figure 4A,B). However, in line with our earlier

results (Figure 3), PAR-3 localisation was strongly inhibited in *Cdh1* KO cells (Figure 4A,B). These results

suggest that AMIS localisation occurs independently of both ECM signalling and of cell division, relying instead

on E-Cadherin.

To test the sufficiency of E-Cadherin adhesions in initiating AMIS localisation, we cultured individual division-

blocked mESCs (ES-E14) onto either E-Cadherin recombinant protein or Fibronectin pre-coated glass, then

topped the cells with N2B27 medium, with or without 20% Matrigel and carried out IHC for PAR-3 after 24

hours in culture. Like results from hepatocytes (Zhang et al., 2020), cells plated on E-Cadherin and topped

with Matrigel localised PAR-3 to the centre of the cell-Cadherin interface (Figure 4Ci,D,E). However, this

central PAR-3 localisation was significantly reduced when cells were plated on Fibronectin (Figure 4Cii,D,E).

Interestingly, cells cultured on E-Cadherin but in the absence of Matrigel still localised PAR-3 to the centre of

the cell-Cadherin interface (Figure 4 Ciii,D,E).

These results demonstrate that E-Cadherin adhesions are both necessary and sufficient for initiating AMIS

localisation, while ECM is not necessary or sufficient for AMIS localisation.

E-Cadherin is necessary for hollowing lumenogenesis

We next wanted to test the importance of E-Cadherin-mediated AMIS localisation in lumenogenesis. We

therefore cultured WT (ES-E14) and *Cdh1* KO mESCs and fixed them at the AMIS 24-hour stage, PAP 48-

hour stage and lumen 72 and 96-hour stage. We then carried out IHC for PAR-3 and ZO-1 to label

AMIS/apicolateral junctions and PODXYL to label apical proteins. Whilst most WT cell clusters had a

centralised apical domain or small lumen after 48 hours in culture, very few *Cdh1* KO cell clusters had made

a centralised apical domain by the 48-hour PAP stage (Figure 5A-D). In line with our earlier findings at the 24-

hour AMIS stage (Figure 3), this provides further evidence that E-Cadherin is necessary for centralised AMIS

localisation. However, we noticed that a small percentage of *Cdh1* KO cell clusters at 48 hours had formed an

open 'cup-shape', with apically localised PODXYL and apico-laterally localised junctional PAR-3 and ZO-1

(e.g. figure 5Aiii). We termed these 'open cavities' (Figure 5C). Surprisingly, by the 72-hour lumen stage,

approximately 75% of *Cdh1* KO cell clusters had formed polarised cavities, approximately 50% of which were

open cavities and 50% were closed (Figure 5A-E, Figure S5A). Over the course of 48-96 hours in culture, the

overall percentage of polarised cavities increased (Figure 5D) as did the proportion of these structures that

were 'closed' (Figure 5E). This suggested that these cavities might form via gradual 'closure' of the tissue,

rather than via hollowing. Both 'open' and 'closed' cavities were surrounded by polarised Golgi apparatus,

demonstrating that the overall apico-basal axis of cells was in-tact. (Figure S5B).

To further assess the morphogenetic mechanism by which cavities form in *Cdh1* KO cells, we generated WT

(ES-E14) and *Cdh1* KO mESC lines labelled with LifeAct-mRuby. We visualised the process of lumenogenesis

within mESCs cultured in Matrigel via live imaging. Whilst the WT cell cluster expanded the already central

lumen, the open cup-shaped cavity within *Cdh1* KO cell clusters gradually closed, eventually generating a

centralised lumen-like structure without hollowing at a later stage of development (Figure 5F and
supplementary movie 4).

These results confirm that, in the absence of E-Cadherin mediated AMIS localisation, cell clusters do not
hollow but instead generate lumen-like cavities via a closure mechanism (Figure 5G). Our results also
demonstrate that E-Cadherin and centralised AMIS localisation are not required for apical membrane formation.
In the absence of E-Cadherin, an apical surface is still formed but this occurs later in development so appears
less efficient.

Discussion

Epithelial cells can polarise *de novo* in the absence of cell division.
Both AMIS associated proteins PAR-3/ZO-1 and apical polarity protein PAR-6B localised similarly in WT and
division-blocked mESCs (Figures 1 and 2). This finding supports our previously published zebrafish
neuroepithelial cell *in vivo* analysis, which demonstrated the division-independent localisation of Pard3 (PAR-
3) and ZO-1 at the neural rod primordial midline (Buckley et al., 2013). Together, this demonstrates that
although division is an important contributor to AMIS formation, a division-independent mechanism of *de novo*
polarisation and AMIS localisation can occur in both *in vivo* and *in vitro* conditions. Whilst disorganised lumen
formation can also occur in the absence of division in the zebrafish neural rod (Buckley et al., 2013), this was
not possible to test within the mESC culture model since Mitomycin treated cell clusters did not survive beyond
30 hours in culture.

E-Cadherin based cell-cell contacts are necessary and sufficient to initiate AMIS localisation
AMIS localisation in *Cdh1* KO cell clusters is strongly inhibited (Figure 3) and individual mESC cells can direct
their AMIS to the central region of the cell-Cadherin interface independently from ECM-signalling (Figure 4).
Together this demonstrates that formation of Cadherin-based cell-cell contacts is both necessary and sufficient
for initiating AMIS localisation and that ECM is insufficient to direct AMIS localisation in the absence of
Cadherin. Our results therefore suggest that Cadherin-based cell-cell adhesion may provide the spatial cue
required for AMIS localisation during *de novo* polarisation. This in turn directs apical proteins such as PAR-6B
to a centralised region of cell-cell contact (Figure 2), determining where the lumen will hollow.

Whilst we demonstrate that AMIS localisation can occur independently from cell division, the importance of
abscission and midbody formation in apical protein targeting has been robustly demonstrated and the
molecules involved are now starting to emerge (Klinkert et al., 2016; Li et al., 2014; Luján et al., 2016; Mangan
et al., 2016; Rathbun et al., 2020; Schlüter et al., 2009; Wang et al., 2021, 2014). Rather than acting as the
initial symmetry breaking step in AMIS localisation, we suggest that tethering of apically directed proteins to
the midbody might instead act to transiently align cell division, cell adhesion and the apical domain, therefore
enabling an organised structure to be generated in the presence of dynamic cell movement and tissue growth.
The transient localisation of scaffolding and tight junction-associated proteins such as PAR-3 and ZO-1 at the
AMIS might aid in this alignment. For example, Cingulin is a tight junctional protein that has been shown to
bind both to the midbody and to FIP5, which is important for the apical targeting of vesicles containing apical
proteins (Mangan et al., 2016). During zebrafish neural rod development, cell adhesion and cell division align

to allow an organised structure to arise from dynamically reorganising cells (Symonds and Buckley 2020) and loss of N-cadherin results in mis-oriented cell divisions and a disrupted apical domain (Zigman et al., 2011). Once apical proteins fuse with the apical membrane, proteins associated with junctions such as Cadherin, PAR-3 and ZO-1 are then cleared from the apical surface and instead form the apicolateral junctions, as demonstrated in several different epithelial systems (Kim et al., 2021; Morais-de-Sa et al., 2010; Symonds et al., 2020).

Whilst we have demonstrated that E-Cadherin directs AMIS localisation, we do not yet have a full explanation for why AMIS proteins localise at the central-most point of cell-cell contact in the absence of divisions, despite E-Cadherin localisation all along the cell-cell interface. PAR-3 and PAR-6 have been found to directly interact with Cadherin proteins (Iden et al., 2006; Renschler et al., 2018). Recently, opposing actin flows in migrating cells as they first encounter each other were found to be responsible for regulating the first AJ deposition via tension-mediated unfolding of a-catenin and further clustering of surface E-Cadherin molecules (Noordstra et al., 2021). Together, this could provide an explanation for how the first contacts between cells could act as an apical 'seed', therefore defining the position of the AMIS within multicellular tissues. This could therefore explain why we have previously seen an upregulation of N-Cadherin at the zebrafish neural rod midline, where cells growing from either side of the organ primordium meet (Symonds et al., 2020). However, it is still unclear how this might regulate the subcellular localisation of the AMIS to the centre of cell-cell contacts. Our FRAP results demonstrate that E-Cadherin is relatively more stable at the central-most point of contact between two adhering cells (Figure 3I,J), which could explain this central AMIS localisation and suggests that E-Cadherin is more stably bound via its downstream partners to the internal actin cytoskeleton at this point. In line with this hypothesis, previous publications have demonstrated an upregulation of phosphorylated MYOSIN-II (pMLC) at the AMIS (Molè et al., 2021), which is suggestive of higher actomyosin-mediated tension. Uncovering the mechanisms directing adhesion-dependent AMIS localisation precisely to the midpoint of cell-cell adhesions will be an interesting area for future studies. In addition, a recent study of chick neural tube polarisation (where N-Cadherin is the dominant Cadherin) has demonstrated that the interaction of β-catenin with pro-N-cadherin in the Golgi apparatus is necessary for the maturation of N-Cadherin, which is in turn important for apicobasal polarity establishment (Herrera et al, 2021). This provides the possibility that the polarised Golgi apparatus that we observe in the mESC clusters might be directionally delivering mature E-Cadherin to the central-most region of cell-cell contact.

In the absence of an AMIS, lumens form via 'closure' rather than hollowing

The centralised localisation of an AMIS appears necessary to enable lumen hollowing within multi-cellular clusters. *Cdh1* KO cells lack AMIS localisation at the 24-hour AMIS stage (Figure 3). However, they still retain their apico-basal polarity axis (as denoted by Golgi apparatus and centrosome localisation, Figure 3F-H & S2C) and form apico-lateral junctions at luminal stages of development (from approximately 72 hours in culture, Figure 5). Therefore, *Cdh1* KO cells do appear to still make an apical membrane but do so more slowly than in WT cells and without going through a centralised AMIS stage. This suggests that the role of E-Cadherin in *de novo* polarisation is specifically to localise the AMIS, which enables the integration of individual cell apical domains to a centralised region preceding lumen hollowing. The lack of a centralised AMIS in Cadherin deficient cells could also explain the multiple-lumen (but otherwise polarised) phenotypes previously seen in E-Cadherin deficient MDCK cells cultured on collagen (Jia et al., 2011).

A surprising observation was the ability of *Cdh1* KO mESC clusters, in the absence of AMIS localisation, to instead form 'lumen-like' structures via a 'closure' process. We do not currently know the mechanism by which such 'closure' occurs. However, the presence of F-actin and p-MLC rich cable-like structures in 'cup'-shaped open cavities is potentially suggestive of a contractile process (Figure S5B). We were unable to live-image earlier stages of *Cdh1* KO cluster development due to the sensitivity of these cells to phototoxicity so we can't exclude the possibility that hollowing lumenogenesis occurs in parallel, although our IHC analysis does not indicate that this is the case. Understanding the relative roles of mechanics in localisation of the AMIS and in 'opening' vs. 'closing' tubes is an important future research goal, as is the potential role of cell geometry in mediating such differences. Additionally, collective cell migration could play a role in this 'closure' mechanism. Collective inwards migration of cells caused lumen formation via a folding mechanism when MDCK monolayers were overlaid with a soft collagen gel (Ishida et al., 2014). A similar collective process could be occurring in the *Cdh1* KO cell clusters from our study, which were cultured in a soft (20%) Matrigel and formed loosely connected cell clusters, which then 'closed' to make a centralised lumen.

In summary, our work suggests that Cadherin-mediated cell-cell adhesion directs AMIS localisation during *de novo* polarisation of epithelial tubes and cavities. Our work also suggests that ECM is insufficient to direct AMIS localisation in the absence of Cadherin. In parallel with the well described role of the midbody in tethering apical proteins, this suggests that there are two, interrelated mechanisms of AMIS localisation: cell adhesion and cell division. The alignment of these cellular processes allows for redundancy in the system and provides an explanation for how an organised epithelial structure can arise within the centre of a proliferating organ primordium.

Material and Methods

Cells

Wild-type mESCs (ES-E14) were purchased from Cambridge Stem Cell Institute. *Cdh1* KO mESCs were gifted from Lionel Larue lab at Institute Curie. mESC carriers were maintained in Feeder Cell Medium in Corning cell culture dishes precoated with 0.1% Gelatin (ES-006-B, Sigma-Aldrich), at 37 ℃ suppled with 5% CO_2 at one atmospheric pressure. To maintain cells in pluripotency, 2i/LIF (1 mM MEK inhibitor PD0325901, 13034, Cayman Chemical; 3 mM GSK3 inhibitor CHIR99021, 13122, Cayman Chemical; and 10 ng/ml LIF, A35933, Gibco) was added to the culture medium to preserve naive pluripotency. The culture medium was renewed every three days. The cells were trypsinized when reaching confluency to be passaged or subjected to experiments.

Mouse PAR-6B CDS cDNA were assembled with mCherry by Gibson assembly, LifeAct-Ruby cDNA were sub-cloned from an existing plasmid. The cDNAs were cloned into pDONR221 plasmid and introduced into the PB-Hyg-Dest plasmid using Gateway technology (Thermo Fisher Scientific). The PB-Hyg-Dest-mCherry-PAR-6B or the PB-Hyg-Dest-LifeAct-Ruby plasmid was co-transfected with the piggyBac plasmid using Lipofectamine 3000 to generate Hygromycin B resistant stable cell lines. The mCherry-PAR-6B or LifeAct-

Ruby expressing mESC stable cell line was created via 10 µg/mL Hygromycin B selection and single cell
colonies expansion.

Cell cultures and treatment

For 3D culture of wild-type and E-cadherin knock-out mESCs, 20µL of Matrigel (356231, Corning, Lot 354230,
354234, 356231) was spread evenly to the bottom of each well in a µ-slide 8 well dish (80821, Ibidi). The dish
was left on ice for 10 minutes to flatten the Matrigel surface, then was left at 37 ℃ for 10 minutes to solidify the
Matrigel. mESCs were trypsinized, pipetted thoroughly and passed through a cell strainer (431750, Corning)
to isolate cells into single cells. Singled mESCs were suspended in the N2B27 medium and seeded onto the
solidified Matrigel. The seeded density was control, 14 cells/mm^2; Mitomycin C treated, 227 cells/mm^2. The
cells were left at 37℃ for 15min when over 95% of the cells attached to the Matrigel, then the culture medium
was renewed to 10% Matrigel/N2B27 medium with or without 2i/Lif.

For control and *Cdh1* KO mESC chimeric cluster cultures, wild-type and *Cdh1* KO mESCs were mixed at 1:4
ratio and co-cultured at 2D in the Feeder Cell Medium. They were then treated with Mitomycin C for 2 hours,
trypsinized and seeded for 3D Matrigel culture at 227 cells/mm^2.

For mESC cultured in Agarose, 5,000 control or 125,000 Mitomycin treated cells were suspended in a 37 ℃
warmed 20 µL 0.5% low melting point Agarose (16520, Invitrogen) droplet at the bottom of the µ-slide 8 well
dish. The dish was left at room temperature for 5 minutes to solidify and topped with the N2B27 medium. The
cells were then cultured at 37 ℃, 5% CO$_2$ until analysis.

For cells cultured on E-cadherin and fibronectin coated glass, the µ-slide 8 well dish was incubated with
nitrocellulose/methanal at 37℃ for 3 hours and then was left air dried. The dish was then incubated with
10µg/mL mouse E-cadherin recombinant protein (8875-EC-050, Bio-Techne) or fibronectin at 4℃ overnight.
The dish was briefly washed with water. Mitomycin C pre-treated ES-E14 cells were seeded onto the dish at
14 cells/mm^2 in N2B27 medium. The cells were allowed to attach to the glass at 37 ℃ for one hour, then the
medium was renewed to N2B27 medium with 20% Matrigel. The cells were fixed 24 hours post the Matrigel
introduction.

For Mitomycin C treatment, the cells were incubated with 10 µg/mL Mitomycin C (J63193, Alfa Aesar) in culture
media at 37 ℃ for two hours. The Mitomycin C contained media were removed and the cells were washed
with PBS briefly. Then the Mitomycin C treated cells were trypsinized and subjected to further experiments.

Compositions of the culture media were as follows. Feeder Cell Medium: DMEM (41966, Thermo Fisher
Scientific), 15% FBS (Stem Cell Institute), penicillin–streptomycin (15140122, Thermo Fisher Scientific),
GlutaMAX (35050061, Thermo Fisher Scientific), MEM non-essential amino acids (11140035, Thermo Fisher
Scientific), sodium pyruvate (11360070, Thermo Fisher Scientific) and 100 µM β-mercaptoethanol (31350-010,
Thermo Fisher Scientific). N2B27 Medium: 1:1 mix of DMEM F12 (21331-020, Thermo Fisher Scientific) and
neurobasal A (10888-022, Thermo Fisher Scientific) supplemented with 2% v/v B27 (10889-038, Thermo
Fisher Scientific), 0.2% v/v N2 (17502048, Gibco), 100 µM β-mercaptoethanol (31350-010, Thermo Fisher

Scientific), penicillin–streptomycin (15140122, Thermo Fisher Scientific) and GlutaMAX (35050061, Thermo Fisher Scientific).

Immunofluorescence

Cells cultured in a μ-slide 8 well dish were fixed with 4% paraformaldehyde (J61899, Alfa Aesar) for 30 minutes at room temperature, then were permeabilised with 0.5% Triton X-100 for 15 minutes at room temperature. The cells were blocked with the incubation buffer (0.5% BSA, 0.1% Tween in PBS) for two hours, then were incubated with primary antibodies diluted in the incubation buffer at 4 ℃ overnight on shaking. The primary antibodies were washed off with PBS, then were incubated with secondary antibodies diluted in the incubation buffer at room temperature for two hours. The secondary antibodies were washed off with PBS. Samples cultured in the Matrigel were kept in PBS; samples cultured in Agarose were sealed in 200 μL 0.5% Agarose. The samples were imaged shortly after. Antibodies and dilutions were as listed below.

Primary antibody list

Protein	Catalog number	Supplier	Type	Specie	Concentrations
E-cadherin	ECCD-2	Invitrogen	Monoclonal	Rat	2 μg/mL
GM130	610822	BD Biosciences	Monoclonal	Mouse	1:300
Nanog	ab80892	Abcam	Polyclonal	Mouse	1:300
OTX2	AF1979	R&D Systems	Polyclonal	Goat	1:300
Podocalyxin	MAB1556	R&D Systems	Monoclonal	Rat	3.3 μg/mL
PAR-3	07-330	Merck Millipore	Polyclonal	Rabbit	1:100
P-cadherin	AF761	R&D Systems	Polyclonal	Goat	1:500
Γ-Tubulin	T6557	Sigma-Aldrich	Monoclonal	Mouse	1:250
ZO-1	61-7300	Invitrogen	Polyclonal	Rabbit	1:500

Secondary antibody list

Name	Supplier	Target	Concentrations
Alexa Fluor 488	Invitrogen	Rat	1:500
Alexa Fluor 488	Invitrogen	Rabbit	1:500
Alexa Fluor 488	Invitrogen	Goat	1:500
Alexa Fluor 546	Invitrogen	Mouse	1:500
DyLight 550	Invitrogen	Rat	1:500
CF 633	Insight Biotechnology	Phalloidin	6.6 pM

Microscope imaging

Live cell imaging was carried out on the PerkinElmer UltraVIEW spinning disk system fitted on an Olympus IX80 confocal microscope with a 37 ℃ and 5% CO_2 chamber. Images were captured with the 40x 1.3 NA (oil) objective, 2x Hamamatsu Orca-R2 CCD camera and Volocity 3.7.1 software. The cells were imaged at 2 μm

z-step size and 30 minutes time intervals. Fixed samples were imaged on the Leica SP8 confocal microscope with the 40x 1.3 NA (oil) or 63x 1.4 NA (oil) objective and LAS X 3.7.4 software. The cells were imaged at 0.3 μm z-step size and 2X line average. FRAP was performed on the Zeiss LSM-900 confocal microscope with 63X, 1.40NA Plan Apo objective and ZEN Blue 2.1 software equipped with a 37 °C heated stage.

Image and data analysis

The central section images were projected from raw images in the Fiji software by maximum-value projection of the whole z stacks to produce the 3D projection images, or of the central three image of the z stacks to produce the central-section images. The whole z stacks projections were applied to count clump percentages. The central-section images were applied for line-scans to determine protein signals.

The mESCs with positive or negative protein centres were manually determined and counted. Based on cell cluster morphologies displayed by F-actin signals and nucleus morphologies displayed by DAPI signals, only cells at the interphase were analysed. The percentage was calculated with the number relative to the number of total clumps captured in each condition. The mean percentages from three independent experiments were compared with student's t-test or one-way ANOVA specified in figure legends using the GraphPad Prism software. Sample sizes are specified in figure legends.

To quantify PAR-3 signal along the cell-cell interface, a 0.8μm width line was drawn alone the cell-cell interface by using F-actin (labelled by Phalloidin) or E-Cadherin signal as the path. PAR-3 and F-actin Pixel values along the path were extracted. The two peaks of F-actin signals at two ends of the path were determined as the start of end of the cell-cell interface and the positions in-between were defined as 1X cell-cell interface. The corresponding PAR-3 pixel to the F-actin peak positions were identified and the PAR-3 line profile between the two positions were sectioned to 20 sections. PAR-3 pixel value in each section was averaged to be the PAR-3 signal in 5% of the cell-cell interface. The values from 10-15 cells were plotted as line graphs.

To generate the PAR-3 signal heatmap, based on the F-actin signal, a squared region of interest was created in the Fiji software for every cell, which the main cell body of a cell was fitted into. The PAR3 image in each squared region was transformed into a 10 x 10-unit matrix by using the R-language. Each unit was created by averaging the PAR3 fluorescence intensity in every 10% length along the X or Y axis of the squared region of interest extracted from the original image. A serial of images from one condition of an experiment was stacked and the averaged pixel value at each position was calculated to generate an averaged 10 x 10 matrix. The final matrix was transformed into a heatmap in the GraphPad Prism software.

To compare the level of PAR-3 in the core areas in the cells culture on the glass, a 6 μm diameter circular area of interest was created in the Fiji software to cover the PAR3 core area in a cell, and the average PAR3 fluorescence values in the circle was measured. The boundary of the cell was drawn by the free-hand tool in Fiji, and the average PAR-3 fluorescence values in the boundary was measured. The ratio between the PAR3 values in the circle and inside the boundary was then calculated.

FRAP

FRAP of GFP-E-Cadherin was performed to measure E-Cadherin dynamics at the cell-cell interface. A GFP-E-Cadherin expressing stable ES-E14 line was used to perform the experiment (Molè et al., 2021). After 24 hours culturing in Matrigel, two-cell clusters with long axis parallel to surface of the culture dish, hence the cell-cell interfaces were aligned to the axis of the objective were used for FRAP. A ROI of 2 X 1 µm along xy-axis was chosen at the approximately centre-most region or side regions at each cell-cell interface. The ROI in a series of z-axis stacks with 0.3 µm intervals cross 2 µm was bleached with 3 iterations of the 488 nm laser with 100% transmission. This resulted in a photobleaching of over 80% in a 2 X 2 µm (xz) region at the cell-cell interface. Time-lapse images were acquired before (3 frames) and after (30 frames) photobleaching with an interval of 10s per frame. Average fluorescence intensity values $F(t)$ in the bleached area within the centre z-stack were analysed with ImageJ. The mean values of the three frames before bleaching was used as the pre-bleached value $F(i)$. The value of the first frame after bleaching was defined as $F(0)$. FRAP values were then calculated and plotted over time in Figure 3I as:

$$\mathrm{FRAP} = \frac{F(t) - F(0)}{F(i) - F(0)}$$

The FRAP values were fitted using a non-linear regression and the exponential one-phase association model using Y0 = 0 and where mobile fraction corresponds to the plateau value in Prism software. For Figure 3J, the mobile fraction from each FRAP profile was pooled and compared between conditions.

Supplementary Material and Methods

Aphidicolin treatment

For Aphidicolin treatment, mESCs cultured on gelatin were incubated with 5 µM Aphidicolin (Sc-201535, Insight Biotechnology) in culture media at 37 ℃ for two hours. The treated cells were trypsinized and seeded into 10% Matrigel in culture media containing 5 µM Aphidicolin till further analysis.

Line-scan analysis of E-cadherin
To quantify E-cadherin level along the cell membrane in Figure S1C, the central z-stack from confocal microscope raw images was extracted. On that image, a line of 6 µm wide were drawn perpendicular against the cell-cell membrane or cell-ECM membrane in Fiji. The peak fluorescence value of E-cadherin was extract from the line-scan profile. Three line-scans were done at different regions of the cell-cell membrane in one 2-cell cluster, and three line-scans were done at the cell-ECM membrane in each cell of the 2-cell clump. The average value of the three or six line-scans was regarded as the E-cadherin level at the cell-cell membrane or the cell-ECM membrane of that 2-cell clump. To quantify E-cadherin level at the central-most or side regions at the cell-cell interface in Figure S3B, a line of 3 µm wide was drawn perpendicular with the cell-cell interface to generate the line-scan profile. The region corresponding to the polarised PAR-3 dot was determined as the central-most region, and regions between the PAR3 centre and cell-free regions was determined as the side region. Two side regions were analysed at each side flanking the centre region, and the average line-scan profile of the two was taken as the line-scan profile at the side region of a 2-cell cluster.

siRNA transfection

To achieve E-cadherin knock-down, siRNA was transfected using Lipofectamine RNAiMAX according to the manufacturer's instruction. mESCs cultured on the gelatin was transfected with 100 nM E-cadherin pre-designed siRNA (s63752, Silencer Select). The siRNA target sequence is GAAGAUCACGUAUCGGAUU. Three days after the transfection, the cells were seeded into 10% Matrigel. The cells were cultured in Matrigel with Lipofectamine RNAiMAX and 100 nM siRNA for 24 hours, then were subjected to further experiments and analysis.

Analysis of nuclear OTX2 or NANOG

To quantify nuclear OTX2 or NANOG level, raw images were captured on the confocal microscope with OTX2 or NANOG and DAPI channels. The raw images that contained z-axis stacks were subjected to maximum value projections in Fiji. The DAPI channel was turned into a binary image and the region under DAPI signals were masked by using the Analyse Particles tool in Fiji. The total OTX2 or NANOG fluorescence values in each cell that were under the correspondent DAPI mask was measured and regarded as the OTX2 or NANOG level of that cell. Based on cell cluster morphologies displayed by F-actin signals and nucleus morphologies displayed by DAPI signals, only cells at the interphase were analysed.

Acknowledgements

We are grateful to Jon Clarke and Ben Steventon for critical reading of the manuscript and other members of the Buckley and Zernicka-Goetz labs for scientific discussion, especially Matteo Molè and Marta Shahbazi. Thank you to the Lionel Larue lab for kindly gifting the *Cdh1* KO cells.

Funding: This research was financially supported by: CEB - the Wellcome Trust and Royal Society (Sir Henry Dale Fellowship grant no. 208758/Z/17/Z and Dorothy Hodgkin Fellowship grant no. DH160086), XL - European Union's Horizon 2020 programme (Marie Skłodowska-Curie Individual Fellowship grant no. 844330), the Issac Newton Trust and Leverhulme Trust (Leverhulme Early Career Fellowship grant no. ECF-2019-175). MZG - The Wellcome Trust (207415/Z/17/Z) and ERC (669198).

Thank you to the Cambridge Advanced Imaging Centre and the Ewa Paluch lab for help and access to confocal microscopy.

Author contributions: Conceptualisation: XL, CEB. Methodology: XL, AW. Formal analysis: XL, CYH. Investigation: XL, AW. Writing: XL, CEB. Visualisation: XL. Supervision: MZG, CEB. Funding acquisition: XL, AW, MZG, CEB.

Conflict of Interests

The authors declare that they have no conflict of interest

References

Akhtar, N., and Streuli, C.H. (2013). An integrin--ILK--microtubule network orients cell polarity and lumen formation in glandular epithelium. Nat Cell Biol *15*, 17–27.

Bedzhov, I., and Zernicka-Goetz, M. (2014). Self-organizing properties of mouse pluripotent cells initiate morphogenesis upon implantation. Cell *156*, 1032–1044.

Blasky, A.J., Mangan, A., and Prekeris, R. (2015). Polarized protein transport and lumen formation during epithelial tissue morphogenesis. Annu Rev Cell Dev Biol *31*, 575–591.

Bryant, D.M., Datta, A., Rodriguez-Fraticelli, A.E., Peranen, J., Martin-Belmonte, F., and Mostov, K.E. (2010). A molecular network for de novo generation of the apical surface and lumen. Nat Cell Biol *12*, 1035–1045.

Bryant, D.M., Roignot, J., Datta, A., Overeem, A.W., Kim, M., Yu, W., Peng, X., Eastburn, D.J., Ewald, A.J., Werb, Z., et al. (2014). A molecular switch for the orientation of epithelial cell polarization. Dev Cell *31*, 171–187.

Buckley, C.E., Ren, X., Ward, L.C., Girdler, G.C., Araya, C., Green, M.J., Clark, B.S., Link, B.A., and Clarke, J.D.W. (2013). Mirror-symmetric microtubule assembly and cell interactions drive lumen formation in the zebrafish neural rod. Embo J *32*, 30–44.

Ciruna, B., Jenny, A., Lee, D., Mlodzik, M., and Schier, A.F. (2006). Planar cell polarity signalling couples cell division and morphogenesis during neurulation. Nature *439*, 220–224.

Herrera, A., Menendez, A., Torroba, B., Ochoa, A., and Pons, S. (2021). Dbnl and β-catenin promote pro-N-cadherin processing to maintain apico-basal polarity. J Cell Biol *220*, e202007055.

Iden, S., Rehder, D., August, B., Suzuki, A., Wolburg-Buchholz, K., Wolburg, H., Ohno, S., Behrens, J., Vestweber, D., and Ebnet, K. (2006). A distinct PAR complex associates physically with VE-cadherin in vertebrate endothelial cells. Embo Rep *7*, 1239–1246.

Ishida, S., Tanaka, R., Yamaguchi, N., Ogata, G., Mizutani, T., Kawabata, K., and Haga, H. (2014). Epithelial Sheet Folding Induces Lumen Formation by Madin-Darby Canine Kidney Cells in a Collagen Gel. Plos One *9*, e99655.

Jia, L., Liu, F., Hansen, S.H., Beest, M.B.A. ter, and Zegers, M.M.P. (2011). Distinct roles of cadherin-6 and E-cadherin in tubulogenesis and lumen formation. Mol Biol Cell *22*, 2031–2041.

Kim, Y.S., Fan, R., Kremer, L., Kuempel-Rink, N., Mildner, K., Zeuschner, D., Hekking, L., Stehling, M., and Bedzhov, I. (2021). Deciphering epiblast lumenogenesis reveals proamniotic cavity control of embryo growth and patterning. Sci Adv *7*, eabe1640.

Klinkert, K., Rocancourt, M., Houdusse, A., and Echard, A. (2016). Rab35 GTPase couples cell division with initiation of epithelial apico-basal polarity and lumen opening. Nat Commun *7*, 11166.

Larue, L., Antos, C., Butz, S., Huber, O., Delmas, V., Dominis, M., and Kemler, R. (1996). A role for cadherins in tissue formation. Development *122*, 3185–3194.

Li, D., Mangan, A., Cicchini, L., Margolis, B., and Prekeris, R. (2014). FIP5 phosphorylation during mitosis regulates apical trafficking and lumenogenesis. Embo Rep *15*, 428–437.

Luján, P., Varsano, G., Rubio, T., Hennrich, M.L., Sachsenheimer, T., Gálvez-Santisteban, M., Martín-Belmonte, F., Gavin, A.-C., Brügger, B., and Köhn, M. (2016). PRL-3 disrupts epithelial architecture by altering the post-mitotic midbody position. J Cell Sci *129*, 4130–4142.

Mangan, A.J., Sietsema, D.V., Li, D., Moore, J.K., Citi, S., and Prekeris, R. (2016). Cingulin and actin mediate midbody-dependent apical lumen formation during polarization of epithelial cells. Nat Commun *7*, 12426.

Molè, M.A., Weberling, A., Fässler, R., Campbell, A., Fishel, S., and Zernicka-Goetz, M. (2021). Integrin β1 coordinates survival and morphogenesis of the embryonic lineage upon implantation and pluripotency transition. Cell Reports *34*, 108834.

Morais-de-Sa, E., Mirouse, V., and Johnston, D.S. (2010). aPKC phosphorylation of Bazooka defines the apical/lateral border in Drosophila epithelial cells. Cell *141*, 509–523.

Nikolopoulou, E., Galea, G.L., Rolo, A., Greene, N.D.E., and Copp, A.J. (2017). Neural tube closure: cellular, molecular and biomechanical mechanisms. Development *144*, 552–566.

Noordstra, I., Hermoso, M.D., Schimmel, L., Bonfim-Melo, A., Kalappurakkal, J.M., Mayor, S., Gordon, E., Cusachs, P.R., and Yap, A.S. (2021). Cortical actin flow activates an α-catenin clutch to assemble adherens junctions. Biorxiv 2021.07.28.454239.

Quesada-Hernandez, E., Caneparo, L., Schneider, S., Winkler, S., Liebling, M., Fraser, S.E., and Heisenberg, C.P.
(2010). Stereotypical Cell Division Orientation Controls Neural Rod Midline Formation in Zebrafish. Curr Biol *20*,
1966–1972.

Rathbun, L.I., Colicino, E.G., Manikas, J., O'Connell, J., Krishnan, N., Reilly, N.S., Coyne, S., Erdemci-Tandogan, G.,
Garrastegui, A., Freshour, J., et al. (2020). Cytokinetic bridge triggers de novo lumen formation in vivo. Nat Commun
11, 1269.

Renschler, F.A., Bruekner, S.R., Salomon, P.L., Mukherjee, A., Kullmann, L., Schütz-Stoffregen, M.C., Henzler, C.,
Pawson, T., Krahn, M.P., and Wiesner, S. (2018). Structural basis for the interaction between the cell polarity proteins
Par3 and Par6. Science Signaling *11*, eaam9899.

Schlüter, M.A., Pfarr, C.S., Pieczynski, J., Whiteman, E.L., Hurd, T.W., Fan, S., Liu, C.-J., and Margolis, B. (2009).
Trafficking of Crumbs3 during cytokinesis is crucial for lumen formation. Mol Biol Cell *20*, 4652–4663.

Shahbazi, M.N., Scialdone, A., Skorupska, N., Weberling, A., Recher, G., Zhu, M., Jedrusik, A., Devito, L.G., Noli, L.,
Macaulay, I.C., et al. (2017). Pluripotent state transitions coordinate morphogenesis in mouse and human embryos.
Nature *552*, 239–243.

Soncin, F., Mohamet, L., Eckardt, D., Ritson, S., Eastham, A.M., Bobola, N., Russell, A., Davies, S., Kemler, R.,
Merry, C.L.R., et al. (2009). Abrogation of E-Cadherin-Mediated Cell–Cell Contact in Mouse Embryonic Stem Cells
Results in Reversible LIF-Independent Self-Renewal. Stem Cells *27*, 2069–2080.

Symonds, A.C., Buckley, C.E., Williams, C.A., and Clarke, J.D.W. (2020). Coordinated assembly and release of
adhesions builds apical junctional belts during de novo polarisation of an epithelial tube. Development *147*, dev191494.

Tawk, M., Araya, C., Lyons, D.A., Reugels, A.M., Girdler, G.C., Bayley, P.R., Hyde, D.R., Tada, M., and Clarke, J.D.
(2007). A mirror-symmetric cell division that orchestrates neuroepithelial morphogenesis. Nature *446*, 797–800.

Wang, L.-T., Rajah, A., Brown, C.M., and McCaffrey, L. (2021). CD13 orients the apical-basal polarity axis necessary
for lumen formation. Nat Commun *12*, 4697.

Wang, T., Yanger, K., Stanger, B.Z., Cassio, D., and Bi, E. (2014). Cytokinesis defines a spatial landmark for
hepatocyte polarization and apical lumen formation. J Cell Sci *127*, 2483–2492.

Yu, W., Datta, A., Leroy, P., O'Brien, L.E., Mak, G., Jou, T.S., Matlin, K.S., Mostov, K.E., and Zegers, M.M.
(2005). Beta1-integrin orients epithelial polarity via Rac1 and laminin. Mol Biol Cell *16*, 433–445.

Zhang, Y., Mets, R.D., Monzel, C., Acharya, V., Toh, P., Chin, J.F.L., Hul, N.V., Ng, I.C., Yu, H., Ng, S.S., et al.
(2020). Biomimetic niches reveal the minimal cues to trigger apical lumen formation in single hepatocytes. Nat Mater
19, 1026–1035.

Zigman, M., le, A.T., Fraser, S.E., and Moens, C.B. (2011). Zebrafish neural tube morphogenesis requires scribble-
dependent oriented cell divisions. Curr Biol *21*, 79–86.

Figures

Figure 1

Figure 1. Cell division is dispensable for mouse embryonic stem cells (mESC) in Matrigel 3D cultures.

(A) Stages of polarisation and lumen formation in mESCs cultured in Matrigel

(B) Timeline of experiment setups to assess AMIS formation.

(C) Immunofluorescence of PAR3. A polarised, concentrated PAR3 centre was formed in 2-cell mESC doublets and 3- or 4-cell mESC in both control and Mitomycin treated conditions.

(D) Quantification of the frequency of cell clusters with a polarised Pard3 centre. N=3 experiments, at least 25 clusters were analysed for each column in every experiment. Data are means ± SEM, one-way ANOVA analysis; n.s., not significant.

(E) Line-scans of Pard3 at the cell-cell interface of 2-cell mESC doublets. Line scans were sections and fitted to each 5% along the cell-cell interface length. 12 lines scans from one experiment were and profiled for each condition.

(F) An example and illustration of the line-scan analysis in E. The line-scan profiles were from the yellow arrow to the orange arrow.

(G) Representative images of centralised ZO-1 puncta and polarised Golgi apparatus (labelled by GM130).
(H) Representative images of polarised centrosomes, labelled with γ-tubulin.
(I) Quantification of the frequency of cell doublets with a polarised ZO-1 or Golgi apparatus centre. N=3 experiments, at least 25 cell doublets were analysed for each column in every experiment. Data are means ± SEM, student's t-test analysis; n.s., not significant.
(J) Distance between centrosomes in cell doublets. The distance was normalised to the length of the long axis of the doubles N = 35 doublets in each condition. Data with the median values, student's t-test analysis; n.s., not significant.

Figure 2

Figure 2. Polarised Pard6B in mESC 3D cultures
(A) Live cell imaging of mCherry-PAR6B mESC control cells cultured from 24 – 96 hours in Matrigel.
(B) Representative live cell images of mCherry-PAR6B in control and Mitomycin division-blocked mESCs upon 24 hours in Matrigel.
(C) Quantification of the frequency of cell clusters with polarised mCherry-Pard6B. N=3 experiments, at least 25 cell clusters were analysed for each column in every experiment. Data are means ± SEM, student's t-test analysis; n.s., not significant.
(D) Movie stills of mCherry-PAR6B in control and Mitomycin C treated mESCs cultured in Matrigel (also Supplementary Movie 2). Control cells divided (i) and two mitomycin treated cells touched (ii) to form 2-cell doublets. Control cells divided twice (iii) and two mitomycin treated cell clusters touched (iv) to form 4-cell clusters. (v), Kymograph of mCherry-PAR6B in the arrow-head cell in (ii) along the path between the arrows; each pixel is the fluoresce values averaged in 0.2 μm sections.

Figure 3. E-cadherin junctions are important for polarisation.

(A) Immunofluorescence of PAR3 in wild-type and E-cadherin knock-out (*Cdh1* KO) cells at 24 hrs in Matrigel.

(B) Proportions of mESC clusters with a positive Pard3 centre. N=3 experiments, at least 25 clusters were analysed for each column in every experiment. Data are means ± SEM, one-way ANOVA; n.s., not significant, ****, P<0.0001.

(C) Line-scan profiles of PAR3 at the cell-cell interface in wild-type control, mitomycin C treated and Cdh1 KO control, mitomycin C treated 2-cell doublets. N = 12 line scans in each condition.

(D) Representative images of PAR3 immunofluorescence in WT homogeneous or WT/*Cdh1* KO chimeric mESC 2-cell doublets.

(E) Line-scan profiles of PAR3 at the cell-cell interface of homogeneous and chimeric mESC doublets. N = 11 doublets in each condition.

(F) Representative images of ZO-1 puncta and Golgi apparatus (labelled by GM130) in WT and Cdh1 KO mESC doublets.

(G) Proportions of mESC doublets with central ZO1 puncta or polarised Golgi apparatus. N=3 experiments, at least 25 doublets were analysed for each column in every experiment. Data are means ± SEM, students' t test; n.s., not significant.

(H) Distance between centrosomes in cell doublets. The distance was normalised to the length of the long axis of the doubles. N = 35-40 doublets in each condition. Data with the median values, one-way ANOVA analysis; n.s., not significant.

(I) Fluorescence recovery after photobleaching (FRAP) of GFP-E-cadherin at the centre-most or side 2 μm regions of division-blocked mESC doublets' cell-cell interfaces.

(J) Mobile fraction of GFP-E-cadherin calculated from the plotting of I. N = 15 doublets. Data with the median values ± SD, students' t test; **, P<0.01.

Figure 4

Figure 4. Cell-ECM interactions in regulating AMIS seeding.

(A) Immunofluorescence of PAR3 in wild-type and E-cadherin knock-out (*Cdh1* KO) cells at 30 hours in Matrigel.

(B) Proportions of mESC doublets with polarised PAR3. N=2 experiments, at least 15 doublets were analysed for each column in every experiment. Data are means ± SEM, one-way ANOVA; n.s., not significant, *, P<0.05.

(C) Immunofluorescence of PAR-3 in cell division blocked mESCs cultured against E-cadherin-FC or Fibronectin coated glass topped with or without Matrigel for 24 hours.

(D) Heatmap of PAR3 in cells from (C). Squared frames were fitted to the main bodies of the cells and average values in each 10% along the width of the frames was calculated as a pixel. The heatmaps were stacks of 15 cells in one experiment. The values were normalised to the peak value in the E-cad-FC plus Matrigel condition.

(E) Ratios between PAR3 in the 2.5 *µ*m diameter central core and whole cell surface. N = 15 cells in one experiment. Data are means ± SEM, one-way ANOVA; n.s., not significant, **, P<0.01, ***, P<0.001.

Figure 5

Figure 5. Lumenogenesis in wild-type and E-cadherin knock-out mESC cultures.

(A) PAR3 immunofluorescence in wild-type and E-cadherin knock-out (*Cdh1* KO) mESCs cultured in Matrigel for 48 and 72 hours. At 48 hours, the majority of WT cell clusters had formed a polarised rosette or small lumen (i). However, the majority of *Cdh1* KO cells had not formed a central AMIS (ii). A small percentage of *Cdh1* KO cells had formed an open 'cup-shape', with apically localised PAR3 (iii). By 72 hours, a significant proportion of *Cdh1* KO cells had formed polarised 'cavity-like' structures, about half of which were configured in an open 'cup-shape' and half as closed 'lumen-like' structures (iv). The percentages were quantified from one experiment.

(B) Podocalyxin (PodxI) immunofluorescence in WT and *Cdh1* KO mESCs cultured in Matrigel from 1-4 days.

(C) Examples of masked *Cdh1* KO cell cluster surfaces from 'open' and 'closed' cavities. The cavities were categorised based on Podocalyxin signals.

(D) Percentage of cell clusters with different cavities relative to total cell clusters in different time points. N=3 experiments, at least 25 clusters were analysed for each column in every experiment. Data are means ± SEM, one-way ANOVA analysis.

(E) Percentage of cell clusters with closed cavities relative to total cell clusters with cavities calculated from (D).

(F) Movies stills of LifeAct-mRuby labelled cell clusters. Images are overlaid with 3D rendering of the cluster surfaces. The images are 50 μm-depth z-projections.

(G) Illustration of lumenogenesis in WT and Cdh1 KO mESCs.

Supplementary Figures

Figure S1

Figure S1. Division blocked mouse embryonic stem cells (mESC) formed E-cadherin cell-cell contacts in Matrigel 3d cultures.

(A) Mitomycin C sufficiently blocked cells divisions. Track of cells experienced cell divisions or did not experience cell divisions for live movies in the first 24 hours upon seeding into Matrigel by live cell imaging. N equals to total numbers of cells tracked at time point zero.

(B) Movie stills of Mitomycin treated cells from Supplemental Movie 1.

(C) Quantification of E-cadherin fluorescence intensity at cell-cell interfaces and cell-matrix interfaces. N = 18 cell clusters in each condition. Data are means ± SEM, student's t-test; ***, P<0.001.

(D) Representative images of mESCs that were negative of PAR3 centres.

(E) Split-channel images of Figure 1C.

(F) Split-channel images of Figure 1G, H.

(G) Aphidicolin sufficiently blocked cells divisions. N equals to total numbers of cells tracked at time point zero.

(H) Immunofluorescence of PAR3 in control and aphidicolin treated cells cultured for 24 hours in Matrigel.

(I) ZO-1 and Golgi apparatus (labelled by GM130) in control and aphidicolin treated cells. The percentage of polarised cells was quantified from one experiment of over 25 cell clusters.

Figure S2

Figure S2. Wild-type, E-cadherin knock-down and knock-out cell clusters.

(A) Both wild-type and E-cadherin knock-out (*Cdh1* KO) mESCs form cell clusters when untreated or after Mitomycin treatment.

(B) Immunofluorescence of P-cadherin in wild-type and *Cdh1* KO mESCs cultured 24 hours in Matrigel.

(C) Immunofluorescence of E-cadherin in wild-type and E-cadherin siRNA knock-down mESCs cultured for 24 hours in Matrigel.

(D) Line-scans of Pard3 at the cell-cell interface of 2-cell mESC doublets. Line scans were sections and fitted to each 5% along the cell-cell interface length. N = 14 lines scans for each condition.

(E) Representative images of centrosomes in wild-type and E-cadherin KO mESCs cultured 24 hours in Matrigel with or without cell divisions.

Figure S3

Region of photo-bleaching Region of analysis

Figure S3. E-cadherin in the centre-most and side regions at two-cell cluster interfaces.
(A)   An example line-scan at the centre-most (blue) and side (red) regions at division blocked two-cell cluster interfaces. The width of line-scans was 3 μm. Two side regions were line-scanned and the average was taken as the line-scan profile at the side regions for a two-cell cluster (See panel B).
(B)   Line-scan profiles of E-cadherin at the centre-most and side regions. The statistical comparison was done between the area under the curves. N = 15 line-scans. Data are means ± SD, student's t-test analysis; ns, not significant.
(C)   Illustrations of E-cadherin-GFP FRAP at a two-cell cluster interface.
(D)   E-cadherin-GFP levels at the photo bleaching regions before bleaching in division blocked cells. N = 15 two-cell clusters. Data are means ± SD, student's t-test analysis; ns, not significant.

Figure S4. AMIS seeding and pluripotency exit in wild-type and E-cadherin knock-out mESCs.

(A)   Timeline of experiment setups to assess *de novo* polarisation when mESCs were cultured with 2i/Lif to remain pluripotent.

(B)   Immunofluorescence of Pard3 in wild-type and E-cadherin knock-out (KO) mESCs cultured in Matrigel for 24 hours with 2i/Lif.

(C)   Quantification of the frequency of cell clusters with a polarised Pard3 centre. N=3 experiments, at least 25 clusters were analysed for each column in every experiment. Data are means ± SEM, one-way ANOVA analysis; ns, not significant, ****, P<0.0001.

(D)   Nuclear levels of Otx2 and Nanog based on immunofluorescence in wild-type and E-cadherin KO mESCs at 12 or 24 hours post seeding into Matrigel. Only cells in interphase were analysed. N = 22-46 cells in each column from one experiment. Data are means ± SD, one-way ANOVA analysis. ns, not significant, *, P<0.05, **, P<0.01, ***, P<0.001, ****, P<0.0001.

Figure S5

A

B

Figure S5. Wild-type and E-cadherin knock-out mESC cultures.
(A)   ZO-1 immunofluorescence in WT and *Cdh1* KO mESCs cultured in Matrigel from 1-4 days.
(B)   The Golgi network, phospho-myosin light chain 2 and F-actin in E-cadherin knock-out cells cultured for 72 hours in Matrigel.

Supplementary Movies
Supplementary Movie 1. Dividing and division-blocked mESCs cultured in Matrigel.
Bright-field live movies of left, control and right Mitomycin C treated mESCs cultures in Matrigel from 0 – 24 hours. The control cells divided once or two times while the treated cells had no divisions.

Supplementary Movie 2. Representative movies of mCherry-Pard6B in dividing and division-blocked mESCs cultured in Matrigel
Top: control and Mitomycin division-blocked mESCs cultured in Matrigel formed 2-cell doublets from 6 – 18 hours in Matrigel. Bottom: control and Mitomycin division-blocked mESCs cultured in Matrigel formed multi-cell clusters from 9 – 19 hours in Matrigel.

Supplementary Movie 3. Representative movies of cysts forming in wild-type and E-cadherin knock-out mESCs cultured in Matrigel
The central 5µm section of wild-type and E-cadherin knock-out (*Cdh1* KO) mESCs cultured in Matrigel. Left, wild-type mESCs cultured from 74 – 84 hours, interval = 1 hour; Right, *Cdh1* KO mESCs cultured from 78 – 85 hours, interval = 30 min. The cells expressed LifeAct-mRuby.

Supplementary Movie 4. Rotation of the 3D rendering in Figure 5F.
Left, Cdh1KO mESCs cultured 78 hours; Middle, 81.5 hours; Right, 85 hours in Matrigel. Central 5 µm of mESC cultures. Scale bar = 20 µm.