Functional analysis of the polar amino acid in the TatA transmembrane helix

Binhan Hao, Wenjie Zhou, Steven M. Theg

Plant Biology Department, University of California, Davis, CA 95616

November 30, 2021

For correspondence e-mail: smtheg@ucdavis.edu

Abstract

The twin-arginine translocation (Tat) pathway utilizes the proton-motive force (pmf) to transport folded proteins across cytoplasmic membranes in bacteria and archaea, as well as across the thylakoid membrane in plants and the inner membrane in mitochondria. In most species, the minimal components required for Tat activity consist of three subunits, TatA, TatB, and TatC. Previous studies have shown that a polar amino acid is present at the N-terminus of the TatA transmembrane helix (TMH) across many different species. In order to systematically assess the functional importance of this polar amino acid in the TatA TMH in *Escherichia coli*, a complete set of 19-amino-acid substitutions was examined. Unexpectedly, although being preferred overall, our experiments suggest that the polar amino acid is not necessary for a functional TatA. Hydrophobicity and helix stabilizing properties of this polar amino acid were found to be highly correlated with the Tat activity. Specifically, change in charge status of the amino acid side chain due to pH resulted in a shift in hydrophobicity, which was demonstrated to impact the Tat transport activity. Furthermore, a four-residue motif at the N-terminus of the TatA TMH was identified by sequence alignment. Using a biochemical approach, the N-terminal motif was found to be functionally significant, with evidence indicating a potential role in the preference for utilizing different pmf components. Taken together, these findings yield new insights into the functionality of TatA and its potential role in the Tat transport mechanism.

Introduction

The twin-arginine translocation (Tat) pathway, a protein transport machinery, is found

in bacteria, archaebacteria, chloroplasts, and plant mitochondria. In bacteria, the Tat

pathway is involved in many critical biological processes, including cell division,

stress tolerance, and electron transport (Berks, 2015; Gohlke et al., 2005; Ize et al.,

2003). In plants, the Tat pathway transports several proteins that are essential for

photosynthesis across thylakoid membranes (Clark and Theg, 1997). In haloarchaea

which normally possess high cytoplasmic salt concentration leading to faster protein

folding, the Tat pathway serves almost 50% of secretome (Ghosh et al., 2019). Unlike

the similarly ubiquitous Sec pathway, the Tat pathway is able to transport folded

proteins (Clark and Theg, 1997) while requiring only the proton motive force (pmf),

with no contribution from NTP hydrolysis (Braun et al., 2007).

The Tat system in most species consists of three subunits, TatA, TatB, and TatC

(Berks, 2015). It has been shown that all three subunits can form a complex which

works as a receptor for signal peptides of Tat substrates. After the receptor complex

binds to the Tat substrate, more TatA subunits are recruited to the translocon. While

TatB and TatC are present in a 1:1 stoichiometry in the complex, TatA is present at

much higher stoichiometries (8-40 folds) (Leake et al., 2008; Mori et al., 2001;

Sargent et al., 2001).

Functional analysis of the polar amino acid in TatA TMH

The structures of the three individual Tat subunits have been determined (Hu et al.,

2010; Pettersson et al., 2018; Rollauer et al., 2012; Zhang et al., 2014). Both TatA and

TatB possess a short transmembrane helix (TMH) followed by a hinge region, one or

more amphipathic helices (APH), and then a relatively unstructured C-terminus.

Conversely, TatC has six TMHs and exhibits a cupped hand shape. Remarkably, the

Tat machinery transports folded proteins of different sizes without introducing

significant ion leakage (Asher and Theg, 2021; Teter and Theg, 1998). Recent work

suggests that this occurs in the absence of channel/pore like structure, unlike the

well-characterized Sec translocon (Tsirigotaki et al., 2017). It has been proposed that

Tat pathway operates through lipid-lined toroidal pores formed in a destabilized

membrane, rather than common proteinaceous pore (Asher and Theg, 2021; Brüser

and Sanders, 2003; Hao et al., 2021).

Previous studies have demonstrated that TatAs have a highly conserved short TMH

(Hao et al., 2021) which could cause membrane destabilization and is critical to the

overall Tat function (Hao et al., 2021; Hou et al., 2018). Such short TMHs make the

TatA protein energetically unfavorable to span on the membrane bilayer. Furthermore,

it has been shown that almost all TatA family proteins contain a polar residue at the

N-terminus of its TMH (Hicks et al., 2003). In *Escherichia coli* (*E. coli*), TatA has a

glutamine (Q) at the 8th position (TMH: 6th – 20th), while in pea chloroplasts, TatA has

a glutamic acid (E) instead at the corresponding position. A cysteine-scanning study in

E. coli TatA illustrated that cysteine substitution for the 8[th] glutamine (Q8C mutant)

led to the complete loss of Tat activity, whereas cysteine substitutions for all the other

residues in the TMH retained Tat transport activity similar to that of the wtTatA

(Greene et al., 2007). Further studies showed that *E. coli* TatA asparagine substitution

(Q8N) and Q8E mutants retained the ability to export TMAO (trimethylamine

N-oxide) reductase (TorA) to the periplasm (Greene et al., 2007), while the alanine

(Q8A) mutant was not able to transport either TorA (Greene et al., 2007) or CueO

(Alcock et al., 2016). However, another study showed a somewhat conflicting result

that *E. coli* Q8A and Q8C mutants retained TorA transport capability when

overexpressed (Warren et al., 2009). On the other hand, in pea or maize chloroplasts,

it was shown that all of the 10[th] glutamic acid substitution variants, E10A, E10D, and

E10Q, severely affected Tat transport in a biochemical complementation assay

(Dabney-Smith et al., 2003; Fincher et al., 2003). Interestingly, although the

chloroplast TatA E10Q mutant exhibited extremely low Tat activity, the *E. coli* TatA

Q8E mutant displayed high Tat transport activity, suggesting that residue preference

was not identical for different species. Such different preference was also shown with

chimeric pea/*E. coli* TatA derivatives (Hauer et al., 2017). Furthermore, the *Bacillus*

subtilis Tat system contains two sets of TatA/TatC subunits (TatAd/TatCd and

TatAy/TatCy), and those TatAs actually lack this "conserved" polar residue in the

TMH. Although extensive effort has been made to investigate the function of this

"conserved" polar amino acid in TatA, the lack of a complete set of mutants and use

of different transport assays has allowed the role of this residue at the analogous

position to remain an open question.

In this study, we analyzed the amino acid requirement for the 8th residue in *E. coli*

TatA through a complete set of Q8 mutants. Both qualitative and quantitative assays

were conducted to compare the activity between TatA mutants. Unexpectedly, we

found that polar amino acids were preferred overall but actually not necessary for a

functional TatA. However, the Tat activity was highly correlated with hydrophobicity

and helix stability of the 8th residue in TatA. Furthermore, by sequence alignment and

biochemical analysis, a four-residue motif at the N-terminus of the TatA TMH was

found to be functionally significant. These results expand our understanding of the

potential function of TatA and provide new insight into the Tat transport mechanism.

Results

1. Polar amino acids at the 8th position in *E. coli* TatA TMH are preferred but not necessary for minimal Tat activity.

To systematically assess the characteristic requirements of the 8th residue in *E. coli* TatA, Q8 was replaced by all the other 19 amino acids and their effects on Tat activity was measured. Membrane stability of TatA Q8 mutants was assessed by membrane isolation followed by carbonate wash, and all mutants displayed an integrated presence in the membrane (Supplemental figure 1). Minimal Tat activity was subsequently screened by SDS growth assays (Ize et al., 2003) for each of the TatA Q8 mutants. All polar amino acid substitutions (Q8N, Q8S, Q8T, Q8D, Q8E, Q8H, Q8K, and Q8R) exhibited sufficient Tat activity to grow in LB medium containing 10% SDS (Figure 1A). Surprisingly, TatA Q8A, Q8C, Q8G, and Q8P mutants, whose substituted amino acids do not possess strong polar side chains, also showed a growth activity in the same medium (Figure 1B, C). The rest of substitutions (Q8F, Q8I, Q8L, Q8M, Q8V, Q8W, and Q8Y) displayed no growth activity in LB medium containing 5% SDS, indicating a complete loss of Tat activity (Figure 1C). An *in vivo* transport assay was implemented to further confirm the SDS growth assay result. In this assay, a FLAG-tagged *E. coli* native Tat substrate, SufI-FLAG, was overexpressed in the TatA Q8 mutants. After 2.5 hours transport at 37°C, a periplasmic fraction was isolated by treated with EDTA/lysozyme and osmatic shock (Petiti et al., 2017a), a harsh method

that minimizes the loss of periplasmic proteins. The precursor and mature forms of

SufI-FLAG were separated by SDS-PAGE according to their size. This assay revealed

that, despite their ability to grow in 10% SDS, the Q8A, Q8C, Q8D, Q8E, Q8G, Q8P,

Q8R, and Q8T mutants displayed significantly lower Tat activity (<25%) compared to

wild-type TatA (Figure 1D).

Pulse-chase assays, representing a more fine-grained assessment of Tat transport

activity (Palmer et al., 2010), were conducted with all the functional TatA Q8 mutants

to compare their transport rates. Figure 2A shows the quantitative analysis of the pulse

chase results (Supplemental figure 2). An exponential plateau model $y = y_{\max} -$

$(y_{\max} - y_0) \times e^{-kt}$ was applied in data fitting. Estimated initial transport rate (V_0)

derived from the fitting curve was used to compare the transport activity between the

TatA Q8 mutants (Figure 2B). Unexpectedly, the Q8H mutant exhibited the highest

V_0 (228% of the V_0[wtTatA]). Q8N, Q8S, and Q8E variants of TatA, exhibited 58%,

37% and 33% of the V_0[wtTatA], respectively. However, Q8A, Q8C, Q8D, Q8G,

Q8K, Q8P and Q8R showed relatively lower transport rate (less than 25% of

V_0[wtTatA]). One unanticipated finding was that the Q8T, Q8D and Q8R mutants,

whose 8[th] glutamine was replaced by a polar amino acid, showed even lower transport

rates than the poorly performing Q8A mutant. The results from pulse-chase assays

were overall consistent with the *in vivo* transport assay except that Q8K performed

relatively worse in the pulse-chase assay than in the *in vivo* transport assay.

In summary, polar amino acids are overall strongly favored at the 8^{th} position in the *E.*

coli TatA but not necessary for a functional Tat translocon. Moreover, not all polar

amino acid substitutions showed higher transport ability than non-polar substitutions.

2. The Tat activity of Q8 mutants is correlated with the

side chain hydrophobicity and helix stabilizing properties

of the substituted amino acid.

Our data clearly show that polar amino acids are preferred overall in the TatA 8^{th}

position, but are not strictly necessary to retain Tat activity (ex. Q8A, Q8C, Q8G,

Q8P). Moreover, some of the polar residue substitutions (i.e., Q8D, Q8R and Q8T),

even though they retained the minimal Tat activity required to grow in LB medium

with SDS, displayed significantly lowered transport activity. Such results could not

simply be explained by the binary categories, polar or non-polar, of amino acids. In

order to understand the functional importance of this position in the *E. coli* TatA, a

further analysis of the amino acids' preference is needed. On one hand, previous

studies showed that the overall hydrophobicity of the TatA TMH might affect TatA

function in thylakoids (Hauer et al., 2017). On the other, a recent study has

demonstrated that the hydrophobic mismatch between the TatA TMH and membrane

bilayer is critical to Tat function (Hao et al., 2021; Mehner-Breitfeld et al., 2021). To

maintain such a hydrophobic mismatch, a relatively stable alpha helix structure is

likely to be required. To assess these properties, the hydrophobicity and α-helix

stability of each amino acid (Monera et al., 1995) were plotted together with the

transport performance of the corresponding TatA Q8 mutants (Figure 1E).

Interestingly, a high correlation between Tat activity and the combination of

hydrophobicity and α-helix stability was observed: 1) the Q8 mutants which

completely lost Tat activity clustered at the top-right corner, suggesting that very high

hydrophobicity and helix stability at this position are detrimental for Tat transport. 2)

all mutants with high Tat activity were clustered in the middle ranges of

hydrophobicity and helix stability (Figure 1E, pink box). These observations draw our

attention to the importance of a moderate hydrophobicity and α-helix stability of the

8[th] amino acid in *E. coli* TatA.

3. Charge status of the side chain of the polar amino acid

affects the TatA function.

Due to the nature of polar amino acids with electrically charged side chains, previous

studies have reported that the hydrophobicity of those amino acids changes

dramatically under different pH conditions (Kovacs et al., 2006; Mant and Hodges,

2002). For instance, the side chain of histidine ($pKa_{(His)}$ = 6.04) is relatively

hydrophobic (similar as Ala) at pH 7 (mostly uncharged), while it becomes one of the

most hydrophilic side chains when the pH is lower than 5 (mostly charged). In this

study, we hypothesized that a moderately hydrophobic side chain is required for the

8[th] residue in the *E. coli* TatA. To further investigate this hypothesis, a comparison of

Tat transport rates between wtTatA and Q8H was studied under different pH

conditions. The pulse-chase experiments in Figure 2 showed that the Q8H mutant

performed approximately 2-fold better than the wild-type TatA at pH = 7.0 where

most of (90%) the histidines in TatAs were presumably deprotonated. To assess

whether a change in histidine hydrophobicity would alter activity of the Q8H mutant,

we measured transport rates by pulse-chase at pH 5.0, 6.0 and 8.0 (Figure 3A). Figure

3 shows that wtTatA and Q8H had different behaviors when transporting SufI at

different pH. First, wtTatA and Q8H mutant exhibited optimal transport rates at

different pH values (Figure 3B and C). wtTatA displayed optimal transport activity at

pH 5.0, while the optimal pH for Q8H mutant was 7.0. However, their highest V_0

values were comparable to each other (Figure 3D). Second, V_0 [wtTatA] was

significantly higher than V_0[Q8H] at pH = 5.0 (Figure 3E) whilst V_0[wtTatA] was

significantly lower than V_0[Q8H] at pH = 7.0 (Figure 3G). Third, wtTatA had a very

similar V_0 as Q8H mutant at pH = 8.0 (Figure 3H) where 99% of the histidines were

presumably deprotonated, a similar status as uncharged glutamine in wtTatA. In

summary, the charged status of the histidine side chain, a model residue for

comparison with the wtTatA, affects its transport.

In the aggregate, our experiments in the pH range 5.0-8.0 are consistent with the

hypothesis that a functional Tat translocon requires that its TatA possesses a

moderately hydrophobic amino acid at 8th position. In addition, it demonstrates that

not only hydrophobicity of different amino acid but also hydrophobicity of same

amino acids under different charged status affects its ability to support Tat transport.

Together these results provide important insights into understanding the function of

the 8th residue in *E. coli* TatA (see discussion).

To further investigate how charged residues in TatA affect the overall Tat transport

machinery, we determined how well the membrane can hold a pmf by measuring the

ΔpH induced by ATP hydrolysis in inverted membrane vesicles (IMVs) membrane

expressing corresponding mutated TatA proteins. According to their charge status at

pH 7.0, Q8E, wtTatA, Q8H and Q8K mutants were selected as examples of negatively,

neutral, partially positively and completely positively charged residues at the 8th

position in *E. coli* TatA (Figure 4A). wtTatA and the Q8E mutant showed similar low

membrane leakage outcomes in the presence or absence of the TatBC complex.

However, Q8H and Q8K mutants exhibited significant different membrane leakage

characteristics when the TatBC complex was present. Q8H, which showed the highest

membrane leakage without TatBC, displayed less membrane leakage when TatBC was

also expressed on the membrane. In contrast, the Q8K mutant showed the opposite

behavior, exhibiting the highest ΔpH without TatBC but developing a dramatically

high membrane leak when TatBC was present. That is to say, even though lysine and

glutamic acid have similar hydrophobicity at pH 7.0 (Figure 1E), their different

charge status might affect how they interact with the TatBC complex, which results in

different membrane leakage outcomes. The cause for this behavior is not clear and

awaits further investigation.

4.   Different species have different preferences for polar

amino acids in the TatA TMH.

Even though in *E. coli* TatA, the polar amino acid in the TMH is the uncharged

glutamine, in plant, the Tat system, this position is usually occupied by a glutamic

acid, a charged amino acid (Mori and Cline, 2001). Moreover, *Bacillus subtilis*, a

gram-positive bacterium, expresses two different kinds of TatA with glycine (TatAc)

or serine (TatAy) at the corresponding position. To investigate if such amino acid

choices are correlated to species, TatA sequences were aligned according to five

categories: proteobacteria (N=7534), actinobacteria and firmicutes (N=6978),

chloroplasts (N=242), cyanobacteria (N=149) and archaea (N=151). A conserved

12-residue hydrophobic core was observed (9^{th} – 20^{th} in Figure 5A-E) before the

invariant "FG" motif at the end of TMH (Hao et al., 2021). In addition, there is a clear

Functional analysis of the polar amino acid in TatA TMH

preference for particular amino acids in the TatA TMH at the 8th position with

different species (Figure 5F) (Here, we use the numbering from *E. coli* for simplicity.

The exact number might be different in different species.) In proteobacteria

(representing *E. coli*), histidine (50%) and glutamine (34%) are the top two choices.

Interestingly, such choice preferences are consistent with our Tat activity results in

that Q8H and wtTatA exhibited the highest Tat transport rates. In contrast, in the other

two groups of bacteria, actinobacteria and firmicutes, even though histidine is still the

top choice for TatA (46%), glutamine is present in only 2% of the species in these

categories, and relatively more of them choose glutamic acid (32%) or glycine (10%)

compared to proteobacteria. However, all of chloroplasts, cyanobacteria and archaea

TatAs have a highly conserved glutamic acid (>90%) at the 8th position. Remarkably,

even though polar amino acids are present in TatA in most species, 1% of

actinobacteria and firmicutes (N = 80) and 0.1% of proteobacteria (N = 13) possesses

an alanine at that position. This finding further confirms our experimental

observations that polar residues at 8th position are not indispensable for a functional

TatA (Supplemental Figure 3A and B). Such different preferences in different species

suggest that a possible evolutionary pressure exists at this position for the Tat pathway

to accommodate various external conditions.

5. A four-residues functional motif at the N-terminus of the TatA TMH is important for Tat activity.

Besides the various preference of amino acids at the 8^{th} position in different species, a

conserved GxPE motif (the 5^{th} to 8^{th} position in Figure 5C, D and E) was also

observed in chloroplasts, cyanobacteria, and archaea TatAs. In contrast, in

proteobacteria, actinobacteria and firmicutes, such "GxPE" motif was not obviously

presented. Instead, a SIW(H/Q) motif was generally conserved in proteobacteria. To

see if these motifs are correlated to different polar amino acids, TatA sequences were

further examined based on the identity of the amino acid at the 8^{th} position

(Supplemental Figure 3). Supplemental Figure 3A clearly shows that proteobacteria

favor "SIWH" and "SIWQ" (i.e., *E. coli*) at the N-terminus of the TatA TMH. In

addition, "GLPG", "SITK" and "S(I/L)WN" motifs were identified in proteobacteria

as well. Interestingly, in *E. coli* TatE, "SITK" was at the corresponding location (the

5^{th} to 8^{th} residues). It has been reported that the TatE gene was found only in

enterobacteria, a subset of proteobacteria (Yen et al., 2002). Accordingly, in this study,

TatA or TatE sequences from enterobacteria were also aligned (Supplemental Figure

3F) to check if such motifs are also present in other TatE genes. Surprisingly, "SIWQ"

and "SITK" were extremely conserved across TatA (N=599) and TatE (N=160),

respectively, in enterobacteria. Additionally, although "SIWH" was the most

conserved motif in proteobacteria, gram-positive bacteria favored "xPWH" in the

same position (Supplemental Figure 3B), These results suggest that the four-residues

motif at the N-terminus of the TatA TMH might play an important role in TatA

function despite minor deviations in different species.

To assess whether the corresponding four-residue motifs are required for specific

amino acids at the 8th position to exhibit higher Tat transport activity, we also

substituted the 5th to 8th amino acids in *E. coli* TatA according to the above conserved

motifs, including "GVPE" (a motif of chloroplasts TatAs), "GLPE" (a motif of

cyanobacteria TatAs) and "SITK" (a motif of enterobacteria TatEs). Moreover, to

comprehensively understand how the 5th to 7th amino acids affects TatA function, the

5th to 8th residues of *E. coli* TatA were also changed to "GVPQ" or "GLPQ" (Figure

6).

A recent study has shown that a shorter TMH negatively affects TatA's membrane

stability (Hao et al., 2021). In terms of our new mutants, GVP(E/Q) and GLP(E/Q), it

is possible that those mutants lost their ability to stably embed into the membrane

since proline (immediately before the 8th residue) usually acts as a helix breaker

(Chou and Fasman, 1978) in protein secondary structure. Figure 6A shows that TatA

in GVP(E/Q) and GLP(E/Q) mutants exhibited much lower membrane abundance

compared to wtTatA or Q8E mutant, indicating that a proline residue severely affects

the membrane insertion, even though it did not cause high membrane leakage in those

mutants (Figure 4B and C). But, very surprisingly, GVPE mutant displayed a better

performance than Q8E in the *in vivo* transport assay (Figure 6B). Pulse-chase assays

with those N-motif mutants were also conducted to quantitatively compare their

transport rates. Both GLPQ and GVPQ displayed somewhat lower transport rates than

wtTatA, suggesting that a disruption of the conserved "SIWQ" motif was deleterious

to *E. coli* TatA function (Figure 6C). However, GVPE and SITK mutants exhibited

significantly higher transport rates than Q8E and Q8K mutants, respectively, which

indicates that the evolutional conserved motifs are functionally more suitable for

certain amino acids at the 8^{th} position in TatA (Figure 6D-F). In spite of that replacing

to conserved N-motifs were generally beneficial for Tat function, we also observed

that the GLPE mutant supported significantly lower transport rates than both the

GVPE and Q8E mutants. Such results demonstrate that changes of a single amino acid

in the N-terminal motif (even from one hydrophobic residue to another hydrophobic

residue) strongly affected the final TatA function. This confirms our hypothesis that

the N-terminus of TatA, and not just the well-known polar amino acid, determines the

Tat transport activity. Further, it implies that the combinations of these amino acids at

the N-terminus of the TatA TMH are potentially tuned to their own membrane

environments, and those different motifs could be a strategy to adjust to diverse and

subtle biological membrane properties.

6. The conserved "GVPE" motif from chloroplast TatA

might be responsible for preference of utilization of ΔpH.

Even though the GVPE mutant performed significantly better than the Q8E mutant

(Figure 6F), it was still less effective than the wild-type SIWQ motif in the *E. coli*

TatA.

This result does not point to an explanation of why most chloroplasts and

cyanobacteria possess a glutamic acid at the 8^{th} position in their TatAs. It was

previously shown that the chloroplast Tat system is highly dependent on ΔpH (Cline

et al., 1992), while *E. coli* Tat system does not require ΔpH for active transport

(Bageshwar and Musser, 2007a). These energetic preferences are also consistent with

their own respective tendencies to partition their pmfs between a ΔpH and a Δψ (Cruz

et al., 2001; Zilberstein et al., 1984). To investigate if the invariant motif in

chloroplast TatA is related to this energetic preference, IMVs, the bacterial

counterpart to chloroplast thylakoids, were made from wtTatA, Q8E and GVPE

mutants. A previous study (Bageshwar and Musser, 2007a) used IMVs to test Tat

transport ability powered by different components of the pmf. In our hands also, the

IMVs made with wild-type *E. coli* Tat displayed a higher transport ability in the

presence of nigericin which presumably dissipated the ΔpH. Figure 7A shows that

wtTatA and the Q8E mutant displayed consistently higher performance in the

presence of nigericin. However, the GVPE mutant exhibited significantly lower

transport capability when the ΔpH was dissipated. These results suggest that GVPE

mutant relied more on ΔpH for transport than the wtTatA or the Q8E mutant.

Discussion

An incomplete set of *E. coli* TatA Q8 mutants was a limitation to a full understanding

of the function of the "conserved" polar amino acid in the TatA TMH. In this study,

we systematically analyzed the Tat transport ability of all the Q8 mutants which the

8^{th} glutamine of *E. coli* TatA was substituted by the other 19 amino acids. Three

assays were conducted to compare the transport rates between different mutants,

qualitatively and quantitatively. The transport ability results of our TatA Q8 mutants

showed that a polar amino acid at the 8^{th} position of TatA is preferred but not

necessary for a functional TatA (Figure 1 and 2). Rather, a strong correlation between

Tat transport activity and the hydrophobicity and helix stability of the 8^{th} residue was

noticed (Figure 1E). It reveals that a residue possessing a moderate range of

hydrophobicity and helix stability was preferred for TatA to retain high transport

capability. Moreover, by altering the environmental pH in our transport assay, we

demonstrated that the Q8H mutant, which contains a protonatable amino acid

substitution, exhibited significantly lower transport ability under conditions in which

this residue was charged, that is, where the hydrophobicity of the 8^{th} residue was

much lower than the hypothesized required range (Figure 3). Furthermore, by

analyzing alignments of native TatA sequences, we found that species from different

kingdoms have different preferred amino acids at the 8^{th} position correlating with

conserved 4-residue motifs at the N-terminus of TMH (Figure 5). Such 4-residue

motifs were further tested in *E. coli* TatA and proved to have significant effect on the

transport ability (Figure 6). Interestingly, these motifs were correlated with the

predominance of usage of the ΔpH or Δψ components of the pmf (Figure 7). In the

congregate, these results provide new perspectives through which to understand the

functional significance of the 8th residue in TatAs.

Our finding that the hydrophobicity, but not polarity, of the 8th amino acid, is critical

for TatA supports the idea that this residue's ability to respond to membrane potential

or to electrically interact with other polar amino acids in other Tat subunits is not

necessary for the fundamental transport mechanism.   Given that *E. coli* TatA lacks a

charged polar amino acid whilst chloroplast TatAs have a charged residue at the

corresponding position, the possibility that bacteria and thylakoids might use different

mechanisms in terms of TatA function has been discussed (New et al., 2018). It has

been debated whether an active shift of protonation states of the glutamic acid in the

chloroplasts TatA is required for a functional transporter (Berks, 2015; New et al.,

2018). However, this cannot be the case since many TatAs do not have this

protonatable residue, and we find that even Ala in this position confers some activity.

Instead, the requirement appears to be for a residue with a moderate hydrophobicity

and helix stability. In that context, the pKa of the side chain of glutamic acid matches

the thylakoid's lumen pH in the light (Kramer et al., 1999), and this changes the

overall hydrophobicity of the amino acid (Kovacs et al., 2006) to the moderate range.

Such an explanation is further supported by two observations. First, it has been shown

that thylakoid Tat transport operates with a delay between the light illumination and

transport event (a sigmoidal transport curve), even at saturating Tha4 concentrations

(Celedon and Cline, 2012). Such a delay cannot be completely attributed to the

translocon assembly step, and may also be caused by the time required to develop a

lumen pH low enough to modulate the Glu hydrophobicity. Second, our sequence

alignments revealed that nine archaea species have TatAs with an aspartic acid

(Supplemental Figure 3E) at the corresponding position whose side chain's pKa is

even lower than that of glutamic acid. Interestingly, we found by analyzing the living

environments for those species that all of them live in extreme acidic environments

(Supplemental Table 1). A random sample of archaea species with E8 did not reveal

any acidic extremophiles. While these observations do not explain why the TatAs

have acidic amino acids that require neutralizing in the first place, they do provide

examples of how Tat system might be altered to accommodate to different

physiological conditions.

Our experiments suggest that the creation of a relatively unstable N-terminus structure

by placing a polar 8th residue within the hydrophobic TatA TMH is likely significant.

Such a residue would be expected to be located at or near the membrane/water

interface, thereby biasing the position the N-terminus of the TMH close to the

membrane surface(Pettersson et al., 2018). Meanwhile, a recent study reported that

the *E. coli* TatA TMH is tuned to be 15-residues long to keep the hydrophobic

mismatch while also maintaining membrane integrity (Hao et al., 2021;

Mehner-Breitfeld et al., 2021) Also, it has been shown that TatAs have a highly

conserved 12-residue hydrophobic core (9^{th}-20^{th} in *E. coli* TatA) in the TMH (Hao et

al., 2021; Mehner-Breitfeld et al., 2021). Together these observations indicate that the

TatA TMH could be as short as 12 amino acids while the 8^{th} residue is transiently

exposed to the aqueous environment.   This might allow membrane bilayer rupture

(i.e., toroidal pore formation), with concomitant transport of a substrate held at the

site.   Our data herein demonstrated that if the 8^{th} residue was too hydrophobic

(Figure 1E), Tat transport was completely blocked, even though the hydrophobic

mismatch between TatA TMH and membrane bilayer was still maintained. This

suggests that a 15-residue long TMH is required but insufficient for active Tat

transport, and a transition from that length to an even shorter TMH might be

functionally important.

We showed that not only the 8^{th} residue is critical, but also found a changing

four-residue motif at the N-terminus of the TatA TMH affects the overall TatA

function (Figure 5 and 6).   This motif was perhaps not identified in previous studies

for two reasons. First, the functional N-terminus motif was not obviously conserved

across different species. For example, in chloroplasts and cyanobacteria, a "GxPE"

motif was nearly invariant at the N-terminus of the TatA TMH. However, this motif is

not present in proteobacterial species. Second, the identified N-terminal motifs

showed strong correlation with the 8th residue in the TatA TMH. For instance, in

proteobacteria, "SIW" was conserved in the 5th to 7th position when the 8th residue

was glutamine or histidine, whereas this motif was not conserved when lysine or

glycine was at the 8th position (Supplemental Figure 3). Even though we still do not

understand the explicit roles of the conserved motifs, the amino acid choices therein

likely provide clues about the function of N-terminus of the TatA TMH. For instance,

a tryptophan was observed immediately before H8 or Q8 in proteobacteria and in two

groups of gram-positive bacteria. One question would be what are the special features

of tryptophan that cause it to be conserved in this context. Tryptophan is normally

considered to be a non-polar amino acid in biochemistry textbooks. However, it

possesses a highly polarizable indole group in the side chain. This results in

tryptophan being highly abundant at the water-lipid interface and acting as an anchor

to stabilize membrane protein segments (de Jesus and Allen, 2013; Sun et al., 2008).

Moreover, the indole group could respond to an electric field and lower the

conformational energy of the TMH were it to algin with the electrical force lines

(Zhang and Li, 2019).   Thus, the tryptophan might help TatA embed in the

membrane and serve as an anchor near the interface. Additionally, we noticed that a

proline residue is highly conserved at position 7 when a glutamic acid is in position 8

in TatAs. Proline is generally considered a helix breaker in soluble proteins (Barlow

and Thornton, 1988). However, it has been suggested that proline would not disrupt

an alpha-helix in a membrane environment (Li et al., 1996; Senes et al., 2004).

Moreover, it has been shown that the hinge-bending motions of proline residue are

likely to play a significant role in catalysis and signal transduction (S.P. Sansom and

Weinstein, 2000; Tieleman et al., 2001). Such a hinge forming function at the

N-terminus of the TatA TMH is consistent with our hypothesis that a transition to an

even shorter TMH might be beneficial for the Tat mechanism. In addition, according

to the TatA NMR structure (Hu et al., 2010; Pettersson et al., 2018), this 4-residue

motif presents in the first coil of the TMH. Ser6 and Gln8 orientate oppositely than

Ile6 and Trp7, suggesting a weak local amphipathic topology. Moreover, other

4-residues motifs in chloroplasts and gram-positive bacteria are also arrayed this way,

even though they have different combinations of four residues at those positions.   It

is possible that this local amphipathic character assists in the reported motion of the

TMH into and out of the membrane during transport (Aldridge et al., 2012).

Incorporating our new findings related to the function of TatA subunits we propose a

speculative new model for the mechanism of Tat pathway (Figure 8). In this model,

presented here to stimulate discussion: (i) Before substrate binding, TatA, TatB and

TatC form an oligomeric structure which serves as the receptor complex for signal

peptide of Tat substrates (Alcock et al., 2016).   At this stage, relatively few TatA

proteins are associated with the TatBC complex. (ii) – (iii) After substrate binding and

in the presence of a pmf, the receptor complex undergoes a conformation change that

attracts more TatA proteins (Aldridge et al., 2014). Unlike previous models

(Rodriguez et al., 2013), we take the view of (Mehner-Breitfeld et al., 2021) and

propose these additional TatA proteins will be initially recruited into a non-circular

structure, perhaps arrayed as the double line illustrated.   (iv) As more TatA proteins

are recruited to the translocon, the hydrophobic mismatch, unstable N-terminus

structure, and the membrane-thinning action of the pmf (Matthew P Johnson et al.,

2011; Matthew P. Johnson et al., 2011; Kirchhoff et al., 2011; Murakami and Packer,

1970) cause the double line to separate and join with subunits in adjacent lines to

form a ring with a tight-fitting toroidal pore around the substrate (v). (vi) The

concomitant movement of protons during substrate translocation (Alder and Theg,

2003) results in a local dissipation of the pmf, and this in turn returns the membrane

to a thickness that favors the bilayer, causing dispersal of TatA from the translocation

site and return of the translocon to its resting condition.   While certain aspects of this

model are admittedly speculative, we believe it captures many essential properties of

Tat transport discerned from experiments.   Future studies probing the biophysical

properties of the membranes at the site of protein translocation are likely to shed

further light on the mechanism of this enigmatic protein transport system.

Methods

1. Strain and Plasmid Constructions

E. coli strain DADE-A (MC4100, *ΔtatABCE*, arabinose resistance) was used in this

study (Wexler et al., 2000). For TatA variants in pTat101 (Kneuper et al., 2012),

which is a low copy plasmid expressing the TatABC constitutively, the corresponding

single and multiple amino acid substitutions were introduced by QuikChange

site-directed mutagenesis (NEB, Phusion High-Fidelity PCR Kit) followed by KLD

(Kinase, Ligase, and DpnI) digestion (NEB). Detailed primers used for TatA variant

construction can be found in Supplemental Table 2. For TatA variants in pBAD22,

which is a vector with an arabinose-inducible *araBAD* operon, indicated TatA alleles,

together with the wild-type TatBC, were amplified from the respective constructs in

pTat101. For TatA single Q8 substitution, the forward primer TatA_gopBAD22_F

(5'-ACC ACA GAG GAA CAT GTA TGG GTG GTA TCA GTA TTT G-3') was used.

For TatA multiple substitutions, the forward primer TatAnmotif_gopBAD22_F (5'-

CTA CCA CAG AGG AAC ATG TAT GGG TGG TAT C-3') was used. Reverse

primer TatABC_gopBAD22_R (5'-TCT AGA GGC GGT TGA ATT TAT TCT TCA

GTT TTT CG C-3') was used in both cases. Empty vectors were constructed using

the primers pBAD22delABC_F (5'-ATT CAA CCG CCT CTA GAG-3') and

pBAD22delABC_R (5'-ACA TGT TCC TCT GTG GTA G-3'). Fragments and vector

were assembled using the Gibson Assembly approach (NEBuilder HiFi DNA

Assembly Master Mix). For TatA variants in pBAD33, which is a vector with an

arabinose-inducible *araBAD* operon, a 6X His tag was first added at the C-terminus of

TatA   by   using   the   primers   pBAD33His_F   (5'-

CATCATCATCACCACCACTAATGGC-3')   and   pBAD33His_R

(5'-TGATGATGACCCACCTGCTCTTTATCGTG-3'). Subsequently, TatA mutations

were introduced by QuikChange site-directed mutagenesis as described above.

pQE80l (SufI-FLAG) for the *in vivo* transport assay was constructed as describe

previously (Huang and Palmer, 2017). pNR14 and pNR42, two plasmids used in pulse

chase experiments, were as described previously (Sargent et al., 1999; Stanley et al.,

2000). All plasmids were confirmed by Sanger sequencing. More detailed information

on plasmid constructions can be found in Supplemental Table 3.

2. Sequence Alignments and Statistical Analyses

A total of 15054 TatA sequences were downloaded from the identical protein groups

in GenBank (NCBI). Sequences were then divided into five sub-groups based on

taxonomy (7534 from proteobacteria, 6978 from firmicutes and actinobacteria, 149

from cyanobacteria, 242 from chloroplast, and 151 from archaea), and were

subsequently subjected to multiple sequence alignments within each group using

MAFFT (Rozewicki et al., 2019). Sequence logos corresponding to each group were

generated using RStudio (Ver. 1.4.1103) with ggseqlogo package (Wagih, 2017). For

the interest of this project, sequences were further grouped by the identity of the

residue aligned to the eighth position in *E. coli* TatA, and the corresponding sequence

logos were generated. A stacked bar chart summarizing the distribution of the identity

of residues in this position in each taxonomical group was plotted using GraphPad

Prism version 8.2.1 for Windows (GraphPad Software, San Diego, California USA).

3. Liquid SDS Growth Assay

558   Overnight bacteria cultures were sub-cultured with an optical density (OD_{600}) of 0.002

559   were grown in the Luria-Bertani (LB) medium containing 0%, 5%, or 10% SDS,

560   respectively. After cells were incubated at 37°C with shaking for 5 hours, the final

561   OD_{600} were measured. For each mutant, survival ratio was calculated by its final OD_{600}

562   at the indicated SDS concentration to the final OD_{600} in the LB medium without SDS.

4. *in vivo* Transport Assay and Cell Fractionation

Corresponding mutant plasmids derived from pTat101 were co-transformed with

pQE80l (SufI-FLAG), which under the control of the T5 promoter, into the DADE-A

strain. Overnight cultures were diluted in a fresh LB medium at a ratio of 1:100 and

grown at 37 °C until the OD_{600} reached 0.5. A final concentration of 1 mM IPTG

(isopropyl β-D-1-thiogalactopyranoside) was added to induce the expression of

SufI-FLAG. After cultured at 37°C for another 2.5 hours, 3 mL of the cells with an

OD_{600} equivalent to 1 were fractionated using the EDTA/lysozyme/cold osmotic shock

method described before (Petiti et al., 2017b). Periplasmic fractions were collected by

adding equal volume of 2X SDS sample buffer, which were then subjected to

SDS-PAGE. Membrane fractions were washed in 10 mM Na_2CO_3 for 1 hour at 4°C

with gentle shaking. Subsequently, samples were ultracentrifuged at 120,000 x g for 45

min at 4°C. Pellets were resuspended with 2X SDS sample buffer and collected as the

membrane fractions.

5. SDS-PAGE and Western Blot

For periplasmic fractions, samples were immunoblotted with α-FLAG (GenScript)

followed by HRP-conjugated α-mouse antibodies (Santa Cruz Biotechnology). For

membrane fractions, after transfer the proteins to the PVDF membranes, PVDF

membranes were cut and immunoblotted with α-TatA and α-TatB antibodies

respectively followed by HRP-conjugated α-rabbit antibody (GenScript). Proteins were

visualized using ProSignal Pico ECL Western Blotting detection kit (Genesee

Scientific).

6. Pulse-Chase Experiment and Autoradiography

The pulse-chase experiment protocol was modified from (Stanley et al., 2000). Tat

variants carrying pNR14 and pNR42 were grown overnight in LB media at 30°C,

which were then sub-cultured in fresh LB media the next day for 1.5 hours at 30°C.

Subsequently, cells were harvested and normalized to the equivalent of 0.5 mL cells

with $OD_{600} = 0.2$. Cells were washed with 0.5 mL 1X M9 medium (diluted from 10X

M9 medium supplemented with 0.1 mM $CaCl_2$, 0.002% thiamine, 2 mM $MgSO_4$, and

0.01% of the 18 amino acids excluding methionine and cysteine, pH = 7.0). Cells were

resuspended in 2.5 mL M9 medium and cultured for another hour at 30°C. Then, cells

were transferred to 42°C for 15 min to induce T7 polymerase from pNR42. 400 µg/mL

of rifampicin was added to inhibit the *E. coli* endogenous RNA polymerase, and cells

were grown at 42°C for another 10 min. Subsequently, cells were grown at 30°C for

another 20 mins before transferred to the indicated temperature until the completion of

the experiment. If needed, pH was adjusted by 1 M HCl or 1 M KOH immediately after

this step. Bacteria cultures were then incubated at indicated temperature for 5 min.

0.025 mCi of [^{35}S] methionine (PerkinElmer Inc. NEG772002MC) was added for the

pulse step, and 750 µg/mL of unlabeled cold methionine was added after 5 min. 300 µL

of the culture was taken at the indicated time point, and transport was quenched by

freezing in liquid nitrogen. Samples were subsequently thawed on ice, centrifuge, and

resuspended with 2X SDS sample buffer followed by SDS-PAGE and autoradiography.

Results were quantified using ImageJ software.

7. Statistical Analysis of the Pulse-Chase Experimental Results

Mature-to-total (sum of precursor and mature) data from the pulse-chase experiment

was plotted against each time point, which were then fitted with the exponential plateau

model deriving from the first-order reaction model, $y = (y_{max} - y_0) \times e^{-kt}$, using

GraphPad Prism. y_{max} was defined as the maximum of the mature-to-total value, and

y_0 represents the initial mature-to-total value at the end of the pulse. t represents time

in minutes. the initial velocity (V_0), representing the transport rate when $y = 0$, was

obtained by $V_0 = y_{max} \times k$, with the units of the reciprocal minutes.

8. IMVs (Inverted-Membrane Vesicles) preparation

Tat proteins expression and IMVs preparation were performed as described

(Bageshwar and Musser, 2007b), with the following modifications. Cells were

resuspended with the same lysis buffer as described, which were then passed through a

French press at ~ 6000 psi once. Subsequently, cells were centrifuged at 30,000 g for 20

mins at 4°C. Supernatants were then centrifuged at 140,000 g for 1 hr. Pellets were

resuspended with IMV storage buffer (10 mM Tris-HCl, pH = 7.5, 1 mM $MgSO_4$, 1

mM KCl, 40% glycerol). IMVs were stored in -80°C freezer.

9. Proton Leakage Measurement

IMVs were made from the DADE-A cells expressing the 6X His-tagged wild-type or

TatA variants in pBAD33 alone or together with the pTat101(*ΔtatA*). Acridine orange

fluorescence-quenching assays were carried out using the Fluorolog-3

spectrofluorometer (HORIBA Scientific, model No. FL3-22). IMVs were added to the

final concentration equivalent to $A_{280} = 0.375$ to the reaction mix containing 1X TE

buffer (25 mM MOPS, 25 mM MES, 5 mM $MgCl_2$, 50 mM KCl, 200 mM sucrose, and

57 µg/mL BSA, pH = 7.0), ATP regeneration system (2.9 mM phosphocreatine, 0.29

Functional analysis of the polar amino acid in TatA TMH

mg/mL creatine kinase), and 2 µM acridine orange. The reaction mix was incubated for

5 mins with gentle stirring at 37°C before measurement. The fluorescence of acridine

orange was recorded at $\lambda_{ex} = 494$ nm (slit = 1 nm) and $\lambda_{em} = 540$ nm (slit = 5 nm) per 0.1

second. 4 mM of ATP was added at 60 seconds, and 10 µM of CCCP was added at 200

seconds to dissipate the proton gradient.

Acknowledgement

We gratefully acknowledge support from the Division of Chemical Sciences,

Geosciences, and Biosciences, Office of Basic Energy Sciences of the U.S.

Department of Energy through Grant DE-SC0020304 to SMT.

References

Alcock F, Stansfeld PJ, Basit H, Habersetzer J, Baker MA, Palmer T, Wallace MI,
Berks BC. 2016. Assembling the Tat protein translocase. *eLife* **5**:e20718.
doi:10.7554/eLife.20718

Alder NN, Theg SM. 2003. Energetics of protein transport across biological
membranes: A study of the thylakoid ΔpH-dependent/cpTat pathway. *Cell*
112:231–242. doi:10.1016/S0092-8674(03)00032-1

Aldridge C, Ma X, Gerard F, Cline K. 2014. Substrate-gated docking of pore subunit
Tha4 in the TatC cavity initiates Tat translocase assembly. *J Cell Biol*
205:51–65. doi:10.1083/jcb.201311057

Aldridge C, Storm A, Cline K, Dabney-Smith C. 2012. The Chloroplast Twin
Arginine Transport (Tat) Component, Tha4, Undergoes Conformational
Changes Leading to Tat Protein Transport. *J Biol Chem* **287**:34752–34763.
doi:10.1074/jbc.M112.385666

Asher AH, Theg SM. 2021. Electrochromic shift supports the membrane
destabilization model of Tat-mediated transport and shows ion leakage during
Sec transport. *Proc Natl Acad Sci U S A* **118**. doi:10.1073/pnas.2018122118

Bageshwar UK, Musser SM. 2007a. Two electrical potential–dependent steps are
required for transport by the Escherichia coli Tat machinery. *J Cell Biol*
179:87–99. doi:10.1083/jcb.200702082

Bageshwar UK, Musser SM. 2007b. Two electrical potential-dependent steps are
required for transport by the Escherichia coli Tat machinery. *J Cell Biol*
179:87–99. doi:10.1083/jcb.200702082

Barlow DJ, Thornton JM. 1988. Helix geometry in proteins. *J Mol Biol* **201**:601–619.
doi:10.1016/0022-2836(88)90641-9

Berks BC. 2015. The Twin-Arginine Protein Translocation Pathway. *Annu Rev*
Biochem **84**:843–864. doi:10.1146/annurev-biochem-060614-034251

Braun NA, Davis AW, Theg SM. 2007. The Chloroplast Tat Pathway Utilizes the
Transmembrane Electric Potential as an Energy Source. *Biophys J*
93:1993–1998. doi:10.1529/biophysj.106.098731

Brüser T, Sanders C. 2003. An alternative model of the twin arginine translocation
system. *Microbiol Res* **158**:7–17. doi:10.1078/0944-5013-00176

Celedon JM, Cline K. 2012. Stoichiometry for binding and transport by the twin
arginine translocation system. *J Cell Biol* **197**:523–534.
doi:10.1083/jcb.201201096

Chou PY, Fasman GD. 1978. Empirical Predictions of Protein Conformation. *Annu*
Rev Biochem **47**:251–276. doi:10.1146/annurev.bi.47.070178.001343

Clark SA, Theg SM. 1997. A folded protein can be transported across the chloroplast
envelope and thylakoid membranes. *Mol Biol Cell* **8**:923–934.
doi:10.1091/mbc.8.5.923

Cline K, Ettinger WF, Theg SM. 1992. Protein-specific energy requirements for

protein transport across or into thylakoid membranes. Two lumenal proteins
are transported in the absence of ATP. *J Biol Chem* **267**:2688–2696.
Cruz JA, Sacksteder CA, Kanazawa A, Kramer DM. 2001. Contribution of Electric
Field (Δψ) to Steady-State Transthylakoid Proton Motive Force (pmf) in Vitro
and in Vivo. Control of pmf Parsing into Δψ and ΔpH by Ionic Strength.
Biochemistry **40**:1226–1237. doi:10.1021/bi0018741
Dabney-Smith C, Mori H, Cline K. 2003. Requirement of a Tha4-conserved
Transmembrane Glutamate in Thylakoid Tat Translocase Assembly Revealed
by Biochemical Complementation. *J Biol Chem* **278**:43027–43033.
doi:10.1074/jbc.M307923200
de Jesus AJ, Allen TW. 2013. The role of tryptophan side chains in membrane protein
anchoring and hydrophobic mismatch. *Biochim Biophys Acta BBA - Biomembr*
1828:864–876. doi:10.1016/j.bbamem.2012.09.009
Fincher V, Dabney☐Smith C, Cline K. 2003. Functional assembly of thylakoid
ΔpH-dependent/Tat protein transport pathway components in vitro. *Eur J*
Biochem **270**:4930–4941. doi:10.1046/j.1432-1033.2003.03894.x
Ghosh D, Boral D, Vankudoth KR, Ramasamy S. 2019. Analysis of haloarchaeal
twin-arginine translocase pathway reveals the diversity of the machineries.
Heliyon **5**:e01587. doi:10.1016/j.heliyon.2019.e01587
Gohlke U, Pullan L, McDevitt CA, Porcelli I, Leeuw E de, Palmer T, Saibil HR,
Berks BC. 2005. The TatA component of the twin-arginine protein transport
system forms channel complexes of variable diameter. *Proc Natl Acad Sci*
102:10482–10486. doi:10.1073/pnas.0503558102
Greene NP, Porcelli I, Buchanan G, Hicks MG, Schermann SM, Palmer T, Berks BC.
2007. Cysteine Scanning Mutagenesis and Disulfide Mapping Studies of the
TatA Component of the Bacterial Twin Arginine Translocase. *J Biol Chem*
282:23937–23945. doi:10.1074/jbc.M702972200
Hao B, Zhou W, Theg SM. 2021. Hydrophobic mismatch effect is a key factor in
protein transport on the Tat pathway. *bioRxiv* 2021.05.26.445883.
doi:10.1101/2021.05.26.445883
Hauer RS, Freudl R, Dittmar J, Jakob M, Klösgen RB. 2017. How to achieve Tat
transport with alien TatA. *Sci Rep* **7**:1–13. doi:10.1038/s41598-017-08818-w
Hicks MG, Leeuw E de, Porcelli I, Buchanan G, Berks BC, Palmer T. 2003. The
Escherichia coli twin-arginine translocase: conserved residues of TatA and
TatB family components involved in protein transport. *FEBS Lett* **539**:61–67.
doi:10.1016/S0014-5793(03)00198-4
Hou B, Heidrich ES, Mehner-Breitfeld D, Brüser T. 2018. The TatA component of the
twin-arginine translocation system locally weakens the cytoplasmic membrane
of Escherichia coli upon protein substrate binding. *J Biol Chem*
293:7592–7605. doi:10.1074/jbc.RA118.002205
Hu Y, Zhao E, Li H, Xia B, Jin C. 2010. Solution NMR Structure of the TatA
Component of the Twin-Arginine Protein Transport System from

Gram-Positive Bacterium Bacillus subtilis. *J Am Chem Soc* **132**:15942–15944.
doi:10.1021/ja053785

Huang Q, Palmer T. 2017. Signal Peptide Hydrophobicity Modulates Interaction with
the   Twin-Arginine   Translocase.   *mBio*   **8**:e00909-17.
doi:10.1128/mBio.00909-17

Ize B, Stanley NR, Buchanan G, Palmer T. 2003. Role of the Escherichia coli Tat
pathway in outer membrane integrity. *Mol Microbiol* **48**:1183–1193.
doi:10.1046/j.1365-2958.2003.03504.x

Johnson Matthew P, Brain AP, Ruban AV. 2011. Changes in thylakoid membrane
thickness associated with the reorganization of photosystem II light harvesting
complexes during photoprotective energy dissipation. *Plant Signal Behav*
6:1386–1390. doi:10.4161/psb.6.9.16503

Johnson Matthew P., Goral TK, Duffy CDP, Brain APR, Mullineaux CW, Ruban AV.
2011. Photoprotective Energy Dissipation Involves the Reorganization of
Photosystem II Light-Harvesting Complexes in the Grana Membranes of
Spinach Chloroplasts. *Plant Cell* **23**:1468–1479. doi:10.1105/tpc.110.081646

Kirchhoff H, Hall C, Wood M, Herbstová M, Tsabari O, Nevo R, Charuvi D, Shimoni
E, Reich Z. 2011. Dynamic control of protein diffusion within the granal
thylakoid   lumen.   *Proc   Natl   Acad   Sci*   **108**:20248–20253.
doi:10.1073/pnas.1104141109

Kneuper H, Maldonado B, Jäger F, Krehenbrink M, Buchanan G, Keller R, Müller M,
Berks BC, Palmer T. 2012. Molecular dissection of TatC defines critical
regions essential for protein transport and a TatB–TatC contact site. *Mol*
Microbiol **85**:945–961. doi:10.1111/j.1365-2958.2012.08151.x

Kovacs JM, Mant CT, Hodges RS. 2006. Determination of intrinsic
hydrophilicity/hydrophobicity of amino acid side chains in peptides in the
absence of nearest-neighbor or conformational effects. *Pept Sci* **84**:283–297.
doi:https://doi.org/10.1002/bip.20417

Kramer DM, Sacksteder CA, Cruz JA. 1999. How acidic is the lumen? *Photosynth*
Res **60**:151–163. doi:10.1023/A:1006212014787

Leake MC, Greene NP, Godun RM, Granjon T, Buchanan G, Chen S, Berry RM,
Palmer T, Berks BC. 2008. Variable stoichiometry of the TatA component of
the twin-arginine protein transport system observed by in vivo single-molecule
imaging.   *Proc   Natl   Acad   Sci*   **105**:15376–15381.
doi:10.1073/pnas.0806338105

Li SC, Goto NK, Williams KA, Deber CM. 1996. Alpha-helical, but not beta-sheet,
propensity of proline is determined by peptide environment. *Proc Natl Acad*
Sci **93**:6676–6681. doi:10.1073/pnas.93.13.6676

Mant CT, Hodges RS. 2002. Reversed-phase liquid chromatography as a tool in the
determination of the hydrophilicity/hydrophobicity of amino acid side-chains
at a ligand–receptor interface in the presence of different aqueous
environments: I. Effect of varying receptor hydrophobicity. *J Chromatogr A*,

773   21st International Symposium on the Separation of Proteins, Peptides and
774   Polynucleotides **972**:45–60. doi:10.1016/S0021-9673(02)00354-0

775 Mehner-Breitfeld D, Ringel MT, Tichy DA, Endter LJ, Stroh KS, Luensdorf H,
776   Risselada HJ, Brueser T. 2021. TatA and TatB generate a hydrophobic
777   mismatch that is important for function and assembly of the Tat translocon in
778   Escherichia coli. *bioRxiv* 2021.05.26.445790. doi:10.1101/2021.05.26.445790

779 Mori H, Cline K. 2001. Post-translational protein translocation into thylakoids by the
780   Sec and ΔpH-dependent pathways. *Biochim Biophys Acta BBA - Mol Cell Res*
781   **1541**:80–90. doi:10.1016/S0167-4889(01)00150-1

782 Mori H, Summer EJ, Cline K. 2001. Chloroplast TatC plays a direct role in thylakoid
783   ΔpH-dependent protein transport. *FEBS Lett* **501**:65–68.
784   doi:10.1016/S0014-5793(01)02626-6

785 Murakami S, Packer L. 1970. Light-induced Changes in the Conformation and
786   Configuration of the Thylakoid Membrane of Ulva and Porphyra Chloroplasts
787   in Vivo. *Plant Physiol* **45**:289–299. doi:10.1104/pp.45.3.289

788 New CP, Ma Q, Dabney-Smith C. 2018. Routing of thylakoid lumen proteins by the
789   chloroplast twin arginine transport pathway. *Photosynth Res* **138**:289–301.
790   doi:10.1007/s11120-018-0567-z

791 Palmer T, Berks BC, Sargent F. 2010. Analysis of Tat Targeting Function and
792   Twin-Arginine Signal Peptide Activity in Escherichia coli In: Economou A,
793   editor. Protein Secretion: Methods and Protocols, Methods in Molecular
794   Biology. Totowa, NJ: Humana Press. pp. 191–216.
795   doi:10.1007/978-1-60327-412-8_12

796 Petiti M, Houot L, Duché D. 2017a. Cell Fractionation In: Journet L, Cascales E,
797   editors. Bacterial Protein Secretion Systems: Methods and Protocols, Methods
798   in Molecular Biology. New York, NY: Springer New York. pp. 59–64.
799   doi:10.1007/978-1-4939-7033-9_3

800 Petiti M, Houot L, Duché D. 2017b. Cell Fractionation In: Journet L, Cascales E,
801   editors. Bacterial Protein Secretion Systems: Methods and Protocols, Methods
802   in Molecular Biology. New York, NY: Springer. pp. 59–64.
803   doi:10.1007/978-1-4939-7033-9_3

804 Pettersson P, Ye W, Jakob M, Tannert F, Klösgen RB, Mäler L. 2018. Structure and
805   dynamics of plant TatA in micelles and lipid bilayers studied by solution NMR.
806   *FEBS J* **285**:1886–1906. doi:10.1111/febs.14452

807 Rodriguez F, Rouse SL, Tait CE, Harmer J, Riso AD, Timmel CR, Sansom MSP,
808   Berks BC, Schnell JR. 2013. Structural model for the protein-translocating
809   element of the twin-arginine transport system. *Proc Natl Acad Sci*
810   **110**:E1092–E1101. doi:10.1073/pnas.1219486110

811 Rollauer SE, Tarry MJ, Graham JE, Jääskeläinen M, Jäger F, Johnson S, Krehenbrink
812   M, Liu S-M, Lukey MJ, Marcoux J, McDowell MA, Rodriguez F, Roversi P,
813   Stansfeld PJ, Robinson CV, Sansom MSP, Palmer T, Högbom M, Berks BC,
814   Lea SM. 2012. Structure of the TatC core of the twin-arginine protein transport

system. *Nature* **492**:210–214. doi:10.1038/nature11683

Rozewicki J, Li S, Amada KM, Standley DM, Katoh K. 2019. MAFFT-DASH:

integrated protein sequence and structural alignment. *Nucleic Acids Res*

47:W5–W10. doi:10.1093/nar/gkz342

Sargent F, Gohlke U, Leeuw E de, Stanley NR, Palmer T, Saibil HR, Berks BC. 2001.

Purified components of the Escherichia coli Tat protein transport system form

a double-layered ring structure. *Eur J Biochem* **268**:3361–3367.

doi:10.1046/j.1432-1327.2001.02263.x

Sargent F, Stanley NR, Berks BC, Palmer T. 1999. Sec-independent Protein

Translocation in Escherichia coli. *J Biol Chem* **274**:36073–36082.

doi:10.1074/jbc.274.51.36073

Senes A, Engel DE, DeGrado WF. 2004. Folding of helical membrane proteins: the

role of polar, GxxxG-like and proline motifs. *Curr Opin Struct Biol*

14:465–479. doi:10.1016/j.sbi.2004.07.007

S.P. Sansom M, Weinstein H. 2000. Hinges, swivels and switches: the role of prolines

in signalling via transmembrane α-helices. *Trends Pharmacol Sci* **21**:445–451.

doi:10.1016/S0165-6147(00)01553-4

Stanley NR, Palmer T, Berks BC. 2000. The twin arginine consensus motif of Tat

signal peptides is involved in Sec-independent protein targeting in Escherichia

coli. *J Biol Chem* **275**:11591–11596. doi:10.1074/jbc.275.16.11591

Sun H, Greathouse DV, Andersen OS, Koeppe RE. 2008. The Preference of

Tryptophan for Membrane Interfaces. *J Biol Chem* **283**:22233–22243.

doi:10.1074/jbc.M802074200

Teter SA, Theg SM. 1998. Energy-transducing thylakoid membranes remain highly

impermeable to ions during protein translocation. *Proc Natl Acad Sci*

95:1590–1594. doi:10.1073/pnas.95.4.1590

Tieleman DP, Shrivastava IH, Ulmschneider MR, Sansom MSP. 2001.

Proline-induced hinges in transmembrane helices: Possible roles in ion

channel gating. *Proteins Struct Funct Bioinforma* **44**:63–72.

doi:10.1002/prot.1073

Tsirigotaki A, De Geyter J, Šoštaric´ N, Economou A, Karamanou S. 2017. Protein

export through the bacterial Sec pathway. *Nat Rev Microbiol* **15**:21–36.

doi:10.1038/nrmicro.2016.161

Wagih O. 2017. ggseqlogo: a versatile R package for drawing sequence logos.

Bioinformatics **33**:3645–3647. doi:10.1093/bioinformatics/btx469

Warren G, Oates J, Robinson C, Dixon AM. 2009. Contributions of the

Transmembrane Domain and a Key Acidic Motif to Assembly and Function of

the TatA Complex. *J Mol Biol* **388**:122–132. doi:10.1016/j.jmb.2009.02.060

Wexler M, Sargent F, Jack RL, Stanley NR, Bogsch EG, Robinson C, Berks BC,

Palmer T. 2000. TatD is a cytoplasmic protein with DNase activity. No

requirement for TatD family proteins in sec-independent protein export. *J Biol*

Chem **275**:16717–16722. doi:10.1074/jbc.M000800200

Yen M-R, Tseng Y-H, Nguyen EH, Wu L-F, Saier MH. 2002. Sequence and
858   phylogenetic analyses of the twin-arginine targeting (Tat) protein export
859   system. *Arch Microbiol* **177**:441–450. doi:10.1007/s00203-002-0408-4

860   Zhang XC, Li H. 2019. Interplay between the electrostatic membrane potential and
861   conformational changes in membrane proteins. *Protein Sci* **28**:502–512.
862   doi:10.1002/pro.3563

863   Zhang Y, Wang L, Hu Y, Jin C. 2014. Solution structure of the TatB component of the
864   twin-arginine translocation system. *Biochim Biophys Acta BBA - Biomembr*
865   **1838**:1881–1888. doi:10.1016/j.bbamem.2014.03.015

866   Zilberstein D, Agmon V, Schuldiner S, Padan E. 1984. Escherichia coli intracellular
867   pH, membrane potential, and cell growth. *J Bacteriol* **158**:246–252.

Figure 1. Polar residues at the eighth position of the *E. coli* TatA TMH are preferred but not necessary for minimal Tat activity. (A-C) Minimal Tat transport abilities of the TatA Q8 variants were evaluated by the growth performance in LB medium with indicated SDS concentrations. wtTatA served as the positive control, while *ΔtatA* acted as the negative control. Survival ratio denotes the ratio of the OD_{600} of the cells grown in the presence of the corresponding SDS concentration to the OD_{600} of the cells grown without SDS. Three biological replicates for each mutant were included in the experiments. (D) Western blot results of the *in vivo* transport assays with the TatA Q8 variants. Dilutions of the amount of transported SufI-FLAG in the wtTat cells were shown for the comparison of the transport performance in the TatA Q8 variants. p, precursor and m, mature. (E) Plot of helix stability (Δt_m) versus side chain hydrophobicity (Δt_R) of amino acids colored according to their activity in the 8[th] position of TatA.   Different levels of the Tat transport activities are denoted with the following rules: no Tat activity, no growth in the LB medium containing 5% SDS; low Tat activity, mutants who could grow in the LB with SDS but showed less than 50% transport activity than wtTat in the *in vivo* transport assay in (D); high Tat activity, mutants with more than 50% of the wtTat activity in (D). Identities of the amino acids with high Tat activity were clustered in the ranges of moderate helix stability and side chain hydrophobicity, which is boxed in pink.

Figure 2. Quantitative analyses of the Tat activity in the TatA Q8 variants. (A)
Plot of the transport kinetics of the TatA Q8 variants in pulse-chase experiments.
Identities of wtTatA and TatA Q8 variants are shown in the right. Mature/total
denotes the ratio of the mature band to the sum of the precursor and mature bands
revealed by autoradiography of the samples from the pulse-chase experiment
transporting SufI; chase time is denoted on the abscissa.   Lines show best fits of an
exponential model to the points arising from at least two biological replicates for each
mutant.   (B) Plot of the initial velocities (V_0) in the TatA Q8 variants transporting
SufI.   Exact values of the V_0 are shown above the bars.   TatA Q8 variants are
grouped based classification of the amino acids.

Figure 3. Comparison of the Tat activity of wtTat and Q8H mutant at different
pH conditions. (A) Radioautography of the wtTatA and Q8H variants transporting
SufI at different pHs. Samples from the pulse-chase experiment were subjected to
SDS-PAGE. 8-16% polyacrylamide gels were used to separate the mature SufI (m)
from the precursor (p). Corresponding times of the chase steps are shown on top. Two
replicates for each variant were included in this experiment. (B-C) Quantitative
analyses of the transport performance of the wtTatA (B) and Q8H (C) variants based
on the radioautography in (A). Same analysis approach was used as mentioned earlier.
The pH condition in which the experiment was performed is indicated in the legend of

the plot. (D) Plot of the initial velocities under different pH conditions between the

wtTat and Q8H variants.   Initial velocities of the wtTatA under different pH are

denoted in black, and initial velocities of the Q8H variant under different pH are

denoted in grey. (E-H) Comparison of the transport kinetics between the wtTatA and

Q8H variants under the same pH (E, pH = 5.0; F, pH = 6.0; G, pH = 7.0; H, pH = 8.0),

with the filled symbols representing the wtTatA, and empty circles representing the

Q8H. Other details as described in the legend of Fig. 2.

Figure 4. Membrane leakage profiles of TatA mutants. The ΔpH developed
across the membrane was measured by the acridine orange quenching in the IMVs. 4
mM ATP was added at the 60 secs to energize the IMV, and 10 μM CCCP was added
at 200 secs to dissipate the proton gradient, resulting the recovery of the fluorescence.
(A) Measurement of the ΔpH in the DADE (Tat knockout variant) IMVs with
indicated overexpressed TatA Q8 variants alone or together with the constitutively
expressed TatBC. (B) Measurement of the ΔpH in the DADE (Tat knockout variant)
IMVs with indicated overexpressed TatA N-terminal motif variants alone. (C)
Measurement of the ΔpH in the DADE (Tat knockout variant) IMVs with indicated

overexpressed TatA N-terminal motif variants together with the constitutively
expressed TatBC. AU, arbitrary unit.
934

935
936 **Figure 5. Different species have different preferences at the 8th position in TatA.**

Figure 5. Different species have different preferences at the 8^{th} position in TatA.
Sequence logos of the TatA sequences alignment across (A) proteobacteria, (B)
actinobacteria and firmicutes, (C) chloroplasts, (D) cyanobacteria, and (E) archaea.
Numbers of sequences included in the corresponding categories were denoted as N.
The N-terminal motifs were boxed as indicated. (F) Distributions of amino acids
selected in the location equivalent to the 8^{th} position of the *E. coli* TatA across
different species. Percentage of species with the indicated residues were shown.
Amino acids were colored as indicated to the right of the plot.
944
945

Figure 6. Comparison of the Tat transport activity with the TatA N-terminal motif mutants. (A) Membrane after carbonate washing and total extract samples were subjected to SDS-PAGE and immunoblotting with the α-TatA antibody. Molecular weight is shown on the left. (B) Periplasmic fractions (top) and total extract (bottom) of the cells in the *in vivo* transport assays were isolated. Precursor (p) and mature (m) proteins were labeled on the right, and molecular weight is shown on the left. (C-E) Plots of the transport kinetics for the N-terminal motif mutants in the

954   pulse-chase experiment. At least two replicates were included for each mutant. (F)

955   Plot of the initial velocities (V_0) of the N-terminal motif mutants in the pulse-chase

956   experiments. V_0 was obtained from the kinetic plots in (C-E). Same methodology was

957   implemented as described before. Exact values of the initial velocities are displayed

958   on top of bar graph. Other details as described in the legend of Fig. 2.

Figure 7. TatA N-terminus mutants display differences in energy utilization preference and thermostability. (A) Comparison of the IMV transport of SufI among wtTatA, TatA Q8E (SIWE), and TatA GVPE mutants in response to 10 μM nigericin or 10 μM valinomycin treatment.   The control groups received equivalent amounts of EtOH and were set to 100%. The exact percentage of transport in the experimental groups are shown above the bars.   Each treatment group per mutant was performed in four replicates, see Supplemental Figure 4.

969
970 **Figure 8. Schematic diagram for the proposed Tat transport mechanism.** In this
971 diagram, a view of the proposed Tat transport process is shown from above the
972 membrane.   Side view of the TatA interfaces is shown on the right of the step 4.
973 Blue/grey circles represent lipid head group.