The influence of sponge-dwelling gobies (*Elacatinus horsti*) on the
feeding efficiency of their Caribbean tube sponge (*Aplysina* sp.)
hosts

Megan J. Siemann[1,2,3], Aldo Turco[2,4], Shannon D. Brown[2,5], Rita B.J. Peachey[2,6]

[1]Biosciences Department, Rice University, Houston, TX USA;
[2]CIEE Research Station, Kralendijk, Bonaire, Caribbean Netherlands;
[3]Department of Translational Molecular Pathology, The University of Texas MD Anderson
Cancer Center, Houston TX USA;
[4]Golder Associates, Melbourne, Australia;
[5]Red Sea Research Center, King Abdullah University of Science and Technology, Thuwal,
Kingdom of Saudi Arabia;
[6]Institute of Sustainable Technology, Kaya Grandi #77, Kralendijk, Bonaire, Caribbean
Netherlands

Correspondence:
Megan J. Siemann
megan.siemann@gmail.com
+1-713-865-6297

Keywords: *Aplysina, Elacatinus,* mutualism, yellowline goby

Abstract Mutualistic associations between benthic marine invertebrates and reef taxa are common.

Sponge-dwelling gobies benefit from protection within sponge tubes and greater food

availability. Sponge-dwelling gobies are hypothesized to increase sponge pump rates by

consuming polychaete parasites, but such increases have not yet been demonstrated. We

investigated the association between sponge-dwelling gobies (*Elacatinus horsti*) and two

species of tube sponge (*Aplysina lacunosa* and *Aplysina archeri*) in Bonaire, Caribbean

Netherlands. We visually assessed goby presence in sponges and used *in situ* methods with

fluorescein dye to measure estimate feeding rates via pump rates. *Aplysina archeri* were more

likely to host a goby than *A. lacunosa.* For both sponge species, pump rates of tubes with

gobies were higher on average than those of tubes without gobies. Our observations,

therefore, suggest that *E. horsti* associations with *Aplysina* are likely mutualistic relationships

in which sponges benefit from higher feeding rates when gobies are present.

Introduction

Sponges are found in most marine habitats from equatorial to Arctic waters (Bell 2008). As active suspension feeders, sponges are recognized as important planktonic feeders in coral reef food webs (Lesser 2006) with the potential to alter the environment around them. Investigating the magnitude of sponge filtration and the factors which influence their filtration ability will help clarify their influence on local ecosystems. At high biomasses, sponges reduce picoplankton availability in the water column which impacts food availability to other planktivores (Savarese et al. 1997). Oysters and brittle stars have a symbiotic relationship with sponges in which the sponge benefits from increased pump rates (Hendler 1984; Burns and Bingham 2002). Here we assumed that higher pump rates enable greater feeding rates. Reviews have categorized interspecific interactions of sponges as predation, competition, or utilization of sponges as microhabitat (Wulff 2006; Bell 2008). For instance, invertebrates and fish dwell inside sponge microhabitats (Wulff 2006; Bell 2008).

Gobies associated with sponges include free-swimming gobies and obligate sponge-dwelling gobies, such as yellowline goby (*Elacatinus horsti*) (Tyler and Böhlke 1972; Colin 1975; D'Aloia et al. 2011). Some cleaner gobies spend time in sponges but are not considered sponge-dwelling (Whiteman and Côté 2002). *Elacatinus horsti* occupy convoluted barrel sponges (*Aplysina lacunosa*) and stove-pipe sponges (*Aplysina archeri*), two species of tube sponge found in the Caribbean (Villamizar and Laughlin 1991). Both *A. archeri* and *A. lacunosa* grow in clumps of tubes with individual tubes reaching maximum lengths of 180 cm and 90 cm, respectively (Humann and DeLoach 1992).

While many organisms have been documented inhabiting sponges (Tyler and Böhlke 1972; Villamizar and Laughlin 1991; Meroz and Ilan 1995; Bell 2008; D'Aloia et al. 2011), little is known regarding the effects of the inhabitants on the sheltering sponge. Hendler (1984) found that sponges benefit from resident brittle stars that clear sediment from their

tubes. The relationship between sponge-dwelling gobies and their sponge hosts was originally described as obligate commensalism with gobies benefitting from the shelter (Böhlke and Robins 1969; Majoris et al. 2018). Other research has recognized sponge-dwelling invertebrates as an additional benefit to *Elacatinus* goby sp.. Colin (1975) concluded that *E. horsti* were eating parasitic polychaetes because sponge tissue was found in the gut content of *E. horsti*.

Furthermore, it has been proposed that sponges with sponge-dwelling gobies benefit from lower parasite loads (Whiteman and Côté 2002; D'Aloia et al. 2011), which may increase sponge feeding efficiency. However, directly measuring a sponges' feeding efficiency *in situ* is considered difficult (Yahel et al. 2005). Therefore, in numerous studies, the pump rate has been considered a proxy for feeding efficiency on the assumption that the more water a sponge can pump, the more food can be filtered (Gerrodette and Flechsig 1979; Hendler 1984; Savarese et al. 1997). The pump rate of a species is also affected by the size of sponges and depth, which must be considered potential sources of variation (Lesser 2006; McMurray et al. 2014), and therefore, accounted for in studies on the filtration rates of sponge.

In this study, we quantified the effect of *E. horsti* gobies on *Aplysina* sponges using pump rate as a proxy for feeding rate. We predicted that tubes of *Aplysina lacunosa* and *Aplysina archeri* containing *E. horsti* gobies would have a higher pump rates (i.e., greater feeding efficiency).

Methods

Study site

We conducted this study from September - October 2017 at Playa Lechi dive site (12°09'36.6"N, 68°16'55.1"W) on the leeward side of the island of Bonaire, Caribbean Netherlands. The study area is characterized by a reef slope starting at 5 m depth which continues until ~35 m deep where it flattens to sandy substrate. While *A. archeri* and *A.*

lacunosa are found in depths of 3-45 m (Villamizar and Laughlin 1991), on SCUBA, we

conducted this study between 5-15 m deep to limit depth-dependent effects on pump rate

(Bell 2008). Short study duration and daytime dive times were chosen to limit seasonal and

daily temporal cycles in sponge pump rates (Morganti et al. 2019).

Investigation of sponge abundance and symbiont presence

We performed 50 m transects at 10 m depth to investigate the abundance of both *Aplysina*

species and the frequency of *E. horsti*. Only presence and absence information was recorded

for the gobies. To limit the effect of the divers, a 3-min acclimation period was allotted after

laying the transect.

Measuring pump rate

In situ methods

We randomly selected *A. archeri* and *A. lacunosa* and checked tubes for *E. horsti*. At each

sponge, we measured tube length and pump rate on the most landward tube with *E. horsti*

and, where possible, the most landward tube without *E. horsti*. We measured pump rates

following the approach of Gerrodette and Flechsig (1979). We taped a flexible, clear tube (1

cm diameter) with scale marks to a hole in a 700 ml resealable, plastic bag. Then, we secured

the bag to the sponge osculum using rubber bands. One diver injected 2 ml of 100 mg L^{-1}

fluorescein dyed seawater (Yahel et al. 2005) through a hole near the base of the tube, while

the other stabilized the tube and filmed dye movement at 30 frames s^{-1}. For each sponge tube,

we released dye three times then removed the bag.

Video processing

We used linear regressions to calculate the velocity of the dye front on a fixed 2D plane

based on scale marks on the tube and frame rate (Tracker 4.11.0 software) and used the tube's

cross-section area to convert these to pump rates.

Statistical analysis

We performed separate repeated measures (multiple measurements per tube) ANCOVAs for

each species of sponge to determine whether pump rate depended on the presence of *E.*

horsti (categorical predictor) and tube length (continuous predictor). *Elacatinus horsti*

presence was treated as a binary variable as only 3 of 72 tubes tested for pump rate contained

more than one *E. horsti*. We used a z-test (two-tail) to test for differences between the

proportions of tubes containing *E. horsti* for the two species of sponge.

Results and discussion

Mutualistic symbiosis

Pump rates of both sponge species increased when *E. horsti* were present (*A. lacunosa*:

$F_{1,32}$=10.56, p=0.0027, +2.35 cm^3 s^{-1}; *A. archeri*: $F_{1,34}$=8.39, p=0.0458, +1.78 cm^3 s^{-1}; Fig. 1).

The significantly higher pump rates of tubes with *E. horsti* supported the hypothesis of a

conferred feeding benefit to sponges. In nature, the symbiosis is thought to be obligatory for

the sponge-dwelling goby that is dependent on the sponge for both food and shelter (Colin

1975; Whiteman and Côté 2004a; D'Aloia et al. 2011). *Elacatinus horsti* benefit from the

interaction through nutrition supplied from eating parasites that feed on sponge tissue plus

gobies are readily attacked by predators when disturbed from their sponge tubes or held in

open, experimental tanks (Colin 1975). Nevertheless, it may also be that gobies choose tubes

with higher pump rates, a possibility that would require experimental introductions or

removals of gobies to test. Our results are consistent with a benefit to the sponge and,

therefore, a mutually beneficial symbiosis.

We did not explicitly test parasite reduction as a potential mechanism driving the

difference in pump rates But other studies have shown that parasitic polychaetes, which live

inside sponge tissue and consume choanocytes, have a negative impact on the sponge

(Magnino and Gaino 1998). Colin (1975) found that *E. horsti* consume *Haplosyllis* sp., but

this conclusion was based on gut content analysis of a small number of fish. In addition, the

relationship between *Haplosyllis* sp. and the pump rate of a sponge remains poorly

understood (Lattig and Martin 2011). To fully understand the interaction among gobies,

sponges, and polychaetes, it is important to examine the relationship between parasite loads

and the pump rate of sponges.

The clearing of sediment or algae from sponge tissue has also been suggested as the

mechanism of pump rate increases in invertebrate-sponge symbioses because the feeding

ability of a sponge decreases when sediment covers their collar cells (Gerrodette and Flechsig

1979; Burns and Bingham 2002). Decreases in pumping rate due to increased sediment has

been documented for *A. lacunosa* sponges (Gerrodette and Flechsig 1979). It is possible that

E. horsti movement and feeding clears sediment from sponge choanocytes in a relationship

similar to one seen between brittle-stars and sponges (Hendler 1984). Regardless of the

underlying mechanism, we found that goby presence is associated with an improved pumping

efficiency which likely increases the feeding ability of the host sponge. These results may

apply to other sponge species that host microhabitat dwellers (Tyler and Böhlke 1972;

Villamizar and Laughlin 1991; Bell 2008) and is likely to apply to other sponge-dwelling

Elacatinus sp. with similar biological and ecological traits.

Interspecific differences

We found that pump rates increased with tube length for *A. archeri* ($F_{1,34}$=24.26, p<0.0001,

slope: 0.097 cm^2 s^{-1}) but did not depend on tube length for *A. lacunosa* ($F_{1,32}$=0.09, p=0.7664;

Fig 1). Other studies have shown that the pump rates of sponges are higher for longer tubes

(Lesser 2006). The pump rates we found here are comparable to those reported previously for

sponges of similar size (Gerrodette and Flechsig 1979; Morganti et al. 2019). The underlying

differences between the sponge species, such as the bumpy outer surface seen in *A. lacunosa*

versus the smoother textured *A. archeri* tube, may be the cause of our observed lack of

correlation between tube lengths and pump rate in *A. lacunosa* (Villamizar and Laughlin

1991). To more clearly understand the effect of tube length on pumping, other factors

affecting interspecific differences need to be examined.

Aplysina archeri tubes were more likely to have an *E. horsti* present than were *A.*

lacunosa tubes (z=3.32, p<0.001; Fig. 2). In other southern Caribbean reefs, Villamizar and

Laughlin (1991) found a lower frequency of *E. horsti* in *A. lacunosa* tubes compared to *A.*

archeri tubes. Differential settlement patterns and persistent recruitment between sponge

species and sizes has been recorded for another sponge-dwelling goby species, *Elacatinus*

lori (Majoris et al. 2018). The study conducted by Majoris et al. (2018) indicated that habitat

preferences in sponge-dwelling gobies are persistent between generations and geographical

area, and there is likely a genetic basis for settlement preference. Little is known about the

movement of sponge-dwelling gobies post-recruitment (Majoris et al. 2018). Habitat qualities

such as food availability and parasite density may influence the settlement of gobies;

however, parasite density has not been measured between sponge species or size classes

(D'Aloia et al. 2011). Such habitat preferences have been shown for sponge-dwelling

cleaning gobies that preferentially maintain territories on barrel sponges with higher parasite

densities (Whiteman and Côté 2004b). The movement of these gobies is critical for

understanding the relationships between sponges and gobies.

Applications to monitoring reef health

As active suspension-feeders, sponges can impact overall nutrient availability in the water

column and, in turn, impact reef health overall (Lesser 2006). As such, sponge health should

be considered when evaluating reef health and plans for monitoring reefs. With the

relationship between *E. horsti* and *Aplysina* sp. resulting in greater pump rate, goby presence

may be an indicator of sponge health. The availability of sponge microhabitat can also

provide insights into sponge-dwelling goby distributions and goby population health

(D'Aloia et al. 2011; Majoris et al. 2018). Taking into account these interspecific

relationships can help to make reef monitoring more comprehensive (Bell et al. 2017). With

current climate trends, accounting for factors that may impact the competitive relationship

between corals and sponges may also help scientists to understand and predict future shifts

towards or away from sponge-dominated reefs (Bell et al. 2017, 2018).

Acknowledgements

We would like to thank the Council on International Educational Exchange (CIEE) Research

Station Bonaire for support; CIEE station staff and students for dive support and helpful

comments; and E. Siemann for statistical advice and comments on this manuscript.

Data availability

All data are available from the corresponding author upon reasonable request.

Declarations

Conflict of interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

References

Bell J, Rovellini A, Davy S, W. Taylor M, Fulton E, R. Dunn M, Bennett H, M. Kandler N,
Luter H, Webster N (2018) Climate change alterations to ecosystem dominance: How
might sponge-dominated reefs function?

Bell JJ (2008) The functional roles of marine sponges. Estuar Coast Shelf Sci 79:341–353

Bell JJ, Biggerstaff A, Bates T, Bennett H, Marlow J, McGrath E, Shaffer M (2017) Sponge
monitoring: moving beyond diversity and abundance measures. Ecol Indic 78:470–488

Burns D, Bingham B (2002) Epibiotic sponges on the scallops *Chlamys hastata* and *Chlamys*
rubida: increased survival in a high-sediment environment. J Mar Biol Assoc United
Kingdom 82:961–966

Böhlke JE, Robins CR (1969) Western Atlantic sponge-dwelling gobies of the genus

Evermannichthys: their taxonomy, habits and relationships. Proc Acad Nat Sci
Philadelphia 121:1–24

Colin P (1975) The neon gobies: the comparitive biology of the gobies of genus *Gobiosoma*,
subgenus *Elacatinus*, (Pisces:Gobiidae) in the tropical western north Atlantic Ocean.
TFH Publications, Neptune City

D'Aloia CC, Majoris JE, Buston PM (2011) Predictors of the distribution and abundance of a
tube sponge and its resident goby. Coral Reefs 30:777

Gerrodette T, Flechsig AO (1979) Sediment-induced reduction in the pumping rate of the
tropical sponge *Verongia lacunosa*. Mar Biol 55:103–110

Hendler G (1984) The association of *Ophiothrix lineata* and *Callyspongia vaginalis*: a
brittlestar-sponge cleaning symbiosis? Mar Ecol 5:9–27

Humann P, DeLoach N (1992) Reef creature identification: Florida, Caribbean, Bahama.
New World Publications, Inc., Jacksonville

Lattig P, Martin D (2011) Sponge-associated *Haplosyllis* (Polychaeta: Syllidae: Syllinae)
from the Caribbean Sea, with the description of four new species. Sci Mar 75:733–758

Lesser MP (2006) Benthic–pelagic coupling on coral reefs: feeding and growth of Caribbean
sponges. J Exp Mar Bio Ecol 328:277–288

Magnino G, Gaino E (1998) *Haplosyllis spongicola* (GRUBE) (Polychaeta, Syllidae)
associated with two species of sponges from East Africa (Tanzania, Indian Ocean). Mar
Ecol Della Stn Zool Di Napoli I 19:77–87

Majoris JE, D'Aloia CC, Francis RK, Buston PM (2018) Differential persistence favors
habitat preferences that determine the distribution of a reef fish. Behav Ecol 29:429–439

McMurray SE, Pawlik JR, Finelli CM (2014) Trait-mediated ecosystem impacts: How
morphology and size affect pumping rates of the Caribbean giant barrel sponge. Aquat
Biol 23:

Meroz E, Ilan M (1995) Cohabitation of a coral reef sponge and a colonial scyphozoan. Mar
Biol 124:453–459

Morganti TM, Ribes M, Yahel G, Coma R (2019) Size Is the Major Determinant of Pumping
Rates in Marine Sponges. Front Physiol 10:

Savarese M, Patterson MR, Chernykh VI, Fialkov V a (1997) Trophic effects of sponge

feeding within Lake Baikal's littoral zone .1. In situ pumping rates. Limnol Oceanogr
42:171–178

Tyler JC, Böhlke JE (1972) Records of sponge-dwelling fishes, primary of the Caribbean.
Bull Mar Sci 22:601–642

Villamizar E, Laughlin RA (1991) Fauna associated with the sponges Aplysina archeri and
Aplysina lacunosa in a coral reef of the Archipiélago de Los Roques, National Park,
Venezuela. In: Reitner J., Keupp H. (eds) Fossil and recent sponges. Springer Berlin
Heidelberg, Berlin, Heidelberg, pp 522–542

Whiteman EA, Côté IM (2002) Cleaning activity of two Caribbean cleaning gobies: intra-
and interspecific comparisons. J Fish Biol 60:1443–1458

Whiteman EA, Côté IM (2004a) Individual differences in microhabitat use in a Caribbean
cleaning goby: a buffer effect in a marine species? J Anim Ecol 73:831–840

Whiteman EA, Côté IM (2004b) Dominance hierarchies in group-living cleaning gobies:
causes and foraging consequences. Anim Behav 67:239–247

Wulff JL (2006) Ecological interactions of marine sponges. Can J Zool 84:146–166

Yahel G, Marie D, Genin A (2005) InEx - a direct in situ method to measure filtration rates,
nutrition and metabolism of active suspension feeders. Limnol Oceanogr Methods 3:46–
58

Figure legends

Fig. 1 The dependence of average pump rate on tube length and goby presence for a)

Aplysina lacunosa and b) *Aplysina archeri*. Black symbols and lines indicate a sponge tube

with a *Elacatinus horsti* . Gray symbols and lines indicate a sponge tube with no *E. horsti*.

Fig. 2 The percent of *Aplysina lacunosa* (n = 226) and *Aplysina archeri* (n = 85) sponge

tubes with *Elacatinus horsti* (black bar) or no *E. horsti* (gray bar).

Goby present | Gobies absent

Percent of sponge tubes