1 **Image recognition based on deep learning in *Haemonchus contortus* motility**

2 **assays**

Martin Žofka, Linh Thuy Nguyen, Eva Mašátová, Petra Matoušková*

Department of Biochemical Sciences, Faculty of Pharmacy, Charles University, Heyrovského

1203, 500 05 Hradec Králové, Czech Republic.

Correspondence:

*Corresponding Author

matousp7@faf.cuni.cz

12 Abstract

Poor efficacy of some anthelmintics and rising concerns about the widespread drug resistance

have highlighted the need for new drug discovery. The parasitic nematode *Haemonchus*

contortus is an important model organism widely used for studies of drug resistance and drug

screening with the current gold standard being the motility assay. We applied a deep learning

approach Mask R-CNN for analysing motility videos containing varying rates of motile worms

and compared it to other commonly used algorithms with different levels of complexity, namely

the Wiggle Index and the Wide Field-of-View Nematode Tracking Platform. Mask R-CNN

consistently outperformed the other algorithms in terms of the detection of worms as well as

the precision of motility forecasts, having a mean absolute percentage error of 7.6% and a mean

absolute error of 5.6% for the detection and motility forecasts, respectively. Using Mask R-

CNN for motility assays confirmed the common problem with algorithms that use non-

maximum suppression in detecting overlapping objects, which negatively impacts the overall

precision. The use of intersect over union as a measure of the classification of motile / non-

motile instances had an overall accuracy of 89%, indicating that it is a viable alternative to previously used methods based on movement characteristics, such as body bends. In comparison to the existing methods evaluated here, Mask R-CNN performed better and we anticipate that this method will broaden the number of possible approaches to video analysis of worm motility.

Keywords: parasite; nematode; instance segmentation; object detection; Mask R-CNN

1. Introduction

Parasitic nematode infections present a substantial problem to human and veterinary medicine. Anthelmintics are often used as a weapon of choice to control or eliminate parasitic worms. With increasing reports of drug resistance, there is an ongoing effort to develop new anthelmintics [1, 2]. In view of this, drug screening and drug testing methods are clearly necessary. A number of important human and animal nematodes were already studied in large chemical library-scale screenings [3, 4], although nearly all of the studies were aimed at well-established non-parasitic *Caenorhabditis elegans* because of the scientific interest and medical applications as potential human disease models [5-9].

Nowadays, *in vitro* assays can measure the effects of compounds on development, growth, behaviour, and motility. Several approaches have been pursued in attempts to deliver robust, automated assays on the nematode phenotype [10-14]. The current gold standard for measuring drug effectiveness is *in vitro* assessment of worm motility, as measured via microscopy. The automated phenotyping from short video recordings offers a useful assay for high-throughput whole-organism phenotypic screening and avoids manual scoring. In assays with parasitic worms under experimental conditions, several challenges need to be overcome to enhance the performance of motility measurements. These challenges are associated with the

different worm lengths, their tendency to clump, and also the range of diverse, complex
movement patterns different from that seen in *C. elegans*.

Focusing on veterinary importance, the parasitic nematode *Haemonchus contortus*,
known as the barber's pole worm, is a model organism often used for drug screening and
studying drug resistance; due to its economic importance, known genome and transcriptome
[15, 16]. *H. contortus* infects predominantly small ruminants. It has a direct life cycle that starts
with eggs being shed onto a pasture in the faeces of the infected animals. The cycle continues
through three free-living larval stages to the parasitic fourth larval stage and lastly adults, which
reside in the abomasum where they feed on blood. The ability to maintain *H. contortus* infection
in a natural host animal and produce different larval stages *in vitro* provides experimental
advantage. The infective third-stage larvae (L3s) can be stored, and researchers commonly use
L3s or *in vitro* fourth larval stage for motility studies. Many automated or semi-automated
approaches to measure motility have already been described such as via electrical impedance
[17] or infrared light beam-interference [7]. Nevertheless, in our study of analysing the video
recordings in *H. contortus* larvae to assess the motility, we have compared various
(representative) algorithms with different levels of complexity in terms of model preparation as
well as computational cost. We were interested to assess if the additional complexity of the
methods leads to a sufficient increase in precision, that would justify the time-performance
trade-off of these methods.

The first one mentioned is an algorithm which calculates a motility index value based
on measuring the standard deviation of the pixel's light intensity averaged for a number of
frames [18]. This Wiggle Index (WI) does not allow for the detection of the phenotype of
individual worms; still, this assay led to many "hit compounds" in screening of a large variety
of compound libraries [19, 20]. The second approach called Wide Field-of-View Nematode
Tracking Platform (WF-NTP) is built on the use of the Gaussian adaptive threshold of the image

frames to identify worms and was applied in model free-living nematode *C. elegans* [21]. It

was initially developed for drug screening purposes; nevertheless, it has also found applications

in other studies such as detailed genetic or behavioural studies. Nonetheless, WF-NTP has a

major limitation for images with overlapping objects as these objects get discarded by the

algorithm based on an upper pixel size threshold per object, resulting in underestimation of the

the total number of detected objects in an image [21]. In our study, we customized the associated

WF-NTP software written in Python and compared it to the last approach which involves the

application of deep learning.

The advent of machine learning has revolutionized the surveying and classifying of

biological data including image recognition, enabling the automation of many tasks. The most

important aspect of large-scale computerization is the possibility to automate *in vitro* assays

and scale them, allowing researchers to avoid tedious repetitive tasks and focus on activities

where they can use their knowledge to provide added value, leaving the analysis and

interpretation of the data as the only rate-limiting step. In the field of computer vision, deep

learning algorithms with a convolutional neural network (CNN) architecture have made rapid

advancements on a variety of image classification tasks in cell cultures [22], and the model

organism *C. elegans* [23, 24], even parasitic nematode in plants [25], yet none of them have

been applied to parasitic gastro-intestinal nematodes of livestock, including *H. contortus*.

Object-detection-based deep learning methods have also been adapted for instance

segmentation. Here we used a state-of-the-art region-based CNN called Mask R-CNN [26],

which can be trained based on a set of annotated images (Fig 1a) to detect the objects of interest,

in our case the larval stages of *H. contortus*. In the first step, the algorithm uses a regional

proposal network (RPN), i.e., a binary classifier that identifies rectangular subsets of the image

that are of interest for the given task and are called regions of interest (ROI). In our case that

would be parts of the image that potentially contain worms. These ROIs are then refined to

cover the whole object (Fig 1b) and filtered using non-maximum suppression (NMS). NMS is

a process of removing highly overlapping ROIs, to avoid detecting the same object multiple

times before passing the filtered ROIs as proposals for classification. The classifier performs a

refinement of the ROI and outputs a final rectangle surrounding the detected object called a

bounding box, the class of the predicted instance along with the detection confidence and the

object mask, which are the pixels that make up the object (Fig 1c).

The aim of the study was to train a Mask R-CNN model for image recognition of

different life cycle stages of *H. contortus* and evaluate the application of this deep learning

algorithm to the field of drug discovery of anthelmintics. Thereby, we compared the Mask R-

CNN detections and motility predictions to the WI and WF-NTP methods.. Additionally for the

motility, we tested a metric called intersection over union (IoU) instead of body bends from

WF-NTP readout. IoU is used regularly in image recognition to quantify the extent of overlap

between two objects, the intersection of the objects is divided by the union of the objects, which

results in a numerical value between 0 and 1 (see Graphical abstract). Furthermore, we

discussed the possible application of automated Mask R-CNN in other assays related to the

veterinary practice.

Fig 1: Mask R-CNN training and detection process **(a)** annotated images used for training the

algorithm, **(b)** regions of interest (ROI) detected in the first step by the regional proposal

network (solid boxes) and the refined regions (dashed boxes), **(c)** Mask R-CNN predictions

containing the object boundaries, predicted object classes (i.e., L3 – third-stage larva) with an

associated confidence level and the mask of the object (coloured contour).

2. Materials and methods

2.1. **Procurement of L3s of *H. contortus***

Six-month-old sheep were orally infected with 8,000 infective L3s of MHco3 strain (Inbred Susceptible Edinburgh, ISE) [27]. Four weeks post-infection faecal samples were collected. L3s were produced from eggs by incubating faeces in a plastic box covered with foil at 27 °C for 7 days. Then, the faeces were rinsed twice in tap water which was poured into 1,000mL conical measuring cups, in which the larvae sank to bottom. To remove dirt or dead individuals, the pellet of larvae was pipetted out and filtered through a 20μm sieve submerged in water (27 °C). Clean L3s at a concentration of approximately 4,000 L3 per mL were stored in culture flasks in water at 10 °C for several months.

During the storage, some larvae naturally die. Prior to the experiment, we filtered larvae through a 20μm sieve (for 12 h), this time collecting both live larvae from the bottom of the sieve and dead individuals remaining on the sieve.

All experimental procedures were examined and approved by the Ethics Committee of the Ministry of Education, Youth and Sports (Protocol MSMT-25908/2014-9).

2.2. Video processing and manual counting

Videos from a microscope camera were recorded for a duration of 10 seconds with a framerate of 20 frames per second (fps) with a pixel quality of 2560 by 2160 (see Supplementary Video). The videos (.avi with MJPEG compression, ~ 0.45GB) were acquired using NIS-Elements Imaging Software (version 4.20) using a Nikon Eclipse Ti microscope (4x magnification) with the camera (Andor Zyla 5.5 sCMOS, 12bit: 2560 x 2160 pixels). The videos were recorded for different ratios of live and dead larvae (100:0, 75:25, 50:50, 25:75, 0:100; denoted as motility group 100, 75, 50, 25 and 0, respectively) in 9 replicates and they contained on average 60 worms. The live larvae were kept at 37 °C to maintain their motility.

The video recordings were processed so that each worm was labelled with a numeric

identifier. A trained human with prior experience of manual motility assays evaluation assessed

each worm from each video and information about motility and the dead phenotype was

recorded. The discrepancies were dealt with as follows (i) a dead worm which was moving due

to the interference of a motile worm was counted as motile; (ii) a live coil non-motile worm

was counted as non-motile; (iii) the worms which were not labelled by an algorithm were given

an identifier manually; (iv) worms were excluded from the counts when they appeared in less

than 20 frames; (v) dead phenotype was recorded when overt microscopical pathologies, such

as the presence of large vacuoles in the tegument and straight phenotype, were observed.

This manual counting was taken as the referential method for comparison of accuracy

among the three models.

2.3. Wiggle Index analysis

We used an ImageJ implementation of the WI for the analysis [28]. We did not apply

any scaling of the videos, the Gaussian image blur was set to 2, the number of frames to be

averaged was set to 50 and the motility output was based on the standard deviation. Since the

values from the WI are relative, we used motility group 100 as the control group to be able to

convert these relative values to percentages. We calculated the mean value for motility group

100 and divided all values by the mean to obtain normalized motility values.

2.4. WF-NTP analysis

WF-NTP presents an automated evaluation of motility in *C. elegans*. For its successful

implementation, we adopted the method for the *H. contortus* motility assay. The algorithm was

based on three steps;in the first instance all worms in each frame were detected, in the second

step, these detections were linked based on a neighbourhood search, and in the final step, each

worm was classified as motile or non-motile based on bends per second and the speed of the

worm. However, as these parasites differ in their sizes as well as their movement, some

modifications to the original configuration and algorithm were necessary. These modifications

belonged predominantly to two categories, the first was a recalibration of the available input

parameters of the WF-NTP method. This included aspects such as the image size, the size of

the worms that we want to detect and other configurations (see Data availability). The other

modifications were changes in the evaluation of the conditions for worm motility and a limit

on the minimum number of frames that a worm needed to be detected in for it to be included in

the evaluation.

The original method relied on bends per second and the speed of individual worms to

classify them as either motile or non-motile [21]. When a worm was below a certain threshold

of bends per second and below a certain speed threshold, it was classified as non-motile. In our

settings, the speed threshold was obtained from the motility group 0. From the initial test for

the motility group 50, the result was that out of 169 worms classified as motile, 23% exceeded

the speed threshold, but no bends were recorded. This failure to measure bends per second for

motile worms, was most likely caused by the more complex movement of *H. contortus.*, This

led us to replace the original approach with an evaluation metric, IoU, that is regularly used in

image recognition [29]. The use of IoU allowed us to use the capabilities of WF-NTP in terms

of the detection and tracking of individual worm instances in an image and then apply the IoU

metric to determine the worm motility. The calculation of IoU was done for the same worm

instance in neighbouring frames to detect changes on a frame-by-frame basis. Afterwards, the

mean was calculated and worms above a certain IoU threshold were labelled as non-motile. To

ensure that the comparison of WF-NTP and Mask R-CNN was aimed at the quality of the

detection of individual worm instances rather than the approach to determine the motility, we

used the same tracking for detections across frames and the same IoU conditions for both

methods.

Another modification made to the original method was the introduction of a threshold

for the minimum number of frames that a worm needed to be detected in, in order to be included

in the analysis. The reason for this restriction was that as the individual worms move, they have

a tendency to change shape or overlap with other worms, therefore in terms of the detection the

individual worm can be lost for a certain number of frames and reappear later, however this can

lead to counting some instances multiple times. To limit these potential duplicates

(Supplementary Fig S1) as much as possible and not bias the total number of worms detected

in the video, we selected the highest number of worms detected in a given video across all

frames and added a buffer of 10% in order to account for situations where some worms were

not detected, for example due to their overlaps. This maximum number then served as a limit

for selecting the number of worms to be evaluated as motile or non-motile. The instances were

selected based on the number of frames in a decreasing order, therefore ensuring that we had

data from enough frames to classify the worm correctly. Again, this restriction was applied to

detections from Mask R-CNN to enable a comparison under the same conditions.

2.5. Mask R-CNN analysis

For Mask R-CNN we used the Matterport implementation

(https://github.com/matterport/Mask_RCNN) of the algorithm, including a trained model using

a backbone architecture of Resnet101. The pre-trained model weights were obtained from

training the model on the MS COCO dataset [30]. The process of using a pre-trained model is

called Transfer Learning, which is an approach where the knowledge gained from solving one

problem is applied to a related problem. The motivation for selecting a pre-trained model is that

in general deep neural networks require a lot of annotated data for training and using an existing

model and re-training the top layer of the network allows us to have a smaller dataset while

maintaining good performance. We manually annotated a total of 95 images and divided them

into training (69) and validation (26) using the VGG (Visual Geometry Group) Image

Annotator [31], these images contained three categories of objects; eggs, first-stage larvae (L1),

and L3, to fully utilise the capabilities of Mask R-CNN which is able to do instance

segmentation and classification. The total number of annotated objects was around 10,400

divided into training (6,800 objects) and validation (3,600 objects). Then we ran the training on

a GPU instance until the loss function showed no signs of improvement. To avoid overfitting,

we used data augmentation and applied one or multiple of the following augmentations:

flipping, contrast normalization, additive gaussian noise, multiplying the pixels by a number

within a given interval to make the whole image lighter or darker. The final model had a mean

average precision at 0.5 IoU of 64.1% on the validation set.

The trained model was then used on the individual frames of the video to detect the

instances of parasites and we calculated the centroid of the detected object in terms of the

position on the x and y axis using the scikit-image regionProps functionality [32]. These

detected instances were then linked together using the trackpy library [33], which detected the

trajectories of objects based on certain conditions. In our case we restricted the distance that the

centroid could move a maximum of 100 pixels in each dimension. As a result of detecting the

trajectories, we were able to identify the same parasite instance across multiple frames and

assign it with a unique identifier. In cases where the tracking did not meet the above conditions,

a new identifier was assigned, and therefore a single worm could be detected multiple times,

which occurred for complex cases with extensive movements and a high number of overlaps.

Subsequently, we calculated the IoU and the threshold for the number of parasites to select for

the evaluation as described in the WF-NTP section. We observed a decreasing mean percentage

error (MPE) with an increase in the length of the video, and therefore we decided on using the

full length of the videos for the motility assay (Supplementary Fig S2).

2.6. Error metrics

For the evaluation of the performance of the individual algorithms, we used standard

metrics for the quantification of the forecast precision. These metrics work with the error term,

which is defined as the difference between the actual and the forecasted value.In our case, the

actual value was obtained by manually processing the videos. Two aspects were measured; the

first one was the bias of the forecasts to determine if the algorithm is systematically over or

under-estimating, for that purpose, we used mean error (ME) or mean percentage error (MPE),

which both work with the mean of the error term. The second aspect was the size of the error

term irrespective of the error term being positive or negative to compare the overall

performance of the algorithms. For that purpose, we used mean absolute error (MAE) or mean

absolute percentage error (MAPE), which both work with the mean absolute value of the error

term.

2.7. Monte Carlo drug screening simulation

To assess the impact of the differences in detection accuracy we ran a Monte Carlo drug

screening simulation [34] for all three algorithms. To remove the effect of individual motility

groups we worked with the differences between the predicted motility rates and the manually

obtained motility rates. Using these differences we were able to obtain the correlations between

the algorithms as well as the means and standard deviations. A multivariate normal distribution

was used as the data generation process for the simulation to ensure that the correlation among

the algorithms was taken into account. The SciPy package (version 1.3.3) implementation of

the multivariate normal distribution was used.

For the simulation, we generated 1 million random motility rates from a continuous uniform

distribution within the range of zero to one. Similarly, we generated 1 million observations of

the multivariate normal distribution. These observations were the detection errors of each of

the algorithms. By adding this error to the randomly generated motility rate, which served as

our true observation, we received simulated motility predictions from the three algorithms. A

"hit" in our simulation was defined as having a motility rate lower than 50%.

3. Results

3.1. Detection of motility

The observed typical cases for detection scenarios in the motility videos were depicted in Fig

2, where the results of Mask R-CNN were used for illustrating these examples. The most

common situation, which did not pose a problem to the algorithms, were worms with few or no

overlaps (Fig 2a). Other common occurrences were collisions between worms that caused a

non-motile worm to be moved and therefore being perceived by the algorithms as motile (Fig

2b). Lastly, the most challenging scenarios contained a high number of worms with significant

overlaps with mostly motile worms, which resulted in problematic detections and negatively

impacted the ability to track worms across multiple frames (Fig 2c).

Fig 2: Sample frames of common detection scenarios in motility videos. The blue outlines are

the masks detected by Mask R-CNN. Each detected worm has an associated identification

number displayed in white based on the tracking across the frames. **(a) c**ommon scenario with

low or no overlaps of individual worms. **(b) e**xample of a motile worm (id 32) colliding with a

non-motile worm (id 26) causing the later to be perceived as motile. **(c) c**omplex detection

scenario with a high number of worms with significant overlaps

3.2. Motility forecast errors

In the manual video processing, we observed the motility and the live rate. To measure the performance of the algorithms, we used the motility rate rather than the live rate even though the stillness alone does not denote a dead worm. The reason for this choice was that the algorithms detect movement and are not able to differentiate if the movement was caused by the object itself or by an interaction with another object. The general manual motility rate was always higher than the manual live rate (Supplementary Table S1). We measured the performance of the individual algorithms by evaluating the errors, defined as the differences between the manual processing of the file and the respective algorithm. Out of the three methods, Mask R-CNN had the lowest MAE with a value of 5.6%, followed by WF-NTP (8.76%) and the WI (14.2%). In terms of the bias of the predictions, the WI had the lowest ME of -0.71%, therefore slightly overestimating the motility rate on average, while both WF-NTP and Mask R-CNN underestimated the motility rate with values of 6.52% and 1.95% respectively (Table 1).

Shown in Fig 3, the WI had an increasing volatility with an increase in the ratio of alive worms in the group. Mask R-CNN overestimated the motility rate for the motility group 0; while four videos had a zero motility rate, showing that the overestimation is not systematic, two videos had a motility rate error below 3.6%, and the remaining three videos had high motility rates without any apparent reason. WF-NTP and Mask R-CNN had a tendency to systematically underestimate for the motility group 100 (Fig 3). The mean and standard deviation of motility rates per motility group and algorithm is shown also in Supplementary Table S2 and the same statistics are reported as differences between the manual count and the respective algorithm in Supplementary Table S3.

Table 1: Motility error metrics for algorithms

Algorithm	MAE, %	ME, %
Wiggle Index	14.2	-0.71
WF-NTP	8.76	6.52
Mask R-CNN	5.61	1.95

The differences between the manual processing and the respective algorithm were expressed as the mean absolute error (MAE) and mean error (ME).

Fig 3: Motility rates comparisons for individual motility groups per algorithm. The number of the motility group indicates the percentage of live larvae. The motility rate is visualized as box plots.

3.3. Instance detection performance

The motility rates are an indicator of the degree of correctly classified worms as motile or non-motile based on IoU. However, they do not provide the information of the number of detected worm instances in the image. In order to ensure that the motility rates were estimated reliably and not only based on a small subset of detected worms, the detection performance was validated. To achieve this, we compared Mask R-CNN and WF-NTP for the total worm count (WI was excluded as the algorithm does not detect individual instances of worms, therefore this information was not available). The precision was measured based upon the error term between the manual counts and the respective algorithm. In terms of the size of the error, the MAPE was used and we observed that the error was significantly lower ($P < 0.0001$, Wilcoxon signed-rank test) for Mask R-CNN (7.6%) than for WF-NTP (40.23%). Both algorithms had a tendency to underestimate the number of worms on average, therefore not detecting all the worm instances

in a video, mainly due to the challenges of detecting overlapping or clustered worms. This was confirmed by the MPE with values of 5.61% for Mask R-CNN and 40.23% for WF-NTP (Table 2). The reported MAPE and MPE for WF-NTP had the same value because all the detections underestimated the total number of worms. WF-NTP underestimated the worm count for all motility groups, while Mask R-CNN had similar box plots in comparison to the manual counts apart from underestimating the motility group 0, as shown in Fig 4. The mean error and standard deviation of the differences between the manual count and the respective motility algorithm is shown in Supplementary Table S4.

Table 2: Worm count error metrics for algorithms

Algorithm / Metric	MAPE, %	MPE, %
WF-NTP	40.23	40.23
Mask R-CNN	7.6	5.61

The differences between the manual counts and the respective algorithm were expressed as the mean absolute percentage error (MAPE) and mean percentage error (MPE).

Fig 4: Worm count comparisons for individual motility groups per algorithm. The number of the motility group indicates the percentage of live larvae. The number of worms is visualized in box plots.

3.4. Mask R-CNN detection and classification precision

The precision of detecting worm instances in a video was on average 98.6%. This was calculated based on all the tracked worms across all the motility groups without setting a minimum number of frames for which a worm needed to be tracked. The detections were

manually validated and the detection was considered correct despite only a part of the worm

being detected in some frames. The majority of cases where the detection was incorrect was

due to debris that had a similar size and shape as a worm. For the detected worms across all the

motility groups, there was an overall accuracy of 89% to correctly label them as either motile

or non-motile based on their mean IoU values. In Fig 5, the distribution for different IoU values

obtained from all the worms across all the motility groups clearly showed that most of

misclassified cases were located around the 0.8 threshold, while the classification accuracy

increased for values further away from the threshold. The precision and recall values (Table 3)

confirm that the prevalent case was the misclassification of motile worms as non-motile.

The errors defined as differences between Mask R-CNN and the manually evaluated

motility rates were tested for normality using the D'Agostino-Pearson test as well as Shapiro-

Wilk test. In both cases we could not reject the null hypothesis that the data is from a normal

distribution with p-values 0.85 and 0.35 respectively (Table 4).

Fig 5: The distribution for different Intersection over Union (IoU). Both **(a) a** histogram of IoU

and the associated number of worms per bin classified into motile / non-motile and **(b)** a

histogram of IoU and the associated probability per class (motile / non-motile) show that the

majority of misclassified cases were located around the 0.8 IoU.

Table 3: Classification performance metrics

Classification	Precision, %	Recall, %
Motile	94	88
Non-Motile	84	92

385 Precision and recall metrics for classification performance

Table 4: Normality tests for the motility rate error terms between Mask R-CNN and the manual processing

Normality test	Statistic	p-value
D'Agostino-Pearson	0.32	0.85
Shapiro-Wilk	0.97	0.35

For both normality tests we cannot reject the null hypothesis that the data is from a normal distribution.

3.5. Monte Carlo drug screening simulation

The Monte Carlo simulation was based on the results from the motility forecasts and the obtained forecast errors. While WF-NTP and Mask R-CNN had a positive correlation of forecast errors (0.65), the forecast errors for WI were not correlated with the other algorithms (Table 5).

Table 5: Correlation matrix for the prediction errors of the individual algorithms

	Wiggle Index	WF-NTP	Mask R-CNN
Wiggle Index	1		
WF-NTP	-0.04	1	
Mask R-CNN	0.08	0.65	1

Mask R-CNN achieved the highest accuracy in the simulation with a value of 94% followed by WF-NTP 90% and the WI 85% (full classification performance in Supplementary Table S5). The simulation estimated that an improved detection performance of 3.15% (MAE) between Mask R-CNN and WF-NTP led to a 3.5% improvement of drug screening accuracy, while an improvement of 8.59% (difference of MAE between Mask R-CNN and WI) led to a 9% improvement of drug screening accuracy.

4. Discussion

The motility phenotype is often used for drug screening and drug testing of anthelmintics. In our study, we deployed and compared three algorithms to measure motility in the parasitic nematode *H. contortus*. Apart from evaluating motility for both WF-NTP and Mask R-CNN, we were also interested in the detection precision of worms by these algorithms. The reason for that was the fact that for both algorithms, the detection was the first step followed by the tracking across the frames and the classification into motile and non-motile. Therefore, accurate detection was the pre-requisite for reliable motility quantification.

The results showed that the performance increased with an increased complexity of algorithm. The WI, despite requiring minimal configuration and being the fastest algorithm, had some limitations in comparison to the other methods; it was by design not able to detect individual instances of worms and a control group with only motile worms needed to be created in order to be able to translate the WI into a percentage for the motility rate for each video, otherwise it was hard to interpret the numerical value of the WI. Because of these limitations and the lack of precision, it would be advisable to use WI only for initial studies to quickly test experimental results.

For WF-NTP, overlapping parasites had the highest negative impact on the detection ability, as the algorithm filters out objects that are above or below a certain pixel size. This resulted in detecting overlapping worms as a single object and these objects were then excluded based on the size threshold. As a consequence, the total number of detected instances in an image was significantly lower than the actual number obtained by manually processing the videos. Although WF-NTP struggled with detecting all instances, the classification of motile / non-motile performed well once an instance was detected.

Mask R-CNN had the lowest MAE. However, similarly to WF-NTP, it struggled with overlapping objects, primarily due to the fact that the algorithm uses NMS. As a result, some

instances were not detected, although in contrast to WF-NTP, the worm clusters were not

discarded as a whole. The problem of detecting overlapping objects is currently an area of active

research in object detection, particularly for biological data. For instance, Böhm et al. [35] are

working on algorithms that can handle overlapping objects and have had success with applying

their algorithm to *C. elegans*. Motile worms had a higher chance of being detected, because

they had higher variability across the frames due to their movement. In terms of precision, the

algorithm was slightly biased towards underestimating the motility rate; however, the errors

were normally distributed enabling us to increase the number of acquired videos in order to get

to a desired level of confidence and therefore successfully replace manual processing.

The results confirmed that IoU could be used as a metric for determining the motility of

larvae, although some modifications might be required in order to deal with slow-moving

worms, such as including the total distance travelled during the video to complement the frame-

by-frame information. Furthermore, the IoU threshold should be considered a parameter to be

optimized as it can vary based on the age of the larvae, the temperature of the environment and

other factors. The downside of using Mask R-CNN is that it requires a quite big amount of

annotated data (provided in the Data availability), which is time consuming to create and the

training runs for 1-2 days on a GPU instance (an Amazon Web Services p2.xlarge compute

instance with 1 GPU, 4 vCPUs and 61GB RAM was used).

The drug screening simulation showed that the improved detection performance

resulted in increased accuracy for drug screening of roughly the same magnitude as the MAE

improvement, illustrating that the improvement is not negligible for practical purposes even

though some of the forecast errors of the algorithms were correlated. For applications with

multiple observations for a given category / group, the impact would be most likely lower,

because with an increase in the number of observations per category / group, the sample mean

of the forecast errors would converge to the expected value of the forecast errors. In this

situation, the bias of the estimate measured in our case by ME would then have a higher impact

than the variance which would make WF-NTP, with the highest bias, the least accurate

algorithm.

In general, for quick low-precision experiments, such as primary screening of large

compound libraries, the best option is to use the WI as it can be used out-of-the-box with slight

modifications to a few parameters. If the detection of individual instances is desired, Mask R-

CNN is the best choice, although it is more time-intensive to prepare the data and train the

model. Moreover, the ability to precisely detect larvae in an image then allows the researchers

to perform multiple types of analyses that involve microscopy of the parasites and are not only

limited to motility assays, but also include other common *in vitro* assays for the diagnosis of

drug resistance, e.g. larval development tests and egg hatch tests [36]. In the aspect of

potentially repurposing the algorithms for other assays, Mask R-CNN is the most flexible of

the models, because it is able to detect multiple classes of objects. On contrary, WF-NTP

decides solely on the size of the objects, making a distinction between classes of similar sizes

impossible.

5. Conclusions

Our study confirmed that the state-of-the-art machine learning algorithm Mask R-CNN was

able to outperform less computationally complex algorithms that are based on pixel intensity

or Gaussian threshold. The algorithm performed better in the detection of individual worm

instances as well as the subsequent motility classification. This was consistent with how the

algorithm is designed as the tracking and motility classification depend on the detection

accuracy. For the motility classification, we used IoU as a new metric, that successfully

replaced previous ones based on locomotion specific to the studied organisms. Whilst IoU is

very flexible, as it only requires a single parameter, it would need further extension to be able

to deal with cases of heterogeneous groups of viable worms that have a high variance in their

speed of movement.

The gain in precision of Mask R-CNN came at the cost of requiring an extensive annotated

dataset as well as computational resources for training the model. Therefore, the time-precision

trade-off needs to be evaluated by the researchers on a case-by-case basis, unless the study

allows reapplying an already existing trained model. The output of the present study provides

an annotated dataset containing various developmental stages of *H. contortus* that can be

utilised by researchers for future machine learning applications.

Data availability

491 All underlying data for training the model, the trained model itself as well as the input and

492 output of the motility analysis for all three algorithms is available at

493 https://doi.org/10.5281/zenodo.5734143. The source code for Mask R-CNN is available in a

494 repository on GitHub https://github.com/zofkam/mask_rcnn_motility. We have also used

495 Zenodo to assign a DOI to the repository https://doi.org/10.5281/zenodo.5735068.

Additional files

Supplementary Table S1: Manual counts of motility and live rates per motility group

Supplementary Table S2: Motility rates per motility group and algorithm

Supplementary Table S3: Mean error between the manual count and the respective motility

algorithm

Supplementary Table S4: Mean of differences in worm counts between the manual processing

and the respective algorithm

Supplementary Table S5: Drug screening simulation classification performance

Supplementary Fig S1: Average number of worms in the videos per motility group for WF-NTP and Mask R-CNN.

Supplementary Fig S2: The impact of the length of the video in frames on the percentage error of the detected worms for Mask R-CNN.

Supplementary Video: Mask R-CNN sample detection video of motility group 50.

Abbreviations

CNN: convolutional neural network; fps: frames per second; GPU: graphics processing unit; IoU: intersection over union; L3: third-stage larva; MAE: mean absolute error; mAP: mean average precision; MAPE: mean absolute percentage error; Mask R-CNN: region based convolutional neural network; ME: mean error; MPE: mean percentage error; NMS: non-maximum suppression; ROI: region of interest; RPN: regional proposal network; WF-NTP: Wide Field-of-View Nematode Tracking Platform; WI: Wiggle Index

Competing Interests

The authors declare that they have no competing interests.

Funding

This work was supported by Charles University UNCE/18/SCI/012 and SVV 260 550 as well as by the project EFSA-CDN [CZ.02.1.01/0.0/0.0/16_019/0000841], co-funded by ERDF.

Authors' contributions

M.Ž. and L.T.N. designed this study and wrote the manuscript. M.Ž. implemented the deep learning model, wrote the code, ran the statistical analyses and generated the figures. L.T.N. created the graphical abstract. L.T.N., E.M. acquired and annotated the images used to train the

530 deep learning model. M.Ž. and L.T.N. conducted the experiments and analyzed the results. P.M.

531 managed the project. All authors contributed to literature review and critical revision of the

532 manuscript.

Acknowledgements

535 We thank MVDr. Tomáš Kaňka for assisting with the sheep infection. The authors thank prof.

536 Lenka Skálová for critical discussion and comments. We would like to express our gratitude to

537 Jon Nolan for proofreading and revisions.

References

[1] Jiao Y, Preston S, Hofmann A, Taki A, Baell J, Chang BCH, et al. A perspective on the
discovery of selected compounds with anthelmintic activity against the barber's pole worm-
Where to from here? Adv Parasitol. 2020;108:1-45. doi: 10.1016/bs.apar.2019.12.003.
[2] Geary TG, Sakanari JA, Caffrey CR. Anthelmintic Drug Discovery: Into the Future. Journal
of Parasitology. 2015;101(2):125-33. doi: 10.1645/14-703.1.
[3] Partridge FA, Murphy EA, Willis NJ, Bataille CJ, Forman R, Heyer-Chauhan N, et al.
Dihydrobenz[e][1,4]oxazepin-2(3H)-ones, a new anthelmintic chemotype immobilising
whipworm and reducing infectivity in vivo. PLoS Negl Trop Dis. 2017;11(2):e0005359. doi:
10.1371/journal.pntd.0005359.
[4] Abriola L, Hoyer D, Caffrey CR, Williams DL, Yoshino TP, Vermeire JJ. Development and
optimization of a high-throughput screening method utilizing *Ancylostoma ceylanicum* egg
hatching to identify novel anthelmintics. PLoS One. 2019;14(6):e0217019. doi:
10.1371/journal.pone.0217019.
[5] Buckingham SD, Partridge FA, Sattelle DB. Automated, high-throughput, motility analysis
in *Caenorhabditis elegans* and parasitic nematodes: Applications in the search for new
anthelmintics. Int J Parasitol Drugs Drug Resist. 2014;4(3):226-32. doi:
10.1016/j.ijpddr.2014.10.004.
[6] Burns AR, Luciani GM, Musso G, Bagg R, Yeo M, Zhang Y, et al. *Caenorhabditis elegans*
is a useful model for anthelmintic discovery. Nat Commun. 2015;6. doi: 10.1038/ncomms8485.
[7] Taki AC, Byrne JJ, Wang T, Sleebs BE, Nguyen N, Hall RS, et al. High-Throughput
Phenotypic Assay to Screen for Anthelmintic Activity on *Haemonchus contortus*.
Pharmaceuticals (Basel). 2021;14(7). doi: 10.3390/ph14070616.
[8] Elfawal MA, Savinov SN, Aroian RV. Drug Screening for Discovery of Broad-spectrum
Agents for Soil-transmitted Nematodes. Scientific Reports. 2019;9. doi: 10.1038/s41598-019-
48720-1.
[9] Pulak R. Techniques for analysis, sorting, and dispensing of *C. elegans* on the COPAS flow-
sorting system. Methods Mol Biol. 2006;351:275-86. doi: 10.1385/1-59745-151-7:275.
[10] Ramot D, Johnson BE, Berry TL, Carnell L, Goodman MB. The Parallel Worm Tracker:
A Platform for Measuring Average Speed and Drug-Induced Paralysis in Nematodes. PLoS
One. 2008;3(5). doi: 10.1371/journal.pone.0002208.

[11] Javer A, Ripoll-Sanchez L, Brown AEX. Powerful and interpretable behavioural features
for quantitative phenotyping of *Caenorhabditis elegans*. Philos Trans R Soc Lond B Biol Sci.
2018;373(1758). doi: 10.1098/rstb.2017.0375.
[12] Partridge FA, Brown AE, Buckingham SD, Willis NJ, Wynne GM, Forman R, et al. An
automated high-throughput system for phenotypic screening of chemical libraries on *C. elegans*
and parasitic nematodes. Int J Parasitol-Drug. 2018;8(1):8-21. doi:
10.1016/j.ijpddr.2017.11.004.
[13] Puckering T, Thompson J, Sathyamurthy S, Sukumar S, Shapira T, Ebert P. Automated
Wormscan. F1000Res. 2019;6:192. doi: 10.12688/f1000research.10767.3.
[14] McDermott-Rouse A, Minga E, Barlow I, Feriani L, Harlow PH, Flemming AJ, et al.
Behavioral fingerprints predict insecticide and anthelmintic mode of action. Mol Syst Biol.
2021;17(5):e10267. doi: 10.15252/msb.202110267.
[15] Gilleard JS. *Haemonchus contortus* as a paradigm and model to study anthelmintic drug
resistance. Parasitology. 2013;140(12):1506-22. doi: 10.1017/S0031182013001145.
[16] Preston S, Jabbar A, Gasser RB. A perspective on genomic-guided anthelmintic discovery
and repurposing using *Haemonchus contortus*. Infect Genet Evol. 2016;40:368-73. doi:
10.1016/j.meegid.2015.06.029.
[17] Smout MJ, Kotze AC, McCarthy JS, Loukas A. A Novel High Throughput Assay for
Anthelmintic Drug Screening and Resistance Diagnosis by Real-Time Monitoring of Parasite
Motility. Plos Neglected Tropical Diseases. 2010;4(11). doi: ARTN e885
10.1371/journal.pntd.0000885.
[18] Preston S, Jabbar A, Nowell C, Joachim A, Ruttkowski B, Baell J, et al. Low cost whole-
organism screening of compounds for anthelmintic activity. Int J Parasitol. 2015;45(5):333-43.
doi: 10.1016/j.ijpara.2015.01.007.
[19] Jiao Y, Preston S, Koehler AV, Stroehlein AJ, Chang BCH, Simpson KJ, et al. Screening
of the 'Stasis Box' identifies two kinase inhibitors under pharmaceutical development with
activity against *Haemonchus contortus*. Parasit Vectors. 2017;10(1):323. doi: 10.1186/s13071-
017-2246-x.
[20] Preston S, Jiao Y, Jabbar A, McGee SL, Laleu B, Willis P, et al. Screening of the 'Pathogen
Box' identifies an approved pesticide with major anthelmintic activity against the barber's pole
worm. Int J Parasitol Drugs Drug Resist. 2016;6(3):329-34. doi: 10.1016/j.ijpddr.2016.07.004.
[21] Perni M, Challa PK, Kirkegaard JB, Limbocker R, Koopman M, Hardenberg MC, et al.
Massively parallel *C. elegans* tracking provides multi-dimensional fingerprints for phenotypic
discovery. J Neurosci Methods. 2018;306:57-67. doi: 10.1016/j.jneumeth.2018.02.005.
[22] Moen E, Bannon D, Kudo T, Graf W, Covert M, Van Valen D. Deep learning for cellular
image analysis. Nat Methods. 2019;16(12):1233-46. doi: 10.1038/s41592-019-0403-1.
[23] Bates K, Le K, Lu H. Deep learning for robust and flexible tracking in behavioral studies
for *C. elegans*. bioRxiv. 2021:2021.02.08.430359. doi: 10.1101/2021.02.08.430359.
[24] Fudickar S, Nustede EJ, Dreyer E, Bornhorst J. Mask R-CNN Based *C. Elegans* Detection
with a DIY Microscope. Biosensors-Basel. 2021;11(8). doi: ARTN 257
10.3390/bios11080257.
[25] Divya V Naik PS. Leaf disease detection using Mask R-CNN. International journal of
current engineering and scientific research. 2020;7(7):52-7.
[26] He K, Gkioxari G, Dollar P, Girshick R. Mask R-CNN. IEEE Trans Pattern Anal Mach
Intell. 2020;42(2):386-97. doi: 10.1109/TPAMI.2018.2844175.
[27] Roos MH, Otsen M, Hoekstra R, Veenstra JG, Lenstra JA. Genetic analysis of inbreeding
of two strains of the parasitic nematode *Haemonchus contortus*. Int J Parasitol. 2004;34(1):109-
15. doi: 10.1016/j.ijpara.2003.10.002.

[28] Preston S, Jabbar A, Nowell C, Joachim A, Ruttkowski B, Cardno T, et al. Practical and
low cost whole-organism motility assay: A step-by-step protocol. Mol Cell Probes.
2016;30(1):13-7. doi: 10.1016/j.mcp.2015.08.005.
[29] Rezatofighi H, Tsoi N, Gwak J, Sadeghian A, Reid I, Savarese S. Generalized Intersection
over Union: A Metric and A Loss for Bounding Box Regression. 2019 Ieee/Cvf Conference on
Computer Vision and Pattern Recognition (Cvpr 2019). 2019:658-66. doi:
10.1109/Cvpr.2019.00075.
[30] Lin T-Y, Maire M, Belongie S, Hays J, Perona P, Ramanan D, et al. Microsoft COCO:
Common Objects in Context. Computer Vision – ECCV 2014. 2014;8693:740-55. doi:
10.1007/978-3-319-10602-1_48.
[31] Dutta A, Zisserman A. The VIA annotation software for images, audio and video.
Proceedings of the 27th ACM international conference on multimedia. 2019:2276-9. doi:
10.1145/3343031.3350535.
[32] van der Walt S, Schönberger JL, Nunez-Iglesias J, Boulogne F, Warner JD, Yager N, et
al. scikit-image: Image processing in Python. PeerJ. 2014;2:e453. doi: 10.7717/peerj.453.
[33] Allan D, Caswell T, Keim N, van der Wel C. trackpy: Trackpy v0.3.2. 2016. doi:
10.5281/zenodo.1213240.
[34] Von Neumann J, Ulam S. Monte carlo method. National Bureau of Standards Applied
Mathematics Series. 1951;12(1951):36.
[35] Böhm A, Mayer N, Brox T. Diskmask: Focusing Object Features for Accurate Instance
Segmentation of Elongated or Overlapping Objects. 2020 IEEE 17th International Symposium
on Biomedical Imaging (ISBI). 2020:230-4. doi: 10.1109/ISBI45749.2020.9098435.
[36] Roeber F, Jex AR, Gasser RB. Advances in the diagnosis of key gastrointestinal nematode
infections of livestock, with an emphasis on small ruminants. Biotechnol Adv.
2013;31(8):1135-52. doi: 10.1016/j.biotechadv.2013.01.008.
643