1 **Image recognition based on deep learning in *Haemonchus contortus* motility**

2 **assays**

Martin Žofka, Linh Thuy Nguyen, Eva Mašátová, Petra Matoušková*

Department of Biochemical Sciences, Faculty of Pharmacy, Charles University, Heyrovského

1203, 500 05 Hradec Králové, Czech Republic.

Correspondence:

*Corresponding Author

matousp7@faf.cuni.cz

12 Abstract

Poor efficacy of some anthelmintics and rising concerns about the widespread drug resistance

have highlighted the need for new drug discovery. The parasitic nematode *Haemonchus*

contortus is an important model organism widely used for studies of drug resistance and drug

screening with the current gold standard being the motility assay. We applied a deep learning

approach Mask R-CNN for analysing motility videos and compared it to other commonly used

algorithms with different levels of complexity, namely Wiggle Index and Wide Field-of-View

Nematode Tracking Platform. Mask R-CNN consistently outperformed the other algorithms in

terms of the forecast precision across the videos containing varying rates of motile worms with

a mean absolute error of 5.6%. Using Mask R-CNN for motility assays confirmed the common

problem of algorithms that use Non-Maximum Suppression in detecting overlapping objects,

which negatively impacted the overall precision. The use of intersect over union (IoU) as a

measure of the classification of motile / non-motile instances had an overall accuracy of 89%.

In comparison to the existing methods evaluated here, Mask R-CNN performed better and we

can anticipate that this method will broaden the number of possible approaches to video analysis

of worm motility. IoU has shown promise as a good metric for evaluating motility of individual

worms.

Author summary

Deep Learning has made significant improvements over the past years reaching a level where

models are able to successfully detect heterogeneous object instances in an image. The ability

to precisely detect larvae in an image then allows the researchers to perform multiple types of

analyses that work with the viability of the parasites and are not only limited to motility assays,

but also include larval development tests and egg hatch tests which are common *in vitro* assays

for diagnosis of drug resistance. Although the other types of tests were not the main subject of

this study, they show the versatile applications that are at hand with the use of Deep Learning

models. The most important aspect of these applications is the possibility to automate these *in*

vitro assays and scale them, therefore allowing researchers to avoid tedious repetitive tasks and

focus on activities where they provide a higher added value.

Introduction

Parasitic nematode infections present a substantial problem to human and veterinary

medicine. Anthelmintics are often used as a weapon of choice to control or eliminate parasitic

worms. With increasing reports of drug resistance, there is an ongoing effort put into the

development of new anthelmintics [1, 2]. In view of this, drug screening and drug testing

methods are clearly necessary. A number of important human and animal nematodes were

already studied in large chemical library-scale screenings, although, nearly all of the studies

were aimed at well-established non-parasitic *Caenorhabditis elegans* because of the scientific

interest and medical applications as potential human disease models [3-7].

Nowadays, *in vitro* assays can measure the effects of compounds on development,

growth, behaviour, and motility. Several approaches have been pursued in attempts to deliver

robust, automated assays on nematode phenotype [8-11]. The current gold standard for

measuring drug effectiveness is *in vitro* assessment of worm motility, as measured via

microscopy. The automated phenotyping from short video recordings offers a useful assay for

high-throughput whole-organism phenotypic screening and avoids manual scoring. In assays

with parasitic worms under experimental conditions, several challenges need to be overcome

to enhance performance of motility measurements. These challenges are associated with the

different worm lengths, their tendency to clump, and also the range of diverse complex patterns

of movement different from that seen in *C. elegans*.

Focusing on veterinary importance, the parasitic nematode *Haemonchus contortus*,

known as barber's pole worm, is a model organism often used for drug screening and studying

drug resistance; due to its socio-economic importance, known genome and transcriptome [12,

13]. *H. contortus* infects predominantly small ruminants, it resides in abomasum where it feeds

on blood. It has a direct life cycle, the third-larval stage (L3s) can be stored, and researchers

commonly use L3s or *in vitro* fourth larval stage for motility studies. Many automated or semi-

automated approaches to measure motility have already been described such as via electrical

impedance [14] or infrared light beam-interference [5]. Nevertheless, for our study, we have

compared various (representative) algorithms on how to analyse the video recordings in *H.*

contortus with a different level of complexity in terms of model preparation as well as

computational cost. Therefore, we were interested to assess if the additional complexity of the

methods leads to a sufficient increase in precision, that would justify the time/performance

trade-off of these methods.

The first one mentioned is an algorithm which calculates a motility index value based

on measuring the standard deviation of the pixel's light intensity averaged for a number of

frames. This Wiggle Index (WI) assay led to many "hit compounds" in screening of a large

variety of compound libraries [15, 16]. The second approach called Wide Field-of-View

Nematode Tracking Platform (WF-NTP) is built on the use of the Gaussian adaptive threshold

of the image frames to identify worms and was applied in model free-living nematode *C.*

elegans. It was initially developed for drug screening purposes, nevertheless, it has also found

applications in other studies such as detailed genetic or behavioural studies [17]. In our study,

we customized the associated WF-NTP software written in Python and compared it to the last

approach which involves the application of deep learning.

The advent of machine learning has revolutionized the surveying and classifying of

biological data including image recognition. Large-scale computerization reduced the time that

needs to be spent in the laboratory, however the rate-limiting step is the analysis and

interpretation of data. In the field of computer vision, deep learning algorithms with a

convolutional neural network (CNN) architecture have made rapid advancements on a variety

of image classification tasks in cell cultures [18], yet none of them have been applied to parasitic

nematodes. Object-detection-based deep learning methods have also been adapted for instance

segmentation. They work on the principle of predicting bounding boxes for all objects in an

image and use non-maximum suppression to remove redundant bounding-box predictions. Here

we used a state-of-the-art region-based CNN called Mask R-CNN [19] which predicts an object

mask for each bounding box, and compared it to the WI and WF-NTP methods. The aim was

to evaluate the application of the Mask R-CNN method to the field of drug discovery of

anthelmintics via measuring motility of *H. contortus*, using a metric called intersection over

union (IoU).

Results

Motility forecast errors

In the manual video processing, we observed the motility and the live rate. To measure the performance of the algorithms, we used the motility rate rather than the live rate even though the stillness does not implicate a dead worm. The reason for this choice was that the algorithms detect movement and are not able to differentiate if the movement was caused by the object itself or by an interaction with another object. The general manual motility rate was always higher than the manual live rate (Supplementary Table S1). We measured the performance of the individual algorithms by evaluating the errors, defined as the differences between the manual processing of the file and the respective algorithm. Out of the three methods, Mask R-CNN had the lowest mean absolute error (MAE) with a value of 5.6%, followed by WF-NTP (8.76%) and the WI (14.2%). For WI we can apply a correction to limit the maximum motility value to 1, then the capped MAE is reduced to 11.8%. In terms of the bias of the predictions, the WI had the lowest mean error (ME) of -0.71% therefore slightly overestimating the motility rate on average, while both WF-NTP and Mask R-CNN underestimated the motility rate with values of 6.52% and 1.95%, respectively (Table 1).

WI had an increasing volatility with an increase in the ratio of alive worms in the group. WF-NTP and Mask R-CNN had a tendency to systematically underestimate for the motility group 100 as can be seen in Fig 1. The mean and standard deviation of motility rates per motility group and algorithm is shown also in Supplementary Table S2.

Table 1: Motility error metrics for algorithms

Algorithm	MAE, %	ME, %
Wiggle Index	14.2	-0.71
WF-NTP	8.76	6.52

Mask R-CNN	5.61	1.95

The differences between the manual processing and the respective algorithm were expressed as the mean absolute error (MAE) and mean error (ME).

Fig 1: Motility rates comparisons for individual motility groups per algorithm. The number of the motility group indicates the percentage of live larvae. The motility rate is visualized as box plots.

Instance detection performance

We only compared Mask R-CNN and WF-NTP for the total worm count as the WI does not detect individual instances of worms and therefore this information was not available. Similarly as for the motility rate, the precision was measured based on the error term between the manual counts and the respective algorithm. In terms of the size of the error, the ean absolute percentage error (MAPE) was used and we observed that the error was significantly lower ($P <$ 0.0001, Wilcoxon signed-rank test) for Mask R-CNN (7.6%) than for WF-NTP (40.23%). Both algorithms had a tendency to underestimate the number of worms on average, therefore not detecting all the worm instances in a video. This was confirmed by the mean percentage error (MPE) with values of 5.61% for Mask R-CNN and 40.23% for WF-NTP (Table 2). The reported MAPE and MPE for WF-NTP had the same value because all the detections underestimated the total number of worms. WF-NTP underestimated the worm count similarly for all motility groups, while Mask R-CNN had similar box plots in comparison to the manual counts apart from underestimating the motility group 0 as shown in Fig 2. The mean error and standard deviation of the differences between the manual count and the respective motility algorithm is shown in Supplementary Table S3.

Table 2: Worm count error metrics for algorithms

Algorithm / Metric	MAPE, %	MPE, %
WF-NTP	40.23	40.23
Mask R-CNN	7.6	5.61

The differences between the manual counts and the respective algorithm were expressed as the mean absolute percentage error (MAPE) and mean percentage error (MPE).

Fig 2: Worm count comparisons for individual motility groups per algorithm. The number of the motility group indicates the percentage of live larvae. The number of worms is visualized in box plots.

Mask R-CNN detection and classification precision

The precision of detecting worm instances in a video was on average 98.6%. The majority of cases where the detection was incorrect was due to debris that had a similar size and shape as a worm. For the detected worms there was an overall accuracy of 89% to correctly label them as either motile or non-motile based on their mean IoU values. In Fig 3, the distribution for different IoU values clearly showed that most of misclassified cases were located around the threshold 0.8, while the classification accuracy increased for values further away from the threshold. The precision and recall values (Table 3) confirm that the prevalent case was the misclassification of motile worms as non-motile.

The errors defined as differences between Mask R-CNN and the manually evaluated motility rates were tested for normality using the D'Agostino-Pearson test as well as Shapiro-

Wilk test. In both cases we could not reject the null hypothesis that the data were from a normal

distribution with p-values 0.85 and 0.35, respectively (Table 4).

Fig 3: The distribution for different Intersection over Union (IoU). Both **(a)** histogram of IoU

and the associated number of worms per bin classified into motile / non-motile and **(b)**

histogram of IoU and the associated probability per class (motile / non-motile) show that the

majority of misclassified cases were located around the 0.8 IoU.

Table 3: Classification performance metrics

Classification	Precision, %	Recall, %
Motile	94	88
Non-Motile	84	92

176 Precision and recall metrics for classification performance

178 **Table 4:** Normality tests for the motility rate error terms between Mask R-CNN and the manual

179 processing

Normality test	Statistic	p-value
D'Agostino-Pearson	0.32	0.85
Shapiro-Wilk	0.97	0.35

181 For both normality tests we cannot reject the null hypothesis that the data were from a normal

182 distribution.

Discussion

The motility phenotype is often used for drug screening and drug testing of anthelmintics. In our study, we deployed and compared three algorithms to measure motility in the parasitic nematode *H. contortus*. The results showed that the performance increased with an increasing complexity of the algorithm. WI apart from requiring very little configuration and being the fastest algorithm had some limitations in comparison to the other methods; it was by design not able to detect individual instances of worms and a control group with only motile worms needed to be created in order to be able to translate the WI into a percentage for the motility rate for each video, else it was hard to interpret the numerical value of the WI. Because of these limitations and the worse precision, it would be advisable to use WI only for initial studies to quickly test experimental results as the method is the easiest to set up.

For WF-NTP, overlapping parasites had the highest impact on the decline in the detection ability, as the algorithm filters out objects that are above or below a certain pixel size. This resulted in detecting overlapping worms as a single object, where these objects were then excluded based on the size threshold. In consequence, the total number of detected instances in an image was significantly lower than the actual number obtained by manually processing the videos. Although WF-NTP struggled with detecting all instances, the classification motile / non-motile worked well, once an instance was detected.

Mask R-CNN had the lowest MAE, nonetheless similarly to WF-NTP it struggled with overlapping objects, primarily due to the fact that the algorithm uses Non-Maximum Suppression. As a result, some instances were not detected, although in contrast to WF-NTP the worm clews were not discarded as a whole. The problem of detecting overlapping objects is currently an area of active research in object detection, particularly for biological data. For instance, Böhm et al. [20] are working on algorithms that can handle overlapping objects and have had success with applying their algorithm to *C. elegans*. Motile worms had a higher chance

of being detected, because they had higher variability across the frames due to their movement.

In terms of precision, the algorithm was slightly biased towards underestimating the motility

rate, however the errors were normally distributed enabling us to increase the number of

acquired videos in order to get to a desired level of confidence and therefore successfully

replace manual processing.

The results confirmed that IoU can be used as a metric for determining the motility of

larvae, although some modifications might be required in order to deal with slow-moving

worms, such as including the total distance travelled during the video to complement the frame-

by-frame information. Furthermore, the IoU threshold should be considered a parameter to be

optimized as it can vary based on the age of the larvae, the temperature of the environment and

other factors. The downside of using Mask R-CNN is that it requires a quite big amount of

annotated data (provided in the Data availability), which is time demanding to create and that

the training runs 1-2 days on a GPU instance.

In general, for quick low-precision experiments, the best option is to use the WI as it

can be used out-of-the-box with slight modifications to a few parameters. If we need to detect

individual instances, Mask R-CNN is the best choice, although it is more time-demanding to

prepare the data and train the model.

Materials and methods

Procurement of L3s of *H. contortus*

Six-month-old sheep was orally infected with 8,000 infective L3s of MHco3 strain

(Inbred Susceptible Edinburgh, ISE) [21]. Four weeks post-infection, faecal samples were

collected. L3s were produced from eggs by incubating faeces in a plastic box covered with foil

at 27 °C for 7 days. Then, faeces were rinsed twice in tap water which was poured into 1,000mL

conical measuring cups in which the larvae sank to bottom. To remove dirt or dead individuals,

the pellet of larvae was pipetted out and filtered through a 20µm sieve submerged in water (27

°C). Clean L3s at a concentration of approximately 4,000 L3 per mL were stored in culture

flasks in water at 10 °C for several months.

During the storage, some larvae naturally die. Prior to the experiment, we filtered larvae

through a 20µm sieve (for 12 h), this time collecting both live larvae from the bottom of the

sieve and dead individuals remaining on the sieve.

All experimental procedures were examined and approved by the Ethics Committee of

the Ministry of Education, Youth and Sports (Protocol MSMT-25908/2014-9).

Video processing and manual counting

Videos from a microscope camera were recorded for a duration of 10 seconds with a

framerate of 20 frames per second (fps) in a quality of 2560 by 2160 pixels. The videos (.avi

with MJPEG compression, ~ 0.45GB) were acquired using NIS-Elements Imaging Software

(version 4.20) using a Nikon Eclipse Ti microscope (4x magnification) with camera (Andor

Zyla 5.5 sCMOS, 12bit: 2560 x 2160 pixels). The videos were recorded for different ratios of

live and dead larvae (100:0, 75:25, 50:50, 25:75, 0:100; denoted as motility group 100, 75, 50,

25 and 0, respectively) in 9 replicates and they contained on average 60 worms. The live larvae

were kept in 37 °C to maintain their motility.

The video recordings were processed so that each worm was labelled with a numeric

identifier. A trained human accessed each worm from each video and information about motility

and dead phenotype was recorded. The discrepancies were dealt with as following (i) a dead

worm which was moving due to interference of a motile worm was counted as motile; (ii) a live

coil non-motile worm was counted as non-motile; (iii) the worms which were not labelled by

an algorithm were given an identifier manually; (iv) worms were excluded from the counts

when they appeared in less than 20 frames; (v) dead phenotype was evaluated when overt

microscopical pathologies, such as the presence of large vacuoles in the tegument and straight

phenotype, were observed.

This manual counting was taken as a referential method for comparison of accuracy

among the three models.

Wiggle Index analysis

We used an ImageJ implementation of the WI for the analysis [22]. We did not apply

any scaling of the videos, the Gaussian image blur was set to 2, the number of frames to be

averaged was set to 50 and the motility output was based on the standard deviation. Since the

values from the WI are relative, we used motility group 100 as a control group to be able to

convert these relative values to percentages. We calculated the mean value for the group 100

and divided all values by the obtained mean. However, after the conversion some values may

be higher than 1, which in the context of motility does not make sense, therefore those values

can be capped to 1 for the calculation of the error metrics.

WF-NTP analysis

WF-NTP presents an automated evaluation of motility in *C. elegans*. For its successful

implementation, we adopted the method for the *H. contortus* motility assay. However, as these

parasites differ in their sizes as well as their movement, some modifications to the original

configuration and algorithm were necessary. These modifications belonged generally to two

categories, where the first was a recalibration of the available input parameters of the WF-NTP

method. This included aspects such as the image size, the size of the worms that we want to

detect and other configurations (see Data availability). The other modifications were changes

in the evaluation of the conditions for worm motility and a limit on the minimum number of

frames that a worm needs to be detected in for it to be included in the evaluation.

The original method relied on bends per second of individual worms to classify them as

either motile or non-motile, however as the movement of *H. contortus* is more complex, we

replaced the bends per second with an evaluation metric that is regularly used in image

recognition, namely Intersection over union (IoU) which quantifies the percent overlap between

the target mask and our prediction output [23]. The use of IoU allowed us to use the capabilities

of WF-NTP in terms of the detection of individual worm instances in an image and then apply

the IoU metric to determine the worm motility. The calculation of IoU was done for the same

worm instance in neighbouring frames to detect the change on a frame-by-frame basis and then

the mean was taken and worms above a certain IoU threshold were labelled as non-motile. To

ensure that the comparison of WF-NTP and Mask R-CNN was aimed at the quality of the

detection of individual worm instances rather than the approach to determine the motility, we

used the same IoU conditions for both methods.

Another modification of the original method was the introduction of a threshold for the

minimum number of frames, that a worm needed to be detected in order to be included in the

analysis. The reason for this restriction was that as the individual worms move, they have a

tendency to either change the shape or overlap with other worms, therefore in terms of the

detection the individual worm can be lost for a certain number of frames and reappear later,

however this can lead to counting some instances multiple times. To limit these potential

duplicities (Supplementary Fig S1) as much as possible and not bias the total number of worms

detected in the video, we selected the highest number of worms detected in a given video across

all frames and added a buffer of 10% in order to account for situations where some worms were

not detected, for example due to their overlaps. This maximum number then served as a limit

for selecting the number of worms to be evaluated as motile or non-motile. The instances were

selected based on the number of frames in a decreasing order, therefore ensuring that we had

data from enough frames to classify the worm correctly. Again, this restriction was applied to

detections from Mask R-CNN as well to enable a comparison under the same conditions.

Mask R-CNN analysis

For   Mask   R-CNN   we   used   the   Matterport   implementation

(https://github.com/matterport/Mask_RCNN) of the algorithm including a trained model using

a backbone architecture of Resnet101. The pre-trained model weights were obtained from

training the model on the MS COCO dataset [24]. We selected a pre-trained model because in

general deep neural networks require a lot of annotated data for training and using an existing

model and re-training the top layer of the network allows us to have a smaller dataset for the

training of the model, while maintaining a good performance. We manually annotated a total

of 95 images and divided them into training (69) and validation (26) using the VGG (Visual

Geometry Group) Image Annotator [25], these images contained three categories of objects

eggs, second-stage larvae (L2) and third-stage larvae (L3), to fully use the capabilities of Mask

R-CNN which is able to do instance segmentation as well as classification of those instances.

The total number of annotated objects was around 10,400 divided into training (6,800 objects)

and validation (3,600 objects). Then we ran the training on a GPU instance until the loss

function showed no signs of improvement. To avoid overfitting, we used data augmentation

and applied one or multiple of the following augmentations: flipping, contrast normalization,

additive gaussian noise, multiplying the pixels by a number within a given interval to make the

whole image lighter or darker. The final model had a mAP IoU of 64.1% on the validation set.

The trained model was then used on the individual frames of the video to detect the

instances of parasites and we calculated the centroid of the detected object in terms of the

position on the x and y axis using the scikit-image regionProps functionality [26]. These

detected instances were then linked together using the trackpy library [27], which detected the

trajectories of objects based on certain conditions, in our case we restricted the distance that the

centroid could move a maximum of 100 pixels in each dimension. As a result of detecting the

trajectories, we were now able to identify the same parasite instance across the frames.

Subsequently, we calculated the IoU and the threshold for the number of parasites to select for

the evaluation as described in the WF-NTP section. We observed a decreasing MPE with an

increase in the length of the video and therefore we decided on using the full length of the

videos for the motility assay (Supplementary Fig S2).

Error metrics

For the evaluation of the performance of the individual algorithms we used standard

metrics for the quantification of the forecast precision. These metrics work with the error term,

which is defined as the difference between the actual and the forecasted value, where in our

case the actual value was obtained by manually processing the videos. Two aspects were

measured, the first one was the bias of the forecasts to determine if the algorithm is

systematically over or under-estimating, for that purpose we used ME or MPE, which both work

with the mean of the error term. The second aspect was the size of the error term irrespective if

the error term was positive or negative to compare the overall performance of the algorithms,

for that purpose we used MAE or MAPE, which both work with the mean absolute value of the

error term.

Additional files

Supplementary Table S1: Manual counts of motility and live rates per motility group

Supplementary Table S2: Motility rates per motility group and algorithm

Supplementary Table S3: Mean error between the manual count and the respective motility

algorithm

Supplementary Table S4: Mean of differences in worm counts between the manual processing

and the respective algorithm

Supplementary Fig S1: Average number of worms in the videos per motility group for WF-

NTP and Mask R-CNN.

Supplementary Fig S2: The impact of the length of the video in frames on the percentage error

of the detected worms for Mask R-CNN.

Abbreviations

365 CNN: convolutional neural network; fps: frames per second; GPU: graphics processing unit;

366 IoU: intersection over union; MAE: mean absolute error; MAPE: mean absolute percentage

367 error; Mask R-CNN: region based convolutional neural network; ME: mean error; MPE: mean

368 percentage error; WF-NTP: Wide Field-of-View Nematode Tracking Platform; WI: Wiggle

369 Index

Competing Interests

372 The authors declare that they have no competing interests.

Funding

375 This work was supported by Charles University UNCE/18/SCI/012 and SVV 260 550 as well

376 as by the project EFSA-CDN [CZ.02.1.01/0.0/0.0/16_019/0000841], co-funded by ERDF.

Authors' contributions

M.Ž. and L.T.N. designed this study and wrote the manuscript. M.Ž. implemented the deep

learning model, wrote the code, ran the statistical analyses and generated the figures. L.T.N.,

E.M. acquired and annotated the images used to train the deep learning model. M.Ž. and L.T.N.

conducted the experiments and analyzed the results. P.M. managed the project. All authors

contributed to literature review and critical revision of the manuscript.

Acknowledgements

We thank MVDr. Tomáš Kaňka for assisting with the sheep infection. The authors thank prof.

Lenka Skálová for critical discussion and comments.

References

1.   Jiao Y, Preston S, Hofmann A, Taki A, Baell J, Chang BCH, et al. A perspective on the
discovery of selected compounds with anthelmintic activity against the barber's pole worm-
Where to from here? Adv Parasitol. 2020;108:1-45. Epub 2020/04/16. doi:
10.1016/bs.apar.2019.12.003. PubMed PMID: 32291083.
2.   Geary TG, Sakanari JA, Caffrey CR. Anthelmintic drug discovery: into the future. J
Parasitol. 2015;101(2):125-33. Epub 2015/01/15. doi: 10.1645/14-703.1. PubMed PMID:
25584662.
3.   Buckingham SD, Partridge FA, Sattelle DB. Automated, high-throughput, motility
analysis in Caenorhabditis elegans and parasitic nematodes: Applications in the search for new
anthelmintics. Int J Parasitol Drugs Drug Resist. 2014;4(3):226-32. Epub 2014/12/18. doi:
10.1016/j.ijpddr.2014.10.004. PubMed PMID: 25516833; PubMed Central PMCID:
PMCPMC4266775.
4.   Burns AR, Luciani GM, Musso G, Bagg R, Yeo M, Zhang Y, et al. Caenorhabditis
elegans is a useful model for anthelmintic discovery. Nat Commun. 2015;6. doi: ARTN 7485
10.1038/ncomms8485. PubMed PMID: WOS:000357178100005.
5.   Taki AC, Byrne JJ, Wang T, Sleebs BE, Nguyen N, Hall RS, et al. High-Throughput
Phenotypic Assay to Screen for Anthelmintic Activity on Haemonchus contortus.
Pharmaceuticals (Basel). 2021;14(7). Epub 2021/07/03. doi: 10.3390/ph14070616. PubMed
PMID: 34206910; PubMed Central PMCID: PMCPMC8308562.
6.   Elfawal MA, Savinov SN, Aroian RV. Drug Screening for Discovery of Broad-
spectrum Agents for Soil-transmitted Nematodes. Sci Rep-Uk. 2019;9. doi: ARTN 12347
10.1038/s41598-019-48720-1. PubMed PMID: WOS:000482564800003.
7.   Pulak R. Techniques for analysis, sorting, and dispensing of C. elegans on the COPAS
flow-sorting system. Methods Mol Biol. 2006;351:275-86. Epub 2006/09/22. doi: 10.1385/1-
59745-151-7:275. PubMed PMID: 16988441.
8.   Ramot D, Johnson BE, Berry TL, Jr., Carnell L, Goodman MB. The Parallel Worm
Tracker: a platform for measuring average speed and drug-induced paralysis in nematodes.
PLoS One. 2008;3(5):e2208. Epub 2008/05/22. doi: 10.1371/journal.pone.0002208. PubMed
PMID: 18493300; PubMed Central PMCID: PMCPMC2373883.
9.   Javer A, Ripoll-Sanchez L, Brown AEX. Powerful and interpretable behavioural
features for quantitative phenotyping of Caenorhabditis elegans. Philos Trans R Soc Lond B
Biol Sci. 2018;373(1758). Epub 2018/09/12. doi: 10.1098/rstb.2017.0375. PubMed PMID:
30201839; PubMed Central PMCID: PMCPMC6158219.

10. Partridge FA, Brown AE, Buckingham SD, Willis NJ, Wynne GM, Forman R, et al. An
automated high-throughput system for phenotypic screening of chemical libraries on C. elegans
and parasitic nematodes. Int J Parasitol Drugs Drug Resist. 2018;8(1):8-21. Epub 2017/12/11.
doi: 10.1016/j.ijpddr.2017.11.004. PubMed PMID: 29223747; PubMed Central PMCID:
PMCPMC5734697.
11. Puckering T, Thompson J, Sathyamurthy S, Sukumar S, Shapira T, Ebert P. Automated
Wormscan. F1000Res. 2019;6:192. doi: 10.12688/f1000research.10767.3. PubMed PMID:
28413617; PubMed Central PMCID: PMCPMC5365223.
12. Gilleard JS. Haemonchus contortus as a paradigm and model to study anthelmintic drug
resistance.   Parasitology.   2013;140(12):1506-22.   Epub   2013/09/04.   doi:
10.1017/S0031182013001145. PubMed PMID: 23998513.
13. Preston S, Jabbar A, Gasser RB. A perspective on genomic-guided anthelmintic
discovery and repurposing using Haemonchus contortus. Infect Genet Evol. 2016;40:368-73.
Epub 2015/07/07. doi: 10.1016/j.meegid.2015.06.029. PubMed PMID: 26144657.
14. Smout MJ, Kotze AC, McCarthy JS, Loukas A. A novel high throughput assay for
anthelmintic drug screening and resistance diagnosis by real-time monitoring of parasite
motility.   PLoS   Negl   Trop   Dis.   2010;4(11):e885.   Epub   2010/11/26.   doi:
10.1371/journal.pntd.0000885. PubMed PMID: 21103363; PubMed Central PMCID:
PMCPMC2982823.
15. Jiao Y, Preston S, Koehler AV, Stroehlein AJ, Chang BCH, Simpson KJ, et al.
Screening of the 'Stasis Box' identifies two kinase inhibitors under pharmaceutical development
with activity against Haemonchus contortus. Parasit Vectors. 2017;10(1):323. Epub
2017/07/07. doi: 10.1186/s13071-017-2246-x. PubMed PMID: 28679424; PubMed Central
PMCID: PMCPMC5499055.
16. Preston S, Jiao Y, Jabbar A, McGee SL, Laleu B, Willis P, et al. Screening of the
'Pathogen Box' identifies an approved pesticide with major anthelmintic activity against the
barber's pole worm. Int J Parasitol Drugs Drug Resist. 2016;6(3):329-34. Epub 2016/08/16. doi:
10.1016/j.ijpddr.2016.07.004. PubMed PMID: 27524698; PubMed Central PMCID:
PMCPMC5196485.
17. Perni M, Challa PK, Kirkegaard JB, Limbocker R, Koopman M, Hardenberg MC, et al.
Massively parallel C. elegans tracking provides multi-dimensional fingerprints for phenotypic
discovery.   J   Neurosci   Methods.   2018;306:57-67.   Epub   2018/02/17.   doi:
10.1016/j.jneumeth.2018.02.005. PubMed PMID: 29452179.
18. Moen E, Bannon D, Kudo T, Graf W, Covert M, Van Valen D. Deep learning for
cellular image analysis. Nat Methods. 2019;16(12):1233-46. doi: 10.1038/s41592-019-0403-1.
PubMed PMID: WOS:000499653100023.
19. He K, Gkioxari G, Dollar P, Girshick R. Mask R-CNN. IEEE Trans Pattern Anal Mach
Intell. 2020;42(2):386-97. Epub 2018/07/12. doi: 10.1109/TPAMI.2018.2844175. PubMed
PMID: 29994331.
20. Böhm A, Mayer N, Brox T. Diskmask: Focusing Object Features for Accurate Instance
Segmentation of Elongated or Overlapping Objects. I S Biomed Imaging. 2020:230-4. PubMed
PMID: WOS:000578080300033.
21. Roos MH, Otsen M, Hoekstra R, Veenstra JG, Lenstra JA. Genetic analysis of
inbreeding of two strains of the parasitic nematode Haemonchus contortus. International
Journal for Parasitology. 2004;34(1):109-15. doi: 10.1016/j.ijpara.2003.10.002. PubMed
PMID: WOS:000188377300013.
22. Preston S, Jabbar A, Nowell C, Joachim A, Ruttkowski B, Cardno T, et al. Practical and
low cost whole-organism motility assay: A step-by-step protocol. Mol Cell Probes.
2016;30(1):13-7. Epub 2015/09/15. doi: 10.1016/j.mcp.2015.08.005. PubMed PMID:
26365227.

23.   Rezatofighi H, Tsoi N, Gwak J, Sadeghian A, Reid I, Savarese S. Generalized
Intersection over Union: A Metric and A Loss for Bounding Box Regression. Proc Cvpr Ieee.
2019:658-66. doi: 10.1109/Cvpr.2019.00075. PubMed PMID: WOS:000529484000067.
24.   Lin T-Y, Maire M, Belongie S, Hays J, Perona P, Ramanan D, et al., editors. Microsoft
coco: Common objects in context. European conference on computer vision; 2014: Springer.
25.   Dutta A, Zisserman A, editors. The VIA annotation software for images, audio and
video. Proceedings of the 27th ACM international conference on multimedia; 2019.
26.   van der Walt S, Schonberger JL, Nunez-Iglesias J, Boulogne F, Warner JD, Yager N, et
al. scikit-image: image processing in Python. PeerJ. 2014;2:e453. Epub 2014/07/16. doi:
10.7717/peerj.453. PubMed PMID: 25024921; PubMed Central PMCID: PMCPMC4081273.
27.   Allan D, Caswell T, Keim N, van der Wel C. TrackPy. DOI: https://doi
org/105281/zenodo. 2014;9971.
485

Figure 1

Figure 2

Figure 3