Experimental sound exposure modifies swimming activities and increases food handling error in adult zebrafish (*Danio rerio*)

Reza Mohsenpour[a], Saeed Shafiei Sabet[a,*]

[a] Fisheries Department, Faculty of Natural Resources, University of Guilan, Sowmeh Sara, Iran.

Abstract:

Anthropogenic sound is globally increasing and is recognized as a source of environmental pollution in terrestrial and aquatic habitats. Elevated sound levels may cause a broad range of impacts on aquatic organisms among taxa. Sound is an important sensory stimulus for aquatic organisms and it may cause fluctuations in stress-related physiological indices and in a broader extent induce behavioural effects such as driving as a distracting stimulus, masking important relevant acoustic signals and cues in a range of marine and freshwater species. However sound exposure may also induce changes in swimming activities, feed efficiency and productivity of available food sources in fish. Here, we experimentally tested sound effects on swimming activities and foraging performance in thirty adult Zebrafish (*Danio rerio*) individually in captivity. We used adult zebrafish and water flea (*Daphnia magna*) as model predator and prey, respectively. In terms of acoustic stimuli, we used four sound treatments with different temporal patterns (all in the same frequency range and moderate exposure level). As in a previous study, our results constitute strong evidence for sound-related effects on zebrafish behaviour. All sound treatments led to a significant increase in the number of startle response, brief and prolonged swimming speed for zebrafish ($P<0.05$). Zebrafish reached to the baseline swimming speed after 60 seconds in all treatments. We found partially brief and prolonged sound effects on the spatial distribution of zebrafish; Although there were not any significant sound-related behavioural changes for horizontal spatial displacement in all treatments ($P>0.05$), zebrafish swam significantly more in the lower layer of the tank except for irregular intermittent 1:1-7 in brief sound exposure ($P<0.05$). The results of foraging performance showed that food discrimination error unaffected by sound treatments and was low for the zebrafish ($P>0.05$). However, food handling error was affected by sound treatments; all treatments induced a significant rise in handling error ($P<0.001$). This study highlights the impact of sound on zebrafish swimming activities, and that more attacks are needed to consume the same number of food items under noisy conditions.

Keywords: Anthropogenic sound, Behaviour, Foraging performance, Sound impact, Zebrafish

1. Introduction

Nowadays, due to the increase in human activities and the advancement of technology globally since the Industrial Revolution, the living environment has undergone extensive changes (Normandeau Associates, 2012). These environmental changes can affect the planet and living organisms, and that can be a major threat to the biodiversity inhabit Earth (Kunc et al., 2016). The rapid growth of these changes poses many environmental challenges (Tuomainen and Candolin, 2011) in both terrestrial and aquatic habitats. Environmental pollutions (including chemical, light, and sound) are introduced by human activities in different time, scale and space ranges and have elevated underwater ambient noise levels with alternating intensities which may affect aquatic organisms in their habitats (Popper and Hastings, 2009; Slabbekoorn et al., 2010). Among these, one of the main sources of environmental pollution which may also can be recognized as an environmental stress stimulus is anthropogenic sound that in addition to affecting terrestrial animals, also have many consequences on aquatic organisms (Popper et al., 2020; Slabbekoorn et al., 2010; Slabbekoorn and Ripmeester, 2008).

Sound sources in aquatic habitats, such as merchant shipping, recreational boating, wind turbines, pile-driving, underwater mining explorations, and explosions related to geological and research experiments, are frequent in the number of events and widespread geographically. (McDonald et al., 2006; Normandeau Associates, 2012). Consequently, anthropogenic sound has changed underwater soundscapes worldwide and represents a very subtle driver of environmental change and novel challenge to aquatic organisms. Moreover, the high speed of sound underwater, which is about 5 times faster than the speed of sound in air, shows the importance and priority of examining the role and applications of acoustic stimuli and their effects on aquatic organisms.

Recent studies have investigated impacts of anthropogenic sound on a wide variety of taxa and across a range of scales (Barber et al., 2009; Morley et al., 2014; Normandeau Associates, 2012; Slabbekoorn et al., 2010; Thomsen et al., 2021; Tyack, 2008). Anthropogenic sound can cause physical, physiological, and behavioural disorders in aquatic organisms, including marine mammals (Erbe et al., 2018; Moore et al., 2012; Southall et al., 2008), seabirds (Bermúdez-Cuamatzin et al., 2018; Green et al., 2016; Hansen et al., 2020),

reptiles (Injaian et al., 2020; Simmons and Narins, 2018), fish (Hastings and Popper, 2005;

Hawkins, 1986; Mills et al., 2020; Popper et al., 2003), and invertebrates (Carroll et al., 2017;

Coquereau et al., 2016; Murchy et al., 2019).

Because of the high propagation rate and low attenuation over large distances in underwater

environments acoustic stimuli play an important role in aquatic environments (Tyack, 1998;

Popper and Hastings, 2009; Slabbekoorn et al., 2010). Depending distance from the sound

source, recent studies have shown dramatic effects of sound such as physical damages, sever

injury or even death (Budelmann, 2011; Halvorsen et al., 2012; Keevin and Hempen, 1997).

Further distance from the sound source, there may be physiological responses such as

permanent and temporary hearing threshold shifts, fluctuations in physiological indices

(André et al., 2011; Casper et al., 2013; McCauley et al., 2003; Popper et al., 2007; Popper et

al., 2005; Scholik and Yan, 2002; Smith et al., 2004; Wysocki et al., 2007; Wysocki et al.,

2006) such as elevated cortisol levels, the classical stress-related hormone (Johansson et al.,

2016; Nichols et al., 2015; Santulli et al., 1999; Smith et al., 2004; Wysocki et al., 2006) and

increased hear rates (Graham and Cooke, 2008; Simpson et al., 2015). Furthest distance from

sound source, in a broader extent, behavioural effects are the most likely to occur and thus

play as a stress driver (Smith et al., 2004; Popper and Hawkins, 2019) as a distracting

stimulus (Popper and Carlson, 1998; Chan et al., 2010), interfere with detecting prey and

antipredator behaviour (Hawkins and Myrberg, 1983), compromise foraging performance

(Neo et al., 2015; Purser and Radford, 2011; Shafiei Sabet et al., 2015; Voellmy et al., 2016),

disrupt reproductive behaviour (McCloskey et al., 2020) or mask important acoustic signals

and cues for conspecific recognition and communication purposes (Codarin et al., 2009;

Amorim et al., 2015; De Jong et al., 2018b; Hawkins and Picciulin, 2019) in a range of

marine and freshwater species.

Many marine and freshwater fishes have well-developed hearing abilities that provide them a

key biological privilege to detect sound and perceive a broad range of frequencies (Hawkins,

1986; Heath et al., 2021; Popper et al., 2019; Wahlberg and Westerberg, 2005; Wysocki et

al., 2006). While there are well-documented studies regarding the effects of sound on the

behaviour of marine fishes (de Jong et al., 2018a; Mortensen et al., 2021; Peng et al., 2015),

and its adverse repercussions in many marine fish species at the individual and community

level, yet much less is known across the current literature about these effects on the behaviour

of freshwater fish and has not received sufficient interest so far (Fedoroff, 2021; Mickle and

Higgs, 2018; Pieniazek et al., 2020). Moreover, sound exposure can also change spatial

distribution and swimming behaviour of fish which may consequently affect ecologically on

their avoiding to forage in noisy food areas and their navigations but also change biologically

their swimming activities and foraging performance (de Vincenzi et al., 2021; Hanache et al.,

2020; Hubert et al., 2021; Shafiei Sabet et al., 2016a). There is growing evidence that

foraging performance is likely to be modulated by anthropogenic sound that can induce shift

from successful feeding to typical behavioural stress responses in fish (Purser and Radford,

2011; Shafiei Sabet et al., 2015; Voellmy et al., 2016. Currently, little is known about the

effects of sound exposure on swimming activity and foraging performance of fish, although

there are some well-documented studies. It has been shown that increased boating activity

was associated with a reduction in activity rates, changed vertical distribution and

compromised foraging success of free-ranging mulloway (*Argyrosomus japonicus*) (Payne et

al., 2015) and Mediterranean Damselfish (*Chromis chromis*) (Bracciali et al., 2012).

Other studies have shown that experimental sound exposure increase performance errors and

therefore displayed a negative impacts on foraging efficiency in both the three-spined stickle

backs (*Gasterosteus aculeatus*) (Purser and Radford, 2011) and the European minnow,

(*Phoxinus phoxinus*) (Voellmy et al., 2014a). More recently our previous study also have

shown a clear sound impact on zebrafish foraging performance; more food handling errors

under noisy conditions (Shafiei Sabet et al., 2015). A primary consequence of sound exposure

would appear to be shifts in the spatial displacement. The resulting disturbance might induce

modifications in allocated foraging time budget, foraging patterns and the relative abundance

of prey items and predatory species. Such changes in turn may increase foraging energy

demand and the amount of time allocated by fish to foraging which, subsequently induce a

number of major changes such as affect food searching, discriminating and handling.

In general, *Danio rerio* is known as a model fish species in behavioural studies and

responding to environmental conditions (Cachat et al., 2010; Egan et al., 2009; Whitfield,

2002). Zebrafish is a member of the Cypriniformes order and the Cyprinidae family and

acclimates well in captivity (Detrich et al., 2011). This fish naturally lives in the tropical

freshwater (Spence et al., 2008). Morphologically, the zebrafish's body is narrow and

elongated, with golden and blue stripes that stretch along the body and tail (Detrich et al.,

2004). Males and females are easily separable, so that females having a more prominent

abdomen and body than males, and males have a spindle-shaped body (Spence et al., 2008).

Zebrafish live in the temperature range of 24±2 ℃ and in the pH range of 6.8 to 7.5

(Cortemeglia and Beitinger, 2005). The adult size of the zebra fish is approximately 4.5 to 6.5

cm (Gerhard et al., 2002; Spence et al., 2008). Moreover, zebrafish have specialized hearing
structures, the Weberian ossicles, with an auditory system that has homologies to the
mammalian auditory system (Weber, 1820; Alexander, 1964; Whitfield, 2002). In recent
years, zebrafish have been used as biological models in genetic, physiological, toxicological,
behavioural, ecological, and other studies (Detrich et al., 2004; Gerlai, 2019; Gerlai et al.,
2000; Kalueff and Cachat, 2011; Lieschke et al., 2009). The high genetic, physiological, and
pharmacological similarities of this species with humans could be a reason for the use of this
species as a biological model in research (Crawford et al., 2008).

Progress in the field of behavioural biology over conservation ecology and findings about the
potential impacts pollutants on organisms is also to a large extent linked with the study of
invertebrates. Daphnia is a small crustacean and inhabits in open and light waters, also they
are an important part of the food web in freshwater habitats and inhabits many types of
shallow water bodies (Ebert, 2005; Parejko and Dodson, 1991; Reynolds, 2011). This
invertebrate is the first crustacean to have its genome sequenced (Stollewerk, 2010) and
because of features such as easy cultivation, small size and short generation time it is a
popular model organism in various biological disciplines from aquatic ecology to biomedical
sciences (Seda and Petrusek, 2011). Daphnia may also be a useful species to study
behavioural studies such as sound impacts on invertebrates.

In addition, it has been reported that in the larval stage, marine crustaceans respond to reef
sounds (Radford et al., 2008). Also, the aquatic invertebrate larvae, which were the same size
as the Daphnia, reacted to natural sounds and sounds from human activities, and their
swimming activities have changed accordingly (Morley et al., 2014). Daphnia is used as a
model in ecological (Stollewerk, 2010), physiological (Altshuler et al., 2011), genetic (Harris
et al., 2012; Miner et al., 2012), toxicological (Shaw et al., 2008), and parasitological (Ebert,
2008) studies.

In the present study, as a follow up or our recent study (Shafiei Sabet et al., 20215), we
experimentally tested the hypothesis that whether experimental sound exposure ensonified by
an underwater speaker affect the general swimming activities and foraging behaviour of
zebrafish as predator upon water flea as prey item under laboratory conditions. Our specific
goals were: firstly, to assess the effect of experimental sound exposure on zebrafish activities
including swimming speed and spatial displacement. Secondly, to estimate whether the
temporal pattern of sound exposure matters and affects differently zebrafish behaviour. And

thirdly, to verify our recent laboratory-based findings of sound impacts on zebrafish
swimming activity and foraging behaviour.

2. Materials and Methods

This study was performed in the ornamental fish breeding facility center at Fisheries
Department, Faculty of Natural Resources, University of Guilan, located in Sowmeh Sara
city, Guilan province, Iran (37°17□39□N, 49°19□55□E), using an aquarium with
dimensions of 50 cm length ×15 cm wide ×20 cm high with a volume of water intake of
112.5 liters in the period of 10:00 to 14:00 every day. Zebrafish (approximate age of 45 days
old and of the wild-type, short-fin variety weight (± standard deviation) of 1.23 ± 0.02 g)
were obtained from an ornamental fish breeding center located in Bazar-Jomeh in Sowmeh
Sara county, Guilan province, Iran.

Zebrafish were stored in a stock glass tank with dimensions 50 cm long ×30 cm wide ×40 cm
high for two weeks and adapted to environmental conditions to reduce possible stress and
hormonal changes due to transportation, captivity conditions and animal welfare issues
(Deakin et al., 2019). The fish were fed 0.8 mm commercial Biomar® feed until the day
before the experiment (Neo et al., 2015). Waterfleas were caught every morning, during the
whole experiment days, from the surface layer of the pool of the faculty, by a plankton net
with a net mesh size of 0.2 mm. This was done by horizontal twisting at a depth of 10 cm and
a length of 30 cm and was kept overnight in a separate tank at the same temperature as the
test tank to adapt similarly mimicking to the temperature conditions. According to the
previous study of Shafiei Sabet et al. (2019) as well as the same sampling location and depth
in the same time period with present study and the identification key used in the previous
Daphnia species study, *Daphnia magna* was identified (Shafiei Sabet et al., 2019).

2.1. Sound treatments

In the present experiment, four sound treatments with different temporal patterns along with
control treatment were used, including the first treatment as control treatment in which the
fish were exposed to ambient noise (AN). Second treatment, Continuous sound (CS) (Fig. 1
(a)), third treatment, regular intermittent noise (IN) with fast pulse rate (1:1), fourth
treatment, regular intermittent sound with slow pulse rate (1:4) and Fifth treatment is the

irregular intermittent sound (1:1-7). All three intermittent treatments include one second of

sound, but the difference between these sound treatments is the intervals between these

sounds (silence time), which are described in detail below. Regular intermittent sound with a

fast pulse rate (1:1) involves one second of sound and one second of silence (Fig. 1 (b)).

Regular intermittent sound with slow pulse rate (1:4) consists of one second of sound and

four seconds of silence (Fig. 1 (c)) and irregular intermittent sound (1: 1-7) includes one

second of sound and interval 1, 2, 3,4 ,5, 6 or 7 seconds of silence is random (Fig. 1 (d)).

Also, the sound treatments that are broadcast for different fish on different days were

identified quite randomly.

The sound treatments were generated and the sound files modified using Audacity® software

(2.3.1) at the sound frequency that can be detected and heard for zebrafish (300-1500 Hz)

(Higgs et al., 2002) as well as the bandwidth of anthropogenic sounds, including vehicles,

pump systems, and similar pile driving that overlap (Slabbekoorn et al., 2010). The designed

sound was produced by the software in the same sound frequency range of 400-2000 Hz

based on our earlier study (Shafiei Sabet et al., 202).

2.2. Experiment Tank

The experimental tank with dimensions of 50×20×15 cm with black background was

prepared to increase the contrast between Daphnia and fish in the video file. The

experimental fish tank was filled with water to a level of 15 ± 0.5 cm of filtered and aerated

dechlorinated tap water after overnighting. During the broadcast of the sound treatments, the

experiment tank was filmed by a video camera (Panasonic HC-V180 Full HD 28 mm Wide

Lens Camcorder) at a distance of about 50 cm from in front of the test tank. After production

of sound treatments, they were played back as stereo WAV files using a player connected

(Sony Vaio SVF1421A4E Laptop) to an underwater speaker (custom-build speaker in Iran,

8Ω, 30 W, 10 Hz- 10 KHz).

In this experiment, a divider plate was placed transversely in the tank and the tank length was

halved (25×20×15 cm) in order to increase the enclosure on the zebrafish swimming

environment and make the entire fish swimming space visible for video-recording (Fig. 2). In

order to reduce the stress of the fish and also to reduce the effects of people moving to the

test site, the test tank was surrounded by black plastic so that the fish's behaviour was not

affected by other factors and only the camcorder lens passed through the plastic and was the

same for all treatments and repetitions. In such a way that when disconnecting and
connecting the camcorder between sound treatments, the fish is not seen and is not affected.
To isolate the experiments from external sounds and reduce unwanted sounds, we placed and
covered all walls at the entrance to the experimental room and also inside walls using egg
boxes.

The physicochemical properties of the water were the same and monitored regularly on a
daily basis. The photoperiod used in the experiment was 12 hours of light and 12 hours of
darkness (Higgs et al., 2002; Villamizar et al., 2014) and the light intensity measured in this
experiment by the light meter model (TES_1336A – TES Electrical Electronic Corp. Taiwan)
averaged 62 lux. Water temperature was measured during the experiment and was maintained
at 26±1 °C. And also, the amount of dissolved oxygen (DO) in the water was measured and
was maintained at 8±1 mg / L.

The underwater speaker used in the experiment was placed horizontally on the other side of
the separator plate (See Fig. 2). In this experiment, after introducing the fish to the test tank
during the night for about 20 hours, the fish was given the opportunity to adapt to the
environment so that it could use the entire tank space for swimming and display natural
swimming behaviour (Shafiei Sabet et al., 2015) and have the normal conditions (Neo et al.,
2015; Shafiei Sabet et al., 2015). Then, test was performed with a video camera located in
front of the tank. After ten minutes, the sound treatment was played by a speaker and a sound
player for 20 minutes, However, the food item (Daphnia) and non-food item (Duckweed)
added to the experiment tank after ten minutes from sound playback. The nutrition of the
predatory species was investigated. This same process was performed for the other sound
treatments with a 15-minute interval between treatments and the fish was exposed to all five
acoustic treatments and repeated the next day for the next fish (Fig. 3). The order of broadcast
of sound treatments on a daily basis was randomly balanced. We used a similar approach
previously described to quantify behavioural indices of zebrafish and acoustic measurements
of tank in this system (Shafiei Sabet et al., 2015).

2.3. Underwater sound measurement

In order to check the intensity of the sound that can be played by the underwater speaker and to understand whether the tested fish was able to detect and perceive the sound treatments or not, first the continuous treatment sound file was played back using a laptop sound player (Sony Vaio SVF1421A4E) connected to a custom-build underwater speaker which a custom-build sound tuning amplifier was attached. The level of sound intensity under water was recorded by a hydrophone model (Aquarian Scientific AS-1) which connected to the amplifier model (PA-4) and a Tascam linear PCM recorder model (DR-100MKII). The recorded sound file was evaluated in R studio software (Version 1.1.456 - © 2009-2018 RStudio, Inc.).

According to the Fig. 4, during continuous sound playback, the frequency range emitted from the underwater speaker was completely in the zebrafish's hearing range in the range of 300-1500 Hz (Higgs et al., 2002) and far above the ambient noise playback. The sound pressure of continuous sound treatment during playback was average 121 dB re 1 μPa^2/Hz for 5 seconds and the ambient sound pressure was average 96 dB re 1 μPa^2/Hz for 5 seconds.

2.4. Effect of sound on Zebrafish swimming behaviour

To investigate the effect of sound on the behaviour of Zebrafish, 30 fishes (15 males and 15 females) were introduced individually in the experimental tank after biometrics. The fish were introduced to the test tank in last day (Overnighting) and the fish were given the opportunity to get used to the environment so that they could use the entire space of the tank for their swimming and have normal conditions. Also, all fish were given 30 minutes to relax. Since then, the fish has undergone five sound treatments.

Zebrafish behavioural response to five sound treatments was video recorded for a maximum of 30 minutes for each treatment (maximum 20 minutes for exposure and a maximum of 10 minutes before exposure) (Fig. 3). Swimming behaviour parameters such as startle response (which is the peak of swimming speed of fish more than 10 cm per second that occurs immediately after the sound was played for one minute (See Shafiei Sabet et al. (2015)), brief swimming speed (5 seconds before and 5 seconds during sound) and prolonged swimming speed (one minute before and one minute during sound) were evaluated for all treatments.

Also, to explore spatial distribution of fish in the tank, in the vertical/column profile,
according to the dewatering height of 15 cm of the test tank, tank height during inspection
and analysis behavioural data was divided into two parts: zero to 7.5 cm and 7.5 to 15 cm. To
check the distance from the sound source in the horizontal profile, the length of the tank was
divided into three parts: zero to 8.33 cm, 8.33 to 16.66 cm and 16.66 to 25 cm.

2.5. Effect of sound on foraging behaviour of zebrafish on water fleas

As mentioned earlier (Fig. 3), the effect of sound on the foraging behaviour of zebrafish was
investigated in such a way that 5 waterfleas (about three millimeters) as a prey species
(target) and 5 non-food substances as non-food item in the same size as Daphnia (about three
millimeters) was mixed in 25 ml beaker and added to the fish tank in the same manner for all
treatments. The waterfleas were in the same sizes that caught with plastic Pasteur pipettes to
decrease damaging water, which is suitable for feeding this species of fish at puberty and can
be received by the mouth of the fish (Shafiei Sabet et al., 2015). Naturally, both of food and
non-food items are present in the habitat of this fish.

Foraging performance was assessed as follows: To investigate the effect of sound treatments
on the foraging power of zebrafish, the parameters of food discrimination error (Formula 1)
and food handling error (Formula 2) were measured.

$$\text{Food Discrimination Error} = \frac{\text{number of non-food item attacks}}{\text{total number of food and non-food attacks}} \qquad \text{(Formula 1)}$$

$$\text{Food Handling Error} = \frac{\text{number of unsuccessful attacks on water fleas}}{\text{total number of successful and unsuccessful attacks on water fleas}} \qquad \text{(Formula 2)}$$

2.6. Behavioural information processing and statistical analysis

Recorded videos of zebrafish behaviour were converted to 10 frames per second by Xilisoft
Video Converter Ultimate software to reduce the magnification of time, in order to increase
the accuracy of the fish swimming survey and also to reduce the fish speed for spatial
inspection per second. Logger Pro® video tracking software (Vernier Software, Beaverton,
OR, USA, version 3.10.1) was used to examine behavioural responses including the number
of explosive movements, swimming speed, and spatial distribution of the fish. Entering
information and data in M.S. Excel 2016 and data analysis was performed using SPSS 25
software. The normality of the data was evaluated by Kolmogorov-Smirnov test and the

homogeneity of the data by Levene test. Then, the presence or absence of significant differences between the mean of the data was assessed by repeated measures ANOVA analysis and using Tukey multi-range test. A HuynheFeldt correction was performed when sphericity could not be assumed in the repeated measures ANOVA. Bonferroni corrected post hoc tests were performed when ANOVA test results were significant. The level of significance in this study was considered P<0.05. A custom-written acoustic calibration script in R studio software (Version 1.1.456 - © 2009-2018 RStudio, Inc.) was also used to evaluate sound pressure levels and power spectral density that were played by the underwater speaker.

3. Ethical note

We considered the 3Rs in behavioural research, the guidelines with respect to Good Laboratory Practice (GLP). All housing, handling and experimental conditions were in accordance with the guidelines for the treatment of animals in behavioural research and teaching (ASAB, 2020). Water fleas and zebrafish were allowed to acclimatize gradually to the laboratory conditions before they were used in any of the experiments and showed no signs of adverse effects of the experimental conditions. Zebrafish showed only a brief startle response with the onset of the moderate sound playbacks and no mortalities or physical damages were observed during experiments (Neo et al., 2015; Shafiei Sabet et al., 2015). There are no legal requirements for studies involving invertebrates such as waterfleas (Daphnia) in Iran. The principal investigator (S.SH.S) passed successfully the exam for the course on laboratory animal science at Leiden University, the Netherlands and holds animal testing act certificate "as an Article 9 researcher".

4. Result

4.1. Impact of sound on swimming behaviour of Zebrafish

Experimental sound exposure has changed zebrafish swimming activities in different ways. Sound treatments significantly increased the parameters of startle response and swimming speed of fish (Fig. 5a, b, c). The number of startle response instantly increased in exposure to sound treatments so that all sound treatments showed a significant difference compared to the ambient condition (repeated measures ANOVA: $F_{3.23, 93.68}$=6.31, P=0.000435), But no significant differences were observed in terms of temporal patterns between sound treatments (P>0.05) (Fig 5a). Also, in the brief swimming speed difference (5 seconds before the sound

and 5 seconds during the sound) there was a significant difference compared to the ambient

condition (repeated measures ANOVA: $F_{3.06,88.70}=11.17$, $P=0.000002$), although there was no

significant difference between the sound treatments with different temporal patterns in this

time period ($P>0.05$) (Fig 5b). This difference in swimming speed in exposure to sound

treatments was also true in prolonged (60 seconds before the sound and 60 seconds during the

sound), so that there was a significant affect between sound treatments compared to ambient

conditions (repeated measures ANOVA: $F_{3.39,98.34}=7.72$, $P=0.000054$), however again there

was nonsignificant difference between sound treatments ($P>0.05$) (Fig 5b).

According to Figure 6, in four sound treatments (CS, 1:1, 1:4, 1:1-7), a sudden increase in

fish swimming speed was observed once sound treatments were played at 60 seconds.

Observations also showed that in all sound treatments, fish reached the baseline after 60

seconds during sound. In ambient condition (Fig 6a), no significant difference in swimming

speed was observed in any of the time periods. While in continuous sound and regular

intermittent 1:1 treatment (Fig 6b, c), a significant difference was observed in the comparison

of 5 seconds before sound playback and the first 5 seconds of sound playback (CS= repeated

measure ANOVA: $F_{4.54,131.66}=9.53$, $P\leq0.001$, 1:1= repeated measure ANOVA:

$F_{3.85,111.72}=11.72$, $P\leq0.001$). This is while in this treatment, compared to the last 5 seconds

before the sound and the last 5 seconds of the sound, fish reached the base line and there was

no difference ($P>0.05$). Also, this difference was observed in the two treatments regular

intermittent 1:4 and irregular intermittent 1:1-7 (Fig 6d, e) with more intensity compared to

before and during the sound (1:4= repeated measure ANOVA: $F_{6.24,181.01}=14.17$, $P\leq0.0001$,

1:1-7= repeated measure ANOVA: $F_{7.14,207.15}=15.01$, $P\leq0.0001$). But in comparing the

baseline time in these two treatments, no significant difference was observed and the fish

reached the baseline ($P>0.05$).

Changes in the spatial distribution of Zebrafish were investigated when exposed to sound

treatments in vertical scale (lower layer) (Fig. 7a, b) and horizontal scale (X position) (Fig.

8a, b). According to Figure 7a, the average percentage of fish time spent in the lower layer of

the tank during sound treatments exposure over a brief time (15 seconds before sound and 15

seconds during sound exposure) there was no treatment effect ($F_{3.77,109.47}=1.486$, $P=0.217$) or

interaction for treatment*times ($F_{3.55,102.92}=0.634$, $P=0.621$). But there was effect of time

($F_{1,29}=28.274$, $P=0.000011$). In two treatments of ambient (AN) and irregular intermittent

sound (1:1-7) with before sound exposure was not significant effect (AN= $F_{1,29}=28.274$,

$P=0.104$, 1:1-7= $F_{1,29}=28.274$, $P=0.051$). But there was a significant difference between two

sound treatments: continuous sound (CS) and 1:4 during and before exposure to sound (CS= 378

$F_{1,29}$=28.274, P=0.007, 1:4= $F_{1,29}$=28.274, P=0.004) and also 1:1 treatment was a significant 379

difference with before the sound exposure ($F_{1,29}$=28.274, P=0.011). According to Figure 7b, 380

the average percentage of fish time spent in the lower layer of the tank during sound 381

treatments over a prolonged time (60 seconds before sound and 60 seconds during sound 382

exposure), there was no treatment effect ($F_{3.65,105.92}$=1.837, P=0.133) or interaction for 383

treatment*times ($F_{4,116}$=1.780, P=0.137). But there was effect of times ($F_{1,29}$=35.398, 384

P=0.000002). In control treatment (AN) between before and during sound exposure do not 385

show a significant difference ($F_{1,29}$=35.398, P=0.975). While in continuous treatment (CS) 386

and 1:4 between before and during sound exposure showed a significant difference (CS= 387

$F_{1,29}$=35.398, P=0.000411, 1:4= $F_{1,29}$=35.398, P=0.000360) and 1:1 treatment was significant 388

difference between during exposure and before time ($F_{1,29}$=35.398, P=0.008). Also, irregular 389

intermittent sound treatment (1:1-7) between during and before sound exposure was 390

significant difference ($F_{1,29}$=35.398, P=0.042). But in the horizontal profile (X position) 391

there was no significant difference between sound treatments and control conditions, both in 392

the brief and prolonged time ($F_{4,116}$=1.369, P=0.249) (Fig 8a) ($F_{3.78,109.71}$=1.810, P=0.136) 393

(Fig 8b). Also, according to Figure 9a, b, the effects of different sound patterns on the spatial 394

distribution of fish can be seen as a heat map in brief and prolonged duration. Acoustic 395

treatments affected changes in swimming pattern from the top layer to the bottom layer but 396

had no effect on the distance from the sound source (right side of the tank). 397

4.2. Impact of sound on foraging performance of Zebrafish 399

According to Figure 10a, none of the acoustic treatments showed a significant difference 400

compared to the silent treatment on zebrafish food discrimination error (repeated measure 401

ANOVA: F4,116=1.339, P=0.260). In fact, there was no food discrimination error between 402

food and non-food item by broadcasting sound treatments compared to control treatment. 403

However, all acoustic treatments showed a significant difference compared to the control 404

treatment (repeated measure ANOVA: F2.82,81.91=26.023, P≤0.001) but no significant 405

difference was observed between sound treatments (P>0.05) (Fig. 10b). This means that with 406

the broadcast of acoustic treatments compared to the control treatment, there was a significant 407

handling error in food intake in the food. 408

5. Discussion

Our results unequivocally demonstrate that acoustic stimuli affect zebrafish behaviour in different ways. In this experiment, zebrafish swimming behaviour indices such as the number of startle response, the difference in brief swimming speed (within 5 seconds), the difference in prolonged swimming speed (within 60 seconds) and the spatial distribution of zebrafish such as the percentage of fish in the lower layer of the tank and horizontal spatial distribution in response to continuous and intermittent sound patterns were examined. The results showed that the number of startle response indicating anxiety in zebrafish and other aquatic species (Blaser et al., 2010; Maximino et al., 2010), in different sound treatments, showed a significant difference compared to control condition. However, these changes did not show a significant difference between sound treatments. Also, in another part of the results, the difference between brief and prolonged swimming speeds in all sound treatments compared to the control condition was significant. However, these changes did not show a significant difference between sound treatments. Moreover, interestingly like what we have shown in our earlier study (Shafiei Sabet et al., 2015), we have seen the same impact of sound exposure on foraging performance in zebrafish as all will be discussed further as follows in the next sections.

5.1. Startle responses as a specific indicator of moderate anxiety in zebrafish?

Startle response is an involuntary action that is controlled by a pair of brain neurons in the Mauthner (M-) cells in the mesencephalon and play a major role in the decision-making process (Eaton et al., 1977; Eaton et al., 1991; Mirjany et al., 2011; Zottoli, 1977). Increasing the intensity of the sound triggers a response by involuntary Mauthner cells in the mesencephalon, leading to an involuntary escape response in fish (Eaton et al., 1977; Eaton et al., 1991). Increasing the number of startle response and brief swimming speed of fish by playing sound treatments, causes behavioural responses related to fear and anxiety in this species. Previous studies have shown that sounds increase motor acceleration and startle responses in zebrafish (Neo et al., 2015; Shafiei Sabet et al., 2015). However, Shafiei Sabet et al. (2015) using an in-air speaker as a sound source reported that the difference in the number of startle response between continuous and intermittent (1:1) treatments with the ambient treatment was not significant which is not consistent with the results of this study. The reason of this difference in the occurrence of stress-related swimming behaviour could be due to differences in the background sound intensity before the test, differences in the method of

ensonifiying of the fish tank; the sound source (speaker) under water or in-air, different

storage conditions of fish and also differences Genetics and individual characteristics in

zebrafish.

Studies by other researchers have shown that other species of fish respond similarly to sudden

swimming behaviour in response to sound exposure. European minnow (*Phoxinus phoxinus*)

and sticklebacks (*Gasterosteus aculeatus*) also showed a significant increase in the number of

startle response in the face of sound, which is similar to the results of this study (Purser and

Radford, 2011; Voellmy et al., 2014a). Startle response at the onset of sudden sound exposure

is a common behavioural feature in fish kept in captivity and in the laboratory conditions. Of

course, fish in the open and natural conditions can also show behavioural responses related to

fear and anxiety (Neo et al., 2016; Staaterman et al., 2020). Spiga et al. (2017) stated that

European seabass (*Dicentrarchus labrax*) also showed a higher number of startle response in

continuous and intermittent sound treatment against ambient (control) treatment.

Startle response by prey fish is a behavioural response to increase survival in predator-prey

relationships (Webb, 1986). By hearing the sounds of predator fish and receiving sound

signals related to the attack, the prey fish starts swimming at high speed and explosively in

the opposite direction of the perceived sound in order to increase the success rate of escaping

and staying away from the predatory species. Sounds can affect the prey fish's decision-

making power against sound sources or danger, the way in which prey assesses risk (Dukas,

2004), and the loss of focus and lack of appropriate response to danger (Chan et al., 2010;

Simpson et al., 2015). It has been suggested that increasing sound levels can potentially

impair the perception of danger by predatory fish species (Slabbekoorn et al., 2010).

Involuntary and acquired behavioural responses related to fear and anxiety are associated

with the potential for the presence of danger (Blaser et al., 2010; Maximino et al., 2010). The

quality and quantity of behavioural responses of fish in captivity and in vitro to brief and

severe stress stimuli are different from those of fish living in the habitat, reducing the

behavioural responses in the habitat and in the wild (Malavasi et al., 2004). One of the

reasons for these differences could be due to the most ability of fish to respond in the wild

before reaching the stimulus threshold, so that these fish have a longer time to make decisions

and escape from the danger zone by hearing and perceiving closely the sounds associated

with the predator species and they have more space available than controlled laboratory

environments. Another reason could be the high level of basal stress potential in controlled

laboratory environments, which with additional stress due to the perception of the predator

species leads to an increase and intensification of total stress in the prey species and more
intense responses are shown.

In addition to behavioural responses, increasing sound levels can also affect physiological
responses in the laboratory and in the natural habitat of fish. The study of Spiga et al. (2017)
and Radford et al. (2016) showed that sound exposure had a significant effect on the number
of opening and closing of gills and thus on the gill ventilation of European seabass
(*Dicentrarchus labrax*) compared to the control treatment. This increase in oxygen demand
by European bass (*Dicentrarchus labrax*), which is accompanied by an increase in gill
ventilation and the opening and closing of gill operculum, indicates an increase in stress
levels. Santulli et al. (1999) showed that blood biochemical parameters including cortisol and
glucose in European seabass (*Dicentrarchus labrax*) increased in sound treatments compared
to the ambient treatment. Staaterman et al. (2020) stated that anthropogenic sound treatments
in the natural environment also have the potential to affect stress-related physiology in coral
reef fish, so that the amount of cortisol in sound treatments was significantly increased
compared to the control treatment.

5.2. Sound impacts on the behavioural tolerance in swimming activities

In the experiment performed on zebrafish, different sound patterns had a significant effect on
the brief and prolonged swimming speed of the fish compared to the ambient treatment,
which is consistent with the observations of the present study (Neo et al., 2015; Shafiei Sabet
et al., 2015; Shafiei Sabet et al., 2016a). In the experiment of Neo et al. (2015) with a group
of 5 zebrafish in each treatment, in comparison with increasing the swimming speed of fish
with the ambient treatment, as in the results of this study, the intermittent (1:1) treatment had
a higher mean speed and significant difference than other treatments. Also, with onset of
sound exposure, a significant difference in swimming speed was observed in cod fish (*Gadus*
morhua), which was consistent with the results of this study (Handegard et al., 2003). There
was also a significant difference in the swimming speed of European seabass (*Dicentrarchus*
labrax) compared to the silent conditions (Neo et al., 2018). The prolonged swimming speed
of the zebrafish weakened with repeated exposure suggesting a habituation process. Zebrafish
might experience negative consequences if they are disturbed and do not habituate contrary to
sound exposure. Shafiei Sabet et al. (2016b) by comparing the effect of sound on the
swimming behaviour of two species of The Lake Victoria cichlids (*Haplochromis piceatus*)

and zebrafish (*Danio rerio*) have showed that the application of sound treatments reduced the

swimming speed of cichlids (*Haplochromis piceatus*) and increased the swimming speed of

zebrafish (Shafiei Sabet et al., 2016b). The reason of the difference in the swimming speed of

cichlids (*Haplochromis piceatus*) can be related to species-specific behavioural responses in

response to acoustic stimuli, genetic characteristics and habitat conditions.

5.3. The effect of sound on the spatial distribution of zebrafish (Vertical/Horizontal)

Studies on the stress indices of zebrafish in the face of different sound patterns showed that

with the onset of sound treatments, the spatial distribution of fish changes and the fish shows

a greater tendency to swim in the lower layer of the aquarium environment. Also, the study

on the percentage of fish in the lower layer of the test tank showed that in some sound

treatments in the brief time there was a significant difference and in the prolonged time in all

sound treatments except the ambient treatment, this means Which had a greater tendency to

be present in the lower layer when playing the sound of fish. In another part of the results of

this study and the study of the spatial distribution of the presence of fish in the horizontal

profile (X position) also showed that in none of the sound treatments the fish did not tend to

distance from the sound source and did not show significant differences.

Neo et al. (2015) in the study of the effect of sound on the spatial distribution of zebrafish,

found that zebrafish with the beginning of broadcasting sound treatments showed startle

response and increased brief swimming speed and the spatial distribution of zebrafish

changed so that fish was more inclined to and they swam in the top and surface layers of the

test tank. Also, in this study (Neo et al. 2015), there were no observations of the freezing and

standing of the fish in the lower layer of the tank. However, the results of the present

experiment in the spatial distribution of zebrafish, the percentage of time fish staying in the

lower layer of the test tank was higher, which is contrary to the report of Neo et al. (2015).

One of the reasons for this difference in the results of vertical spatial distribution could be the

amount of sound intensity emitted in the treatments used in these two studies. The intensity of

sound emitted in acoustic treatments was equal to 112 dB re 1µ Pa, which is less than the

intensity of sound in this study (121 dB 1µ Pa). Therefore, a significant increase in sound

intensity and high sound level difference between sound treatments and ambient treatments

can lead to different responses in fish. In another study, Neo et al. (2018) designed an

experiment to investigate the effect of different sound patterns on the behaviour of European

seabass and found that anthropogenic sounds increase the swimming depth of European

seabass and distance from the sound source, which results in behavioural responses is
consistent with present study in lower layer results.

Other fishes have also shown spatial distribution changes in response to acoustic stimuli. In a
field study (Kok et al., 2021) have shown that bottom-moored echosounders, representative
of a high intensity impulsive intermittent anthropogenic sound, affect the abundance,
schooling cohesion behaviour and swimming depth of pelagic fish. Two recent telemetry
tagging studies demonstrated the effects of another intermittent source of sound, seismic
surveys, on free-ranging benthic fish species. Bruce et al. (2018) showed shift of diurnal
activity patterns and general swimming speed in eight tiger flatheads (*Neoplatycephalus*
richardsoni). van der Knaap et al. (2021) revealed also demonstrated in a field study that
Atlantic cod (*Gadus morhua)* left noisy area more quickly and decrease their swimming
activity rather than ambient sound condition as control treatment. They also have shown that
seismic surveys can disrupt diurnal activity cycles which may have long term negative
consequences on this species foraging success and fitness.

The swimming of zebrafish towards the upper layer at the beginning of the sound
transmission has been interpreted as curiosity and searching behaviour, as the authors'
experimental observations have shown that by opening the door of the zebrafish test saloon
and walking the staff to perform feeding fish usually produce low-pitched sounds, which
attract the attention and curiosity of zebrafish and show the highest distribution at the water
column level for feeding activities (Shafiei Sabet et al., 2015), However, the response of
changing the spatial distribution of fish to the depth and lower layer with the beginning of
sound treatments indicates the occurrence of stress and fear in fish, which is similar and
expressed in studies of other researchers on other fish species (Neo et al., 2018; Sarà et al.,
2007). Examination of fish behaviour to other stimuli including chemicals and fear extract
has also shown that with the release of chemicals and fear extract, fish move to the lower
layer (deep) column and this pattern of spatial distribution is a behavioural indicator of fear in
many interpreted fish species (Gerlai et al., 2000; Gerlai et al., 2006).

In addition, another reason for the difference in the vertical distribution behavioural results
observed in the study of Neo et al. (2015) and the present study could be the difference in the
use of speakers, such as the use of in air speaker in the previous study (Neo et al., 2015) and
the use of underwater speaker in the present study. The use of speakers in air to broadcast
sound treatments leads to the production of more sound intensity in the deeper parts than in

the middle and the surface of the water in the aquarium tank, which may cause the fish to
move and escape towards the upper and surface layer where less sound intensity is felt
(Shafiei Sabet et al., 2015).

In the study of Shafiei Sabet et al. (2015), they investigated the effect of acoustic treatments
with sound intensity almost similar to this study (122 dB re 1μ Pa) on the spatial distribution
of zebrafish showed that it is not consistent with the present study. The reason for this
difference in spatial distribution behavioural response could be the use of in air speaker in the
study by Shafiei Sabet et al. (2015) and underwater speaker in this study, as well as the
complexity of sound distribution patterns and sound gradients in aquarium environments
(Campbell et al., 2019) and other factors include the size of the test tank, the life cycle and
location of the fish storage tank, as well as differences in the species and genetics of the fish
species. Also in the study of Shafiei Sabet et al. (2015) in the spatial distribution of fish
horizontally, a similar result was shown with the results of the present study and no
significant difference was observed in the x position.

According to the available sources for measuring sound intensity and scattering patterns in
aquarium environments, an experiment was designed by Parvulescu (1967) and Akamatsu et
al. (2002) which shows the complexity and variability of sound scattering patterns and sound
gradients in enclosed aquarium tank environments. This fact indicates the limitations of
studying the spatial distribution of aquatic animals in enclosed and controlled environments
that must be considered. Therefore, in order to study the distribution patterns of fish and other
aquatic species more accurately, it is recommended to conduct field studies in natural
environments of animal species in order to obtain a more accurate and complete
understanding of the manner and patterns of sound-dependent distribution in aquatic species.

5.4. The importance of particle motion in fish tanks; behavioural observations for future
works

In order to understand the behavioural changes of zebrafish in response to sound, first of all,
it is very important to understand how the species detects and processes, and how it
behaviourally responds to sound (Hawkins and Popper, 2020). Because the auditory system
of fishes evolved primarily to detect particle motion, many fishes are most sensitive to
particle motion and they can use it to determine the direction of a sound source (Hawkins and
Popper, 2018; Popper and Hawkins, 2018; Sand and Bleckmann, 2008; Sisneros and Rogers,
2016). Only some of them, including the zebrafish, are sensitive to sound pressure as well as

the particle motion (Popper and Fay, 2011; Popper and Hawkins, 2018). There are some
studies revealing directional hearing and sound source localization in fish under laboratory
conditions and in free sound fields. Schuijf (1975) proposed that the cod determined sound
direction by monitoring the particle motion of the sound field, presumably employing the
directional orientation of the inner ear sensory cells (Dale, 1976). Although, Schuijf (1975)
also concluded that the direction of only particle motion may be insufficient to determine the
direction of a sound source. It has already been shown that cod could discriminate between
signals coming towards the head as compared to those coming towards the tail (Buwalda et
al., 1983; Schuijf and Buwalda, 1975). They argued that directional hearing might involve
both comparing the responses of hear cells oriented in different directions and also analysis of
the phase relationship between the sound pressure and particle motion to eliminate any
remaining 180^0 ambiguities (Schuijf, 1976).

5.5. How important it is particle motion to fishes and invertebrates

We did not mention the levels and direction of the particle motion that is generated within the
fish tank. Therefore, we believe it is premature to conclude that zebrafish cannot localize
sound source in our experimental set up. One might be because we know very little about
hearing in fishes only over 120 species of the more than 33000 known fish species (Ladich
and Fay, 2013) and that the empirical and theoretical work on sound source localization and
directional hearing in fishes have been contradictory and obscure for decades (Sisneros and
Rogers, 2016). Moreover, some explanations would be that practically because it is difficult
to monitor particle motion in fish tank, the lack of easily used and reasonably priced
instrumentation to measure particle motion, lack of sound exposure criteria for particle
motion and finally lack of particle motion measurement standards (Popper et al., 2014).

Within an aquarium tank the levels of particle motion are often highest at the water surface,
and close to the tank walls, when an underwater loudspeaker is used (Jones et al., 2019).
Although, resonant frequencies and reverberation may influence propagation and spectro-
temporal structure of received acoustic stimuli in fish ranks (Jones et al., 2019). Our fish
moved towards the lower levels of the tank, which may be because the particle motion levels
were highest close to the water surface, and lower at the bottom of the tank. It is always
important to monitor the particle motion when examining the effects of sounds upon fishes
and invertebrates (Nedelec et al., 2016; Popper and Hawkins, 2018). Moreover, Invertebrates
are especially sensitive to substrate vibration (Aimon et al., 2021; Hawkins et al., 2021;

Morley et al., 2014; Roberts et al., 2016), and some fish are too. Particle motion measurement

may play an important role in answering crucial biological and ecological questions relating

to fishes and other species among taxa (Nedelec et al., 2016). Thus, in doing future

experiments to explore anthropogenic sound impacts on the behaviour of fishes or

invertebrates under laboratory conditions it is necessary to develop open source and

accessible protocols for monitor both particle motion on three axes and sound pressure.

5.6. Acoustic stimuli trigger foraging performance modifications negatively in zebrafish

In the present study, the parameters of fish foraging behaviour such as food discrimination

error and food handling error were examined. The results of this experiment showed that the

zebrafish did not show any significant difference in the food discrimination error when

exposed to sound treatments compared to the ambient treatment. Also, the results of food

handling error showed that all sound treatments showed a significant difference compared to

the ambient treatment, but these sound treatments did not show a significant difference

compared to each other. Here we confirmed our earlier findings (Shafiei Sabet et al., 2015)

that sound impacts may goes beyond single species. Experimental sound exposure causes

more food handling errors and foraging in zebrafish as predator, which led to more survival

in waterflea as prey and avoiding from being eaten by a predator in noisy conditions.

In a study by Purser and Radford (2011), they found that the broadcasting of sound

treatments significantly affected the foraging performance of stickleback and food

discrimination error and food handling error increased significantly compared to the ambient

treatment and reduced foraging performance, which in the food discrimination error, it is not

consistent with the results of the present experiment, and one of the reasons for this

difference is the difference in the physiology of fish and its diet. Also, the possible difference

in the physiology of fish visual sense can be one of the factors influencing these differences,

but in the food handling error, a similar result was observed with the results of the present

experiment.

In other studies, by Voellmy et al. (2014b), the results showed a significant difference in the

number of unsuccessful takes of Daphnia in stickleback and no significant difference in

minnow fish, which can generally indicate the physiological difference between the two

species and possibly the difference in visual sense between the two species. The minnow fish,

which belongs to the Ciprinidae family which similar to zebrafish family, did not show a

significant difference in the unsuccessful takes of Daphnia, but in another species, a significant difference was observed.

In a study almost similar to the present experiment, Shafiei Sabet et al. (2015) designed an experiment to investigate the effect of different sound patterns on the swimming and foraging behaviour of zebrafish and found that the application of sound treatments caused a significant difference in fish handling error. There was no significant difference in food discrimination error that was consistent with the results of this experiment. Sound playback can be effective in identifying a substance as food and making decisions and attacking it. Since zebrafish are among the fish that have strong visual sense and use this sense to catch and hunt, so sounds can affect the ability to perceive potential vision.

It is also important to consider the potential impacts of acoustic stimuli in real field conditions. Although to date, there is a gap of knowledge and very few studies have investigated soundscapes in freshwater environments and the effects of elevated sound levels on fish in freshwater habitats (Bolgan et al., 2016; Linke et al., 2018; Rountree et al., 2020; Putland and Mensinger, 2020). Other studies under field conditions have also focused on the effects of sound on fish species. In a comparative study both in a field study in the Laurentian Great Lakes also under laboratory conditions Pieniazek et al., 2020 shown that experimental generated white noise and boat noise have changed swimming patterns and foraging behaviour of freshwater fish species in different ways and with highly variable patterns by family. They demonstrated that captive black bullhead, *Ameiurus melas*, foraged less and startled more when exposed to both white noise and boat noise treatments.

Our laboratory experiments were conducted to verify that acoustic stimuli affect swimming behaviour and foraging performance of zebrafish and establish hypothesis for future exploration to be conducted on other freshwater fish species and in the field.

6. Conclusion

The results of this study highlighted impacts of acoustic stimuli on a freshwater fish species and confirmed our earlier study on the same fish species (zebrafish) under laboratory conditions. Our findings show that the parameters of zebrafish swimming and foraging behaviour in different sound patterns were significantly different compared to control or ambient treatment. The results showed that sound treatments compared to the ambient

treatment caused a significant difference in the parameters of swimming behaviour including

the number of startle response, swimming speed of fish in the brief and prolonged time,

spatial distribution of fish in vertical and horizontal profiles. In general, the observed

behavioural patterns in response to sound treatments as a stressor, especially in the brief time

of this study have been observed in natural environments and in other fish species. Sound

pollution as a stressor in the brief and prolonged time can cause behavioural changes and

disturbances in the individual levels of aquatic species and have broad and important

repercussions on the communities of an ecosystem.

Also, the results obtained in the foraging behaviour showed that different sound patterns in

comparison with the ambient treatment caused a significant difference in food handling error

but no significant difference was observed in food discrimination error. Depending on the

species characteristics, these behavioural responses include escaping the predator species,

hiding in a shelter to avoid being hunted, forming crowded clusters, and even approaching the

predator species aggressively. The movement towards the lower layer observed in the

experimental tank by zebrafish can be due to the stress caused by the potential for the

presence of a predatory species, that in their natural habitats, where clear water is permeable

to light, to the deeper parts of their habitat, which have less light and have vegetation that

reduces the visibility of the predator species, and thus ensure its survival. In general, due to

the simultaneous presence of the species used in this experiment in their main habitat

(zebrafish, daphnia and duckweed) and the results obtained can be understood that sound

causes overlap in the water particles motion, which is one of the factors in the perception of

the prey species by the predator species, and the zebrafish cannot have a proper

understanding of the presence of the prey species. On the other hand, creating a state of

distress and anxiety after playing sound treatments has caused a lack of proper vision of the

prey species and the zebrafish, which is an omnivorous and visually impaired fish in prey,

cannot see and hunt well.

Increased background sound levels by human activities can be recognized as a stressor and

lead to a series of changes in the activities and swimming patterns of aquatic species. It

should be noted that the results of this study are obtained in captivity and under laboratory

conditions therefore the interpretation of the results should be done with caution and attention

on the conditions of natural environments in the various habitat and behavioural limitations

of any species.

Funding:

This work was funded by the Iranian Ministry of Science, Research and Technology, University of Guilan with the grant number 971511210004.

Credit author statement

Reza Mohsenpour: Conceptualization, Methodology, Software, Formal analysis, Visualization, Data curation, writing - original draft.

Saeed Shafiei Sabet: Conceptualization, Methodology, Supervision, Project administration, Writing - review & editing.

Data accessibility:

All data used for the analyses reported in this article and some videos are available from the figshare. Moreover, the data and some videos that support the findings of this article are available from the first author upon request.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments:

Thanks to Jeroen Hubert and Ozkan Sertlek for their guide and advice on hydrophone calibrations, editing R script versions and acoustic measurements. Special tanks to Hans

755   Slabbekoorn for providing accessibility to acoustic measurement devices. We also thank
756   Narjes Karimi for helping us in reviewing the results and graphs.

References:

759   Aimon, C., Simpson, S.D., Hazelwood, R.A., Bruintjes, R., Urbina, M.A., 2021. Anthropogenic
760   underwater vibrations are sensed and stressful for the shore crab Carcinus maenas. Environmental
761   Pollution 285, 117148. https://doi.org/10.1016/j.envpol.2021.117148.
762
763   Akamatsu, T., Okumura, T., Novarini, N., Yan, H.Y., 2002. Empirical refinements applicable to the
764   recording of fish sounds in small tanks. The Journal of the Acoustical Society of America 112, 3073-
765   3082. https://doi.org/10.1121/1.1515799.
766
Alexander, R. 1964. The structure of the Weberian apparatus in the Siluri, in Proceedings of the
Zoological Society ofLondon, Wiley Online Library, pp. 419–440. https://doi.org/10.1111/j.1469-
7998.1964.tb04507.x.
Altshuler, I., Demiri, B., Xu, S., Constantin, A., Yan, N.D., Cristescu, M.E., 2011. An integrated multi-
disciplinary approach for studying multiple stressors in freshwater ecosystems: Daphnia as a model
organism. Integrative and comparative biology 51, 623-633. https://doi.org/10.1093/icb/icr103.

Amorim, M.C.P., Vasconcelos, R.O., Fonseca, P.J., 2015. Fish sounds and mate choice, Sound
communication in fishes. Springer, pp. 1-33. https://doi.org/10.1007/978-3-7091-1846-7_1
André, M., Solé, M., Lenoir, M., Durfort, M., Quero, C., Mas, A., Lombarte, A., Van Der Schaar, M.,
López-Bejar, M., Morell, M., 2011. Low-frequency sounds induce acoustic trauma in cephalopods.
Frontiers in Ecology and the Environment 9, 489-493. https://doi.org/10.1890/100124.
778
779   Animal Behaviour. 2020. Guidelines for the treatment of animals in behavioural research and
780   teaching. Animal Behaviour. 159:1–11. https://doi.org/10.1006/anbe.1996.0293.
781
782   Barber, J.R., Fristrup, K.M., Brown, C.L., Hardy, A.R., Angeloni, L.M., Crooks, K.R., 2009. Conserving
783   the wild life therein: Protecting park fauna from anthropogenic noise.   26, 26-31.
784   http://www.nature.nps.gov/ParkScience/index.cfm?ArticleID=370&Page=1
785
786   Bermúdez-Cuamatzin, E., López-Hernández, M., Campbell, J., Zuria, I., Slabbekoorn, H., 2018. The
787   role of singing style in song adjustments to fluctuating sound conditions: a comparative study on
788   Mexican birds. Behavioural processes 157, 645-655. https://doi.org/10.1016/j.beproc.2018.04.005.
789
790   Blaser, R., Chadwick, L., McGinnis, G., 2010. Behavioral measures of anxiety in zebrafish (*Danio
791   rerio*). Behavioural brain research 208, 56-62. https://doi.org/10.1016/j.bbr.2009.11.009.
792
Bolgan, M., Chorazyczewska, E., Winfield, I.J., Codarin, A., O'Brien, J. and Gammell, M., 2016. First
observations of anthropogenic underwater noise in a large multi-use lake. Journal of
Limnology, 75(3), pp.644-651. doi: 10.4081/jlimnol.2016.1405.

Bracciali, C., Campobello, D., Giacoma, C., Sara, G., 2012. Effects of nautical traffic and noise on
foraging patterns of Mediterranean damselfish (*Chromis chromis*). PloS one 7, e40582.
https://doi.org/10.1371/journal.pone.0040582.

Bruce, B., Bradford, R., Foster, S., Lee, K., Lansdell, M., Cooper, S., Przeslawski, R., 2018. Quantifying
fish behaviour and commercial catch rates in relation to a marine seismic survey. Marine
environmental research 140, 18-30. https://doi.org/10.1016/j.marenvres.2018.05.005.
802
803 Budelmann, B.-U., 2011. Morphological diversity of equilibrium receptor systems in aquatic
804 invertebrates, Sensory biology of aquatic animals. Springer, pp. 757-782.
805 https://doi.org/10.1007/978-1-4612-3714-3_30 .
806
807 Buwalda, R., Schuijf, A., Hawkins, A., 1983. Discrimination by the cod of sounds from opposing
808 directions. Journal of Comparative Physiology 150, 175-184. https://doi.org/10.1007/BF00606367.
809
810 Cachat, J., Canavello, P., Elegante, M., Bartels, B., Hart, P., Bergner, C., Egan, R., Duncan, A., Tien, D.,
811 Chung, A., 2010. Modeling withdrawal syndrome in zebrafish. Behavioural brain research 208, 371-
812 376. https://doi.org/10.1016/j.bbr.2009.12.004.
813
814 Campbell, J., Shafiei Sabet, S., Slabbekoorn, H., 2019. Particle motion and sound pressure in fish
815 tanks: a behavioural exploration of acoustic sensitivity in the zebrafish. Behavioural processes 164,
816 38-47. https://doi.org/10.1016/j.beproc.2019.04.001.
817
818 Carroll, A., Przeslawski, R., Duncan, A., Gunning, M., Bruce, B., 2017. A critical review of the potential
819 impacts of marine seismic surveys on fish & invertebrates. Marine Pollution Bulletin 114, 9-24.
820 https://doi.org/10.1016/j.marpolbul.2016.11.038.
821
822 Casper, B.M., Smith, M.E., Halvorsen, M.B., Sun, H., Carlson, T.J., Popper, A.N., 2013. Effects of
823 exposure to pile driving sounds on fish inner ear tissues. Comparative Biochemistry and Physiology
824 Part A: Molecular & Integrative Physiology 166, 352-360.
825 https://doi.org/10.1016/j.cbpa.2013.07.008.
826
827 Chan, A.A.Y.-H., Giraldo-Perez, P., Smith, S., Blumstein, D.T., 2010. Anthropogenic noise affects risk
828 assessment and attention: the distracted prey hypothesis. Biology letters 6, 458-461.
829 https://doi.org/10.1098/rsbl.2009.1081.
830
831 Coquereau, L., Grall, J., Chauvaud, L., Gervaise, C., Clavier, J., Jolivet, A., Di Iorio, L., 2016. Sound
832 production and associated behaviours of benthic invertebrates from a coastal habitat in the north-
833 east Atlantic. Marine biology 163, 127. https://doi.org/10.1007/s00227-016-2902-2.
834
835 Cortemeglia, C., Beitinger, T.L., 2005. Temperature tolerances of wild-type and red transgenic zebra
836 danios. Transactions of the American Fisheries Society 134, 1431-1437. https://doi.org/10.1577/T04-
837 197.1.
838
839 Crawford, A.D., Esguerra, C.V., de Witte, P.A., 2008. Fishing for drugs from nature: zebrafish as a
840 technology platform for natural product discovery. Planta medica 74, 624-632. DOI: 10.1055/s-2008-
841 1034374.
842
843 Dale, T., 1976. The labyrinthine mechanoreceptor organs of the cod *Gadus morhua* l-(teleostei:
844 gadidae). a scanning electron microscopical study, with special reference to the morphological
845 polarization of the macular sensory cells. http://pascal-
846 francis.inist.fr/vibad/index.php?action=getRecordDetail&idt=PASCAL7650248453.
847

de Jong, K., Amorim, M., Fonseca, P.J., Heubel, K.U., 2018a. Noise affects multimodal communication
during courtship in a marine fish. Frontiers in Ecology and Evolution 6, 113.
https://doi.org/10.3389/fevo.2018.00113.

De Jong, K., Amorim, M.C.P., Fonseca, P.J., Fox, C.J., Heubel, K.U., 2018b. Noise can affect acoustic
communication and subsequent spawning success in fish. Environmental Pollution 237, 814-823.
https://doi.org/10.1016/j.envpol.2017.11.003.

de Vincenzi, G., Micarelli, P., Viola, S., Buffa, G., Sciacca, V., Maccarrone, V., Corrias, V., Reinero, F.R.,
Giacoma, C., Filiciotto, F., 2021. Biological sound vs. anthropogenic noise: assessment of behavioural
changes in *Scyliorhinus canicula* exposed to boats noise. Animals 11, 174.
https://doi.org/10.3390/ani11010174.

Deakin, A.G., Spencer, J.W., Cossins, A.R., Young, I.S., Sneddon, L.U., 2019. Welfare challenges
influence the complexity of movement: fractal analysis of behaviour in zebrafish. Fishes 4, 8.
https://doi.org/10.3390/fishes4010008.

Detrich, H.W., Detrich III, H.W., Westerfield, M., Zon, L.I., 2011. The zebrafish: genetics, genomics
and informatics. Academic Press.

Detrich, H.W., Westerfield, M., Zon, L.I., 2004. The zebrafish: Biology. ACADEMIC PRESS.

Dukas, R., 2004. Causes and consequences of limited attention. Brain, Behavior and Evolution 63,
197-210. https://doi.org/10.1159/000076781.

Eaton, R.C., Bombardieri, R.A., Meyer, D.L., 1977. The Mauthner-initiated startle response in teleost
fish. Journal of Experimental Biology 66, 65-81. https://doi.org/10.1242/jeb.66.1.65.

Eaton, R.C., Didomenico, R., Nissanov, J., 1991. Role of the Mauthner cell in sensorimotor integration
by the brain stem escape network. Brain, Behavior and Evolution 37, 272-285.
https://doi.org/10.1159/000114365.

Ebert, D., 2005. Ecology, epidemiology, and evolution of parasitism in Daphnia. National Library of
Medicine. https://edoc.unibas.ch/13505/1/Ebert-Parasitism_in_Daphnia-2005-A4-print.pdf.

Ebert, D., 2008. Host–parasite coevolution: insights from the Daphnia–parasite model system.
Current opinion in microbiology 11, 290-301. https://doi.org/10.1016/j.mib.2008.05.012.

Egan, R.J., Bergner, C.L., Hart, P.C., Cachat, J.M., Canavello, P.R., Elegante, M.F., Elkhayat, S.I.,
Bartels, B.K., Tien, A.K., Tien, D.H., 2009. Understanding behavioral and physiological phenotypes of
stress and anxiety in zebrafish. Behavioural brain research 205, 38-44.
https://doi.org/10.1016/j.bbr.2009.06.022.

Erbe, C., Dunlop, R., Dolman, S., 2018. Effects of noise on marine mammals, Effects of anthropogenic
noise on animals. Springer, pp. 277-309. https://doi.org/10.1007/978-1-4939-8574-6_10.

Fedoroff, K.J., 2021. Sound bites: the impact of boat noise on freshwater fish. University of
Saskatchewan. http://hdl.handle.net/10388/13300.

Gerhard, G.S., Kauffman, E.J., Wang, X., Stewart, R., Moore, J.L., Kasales, C.J., Demidenko, E., Cheng,
K.C., 2002. Life spans and senescent phenotypes in two strains of Zebrafish (*Danio rerio*).
Experimental gerontology 37, 1055-1068. https://doi.org/10.1016/S0531-5565(02)00088-8.

Gerlai, R., 2019. Reproducibility and replicability in zebrafish behavioral neuroscience research.
Pharmacology Biochemistry and Behavior 178, 30-38. https://doi.org/10.1016/j.pbb.2018.02.005.

Gerlai, R., Lahav, M., Guo, S., Rosenthal, A., 2000. Drinks like a fish: zebra fish (Danio rerio) as a
behavior genetic model to study alcohol effects. Pharmacology Biochemistry and Behavior 67, 773-
782. https://doi.org/10.1016/S0091-3057(00)00422-6.

Gerlai, R., Lee, V., Blaser, R., 2006. Effects of acute and chronic ethanol exposure on the behavior of
adult zebrafish (*Danio rerio*). Pharmacology Biochemistry and Behavior 85, 752-761.
https://doi.org/10.1016/j.pbb.2006.11.010.

Graham, A.L., Cooke, S.J., 2008. The effects of noise disturbance from various recreational boating
activities common to inland waters on the cardiac physiology of a freshwater fish, the largemouth
bass (*Micropterus salmoides*). Aquatic Conservation: Marine and Freshwater Ecosystems 18, 1315-
1324. https://doi.org/10.1002/aqc.941.

Green, R.E., Langston, R.H., McCluskie, A., Sutherland, R., Wilson, J.D., 2016. Lack of sound science in
assessing wind farm impacts on seabirds. Journal of Applied Ecology 53, 1635-1641. doi:
10.1111/1365-2664.12731.

Halvorsen, M.B., Casper, B.M., Matthews, F., Carlson, T.J., Popper, A.N., 2012. Effects of exposure to
pile-driving sounds on the lake sturgeon, Nile tilapia and hogchoker. Proceedings of the Royal
Society B: Biological Sciences 279, 4705-4714. https://doi.org/10.1098/rspb.2012.1544.

Hanache, P., Spataro, T., Firmat, C., Boyer, N., Fonseca, P., Médoc, V., 2020. Noise-induced reduction
in the attack rate of a planktivorous freshwater fish revealed by functional response analysis.
Freshwater Biology 65, 75-85. https://doi.org/10.1111/fwb.13271.

Handegard, N.O., Michalsen, K., Tjøstheim, D., 2003. Avoidance behaviour in cod (Gadus morhua) to
a bottom-trawling vessel. Aquatic Living Resources 16, 265-270. https://doi.org/10.1016/S0990-
7440(03)00020-2.

Hansen, K.A., Hernandez, A., Mooney, T.A., Rasmussen, M.H., Sørensen, K., Wahlberg, M., 2020. The
common murre (*Uria aalge*), an auk seabird, reacts to underwater sound. J. Acoust. Soc. Am 147,
4069-4074. https://doi.org/10.1121/10.0001400.

Harris, K.D., Bartlett, N.J., Lloyd, V.K., 2012. Daphnia as an emerging epigenetic model organism.
Genetics research international 2012. doi:10.1155/2012/147892.

Hastings, M.C., Popper, A.N., 2005. Effects of sound on fish. California Department of
Transportation. https://www.researchgate.net/profile/Arthur-
Popper/publication/253413309_Update_on_exposure_metrics_for_evaluation_of_effects_of_sound
_on_fish/links/0c9605295e4f7183f7000000/Update-on-exposure-metrics-for-evaluation-of-effects-
of-sound-on-fish.pdf.

Hawkins, A., Myrberg, A., 1983. Hearing and sound communication under water. Bioacoustics, a
comparative approach. Academic Press, London, 347-405.

Hawkins, A.D., 1986. Underwater sound and fish behaviour, The behaviour of teleost fishes.
Springer, pp. 114-151. https://doi.org/10.1007/978-1-4684-8261-4_5.

Hawkins, A.D., Hazelwood, R.A., Popper, A.N., Macey, P.C., 2021. Substrate vibrations and their
potential effects upon fishes and invertebrates. The Journal of the Acoustical Society of America 149,
2782-2790. https://doi.org/10.1121/10.0004773.

Hawkins, A.D., Picciulin, M., 2019. The importance of underwater sounds to gadoid fishes. The
Journal of the Acoustical Society of America 146, 3536-3551. https://doi.org/10.1121/1.5134683.

Hawkins, A.D., Popper, A.N., 2018. Directional hearing and sound source localization by fishes. The
Journal of the Acoustical Society of America 144, 3329-3350. https://doi.org/10.1121/1.5082306.

Hawkins, A.D., Popper, A.N., 2020. Sound detection by Atlantic cod: An overview. The Journal of the
Acoustical Society of America 148, 3027-3041. https://doi.org/10.1121/10.0002363.

Heath, V.L., Miehls, S., Johnson, N., Higgs, D.M., 2021. Behavioural response of sea lamprey
(*Petromyzon marinus*) to acoustic stimuli in a small stream. Canadian Journal of Fisheries and
Aquatic Sciences 78, 341-348. https://doi.org/10.1139/cjfas-2020-0301.

Higgs, D.M., Souza, M.J., Wilkins, H.R., Presson, J.C., Popper, A.N., 2002. Age-and size-related
changes in the inner ear and hearing ability of the adult zebrafish (*Danio rerio*). JARO: Journal of the
Association for Research in Otolaryngology 3, 174. 10.1007/s101620020035.

Hubert, J., van Bemmelen, J., Slabbekoorn, H., 2021. No negative effects of boat sound playbacks on
olfactory-mediated food finding behaviour of shore crabs in a T-maze. Environmental Pollution 270,
116184. https://doi.org/10.1016/j.envpol.2020.116184.

Injaian, A.S., Francis, C.D., Ouyang, J.Q., Dominoni, D.M., Donald, J.W., Fuxjager, M.J., Goymann, W.,
Hau, M., Husak, J.F., Johnson, M.A., 2020. Baseline and stress-induced corticosterone levels across
birds and reptiles do not reflect urbanication levels. Conservation Physiology 8, 1.
10.1093/conphys/coz110.

Johansson, K., Sigray, P., Backström, T., Magnhagen, C., 2016. Stress response and habituation to
motorboat noise in two coastal fish species in the Bothnian Sea, The effects of noise on aquatic life
II. Springer, pp. 513-521. https://doi.org/10.1007/978-1-4939-2981-8_62.

Jones, I.T., Stanley, J.A., Bonnel, J., Mooney, T.A., 2019. Complexities of tank acoustics warrant
direct, careful measurement of particle motion and pressure for bioacoustic studies, Proceedings of
Meetings on Acoustics 5ENAL. Acoustical Society of America, p. 010005.
https://doi.org/10.1121/2.0001073.

Kalueff, A.V., Cachat, J.M., 2011. Zebrafish neurobehavioral protocols.
https://link.springer.com/book/10.1007%2F978-1-60761-953-6.

Keevin, T.M., Hempen, G.L., 1997. The environmental effects of underwater explosions with
methods to mitigate impacts. Corps of engineering st louis mo st louis district.
https://apps.dtic.mil/sti/citations/ADA575523.

Kok, A.C., Bruil, L., Berges, B., Sakinan, S., Debusschere, E., Reubens, J., de Haan, D., Norro, A.,
Slabbekoorn, H., 2021. An echosounder view on the potential effects of impulsive noise pollution on
pelagic fish around windfarms in the North Sea. Environmental Pollution, 118063.
https://doi.org/10.1016/j.envpol.2021.118063.
1002
1003 Kunc, H.P., McLaughlin, K.E., Schmidt, R., 2016. Aquatic noise pollution: implications for individuals,
1004 populations, and ecosystems. Proceedings of the Royal Society B: Biological Sciences 283, 20160839.
1005 https://doi.org/10.1098/rspb.2016.0839.
1006
1007 Ladich, F., Fay, R.R., 2013. Auditory evoked potential audiometry in fish. Reviews in Fish Biology and
1008 Fisheries 23, 317-364. 10.1007/s11160-012-9297-z.
1009
1010 Lieschke, G.J., Oates, A.C., Kawakami, K., 2009. Zebrafish: methods and protocols. Springer.
1011 https://link.springer.com/book/10.1007/978-1-4939-3771-4.
1012
1013 Linke, S., Gifford, T., Desjonquères, C., Tonolla, D., Aubin, T., Barclay, L., Karaconstantis, C., Kennard,
1014 M.J., Rybak, F. and Sueur, J., 2018. Freshwater ecoacoustics as a tool for continuous ecosystem
1015 monitoring. Frontiers in Ecology and the Environment, 16(4), pp.231-238.
1016 https://doi.org/10.1002/fee.1779.
1017
1018 Malavasi, S., Georgalas, V., Lugli, M., Torricelli, P., Mainardi, D., 2004. Differences in the pattern of
1019 antipredator behaviour between hatchery-reared and wild European sea bass juveniles. Journal of
1020 fish biology 65, 143-155. https://doi.org/10.1111/j.0022-1112.2004.00545.x.
1021
1022 Maximino, C., de Brito, T.M., da Silva Batista, A.W., Herculano, A.M., Morato, S., Gouveia Jr, A., 2010.
1023 Measuring anxiety in zebrafish: a critical review. Behavioural brain research 214, 157-171.
1024 https://doi.org/10.1016/j.bbr.2010.05.031.
1025
1026 McCauley, R.D., Fewtrell, J., Popper, A.N., 2003. High intensity anthropogenic sound damages fish
1027 ears. The Journal of the Acoustical Society of America 113, 638-642.
1028 https://doi.org/10.1121/1.1527962.
1029
1030 McCloskey, K.P., Chapman, K.E., Chapuis, L., McCormick, M.I., Radford, A.N., Simpson, S.D., 2020.
1031 Assessing and mitigating impacts of motorboat noise on nesting damselfish. Environmental Pollution
1032 266, 115376. https://doi.org/10.1016/j.envpol.2020.115376.
1033
1034 McDonald, M.A., Hildebrand, J.A., Wiggins, S.M., 2006. Increases in deep ocean ambient noise in the
1035 Northeast Pacific west of San Nicolas Island, California. The Journal of the Acoustical Society of
1036 America 120, 711-718. https://doi.org/10.1121/1.2216565.
1037
1038 Mickle, M.F., Higgs, D.M., 2018. Integrating techniques: a review of the effects of anthropogenic
1039 noise on freshwater fish. Canadian Journal of Fisheries and Aquatic Sciences 75, 1534-1541.
1040 https://doi.org/10.1139/cjfas-2017-0245.
1041
1042 Mills, S.C., Beldade, R., Henry, L., Laverty, D., Nedelec, S.L., Simpson, S.D., Radford, A.N., 2020.
1043 Hormonal and behavioural effects of motorboat noise on wild coral reef fish. Environmental
1044 Pollution 262, 114250. https://doi.org/10.1016/j.envpol.2020.114250.
1045
1046 Miner, B.E., De Meester, L., Pfrender, M.E., Lampert, W., Hairston Jr, N.G., 2012. Linking genes to
1047 communities and ecosystems: Daphnia as an ecogenomic model. Proceedings of the Royal Society B:
1048 Biological Sciences 279, 1873-1882. https://doi.org/10.1098/rspb.2011.2404.

Mirjany, M., Preuss, T., Faber, D.S., 2011. Role of the lateral line mechanosensory system in directionality of goldfish auditory evoked escape response. Journal of Experimental Biology 214, 3358-3367. https://doi.org/10.1242/jeb.052894.

Moore, S.E., Reeves, R.R., Southall, B.L., Ragen, T.J., Suydam, R.S., Clark, C.W., 2012. A new framework for assessing the effects of anthropogenic sound on marine mammals in a rapidly changing Arctic. BioScience 62, 289-295. https://doi.org/10.1525/bio.2012.62.3.10.

Morley, E.L., Jones, G., Radford, A.N., 2014. The importance of invertebrates when considering the impacts of anthropogenic noise. Proceedings of the Royal Society B: Biological Sciences 281, 20132683. https://doi.org/10.1098/rspb.2013.2683.

Mortensen, L.O., Chudzinska, M.E., Slabbekoorn, H., Thomsen, F., 2021. Agent-based models to investigate sound impact on marine animals: bridging the gap between effects on individual behaviour and population level consequences. Oikos. https://doi.org/10.1111/oik.08078.

Murchy, K.A., Davies, H., Shafer, H., Cox, K., Nikolich, K., Juanes, F., 2019. Impacts of noise on the behavior and physiology of marine invertebrates: A meta-analysis. Proceedings of Meetings on Acoustics 5ENAL. Acoustical Society of America, p. 040002. https://doi.org/10.1121/2.0001217.

Nedelec, S.L., Campbell, J., Radford, A.N., Simpson, S.D., Merchant, N.D., 2016. Particle motion: the missing link in underwater acoustic ecology. Methods in Ecology and Evolution 7, 836-842.

https://doi.org/10.1111/2041-210X.12544.

Neo, Y., Hubert, J., Bolle, L., Winter, H., Slabbekoorn, H., 2018. European seabass respond more strongly to noise exposure at night and habituate over repeated trials of sound exposure. Environmental Pollution 239, 367-374. https://doi.org/10.1016/j.envpol.2018.04.018.

Neo, Y., Hubert, J., Bolle, L., Winter, H.V., Ten Cate, C., Slabbekoorn, H., 2016. Sound exposure changes European seabass behaviour in a large outdoor floating pen: Effects of temporal structure and a ramp-up procedure. Environmental Pollution 214, 26-34. https://doi.org/10.1016/j.envpol.2016.03.075.

Neo, Y.Y., Parie, L., Bakker, F., Snelderwaard, P., Tudorache, C., Schaaf, M., Slabbekoorn, H., 2015. Behavioral changes in response to sound exposure and no spatial avoidance of noisy conditions in captive zebrafish. Frontiers in behavioral neuroscience 9, 28. https://doi.org/10.3389/fnbeh.2015.00028.

Nichols, T.A., Anderson, T.W., Širović, A., 2015. Intermittent noise induces physiological stress in a coastal marine fish. PloS one 10, e0139157. https://doi.org/10.1371/journal.pone.0139157.

Normandeau Associates, I., 2012. Effects of Noise on Fish, Fisheries, and Invertebrates in the U.S. Atlantic and Arctic from Energy Industry Sound-Generating Activities., in: Interior, A.L.S.f.t.U.S.D.o.t. (Ed.), p. 135. https://www.cbd.int/doc/meetings/mar/mcbem-2014-01/other/mcbem-2014-01-submission-boem-04-en.pdf.

Parejko, K., Dodson, S.I., 1991. The evolutionary ecology of an antipredator reaction norm: *Daphnia pulex* and *Chaoborus americanus*. Evolution 45, 1665-1674. doi: 10.1111/j.1558-5646.1991.tb02671.x..

Parvulescu, A., 1967. The acoustics of small tanks. Marine Bio Acoustics 2, 7-13. https://www.scirp.org/(S(i43dyn45teexjx455qlt3d2q))/reference/ReferencesPapers.aspx?ReferenceID=55337.

Payne, N.L., van der Meulen, D.E., Suthers, I.M., Gray, C.A., Taylor, M.D., 2015. Foraging intensity of wild mulloway *Argyrosomus japonicus* decreases with increasing anthropogenic disturbance. Marine biology 162, 539-546. https://doi.org/10.1007/s00227-014-2603-7.

Peng, C., Zhao, X., Liu, G., 2015. Noise in the sea and its impacts on marine organisms. International journal of environmental research and public health 12, 12304-12323. https://doi.org/10.3390/ijerph121012304.

Pieniazek, R.H., Mickle, M.F., Higgs, D.M., 2020. Comparative analysis of noise effects on wild and captive freshwater fish behaviour. Animal Behaviour 168, 129-135. https://doi.org/10.1016/j.anbehav.2020.08.004.

Popper, A.N., Carlson, T.J., 1998. Application of sound and other stimuli to control fish behavior. Transactions of the American Fisheries Society 127, 673-707. https://doi.org/10.1577/1548-8659(1998)127<0673:AOSAOS>2.0.CO;2.

Popper, A.N., Fay, R.R., 2011. Rethinking sound detection by fishes. Hearing research 273, 25-36. https://doi.org/10.1016/j.heares.2009.12.023.

Popper, A.N., Fewtrell, J., Smith, M.E., McCauley, R.D., 2003. Anthropogenic sound: effects on the behavior and physiology of fishes. Marine Technology Society Journal 37, 35-40. https://doi.org/10.4031/002533203787537050.

Popper, A.N., Halvorsen, M.B., Kane, A., Miller, D.L., Smith, M.E., Song, J., Stein, P., Wysocki, L.E., 2007. The effects of high-intensity, low-frequency active sonar on rainbow trout. The Journal of the Acoustical Society of America 122, 623-635. https://doi.org/10.1121/1.2735115.

Popper, A.N., Hawkins, A.D., 2018. The importance of particle motion to fishes and invertebrates. The Journal of the Acoustical Society of America 143, 470-488. https://doi.org/10.1121/1.5021594.

Popper, A.N., Hawkins, A.D., 2019. An overview of fish bioacoustics and the impacts of anthropogenic sounds on fishes. Journal of fish biology 94, 692-713. https://doi.org/10.1111/jfb.13948.

Popper, A.N., Hawkins, A.D., Fay, R.R., Mann, D.A., Bartol, S., Carlson, T.J., Coombs, S., Ellison, W.T., Gentry, R.L., Halvorsen, M.B., 2014. Sound exposure guidelines, ASA S3/SC1. 4 TR-2014 Sound Exposure Guidelines for Fishes and Sea Turtles: A Technical Report prepared by ANSI-Accredited Standards Committee S3/SC1 and registered with ANSI. Springer, pp. 33-51. https://doi.org/10.1007/978-3-319-06659-2_7.

Popper, A.N., Hawkins, A.D., Sand, O., Sisneros, J.A., 2019. Examining the hearing abilities of fishes. The Journal of the Acoustical Society of America 146, 948-955. https://doi.org/10.1121/1.5120185.

Popper, A.N., Hawkins, A.D., Thomsen, F., 2020. Taking the animals' perspective regarding anthropogenic underwater sound. Trends in ecology & evolution 35, 787-794. https://doi.org/10.1016/j.tree.2020.05.002.

1146

Popper, A.N., Smith, M.E., Cott, P.A., Hanna, B.W., MacGillivray, A.O., Austin, M.E., Mann, D.A., 2005.
Effects of exposure to seismic airgun use on hearing of three fish species. The Journal of the
Acoustical Society of America 117, 3958-3971. https://doi.org/10.1121/1.1904386.

Purser, J., Radford, A.N., 2011. Acoustic noise induces attention shifts and reduces foraging
performance in three-spined sticklebacks (*Gasterosteus aculeatus*). PloS one 6, e17478.
https://doi.org/10.1371/journal.pone.0017478.

Putland, R.L. and Mensinger, A.F., 2020. Exploring the soundscape of small freshwater
lakes. Ecological Informatics, 55, p.101018. https://doi.org/10.1016/j.ecoinf.2019.101018.

Radford, A.N., Lèbre, L., Lecaillon, G., Nedelec, S.L., Simpson, S.D., 2016. Repeated exposure reduces
the response to impulsive noise in European seabass. Global change biology 22, 3349-3360.
https://doi.org/10.1111/gcb.13352.

Radford, C.A., Jeffs, A.G., Montgomery, J.C., 2008. Orientated swimming behaviour of crab
postlarvae in response to reef sound. Bioacoustics 17, 87-89.
https://doi.org/10.1080/09524622.2008.9753776.

Reynolds, C., 2011. Daphnia: Development of Model Organism in Ecology and Evolution-2011.
Freshwater Reviews 4, 85-87. https://doi.org/10.1608/FRJ-4.1.425.
1163
Roberts, L., Harding, H.R., Voellmy, I., Bruintjes, R., Simpson, S.D., Radford, A.N., Breithaupt, T.,
Elliott, M., 2016. Exposure of benthic invertebrates to sediment vibration: from laboratory
experiments to outdoor simulated pile-driving, Proceedings of Meetings on Acoustics 4ENAL.
Acoustical Society of America, p. 010029. https://doi.org/10.1121/2.0000324.

Rountree, R.A., Juanes, F. and Bolgan, M., 2020. Temperate freshwater soundscapes: A cacophony of
undescribed biological sounds now threatened by anthropogenic noise. Plos one, 15(3), p.e0221842.
https://doi.org/10.1371/journal.pone.0221842.

Sand, O., Bleckmann, H., 2008. Orientation to auditory and lateral line stimuli, Fish bioacoustics.
Springer, pp. 183-231. https://doi.org/10.1007/978-0-387-73029-5_6.
1173
1174   Santulli, A., Modica, A., Messina, C., Ceffa, L., Curatolo, A., Rivas, G., Fabi, G., D'amelio, V., 1999.
1175   Biochemical responses of European sea bass (*Dicentrarchus labrax* L.) to the stress induced by off
1176   shore experimental seismic prospecting. Marine Pollution Bulletin 38, 1105-1114.
1177   https://doi.org/10.1016/S0025-326X(99)00136-8.
1178
Sarà, G., Dean, J., d'Amato, D., Buscaino, G., Oliveri, A., Genovese, S., Ferro, S., Buffa, G., Martire,
M.L., Mazzola, S., 2007. Effect of boat noise on the behaviour of bluefin tuna Thunnus thynnus in the
Mediterranean Sea. Marine Ecology Progress Series 331, 243-253. https://www.int-
res.com/abstracts/meps/v331/p243-253/.

Scholik, A.R., Yan, H.Y., 2002. Effects of noise on auditory sensitivity of fishes.
Bioacoustics 12, 186-188. https://doi.org/10.1080/09524622.2002.9753690

Schuijf, A., 1975. Directional hearing of cod (*Gadus morhua*) under approximate free field conditions.
Journal of Comparative Physiology 98, 307-332. https://doi.org/10.1007/BF00709803.
1187

Schuijf, A., 1976. The phase model of directional hearing in fish. Sound reception in fish, 63-86.

Schuijf, A., Buwalda, R., 1975. On the mechanism of directional hearing in cod (*Gadus morhua* L.).
Journal of Comparative Physiology 98, 333-343. https://doi.org/10.1007/BF00709804.

Seda, J., Petrusek, A., 2011. Daphnia as a model organism in limnology and aquatic biology:
introductory remarks. Journal of Limnology 70, 337-344. DOI: 10.3274/JL11-70-2-17.

Shafiei Sabet, S., Karnagh, S.A., Azbari, F.Z., 2019. Experimental test of sound and light exposure on
water flea swimming behaviour, Proceedings of Meetings on Acoustics 5ENAL. Acoustical Society of
America, p. 010015. https://doi.org/10.1121/2.0001270.

Shafiei Sabet, S., Neo, Y.Y., Slabbekoorn, H., 2015. The effect of temporal variation in sound
exposure on swimming and foraging behaviour of captive zebrafish. Animal Behaviour 107, 49-60.
https://doi.org/10.1016/j.anbehav.2015.05.022.

Shafiei Sabet, S., Van Dooren, D., Slabbekoorn, H., 2016a. Son et lumière: Sound and light effects on
spatial distribution and swimming behavior in captive zebrafish. Environmental Pollution 212, 480-
488. https://doi.org/10.1016/j.envpol.2016.02.046.

Shafiei Sabet, S., Wesdorp, K., Campbell, J., Snelderwaard, P., Slabbekoorn, H., 2016b. Behavioural
responses to sound exposure in captivity by two fish species with different hearing ability. Animal
Behaviour 116, 1-11. https://doi.org/10.1016/j.anbehav.2016.03.027.

Shaw, J.R., Pfrender, M.E., Eads, B.D., Klaper, R., Callaghan, A., Sibly, R.M., Colson, I., Jansen, B.,
Gilbert, D., Colbourne, J.K., 2008. Daphnia as an emerging model for toxicological genomics.
Advances in Experimental Biology 2, 165-328. https://doi.org/10.1016/S1872-2423(08)00005-7.

Simmons, A.M., Narins, P.M., 2018. Effects of anthropogenic noise on amphibians and reptiles,
Effects of anthropogenic noise on animals. Springer, pp. 179-208. https://doi.org/10.1007/978-1-
4939-8574-6_7.

Simpson, S.D., Purser, J., Radford, A.N., 2015. Anthropogenic noise compromises antipredator
behaviour in European eels. Global change biology 21, 586-593. https://doi.org/10.1111/gcb.12685.

Sisneros, J.A., Rogers, P.H., 2016. Directional hearing and sound source localization in fishes. Fish
hearing and bioacoustics, 121-155. https://doi.org/10.1007/978-3-319-21059-9_7.

Slabbekoorn, H., Bouton, N., van Opzeeland, I., Coers, A., ten Cate, C., Popper, A.N., 2010. A noisy
spring: the impact of globally rising underwater sound levels on fish. Trends in ecology & evolution
25, 419-427. https://doi.org/10.1016/j.tree.2010.04.005.

Slabbekoorn, H., Ripmeester, E.A.P., 2008. Birdsong and anthropogenic noise: implications and
applications for conservation. Molecular ecology 17, 72-83. https://doi.org/10.1111/j.1365-
294X.2007.03487.x.

Smith, A.L., Parsons, C.H., Lanyon, R.G., Bizley, J.K., Akerman, C.J., Baker, G.E., Dempster, A.C.,
Thompson, I.D., King, A.J., 2004. An investigation of the role of auditory cortex in sound localization
using muscimol-releasing Elvax. European Journal of Neuroscience 19, 3059-3072.
https://doi.org/10.1111/j.0953-816X.2004.03379.x.

Southall, B.L., Bowles, A.E., Ellison, W.T., Finneran, J.J., Gentry, R.L., Greene Jr, C.R.,
Kastak, D., Ketten, D.R., Miller, J.H., Nachtigall, P.E., 2008. Marine mammal noise-
exposure criteria: initial scientific recommendations. Bioacoustics 17, 273-275.
https://doi.org/10.1080/09524622.2008.9753846.

Spence, R., Gerlach, G., Lawrence, C., Smith, C., 2008. The behaviour and ecology of the zebrafish,
Danio rerio. Biological Reviews 83, 13-34. https://doi.org/10.1111/j.1469-185X.2007.00030.x.
1244
1245 Spiga, I., Aldred, N., Caldwell, G.S., 2017. Anthropogenic noise compromises the anti-predator
1246 behaviour of the European seabass, *Dicentrarchus labrax* (L.). Marine Pollution Bulletin 122, 297-
1247 305. https://doi.org/10.1016/j.marpolbul.2017.06.067.
1248
1249 Staaterman, E., Gallagher, A., Holder, P., Reid, C., Altieri, A., Ogburn, M., Rummer, J., Cooke, S., 2020.
1250 Exposure to boat noise in the field yields minimal stress response in wild reef fish. Aquatic Biology
1251 29, 93-103. https://doi.org/10.3354/ab00728.
1252
1253 Stollewerk, A., 2010. The water flea Daphnia-a'new'model system for ecology and evolution? Journal
1254 of biology 9, 1-4. https://doi.org/10.1186/jbiol212.
1255
1256 Thomsen, F., Popper, A.N., Hawkins, A.D., 2021. Sound Impact Studies: A Reply to Risch et al. Trends
1257 in ecology & evolution 36, 382-384. https://doi.org/10.1016/j.tree.2021.01.012.
1258
1259 Tuomainen, U., Candolin, U., 2011. Behavioural responses to human-induced environmental change.
1260 Biological Reviews 86, 640-657. https://doi.org/10.1111/j.1469-185X.2010.00164.x.
1261
1262 Tyack, P.L., 1998. Acoustic communication under the sea. Animal acoustic communication, pp.163-
1263 220. https://doi.org/10.1007/978-3-642-76220-8_6.
1264
1265 Tyack, P.L., 2008. Implications for marine mammals of large-scale changes in the marine acoustic
1266 environment. Journal of Mammalogy 89, 549-558. https://doi.org/10.1644/07-MAMM-S-307R.1.
1267
1268 van der Knaap, I., Reubens, J., Thomas, L., Ainslie, M.A., Winter, H.V., Hubert, J., Martin, B.,
1269 Slabbekoorn, H., 2021. Effects of a seismic survey on movement of free-ranging Atlantic cod. Current
1270 Biology 31, 1555-1562. e1554. https://doi.org/10.1016/j.cub.2021.01.050.
1271
1272 Villamizar, N., Vera, L.M., Foulkes, N.S., Sánchez-Vázquez, F.J., 2014. Effect of lighting conditions on
1273 zebrafish growth and development. Zebrafish 11, 173-181. https://doi.org/10.1089/zeb.2013.0926.
1274
Voellmy, I.K., Purser, J., Flynn, D., Kennedy, P., Simpson, S.D., Radford, A.N., 2014a. Acoustic noise
reduces foraging success in two sympatric fish species via different mechanisms. Animal Behaviour
89, 191-198. https://doi.org/10.1016/j.anbehav.2013.12.029.

Voellmy, I.K., Purser, J., Simpson, S.D., Radford, A.N., 2014b. Increased noise levels have different
impacts on the anti-predator behaviour of two sympatric fish species. PloS one 9, e102946.
https://doi.org/10.1371/journal.pone.0102946.

Voellmy, I.K., Purser, J., Simpson, S.D., Radford, A.N., 2016. Effects of previous acoustic experience
on behavioral responses to experimental sound stimuli and implications for research, The effects of
noise on aquatic life II. Springer, pp. 1191-1196. https://doi.org/10.1007/978-1-4939-2981-8_149.
1284

1285 Wahlberg, M., Westerberg, H., 2005. Hearing in fish and their reactions to sounds from offshore
1286 wind farms. Marine Ecology Progress Series 288, 295-309. 10.3354/meps288295.
1287
1288 Webb, P., 1986. Effect of body form and response threshold on the vulnerability of four species of
1289 teleost prey attacked by largemouth bass (*Micropterus salmoides*). Canadian Journal of Fisheries and
1290 Aquatic Sciences 43, 763-771. https://doi.org/10.1139/f86-094.
1291
1292 Whitfield, T.T., 2002. Zebrafish as a model for hearing and deafness. Journal of neurobiology 53,
1293 157-171. https://doi.org/10.1002/neu.10123.
1294 Weber, E. H. 1820. De aure et auditu hominis et animalium. Pars I (The Ear and Hearing of Humans
1295 and Animals) (Gerhard Fleischer, Leipzig).
1296
1297 Wysocki, L.E., Davidson III, J.W., Smith, M.E., Frankel, A.S., Ellison, W.T., Mazik, P.M., Popper, A.N.,
1298 Bebak, J., 2007. Effects of aquaculture production noise on hearing, growth, and disease resistance
1299 of rainbow trout *Oncorhynchus mykiss*. Aquaculture 272, 687-697.
1300 https://doi.org/10.1016/j.aquaculture.2007.07.225.
1301
1302 Wysocki, L.E., Dittami, J.P., Ladich, F., 2006. Ship noise and cortisol secretion in European freshwater
1303 fishes. Biological conservation 128, 501-508. https://doi.org/10.1016/j.biocon.2005.10.020.
1304
1305 Zottoli, S.J., 1977. Correlation of the startle reflex and Mauthner cell auditory responses in
1306 unrestrained goldfish. Journal of Experimental Biology 66, 243-254.
1307 https://doi.org/10.1242/jeb.66.1.243.

1308

1309

1310 Figure legends

1311

1312 **Figure 1:** Waveforms and spectrograms (for 20 s stime slot) of all intermittent and continuous sound patterns
1313 used in the experiment (see also Shafiei Sabet et al., 2015). (a) Regular intermittent sound treatment with fast
1314 pulse rate (1:1), (b) Regular intermittent sound treatment with slow pulse rate (1: 4), (c) Irregular intermittent
1315 sound (1:1-7), (d) Continuous sound treatments accordingly.

1316

1317

1318 **Figure 2:** Experimental set-up. Schematic view of the test tank. A: The obscured pages of the test tank to
1319 enhance the fish's visual contrast in the film. B: Underwater speaker space separator screen with fish swimming
1320 space. C: Underwater speaker holder box. D: Underwater speaker.

1321

1322 **Figure 3:** Schematic of the time scale of exposure to the sound of each Zebrafish. Each fish underwent five
1323 sound treatments during the experiment (See also Shafiei Sabet et al., 2015): continuous sound (CN), 1:1, 1:4, 1:
1324 1-7, and ambient sound (AN) as control evidence. Each sound treatment was played for 20 minutes on an
1325 underwater speaker, and 10 minutes before the sound treatment was filmed by the camera. In this experiment,

we examined behavioural changes by comparing two time periods in two moments: 1. Up to 10 minutes before and after exposure to sound treatments to check the effect of sound and 2. Up to 10 minutes before and after the introduction of Daphnia to examine the effect Sound on foraging behaviour. Duration of sound treatment pulses is not to scale.

Figure 4: Spectral distribution of continuous sound pressure level compared to silent treatment (dB ref 1 $\mu Pa^2/Hz$). Silence conditions (blue) and continuous sound playback (red). The diagram shows that the sound intensity level has increased significantly in the range of hearing frequencies of zebrafish. For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.

Figure 5. Effect of sound exposure treatment on swimming behaviour of Zebrafish. (a) Number of startle responses expressed as the difference between the first 60 seconds during sound and the last 60 seconds before sound exposure onset: Continuous sound (CS) and three intermittent sound (1:1, 1:4, 1:1-7) and Ambient noises (AN) as control treatment (N=30, F=6.312, P=0.000435, Standard error changes ($\pm$1)). (b) Brief swimming speed difference of Zebrafish between the first 5 seconds during sound and the last 5 seconds before sound exposure on each four sound treatments and the ambient (N=30, F=11.172, P=0.000002, Standard error changes ($\pm$1)). (c) Prolonged swimming speed difference of Zebrafish between the last 60 seconds before sound and the first 60 seconds during sound exposure on five treatments (N=30, F=7.725, P=0.000054 Standard error changes ($\pm$1)).

Figure 6. Effect of sound treatments on zebrafish prolonged swimming speed. (a) Ambient condition. (b) Continuous sound. (c) regular intermittent 1:1. (d) regular intermittent 1:4. (e) irregular intermittent 1:1-7. The time was divided into three period bins for formal statistical analysis: the last 5 seconds before sound exposure, the first 5 seconds during sound exposure and the last 5 seconds during sound exposure. Prolonged swimming speed weakened with repeated exposure in all sound treatments. (NS= no significance, ***= $P \leq 0.001$, ****= $P \leq 0.0001$).

Figure 7. Average percentage of fish time spent in bottom layer of tank (N=30). (a) Brief time (15 seconds before and 15 seconds exposed to sound) (NS= no significance, *= $P \leq 0.05$, **= $P \leq 0.01$). (b) Prolonged time (60 seconds before and 60 seconds exposed to sound) (NS= no significance, *= $P \leq 0.05$, **= $P \leq 0.01$, ***= $P \leq 0.001$). Bottom layer area for spatial displacement was defined as the bottom layer with a vertical distance of 10 cm from the bottom of the tank. (df= 1) Standard error changes ($\pm$1).

Figure 8. Effect of sound exposure on horizontal spatial distribution of Zebrafish. (a) Brief time (15 seconds before and 15 seconds exposed to sound). (b) Prolonged time (60 seconds before and 60 seconds exposed to sound). The underwater speaker played back from the right tank. Bars show Means ± SE.

1362

Figure 9. Heat map of fish swimming in the tank environment. (a) Brief time (15 seconds before and 15 seconds
exposed to sound). (b) Prolonged time (60 seconds before and 60 seconds exposed to sound). The blue color
(#0000FE) indicates the 0-20 % of the fish in the tank. The aqua color (#01FFFF) indicates the 20-40 % of the
fish in the tank. The lime color (#00FF01) indicates the 40-60 % of the fish in the tank. The yellow color
(#FFFF01) indicates the 60-80 % of the fish in the tank. The red color (#FE0000) indicates the 80-100 % of the
fish in the tank. The underwater speaker played back from the right tank. For interpretation of the references to
colour in this figure legend, the reader is referred to the web version of this article.

1370

Figure 10. Effect of sound treatment on foraging behaviour of Zebrafish. (a) Food discrimination error as the
proportion of duckweed particles attacked relative to the total number of attacks to both duckweed particles and
water fleas from the introduction of food items until the end of sound exposure in sequence for each zebrafish
individual. (N=30, df=3.756, F=1.339, P=0.226) (b) Food handling error as the proportion of the total of water
fleas attacked that were missed or released again after initial grasping from onset of food introduction until the
end of sound exposure in sequence for each zebrafish individual. (N=30, df=2.825, F=26.023, P=0.000019).
Standard error changes ($\pm$1)

1378

1379

1380

1381

1382

1383   Figure 1.1

1384

1410

1411

1412

1413   Figure 1.2

1414

1415

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425

1426

1427

1428

1429

1430

1431

1432

1433

1434

1435

1436

1437

1438

1439

1440

1441

1442

1443   Figure 2

1444

1445

1446

1447

1448   Figure3

1449

1450

1451

1452

1453

1454

1455

1456

1457

1458

1459

1460

1461

1462   Figure 4

1463

1464

1465

1466   Figure 5

1467

1468

1469

1470

1471

1472

1473

1474

1475

1476

1477

1478

1479

1480   Figure 6

1481

1482

1483

1484

1485

1486   Figure 7

1487

1489

1490

1491   Figure 8

1492

1493

1494

1495   Figure 9

1496

1497

1498

1499

1500

1501

1502

1503

1504

1505

1506

1507

1508

1509

1510

1511

1512 Figure 10

1513

1514

1515

1516

1517