Experimental sound exposure modifies swimming activities and increases food handling error in adult zebrafish (*Danio rerio*)

Reza Mohsenpour[a], Saeed Shafiei Sabet[a,*]

[a] Fisheries Department, Faculty of Natural Resources, University of Guilan, Sowmeh Sara, Iran.

Abstract:

Anthropogenic sound is globally increasing and is recognized as a source of environmental pollution in terrestrial and aquatic habitats. Elevated sound levels may cause a broad range of impacts on aquatic organisms among taxa. Sound is an important sensory stimulus for aquatic organisms and it may cause fluctuations in stress-related physiological indices and in a broader extent induce behavioural effects such as driving as a distracting stimulus, masking important relevant acoustic signals and cues in a range of marine and freshwater species. However sound exposure may also induce changes in swimming activities, feed efficiency and productivity of available food sources in fish. Here, as a follow-up study, we experimentally tested sound effects on swimming activities and foraging performance in thirty adult Zebrafish (*Danio rerio*) individually in captivity. We used adult zebrafish and water flea (*Daphnia magna*) as model predator and prey, respectively. In terms of acoustic stimuli, we used four sound treatments with different temporal patterns (all in the same frequency range and moderate exposure level). As in a previous study, our results constitute strong evidence for sound-related effects on zebrafish behaviour. All sound treatments led to a significant increase in the number of startle response, brief and prolonged swimming speed for zebrafish ($P<0.05$). Zebrafish reached to the baseline swimming speed after 60 seconds in all treatments. We found partially brief and prolonged sound effects on the spatial distribution of zebrafish; Although there were not any significant sound-related behavioural changes for horizontal spatial displacement in all treatments ($P>0.05$), zebrafish swam significantly more in the lower layer of the tank except for irregular intermittent 1:1-7 in brief sound exposure ($P<0.05$). The results of foraging performance showed that food discrimination error unaffected by sound treatments and was low for the zebrafish ($P>0.05$). However, food handling error was affected by sound treatments; all treatments induced a significant rise in handling error ($P<0.001$). This study highlights the impact of sound on zebrafish swimming activities, and that more attacks are needed to consume the same number of food items under noisy conditions.

Keywords: Anthropogenic sound, Behaviour, Foraging performance, Sound impact, Zebrafish

1. Introduction

Nowadays, due to the increase in human activities and the advancement of technology globally since the Industrial Revolution, the living environment has undergone extensive changes (Normandeau Associates, 2012). These environmental changes can affect the planet and living organisms, and that can be a major threat to the biodiversity inhabit Earth (Kunc et al., 2016). The rapid growth of these changes poses many environmental challenges (Tuomainen and Candolin, 2011) in both terrestrial and aquatic habitats. Environmental pollutions (including chemical, light, and sound) are introduced by human activities in different time, scale and space ranges and have elevated underwater ambient noise levels with alternating intensities which may affect aquatic organisms in their habitats (Popper and Hastings, 2009; Slabbekoorn et al., 2010). Among these, one of the main sources of environmental pollution which may also can be recognized as an environmental stress stimulus is anthropogenic sound that in addition to affecting terrestrial animals, also have many consequences on aquatic organisms (Popper et al., 2020; Slabbekoorn et al., 2010; Slabbekoorn and Ripmeester, 2008).

Sound sources in aquatic habitats, such as merchant shipping, recreational boating, wind turbines, pile-driving, underwater mining explorations, and explosions related to geological and research experiments, are frequent in the number of events and widespread geographically. (McDonald et al., 2006; Normandeau Associates, 2012). Consequently, anthropogenic sound has changed underwater soundscapes worldwide and represents a very subtle driver of environmental change and novel challenge to aquatic organisms. Moreover, the high speed of sound underwater, which is about 5 times faster than the speed of sound in air, shows the importance and priority of examining the role and applications of acoustic stimuli and their effects on aquatic organisms.

Recent studies have investigated impacts of anthropogenic sound on a wide variety of taxa and across a range of scales (Barber et al., 2009; Morley et al., 2014; Normandeau Associates, 2012; Slabbekoorn et al., 2010; Thomsen et al., 2021; Tyack, 2008). Anthropogenic sound can cause physical, physiological, and behavioural disorders in aquatic organisms, including marine mammals (Erbe et al., 2018; Moore et al., 2012; Southall et al., 2008), seabirds (Bermúdez-Cuamatzin et al., 2018; Green et al., 2016; Hansen et al., 2020), reptiles (Injaian et al., 2020; Simmons and Narins, 2018), fish (Hastings and Popper, 2005; Hawkins, 1986; Mills et al., 2020; Popper et al., 2003), and invertebrates (Carroll et al., 2017; Coquereau et al., 2016; Murchy et al., 2019).

Because of the high propagation rate and low attenuation over large distances in underwater environments acoustic stimuli play an important role in aquatic environments (Tyack, 1998; Popper and Hastings, 2009; Slabbekoorn et al., 2010). Depending distance from the sound source, recent studies have shown dramatic effects of sound such as physical damages, sever injury or even death (Budelmann, 2011; Halvorsen et al., 2012; Keevin and Hempen, 1997). Further distance from the sound source, there may be physiological responses such as permanent and temporary hearing threshold shifts, fluctuations in physiological indices (André et al., 2011; Casper et al., 2013; McCauley et al., 2003; Popper et al., 2007; Popper et al., 2005; Scholik and Yan, 2002; Smith et al., 2004; Wysocki et al., 2007; Wysocki et al., 2006) such as elevated cortisol levels, the classical stress-related hormone (Johansson et al., 2016; Nichols et al., 2015; Santulli et al., 1999; Smith et al., 2004; Wysocki et al., 2006) and increased hear rates (Graham and Cooke, 2008; Simpson et al., 2015). Furthest distance from sound source, in a broader extent, behavioural effects are the most likely to occur and thus play as a stress driver (Smith et al., 2004; Popper and Hawkins, 2019) as a distracting stimulus (Popper and Carlson, 1998; Chan et al., 2010), interfere with detecting prey and antipredator behaviour (Hawkins and Myrberg, 1983), compromise foraging performance (Neo et al., 2015; Purser and Radford, 2011; Shafiei Sabet et al., 2015; Voellmy et al., 2016), disrupt reproductive behaviour (McCloskey et al., 2020) or mask important acoustic signals and cues for conspecific recognition and communication purposes (Codarin et al., 2009; Amorim et al., 2015; De Jong et al., 2018b; Hawkins and Picciulin, 2019) in a range of marine and freshwater species.

Many marine and freshwater fishes have well-developed hearing abilities that provide them a key biological privilege to detect sound and perceive a broad range of frequencies (Hawkins, 1986; Heath et al., 2021; Popper et al., 2019; Wahlberg and Westerberg, 2005; Wysocki et al., 2006). While there are well-documented studies regarding the effects of sound on the behaviour of marine fishes (de Jong et al., 2018a; Mortensen et al., 2021; Peng et al., 2015), and its adverse repercussions in many marine fish species at the individual and community level, yet much less is known across the current literature about these effects on the behaviour of freshwater fish and has not received sufficient interest so far (Fedoroff, 2021; Mickle and Higgs, 2018; Pieniazek et al., 2020). Moreover, sound exposure can also change spatial distribution and swimming behaviour of fish which may consequently affect ecologically on their avoiding to forage in noisy food areas and their navigations but also change biologically their swimming activities and foraging performance (de Vincenzi et al., 2021; Hanache et al., 2020; Hubert et al., 2021; Shafiei Sabet et al., 2016a). There is growing evidence that foraging

performance is likely to be modulated by anthropogenic sound that can induce shift from

successful feeding to typical behavioural stress responses in fish (Purser and Radford, 2011;

Shafiei Sabet et al., 2015; Voellmy et al., 2016. Currently, little is known about the effects of

sound exposure on swimming activity and foraging performance of fish, although there are

some well-documented studies. It has been shown that increased boating activity was

associated with a reduction in activity rates, changed vertical distribution and compromised

foraging success of free-ranging mulloway (*Argyrosomus japonicus*) (Payne et al., 2015) and

Mediterranean Damselfish (*Chromis chromis*) (Bracciali et al., 2012).

Other studies have shown that experimental sound exposure increase performance errors and

therefore displayed a negative impacts on foraging efficiency in both the three-spined stickle

backs (*Gasterosteus aculeatus*) (Purser and Radford, 2011) and the European minnow,

(*Phoxinus phoxinus*) (Voellmy et al., 2014a). More recently our previous study also have

shown a clear sound impact on zebrafish foraging performance; more food handling errors

under noisy conditions (Shafiei Sabet et al., 2015). A primary consequence of sound exposure

would appear to be shifts in the spatial displacement. The resulting disturbance might induce

modifications in allocated foraging time budget, foraging patterns and the relative abundance

of prey items and predatory species. Such changes in turn may increase foraging energy

demand and the amount of time allocated by fish to foraging which, subsequently induce a

number of major changes such as affect food searching, discriminating and handling.

In general, *Danio rerio* is known as a model fish species in behavioural studies and responding

to environmental conditions (Cachat et al., 2010; Egan et al., 2009; Whitfield, 2002). Zebrafish

is a member of the Cypriniformes order and the Cyprinidae family and acclimates well in

captivity (Detrich et al., 2011). This fish naturally lives in the tropical freshwater (Spence et

al., 2008). Morphologically, the zebrafish's body is narrow and elongated, with golden and blue

stripes that stretch along the body and tail (Detrich et al., 2004). Males and females are easily

separable, so that females having a more prominent abdomen and body than males, and males

have a spindle-shaped body (Spence et al., 2008). Zebrafish live in the temperature range of

24±2 ℃ and in the pH range of 6.8 to 7.5 (Cortemeglia and Beitinger, 2005). The adult size of

the zebra fish is approximately 4.5 to 6.5 cm (Gerhard et al., 2002; Spence et al., 2008).

Moreover, zebrafish have specialized hearing structures, the Weberian ossicles, with an

auditory system that has homologies to the mammalian auditory system (Weber, 1820;

Alexander, 1964; Whitfield, 2002). In recent years, zebrafish have been used as biological

models in genetic, physiological, toxicological, behavioural, ecological, and other studies

(Detrich et al., 2004; Gerlai, 2019; Gerlai et al., 2000; Kalueff and Cachat, 2011; Lieschke et al., 2009). The high genetic, physiological, and pharmacological similarities of this species with humans could be a reason for the use of this species as a biological model in research (Crawford et al., 2008).

Progress in the field of behavioural biology over conservation ecology and findings about the potential impacts pollutants on organisms is also to a large extent linked with the study of invertebrates. Daphnia is a small crustacean and inhabits in open and light waters, also they are an important part of the food web in freshwater habitats and inhabits many types of shallow water bodies (Ebert, 2005; Parejko and Dodson, 1991; Reynolds, 2011). This invertebrate is the first crustacean to have its genome sequenced (Stollewerk, 2010) and because of features such as easy cultivation, small size and short generation time it is a popular model organism in various biological disciplines from aquatic ecology to biomedical sciences (Seda and Petrusek, 2011). Daphnia may also be a useful species to study behavioural studies such as sound impacts on invertebrates.

In addition, it has been reported that in the larval stage, marine crustaceans respond to reef sounds (Radford et al., 2008). Also, the aquatic invertebrate larvae, which were the same size as the Daphnia, reacted to natural sounds and sounds from human activities, and their swimming activities have changed accordingly (Morley et al., 2014). Daphnia is used as a model in ecological (Stollewerk, 2010), physiological (Altshuler et al., 2011), genetic (Harris et al., 2012; Miner et al., 2012), toxicological (Shaw et al., 2008), and parasitological (Ebert, 2008) studies.

In the present study, as a follow up of our recent study (Shafiei Sabet et al., 20215), we experimentally tested the hypothesis that whether experimental sound exposure ensonified by an underwater speaker affect the general swimming activities and foraging behaviour of zebrafish as predator upon water flea as prey item under laboratory conditions. Comparing to Shafiei Sabet er al., 2015 that we used an in-ear speaker, in this study we adopted an underwater speaker to increase sound pressure levels and potential gradient of it in the arena. We also extended time of behavioural analysis to explore more long-term effects of sound on the zebrafish. Our specific goals were: firstly, to assess the effect of experimental sound exposure on zebrafish activities including swimming speed and spatial displacement. Secondly, to estimate whether the temporal pattern of sound exposure matters and affects differently

zebrafish behaviour. And thirdly, to verify our recent laboratory-based findings of sound
impacts on zebrafish swimming activity and foraging behaviour.

2. Materials and Methods

This study was performed in the ornamental fish breeding facility center at Fisheries
Department, Faculty of Natural Resources, University of Guilan, located in Sowmeh Sara city,
Guilan province, Iran (37°17′39″N, 49°19′55″E), using an aquarium with dimensions of 50 cm
length ×15 cm wide ×20 cm high with a volume of water intake of 112.5 liters in the period of
10:00 to 14:00 every day. Zebrafish (approximate age of 45 days old and of the wild-type,
short-fin variety weight (± standard deviation) of 1.23 ± 0.02 g) were obtained from an
ornamental fish breeding center located in Bazar-Jomeh in Sowmeh Sara county, Guilan
province, Iran.

Zebrafish were stored in a stock glass tank with dimensions 50 cm long ×30 cm wide ×40 cm
high for two weeks and adapted to environmental conditions to reduce possible stress and
hormonal changes due to transportation, captivity conditions and animal welfare issues (Deakin
et al., 2019). The fish were fed 0.8 mm commercial Biomar® feed until the day before the
experiment (Neo et al., 2015). Waterfleas were caught every morning, during the whole
experiment days, from the surface layer of the pool of the faculty, by a plankton net with a net
mesh size of 0.2 mm. This was done by horizontal twisting at a depth of 10 cm and a length of
30 cm and was kept overnight in a separate tank at the same temperature as the test tank to
adapt similarly mimicking to the temperature conditions. According to the previous study of
Shafiei Sabet et al. (2019) as well as the same sampling location and depth in the same time
period with present study and the identification key used in the previous Daphnia species study,
Daphnia magna was identified (Shafiei Sabet et al., 2019).

2.1. Sound treatments

In the present experiment, four sound treatments with different temporal patterns along with
control treatment were used, including the first treatment as control treatment in which the fish
were exposed to ambient noise (AN). Second treatment, Continuous sound (CS) (Fig. 1 (a)),
third treatment, regular intermittent noise (IN) with fast pulse rate (1:1), fourth treatment,
regular intermittent sound with slow pulse rate (1:4) and Fifth treatment is the irregular
intermittent sound (1:1-7). All three intermittent treatments include one second of sound, but
the difference between these sound treatments is the intervals between these sounds (silence
time), which are described in detail below. Regular intermittent sound with a fast pulse rate

(1:1) involves one second of sound and one second of silence (Fig. 1 (b)). Regular intermittent

sound with slow pulse rate (1:4) consists of one second of sound and four seconds of silence

(Fig. 1 (c)) and irregular intermittent sound (1: 1-7) includes one second of sound and interval

1, 2, 3,4 ,5, 6 or 7 seconds of silence is random (Fig. 1 (d)). Also, the sound treatments that are

broadcast for different fish on different days were identified quite randomly.

The sound treatments were generated and the sound files modified using Audacity® software

(2.3.1) at the sound frequency that can be detected and heard for zebrafish (300-1500 Hz)

(Higgs et al., 2002) as well as the bandwidth of anthropogenic sounds, including vehicles,

pump systems, and similar pile driving that overlap (Slabbekoorn et al., 2010). The designed

sound was produced by the software in the same sound frequency range of 400-2000 Hz based

on our earlier study (Shafiei Sabet et al., 202).

2.2. Experiment Tank

The experimental tank with dimensions of 50×20×15 cm with black background was prepared

to increase the contrast between Daphnia and fish in the video file. The experimental fish tank

was filled with water to a level of 15 ± 0.5 cm of filtered and aerated dechlorinated tap water

after overnighting. During the broadcast of the sound treatments, the experiment tank was

filmed by a video camera (Panasonic HC-V180 Full HD 28 mm Wide Lens Camcorder) at a

distance of about 50 cm from in front of the test tank. After production of sound treatments,

they were played back as stereo WAV files using a player connected (Sony Vaio SVF1421A4E

Laptop) to an underwater speaker (custom-build speaker in Iran, 8Ω, 30 W, 10 Hz- 10 KHz).

In this experiment, a divider plate was placed transversely in the tank and the tank length was

halved (25×20×15 cm) in order to increase the enclosure on the zebrafish swimming

environment and make the entire fish swimming space visible for video-recording (Fig. 2). In

order to reduce the stress of the fish and also to reduce the effects of people moving to the test

site, the test tank was surrounded by black plastic so that the fish's behaviour was not affected

by other factors and only the camcorder lens passed through the plastic and was the same for

all treatments and repetitions. In such a way that when disconnecting and connecting the

camcorder between sound treatments, the fish is not seen and is not affected. To isolate the

experiments from external sounds and reduce unwanted sounds, we placed and covered all

walls at the entrance to the experimental room and also inside walls using egg boxes.

The physicochemical properties of the water were the same and monitored regularly on a daily

basis. The photoperiod used in the experiment was 12 hours of light and 12 hours of darkness

(Higgs et al., 2002; Villamizar et al., 2014) and the light intensity measured in this experiment

by the light meter model (TES_1336A – TES Electrical Electronic Corp. Taiwan) averaged 62

lux. Water temperature was measured during the experiment and was maintained at 26±1 °C.

And also, the amount of dissolved oxygen (DO) in the water was measured and was maintained

at 8±1 mg / L.

The underwater speaker used in the experiment was placed horizontally on the other side of

the separator plate (See Fig. 2). In this experiment, after introducing the fish to the test tank

during the night for about 20 hours, the fish was given the opportunity to adapt to the

environment so that it could use the entire tank space for swimming and display natural

swimming behaviour (Shafiei Sabet et al., 2015) and have the normal conditions (Neo et al.,

2015; Shafiei Sabet et al., 2015). Then, test was performed with a video camera located in front

of the tank. After ten minutes, the sound treatment was played by a speaker and a sound player

for 20 minutes, However, the food item (Daphnia) and non-food item (Duckweed) added to the

experiment tank after ten minutes from sound playback. The nutrition of the predatory species

was investigated. This same process was performed for the other sound treatments with a 15-

minute interval between treatments and the fish was exposed to all five acoustic treatments and

repeated the next day for the next fish (Fig. 3). The order of broadcast of sound treatments on

a daily basis was randomly balanced. We used a similar approach previously described to

quantify behavioural indices of zebrafish and acoustic measurements of tank in this system

(Shafiei Sabet et al., 2015).

2.3. Underwater sound measurement

In order to check the intensity of the sound that can be played by the underwater speaker and

to understand whether the tested fish was able to detect and perceive the sound treatments or

not, first the continuous treatment sound file was played back using a laptop sound player (Sony

Vaio SVF1421A4E) connected to a custom-build underwater speaker which a custom-build

sound tuning amplifier was attached. The level of sound intensity under water was recorded by

a hydrophone model (Aquarian Scientific AS-1) which connected to the amplifier model (PA-

4) and a Tascam linear PCM recorder model (DR-100MKII). The recorded sound file was

evaluated in R studio software (Version 1.1.456 - © 2009-2018 RStudio, Inc.).

According to the Fig. 4, during continuous sound playback, the frequency range emitted from

the underwater speaker was completely in the zebrafish's hearing range in the range of 300-

1500 Hz (Higgs et al., 2002) and far above the ambient noise playback. The sound pressure of

continuous sound treatment during playback was average 121 dB re 1 μPa^2/Hz for 5 seconds and the ambient sound pressure was average 96 dB re 1 μPa^2/Hz for 5 seconds.

2.4. Effect of sound on Zebrafish swimming behaviour

To investigate the effect of sound on the behaviour of Zebrafish, 30 fishes (15 males and 15 females) were introduced individually in the experimental tank after biometrics. The fish were introduced to the test tank in last day (Overnighting) and the fish were given the opportunity to get used to the environment so that they could use the entire space of the tank for their swimming and have normal conditions. Also, all fish were given 30 minutes to relax. Since then, the fish has undergone five sound treatments.

Zebrafish behavioural response to five sound treatments was video recorded for a maximum of 30 minutes for each treatment (maximum 20 minutes for exposure and a maximum of 10 minutes before exposure) (Fig. 3). Swimming behaviour parameters such as startle response (which is the peak of swimming speed of fish more than 10 cm per second that occurs immediately after the sound was played for one minute (See Shafiei Sabet et al. (2015)), brief swimming speed (5 seconds before and 5 seconds during sound) and prolonged swimming speed (one minute before and one minute during sound) were evaluated for all treatments.

Also, to explore spatial distribution of fish in the tank, in the vertical/column profile, according to the dewatering height of 15 cm of the test tank, tank height during inspection and analysis behavioural data was divided into two parts: zero to 7.5 cm and 7.5 to 15 cm. To check the distance from the sound source in the horizontal profile, the length of the tank was divided into three parts: zero to 8.33 cm, 8.33 to 16.66 cm and 16.66 to 25 cm.

2.5. Effect of sound on foraging behaviour of zebrafish on water fleas

As mentioned earlier (Fig. 3), the effect of sound on the foraging behaviour of zebrafish was investigated in such a way that 5 waterfleas (about three millimeters) as a prey species (target) and 5 non-food substances as non-food item in the same size as Daphnia (about three millimeters) was mixed in 25 ml beaker and added to the fish tank in the same manner for all treatments. The waterfleas were in the same sizes that caught with plastic Pasteur pipettes to decrease damaging water, which is suitable for feeding this species of fish at puberty and can be received by the mouth of the fish (Shafiei Sabet et al., 2015). Naturally, both of food and non-food items are present in the habitat of this fish.

Foraging performance was assessed as follows: To investigate the effect of sound treatments
on the foraging power of zebrafish, the parameters of food discrimination error (Formula 1)
and food handling error (Formula 2) were measured.

$\text{Food Discrimination Error} = \dfrac{\text{number of non-food item attacks}}{\text{total number of food and non-food attacks}}$ (Formula 1)

$\text{Food Handling Error} = \dfrac{\text{number of unsuccessful attacks on water fleas}}{\text{total number of successful and unsuccessful attacks on water fleas}}$ (Formula 2)

2.6. Behavioural information processing and statistical analysis

Recorded videos of zebrafish behaviour were converted to 10 frames per second by Xilisoft
Video Converter Ultimate software to reduce the magnification of time, in order to increase
the accuracy of the fish swimming survey and also to reduce the fish speed for spatial inspection
per second. Logger Pro® video tracking software (Vernier Software, Beaverton, OR, USA,
version 3.10.1) was used to examine behavioural responses including the number of explosive
movements, swimming speed, and spatial distribution of the fish. Entering information and
data in M.S. Excel 2016 and data analysis was performed using SPSS 25 software. The
normality of the data was evaluated by Kolmogorov-Smirnov test and the homogeneity of the
data by Levene test. Then, the presence or absence of significant differences between the mean
of the data was assessed by repeated measures ANOVA analysis and using Tukey multi-range
test. A HuynheFeldt correction was performed when sphericity could not be assumed in the
repeated measures ANOVA. Bonferroni corrected post hoc tests were performed when
ANOVA test results were significant. The level of significance in this study was considered
$P<0.05$. A custom-written acoustic calibration script in R studio software (Version 1.1.456 - ©
2009-2018 RStudio, Inc.) was also used to evaluate sound pressure levels and power spectral
density that were played by the underwater speaker.

3. Ethical note

We considered the 3Rs in behavioural research, the guidelines with respect to Good Laboratory
Practice (GLP). All housing, handling and experimental conditions were in accordance with
the guidelines for the treatment of animals in behavioural research and teaching (ASAB, 2020).
Water fleas and zebrafish were allowed to acclimatize gradually to the laboratory conditions
before they were used in any of the experiments and showed no signs of adverse effects of the
experimental conditions. Zebrafish showed only a brief startle response with the onset of the
moderate sound playbacks and no mortalities or physical damages were observed during

experiments (Neo et al., 2015; Shafiei Sabet et al., 2015). There are no legal requirements for
studies involving invertebrates such as waterfleas (Daphnia) in Iran. The principal investigator
(S.SH.S) passed successfully the exam for the course on laboratory animal science at Leiden
University, the Netherlands and holds animal testing act certificate "as an Article 9 researcher".

4. Result

4.1. Impact of sound on swimming behaviour of Zebrafish

Experimental sound exposure has changed zebrafish swimming activities in different ways.
Sound treatments significantly increased the parameters of startle response and swimming
speed of fish (Fig. 5a, b, c). The number of startle response instantly increased in exposure to
sound treatments so that all sound treatments showed a significant difference compared to the
ambient condition (repeated measures ANOVA: $F_{3.23,93.68}$=6.31, P=0.000435), But no
significant differences were observed in terms of temporal patterns between sound treatments
(P>0.05) (Fig 5a). Also, in the brief swimming speed difference (5 seconds before the sound
and 5 seconds during the sound) there was a significant difference compared to the ambient
condition (repeated measures ANOVA: $F_{3.06,88.70}$=11.17, P=0.000002), although there was no
significant difference between the sound treatments with different temporal patterns in this time
period (P>0.05) (Fig 5b). This difference in swimming speed in exposure to sound treatments
was also true in prolonged (60 seconds before the sound and 60 seconds during the sound), so
that there was a significant affect between sound treatments compared to ambient conditions
(repeated measures ANOVA: $F_{3.39,98.34}$=7.72, P=0.000054), however again there was
nonsignificant difference between sound treatments (P>0.05) (Fig 5b).

According to Figure 6, in four sound treatments (CS, 1:1, 1:4, 1:1-7), a sudden increase in fish
swimming speed was observed once sound treatments were played at 60 seconds. Observations
also showed that in all sound treatments, fish reached the baseline after 60 seconds during
sound. In ambient condition (Fig 6a), no significant difference in swimming speed was
observed in any of the time periods. While in continuous sound and regular intermittent 1:1
treatment (Fig 6b, c), a significant difference was observed in the comparison of 5 seconds
before sound playback and the first 5 seconds of sound playback (CS= repeated measure
ANOVA: $F_{4.54,131.66}$=9.53, P≤0.001, 1:1= repeated measure ANOVA: $F_{3.85,111.72}$=11.72,
P≤0.001). This is while in this treatment, compared to the last 5 seconds before the sound and
the last 5 seconds of the sound, fish reached the base line and there was no difference (P>0.05).
Also, this difference was observed in the two treatments regular intermittent 1:4 and irregular

intermittent 1:1-7 (Fig 6d, e) with more intensity compared to before and during the sound

(1:4= repeated measure ANOVA: $F_{6.24,181.01}$=14.17, $P\leq0.0001$, 1:1-7= repeated measure

ANOVA: $F_{7.14,207.15}$=15.01, $P\leq0.0001$). But in comparing the baseline time in these two

treatments, no significant difference was observed and the fish reached the baseline ($P>0.05$).

Changes in the spatial distribution of Zebrafish were investigated when exposed to sound

treatments in vertical scale (lower layer) (Fig. 7a, b) and horizontal scale (X position) (Fig.

8a, b). According to Figure 7a, the average percentage of fish time spent in the lower layer of

the tank during sound treatments exposure over a brief time (15 seconds before sound and 15

seconds during sound exposure) there was no treatment effect ($F_{3.77,109.47}$=1.486, P=0.214) or

interaction for treatment*times ($F_{3.55,102.92}$=0.634, P=0.621). But there was effect of times

($F_{1,29}$=28.274, P=0.000011). In two treatments of ambient (AN) and irregular intermittent

sound (1:1-7) with before sound exposure was not significant effect (AN= $F_{1,29}$=28.274,

P=0.104, 1:1-7= $F_{1,29}$=28.274, P=0.051). But there was a significant difference between two

sound treatments: continuous sound (CS) and 1:4 during and before exposure to sound (CS=

$F_{1,29}$=28.274, P=0.007, 1:4= $F_{1,29}$=28.274, P=0.004) and also 1:1 treatment was a significant

difference with before the sound exposure ($F_{1,29}$=28.274, P=0.011). According to Figure 7b,

the average percentage of fish time spent in the lower layer of the tank during sound

treatments over a prolonged time (60 seconds before sound and 60 seconds during sound

exposure), there was no treatment effect ($F_{3.65,105.92}$=1.837, P=0.133) or interaction for

treatment*times ($F_{4,116}$=1.780, P=0.137). But there was effect of times ($F_{1,29}$=35.398,

P=0.000002). In control treatment (AN) between before and during sound exposure do not

show a significant difference ($F_{1,29}$=35.398, P=0.975). While in continuous treatment (CS)

and 1:4 between before and during sound exposure showed a significant difference (CS=

$F_{1,29}$=35.398, P=0.000411, 1:4= $F_{1,29}$=35.398, P=0.000360) and 1:1 treatment was significant

difference between during exposure and before time ($F_{1,29}$=35.398, P=0.008). Also, irregular

intermittent sound treatment (1:1-7) between during and before sound exposure was

significant difference ($F_{1,29}$=35.398, P=0.042). But in the horizontal profile (X position),

there was no significant difference between sound treatments and control conditions, both in

the brief and prolonged time ($F_{4,116}$=1.369, P=0.249) (Fig 8a) ($F_{3.78,109.71}$=1.810, P=0.136)

(Fig 8b). Also, according to Figure 9a, b, the effects of different sound patterns on the spatial

distribution of fish can be seen as a heat map in brief and prolonged duration. Acoustic

treatments affected changes in swimming pattern from the top layer to the bottom layer but

had no effect on the distance from the sound source (right side of the tank).

4.2. Impact of sound on foraging performance of Zebrafish

According to Figure 10a, none of the acoustic treatments showed a significant difference

compared to the silent treatment on zebrafish food discrimination error (repeated measure

ANOVA: $F_{4,116}=1.339$, $P=0.260$). In fact, there was no food discrimination error between

food and non-food item by broadcasting sound treatments compared to control treatment.

However, all acoustic treatments showed a significant difference compared to the control

treatment (repeated measure ANOVA: $F_{2.82,81.91}=26.023$, $P\leq0.001$) but no significant

difference was observed between sound treatments ($P>0.05$) (Fig. 10b). This means that with

the broadcast of acoustic treatments compared to the control treatment, there was a significant

handling error in food intake in the food.

5. Discussion

Our results unequivocally demonstrate that acoustic stimuli affect zebrafish behaviour in

different ways. In this experiment, zebrafish swimming behaviour indices such as the number

of startle response, the difference in brief swimming speed (within 5 seconds), the difference

in prolonged swimming speed (within 60 seconds) and the spatial distribution of zebrafish such

as the percentage of fish in the lower layer of the tank and horizontal spatial distribution in

response to continuous and intermittent sound patterns were examined. The results showed that

the number of startle response indicating anxiety in zebrafish and other aquatic species (Blaser

et al., 2010; Maximino et al., 2010), in different sound treatments, showed a significant

difference compared to control condition. However, these changes did not show a significant

difference between sound treatments. Also, in another part of the results, the difference

between brief and prolonged swimming speeds in all sound treatments compared to the control

condition was significant. However, these changes did not show a significant difference

between sound treatments. Moreover, interestingly like what we have shown in our earlier

study (Shafiei Sabet et al., 2015), we have seen the same impact of sound exposure on foraging

performance in zebrafish as all will be discussed further as follows in the next sections.

5.1. Startle responses as a specific indicator of moderate anxiety in zebrafish?

Startle response is an involuntary action that is controlled by a pair of brain neurons in the

Mauthner (M-) cells in the mesencephalon and play a major role in the decision-making process

(Eaton et al., 1977; Eaton et al., 1991; Mirjany et al., 2011; Zottoli, 1977). Increasing the

intensity of the sound triggers a response by involuntary Mauthner cells in the mesencephalon,

leading to an involuntary escape response in fish (Eaton et al., 1977; Eaton et al., 1991).

Increasing the number of startle response and brief swimming speed of fish by playing sound treatments, causes behavioural responses related to fear and anxiety in this species. Previous studies have shown that sounds increase motor acceleration and startle responses in zebrafish (Neo et al., 2015; Shafiei Sabet et al., 2015). However, Shafiei Sabet et al. (2015) using an in-air speaker as a sound source reported that the difference in the number of startle response between continuous and intermittent (1:1) treatments with the ambient treatment was not significant which is not consistent with the results of this study. The reason of this difference in the occurrence of stress-related swimming behaviour could be due to differences in the background sound intensity before the test, differences in the method of ensonifiying of the fish tank; the sound source (speaker) under water or in-air, different storage conditions of fish and also differences Genetics and individual characteristics in zebrafish.

Studies by other researchers have shown that other species of fish respond similarly to sudden swimming behaviour in response to sound exposure. European minnow (*Phoxinus phoxinus*) and sticklebacks (*Gasterosteus aculeatus*) also showed a significant increase in the number of startle response in the face of sound, which is similar to the results of this study (Purser and Radford, 2011; Voellmy et al., 2014a). Startle response at the onset of sudden sound exposure is a common behavioural feature in fish kept in captivity and in the laboratory conditions. Of course, fish in the open and natural conditions can also show behavioural responses related to fear and anxiety (Neo et al., 2016; Staaterman et al., 2020). Spiga et al. (2017) stated that European seabass (*Dicentrarchus labrax*) also showed a higher number of startle response in continuous and intermittent sound treatment against ambient (control) treatment.

Startle response by prey fish is a behavioural response to increase survival in predator-prey relationships (Webb, 1986). By hearing the sounds of predator fish and receiving sound signals related to the attack, the prey fish starts swimming at high speed and explosively in the opposite direction of the perceived sound in order to increase the success rate of escaping and staying away from the predatory species. Sounds can affect the prey fish's decision-making power against sound sources or danger, the way in which prey assesses risk (Dukas, 2004), and the loss of focus and lack of appropriate response to danger (Chan et al., 2010; Simpson et al., 2015). It has been suggested that increasing sound levels can potentially impair the perception of danger by predatory fish species (Slabbekoorn et al., 2010). Involuntary and acquired behavioural responses related to fear and anxiety are associated with the potential for the presence of danger (Blaser et al., 2010; Maximino et al., 2010). The quality and quantity of behavioural responses of fish in captivity and in vitro to brief and severe stress stimuli are

different from those of fish living in the habitat, reducing the behavioural responses in the habitat and in the wild (Malavasi et al., 2004). One of the reasons for these differences could be due to the most ability of fish to respond in the wild before reaching the stimulus threshold, so that these fish have a longer time to make decisions and escape from the danger zone by hearing and perceiving closely the sounds associated with the predator species and they have more space available than controlled laboratory environments. Another reason could be the high level of basal stress potential in controlled laboratory environments, which with additional stress due to the perception of the predator species leads to an increase and intensification of total stress in the prey species and more intense responses are shown.

In addition to behavioural responses, increasing sound levels can also affect physiological responses in the laboratory and in the natural habitat of fish. The study of Spiga et al. (2017) and Radford et al. (2016) showed that sound exposure had a significant effect on the number of opening and closing of gills and thus on the gill ventilation of European seabass (*Dicentrarchus labrax*) compared to the control treatment. This increase in oxygen demand by European bass (*Dicentrarchus labrax*), which is accompanied by an increase in gill ventilation and the opening and closing of gill operculum, indicates an increase in stress levels. Santulli et al. (1999) showed that blood biochemical parameters including cortisol and glucose in European seabass (*Dicentrarchus labrax*) increased in sound treatments compared to the ambient treatment. Staaterman et al. (2020) stated that anthropogenic sound treatments in the natural environment also have the potential to affect stress-related physiology in coral reef fish, so that the amount of cortisol in sound treatments was significantly increased compared to the control treatment.

5.2. Sound impacts on the behavioural tolerance in swimming activities

In the experiment performed on zebrafish, different sound patterns had a significant effect on the brief and prolonged swimming speed of the fish compared to the ambient treatment, which is consistent with the observations of the present study (Neo et al., 2015; Shafiei Sabet et al., 2015; Shafiei Sabet et al., 2016a). In the experiment of Neo et al. (2015) with a group of 5 zebrafish in each treatment, in comparison with increasing the swimming speed of fish with the ambient treatment, as in the results of this study, the intermittent (1:1) treatment had a higher mean speed and significant difference than other treatments. Also, with onset of sound exposure, a significant difference in swimming speed was observed in cod fish (*Gadus morhua*), which was consistent with the results of this study (Handegard et al., 2003). There

was also a significant difference in the swimming speed of European seabass (*Dicentrarchus*
labrax) compared to the silent conditions (Neo et al., 2018). The prolonged swimming speed
of the zebrafish weakened with repeated exposure suggesting a habituation process. Zebrafish
might experience negative consequences if they are disturbed and do not habituate contrary to
sound exposure. Shafiei Sabet et al. (2016b) by comparing the effect of sound on the swimming
behaviour of two species of The Lake Victoria cichlids (*Haplochromis piceatus*) and zebrafish
(*Danio rerio*) have showed that the application of sound treatments reduced the swimming
speed of cichlids (*Haplochromis piceatus*) and increased the swimming speed of zebrafish
(Shafiei Sabet et al., 2016b). The reason of the difference in the swimming speed of cichlids
(*Haplochromis piceatus*) can be related to species-specific behavioural responses in response
to acoustic stimuli, genetic characteristics and habitat conditions.

5.3. The effect of sound on the spatial distribution of zebrafish (Vertical/Horizontal)

Studies on the stress indices of zebrafish in the face of different sound patterns showed that
with the onset of sound treatments, the spatial distribution of fish changes and the fish shows
a greater tendency to swim in the lower layer of the aquarium environment. Also, the study on
the percentage of fish in the lower layer of the test tank showed that in some sound treatments
in the brief time there was a significant difference and in the prolonged time in all sound
treatments except the ambient treatment, this means Which had a greater tendency to be present
in the lower layer when playing the sound of fish. In another part of the results of this study
and the study of the spatial distribution of the presence of fish in the horizontal profile (X
position) also showed that in none of the sound treatments the fish did not tend to distance from
the sound source and did not show significant differences.

Neo et al. (2015) in the study of the effect of sound on the spatial distribution of zebrafish,
found that zebrafish with the beginning of broadcasting sound treatments showed startle
response and increased brief swimming speed and the spatial distribution of zebrafish changed
so that fish was more inclined to and they swam in the top and surface layers of the test tank.
Also, in this study (Neo et al. 2015), there were no observations of the freezing and standing
of the fish in the lower layer of the tank. However, the results of the present experiment in the
spatial distribution of zebrafish, the percentage of time fish staying in the lower layer of the
test tank was higher, which is contrary to the report of Neo et al. (2015). One of the reasons for
this difference in the results of vertical spatial distribution could be the amount of sound
intensity emitted in the treatments used in these two studies. The intensity of sound emitted in

acoustic treatments was equal to 112 dB re 1μ Pa, which is less than the intensity of sound in this study (121 dB re 1μ Pa). Therefore, a significant increase in sound intensity and high sound level difference between sound treatments and ambient treatments can lead to different responses in fish. In another study, Neo et al. (2018) designed an experiment to investigate the effect of different sound patterns on the behaviour of European seabass and found that anthropogenic sounds increase the swimming depth of European seabass and distance from the sound source, which results in behavioural responses is consistent with present study in lower layer results.

Other fishes have also shown spatial distribution changes in response to acoustic stimuli. In a field study (Kok et al., 2021) have shown that bottom-moored echosounders, representative of a high intensity impulsive intermittent anthropogenic sound, affect the abundance, schooling cohesion behaviour and swimming depth of pelagic fish. Two recent telemetry tagging studies demonstrated the effects of another intermittent source of sound, seismic surveys, on free-ranging benthic fish species. Bruce et al. (2018) showed shift of diurnal activity patterns and general swimming speed in eight tiger flatheads (*Neoplatycephalus richardsoni*). van der Knaap et al. (2021) revealed also demonstrated in a field study that Atlantic cod (*Gadus morhua)* left noisy area more quickly and decrease their swimming activity rather than ambient sound condition as control treatment. They also have shown that seismic surveys can disrupt diurnal activity cycles which may have long term negative consequences on this species foraging success and fitness.

The swimming of zebrafish towards the upper layer at the beginning of the sound transmission has been interpreted as curiosity and searching behaviour, as the authors' experimental observations have shown that by opening the door of the zebrafish test saloon and walking the staff to perform feeding fish usually produce low-pitched sounds, which attract the attention and curiosity of zebrafish and show the highest distribution at the water column level for feeding activities (Shafiei Sabet et al., 2015), However, the response of changing the spatial distribution of fish to the depth and lower layer with the beginning of sound treatments indicates the occurrence of stress and fear in fish, which is similar and expressed in studies of other researchers on other fish species (Neo et al., 2018; Sarà et al., 2007). Examination of fish behaviour to other stimuli including chemicals and fear extract has also shown that with the release of chemicals and fear extract, fish move to the lower layer (deep) column and this pattern of spatial distribution is a behavioural indicator of fear in many interpreted fish species (Gerlai et al., 2000; Gerlai et al., 2006).

In addition, another reason for the difference in the vertical distribution behavioural results observed in the study of Neo et al. (2015) and the present study could be the difference in the use of speakers, such as the use of in air speaker in the previous study (Neo et al., 2015) and the use of underwater speaker in the present study. The use of speakers in air to broadcast sound treatments leads to the production of more sound intensity in the deeper parts than in the middle and the surface of the water in the aquarium tank, which may cause the fish to move and escape towards the upper and surface layer where less sound intensity is felt (Shafiei Sabet et al., 2015).

In the study of Shafiei Sabet et al. (2015), they investigated the effect of acoustic treatments with sound intensity almost similar to this study (122 dB re 1μ Pa) on the spatial distribution of zebrafish showed that it is not consistent with the present study. The reason for this difference in spatial distribution behavioural response could be the use of in air speaker in the study by Shafiei Sabet et al. (2015) and underwater speaker in this study, as well as the complexity of sound distribution patterns and sound gradients in aquarium environments (Campbell et al., 2019) and other factors include the size of the test tank, the life cycle and location of the fish storage tank, as well as differences in the species and genetics of the fish species. Also in the study of Shafiei Sabet et al. (2015) in the spatial distribution of fish horizontally, a similar result was shown with the results of the present study and no significant difference was observed in the x position.

According to the available sources for measuring sound intensity and scattering patterns in aquarium environments, an experiment was designed by Parvulescu (1967) and Akamatsu et al. (2002) which shows the complexity and variability of sound scattering patterns and sound gradients in enclosed aquarium tank environments. This fact indicates the limitations of studying the spatial distribution of aquatic animals in enclosed and controlled environments that must be considered. Therefore, in order to study the distribution patterns of fish and other aquatic species more accurately, it is recommended to conduct field studies in natural environments of animal species in order to obtain a more accurate and complete understanding of the manner and patterns of sound-dependent distribution in aquatic species.

5.4. The importance of particle motion in fish tanks; behavioural observations for future works

In order to understand the behavioural changes of zebrafish in response to sound, first of all, it is very important to understand how the species detects and processes, and how it behaviourally responds to sound (Hawkins and Popper, 2020). Because the auditory system of fishes evolved

primarily to detect particle motion, many fishes are most sensitive to particle motion and they

can use it to determine the direction of a sound source (Hawkins and Popper, 2018; Popper and

Hawkins, 2018; Sand and Bleckmann, 2008; Sisneros and Rogers, 2016). Only some of them,

including the zebrafish, are sensitive to sound pressure as well as the particle motion (Popper

and Fay, 2011; Popper and Hawkins, 2018). There are some studies revealing directional

hearing and sound source localization in fish under laboratory conditions and in free sound

fields. Schuijf (1975) proposed that the cod determined sound direction by monitoring the

particle motion of the sound field, presumably employing the directional orientation of the

inner ear sensory cells (Dale, 1976). Although, Schuijf (1975) also concluded that the direction

of only particle motion may be insufficient to determine the direction of a sound source. It has

already been shown that cod could discriminate between signals coming towards the head as

compared to those coming towards the tail (Buwalda et al., 1983; Schuijf and Buwalda, 1975).

They argued that directional hearing might involve both comparing the responses of hear cells

oriented in different directions and also analysis of the phase relationship between the sound

pressure and particle motion to eliminate any remaining 180^0 ambiguities (Schuijf, 1976).

5.5. How important it is particle motion to fishes and invertebrates

We did not mention the levels and direction of the particle motion that is generated within the

fish tank. Therefore, we believe it is premature to conclude that zebrafish cannot localize sound

source in our experimental set up. One might be because we know very little about hearing in

fishes only over 120 species of the more than 33000 known fish species (Ladich and Fay, 2013)

and that the empirical and theoretical work on sound source localization and directional hearing

in fishes have been contradictory and obscure for decades (Sisneros and Rogers, 2016).

Moreover, some explanations would be that practically because it is difficult to monitor particle

motion in fish tank, the lack of easily used and reasonably priced instrumentation to measure

particle motion, lack of sound exposure criteria for particle motion and finally lack of particle

motion measurement standards (Popper et al., 2014).

Within an aquarium tank the levels of particle motion are often highest at the water surface,

and close to the tank walls, when an underwater loudspeaker is used (Jones et al., 2019).

Although, resonant frequencies and reverberation may influence propagation and spectro-

temporal structure of received acoustic stimuli in fish ranks (Jones et al., 2019). Our fish moved

towards the lower levels of the tank, which may be because the particle motion levels were

highest close to the water surface, and lower at the bottom of the tank. It is always important

to monitor the particle motion when examining the effects of sounds upon fishes and
invertebrates (Nedelec et al., 2016; Popper and Hawkins, 2018). Moreover, Invertebrates are
especially sensitive to substrate vibration (Aimon et al., 2021; Hawkins et al., 2021; Morley et
al., 2014; Roberts et al., 2016), and some fish are too. Particle motion measurement may play
an important role in answering crucial biological and ecological questions relating to fishes and
other species among taxa (Nedelec et al., 2016). Thus, in doing future experiments to explore
anthropogenic sound impacts on the behaviour of fishes or invertebrates under laboratory
conditions it is necessary to develop open source and accessible protocols for monitor both
particle motion on three axes and sound pressure.

5.6. Acoustic stimuli trigger foraging performance modifications negatively in zebrafish

In the present study, the parameters of fish foraging behaviour such as food discrimination
error and food handling error were examined. The results of this experiment showed that the
zebrafish did not show any significant difference in the food discrimination error when exposed
to sound treatments compared to the ambient treatment. Also, the results of food handling error
showed that all sound treatments showed a significant difference compared to the ambient
treatment, but these sound treatments did not show a significant difference compared to each
other. Here we confirmed our earlier findings (Shafiei Sabet et al., 2015) that sound impacts
may goes beyond single species. Experimental sound exposure causes more food handling
errors and foraging in zebrafish as predator, which led to more survival in waterflea as prey
and avoiding from being eaten by a predator in noisy conditions.

In a study by Purser and Radford (2011), they found that the broadcasting of sound treatments
significantly affected the foraging performance of stickleback and food discrimination error
and food handling error increased significantly compared to the ambient treatment and reduced
foraging performance, which in the food discrimination error, it is not consistent with the
results of the present experiment, and one of the reasons for this difference is the difference in
the physiology of fish and its diet. Also, the possible difference in the physiology of fish visual
sense can be one of the factors influencing these differences, but in the food handling error, a
similar result was observed with the results of the present experiment.

In other studies, by Voellmy et al. (2014b), the results showed a significant difference in the
number of unsuccessful takes of Daphnia in stickleback and no significant difference in
minnow fish, which can generally indicate the physiological difference between the two species
and possibly the difference in visual sense between the two species. The minnow fish, which

belongs to the Cyprinidae family which similar to zebrafish family, did not show a significant

difference in the unsuccessful takes of Daphnia, but in another species, a significant difference

was observed.

In a study almost similar to the present experiment, Shafiei Sabet et al. (2015) designed an

experiment to investigate the effect of different sound patterns on the swimming and foraging

behaviour of zebrafish and found that the application of sound treatments caused a significant

difference in fish handling error. There was no significant difference in food discrimination

error that was consistent with the results of this experiment. Sound playback can be effective

in identifying a substance as food and making decisions and attacking it. Since zebrafish are

among the fish that have strong visual sense and use this sense to catch and hunt, so sounds

can affect the ability to perceive potential vision.

It is also important to consider the potential impacts of acoustic stimuli in real field conditions.

Although to date, there is a gap of knowledge and very few studies have investigated

soundscapes in freshwater environments and the effects of elevated sound levels on fish in

freshwater habitats (Bolgan et al., 2016; Linke et al., 2018; Rountree et al., 2020; Putland and

Mensinger, 2020). Other studies under field conditions have also focused on the effects of

sound on fish species. In a comparative study both in a field study in the Laurentian Great

Lakes also under laboratory conditions Pieniazek et al., 2020 shown that experimental

generated white noise and boat noise have changed swimming patterns and foraging behaviour

of freshwater fish species in different ways and with highly variable patterns by family. They

demonstrated that captive black bullhead, *Ameiurus melas*, foraged less and startled more when

exposed to both white noise and boat noise treatments. Our laboratory experiments were

conducted to verify that acoustic stimuli affect swimming behaviour and foraging performance

of zebrafish and establish hypothesis for future exploration to be conducted on other freshwater

fish species and in the field.

6. Conclusion

The results of this study highlighted impacts of acoustic stimuli on a freshwater fish species

and confirmed our earlier study on the same fish species (zebrafish) under laboratory

conditions. Our findings show that the parameters of zebrafish swimming and foraging

behaviour in different sound patterns were significantly different compared to control or

ambient treatment. The results showed that sound treatments compared to the ambient

treatment caused a significant difference in the parameters of swimming behaviour including

the number of startle response, swimming speed of fish in the brief and prolonged time, spatial

distribution of fish in vertical and horizontal profiles. In general, the observed behavioural

patterns in response to sound treatments as a stressor, especially in the brief time of this study

have been observed in natural environments and in other fish species. Sound pollution as a

stressor in the brief and prolonged time can cause behavioural changes and disturbances in the

individual levels of aquatic species and have broad and important repercussions on the

communities of an ecosystem.

Also, the results obtained in the foraging behaviour showed that different sound patterns in

comparison with the ambient treatment caused a significant difference in food handling error

but no significant difference was observed in food discrimination error. Depending on the

species characteristics, these behavioural responses include escaping the predator species,

hiding in a shelter to avoid being hunted, forming crowded clusters, and even approaching the

predator species aggressively. The movement towards the lower layer observed in the

experimental tank by zebrafish can be due to the stress caused by the potential for the presence

of a predatory species, that in their natural habitats, where clear water is permeable to light, to

the deeper parts of their habitat, which have less light and have vegetation that reduces the

visibility of the predator species, and thus ensure its survival. In general, due to the

simultaneous presence of the species used in this experiment in their main habitat (zebrafish,

daphnia and duckweed) and the results obtained can be understood that sound causes overlap

in the water particles motion, which is one of the factors in the perception of the prey species

by the predator species, and the zebrafish cannot have a proper understanding of the presence

of the prey species. On the other hand, creating a state of distress and anxiety after playing

sound treatments has caused a lack of proper vision of the prey species and the zebrafish, which

is an omnivorous and visually impaired fish in prey, cannot see and hunt well.

Increased background sound levels by human activities can be recognized as a stressor and lead

to a series of changes in the activities and swimming patterns of aquatic species. It should be

noted that the results of this study are obtained in captivity and under laboratory conditions

therefore the interpretation of the results should be done with caution and attention on the

conditions of natural environments in the various habitat and behavioural limitations of any

species.

Funding:

This work was funded by the Iranian Ministry of Science, Research and Technology, University of Guilan with the grant number 971511210004.

Credit author statement

Reza Mohsenpour: Conceptualization, Methodology, Software, Formal analysis, Visualization, Data curation, writing - original draft.

Saeed Shafiei Sabet: Conceptualization, Methodology, Supervision, Project administration, Writing - review & editing.

Data accessibility:

All data used for the analyses reported in this article and some videos are available from the figshare. Moreover, the data and some videos that support the findings of this article are available from the first author upon request.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments:

Thanks to Jeroen Hubert and Ozkan Sertlek for their guide and advice on hydrophone calibrations, editing R script versions and acoustic measurements. Special tanks to Hans Slabbekoorn for providing accessibility to acoustic measurement devices. We also thank Narjes Karimi for helping us in reviewing the results and graphs.

References:

Aimon, C., Simpson, S.D., Hazelwood, R.A., Bruintjes, R., Urbina, M.A., 2021. Anthropogenic underwater vibrations are sensed and stressful for the shore crab Carcinus maenas. Environmental Pollution 285, 117148. https://doi.org/10.1016/j.envpol.2021.117148.

Akamatsu, T., Okumura, T., Novarini, N., Yan, H.Y., 2002. Empirical refinements applicable to the recording of fish sounds in small tanks. The Journal of the Acoustical Society of America 112, 3073-3082. https://doi.org/10.1121/1.1515799.

Alexander, R. 1964. The structure of the Weberian apparatus in the Siluri, in Proceedings of the Zoological Society ofLondon, Wiley Online Library, pp. 419–440. https://doi.org/10.1111/j.1469-7998.1964.tb04507.x.

Altshuler, I., Demiri, B., Xu, S., Constantin, A., Yan, N.D., Cristescu, M.E., 2011. An integrated
multi-disciplinary approach for studying multiple stressors in freshwater ecosystems: Daphnia as a
model organism. Integrative and comparative biology 51, 623-633. https://doi.org/10.1093/icb/icr103.

Amorim, M.C.P., Vasconcelos, R.O., Fonseca, P.J., 2015. Fish sounds and mate choice, Sound
communication in fishes. Springer, pp. 1-33. https://doi.org/10.1007/978-3-7091-1846-7_1
André, M., Solé, M., Lenoir, M., Durfort, M., Quero, C., Mas, A., Lombarte, A., Van Der Schaar, M.,
López-Bejar, M., Morell, M., 2011. Low-frequency sounds induce acoustic trauma in cephalopods.
Frontiers in Ecology and the Environment 9, 489-493. https://doi.org/10.1890/100124.
745
746 Animal Behaviour. 2020. Guidelines for the treatment of animals in behavioural research and teaching.
747 Animal Behaviour. 159:1–11. https://doi.org/10.1006/anbe.1996.0293.
748
749 Barber, J.R., Fristrup, K.M., Brown, C.L., Hardy, A.R., Angeloni, L.M., Crooks, K.R., 2009.
750 Conserving the wild life therein: Protecting park fauna from anthropogenic noise. 26, 26-31.
751 http://www.nature.nps.gov/ParkScience/index.cfm?ArticleID=370&Page=1
752
753 Bermúdez-Cuamatzin, E., López-Hernández, M., Campbell, J., Zuria, I., Slabbekoorn, H., 2018. The
754 role of singing style in song adjustments to fluctuating sound conditions: a comparative study on
755 Mexican birds. Behavioural processes 157, 645-655. https://doi.org/10.1016/j.beproc.2018.04.005.
756
757 Blaser, R., Chadwick, L., McGinnis, G., 2010. Behavioral measures of anxiety in zebrafish (*Danio
758 rerio*). Behavioural brain research 208, 56-62. https://doi.org/10.1016/j.bbr.2009.11.009.
759
Bolgan, M., Chorazyczewska, E., Winfield, I.J., Codarin, A., O'Brien, J. and Gammell, M., 2016. First
observations of anthropogenic underwater noise in a large multi-use lake. Journal of Limnology, 75(3),
pp.644-651. doi: 10.4081/jlimnol.2016.1405.

Bracciali, C., Campobello, D., Giacoma, C., Sara, G., 2012. Effects of nautical traffic and noise on
foraging patterns of Mediterranean damselfish (*Chromis chromis*). PloS one 7, e40582.
https://doi.org/10.1371/journal.pone.0040582.

Bruce, B., Bradford, R., Foster, S., Lee, K., Lansdell, M., Cooper, S., Przeslawski, R., 2018.
Quantifying fish behaviour and commercial catch rates in relation to a marine seismic survey. Marine
environmental research 140, 18-30. https://doi.org/10.1016/j.marenvres.2018.05.005.
769
770 Budelmann, B.-U., 2011. Morphological diversity of equilibrium receptor systems in aquatic
771 invertebrates, Sensory biology of aquatic animals. Springer, pp. 757-782. https://doi.org/10.1007/978-
772 1-4612-3714-3_30 .
773
774 Buwalda, R., Schuijf, A., Hawkins, A., 1983. Discrimination by the cod of sounds from opposing
775 directions. Journal of Comparative Physiology 150, 175-184. https://doi.org/10.1007/BF00606367.
776
777 Cachat, J., Canavello, P., Elegante, M., Bartels, B., Hart, P., Bergner, C., Egan, R., Duncan, A., Tien,
778 D., Chung, A., 2010. Modeling withdrawal syndrome in zebrafish. Behavioural brain research 208, 371-
779 376. https://doi.org/10.1016/j.bbr.2009.12.004.
780
781 Campbell, J., Shafiei Sabet, S., Slabbekoorn, H., 2019. Particle motion and sound pressure in fish tanks:
782 a behavioural exploration of acoustic sensitivity in the zebrafish. Behavioural processes 164, 38-47.
783 https://doi.org/10.1016/j.beproc.2019.04.001.
784
785 Carroll, A., Przeslawski, R., Duncan, A., Gunning, M., Bruce, B., 2017. A critical review of the
786 potential impacts of marine seismic surveys on fish & invertebrates. Marine Pollution Bulletin 114, 9-
787 24. https://doi.org/10.1016/j.marpolbul.2016.11.038.
788

Casper, B.M., Smith, M.E., Halvorsen, M.B., Sun, H., Carlson, T.J., Popper, A.N., 2013. Effects of exposure to pile driving sounds on fish inner ear tissues. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology 166, 352-360. https://doi.org/10.1016/j.cbpa.2013.07.008.

Chan, A.A.Y.-H., Giraldo-Perez, P., Smith, S., Blumstein, D.T., 2010. Anthropogenic noise affects risk assessment and attention: the distracted prey hypothesis. Biology letters 6, 458-461. https://doi.org/10.1098/rsbl.2009.1081.

Coquereau, L., Grall, J., Chauvaud, L., Gervaise, C., Clavier, J., Jolivet, A., Di Iorio, L., 2016. Sound production and associated behaviours of benthic invertebrates from a coastal habitat in the north-east Atlantic. Marine biology 163, 127. https://doi.org/10.1007/s00227-016-2902-2.

Cortemeglia, C., Beitinger, T.L., 2005. Temperature tolerances of wild-type and red transgenic zebra danios. Transactions of the American Fisheries Society 134, 1431-1437. https://doi.org/10.1577/T04-197.1.

Crawford, A.D., Esguerra, C.V., de Witte, P.A., 2008. Fishing for drugs from nature: zebrafish as a technology platform for natural product discovery. Planta medica 74, 624-632. DOI: 10.1055/s-2008-1034374.

Dale, T., 1976. The labyrinthine mechanoreceptor organs of the cod *Gadus morhua* l-(teleostei: gadidae). a scanning electron microscopical study, with special reference to the morphological polarization of the macular sensory cells. http://pascal-francis.inist.fr/vibad/index.php?action=getRecordDetail&idt=PASCAL7650248453.

de Jong, K., Amorim, M., Fonseca, P.J., Heubel, K.U., 2018a. Noise affects multimodal communication during courtship in a marine fish. Frontiers in Ecology and Evolution 6, 113. https://doi.org/10.3389/fevo.2018.00113.

De Jong, K., Amorim, M.C.P., Fonseca, P.J., Fox, C.J., Heubel, K.U., 2018b. Noise can affect acoustic communication and subsequent spawning success in fish. Environmental Pollution 237, 814-823. https://doi.org/10.1016/j.envpol.2017.11.003.

de Vincenzi, G., Micarelli, P., Viola, S., Buffa, G., Sciacca, V., Maccarrone, V., Corrias, V., Reinero, F.R., Giacoma, C., Filiciotto, F., 2021. Biological sound vs. anthropogenic noise: assessment of behavioural changes in *Scyliorhinus canicula* exposed to boats noise. Animals 11, 174. https://doi.org/10.3390/ani11010174.

Deakin, A.G., Spencer, J.W., Cossins, A.R., Young, I.S., Sneddon, L.U., 2019. Welfare challenges influence the complexity of movement: fractal analysis of behaviour in zebrafish. Fishes 4, 8. https://doi.org/10.3390/fishes4010008.

Detrich, H.W., Detrich III, H.W., Westerfield, M., Zon, L.I., 2011. The zebrafish: genetics, genomics and informatics. Academic Press.

Detrich, H.W., Westerfield, M., Zon, L.I., 2004. The zebrafish: Biology. ACADEMIC PRESS.

Dukas, R., 2004. Causes and consequences of limited attention. Brain, Behavior and Evolution 63, 197-210. https://doi.org/10.1159/000076781.

Eaton, R.C., Bombardieri, R.A., Meyer, D.L., 1977. The Mauthner-initiated startle response in teleost fish. Journal of Experimental Biology 66, 65-81. https://doi.org/10.1242/jeb.66.1.65.

Eaton, R.C., Didomenico, R., Nissanov, J., 1991. Role of the Mauthner cell in sensorimotor integration
by the brain stem escape network. Brain, Behavior and Evolution 37, 272-285.
https://doi.org/10.1159/000114365.

Ebert, D., 2005. Ecology, epidemiology, and evolution of parasitism in Daphnia. National Library of
Medicine. https://edoc.unibas.ch/13505/1/Ebert-Parasitism_in_Daphnia-2005-A4-print.pdf.

Ebert, D., 2008. Host–parasite coevolution: insights from the Daphnia–parasite model system. Current
opinion in microbiology 11, 290-301. https://doi.org/10.1016/j.mib.2008.05.012.

Egan, R.J., Bergner, C.L., Hart, P.C., Cachat, J.M., Canavello, P.R., Elegante, M.F., Elkhayat, S.I.,
Bartels, B.K., Tien, A.K., Tien, D.H., 2009. Understanding behavioral and physiological phenotypes of
stress and anxiety in zebrafish. Behavioural brain research 205, 38-44.
https://doi.org/10.1016/j.bbr.2009.06.022.

Erbe, C., Dunlop, R., Dolman, S., 2018. Effects of noise on marine mammals, Effects of anthropogenic
noise on animals. Springer, pp. 277-309. https://doi.org/10.1007/978-1-4939-8574-6_10.

Fedoroff, K.J., 2021. Sound bites: the impact of boat noise on freshwater fish. University of
Saskatchewan. http://hdl.handle.net/10388/13300.

Gerhard, G.S., Kauffman, E.J., Wang, X., Stewart, R., Moore, J.L., Kasales, C.J., Demidenko, E.,
Cheng, K.C., 2002. Life spans and senescent phenotypes in two strains of Zebrafish (*Danio rerio*).
Experimental gerontology 37, 1055-1068. https://doi.org/10.1016/S0531-5565(02)00088-8.

Gerlai, R., 2019. Reproducibility and replicability in zebrafish behavioral neuroscience research.
Pharmacology Biochemistry and Behavior 178, 30-38. https://doi.org/10.1016/j.pbb.2018.02.005.

Gerlai, R., Lahav, M., Guo, S., Rosenthal, A., 2000. Drinks like a fish: zebra fish (Danio rerio) as a
behavior genetic model to study alcohol effects. Pharmacology Biochemistry and Behavior 67, 773-
782. https://doi.org/10.1016/S0091-3057(00)00422-6.

Gerlai, R., Lee, V., Blaser, R., 2006. Effects of acute and chronic ethanol exposure on the behavior of
adult zebrafish (*Danio rerio*). Pharmacology Biochemistry and Behavior 85, 752-761.
https://doi.org/10.1016/j.pbb.2006.11.010.

Graham, A.L., Cooke, S.J., 2008. The effects of noise disturbance from various recreational boating
activities common to inland waters on the cardiac physiology of a freshwater fish, the largemouth bass
(*Micropterus salmoides*). Aquatic Conservation: Marine and Freshwater Ecosystems 18, 1315-1324.
https://doi.org/10.1002/aqc.941.

Green, R.E., Langston, R.H., McCluskie, A., Sutherland, R., Wilson, J.D., 2016. Lack of sound science
in assessing wind farm impacts on seabirds. Journal of Applied Ecology 53, 1635-1641. doi:
10.1111/1365-2664.12731.

Halvorsen, M.B., Casper, B.M., Matthews, F., Carlson, T.J., Popper, A.N., 2012. Effects of exposure
to pile-driving sounds on the lake sturgeon, Nile tilapia and hogchoker. Proceedings of the Royal
Society B: Biological Sciences 279, 4705-4714. https://doi.org/10.1098/rspb.2012.1544.

Hanache, P., Spataro, T., Firmat, C., Boyer, N., Fonseca, P., Médoc, V., 2020. Noise-induced reduction
in the attack rate of a planktivorous freshwater fish revealed by functional response analysis. Freshwater
Biology 65, 75-85. https://doi.org/10.1111/fwb.13271.

Handegard, N.O., Michalsen, K., Tjøstheim, D., 2003. Avoidance behaviour in cod (Gadus morhua) to
a bottom-trawling vessel. Aquatic Living Resources 16, 265-270. https://doi.org/10.1016/S0990-
7440(03)00020-2.

Hansen, K.A., Hernandez, A., Mooney, T.A., Rasmussen, M.H., Sørensen, K., Wahlberg, M., 2020.
The common murre (*Uria aalge*), an auk seabird, reacts to underwater sound. J. Acoust. Soc. Am 147,
4069-4074. https://doi.org/10.1121/10.0001400.

Harris, K.D., Bartlett, N.J., Lloyd, V.K., 2012. Daphnia as an emerging epigenetic model organism.
Genetics research international 2012. doi:10.1155/2012/147892.

Hastings, M.C., Popper, A.N., 2005. Effects of sound on fish. California Department of Transportation.
https://www.researchgate.net/profile/Arthur-
Popper/publication/253413309_Update_on_exposure_metrics_for_evaluation_of_effects_of_sound_o
n_fish/links/0c9605295e4f7183f7000000/Update-on-exposure-metrics-for-evaluation-of-effects-of-
sound-on-fish.pdf.

Hawkins, A., Myrberg, A., 1983. Hearing and sound communication under water. Bioacoustics, a
comparative approach. Academic Press, London, 347-405.

Hawkins, A.D., 1986. Underwater sound and fish behaviour, The behaviour of teleost fishes. Springer,
pp. 114-151. https://doi.org/10.1007/978-1-4684-8261-4_5.

Hawkins, A.D., Hazelwood, R.A., Popper, A.N., Macey, P.C., 2021. Substrate vibrations and their
potential effects upon fishes and invertebrates. The Journal of the Acoustical Society of America 149,
2782-2790. https://doi.org/10.1121/10.0004773.

Hawkins, A.D., Picciulin, M., 2019. The importance of underwater sounds to gadoid fishes. The Journal
of the Acoustical Society of America 146, 3536-3551. https://doi.org/10.1121/1.5134683.

Hawkins, A.D., Popper, A.N., 2018. Directional hearing and sound source localization by fishes. The
Journal of the Acoustical Society of America 144, 3329-3350. https://doi.org/10.1121/1.5082306.

Hawkins, A.D., Popper, A.N., 2020. Sound detection by Atlantic cod: An overview. The Journal of the
Acoustical Society of America 148, 3027-3041. https://doi.org/10.1121/10.0002363.

Heath, V.L., Miehls, S., Johnson, N., Higgs, D.M., 2021. Behavioural response of sea lamprey
(*Petromyzon marinus*) to acoustic stimuli in a small stream. Canadian Journal of Fisheries and Aquatic
Sciences 78, 341-348. https://doi.org/10.1139/cjfas-2020-0301.

Higgs, D.M., Souza, M.J., Wilkins, H.R., Presson, J.C., Popper, A.N., 2002. Age-and size-related
changes in the inner ear and hearing ability of the adult zebrafish (*Danio rerio*). JARO: Journal of the
Association for Research in Otolaryngology 3, 174. 10.1007/s101620020035.

Hubert, J., van Bemmelen, J., Slabbekoorn, H., 2021. No negative effects of boat sound playbacks on
olfactory-mediated food finding behaviour of shore crabs in a T-maze. Environmental Pollution 270,
116184. https://doi.org/10.1016/j.envpol.2020.116184.

Injaian, A.S., Francis, C.D., Ouyang, J.Q., Dominoni, D.M., Donald, J.W., Fuxjager, M.J., Goymann,
W., Hau, M., Husak, J.F., Johnson, M.A., 2020. Baseline and stress-induced corticosterone levels across
birds and reptiles do not reflect urbanization levels. Conservation Physiology 8, 1.
10.1093/conphys/coz110.

Johansson, K., Sigray, P., Backström, T., Magnhagen, C., 2016. Stress response and habituation to
motorboat noise in two coastal fish species in the Bothnian Sea, The effects of noise on aquatic life II.
Springer, pp. 513-521. https://doi.org/10.1007/978-1-4939-2981-8_62.

Jones, I.T., Stanley, J.A., Bonnel, J., Mooney, T.A., 2019. Complexities of tank acoustics warrant direct,
careful measurement of particle motion and pressure for bioacoustic studies, Proceedings of Meetings
on Acoustics 5ENAL. Acoustical Society of America, p. 010005. https://doi.org/10.1121/2.0001073.

Kalueff, A.V., Cachat, J.M., 2011. Zebrafish neurobehavioral protocols.
https://link.springer.com/book/10.1007%2F978-1-60761-953-6.

Keevin, T.M., Hempen, G.L., 1997. The environmental effects of underwater explosions with methods
to mitigate impacts. Corps of engineering st louis mo st louis district.
https://apps.dtic.mil/sti/citations/ADA575523.

Kok, A.C., Bruil, L., Berges, B., Sakinan, S., Debusschere, E., Reubens, J., de Haan, D., Norro, A.,
Slabbekoorn, H., 2021. An echosounder view on the potential effects of impulsive noise pollution on
pelagic fish around windfarms in the North Sea. Environmental Pollution, 118063.
https://doi.org/10.1016/j.envpol.2021.118063.

Kunc, H.P., McLaughlin, K.E., Schmidt, R., 2016. Aquatic noise pollution: implications for individuals,
populations, and ecosystems. Proceedings of the Royal Society B: Biological Sciences 283, 20160839.
https://doi.org/10.1098/rspb.2016.0839.

Ladich, F., Fay, R.R., 2013. Auditory evoked potential audiometry in fish. Reviews in Fish Biology and
Fisheries 23, 317-364. 10.1007/s11160-012-9297-z.

Lieschke, G.J., Oates, A.C., Kawakami, K., 2009. Zebrafish: methods and protocols. Springer.
https://link.springer.com/book/10.1007/978-1-4939-3771-4.

Linke, S., Gifford, T., Desjonquères, C., Tonolla, D., Aubin, T., Barclay, L., Karaconstantis, C.,
Kennard, M.J., Rybak, F. and Sueur, J., 2018. Freshwater ecoacoustics as a tool for continuous
ecosystem monitoring. Frontiers in Ecology and the Environment, 16(4), pp.231-238.
https://doi.org/10.1002/fee.1779.

Malavasi, S., Georgalas, V., Lugli, M., Torricelli, P., Mainardi, D., 2004. Differences in the pattern of
antipredator behaviour between hatchery-reared and wild European sea bass juveniles. Journal of fish
biology 65, 143-155. https://doi.org/10.1111/j.0022-1112.2004.00545.x.

Maximino, C., de Brito, T.M., da Silva Batista, A.W., Herculano, A.M., Morato, S., Gouveia Jr, A.,
2010. Measuring anxiety in zebrafish: a critical review. Behavioural brain research 214, 157-171.
https://doi.org/10.1016/j.bbr.2010.05.031.

McCauley, R.D., Fewtrell, J., Popper, A.N., 2003. High intensity anthropogenic sound damages fish
ears. The Journal of the Acoustical Society of America 113, 638-642.
https://doi.org/10.1121/1.1527962.

McCloskey, K.P., Chapman, K.E., Chapuis, L., McCormick, M.I., Radford, A.N., Simpson, S.D., 2020.
Assessing and mitigating impacts of motorboat noise on nesting damselfish. Environmental Pollution
266, 115376. https://doi.org/10.1016/j.envpol.2020.115376.

McDonald, M.A., Hildebrand, J.A., Wiggins, S.M., 2006. Increases in deep ocean ambient noise in the
Northeast Pacific west of San Nicolas Island, California. The Journal of the Acoustical Society of
America 120, 711-718. https://doi.org/10.1121/1.2216565.
1002

Mickle, M.F., Higgs, D.M., 2018. Integrating techniques: a review of the effects of anthropogenic noise on freshwater fish. Canadian Journal of Fisheries and Aquatic Sciences 75, 1534-1541. https://doi.org/10.1139/cjfas-2017-0245.

Mills, S.C., Beldade, R., Henry, L., Laverty, D., Nedelec, S.L., Simpson, S.D., Radford, A.N., 2020. Hormonal and behavioural effects of motorboat noise on wild coral reef fish. Environmental Pollution 262, 114250. https://doi.org/10.1016/j.envpol.2020.114250.

Miner, B.E., De Meester, L., Pfrender, M.E., Lampert, W., Hairston Jr, N.G., 2012. Linking genes to communities and ecosystems: Daphnia as an ecogenomic model. Proceedings of the Royal Society B: Biological Sciences 279, 1873-1882. https://doi.org/10.1098/rspb.2011.2404.

Mirjany, M., Preuss, T., Faber, D.S., 2011. Role of the lateral line mechanosensory system in directionality of goldfish auditory evoked escape response. Journal of Experimental Biology 214, 3358-3367. https://doi.org/10.1242/jeb.052894.

Moore, S.E., Reeves, R.R., Southall, B.L., Ragen, T.J., Suydam, R.S., Clark, C.W., 2012. A new framework for assessing the effects of anthropogenic sound on marine mammals in a rapidly changing Arctic. BioScience 62, 289-295. https://doi.org/10.1525/bio.2012.62.3.10.

Morley, E.L., Jones, G., Radford, A.N., 2014. The importance of invertebrates when considering the impacts of anthropogenic noise. Proceedings of the Royal Society B: Biological Sciences 281, 20132683. https://doi.org/10.1098/rspb.2013.2683.

Mortensen, L.O., Chudzinska, M.E., Slabbekoorn, H., Thomsen, F., 2021. Agent-based models to investigate sound impact on marine animals: bridging the gap between effects on individual behaviour and population level consequences. Oikos. https://doi.org/10.1111/oik.08078.

Murchy, K.A., Davies, H., Shafer, H., Cox, K., Nikolich, K., Juanes, F., 2019. Impacts of noise on the behavior and physiology of marine invertebrates: A meta-analysis. Proceedings of Meetings on Acoustics 5ENAL. Acoustical Society of America, p. 040002. https://doi.org/10.1121/2.0001217.

Nedelec, S.L., Campbell, J., Radford, A.N., Simpson, S.D., Merchant, N.D., 2016. Particle motion: the missing link in underwater acoustic ecology. Methods in Ecology and Evolution 7, 836-842.

https://doi.org/10.1111/2041-210X.12544.

Neo, Y., Hubert, J., Bolle, L., Winter, H., Slabbekoorn, H., 2018. European seabass respond more strongly to noise exposure at night and habituate over repeated trials of sound exposure. Environmental Pollution 239, 367-374. https://doi.org/10.1016/j.envpol.2018.04.018.

Neo, Y., Hubert, J., Bolle, L., Winter, H.V., Ten Cate, C., Slabbekoorn, H., 2016. Sound exposure changes European seabass behaviour in a large outdoor floating pen: Effects of temporal structure and a ramp-up procedure. Environmental Pollution 214, 26-34. https://doi.org/10.1016/j.envpol.2016.03.075.

Neo, Y.Y., Parie, L., Bakker, F., Snelderwaard, P., Tudorache, C., Schaaf, M., Slabbekoorn, H., 2015. Behavioral changes in response to sound exposure and no spatial avoidance of noisy conditions in captive zebrafish. Frontiers in behavioral neuroscience 9, 28. https://doi.org/10.3389/fnbeh.2015.00028.

Nichols, T.A., Anderson, T.W., Širović, A., 2015. Intermittent noise induces physiological stress in a coastal marine fish. PloS one 10, e0139157. https://doi.org/10.1371/journal.pone.0139157.

Normandeau Associates, I., 2012. Effects of Noise on Fish, Fisheries, and Invertebrates in the U.S. Atlantic and Arctic from Energy Industry Sound-Generating Activities., in: Interior, A.L.S.f.t.U.S.D.o.t. (Ed.), p. 135. https://www.cbd.int/doc/meetings/mar/mcbem-2014-01/other/mcbem-2014-01-submission-boem-04-en.pdf.

Parejko, K., Dodson, S.I., 1991. The evolutionary ecology of an antipredator reaction norm: *Daphnia pulex* and *Chaoborus americanus*. Evolution 45, 1665-1674. doi: 10.1111/j.1558-5646.1991.tb02671.x..

Parvulescu, A., 1967. The acoustics of small tanks. Marine Bio Acoustics 2, 7-13. https://www.scirp.org/(S(i43dyn45teexjx455qlt3d2q))/reference/ReferencesPapers.aspx?ReferenceID=55337.

Payne, N.L., van der Meulen, D.E., Suthers, I.M., Gray, C.A., Taylor, M.D., 2015. Foraging intensity of wild mulloway *Argyrosomus japonicus* decreases with increasing anthropogenic disturbance. Marine biology 162, 539-546. https://doi.org/10.1007/s00227-014-2603-7.

Peng, C., Zhao, X., Liu, G., 2015. Noise in the sea and its impacts on marine organisms. International journal of environmental research and public health 12, 12304-12323. https://doi.org/10.3390/ijerph121012304.

Pieniazek, R.H., Mickle, M.F., Higgs, D.M., 2020. Comparative analysis of noise effects on wild and captive freshwater fish behaviour. Animal Behaviour 168, 129-135. https://doi.org/10.1016/j.anbehav.2020.08.004.

Popper, A.N., Carlson, T.J., 1998. Application of sound and other stimuli to control fish behavior. Transactions of the American Fisheries Society 127, 673-707. https://doi.org/10.1577/1548-8659(1998)127<0673:AOSAOS>2.0.CO;2.

Popper, A.N., Fay, R.R., 2011. Rethinking sound detection by fishes. Hearing research 273, 25-36. https://doi.org/10.1016/j.heares.2009.12.023.

Popper, A.N., Fewtrell, J., Smith, M.E., McCauley, R.D., 2003. Anthropogenic sound: effects on the behavior and physiology of fishes. Marine Technology Society Journal 37, 35-40. https://doi.org/10.4031/002533203787537050.

Popper, A.N., Halvorsen, M.B., Kane, A., Miller, D.L., Smith, M.E., Song, J., Stein, P., Wysocki, L.E., 2007. The effects of high-intensity, low-frequency active sonar on rainbow trout. The Journal of the Acoustical Society of America 122, 623-635. https://doi.org/10.1121/1.2735115.

Popper, A.N., Hawkins, A.D., 2018. The importance of particle motion to fishes and invertebrates. The Journal of the Acoustical Society of America 143, 470-488. https://doi.org/10.1121/1.5021594.

Popper, A.N., Hawkins, A.D., 2019. An overview of fish bioacoustics and the impacts of anthropogenic sounds on fishes. Journal of fish biology 94, 692-713. https://doi.org/10.1111/jfb.13948.

Popper, A.N., Hawkins, A.D., Fay, R.R., Mann, D.A., Bartol, S., Carlson, T.J., Coombs, S., Ellison, W.T., Gentry, R.L., Halvorsen, M.B., 2014. Sound exposure guidelines, ASA S3/SC1. 4 TR-2014 Sound Exposure Guidelines for Fishes and Sea Turtles: A Technical Report prepared by ANSI-Accredited Standards Committee S3/SC1 and registered with ANSI. Springer, pp. 33-51. https://doi.org/10.1007/978-3-319-06659-2_7.

Popper, A.N., Hawkins, A.D., Sand, O., Sisneros, J.A., 2019. Examining the hearing abilities of fishes. The Journal of the Acoustical Society of America 146, 948-955. https://doi.org/10.1121/1.5120185.

1106

1107　Popper, A.N., Hawkins, A.D., Thomsen, F., 2020. Taking the animals' perspective regarding
1108　anthropogenic underwater sound. Trends in ecology & evolution 35, 787-794.
1109　https://doi.org/10.1016/j.tree.2020.05.002.

1110

Popper, A.N., Smith, M.E., Cott, P.A., Hanna, B.W., MacGillivray, A.O., Austin, M.E., Mann, D.A.,
2005. Effects of exposure to seismic airgun use on hearing of three fish species. The Journal of the
Acoustical Society of America 117, 3958-3971. https://doi.org/10.1121/1.1904386.

Purser, J., Radford, A.N., 2011. Acoustic noise induces attention shifts and reduces foraging
performance in three-spined sticklebacks (*Gasterosteus aculeatus*). PloS one 6, e17478.
https://doi.org/10.1371/journal.pone.0017478.

Putland, R.L. and Mensinger, A.F., 2020. Exploring the soundscape of small freshwater
lakes. Ecological Informatics, 55, p.101018. https://doi.org/10.1016/j.ecoinf.2019.101018.

Radford, A.N., Lèbre, L., Lecaillon, G., Nedelec, S.L., Simpson, S.D., 2016. Repeated exposure reduces
the response to impulsive noise in European seabass. Global change biology 22, 3349-3360.
https://doi.org/10.1111/gcb.13352.

Radford, C.A., Jeffs, A.G., Montgomery, J.C., 2008. Orientated swimming behaviour of crab
postlarvae in response to reef sound. Bioacoustics 17, 87-89.
https://doi.org/10.1080/09524622.2008.9753776.

Reynolds, C., 2011. Daphnia: Development of Model Organism in Ecology and Evolution-2011.
Freshwater Reviews 4, 85-87. https://doi.org/10.1608/FRJ-4.1.425.

1127

Roberts, L., Harding, H.R., Voellmy, I., Bruintjes, R., Simpson, S.D., Radford, A.N., Breithaupt, T.,
Elliott, M., 2016. Exposure of benthic invertebrates to sediment vibration: from laboratory experiments
to outdoor simulated pile-driving, Proceedings of Meetings on Acoustics 4ENAL. Acoustical Society
of America, p. 010029. https://doi.org/10.1121/2.0000324.

Rountree, R.A., Juanes, F. and Bolgan, M., 2020. Temperate freshwater soundscapes: A cacophony of
undescribed biological sounds now threatened by anthropogenic noise. Plos one, 15(3), p.e0221842.
https://doi.org/10.1371/journal.pone.0221842.

Sand, O., Bleckmann, H., 2008. Orientation to auditory and lateral line stimuli, Fish bioacoustics.
Springer, pp. 183-231. https://doi.org/10.1007/978-0-387-73029-5_6.

1137

1138　Santulli, A., Modica, A., Messina, C., Ceffa, L., Curatolo, A., Rivas, G., Fabi, G., D'amelio, V., 1999.
1139　Biochemical responses of European sea bass (*Dicentrarchus labrax* L.) to the stress induced by off
1140　shore experimental seismic prospecting. Marine Pollution Bulletin 38, 1105-1114.
1141　https://doi.org/10.1016/S0025-326X(99)00136-8.

1142

Sarà, G., Dean, J., d'Amato, D., Buscaino, G., Oliveri, A., Genovese, S., Ferro, S., Buffa, G., Martire,
M.L., Mazzola, S., 2007. Effect of boat noise on the behaviour of bluefin tuna Thunnus thynnus in the
Mediterranean Sea. Marine Ecology Progress Series 331, 243-253. https://www.int-
res.com/abstracts/meps/v331/p243-253/.

Scholik, A.R., Yan, H.Y., 2002. Effects of noise on auditory sensitivity of fishes.
Bioacoustics 12, 186-188. https://doi.org/10.1080/09524622.2002.9753690

Schuijf, A., 1975. Directional hearing of cod (*Gadus morhua*) under approximate free field conditions.
Journal of Comparative Physiology 98, 307-332. https://doi.org/10.1007/BF00709803.

Schuijf, A., 1976. The phase model of directional hearing in fish. Sound reception in fish, 63-86.

Schuijf, A., Buwalda, R., 1975. On the mechanism of directional hearing in cod (*Gadus morhua* L.). Journal of Comparative Physiology 98, 333-343. https://doi.org/10.1007/BF00709804.

Seda, J., Petrusek, A., 2011. Daphnia as a model organism in limnology and aquatic biology: introductory remarks. Journal of Limnology 70, 337-344. DOI: 10.3274/JL11-70-2-17.

Shafiei Sabet, S., Karnagh, S.A., Azbari, F.Z., 2019. Experimental test of sound and light exposure on water flea swimming behaviour, Proceedings of Meetings on Acoustics 5ENAL. Acoustical Society of America, p. 010015. https://doi.org/10.1121/2.0001270.

Shafiei Sabet, S., Neo, Y.Y., Slabbekoorn, H., 2015. The effect of temporal variation in sound exposure on swimming and foraging behaviour of captive zebrafish. Animal Behaviour 107, 49-60. https://doi.org/10.1016/j.anbehav.2015.05.022.

Shafiei Sabet, S., Van Dooren, D., Slabbekoorn, H., 2016a. Son et lumière: Sound and light effects on spatial distribution and swimming behavior in captive zebrafish. Environmental Pollution 212, 480-488. https://doi.org/10.1016/j.envpol.2016.02.046.

Shafiei Sabet, S., Wesdorp, K., Campbell, J., Snelderwaard, P., Slabbekoorn, H., 2016b. Behavioural responses to sound exposure in captivity by two fish species with different hearing ability. Animal Behaviour 116, 1-11. https://doi.org/10.1016/j.anbehav.2016.03.027.

Shaw, J.R., Pfrender, M.E., Eads, B.D., Klaper, R., Callaghan, A., Sibly, R.M., Colson, I., Jansen, B., Gilbert, D., Colbourne, J.K., 2008. Daphnia as an emerging model for toxicological genomics. Advances in Experimental Biology 2, 165-328. https://doi.org/10.1016/S1872-2423(08)00005-7.

Simmons, A.M., Narins, P.M., 2018. Effects of anthropogenic noise on amphibians and reptiles, Effects of anthropogenic noise on animals. Springer, pp. 179-208. https://doi.org/10.1007/978-1-4939-8574-6_7.

Simpson, S.D., Purser, J., Radford, A.N., 2015. Anthropogenic noise compromises antipredator behaviour in European eels. Global change biology 21, 586-593. https://doi.org/10.1111/gcb.12685.

Sisneros, J.A., Rogers, P.H., 2016. Directional hearing and sound source localization in fishes. Fish hearing and bioacoustics, 121-155. https://doi.org/10.1007/978-3-319-21059-9_7.

Slabbekoorn, H., Bouton, N., van Opzeeland, I., Coers, A., ten Cate, C., Popper, A.N., 2010. A noisy spring: the impact of globally rising underwater sound levels on fish. Trends in ecology & evolution 25, 419-427. https://doi.org/10.1016/j.tree.2010.04.005.

Slabbekoorn, H., Ripmeester, E.A.P., 2008. Birdsong and anthropogenic noise: implications and applications for conservation. Molecular ecology 17, 72-83. https://doi.org/10.1111/j.1365-294X.2007.03487.x.

Smith, A.L., Parsons, C.H., Lanyon, R.G., Bizley, J.K., Akerman, C.J., Baker, G.E., Dempster, A.C., Thompson, I.D., King, A.J., 2004. An investigation of the role of auditory cortex in sound localization using muscimol-releasing Elvax. European Journal of Neuroscience 19, 3059-3072. https://doi.org/10.1111/j.0953-816X.2004.03379.x.

Southall, B.L., Bowles, A.E., Ellison, W.T., Finneran, J.J., Gentry, R.L., Greene Jr, C.R., Kastak, D., Ketten, D.R., Miller, J.H., Nachtigall, P.E., 2008. Marine mammal noise-exposure criteria: initial

scientific recommendations. Bioacoustics 17, 273-275.
https://doi.org/10.1080/09524622.2008.9753846.

Spence, R., Gerlach, G., Lawrence, C., Smith, C., 2008. The behaviour and ecology of the zebrafish, *Danio rerio*. Biological Reviews 83, 13-34. https://doi.org/10.1111/j.1469-185X.2007.00030.x.

Spiga, I., Aldred, N., Caldwell, G.S., 2017. Anthropogenic noise compromises the anti-predator behaviour of the European seabass, *Dicentrarchus labrax* (L.). Marine Pollution Bulletin 122, 297-305. https://doi.org/10.1016/j.marpolbul.2017.06.067.

Staaterman, E., Gallagher, A., Holder, P., Reid, C., Altieri, A., Ogburn, M., Rummer, J., Cooke, S., 2020. Exposure to boat noise in the field yields minimal stress response in wild reef fish. Aquatic Biology 29, 93-103. https://doi.org/10.3354/ab00728.

Stollewerk, A., 2010. The water flea Daphnia-a'new'model system for ecology and evolution? Journal of biology 9, 1-4. https://doi.org/10.1186/jbiol212.

Thomsen, F., Popper, A.N., Hawkins, A.D., 2021. Sound Impact Studies: A Reply to Risch et al. Trends in ecology & evolution 36, 382-384. https://doi.org/10.1016/j.tree.2021.01.012.

Tuomainen, U., Candolin, U., 2011. Behavioural responses to human-induced environmental change. Biological Reviews 86, 640-657. https://doi.org/10.1111/j.1469-185X.2010.00164.x.

Tyack, P.L., 1998. Acoustic communication under the sea. Animal acoustic communication, pp.163-220. https://doi.org/10.1007/978-3-642-76220-8_6.

Tyack, P.L., 2008. Implications for marine mammals of large-scale changes in the marine acoustic environment. Journal of Mammalogy 89, 549-558. https://doi.org/10.1644/07-MAMM-S-307R.1.

van der Knaap, I., Reubens, J., Thomas, L., Ainslie, M.A., Winter, H.V., Hubert, J., Martin, B., Slabbekoorn, H., 2021. Effects of a seismic survey on movement of free-ranging Atlantic cod. Current Biology 31, 1555-1562. e1554. https://doi.org/10.1016/j.cub.2021.01.050.

Villamizar, N., Vera, L.M., Foulkes, N.S., Sánchez-Vázquez, F.J., 2014. Effect of lighting conditions on zebrafish growth and development. Zebrafish 11, 173-181. https://doi.org/10.1089/zeb.2013.0926.

Voellmy, I.K., Purser, J., Flynn, D., Kennedy, P., Simpson, S.D., Radford, A.N., 2014a. Acoustic noise reduces foraging success in two sympatric fish species via different mechanisms. Animal Behaviour 89, 191-198. https://doi.org/10.1016/j.anbehav.2013.12.029.

Voellmy, I.K., Purser, J., Simpson, S.D., Radford, A.N., 2014b. Increased noise levels have different impacts on the anti-predator behaviour of two sympatric fish species. PloS one 9, e102946. https://doi.org/10.1371/journal.pone.0102946.

Voellmy, I.K., Purser, J., Simpson, S.D., Radford, A.N., 2016. Effects of previous acoustic experience on behavioral responses to experimental sound stimuli and implications for research, The effects of noise on aquatic life II. Springer, pp. 1191-1196. https://doi.org/10.1007/978-1-4939-2981-8_149.

Wahlberg, M., Westerberg, H., 2005. Hearing in fish and their reactions to sounds from offshore wind farms. Marine Ecology Progress Series 288, 295-309. 10.3354/meps288295.

Webb, P., 1986. Effect of body form and response threshold on the vulnerability of four species of teleost prey attacked by largemouth bass (*Micropterus salmoides*). Canadian Journal of Fisheries and Aquatic Sciences 43, 763-771. https://doi.org/10.1139/f86-094.

Whitfield, T.T., 2002. Zebrafish as a model for hearing and deafness. Journal of neurobiology 53, 157-171. https://doi.org/10.1002/neu.10123.

Weber, E. H. 1820. De aure et auditu hominis et animalium. Pars I (The Ear and Hearing ofHumans and Animals) (Gerhard Fleischer, Leipzig).

Wysocki, L.E., Davidson III, J.W., Smith, M.E., Frankel, A.S., Ellison, W.T., Mazik, P.M., Popper, A.N., Bebak, J., 2007. Effects of aquaculture production noise on hearing, growth, and disease resistance of rainbow trout *Oncorhynchus mykiss*. Aquaculture 272, 687-697. https://doi.org/10.1016/j.aquaculture.2007.07.225.

Wysocki, L.E., Dittami, J.P., Ladich, F., 2006. Ship noise and cortisol secretion in European freshwater fishes. Biological conservation 128, 501-508. https://doi.org/10.1016/j.biocon.2005.10.020.

Zottoli, S.J., 1977. Correlation of the startle reflex and Mauthner cell auditory responses in unrestrained goldfish. Journal of Experimental Biology 66, 243-254. https://doi.org/10.1242/jeb.66.1.243.

1288

1289 Figure legends

1290

1291 **Figure 1:** Waveforms and spectrograms (for 20 s stime slot) of all intermittent and continuous sound patterns
1292 used in the experiment (see also Shafiei Sabet et al., 2015). (a) Regular intermittent sound treatment with fast
1293 pulse rate (1:1), (b) Regular intermittent sound treatment with slow pulse rate (1: 4), (c) Irregular intermittent
1294 sound (1:1-7), (d) Continuous sound treatments accordingly.

1295

1296

1297 **Figure 2:** Experimental set-up. Schematic view of the test tank. A: The obscured pages of the test tank to enhance
1298 the fish's visual contrast in the film. B: Underwater speaker space separator screen with fish swimming space. C:
1299 Underwater speaker holder box. D: Underwater speaker.

1300

Figure 3: Schematic of the time scale of exposure to the sound of each Zebrafish. Each fish underwent five sound
treatments during the experiment (See also Shafiei Sabet et al., 2015): continuous sound (CN), 1:1, 1:4, 1: 1-7,
and ambient sound (AN) as control evidence. Each sound treatment was played for 20 minutes on an underwater
speaker, and 10 minutes before the sound treatment was filmed by the camera. In this experiment, we examined
behavioural changes by comparing two time periods in two moments: 1. Up to 10 minutes before and after
exposure to sound treatments to check the effect of sound and 2. Up to 10 minutes before and after the introduction
of Daphnia to examine the effect Sound on foraging behaviour. Duration of sound treatment pulses is not to scale.

1308

1309 **Figure 4:** Spectral distribution of continuous sound pressure level compared to silent treatment (dB ref 1
1310 μPa^2/Hz). Silence conditions (blue) and continuous sound playback (red). The diagram shows that the sound
1311 intensity level has increased significantly in the range of hearing frequencies of zebrafish. For interpretation of
1312 the references to colour in this figure legend, the reader is referred to the web version of this article.

1313

Figure 5. Effect of sound exposure treatment on swimming behaviour of Zebrafish. (a) Number of
startle responses expressed as the difference between the first 60 seconds during sound and the last 60 seconds
before sound exposure onset: Continuous sound (CS) and three intermittent sound (1:1, 1:4, 1:1-7) and Ambient
noises (AN) as control treatment (N=30, F=6.312, P=0.000435, Standard error changes ($\pm$1)). (b) Brief swimming
speed difference of Zebrafish between the first 5 seconds during sound and the last 5 seconds before sound
exposure on each four sound treatments and the ambient (N=30, F=11.172, P=0.000002, Standard error changes
($\pm$1)). (c) Prolonged swimming speed difference of Zebrafish between the last 60 seconds before sound and the
first 60 seconds during sound exposure on five treatments (N=30, F=7.725, P=0.000054 Standard error changes
($\pm$1)).

1323

Figure 6. Effect of sound treatments on zebrafish prolonged swimming speed. (a) Ambient condition. (b)
Continuous sound. (c) regular intermittent 1:1. (d) regular intermittent 1:4. (e) irregular intermittent 1:1-7. The
time was divided into three period bins for formal statistical analysis: the last 5 seconds before sound exposure,
the first 5 seconds during sound exposure and the last 5 seconds during sound exposure. Prolonged swimming
speed weakened with repeated exposure in all sound treatments. (NS= no significance, ***= $P \leq 0.001$, ****=
$P \leq 0.0001$).

1330

1331   **Figure 7.** Average percentage of fish time spent in bottom layer of tank (N=30). (a) Brief time (15 seconds before
1332   and 15 seconds exposed to sound) (NS= no significance, *= $P \leq 0.05$, **= $P \leq 0.01$). (b) Prolonged time (60 seconds
1333   before and 60 seconds exposed to sound) (NS= no significance, *= $P \leq 0.05$, **= $P \leq 0.01$, ***= $P \leq 0.001$). Bottom
1334   layer area for spatial displacement was defined as the bottom layer with a vertical distance of 10 cm from the
1335   bottom of the tank. (df= 1) Standard error changes (± 1).

1336

1337   **Figure 8.** Effect of sound exposure on horizontal spatial distribution of Zebrafish. (a) Brief time (15 seconds
1338   before and 15 seconds exposed to sound). (b) Prolonged time (60 seconds before and 60 seconds exposed to
1339   sound). The underwater speaker played back from the right tank. Bars show Means $\pm$ SE.

1340

Figure 9. Heat map of fish swimming in the tank environment. (a) Brief time (15 seconds before and 15 seconds
exposed to sound). (b) Prolonged time (60 seconds before and 60 seconds exposed to sound). The blue color
(#0000FE) indicates the 0-20 % of the fish in the tank. The aqua color (#01FFFF) indicates the 20-40 % of the
fish in the tank. The lime color (#00FF01) indicates the 40-60 % of the fish in the tank. The yellow color
(#FFFF01) indicates the 60-80 % of the fish in the tank. The red color (#FE0000) indicates the 80-100 % of the
fish in the tank. The underwater speaker played back from the right tank. For interpretation of the references to
colour in this figure legend, the reader is referred to the web version of this article.

1348

Figure 10. Effect of sound treatment on foraging behaviour of Zebrafish. (a) Food discrimination error as the
proportion of duckweed particles attacked relative to the total number of attacks to both duckweed particles and
water fleas from the introduction of food items until the end of sound exposure in sequence for each zebrafish
individual. (N=30, df=3.756, F=1.339, P=0.226) (b) Food handling error as the proportion of the total of water
fleas attacked that were missed or released again after initial grasping from onset of food introduction until the
end of sound exposure in sequence for each zebrafish individual. (N=30, df=2.825, F=26.023, P=0.000019).
Standard error changes (± 1)

1356

1357

1358

1359   Figure 1.1

1360

1386

1387

1388

1389   Figure 1.2

1390

1418

1419   Figure 2

1420

1421

1422

1423

1424   Figure3

1425

1426

1427

1428

1429

1430

1431

1432

1433

1434

1435

1436

1437

1438

1439

1440   Figure 4

1441

1442

1443

1444   Figure 5

1445

1446

1447

1448

1449

1450

1451

1452

1453

1454

1455

1456

1457

1458   Figure 6

1459

1460

1461

1462

1463

Figure 7

1465

1466

1467

1468

Figure 8

1470

1471

1472

1473   Figure 9

1474

1475

1476

1477

1478

1479

1480

1481

1482

1483

1484

1485

1486

1487

1488

1489

1490 Figure 10

1491

1492

1493

1494

1495