Experimental sound exposure modifies swimming activities and increases food handling error in adult zebrafish

Abstract:

Anthropogenic noise is increasing globally and is recognized as a source of environmental pollution in terrestrial and aquatic habitats. Sound is an important sensory stimulus for aquatic organisms and it may alter stress-related physiological indices and induce broad behavioural effects in a range of marine and freshwater fishes. Specifically, sound exposure may induce changes in swimming activities, feed efficiency and spatial distribution changes in fish. Here, we experimentally tested sound effects on swimming activities and foraging performance in thirty individually housed, captive adult Zebrafish (*Danio rerio*). Adult zebrafish and water fleas (*Daphnia magna*) were used as model predator and prey species, respectively. Acoustic stimuli consisted of four sound treatments with different temporal patterns. All had the same frequency range and were administered on average 121 dB re 1 μPa^2/Hz. Our results constitute strong evidence for sound-related effects on zebrafish behaviour. All sound treatments led to a significant increase in the number of startle responses, and the brief and prolonged swimming speed for zebrafish. We found sound effects on the spatial distribution of zebrafish; Although there were no significant sound-related changes for horizontal spatial displacement in all treatments, zebrafish swam significantly more in the lower layer of the tank except during the irregular intermittent 1:1-7 in brief sound exposure treatment. The results of foraging performance showed that food discrimination error was unaffected by sound treatments and was low for the zebrafish. However, food handling error was affected by sound treatments; all treatments induced a significant rise in handling error. This study highlights the impact of sound on zebrafish swimming activities, and that more feeding bouts are needed to consume the same number of food items increasing energy demand under noisy conditions.

Keywords: Anthropogenic noise, Behaviour, Foraging performance, Sound impact, Zebrafish

1. Introduction

Currently, due to increased human activities and global technology advancement since the Industrial Revolution, the Earth's environment has undergone extensive changes (Normandeau Associates, 2012). These environmental changes can affect the planet itself and living organisms, which threatens global biodiversity (Kunc et al., 2016). The rapid rate of

change poses many environmental challenges (Tuomainen and Candolin, 2011) in both terrestrial and aquatic habitats. For example, one of the main sources of environmental pollution, which may also can be recognized as an environmental stress stimulus, is anthropogenic noise that not only affects terrestrial animals, but also has many consequences on aquatic organisms (Popper et al., 2020; Slabbekoorn et al., 2010; Slabbekoorn and Ripmeester, 2008). Anthropogenic noise sources in aquatic habitats, such as military sonars, shipping activities, wind turbines, and pile-driving can occur frequently, having varying temporal patterns, and are widespread geographically (McDonald et al., 2006; Normandeau Associates, 2012). Consequently, anthropogenic noise has changed underwater soundscapes worldwide, represents a very subtle driver of environmental change and is a novel challenge to aquatic organisms.

Recent studies have investigated the impacts of anthropogenic noise on a wide range of taxa and across a variety of scales (Barber et al., 2009; Morley et al., 2014; Normandeau Associates, 2012; Slabbekoorn et al., 2010; Thomsen et al., 2021; Tyack, 2008). Anthropogenic noiseee can cause a range of physical, physiological, and behavioural disorders in aquatic organisms, including marine mammals (Erbe et al., 2018; Moore et al., 2012; Southall et al., 2008), seabirds (Bermúdez-Cuamatzin et al., 2018; Green et al., 2016; Hansen et al., 2020), reptiles (Injaian et al., 2020; Simmons and Narins, 2018), fish (Hastings and Popper, 2005; Hawkins, 1986; Mills et al., 2020; Popper et al., 2003), and invertebrates (Carroll et al., 2017; Coquereau et al., 2016; Murchy et al., 2019). Because of the high propagation rate and low attenuation over large distances in underwater environments, acoustic stimuli play an important role in aquatic environments (Tyack, 1998; Popper and Hastings, 2009; Slabbekoorn et al., 2010).

Behavioural effects are the most likely to occur and thus drive stress (Smith et al., 2004; Popper and Hawkins, 2019). Acoustic stimuli can act as a distracting stimulus (Popper and Carlson, 1998; Chan et al., 2010), interfere with detecting prey and antipredator behaviour (Hawkins and Myrberg, 1983), compromise foraging performance (Neo et al., 2015; Purser and Radford, 2011; Shafiei Sabet et al., 2015; Voellmy et al., 2016), disrupt reproductive behaviour (McCloskey et al., 2020), or mask important acoustic signals and cues for conspecific recognition and communication purposes (Codarin et al., 2009; Amorim et al., 2015; De Jong et al., 2018b; Hawkins and Picciulin, 2019) in a range of marine and freshwater species.

Many marine and freshwater fishes have well-developed hearing abilities that provide them with a key biological advantage to detect low intensities and perceive a broad range of frequencies (Hawkins, 1986; Heath et al., 2021; Popper et al., 2019; Wahlberg and Westerberg, 2005; Wysocki et al., 2006). While there are well-documented studies on how sound affects marine fish behaviour (de Jong et al., 2018a; Mortensen et al., 2021; Peng et al., 2015), and its adverse repercussions at the individual and community level, much less is known about how sound affects the behaviour of freshwater fish. To the best of our knowledge, few studies exist on this topic (Fedoroff, 2021; Mickle and Higgs, 2018; Pieniazek et al., 2020). Moreover, sound exposure can change spatial distribution and swimming behaviour of fish which may consequently affect foraging success if they avoid foraging in noisy food areas (de Vincenzi et al., 2021; Hanache et al., 2020; Hubert et al., 2021; Shafiei Sabet et al., 2016a). There is growing evidence that foraging performance is likely to be modulated by anthropogenic noise: instead of performing successful feeding bouts, fish can instead display typical behavioural stress responses (Purser and Radford, 2011; Shafiei Sabet et al., 2015; Voellmy et al., 2016). Currently, little is known about the effects of sound exposure on swimming activity and foraging performance of fish, although there are a few well-documented studies. It has been shown that increased boating activity was associated with a reduction in activity rates, changed vertical distribution and compromised foraging success of free-ranging mulloway (*Argyrosomus japonicus*) (Payne et al., 2015) and Mediterranean Damselfish (*Chromis chromis*) (Bracciali et al., 2012).

Other studies have shown that experimental sound exposure increases performance errors and therefore causes a negative impact on foraging efficiency in both the three-spined stickle back (*Gasterosteus aculeatus*) (Purser and Radford, 2011) and the European minnow, (*Phoxinus phoxinus*) (Voellmy et al., 2014a). More recently our previous study has also shown a clear sound impact on zebrafish foraging performance; more food handling errors occurred under noisy conditions (Shafiei Sabet et al., 2015). A primary consequence of sound exposure would appear to be a shift in the spatial displacement. The resulting disturbance might modify the allocated foraging time budget, foraging patterns, and the relative abundance of prey and predatory species. Such changes in turn may increase foraging energy demand and the amount of time fish must allocate to foraging which, subsequently may affect food searching, discrimination, and handling.

In general, *Danio rerio* is known as a model fish species in behavioural, environmental, and biomedical studies (Cachat et al., 2010; Egan et al., 2009; Whitfield, 2002). Zebrafish is a

member of the Cypriniformes order and acclimates well in captivity (Detrich et al., 2011) and
naturally lives in the tropical freshwater (Spence et al., 2008). Zebrafish have specialized
hearing structures, the Weberian ossicles, with an auditory system that has homologies to the
mammalian auditory system (Weber, 1820; Alexander, 1964; Whitfield, 2002).

Progress in the field of behavioural biology, conservation ecology, and toxicology is also
associated with the study of invertebrates. Daphnia is a small crustacean that is an important
part of the food web in freshwater habitats and inhabits many types of shallow water bodies
(Ebert, 2005; Parejko and Dodson, 1991; Reynolds, 2011).

In the present study, which is a follow up to recent work (Shafiei Sabet et al., 20215), we
experimentally tested the hypothesis that experimental sound exposure produced by an
underwater speaker affects the general swimming activities and foraging behaviour of
predatory zebrafish upon water flea prey under laboratory conditions. Compared to Shafiei
Sabet er al., 2015 where we used an in-ear speaker, in this study we adopted an underwater
speaker to increase sound pressure levels and induce a potential gradient in the arena. We
also extended the time of behavioural observation to explore more long-term effects of sound
on the zebrafish. Our specific goals were: first, to assess the effect of experimental sound
exposure on zebrafish activities including swim speed and spatial displacement; secondly, to
estimate whether the temporal pattern of sound exposure influences zebrafish behaviour; and
third, to verify our recent laboratory-based findings of sound impacts on zebrafish swimming
activity and foraging behaviour.

2. Materials and Methods

This study was performed in the ornamental fish breeding facility center at University of
Guilan, Sowmeh Sara, Iran (37°17′39″N, 49°19′55″E), using a 50 ×15 ×20 cm aquarium
between 10:00 and 14:00 every day. Thirty adult zebrafish (15 F: 15 M) were used and all
zebrafish were approximately 45 days old and of the wild-type, short-fin variety. They
weighed approximately (mean± s.d.) 1.23 ± 0.02 g) and were obtained from an ornamental
fish breeding shop in Sowmeh Sara, Iran. Zebrafish were stored in a stock glass tank
(50×30×40 cm) for two weeks and adapted to environmental conditions to reduce possible
stress and hormonal changes from transportation and captivity conditions (Deakin et al.,
2019). The fish were fed 0.8 mm commercial Biomar® feed until the day before the
experiment (Neo et al., 2015). Water fleas were caught at the campus of the department in

Sowmeh Sara every morning, during the entire experiment (See Shafiei Sabet et al., 2015;
Shafiei Sabet et al., 2019).

2.1. Sound treatments

Four sound treatments with different temporal patterns along with a control treatment were
used. The first treatment served as control treatment in which the fish were exposed to
ambient noise (AN). Experimental treatments consisted of regular intermittent sound (IN)
with a fast pulse rate (1:1) (Figure. 1 (a)), regular intermittent sound with slow pulse rate
(1:4) (Figure. 1 (b)), irregular intermittent sound (1:1-7) (Figure. 1 (c)), and continuous sound
(CS) (Figure. 1 (d)). All three intermittent treatments include one second of sound, but the
difference between these sound treatments is the intervals between these sounds (silence
time) (See also Shafiei Sabet et al., 2015). The sound treatments were generated and modified
using Audacity® software (2.3.1) at the frequency range that zebrafish can detect(300-1500
Hz) (Higgs et al., 2002) as well as encompassing the bandwidth of anthropogenic sounds that
overlap with the zebrafish's hearing range, such as from vehicles, pump systems, and pile
drivers (Slabbekoorn et al., 2010). Sound was produced by the software in the same
frequency range (400-2000 Hz) based on settings from our earlier study (Shafiei Sabet et al.,
2016).

2.2. Experiment Tank

The experimental tank (50×20×15 cm) was prepared with a black background to improve the
contrast between Daphnia and fish in the video file. Zebrafish swimming activities were
filmed by a video camera (Panasonic HC-V180 Full HD) at a distance of 50 cm from the
front of the test tank. Sound treatments were played back as stereo WAV files using a Laptop
(Sony Vaio SVF1421A4E) connected to an underwater speaker (custom-build speaker, 30 W,
10 Hz- 10 KHz). In this experiment, a divider plate was placed in the tank to divide its length
in half (25×20×15 cm) in order to restrict the zebrafish swimming environment and make the
entire swimming space visible for video-recording (Figure. 2). The test tank was surrounded
by black plastic so that the fish's behaviour was not affected by other visual factors. Only the
camcorder lens passed through the plastic, which was the same for all treatments and
repetitions. To isolate the experiments from external, confounding sounds, we covered the
walls at the entrance to the experimental room with egg boxes, both inside the room and in
the hallway.

The underwater speaker used in the experiment was placed horizontally behind a separator
plate (See Figure. 2). Then, each test was performed as follows: after ten minutes, the sound
treatment was played by the speaker and sound player for 20 minutes. The food item
(Daphnia) and a non-food item (Duckweed) were added to the experiment tank after ten
minutes from the onset of each sound playback across all sound treatments. The fish were
exposed to all five sound treatments on the same day, with a 15-minute interval between
treatments. The procedure was repeated the next day with the next fish naïve to any sound
exposure and no experience with sound treatments. The sound treatment playback order was
randomized daily and balanced across days. This approach was previously described to
quantify behavioural indices of zebrafish in response to acoustic measurements in this system
(Shafiei Sabet et al., 2015). The photoperiod used in the experiment was 12:12 (Higgs et al.,
2002; Villamizar et al., 2014) and the light intensity averaged 62 lux, measured by a light
meter (TES_1336A – TES Electrical Electronic Corp. Taiwan). Water temperature and the
amount of dissolved oxygen (DO) in the water were measured during the experiment and
were maintained at 26±1 °C at 8±1 mg / L, respectively.

2.3. Underwater sound measurement in the tank

Sound files were played back using a sound player connected to a custom-build underwater
speaker attached to a custom-build sound tuning amplifier (See Shafiei Sabet et al., 2015).
The level of sound pressure was recorded by a hydrophone (model Aquarian Scientific AS-1)
which was connected to the amplifier (model PA-4) and a Tascam linear PCM recorder
(model DR-100MKII). Sound pressure levels and power spectral density of the sound for the
continuous treatment were evaluated in R studio software (Version. 1.1.456). The sound
pressure of continuous sound treatment during playback was on average 121 dB re 1 μPa^2/Hz
for 5 seconds and the ambient sound pressure was on average 96 dB re 1 μPa^2/Hz for 5
seconds (Figure. 3).

2.4. Effect of sound on Zebrafish swimming behaviour

To investigate the effect of sound on the behaviour of Zebrafish, all animals were introduced
individually in the experimental tank. The fish were introduced to the test tank the day prior
to testing (Overnighting) to acclimate to the environment so that they could clearly use the
entire space of the tank for their swimming and display normal behaviour (Neo et al., 2015;
Shafiei Sabet et al., 2015). All fish were given 30 minutes to acclimate in the experiment
tank.

Zebrafish behavioural responses were video recorded for a maximum of 30 minutes for each
treatment (10 mins before exposure and 20 mins during exposure). Swimming behaviour
parameters such as the number of startle response (defined as the peak swimming speed at a
rate of more than 10 cm per second that occurs immediately after the sound was played for
one minute (See Shafiei Sabet et al. (2015)), brief swimming speed (during the last 5 seconds
before sound exposure and 5 seconds with the onset of sound) and prolonged swimming
speed (during one minute before the start of sound exposure and one minute with the onset of
sound exposure) were evaluated for all treatments. Also, the spatial distribution of fish in the
tank was explored. In the vertical dimension, tank height was divided into two parts: zero to
7.5 cm and 7.5 to 15 cm based on the 15cm dewatering height of the tank. To check the
distance from the sound source in the horizontal dimension, the length of the tank was
divided into three parts: zero to 8.33 cm, 8.33 to 16.66 cm and 16.66 to 25 cm.

2.5. Effect of sound on foraging behaviour of zebrafish on water fleas

The effect of sound on the foraging behaviour of zebrafish was investigated by mixing 5
water fleas (~3 mm; target prey species) and 5 non-food substances (~ 3mm non-food items;
duckweed) were mixed in 25 ml beaker and adding the mixture gently to the fish tank. This
procedure was done in the same manner for all treatments. The water fleas were in the same
sizes that caught with plastic Pasteur pipettes with suitable entrance tube diameter to decrease
damaging water fleas, which is suitable for feeding this species of fish at puberty and can be
orally ingested by the fish (Shafiei Sabet et al., 2015). Both food and non-food items are
naturally present in the habitat of this fish. Foraging performance was assessed as described
in Shafiei Sabet et al., 2015.

2.6. Behavioural information processing and statistical analysis

Zebrafish behaviour videos were converted to 10 frames per second by Xilisoft Video
Converter Ultimate software. This was done to reduce the magnification of time, , which
increases the accuracy of the fish swimming survey by reducing the fish speed for spatial
inspection per second. Logger Pro® video tracking software (Vernier Software, Beaverton,
OR, USA, version 3.10.1) was used to examine behavioural responses including the number
of startle responses, average swimming speeds, and spatial distribution changes of the fish.
Data analysis was performed using SPSS 25 software and Excel 2016. The normality of the
data was evaluated by Kolmogorov-Smirnov test and the homogeneity of the data was
assessed using Levene's test. Then, significant differences between the treatments were

assessed with a repeated measures ANOVA analysis and using Tukey multi-range test. A Huynhe Feldt correction was performed when sphericity could not be confirmed in the repeated measures ANOVA. Bonferroni corrected post hoc tests were performed when ANOVA test results were significant. The level of significance in this study was considered $P<0.05$. A custom-written acoustic calibration script in R studio software (Version 1.1.456) was also used to evaluate sound pressure levels and power spectral density that were played by the underwater speaker.

3. Ethical note

All housing, handling and experimental conditions were in accordance with the guidelines for the treatment of animals in behavioural research and teaching (ASAB, 2020). Water fleas and zebrafish were allowed to acclimate gradually to laboratory conditions before they were used in any of the experiments and showed no signs of adverse effects from the experimental conditions. Zebrafish showed only a brief startle response upon onset of the moderate sound playbacks and no mortalities or physical damage were observed during experiments (Neo et al., 2015; Shafiei Sabet et al., 2015). There are no legal requirements for studies involving invertebrates such as waterfleas (Daphnia) in Iran.

4. Result

4.1. Impact of sound on swimming behaviour of Zebrafish

Experimental sound exposure changed zebrafish swimming activities in several ways. Sound treatments significantly increased the startle responses and swimming speed in fish (Figure. 4). The number of startle responses instantly increased upon exposure to sound so that all treatments showed a significant difference compared to the ambient condition (repeated measures ANOVA: $F_{3.23,93.68}=6.31$, $P<0.001$). However, no significant differences were observed in terms of temporal patterns between sound treatments ($P>0.05$) (Figure 4a). Also, the brief swimming speed (5 seconds before the sound and 5 seconds during the sound) was significantly different in all treatments compared to the ambient condition (repeated measures ANOVA: $F_{3.06,88.70}=11.17$, $P<0.001$), although there was no significant difference between the sound treatments with different temporal patterns for this measure ($P>0.05$) (Figure 4b). This difference in prolonged swimming speed (60 seconds before the sound and 60 seconds during the sound) was also impacted, so that there was a significant effect between sound treatments compared to ambient conditions (repeated measures ANOVA: $F_{3.39,98.34}=7.72$,

$P<0.001$). However, again there was no significant difference between the different sound
treatments ($P>0.05$) (Figure 4C).

Based on Figure 5, in all four sound treatments (CS, 1:1, 1:4, 1:1-7), a sudden increase in
swimming speed was observed once sound treatments were played at the 60 second
observation time. Observations also showed that in all sound treatments, fish swim speed
returned to baseline after 60 seconds of playback. In ambient conditions (Figure 5a), no
significant difference in swimming speed was observed during the entire time period. While
in continuous sound and regular intermittent 1:1 treatment (Figure 5b+ 5c), a significant
difference was observed when comparing the 5 seconds before sound playback and the first 5
seconds after sound playback (CS= repeated measure ANOVA: $F_{4.54,131.66}=9.53$, $P\leq0.001$,
1:1= repeated measure ANOVA: $F_{3.85,111.72}=11.72$, $P\leq0.001$). However, for both treatments,
the fish returned to their baseline speed during the last 5 seconds of sound exposure: there
was no significant difference in swim speed between the 5 seconds preceding sound exposure
and the final 5 seconds of playback ($P>0.05$). This pattern was also observed in the regular
intermittent 1:4 and irregular intermittent 1:1-7 treatments (Figure 5d+ 5e) with a more
intense difference before and during the sound (1:4= repeated measure ANOVA:
$F_{6.24,181.01}=14.17$, $P\leq0.001$, 1:1-7= repeated measure ANOVA: $F_{7.14,207.15}=15.01$, $P\leq0.001$).
Like the continuous and 1:1 treatments, no significant difference was observed between the
swim speed of the 5 seconds preceding sound exposure and the last 5 seconds of the
treatment: the fish returned to their baseline speed ($P>0.05$).

Changes in the spatial distribution of zebrafish were investigated when exposed to sound
treatments in vertical scale (lower layer) (Figure. 6) and horizontal scale (X position) (Figure.
7). According to Figure 6a, there was no effect on the average percent of time spent in the
lower layer of the tank during sound treatments exposure over a brief time (15 seconds before
sound and 15 seconds during sound exposure) ($F_{3.77,109.47}=1.486$, $P=0.214$) or nor was there
an interaction of treatment*times ($F_{3.55,102.92}=0.634$, $P>0.05$). However, there was a main
effect of times ($F_{1,29}=28.274$, $P<0.001$). In the ambient (AN) and irregular intermittent sound
(1:1-7) treatments, there was not a significant effect of time (AN= $F_{1,29}=28.274$, $P>0.05$, 1:1-
7= $F_{1,29}=28.274$, $P>0.05$). However, there was a significant difference in two sound
treatments before and during sound exposure: continuous sound (CS) and intermittent 1:4
(CS= $F_{1,29}=28.274$, $P<0.001$, 1:4= $F_{1,29}=28.274$, $P<0.001$). Also, in the intermittent 1:1
treatment there was a significant difference before and after sound exposure ($F_{1,29}=28.274$,
$P<0.01$). According to Figure 6b, the average percent of time spent in the lower layer of the

tank during sound treatments over a prolonged time (60 seconds before sound and 60 seconds

during sound exposure), was not affected by treatment ($F_{3.65,105.92}=1.837$, $P>0.05$) or the

interaction of treatment*times ($F_{4,116}=1.780$, $P=0.137$). However, there was a significant

effect of times ($F_{1,29}=35.398$, $P<0.001$). In control treatment (AN) there was not a significant

difference before and after sound exposure ($F_{1,29}=35.398$, $P>0.05$). While in the continuous

treatment (CS) and intermittent 1:4 treatment, there was a significant difference before and

after sound exposure (CS= $F_{1,29}=35.398$, $P<0.001$, 1:4= $F_{1,29}=35.398$, $P<0.001$). The

intermittent 1:1 treatment was also significantly different before and after sound exposure

($F_{1,29}=35.398$, $P<0.001$). Also, the irregular intermittent sound treatment (1:1-7) showed a

significant difference before and after sound exposure ($F_{1,29}=35.398$, $P<0.05$).In contrast,

there was no significant difference in the horizontal profile (X position) between sound

treatments and control conditions, both in the brief and prolonged time ($F_{4,116}=1.369$, $P>0.05$)

(Figure 7a), ($F_{3.78,109.71}=1.810$, $P>0.05$) (Figure 7b). Further, according to Figure 8 the effects

of different sound patterns on the spatial distribution of fish can be seen as a heat map in brief

(figure 8a) and prolonged duration (figure 8b). Acoustic treatments affected changes in

swimming pattern from the top layer to the bottom layer but had no effect on the distance

from the sound source located on the right side of the tank.

4.2. Impact of sound on foraging performance of Zebrafish

According to Figure 9a, none of the acoustic treatments had a significant difference compared

to the ambient noise treatment on zebrafish food discrimination error (repeated measure

ANOVA: $F_{4,116}=1.339$, $P>0.05$). In fact, there was no food discrimination error between food

and non-food item by broadcasting sound treatments compared to control treatment.

However, for food handling error, all acoustic treatments showed a significant difference

compared to the control treatment (repeated measure ANOVA: $F_{2.82,81.91}=26.023$, $P\leq0.001$).

There was no significant difference observed between sound treatments though ($P>0.05$)

(Figure. 9b).

5. Discussion

Our results unequivocally demonstrate that acoustic stimuli affect zebrafish behaviour in

different ways. In this experiment, zebrafish swimming behaviour indices such as the number

of startle responses, the difference in brief swimming speed (within 5 seconds), the difference

in prolonged swimming speed (within 60 seconds), and the spatial distribution in response to

continuous and intermittent sound patterns were examined. The number of startle responses,

indicative of anxiety in zebrafish and other aquatic species (Blaser et al., 2010; Maximino et
al., 2010), in different sound treatments were significantly different compared to the control
condition. However, these changes did not show a significant difference between sound
treatments. Also, there was a significant increase in in brief and prolonged swimming speeds
in all sound treatments compared to the control condition. The increased swim speeds were
not significantly different between sound treatments though. Moreover, in support of
previous work (Shafiei Sabet et al., 2015), this data showed the same impact of sound
exposure on foraging performance in zebrafish. All will be discussed further in the following
sections.

5.1. Startle responses as a specific indicator of moderate anxiety in zebrafish

The startle response is an involuntary action that is controlled by a pair of brain neurons in
the Mauthner (M-) cells in the mesencephalon and plays a major role in the decision-making
process (Eaton et al., 1977; Eaton et al., 1991; Mirjany et al., 2011; Zottoli, 1977). Increasing
the intensity of the sound triggers an involuntary and very sudden C-startle response by
Mauthner cells in the mesencephalon, leading to an involuntary escape response in fish
(Eaton et al., 1977; Eaton et al., 1991). The fish's increased number of startle response and
brief swimming speed in response to sound treatments, are due to behavioural responses
related to fear and anxiety in this species. Previous studies have shown that sounds increase
motor acceleration and startle responses in zebrafish (Neo et al., 2015; Shafiei Sabet et al.,
2015). However, Shafiei Sabet et al. (2015), using an in-air speaker as a sound source,
reported no significant differences in the startle response of fish who were exposed to
continuous or intermittent (1:1) sound compared to ambient noise, which is not consistent
with the results of this study. The reason for this discrepancy in stress-related swimming
behaviour could be due to differences in ambient noise intensity before the test, differences in
the method of ensonifiying the fish tank (under water compared to in-air speaker), different
housing conditions for the fish, and genetic differences and individual characteristics of the
zebrafish.

Other species of fish respond similarly in response to sound exposure. European minnow
(*Phoxinus phoxinus*) and sticklebacks (*Gasterosteus aculeatus*) showed a significant increase
in the number of startle responses when exposed to sound, similar to the results of this study
(Purser and Radford, 2011; Voellmy et al., 2014a). A startle response at the onset of sudden
sound exposure is a common behavioural reaction in fish kept in captivity and in laboratory

conditions. Of course, fish in open water and natural conditions can also show behavioural
responses related to fear and anxiety (Neo et al., 2016; Staaterman et al., 2020). Spiga et al.
(2017) stated that European seabass (*Dicentrarchus labrax*) also showed a higher number of
startle responses under continuous and intermittent sound treatments compared to an ambient
treatment.

A startle response by prey fish is a behavioural adaptation to enhance survival rate in the
context of predator-prey relationships (Webb, 1986). When hearing the sounds of predator
fish and those related to an attack, the prey fish will explosively swim in the opposite
direction of the perceived sound in order to increase their chance of escaping from the
predatory species. Sound can affect the prey fish's decision-making power against sources of
danger, and how they assesses risk (Dukas, 2004). However, it can also act as a distraction
and cause inappropriate responses to danger (Chan et al., 2010; Simpson et al., 2015). It has
been suggested that increased sound levels can potentially impair the perception of danger by
predatory fish species (Slabbekoorn et al., 2010). It has been suggested that involuntary and
acquired behavioural responses related to fear and anxiety are associated with the potential
for the presence of danger (Blaser et al., 2010; Maximino et al., 2010). The quality and
quantity of behavioural responses of captive fish to brief and severe stress stimuli are
different from those of fish living in a naturalistic habitat: the behavioural responses can be
amplified in captivity compared to what is seen in the wild (Malavasi et al., 2004). One
reason for these differences could be due to the ability of most wild fish to respond before the
signal reaches the stimulus threshold, so that wild fish have a longer time to make decisions
and escape from potential danger. They also have more space available than in controlled
laboratory environments. Another reason could be the high level of basal stress potential in
controlled laboratory environments, which paired with additional stress from perception of
the predator species can lead to more intense responses.

In addition to behavioural responses, increasing sound levels can also affect physiological
responses in the laboratory and in the natural habitat of fish. The study of Spiga et al. (2017)
and Radford et al. (2016) showed that sound exposure had a significant effect on the number
of opening and closing of gills and thus on the gill ventilation of European seabass
(*Dicentrarchus labrax*) compared to the control treatment. This increase in oxygen demand
by European bass (*Dicentrarchus labrax*), which is accompanied by an increase in gill
ventilation and the opening and closing of gill operculum, indicates an increase in stress
levels. Santulli et al. (1999) showed that blood biochemical parameters including cortisol and

glucose in European seabass (*Dicentrarchus labrax*) increased in sound treatments compared
to the ambient treatment. Staaterman et al. (2020) stated that anthropogenic noise treatments
in the natural environment also have the potential to affect stress-related physiology in coral
reef fish, so that the amount of cortisol in sound treatments was significantly increased
compared to the control treatment.

5.2. Sound impacts on the behavioural tolerance in swimming activities

In past work performed on zebrafish at the group and individual level, different sound
patterns had a significant effect on zebrafish swimming activities such as the brief and
prolonged swimming speed of the fish compared to the ambient treatment, which is
consistent with the observations in the present study (Neo et al., 2015; Shafiei Sabet et al.,
2015; Shafiei Sabet et al., 2016a). In the experiment of Neo et al. (2015), groups of 5
zebrafish had increased mean swimming speed during the intermittent (1:1) treatment
compared to ambient and other sound treatments. Similar results were also previously
observed in cod fish (*Gadus morhua*) and European seabass (*Dicentrarchus labrax*), which
was consistent with the results of this study (Handegard et al., 2003) (Neo et al., 2018). The
prolonged swimming speed of zebrafish weakened with repeated sound exposure suggesting
a habituation process. Zebrafish might experience negative consequences if they are disturbed
and do not habituate accordingly to sound exposure. Shafiei Sabet et al. (2016b) compared
the effect of sound exposure on the swimming behaviour of two species of The Lake Victoria
cichlids (*Haplochromis piceatus*) and zebrafish (*Danio rerio*). That study showed that the
application of sound treatments reduced the swimming speed of cichlids (*Haplochromis
piceatus*) and increased the swimming speed of zebrafish (Shafiei Sabet et al., 2016b). The
reason of the difference in the swimming speed of cichlids (*Haplochromis piceatus*) can be
related to species-specific behavioural traits in response to acoustic stimuli, genetic
characteristics, and habitat conditions.

5.3. The effect of sound on the spatial distribution of zebrafish

Studies on the stress indices of zebrafish in the face of different sound patterns showed that
with the onset of sound treatments, the spatial distribution of fish changes and the fish shows
a greater tendency to swim in the lower layer of the aquarium environment. Neo et al. (2015)
revealed that zebrafish showed startle response and increased brief swimming speed upon the
start of sound treatments, and their spatial distribution changed so that fish was more inclined
to swim in the top layers of the test tank. Also, there were no observations of freezing or

standing in the lower layer of the tank (Neo et al. 2015). However, the results of the present

experiment showed that zebrafish spent more time in the lower layer of the test tank, which is

contrary to the report of Neo et al. (2015). One of the reasons for this vertical distribution

difference could be the amount of sound intensity emitted by the treatments used in these two

studies. The intensity of sound previously emitted in acoustic treatments was equal to 112 dB

re 1μ Pa, which is less than the intensity of sound in this study (121 dB re 1μ Pa). Therefore,

potentially a variety of increase in sound intensity between treatments and ambient noise can

lead to different responses in fish.

It is also important to consider the potential impacts of acoustic stimuli in real field

conditions. Other fishes have also shown spatial distribution changes in response to acoustic

stimuli in field conditions. Neo et al. (2018) found that anthropogenic sounds increase the

swimming depth of European seabass and distance from the sound source, which is consistent

with the present study in terms of vertical distribution results. In a field study (Kok et al.,

2021) have shown that bottom-moored echosounders, representative of a high intensity

impulsive intermittent anthropogenic noise, affect the abundance, schooling cohesion

behaviour, and swimming depth of pelagic fish. Two recent telemetry tagging studies

demonstrated the effects of another intermittent source of sound, seismic surveys, on free-

ranging benthic fish species. Bruce et al. (2018) showed that anthropogenic noise cause

changes in diurnal local activity patterns and general swimming speed shifts of marine fishes

and therefore may negatively affect the catchability of marine fish catchability and the. van

der Knaap et al. (2021) demonstrated in a field study that Atlantic cod (*Gadus morhua)* left

noisy areas more quickly and decrease their swimming activity compared to the ambient

sound treatment. They also have shown that seismic surveys can disrupt diurnal activity

cycles which may have long term negative consequences on this species' foraging success

and fitness. Although to date, there is a gap in knowledge in terms of freshwater soundscape

environments and how elevated sound levels affect fish in freshwater habitats (Bolgan et al.,

2016; Linke et al., 2018; Rountree et al., 2020; Putland and Mensinger, 2020).

Swimming towards the upper layer of the tank at the beginning of the sound treatment has

been considered to be driven by curiosity and searching behaviour. This is based on past

work where low-pitched sounds were emitted during routine feeding by care staff, which

attracted the fish's attention and caused them to gather at the top of the tank in anticipation of

a meal (See Shafiei Sabet et al., 2015), However, the opposite response seen here, where fish

gathered at the bottom layer of the tank during sound treatments, indicates the occurrence of

stress and fear in fish, which is similarly expressed in studies on other fish species (Neo et al.,

2018; Sarà et al., 2007). Responses to other stimuli including chemicals and fear extract have

shown that fish move to the lower layer of the tank upon exposure and this spatial distribution

response is interpreted as a behavioural indicator of fear in many fish species (Gerlai et al.,

2000; Gerlai et al., 2006). Another reason for the contrasting results observed in Neo et al.

(2015) and the present study could be the difference in ensonifying underwater using

speakers: an in-air speaker was used in the previous study (Neo et al., 2015) and an

underwater speaker was used in the present study. The use of in-air speakers to broadcast

sound treatments leads to more slightly intense sound in the deeper parts of the tank than in

the middle and the surface layers, which may cause the fish to move and escape towards the

surface layer where less sound intensity is felt (Shafiei Sabet et al., 2015). However this

observed spatial behaviour change with the onset of sound exposure may be also an

exploratory behaviour and curiosity of fish (c.f. Neo et al., 2015)

The spatial distribution data found in this study is not consistent with what was found

previously (Shafiei Sabet et al. (2015), despite using treatments with similar sound intensity

(122 dB re 1μ Pa). The reason for this difference in spatial distribution could be the use of in

air speaker in the study by Shafiei Sabet et al. (2015) and underwater speaker in this study, as

well as the complexity of sound distribution patterns and sound gradients in aquarium

environments (Campbell et al., 2019). Other factors such as fish tank size and available arena,

fish life stage, and location of the fish storage tank, as well as differences in the species,

genetics, and personality of the fish could contribute to the differing results. However, no

significant differences were observed in the horizontal distribution by both Shafiei Sabet et

al. (2015) and the current study.

Based on available sources for measuring sound intensity and scattering patterns in aquarium

environments, Parvulescu (1967) and Akamatsu et al. (2002) demonstrate the complexity and

variability of sound scattering patterns and sound gradients in enclosed aquarium tank

environments. This indicates a major limitation of studying the spatial distribution of aquatic

animals in enclosed and controlled environments that must be considered. Therefore, in order

to study the distribution patterns of fish and other aquatic species more accurately, it is

recommended to conduct field studies in the species' natural environments in order to obtain

a more comprehensive understanding of sound-dependent distribution patterns in aquatic

species.

5.4. The importance of particle motion in fish tanks; behavioural observations for future
works

In order to understand the behavioural changes of zebrafish in response to acoustic stimuli,
first of all, it is very important to understand the mechanisms behind and that how the species
detects and processes, and how it behaviourally responds to sound (Hawkins and Popper,
2020). Because well-documented studies already showed that the auditory system of fishes
evolved primarily to detect particle motion, many fishes are most sensitive to particle motion
and they can use it to determine the direction of a sound source (Hawkins and Popper, 2018;
Popper and Hawkins, 2018; Sand and Bleckmann, 2008; Sisneros and Rogers, 2016).
Underwater sound consists of both sound pressure and particle motion (Popper and Hawkins,
2018). In fact, all fishes and all invertebrates primarily detect particle motion but not all
detect sound pressure. Only some of fishes, including the zebrafish, are sensitive to sound
pressure as well as the particle motion (Popper and Fay, 2011; Popper and Hawkins, 2018).
There are some studies revealing directional hearing and sound source localization in fish
under laboratory conditions and in free sound fields. Schuijf (1975) proposed that the cod
(*Gadus morhua*) are able to detect sound directionality by monitoring the particle motion of
the sound field, presumably employing the directional orientation of the inner ear sensory
cells (Dale, 1976). Although, Schuijf (1975) also concluded that to determine the direction of
a sound source, the direction of only particle motion may not be sufficient. It has already
been shown that cod could discriminate between signals coming towards the head as
compared to those coming towards the tail (Buwalda et al., 1983; Schuijf and Buwalda,
1975). They argued that to eliminate any remaining 180^0 ambiguities, directional hearing
might involve both comparing the responses of hear cells oriented in different directions and
also analysis of the phase relationship between the sound pressure and particle motion
components (Schuijf, 1976).

5.5. How important particle motion is to fishes and invertebrates

We did not mention the levels and direction of the particle motion that is generated within the
fish tank because of our restriction in acquiring the particle motion measurement devices due
to the COVID-19 pandemic. Therefore, we believe it is premature to conclude that zebrafish
cannot localize sound source in our experimental set up. One reason might be because we
know very little about hearing in only over 120 of the more than 33000 known fish species
(Ladich and Fay, 2013) and that the empirical and theoretical work on sound source

localization and directional hearing in fishes have been contradictory and obscure for decades (Sisneros and Rogers, 2016). Further reasoning can be attributed to practicality: it is difficult to monitor particle motion in a fish tank, there is a lack of easily used and reasonably priced instrumentation to measure particle motion, there is a lack of sound exposure criteria for particle motion, and finally there is a lack of particle motion measurement standards (Popper et al., 2014).

Within an aquarium tank the levels of particle motion are often highest at the water surface, and close to the tank walls, when an underwater loudspeaker is used (Jones et al., 2019). However, resonant frequencies and reverberation may influence propagation and spectro-temporal structure of the received acoustic stimuli in fish ranks (Jones et al., 2019). Our fish moved towards the lower levels of the tank, which may be because the particle motion levels were highest close to the water surface, and lower at the bottom of the tank. It is always important to monitor the particle motion when examining the effects of sounds upon fishes and invertebrates (Nedelec et al., 2016; Popper and Hawkins, 2018). Moreover, invertebrates are especially sensitive to substrate vibration (Aimon et al., 2021; Hawkins et al., 2021; Morley et al., 2014; Roberts et al., 2016), and some fish are too. Particle motion measurement may play an important role in answering crucial biological and ecological questions relating to fishes and other species among taxa (Nedelec et al., 2016). Thus, in doing future experiments to explore anthropogenic noise impacts on the behaviour of fishes or invertebrates under laboratory conditions it is necessary to develop open source and accessible protocols for monitor both particle motion on three axes and sound pressure.

5.6. Acoustic stimuli trigger negative effects in zebrafish foraging performance

In the present study, the parameters of fish foraging behaviour such as food discrimination error and food handling error were examined. The results showed that the zebrafish did not show any significant difference in food discrimination error when exposed to sound treatments compared to ambient noise. Also, the food handling error data showed that all sound treatments were significantly different compared to the ambient treatment, but these sound treatments were not different from each other. Here we confirmed our earlier findings (Shafiei Sabet et al., 2015) that sound impacts go beyond a single species. Experimental sound exposure caused more food handling errors and foraging in zebrafish, which led to more waterflea survival in noisy conditions.

Previously, sound treatments significantly affected the foraging performance of sticklebacks: food discrimination error and food handling error increased significantly and foraging performance reduced significantly compared to the ambient treatment (Purser and Radford (2011). In terms of food discrimination error, this is not consistent with the results of the present experiment. One reason for this difference could be due to physiological differences between the fish species and their diets. Particularly, potential differences in the fish's visual abilities could be one of the factors influencing these differences. In terms of food handling error, a similar result was previously observed with those of the present experiment. In other studies, by Voellmy et al. (2014b), there was a significant difference in the number of unsuccessful takes of Daphnia in stickleback and no significant difference in minnow fish, which can generally indicate the physiological difference between the two species and possibly the difference in visual sense between the two species. The minnow fish, which belongs to the Cyprinidae family and is similar to zebrafish family, did not show a significant difference in the unsuccessful takes of Daphnia.

In a study almost similar to the present experiment, Shafiei Sabet et al. (2015) designed an experiment to investigate the effect of different sound patterns on the swimming and foraging behaviour of zebrafish and found that the application of sound treatments caused a significant difference in fish handling error. There was no significant difference in food discrimination error, which was consistent with the results of this experiment. Sound playback can influence the ability to identify a food substance, and make decisions about attacking it. Since zebrafish have a strong visual sense and use this modality for hunting, sounds can affect the ability to perceive potential vision potentially via attentional shifts and distraction. Field studies have also focused on the effects of sound on other fish species. In a comparative study between locations in the Laurentian Great Lakes and in laboratory conditions, Pieniazek et al., 2020 showed that experimental white noise and boat noise changed swimming patterns and foraging behaviour of freshwater fish species in different ways and with highly variable patterns according to family. They demonstrated that captive black bullhead, *Ameiurus melas*, foraged less and startled more when exposed to both white noise and boat noise treatments. Furthermore, escape behaviour has also been demonstrated in elasmobranches in response to anthropogenic noise (Mickle et al., 2020; Mickle et al., 2022). In a laboratory-based study on two fish species, the spot tail shiner (*Notropis hudsonius*) and the bluegill sunfish (*Lepomis macrochirus*), with different hearing abilities it has been shown that shipping noise caused similar but also specie specific responses (Stasso et al., 2023). Our laboratory experiments

were conducted to explore usefulness of laboratory based experiments to assess anthropogenic noise impacts and address fundamental behavioural research questions. We hypothesised questions to verify that acoustic stimuli affect swimming behaviour and foraging performance of captive zebrafish as a model species and establish insights and new hypotheses for future exploration to be conducted on other freshwater fish species in captivity and in the field.

6. Conclusion

The results of this study highlighted impacts of acoustic stimuli on a freshwater fish species and confirmed our earlier study findings on the same fish species (zebrafish) under laboratory conditions. Our findings show that zebrafish swimming parameters and foraging behaviour in various sound treatments were significantly different compared to ambient noise conditions. Sound pollution as a stressor in the brief and prolonged time can cause behavioural changes and disturbances in individual aquatic species and have broad and important repercussions on the communities of an ecosystem. It should be noted that the results of this study are obtained in captivity and under laboratory conditions therefore the interpretation of the results should be done with caution and attention should be given to the conditions of natural environments and behavioural limitations of any species.

Funding:

The current research was funded by the Iranian Ministry of Science, Research and Technology, University of Guilan with the grant number: 971.511.210004.

Appendices

Data accessibility:

All data used for the analyses reported in this article and some videos are available from the figshare. Moreover, the data and some videos that support the findings of this article are available from the first author upon request.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments:

Thanks to Jeroen Hubert and Ozkan Sertlek for their guide and advice on hydrophone calibrations, editing R script versions and acoustic measurements. Special tanks to Hans Slabbekoorn for providing accessibility to acoustic measurement devices. We also thank Narjes Karimi for helping us in reviewing the results and graphs.

References:

Aimon, C., Simpson, S.D., Hazelwood, R.A., Bruintjes, R., Urbina, M.A., 2021. Anthropogenic underwater vibrations are sensed and stressful for the shore crab Carcinus maenas. Environmental Pollution 285, 117148. https://doi.org/10.1016/j.envpol.2021.117148.

Akamatsu, T., Okumura, T., Novarini, N., Yan, H.Y., 2002. Empirical refinements applicable to the recording of fish sounds in small tanks. The Journal of the Acoustical Society of America 112, 3073-3082. https://doi.org/10.1121/1.1515799.

Alexander, R. 1964. The structure of the Weberian apparatus in the Siluri, in Proceedings of the Zoological Society ofLondon, Wiley Online Library, pp. 419–440. https://doi.org/10.1111/j.1469-7998.1964.tb04507.x.

Amorim, M.C.P., Vasconcelos, R.O., Fonseca, P.J., 2015. Fish sounds and mate choice, Sound communication in fishes. Springer, pp. 1-33. https://doi.org/10.1007/978-3-7091-1846-7_1

Animal Behaviour. 2020. Guidelines for the treatment of animals in behavioural research and teaching. Animal Behaviour. 159:1–11. https://doi.org/10.1006/anbe.1996.0293.

Barber, J.R., Fristrup, K.M., Brown, C.L., Hardy, A.R., Angeloni, L.M., Crooks, K.R., 2009. Conserving the wild life therein: Protecting park fauna from anthropogenic noise. 26, 26-31. http://www.nature.nps.gov/ParkScience/index.cfm?ArticleID=370&Page=1

Bermúdez-Cuamatzin, E., López-Hernández, M., Campbell, J., Zuria, I., Slabbekoorn, H., 2018. The role of singing style in song adjustments to fluctuating sound conditions: a comparative study on Mexican birds. Behavioural processes 157, 645-655. https://doi.org/10.1016/j.beproc.2018.04.005.

Blaser, R., Chadwick, L., McGinnis, G., 2010. Behavioral measures of anxiety in zebrafish (*Danio rerio*). Behavioural brain research 208, 56-62. https://doi.org/10.1016/j.bbr.2009.11.009.

Bolgan, M., Chorazyczewska, E., Winfield, I.J., Codarin, A., O'Brien, J. and Gammell, M., 2016. First observations of anthropogenic underwater noise in a large multi-use lake. Journal of Limnology, 75(3), pp.644-651. doi: 10.4081/jlimnol.2016.1405.

Bracciali, C., Campobello, D., Giacoma, C., Sara, G., 2012. Effects of nautical traffic and noise on foraging patterns of Mediterranean damselfish (*Chromis chromis*). PloS one 7, e40582. https://doi.org/10.1371/journal.pone.0040582.

Bruce, B., Bradford, R., Foster, S., Lee, K., Lansdell, M., Cooper, S., Przeslawski, R., 2018. Quantifying fish behaviour and commercial catch rates in relation to a marine seismic survey. Marine environmental research 140, 18-30. https://doi.org/10.1016/j.marenvres.2018.05.005.

Buwalda, R., Schuijf, A., Hawkins, A., 1983. Discrimination by the cod of sounds from opposing directions. Journal of Comparative Physiology 150, 175-184. https://doi.org/10.1007/BF00606367.

653
654 Cachat, J., Canavello, P., Elegante, M., Bartels, B., Hart, P., Bergner, C., Egan, R., Duncan, A., Tien,
655 D., Chung, A., 2010. Modeling withdrawal syndrome in zebrafish. Behavioural brain research 208,
656 371-376. https://doi.org/10.1016/j.bbr.2009.12.004.
657
658 Campbell, J., Shafiei Sabet, S., Slabbekoorn, H., 2019. Particle motion and sound pressure in fish
659 tanks: a behavioural exploration of acoustic sensitivity in the zebrafish. Behavioural processes 164,
660 38-47. https://doi.org/10.1016/j.beproc.2019.04.001.
661
662 Carroll, A., Przeslawski, R., Duncan, A., Gunning, M., Bruce, B., 2017. A critical review of the
663 potential impacts of marine seismic surveys on fish & invertebrates. Marine Pollution Bulletin 114, 9-
664 24. https://doi.org/10.1016/j.marpolbul.2016.11.038.
665
666 Chan, A.A.Y.-H., Giraldo-Perez, P., Smith, S., Blumstein, D.T., 2010. Anthropogenic noise affects
667 risk assessment and attention: the distracted prey hypothesis. Biology letters 6, 458-461.
668 https://doi.org/10.1098/rsbl.2009.1081.
669
670 Coquereau, L., Grall, J., Chauvaud, L., Gervaise, C., Clavier, J., Jolivet, A., Di Iorio, L., 2016. Sound
671 production and associated behaviours of benthic invertebrates from a coastal habitat in the north-east
672 Atlantic. Marine biology 163, 127. https://doi.org/10.1007/s00227-016-2902-2.
673
674 Dale, T., 1976. The labyrinthine mechanoreceptor organs of the cod *Gadus morhua* l-(teleostei:
675 gadidae). a scanning electron microscopical study, with special reference to the morphological
676 polarization of the macular sensory cells. http://pascal-
677 francis.inist.fr/vibad/index.php?action=getRecordDetail&idt=PASCAL7650248453.
678
679 de Jong, K., Amorim, M., Fonseca, P.J., Heubel, K.U., 2018a. Noise affects multimodal
680 communication during courtship in a marine fish. Frontiers in Ecology and Evolution 6,
681 113. https://doi.org/10.3389/fevo.2018.00113.
682
683 De Jong, K., Amorim, M.C.P., Fonseca, P.J., Fox, C.J., Heubel, K.U., 2018b. Noise can affect
684 acoustic communication and subsequent spawning success in fish. Environmental Pollution 237, 814-
685 823. https://doi.org/10.1016/j.envpol.2017.11.003.
686
687 de Vincenzi, G., Micarelli, P., Viola, S., Buffa, G., Sciacca, V., Maccarrone, V., Corrias, V., Reinero,
688 F.R., Giacoma, C., Filiciotto, F., 2021. Biological sound vs. anthropogenic noise: assessment of
689 behavioural changes in *Scyliorhinus canicula* exposed to boats noise. Animals 11, 174.
690 https://doi.org/10.3390/ani11010174.
691
692 Deakin, A.G., Spencer, J.W., Cossins, A.R., Young, I.S., Sneddon, L.U., 2019. Welfare challenges
693 influence the complexity of movement: fractal analysis of behaviour in zebrafish. Fishes 4, 8.
694 https://doi.org/10.3390/fishes4010008.
695
696 Detrich, H.W., Detrich III, H.W., Westerfield, M., Zon, L.I., 2011. The zebrafish: genetics, genomics
697 and informatics. Academic Press.
698
699 Detrich, H.W., Westerfield, M., Zon, L.I., 2004. The zebrafish: Biology. ACADEMIC PRESS.
700
701 Dukas, R., 2004. Causes and consequences of limited attention. Brain, Behavior and Evolution 63,
702 197-210. https://doi.org/10.1159/000076781.
703
704 Eaton, R.C., Bombardieri, R.A., Meyer, D.L., 1977. The Mauthner-initiated startle response in teleost
705 fish. Journal of Experimental Biology 66, 65-81. https://doi.org/10.1242/jeb.66.1.65.
706

Eaton, R.C., Didomenico, R., Nissanov, J., 1991. Role of the Mauthner cell in sensorimotor integration by the brain stem escape network. Brain, Behavior and Evolution 37, 272-285. https://doi.org/10.1159/000114365.

Ebert, D., 2005. Ecology, epidemiology, and evolution of parasitism in Daphnia. National Library of Medicine. https://edoc.unibas.ch/13505/1/Ebert-Parasitism_in_Daphnia-2005-A4-print.pdf.

Egan, R.J., Bergner, C.L., Hart, P.C., Cachat, J.M., Canavello, P.R., Elegante, M.F., Elkhayat, S.I., Bartels, B.K., Tien, A.K., Tien, D.H., 2009. Understanding behavioral and physiological phenotypes of stress and anxiety in zebrafish. Behavioural brain research 205, 38-44. https://doi.org/10.1016/j.bbr.2009.06.022.

Erbe, C., Dunlop, R., Dolman, S., 2018. Effects of noise on marine mammals, Effects of anthropogenic noise on animals. Springer, pp. 277-309. https://doi.org/10.1007/978-1-4939-8574-6_10.

Fedoroff, K.J., 2021. Sound bites: the impact of boat noise on freshwater fish. University of Saskatchewan. http://hdl.handle.net/10388/13300.

Gerlai, R., Lahav, M., Guo, S., Rosenthal, A., 2000. Drinks like a fish: zebra fish (Danio rerio) as a behavior genetic model to study alcohol effects. Pharmacology Biochemistry and Behavior 67, 773-782. https://doi.org/10.1016/S0091-3057(00)00422-6.

Gerlai, R., Lee, V., Blaser, R., 2006. Effects of acute and chronic ethanol exposure on the behavior of adult zebrafish (*Danio rerio*). Pharmacology Biochemistry and Behavior 85, 752-761. https://doi.org/10.1016/j.pbb.2006.11.010.

Green, R.E., Langston, R.H., McCluskie, A., Sutherland, R., Wilson, J.D., 2016. Lack of sound science in assessing wind farm impacts on seabirds. Journal of Applied Ecology 53, 1635-1641. doi: 10.1111/1365-2664.12731.

Hanache, P., Spataro, T., Firmat, C., Boyer, N., Fonseca, P., Médoc, V., 2020. Noise-induced reduction in the attack rate of a planktivorous freshwater fish revealed by functional response analysis. Freshwater Biology 65, 75-85. https://doi.org/10.1111/fwb.13271.

Handegard, N.O., Michalsen, K., Tjøstheim, D., 2003. Avoidance behaviour in cod (Gadus morhua) to a bottom-trawling vessel. Aquatic Living Resources 16, 265-270. https://doi.org/10.1016/S0990-7440(03)00020-2.

Hansen, K.A., Hernandez, A., Mooney, T.A., Rasmussen, M.H., Sørensen, K., Wahlberg, M., 2020. The common murre (*Uria aalge*), an auk seabird, reacts to underwater sound. J. Acoust. Soc. Am 147, 4069-4074. https://doi.org/10.1121/10.0001400.

Hastings, M.C., Popper, A.N., 2005. Effects of sound on fish. California Department of Transportation.https://www.researchgate.net/profile/Arthur-Popper/publication/253413309_Update_on_exposure_metrics_for_evaluation_of_effects_of_sound_o n_fish/links/0c9605295e4f7183f7000000/Update-on-exposure-metrics-for-evaluation-of-effects-of-sound-on-fish.pdf.

Hawkins, A., Myrberg, A., 1983. Hearing and sound communication under water. Bioacoustics, a comparative approach. Academic Press, London, 347-405.

Hawkins, A.D., 1986. Underwater sound and fish behaviour, The behaviour of teleost fishes. Springer, pp. 114-151. https://doi.org/10.1007/978-1-4684-8261-4_5.

Hawkins, A.D., Hazelwood, R.A., Popper, A.N., Macey, P.C., 2021. Substrate vibrations and their
potential effects upon fishes and invertebrates. The Journal of the Acoustical Society of America 149,
2782-2790. https://doi.org/10.1121/10.0004773.

Hawkins, A.D., Picciulin, M., 2019. The importance of underwater sounds to gadoid fishes. The
Journal of the Acoustical Society of America 146, 3536-3551. https://doi.org/10.1121/1.5134683.

Hawkins, A.D., Popper, A.N., 2018. Directional hearing and sound source localization by fishes. The
Journal of the Acoustical Society of America 144, 3329-3350. https://doi.org/10.1121/1.5082306.

Hawkins, A.D., Popper, A.N., 2020. Sound detection by Atlantic cod: An overview. The Journal of
the Acoustical Society of America 148, 3027-3041. https://doi.org/10.1121/10.0002363.

Heath, V.L., Miehls, S., Johnson, N., Higgs, D.M., 2021. Behavioural response of sea lamprey
(*Petromyzon marinus*) to acoustic stimuli in a small stream. Canadian Journal of Fisheries and
Aquatic Sciences 78, 341-348. https://doi.org/10.1139/cjfas-2020-0301.

Higgs, D.M., Souza, M.J., Wilkins, H.R., Presson, J.C., Popper, A.N., 2002. Age-and size-related
changes in the inner ear and hearing ability of the adult zebrafish (*Danio rerio*). JARO: Journal of the
Association for Research in Otolaryngology 3, 174. 10.1007/s101620020035.

Hubert, J., van Bemmelen, J., Slabbekoorn, H., 2021. No negative effects of boat sound playbacks on
olfactory-mediated food finding behaviour of shore crabs in a T-maze. Environmental Pollution 270,
116184. https://doi.org/10.1016/j.envpol.2020.116184.

Injaian, A.S., Francis, C.D., Ouyang, J.Q., Dominoni, D.M., Donald, J.W., Fuxjager, M.J., Goymann,
W., Hau, M., Husak, J.F., Johnson, M.A., 2020. Baseline and stress-induced corticosterone levels
across birds and reptiles do not reflect urbanication levels. Conservation Physiology 8, 1.
10.1093/conphys/coz110.

Jones, I.T., Stanley, J.A., Bonnel, J., Mooney, T.A., 2019. Complexities of tank acoustics warrant
direct, careful measurement of particle motion and pressure for bioacoustic studies, Proceedings of
Meetings on Acoustics 5ENAL. Acoustical Society of America, p. 010005.
https://doi.org/10.1121/2.0001073.

Kok, A.C., Bruil, L., Berges, B., Sakinan, S., Debusschere, E., Reubens, J., de Haan, D., Norro, A.,
Slabbekoorn, H., 2021. An echosounder view on the potential effects of impulsive noise pollution on
pelagic fish around windfarms in the North Sea. Environmental Pollution, 118063.
https://doi.org/10.1016/j.envpol.2021.118063.

Kunc, H.P., McLaughlin, K.E., Schmidt, R., 2016. Aquatic noise pollution: implications for
individuals, populations, and ecosystems. Proceedings of the Royal Society B: Biological Sciences
283, 20160839. https://doi.org/10.1098/rspb.2016.0839.

Ladich, F., Fay, R.R., 2013. Auditory evoked potential audiometry in fish. Reviews in Fish Biology
and Fisheries 23, 317-364. 10.1007/s11160-012-9297-z.

Linke, S., Gifford, T., Desjonquères, C., Tonolla, D., Aubin, T., Barclay, L., Karaconstantis, C.,
Kennard, M.J., Rybak, F. and Sueur, J., 2018. Freshwater ecoacoustics as a tool for continuous
ecosystem monitoring. Frontiers in Ecology and the Environment, 16(4), pp.231-238.
https://doi.org/10.1002/fee.1779.

Malavasi, S., Georgalas, V., Lugli, M., Torricelli, P., Mainardi, D., 2004. Differences in the pattern of
antipredator behaviour between hatchery-reared and wild European sea bass juveniles. Journal of fish
biology 65, 143-155. https://doi.org/10.1111/j.0022-1112.2004.00545.x.

Maximino, C., de Brito, T.M., da Silva Batista, A.W., Herculano, A.M., Morato, S., Gouveia Jr, A., 2010. Measuring anxiety in zebrafish: a critical review. Behavioural brain research 214, 157-171. https://doi.org/10.1016/j.bbr.2010.05.031.

McCloskey, K.P., Chapman, K.E., Chapuis, L., McCormick, M.I., Radford, A.N., Simpson, S.D., 2020. Assessing and mitigating impacts of motorboat noise on nesting damselfish. Environmental Pollution 266, 115376. https://doi.org/10.1016/j.envpol.2020.115376.

McDonald, M.A., Hildebrand, J.A., Wiggins, S.M., 2006. Increases in deep ocean ambient noise in the Northeast Pacific west of San Nicolas Island, California. The Journal of the Acoustical Society of America 120, 711-718. https://doi.org/10.1121/1.2216565.

Mickle, M.F., Higgs, D.M., 2018. Integrating techniques: a review of the effects of anthropogenic noise on freshwater fish. Canadian Journal of Fisheries and Aquatic Sciences 75, 1534-1541. https://doi.org/10.1139/cjfas-2017-0245.

Mickle, M.F., Pieniazek, R.H. and Higgs, D.M., 2020. Field assessment of behavioural responses of southern stingrays (*Hypanus americanus*) to acoustic stimuli. Royal Society Open Science, 7(1), p.191544. http://dx.doi.org/10.1098/rsos.191544.

Mickle, M.F., Pieniazek, R., Stasso, J.J. and Higgs, D.M., 2022. Anthropogenic sounds induce escape behaviour in southern stingrays Hypanus americanus. Marine Ecology Progress Series, 694, pp.125-132. https://doi.org/10.3354/meps14100.

Mills, S.C., Beldade, R., Henry, L., Laverty, D., Nedelec, S.L., Simpson, S.D., Radford, A.N., 2020. Hormonal and behavioural effects of motorboat noise on wild coral reef fish. Environmental Pollution 262, 114250. https://doi.org/10.1016/j.envpol.2020.114250.

Mirjany, M., Preuss, T., Faber, D.S., 2011. Role of the lateral line mechanosensory system in directionality of goldfish auditory evoked escape response. Journal of Experimental Biology 214, 3358-3367. https://doi.org/10.1242/jeb.052894.

Moore, S.E., Reeves, R.R., Southall, B.L., Ragen, T.J., Suydam, R.S., Clark, C.W., 2012. A new framework for assessing the effects of anthropogenic sound on marine mammals in a rapidly changing Arctic. BioScience 62, 289-295. https://doi.org/10.1525/bio.2012.62.3.10.

Morley, E.L., Jones, G., Radford, A.N., 2014. The importance of invertebrates when considering the impacts of anthropogenic noise. Proceedings of the Royal Society B: Biological Sciences 281, 20132683. https://doi.org/10.1098/rspb.2013.2683.

Mortensen, L.O., Chudzinska, M.E., Slabbekoorn, H., Thomsen, F., 2021. Agent-based models to investigate sound impact on marine animals: bridging the gap between effects on individual behaviour and population level consequences. Oikos. https://doi.org/10.1111/oik.08078.

Murchy, K.A., Davies, H., Shafer, H., Cox, K., Nikolich, K., Juanes, F., 2019. Impacts of noise on the behavior and physiology of marine invertebrates: A meta-analysis. Proceedings of Meetings on Acoustics 5ENAL. Acoustical Society of America, p. 040002. https://doi.org/10.1121/2.0001217.

Nedelec, S.L., Campbell, J., Radford, A.N., Simpson, S.D., Merchant, N.D., 2016. Particle motion: the missing link in underwater acoustic ecology. Methods in Ecology and Evolution 7, 836-842.

https://doi.org/10.1111/2041-210X.12544.

Neo, Y., Hubert, J., Bolle, L., Winter, H., Slabbekoorn, H., 2018. European seabass respond more strongly to noise exposure at night and habituate over repeated trials of sound exposure. Environmental Pollution 239, 367-374. https://doi.org/10.1016/j.envpol.2018.04.018.

Neo, Y., Hubert, J., Bolle, L., Winter, H.V., Ten Cate, C., Slabbekoorn, H., 2016. Sound exposure changes European seabass behaviour in a large outdoor floating pen: Effects of temporal structure and a ramp-up procedure. Environmental Pollution 214, 26-34. https://doi.org/10.1016/j.envpol.2016.03.075.

Neo, Y.Y., Parie, L., Bakker, F., Snelderwaard, P., Tudorache, C., Schaaf, M., Slabbekoorn, H., 2015. Behavioral changes in response to sound exposure and no spatial avoidance of noisy conditions in captive zebrafish. Frontiers in behavioral neuroscience 9, 28. https://doi.org/10.3389/fnbeh.2015.00028.

Normandeau Associates, I., 2012. Effects of Noise on Fish, Fisheries, and Invertebrates in the U.S. Atlantic and Arctic from Energy Industry Sound-Generating Activities., in: Interior, A.L.S.f.t.U.S.D.o.t. (Ed.), p. 135. https://www.cbd.int/doc/meetings/mar/mcbem-2014-01/other/mcbem-2014-01-submission-boem-04-en.pdf.

Parejko, K., Dodson, S.I., 1991. The evolutionary ecology of an antipredator reaction norm: *Daphnia pulex* and *Chaoborus americanus*. Evolution 45, 1665-1674. doi: 10.1111/j.1558-5646.1991.tb02671.x..

Parvulescu, A., 1967. The acoustics of small tanks. Marine Bio Acoustics 2, 7-13. https://www.scirp.org/(S(i43dyn45teexjx455qlt3d2q))/reference/ReferencesPapers.aspx?ReferenceID=55337.

Payne, N.L., van der Meulen, D.E., Suthers, I.M., Gray, C.A., Taylor, M.D., 2015. Foraging intensity of wild mulloway *Argyrosomus japonicus* decreases with increasing anthropogenic disturbance. Marine biology 162, 539-546. https://doi.org/10.1007/s00227-014-2603-7.

Peng, C., Zhao, X., Liu, G., 2015. Noise in the sea and its impacts on marine organisms. International journal of environmental research and public health 12, 12304-12323. https://doi.org/10.3390/ijerph121012304.

Pieniazek, R.H., Mickle, M.F., Higgs, D.M., 2020. Comparative analysis of noise effects on wild and captive freshwater fish behaviour. Animal Behaviour 168, 129-135. https://doi.org/10.1016/j.anbehav.2020.08.004.

Popper, A.N., Carlson, T.J., 1998. Application of sound and other stimuli to control fish behavior. Transactions of the American Fisheries Society 127, 673-707. https://doi.org/10.1577/1548-8659(1998)127<0673:AOSAOS>2.0.CO;2.

Popper, A.N., Fay, R.R., 2011. Rethinking sound detection by fishes. Hearing research 273, 25-36. https://doi.org/10.1016/j.heares.2009.12.023.

Popper, A.N., Fewtrell, J., Smith, M.E., McCauley, R.D., 2003. Anthropogenic sound: effects on the behavior and physiology of fishes. Marine Technology Society Journal 37, 35-40. https://doi.org/10.4031/002533203787537050.

Popper, A.N., Hawkins, A.D., 2018. The importance of particle motion to fishes and invertebrates. The Journal of the Acoustical Society of America 143, 470-488. https://doi.org/10.1121/1.5021594.

Popper, A.N., Hawkins, A.D., 2019. An overview of fish bioacoustics and the impacts of
anthropogenic sounds on fishes. Journal of fish biology 94, 692-713.
https://doi.org/10.1111/jfb.13948.

Popper, A.N., Hawkins, A.D., Fay, R.R., Mann, D.A., Bartol, S., Carlson, T.J., Coombs, S., Ellison,
W.T., Gentry, R.L., Halvorsen, M.B., 2014. Sound exposure guidelines, ASA S3/SC1. 4 TR-2014
Sound Exposure Guidelines for Fishes and Sea Turtles: A Technical Report prepared by ANSI-
Accredited Standards Committee S3/SC1 and registered with ANSI. Springer, pp. 33-51.
https://doi.org/10.1007/978-3-319-06659-2_7.

Popper, A.N., Hawkins, A.D., Sand, O., Sisneros, J.A., 2019. Examining the hearing abilities of
fishes. The Journal of the Acoustical Society of America 146, 948-955.
https://doi.org/10.1121/1.5120185.

Popper, A.N., Hawkins, A.D., Thomsen, F., 2020. Taking the animals' perspective regarding
anthropogenic underwater sound. Trends in ecology & evolution 35, 787-794.
https://doi.org/10.1016/j.tree.2020.05.002.

Popper, A.N., Smith, M.E., Cott, P.A., Hanna, B.W., MacGillivray, A.O., Austin, M.E., Mann, D.A.,
2005. Effects of exposure to seismic airgun use on hearing of three fish species. The Journal of the
Acoustical Society of America 117, 3958-3971. https://doi.org/10.1121/1.1904386.

Purser, J., Radford, A.N., 2011. Acoustic noise induces attention shifts and reduces foraging
performance in three-spined sticklebacks (*Gasterosteus aculeatus*). PloS one 6, e17478.
https://doi.org/10.1371/journal.pone.0017478.

Putland, R.L. and Mensinger, A.F., 2020. Exploring the soundscape of small freshwater
lakes. Ecological Informatics, 55, p.101018. https://doi.org/10.1016/j.ecoinf.2019.101018.

Radford, A.N., Lèbre, L., Lecaillon, G., Nedelec, S.L., Simpson, S.D., 2016. Repeated exposure
reduces the response to impulsive noise in European seabass. Global change biology 22, 3349-3360.
https://doi.org/10.1111/gcb.13352.

Reynolds, C., 2011. Daphnia: Development of Model Organism in Ecology and Evolution-2011.
Freshwater Reviews 4, 85-87. https://doi.org/10.1608/FRJ-4.1.425.

Roberts, L., Harding, H.R., Voellmy, I., Bruintjes, R., Simpson, S.D., Radford, A.N., Breithaupt, T.,
Elliott, M., 2016. Exposure of benthic invertebrates to sediment vibration: from laboratory
experiments to outdoor simulated pile-driving, Proceedings of Meetings on Acoustics 4ENAL.
Acoustical Society of America, p. 010029. https://doi.org/10.1121/2.0000324.

Rountree, R.A., Juanes, F. and Bolgan, M., 2020. Temperate freshwater soundscapes: A cacophony of
undescribed biological sounds now threatened by anthropogenic noise. Plos one, 15(3), p.e0221842.
https://doi.org/10.1371/journal.pone.0221842.

Sand, O., Bleckmann, H., 2008. Orientation to auditory and lateral line stimuli, Fish bioacoustics.
Springer, pp. 183-231. https://doi.org/10.1007/978-0-387-73029-5_6.

Santulli, A., Modica, A., Messina, C., Ceffa, L., Curatolo, A., Rivas, G., Fabi, G., D'amelio, V., 1999.
Biochemical responses of European sea bass (*Dicentrarchus labrax* L.) to the stress induced by off
shore experimental seismic prospecting. Marine Pollution Bulletin 38, 1105-1114.
https://doi.org/10.1016/S0025-326X(99)00136-8.
967

Sarà, G., Dean, J., d'Amato, D., Buscaino, G., Oliveri, A., Genovese, S., Ferro, S., Buffa, G., Martire,
M.L., Mazzola, S., 2007. Effect of boat noise on the behaviour of bluefin tuna Thunnus thynnus in the
Mediterranean Sea. Marine Ecology Progress Series 331, 243-253. https://www.int-
res.com/abstracts/meps/v331/p243-253/.

Schuijf, A., 1975. Directional hearing of cod (*Gadus morhua*) under approximate free field
conditions. Journal of Comparative Physiology 98, 307-332. https://doi.org/10.1007/BF00709803.

Schuijf, A., 1976. The phase model of directional hearing in fish. Sound reception in fish, 63-86.

Schuijf, A., Buwalda, R., 1975. On the mechanism of directional hearing in cod (*Gadus morhua* L.).
Journal of Comparative Physiology 98, 333-343. https://doi.org/10.1007/BF00709804.

Shafiei Sabet, S., Karnagh, S.A., Azbari, F.Z., 2019. Experimental test of sound and light exposure on
water flea swimming behaviour, Proceedings of Meetings on Acoustics 5ENAL. Acoustical Society
of America, p. 010015. https://doi.org/10.1121/2.0001270.

Shafiei Sabet, S., Neo, Y.Y., Slabbekoorn, H., 2015. The effect of temporal variation in sound
exposure on swimming and foraging behaviour of captive zebrafish. Animal Behaviour 107, 49-60.
https://doi.org/10.1016/j.anbehav.2015.05.022.

Shafiei Sabet, S., Van Dooren, D., Slabbekoorn, H., 2016a. Son et lumière: Sound and light effects on
spatial distribution and swimming behavior in captive zebrafish. Environmental Pollution 212, 480-
488. https://doi.org/10.1016/j.envpol.2016.02.046.

Shafiei Sabet, S., Wesdorp, K., Campbell, J., Snelderwaard, P., Slabbekoorn, H., 2016b. Behavioural
responses to sound exposure in captivity by two fish species with different hearing ability. Animal
Behaviour 116, 1-11. https://doi.org/10.1016/j.anbehav.2016.03.027.

Simmons, A.M., Narins, P.M., 2018. Effects of anthropogenic noise on amphibians and reptiles,
Effects of anthropogenic noise on animals. Springer, pp. 179-208. https://doi.org/10.1007/978-1-
4939-8574-6_7.

Simpson, S.D., Purser, J., Radford, A.N., 2015. Anthropogenic noise compromises antipredator
behaviour in European eels. Global change biology 21, 586-593. https://doi.org/10.1111/gcb.12685.

Sisneros, J.A., Rogers, P.H., 2016. Directional hearing and sound source localization in fishes. Fish
hearing and bioacoustics, 121-155. https://doi.org/10.1007/978-3-319-21059-9_7.

Slabbekoorn, H., Bouton, N., van Opzeeland, I., Coers, A., ten Cate, C., Popper, A.N., 2010. A noisy
spring: the impact of globally rising underwater sound levels on fish. Trends in ecology & evolution
25, 419-427. https://doi.org/10.1016/j.tree.2010.04.005.

Slabbekoorn, H., Ripmeester, E.A.P., 2008. Birdsong and anthropogenic noise: implications and
applications for conservation. Molecular ecology 17, 72-83. https://doi.org/10.1111/j.1365-
294X.2007.03487.x.

Southall, B.L., Bowles, A.E., Ellison, W.T., Finneran, J.J., Gentry, R.L., Greene Jr, C.R., Kastak, D.,
Ketten, D.R., Miller, J.H., Nachtigall, P.E., 2008. Marine mammal noise-exposure criteria: initial
scientific recommendations. Bioacoustics 17, 273-275.
https://doi.org/10.1080/09524622.2008.9753846.

Spence, R., Gerlach, G., Lawrence, C., Smith, C., 2008. The behaviour and ecology of the zebrafish,
Danio rerio. Biological Reviews 83, 13-34. https://doi.org/10.1111/j.1469-185X.2007.00030.x.
1019

Spiga, I., Aldred, N., Caldwell, G.S., 2017. Anthropogenic noise compromises the anti-predator behaviour of the European seabass, *Dicentrarchus labrax* (L.). Marine Pollution Bulletin 122, 297-305. https://doi.org/10.1016/j.marpolbul.2017.06.067.

Staaterman, E., Gallagher, A., Holder, P., Reid, C., Altieri, A., Ogburn, M., Rummer, J., Cooke, S., 2020. Exposure to boat noise in the field yields minimal stress response in wild reef fish. Aquatic Biology 29, 93-103. https://doi.org/10.3354/ab00728.

Stasso, J.J., Pieniazek, R.H. and Higgs, D.M., 2023. Interspecific variation in the response of fish to anthropogenic noise. Freshwater Biology, 68(1), pp.25-32. https://doi.org/10.1111/fwb.14006.

Thomsen, F., Popper, A.N., Hawkins, A.D., 2021. Sound Impact Studies: A Reply to Risch et al. Trends in ecology & evolution 36, 382-384. https://doi.org/10.1016/j.tree.2021.01.012.

Tuomainen, U., Candolin, U., 2011. Behavioural responses to human-induced environmental change. Biological Reviews 86, 640-657. https://doi.org/10.1111/j.1469-185X.2010.00164.x.

Tyack, P.L., 1998. Acoustic communication under the sea. Animal acoustic communication, pp.163-220. https://doi.org/10.1007/978-3-642-76220-8_6.

Tyack, P.L., 2008. Implications for marine mammals of large-scale changes in the marine acoustic environment. Journal of Mammalogy 89, 549-558. https://doi.org/10.1644/07-MAMM-S-307R.1.

van der Knaap, I., Reubens, J., Thomas, L., Ainslie, M.A., Winter, H.V., Hubert, J., Martin, B., Slabbekoorn, H., 2021. Effects of a seismic survey on movement of free-ranging Atlantic cod. Current Biology 31, 1555-1562. e1554. https://doi.org/10.1016/j.cub.2021.01.050.

Villamizar, N., Vera, L.M., Foulkes, N.S., Sánchez-Vázquez, F.J., 2014. Effect of lighting conditions on zebrafish growth and development. Zebrafish 11, 173-181. https://doi.org/10.1089/zeb.2013.0926.

Voellmy, I.K., Purser, J., Flynn, D., Kennedy, P., Simpson, S.D., Radford, A.N., 2014a. Acoustic noise reduces foraging success in two sympatric fish species via different mechanisms. Animal Behaviour 89, 191-198. https://doi.org/10.1016/j.anbehav.2013.12.029.

Voellmy, I.K., Purser, J., Simpson, S.D., Radford, A.N., 2014b. Increased noise levels have different impacts on the anti-predator behaviour of two sympatric fish species. PloS one 9, e102946. https://doi.org/10.1371/journal.pone.0102946.

Voellmy, I.K., Purser, J., Simpson, S.D., Radford, A.N., 2016. Effects of previous acoustic experience on behavioral responses to experimental sound stimuli and implications for research, The effects of noise on aquatic life II. Springer, pp. 1191-1196. https://doi.org/10.1007/978-1-4939-2981-8_149.

Wahlberg, M., Westerberg, H., 2005. Hearing in fish and their reactions to sounds from offshore wind farms. Marine Ecology Progress Series 288, 295-309. 10.3354/meps288295.

Webb, P., 1986. Effect of body form and response threshold on the vulnerability of four species of teleost prey attacked by largemouth bass (*Micropterus salmoides*). Canadian Journal of Fisheries and Aquatic Sciences 43, 763-771. https://doi.org/10.1139/f86-094.

Whitfield, T.T., 2002. Zebrafish as a model for hearing and deafness. Journal of neurobiology 53, 157-171. https://doi.org/10.1002/neu.10123.

Weber, E. H. 1820. De aure et auditu hominis et animalium. Pars I (The Ear and Hearing of Humans and Animals) (Gerhard Fleischer, Leipzig).

1072 Wysocki, L.E., Dittami, J.P., Ladich, F., 2006. Ship noise and cortisol secretion in European
1073 freshwater fishes. Biological conservation 128, 501-508.
1074 https://doi.org/10.1016/j.biocon.2005.10.020.

1075
1076 Zottoli, S.J., 1977. Correlation of the startle reflex and Mauthner cell auditory responses in
1077 unrestrained goldfish. Journal of Experimental Biology 66, 243-254.
1078 https://doi.org/10.1242/jeb.66.1.243.

1079

1080

1081

1082

1083

1084

Figure legends

Figure 1: Waveforms and spectrograms (across a 20 s stime period) of all intermittent and continuous sound
patterns used in the experiment (see also Shafiei Sabet et al., 2015). (a) Regular intermittent sound treatment
with fast pulse rate (1:1), (b) Regular intermittent sound treatment with slow pulse rate (1: 4), (c) Irregular
intermittent sound (1:1-7), (d) Continuous sound treatments accordingly.

Figure 2: Experimental set-up. Schematic view of the test tank. A: Obscured pages of the test tank are used to
enhance the fish's visual contrast in the film. B: Underwater speaker space separated by screen from the fish
swimming space. C: Underwater speaker holder box. D: Underwater speaker.

Figure 3: Spectral distribution of continuous sound pressure level compared to silent treatment (dB ref 1
$\mu Pa^2/Hz$). Silence conditions (blue) and continuous sound playback (red). The diagram shows that the sound
intensity level has increased significantly in the range of hearing frequencies of zebrafish. For interpretation of
the references to colour in this figure legend, the reader is referred to the web version of this article.

Figure 4. Effect of sound exposure treatment on swimming behaviour of Zebrafish. (a) Number of
startle responses expressed as the difference between the first 60 seconds during sound exposure and the 60
seconds immediately preceding sound exposure: Fish were exposed to continuous sound (CS), three intermittent
sounds (1:1, 1:4, 1:1-7), and ambient noises (AN) as control treatment (N=30, F=6.312, P<0.001, Standard error
changes ($\pm$1)). (b) Brief swimming speed difference of Zebrafish between the first 5 seconds of sound exposure
and the last 5 seconds preceding sound exposure for each sound treatment and the ambient condition (N=30,
F=11.172, P<0.001, Standard error changes ($\pm$1)). (c) Prolonged swimming speed difference of Zebrafish
between the 60 seconds preceding sound exposure and the first 60 seconds during sound exposure on all five
treatments (N=30, F=7.725, P<0.001 Standard error changes ($\pm$1)).

Figure 5. Effect of sound treatments on zebrafish prolonged swimming speed: (a) Ambient condition, (b) continuous sound, (c) regular intermittent 1:1, (d) regular intermittent 1:4, (e) irregular intermittent 1:1-7. The time was divided into three period bins for formal statistical analysis: the last 5 seconds before sound exposure, the first 5 seconds during sound exposure and the last 5 seconds during sound exposure. Prolonged swimming speed decreased with repeated exposure in all sound treatments. (NS= no significance, ***= $P \leq 0.001$, ****= $P \leq 0.0001$).

Figure 6. Average percentage of fish time spent in bottom layer of tank (N=30). (a) Brief time (15 seconds before and 15 seconds after sound exposure) (NS= no significance, *= $P \leq 0.05$, **= $P \leq 0.01$). (b) Prolonged time (60 seconds before and 60 seconds after sound exposure) (NS= no significance, *= $P \leq 0.05$, **= $P \leq 0.01$, ***= $P \leq 0.001$). Bottom layer area for spatial displacement was defined as the bottom layer up to 10 cm from the bottom of the tank. (df= 1) Standard error changes (±1).

Figure 7. Effect of sound exposure on horizontal spatial distribution of Zebrafish. (a) Brief time (15 seconds before and 15 seconds after sound exposure). (b) Prolonged time (60 seconds before and 60 seconds after sound exposure). The underwater speaker played back from the right side of the tank. Bars show Means ± SE.

Figure 8. Heat map of fish swimming in the tank environment. (a) Brief time (15 seconds before and 15 seconds after sound exposure). (b) Prolonged time (60 seconds before and 60 seconds after sound exposure). The blue color (#0000FE) indicates the 0-20 % of the fish in the tank. The aqua color (#01FFFF) indicates the 20-40 % of the fish in the tank. The lime color (#00FF01) indicates the 40-60 % of the fish in the tank. The yellow color (#FFFF01) indicates the 60-80 % of the fish in the tank. The red color (#FE0000) indicates the 80-100 % of the fish in the tank. The underwater speaker played back from the right tank. For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.

Figure 9. Effect of acoustic stimulus on foraging activities of Zebrafish. (a) Food discrimination error; it is described as the proportion of duckweed particles attacked relative to the total number of attacks to both duckweed particles and water fleas from the introduction of food items until the end of sound exposure in sequence for each zebrafish individual (Shafiei Sabet et al., 2015). (N=30, df=3.756, F=1.339, P=0.226) (b) Food handling error; It is described as the proportion of the total of water fleas attacked that were missed or released again after initial grasping from onset of food introduction until the end of sound exposure in sequence for each zebrafish individual (Shafiei Sabet et al., 2015). (N=30, df=2.825, F=26.023, P=0.000019). Standard error changes (±1)

1142   Figure 1

1143

1169

1170

1171

1172   Figure 1 (continued)

1173

1201

1202   Figure 2

1203

1204

1205

1206   Figure 3

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218   Figure 4

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245   Figure 5

1246

1247

1248

1249

1250   Figure 6

1251

1252

1253

1254

1255   Figure 7

1256

1257

1258

1259

1260   Figure 8

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277   Figure 9

1278

1279

1280

1281

1282