1	Maternal parity affects Day 8 embryo gene expression in old mares
2	Emilie Derisoud ^{1,2} , Luc Jouneau ^{1,2} , Clothilde Gourtay ³ , Anne Margat ³ , Catherine Archilla ^{1,2} , Yan
3	Jaszczyszyn ⁴ , Rachel Legendre ⁵ , Nathalie Daniel ^{1,2} , Nathalie Peynot ^{1,2} , Michèle Dahirel ^{1,2} , Laurie Briot ³ ,
4	Frédéric De Geoffroy ³ , Véronique Duranthon ^{1,2} , Pascale Chavatte-Palmer ^{1,2}
5	
6	¹ Université Paris-Saclay, UVSQ, INRAE, BREED, Jouy-en-Josas, France
7	² Ecole Nationale Vétérinaire d'Alfort, BREED, Maisons-Alfort, France
8	³ IFCE, Plateau technique du Pin, Exmes, France
9	⁴ Institute for Integrative Biology of the Cell (I2BC), UMR 9198 CNRS, CEA, Paris-Sud University, Gif-
10	sur-Yvette, France
11	⁵ Institut Pasteur—Bioinformatics and Biostatistics Hub—Department of Computational Biology, USR
12	3756 IP CNRS, Paris, France
13	
14	Grant support:
15	This work was supported by the "Institut Français du Cheval et de l'Equitation" (grant numbers
16	CS_2018_23, 2018). The National Research Institute for Agriculture, Food and Environment (INRAE)
17	department Animal Physiology and Breeding Systems also supported this research.
18	
19	Correspondence:

- 20 Emilie Derisoud
- 21 <u>emilie.derisoud@gmail.com</u>

- 22 Bâtiment 230 UMR 1198 BREED
- 23 INRAE Domaine de Vilvert
- 24 78350 JOUY EN JOSAS, FRANCE
- 25
- 26 Pascale Chavatte-Palmer
- 27 pascale.chavatte-palmer@inrae.fr
- 28 Bâtiment 230 UMR 1198 BREED
- 29 INRAE Domaine de Vilvert
- 30 78350 JOUY EN JOSAS, FRANCE

31

32 Running title:

33 Embryo gene expression in old nulliparous mare

34

35 Summary sentence:

- 36 Mare's parity in old mares impacts the expression of genes related to development and molecule
- 37 exchanges in ICM and TE of blastocysts suggesting an adaptation to an altered environment.

38

39 Keywords:

40 Blastocyst; RNA sequencing; horse; equine

41 Abstract

42 As sport career is a priority in most of equine breeds, mares are frequently bred for the first time at an 43 advanced age. Both age and first gestation were shown to have a deleterious effect on reproduction 44 outcomes, respectively on fertility and offspring weight but the effect mare's parity in older mares on 45 embryo quality has never been considered. The aim of this project was to determine the effect of old 46 mare's nulliparity on gene expression in embryos. Day 8 post ovulation embryos were collected from 47 old (10-16 years old) nulliparous (ON, N=5) or multiparous (OM, N=6) non-nursing Saddlebred mares, inseminated with the semen of one stallion. Pure (TE part) or inner cell mass enriched (ICMandTE) 48 49 trophoblast were obtained by embryo bisection and paired end, non-oriented RNA sequencing 50 (Illumina, NextSeq500) was performed on each hemi-embryo. To discriminate gene expression in the 51 ICM from that in the TE, deconvolution (DeMixT R package) was used on the ICM and TE dataset. 52 Differential expression was analyzed (DESeq2) with embryo sex and diameter as cofactors using a false 53 discovery rate <0.05 cutoff. Although the expression of only a few genes was altered by mare's 54 nulliparity (33 in ICM and 23 in TE), those genes were related to nutrient exchanges and responses to 55 environment signaling, both in ICM and TE, suggesting that the developing environment from these 56 mares are not optimal for embryo growth. In conclusion, being nulliparous and old does not seem to be the perfect match for embryonic development in mares. 57

58 Introduction

In the equine industry, mares are bred until an advanced age for economic and sentimental reasons. Depending on the breed, mares older than 10 years old represent between 37 and 63% of the broodmares [1–3]. In addition, 4% of Thoroughbred mares in the UK are older than 18 years old at the time of covering [1].

63 Several reproductive parameters are affected by age and mares older than 10 years old can be 64 considered as already old for reproduction as their fertility has already started to progressively decline 65 [for review 4]. Oocytes are particularly affected by maternal age with alterations of spindle stability 66 [5–7], altered gene expression [8,9], and altered metabolism [8,10–12] being reported, all suggesting 67 that oocyte developmental potential is reduced in old mares. The resulting embryos were smaller at 68 the same developmental age in most studies [13-23] with altered gene expression [23] and 69 metabolism [11,12], suggesting impaired development that has mainly been related to the oocyte 70 quality. The reproductive tract, however, is also affected by maternal age. Indeed, more oviductal 71 masses [24], uterine morphological degenerations such as cysts [25-30] and fibrosis [31-33] as well as 72 more endometritis [19,25,34,35] are observed in old mares.

73 In most of farm animals, female parity, defined as the number of pregnancies that reached a viable 74 gestational age (stillbirth and live birth included), is highly correlated with age as to remain profitable 75 and stay in the farm, females must produce offspring regularly. In horses, however, the sport career is 76 prioritized and depending on discipline and breed, can last up to 15 years or even more, as in 77 warmblood dressage and show jumping. In Finn horse and Standardbred, nulliparous mares 78 represented, respectively, 20.5% and 15.5% of mares that were bred in Finland [2]. To the authors' 79 knowledge, the effect of nulliparity/primiparity on oocyte and embryo quality as well as fetal development has not been explored. In any case, the reproductive tract is affected by mare parity. 80 81 Indeed, the ventral position of the uterus [36] and the number of vascular degenerations in the endometrium [37] have been positively correlated with the number of foals. Furthermore, cervical 82

dilatation is poorer in nulliparous vs multiparous mares [38] and it was suggested that uterine 83 clearance was impeded. Primiparity, however, does not affect the prevalence of endometritis [34]. 84 Both age and parity thus affect mare reproductive efficiency but the cumulative effect of nulliparity 85 and aging has not been explored. 86 87 The aim of this study was to determine the effect of maternal nulliparity in old mares on embryo gene expression at the blastocyst stage. Old (>10 years) nulliparous and multiparous mares were 88 89 inseminated with the same stallion semen. Blastocysts were collected and bisected to separate the 90 pure trophoblast (TE_part) from the inner cell mass enriched hemi-embryo (ICMandTE). Gene 91 expression was analyzed by RNA-seg in each compartment.

92

93 Materials and methods

94 Ethics

95 The experiment was performed at the experimental farm of IFCE ("Institut Français du Cheval et de 96 l'Equitation – La jumenterie du Pin" research agreement D61-157-2 valid until November 2023). The 97 protocol was approved by the local animal care and use committee and by the regional ethical 98 committee ("Comité d'Éthique Normand en Matière d'Expérimentation Animale", approved under N° 99 CEEA - 54 in the National Registry of French Ethical Committees for animal experimentation) under 100 protocol number APAFIS#20857-2019051709319621 v3. All experiments were performed in 101 accordance with the European Union Directive 2010/63EU.

102

103 Embryo collection

Twenty-five multiparous mares (mostly Selle Français breed with some French Anglo-Arabian,
 Standardbred and Saddlebred) aged from 10 to 16 years old were included in this study. Multiparous

106 mares were defined as dams that had already foaled at least once while nulliparous mares were 107 defined as mares that have never foaled before the experiment. During the experimental protocol, 108 mares were managed in two herds, independent of mare group, in natural pastures 24h/day with free 109 access to water. The experiments took place from July 8th to August 13th, 2019. All mares remained 100 healthy during this period.

111 Mares were allocated to one of 2 groups according to their parity: nulliparous (ON, n = 11) and 112 multiparous mares (OM, n = 14). During the experimentation, mare's withers' height and weight were 113 measured. Characteristics of all mares and mares that produced an embryo are detailed in Table 1.

114 The mares' estrous period was monitored routinely in the morning by ultrasound with a 50-60Hz trans-115 rectal transducer. During estrus, ovulation was induced with a single injection of human chorionic 116 gonadotropin (i.v.; 1500IU; Chorulon® 5000; MSD Santé animale, France) as soon as one ovarian follicle 117 > 35mm in diameter was observed, together with marked uterine edema. Ovulation usually takes place 118 within 48h, with > 80% occurring 25 to 48h after injection [39,40]. At the same time and one day later, 119 mares were inseminated once with fresh or refrigerated semen containing at least 1 billion motile 120 spermatozoa from a single fertile stallion. Ovulation was checked every 12-24 hours by 121 ultrasonography. If no embryo was recovered, the procedure could be repeated once more.

Embryos were collected by non-surgical uterine lavage using prewarmed (37°C) lactated Ringer's solution (B.Braun, France) and EZ-Way Filter (IMV Technologies, France) on the morning, 8 days post ovulation. At Day 14 post ovulation, a pregnancy diagnosis was performed for each mare and they were treated with luprotiol, an analogue of prostaglandin F2α (i.m; 7.5 mg; Prosolvin, Virbac, France).

When an embryo was collected, a blood sampling was performed at the same time on heparin tube.
Plasma was recovered after centrifugation (3500 rpm, 10min at 4°C). Progesterone was measured in
plasma using ELISA assay as previously described [41,42].

130

131 Embryo bisection and RNA extraction

132 Using a binocular magnifying glass, collected embryos were immediately photographed with a size 133 standard to subsequently determine embryo diameter using ImageJ® software (version 1.52a; National 134 Institutes of Health, Bethesda, MD, USA). Then embryos were washed 4 times in commercially 135 available Embryo holding medium (IMV Technologies, France) at 34°C and bisected with a microscalpel 136 under binocular magnifying glass to obtain a trophoblast (TE_part) and an inner cell mass enriched 137 (ICMandTE) hemi-embryo. At this stage, the TE_part is composed of trophectoderm and endoderm 138 whereas the ICM is composed of epiblast cells. Directly after bisection, RNA extraction of each hemi-139 embryo was started in extraction buffer (PicoPure RNA isolation kit, Applied Biosystems, France) for 140 30 min at 42°C prior to storage at - 80°C. RNA was extracted later on from each hemi-embryo using 141 PicoPure RNA isolation kit (PicoPure RNA isolation kit, Applied Biosystems, France), which included a 142 DNAse treatment, following the manufacturer's instructions. RNA quality and quantity were assessed 143 with the 2100 Bioanalyzer system using RNA 6000 Pico kit (Agilent Technologies, France) according to 144 the manufacturer's instructions.

145

146 RNA sequencing

Five nanograms of total RNA were mixed with ERCC spike-in mix (Thermofisher Scientific, France) according to manufacturer's recommendations. Messenger RNAs were reverse transcribed and amplified using the SMART-Seq V4 ultra low input RNA kit (Clontech, France) according to the manufacturer recommendations. Nine PCR cycles were performed for each hemi-embryo. cDNA quality was assessed on an Agilent Bioanalyzer 2100, using an Agilent High Sensitivity DNA Kit (Agilent Technologies, France). Libraries were prepared from 0.15 ng cDNA using the Nextera XT Illumina library preparation kit (Illumina, France). They were pooled in equimolar proportions and sequenced (Paired

154	end 50-34 pb) on NextSeq500 instrument, using a NextSeq 500 High Output 75 cycles kit (Illumina,
155	France). Demultiplexing was performed with bcl2fastq2 version 2.2.18.12 (Illumina, France) and
156	adapters were trimmed with Cutadapt version 1.15 [43]. Only reads longer than 10pb were kept.
157	
158	RNA mapping and counting
159	As previously described [23], alignment was performed using STAR version 2.6 [44] on previously
160	modified Ensembl 99 EquCab3.0 assembly and annotation. Genes were then counted with
161	FeatureCounts [45] from Subreads package version 1.6.1.
162	
163	Availability of data and materials
164	The RNA sequencing data supporting the conclusions of this article are available in the GEO
165	repository, [accession: GSE188866;
166	https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE188866].
167	
168	
169	Data analysis
170	All statistical analyses were performed by comparing ON to OM (set as reference group) using R version
171	4.0.2 [46] on Rstudio software version 1.3.1056 [47].
172	Embryo were sexed using X Inactive Specific Transcript (XIST) expression as previously described [23].
173	Seven embryos were determined as females (4 in the ON group and 3 in the OM group) while 4 were
174	considered as males (1 in the ON group, and 3 in the OM group).
175	
176	Embryo recovery and fertility rate, embryo diameter and total RNA content analysis

Embryo recovery rates (ERR) were calculated as the number of attempts with at least one embryo collected/total number of attempts. Furthermore, fertility was calculated as the sum of embryo collections with at least one embryo and the number of positive pregnancy checks at Day 11 after a negative embryo collection on the total number of attempts. Both were analyzed using the Exact Fisher test to determine if maternal parity in old mares influenced embryo recovery and the probability of leaving an embryo in the uterus after uterine flush.

For total RNA content analyses, as embryos were bisected without strict equality for each hemiembryo, a separate analysis of ICMandTE and TE_part RNA quantities would not have been meaningful. Thus, ICMandTE and TE_part RNA quantities were summed up. With embryo diameter, both variables were analyzed using a linear model of nlme package version 3.1-148 [48] including maternal age and embryo sex, followed by 1000 permutations using PermTest function from pgirmess package version 1.6.9 [49]. Variables were kept in models when statistically significant differences were observed. Differences were considered as significant for p < 0.05.

190

191 Deconvolution of gene expression in ICMandTE using DeMixT

192 The deconvolution method has already been described in equine embryos [23]. Briefly, this method 193 enables the estimation of the relative gene expression of TE and ICM cell types within the hemi-embryo 194 ICMandTE which is composed of both trophoblast and inner cell mass in unknown relative proportions. 195 After filtering out all genes with at least 3 null count values in at least one group (ON or OM) per hemi-196 embryo (ICM and TE or TE part), removing genes with a null variance in TE part and adding the value "1" to all count values in ICMandTE and TE_part datasets, deconvolution was performed using the 197 198 DeMixT R package version 1.4.0 [50,51]. Output datasets were DeMixT ICM cells and 199 DeMixT TE cells, corresponding to the deconvoluted gene expression in ICM cells and TE cells of 200 ICMandTE, respectively.

At the end of deconvolution, a quality check was automatically performed by the DeMixT R package with the TE_part used as reference for DeMixT_TE_cells. Genes were automatically filtered out if the difference between average deconvoluted expression of reference cells in mixed samples and average expression of reference cells > 4.

205 Outputs of DeMixT_ICM_cells *vs* DeMixT_TE_cells, DeMixT_ICM_cells *vs* TE_part and ICMandTE *vs* 206 TE_part were compared with Deseq2 version 1.28.1 [52] to confirm that the deconvolution was 207 effective at separating gene expression. To check if deconvolution was efficient, as previously 208 described [23], the expression of several genes proper to ICM and TE cells in equine embryos identified 209 using literature search [53] was compared before and after deconvolution. Results of these analyses 210 were represented through manually drawn Venn diagrams as well as principal component analysis 211 graphics of individuals, using ggplot2 version 3.3.3 [54] and factoextra version 1.0.7 [55].

212

213 Maternal parity comparison for gene expression

All genes with an average expression <10 counts in both ON and OM per hemi-embryo (ICM or TE) were filtered out on the DeMixT_ICM_cells and TE_part datasets. Differential analyses were performed with Deseq2 version 1.28.1 [52] with the OM group as reference, without independent filtering and taking into account embryo diameter and sex in the model. Genes were considered differentially expressed (DEG) for FDR < 0.05 after Benjamini-Hochberg correction (also known as false discovery rate, FDR).

Equine Ensembl IDs were converted into Human Ensembl IDs and Entrez Gene names using gorth function in gprofiler2 package version 0.1.9 [56]. Genes without Entrez Gene names using gprofiler2 were manually converted when Entrez Gene names were available, using Ensembl web search function [57]. GO molecular function and GO Biological process annotations were obtained from Uniprot website.

225

226 Gene set enrichment analyses (GSEA)

227	After log transformation using RLOG function of DESeq2 version 1.28.1, gene set enrichment analyses
228	(GSEA) were performed on expressed genes using GSEA software version 4.0.3 (Broad Institute, Inc.,
229	Massachusetts Institute of Technology, and Regents of the University of California) [58,59] to identify
230	biological gene sets disturbed by maternal parity. Molecular Signatures Databases [60] version 7.1 (C2:
231	KEGG: Kyoto Encyclopedia of Genes and Genomes; REACTOME, C5: BP: GO biological process) were
232	used to identify most perturbed pathways. Pathways were considered significantly enriched for
233	FDR< 0.05. When the normalized enrichment score (NES) was positive, the gene set was enriched in
234	the ON group while when NES was negative, the gene set was enriched in the OM group.

235

236 **Results**

237 Embryo recovery rates, diameter, total RNA content and quality and progesterone 238 concentrations

Altogether, 32 embryo collections were performed (14 in ON and 18 in OM, 8 mares being flushed twice) and 15 embryos were obtained (6 from 5 ON mares and 9 from 8 OM mares). Two mares (one in each parity group) produced twin embryos.

Positive embryo collection rate was 36% and 44% in ON and OM, respectively and did not differ between groups (p = 0.72). The embryo recovery rate was 43% and 50% in ON and OM, respectively and did not differ between groups (p = 0.30). At the Day 14 pregnancy check, embryos were found in 3 OM mares (from 2 to 4 foalings) and none was found in the ON group. Fertility, calculated combining positive embryo collections and Day 14 pregnancy diagnosis, was 36% and 61% in ON and OM, respectively, and did not differ between groups (p = 0.29). Altogether, 7 and 11 double ovulations were observed, respectively, in ON and OM. The embryo recovery rate per ovulation at the time of embryo collection and after Day 14 pregnancy check were not different according to group (respectively, 29% and 29% in ON and 31% and 41% in OM, p = 1 and p = 0.39).

All embryos were expanded blastocysts grade I or II according to the embryo classification of McKinnon and Squires [61]. For each twin collection, one embryo was large and the other was small (766µm and 295µm; 829µm and 481µm). For both, as only one embryo per mare was required, only the largest embryo of the twins was chosen for further analysis. Altogether, all ON embryos but only 6 OM embryos out of 8 collected were RNA sequenced. The smallest OM embryo (480µm) and another one randomly chosen (907µm) of diameter were not sequenced.

In embryos selected for RNA sequencing, embryo diameter ranged from 562 μ m to 1426 μ m, with no effect of group on embryo diameter (p = 0.18). Female embryos, however, were significantly smaller than male embryos (in average 764 ± 223 μ m and 1046 ± 287 μ m, p < 0.05) without interaction between maternal parity and embryo sex. RNA yield per embryo ranged from 25.2 ng to 624 ng and was not related to parity (p = 0.43) nor embryo sex (p = 0.08).

The median RNA Integrity Number (RIN) was 9.7 (8.9 - 10 range). Between 34.6 and 54.1 million reads per sample were obtained after trimming. On average, 74.10% of the reads were mapped on the modified EquCab 3.0 using STAR and 67.07% were assigned to genes by featureCounts.

Except one old multiparous mare that had a progesterone plasma concentration of 3.9 ng/ml, progesterone concentrations in plasma were > 4 ng/ml for all mares (range from 8.3 to 25.6ng/ml with an average of 13.7ng/ml) and were not affected by mares' parity (p = 0.66).

269

270 Deconvolution of gene expression to discriminate ICM and TE gene expression in ICMandTE 271 hemi-embryos

After selecting genes with less than 3 non null count values in at least one group (ON or OM) per hemiembryo (ICMandTE or TE_part), 16,803 genes were conserved for deconvolution. In addition, nine genes were removed because their variance was null in the TE_part, as DeMixT does not allow the use of genes with a null variance in the pure sample. For these genes, the mean count in ICMandTE samples was lower or equal to 10 counts. One further gene was removed during the deconvolution because the deconvolution quality for this gene was not sufficient. Therefore, at the end of the deconvolution algorithm, 16,793 genes were available for differential analysis.

279 Before deconvolution, 303 genes were differentially expressed (FDR < 0.05) between the ICMandTE 280 and the TE part (Figure 1A). After deconvolution, the comparison between DeMixT ICM cells and 281 DeMixT_TE_cells yielded 7,116 differentially expressed genes while the comparison DeMixT_ICM_cells 282 vs TE part yielded 5,615 differentially expressed genes, with 5,103 in common (74%). Moreover, all 283 but one of the initially 303 differentially expressed genes before deconvolution were also identified as 284 differentially expressed in both post-deconvolution analyses. On the PCA graph of individuals, Axis 1 285 (21.8% of variance) separated well groups according to data origin. ICMandTE and TE part were 286 separated on axis 1 but very close before deconvolution (Figure 1B). DeMixT TE cells and TE part 287 were partly superposed, indicating that datasets before and after deconvolution have a similar global 288 gene expression; whereas the DeMixT_ICM_cells group is clearly separated from both, indicating that the deconvolution effectively enabled the separation of gene expression in the two cell types. 289

290 Only 5 of the 12 genes previously reported as more expressed in the ICM [53] were also identified 291 more expressed in the ICMandTE vs TE part comparison (Table 2). After deconvolution (comparison 292 DeMixT ICM cells vs TE part), 10 out of 12 of these genes were observed differentially expressed with 293 9 effectively more expressed in the ICM. The expression of Undifferentiated Embryonic Cell 294 Transcription Factor 1, UTF1, however, was identified decreased in the DeMixT_ICM_cells, in contrast 295 to the only published report [53]. In the TE, no gene previously identified was observed differentially 296 expressed in the comparison ICMandTE vs TE_part, i.e., before deconvolution. After deconvolution, 297 the expression of 3 of the 7 reported genes were found increased in TE part compared to DeMixT_ICM_cells. 298

These results indicated that a better qualification of genes expressed by ICM cells was enable by the deconvolution. Thus, for further analyses, TE_part and DeMixT_ICM_cells datasets have been studied.

302 Differential gene expression in deconvoluted ICM cells

303 After the filtering out of genes without an average expression ≥ 10 counts in at least one maternal age 304 group/hemi-embryo, 13,910 genes were considered as expressed in the ICM cells from ON or OM 305 embryos. Only 33 genes were differentially expressed (23 downregulated and 10 upregulated in ON) 306 (Figure 2 and Additional file 1). Respectively, 20 and 5 genes out of the down- and upregulated genes 307 were associated to a protein known and described in human. These 25 genes are presented in Table 308 3. Down regulated genes in the ICM of old nulliparous mares were involved in RNA processing and 309 transcription, immunity, nervous system development, lipid/protein transport, lipid metabolism, 310 chromatin remodeling, DNA repair, cell cycle, signaling and adhesion whereas up-regulated genes 311 were related to different biological processes such as ion transport, regulation of transcription and 312 lipid/protein/carbohydrate catabolism.

313

314 Differential gene expression in the TE part

315 In the TE, 13,322 genes were considered as expressed in OM or ON. Twenty-three were differentially 316 expressed (Additional file 2) with 16 genes being downregulated and 7 being up regulated in ON 317 (Figure 2). Respectively, 14 and 6 out of the down- and upregulated genes were associated to a known 318 protein in human. Moreover, despite the filtering, 2 down-regulated genes in ON (LIM and cysteine 319 rich domains 1, LMCD1; lysophosphatidic acid receptor 4, LPAR4) were only expressed in one embryo 320 and were not considered for further analysis. The remaining 19 genes are presented in Table 4. Down-321 regulated genes in the TE of old nulliparous mares were mainly involved in spindle organization, 322 chromatin remodeling and cellular process, transcription and DNA repair, cell polarity, adhesions, 323 junctions and signaling, extracellular matrix organization, ion homeostasis, glycerol metabolism,

324	glycolysis, immunity, gastrulation and placenta development while up-regulated genes were related
325	to ion binding, cell death, lipid metabolism, protein maturation and membrane organization

326

327 Gene set enrichment analysis in deconvoluted ICM cells

328 After Entrez Gene ID conversion, 12,287 genes were considered expressed in ICM cells. Only one GO

329 Biological Process and one KEGG pathways were disturbed by maternal parity in ICM cells (Additional

file 3 & Table 5). The GO BP "Cytoplasmic microtubule organization" was enriched in the ICM cells from

331 OM embryos (NES = -2.17). The KEGG pathway "Neuroactive ligand receptor interaction" was enriched

in ICM cells from ON embryos (NES = 1.98). Detailed examination indicated that genes involved in this

- enrichment were related to biological regulation, signaling and response.
- 334

335 Gene set enrichment analysis in TE

After Entrez Gene ID conversion, 11,993 genes were considered expressed in TE from ON or OM embryos. All the 2, 2 and 8 perturbed gene sets from GO BP, KEGG and REACTOME, respectively, were enriched in the TE of embryos from OM mares (NES < -1.8; Additional file 4 & Table 5). In the TE, several gene sets were related to extracellular matrix organization and function. Others were involved in NOTCH signaling pathway. The last two pathways were involved in histidine metabolism and cholesterol transport.

342

343 **Discussion**

Maternal parity in mares older than 10 years old did not influence embryo recovery rates. This rate, however, seems lower in old nulliparous compared to old multiparous mares. The effect of mare nulliparity on the fertility in mares older than 10 years old have never been explored.

For 3 mares, a developing embryo was observed at 14 days post ovulation, demonstrating that embryos were not recovered with the flushing. This happened only in multiparous mares and increased 349 occurrence in multiparous mares has not been described to the author's knowledge yet. Uterine size 350 as well as histological and morphological degenerations in uterus could explain this result. After 351 foaling, uterine size decreases quickly during the first week post-partum but the complete involution 352 of both uterine horns seems to end only after twenty to thirty days [62,63] whereas histology of the 353 uterine body endometrium returns to normal by 7 to 10 days post-partum [31]. Ageing and the 354 multiplication of foalings, with repetition of uterine extensions and involutions may affect the uterus's 355 ability to involute. Indeed, clinically the authors have observed that a larger volume of fluid is required 356 to flush the uterus of multiparous mares compared to young mares. Histologically, however, parity is 357 correlated with the presence of elastosis in the myometrial vessels [37]. These alterations may be 358 associated to alterations of uterine contractility and decrease fluid clearance in multiparous mares, 359 both factors that could affect embryo recovery.

360 In both ICM and TE, only a few genes were affected by maternal parity. Some downregulated genes in 361 embryos collected from ON mares were involved in common functions between ICM and TE such as 362 regulation of transcription, cell cycle and development, cell organization and immunity. Up-regulated 363 genes in the ICM of ON embryos were related to ion transport, regulation of transcription and 364 catabolism of lipids, proteins and carbohydrates. In the TE, downregulated genes in ON embryos were 365 related to glycerol metabolism, glycolysis and directly involved in gastrulation and placental formation 366 while upregulated genes were related to cell death, lipid metabolism, protein maturation, membrane 367 organization and as in ICM, ion binding.

In the GSEA analysis, only two gene sets were perturbed in the ICM in relation to maternal parity in old mares with one related to microtubule organization being enriched in ON and the other one being enriched in OM embryos and related membrane receptors. No gene set was enriched in the TE of ON mares but enriched pathways in OM embryos were mainly related to extracellular matrix organization and cell differentiation, mainly related to NOCTH signaling pathway.

373 Of particular interest, the gene SLX4 Structure-Specific Endonuclease Subunit (*SLX4*), the regulatory 374 subunit of structure-specific endonucleases that are required for repair of DNA lesions, is down375 regulated in the ICM of embryos from ON mares. In TE, moreover, part of the post-replicative DNA 376 mismatch repair system, MutS Homolog 3 (MSH3) is downregulated in embryos from ON mares. These 377 results indicate that DNA repair systems in both ICM and TE are affected by maternal parity. 378 Interestingly, the gene Gamma-aminobutyric acid type A receptor subunit rho1, GABRR1, was not 379 expressed in embryos collected from multiparous mares while in nulliparous mares, 4/5 embryos 380 expressed this gene in the ICM. GABRR1 encodes for a Cl⁻ channel receptor involved in the gamma-381 aminobutyric acid (GABA) pathway. Work on mouse embryonic and peripheral neural crest stem cells 382 have shown that GABA receptors negatively affect preimplantation embryonic growth by negatively 383 controlling cell proliferation, being involved in DNA damage checkpoint and by increasing cellular 384 arrest in the S phase [64]. Alterations in the embryo environment because of mare parity could 385 therefore modify DNA lesion repair and therefore, cell proliferation, suggesting that embryo growth is 386 reduced in nulliparous mares.

Several gene sets related to NOTCH signaling pathways were also enriched in the TE of OM embryos and Notch Receptor 1 (*NOTCH1*) always contributed to those enrichments. NOTCH signaling pathway is essential for proper development, with *NOTCH1* being required for cell proliferation in early bovine embryos [65]. The enrichment of this pathway in embryos from multiparous mares therefore suggests that cell proliferation is slowed down in embryos from nulliparous mares

392 The expression of SET domain containing 5 (SETD5) signal transducer and activator of transcription 5A 393 (STAT5A), PCF11 cleavage and polyadenylation factor subunit (PCF11), ribosomal protein L36 (RPL36), 394 exosome component 7 (EXOSC7), H2A clustered histone 20 (H2AC20) were downregulated in the ICM 395 and H4 clustered histone 3 (H4C3) were downregulated in the TE of embryos collected from nulliparous 396 mares. These genes are all involved in transcription and/or translation, suggesting that the expression 397 of genes in embryos from nulliparous mares was altered. Of particular interest, SETD5 et STAT5A are known to be, respectively, key regulators of methylation and signaling via cytokines, both gene 398 expressions being essential for embryo development [77,78]. 399

In the TE, the gene encoding for eomesodermin (*EOMES*) is downregulated in embryos collected from ON mares. This gene controls the formation of germ cell layers and is involved in the differentiation of the trophoblast in the mouse [66] while in human, cattle and pigs, *EOMES* is not expressed in the preimplantation embryos [67–69]. In horses, *EOMES* is suggested to also be a marker of induced trophoblast cells but its role has never been explored [70]. If its role is similar as in mouse, this downregulation could lead to a reduced differentiation of cells in the trophoblast, that could impair its principal function, i.e., the regulation of exchanges with the maternal environment.

407 As the external part of the embryo is exclusively composed of trophoblast in the mare at the studied 408 developmental stage, poorer maternal-embryo exchanges through the trophoblast in nulliparous 409 mares could explain the defects observed in both compartments. Indeed, the reduced gene expression 410 of TRIO and F-Actin Binding Protein (TRIOBP) in the ICM as well as actin beta (ACTB) and microtubule 411 associated protein 1B (MAP1B) in the TE of equine embryos seems to fit the hypothesis that molecule 412 transfer is altered in embryos from ON mares. In polarized epithelial cells, such as the trophoblast, 413 cytoskeleton is essential for the communication with the extracellular environment (for review [71]). 414 The protein encoded by ACTB is a direct component of the cytoskeleton and the one encoded by 415 MAP1B is a molecule responsible for the stabilization of microtubules [72]. TRIOBP regulates actin 416 cytoskeletal organization and the formation of a Tara and TRIO complex coordinates actin remodeling 417 which is essential for exchanges [73]. In addition, the extracellular matrix (ECM) is very important for 418 embryo development and embryo-maternal exchanges (for a review [74]). ECM relative gene sets 419 appeared to be altered by maternal parity in old mares with several gene sets enriched in the TE of 420 embryos from ON mares. The expression of the ADAM metallopeptidase domain 19 gene (ADAM19), 421 moreover, was reduced in the TE of ON embryos. This gene encodes for a transmembrane glycoprotein 422 that is essential for tight junction formation. Tight junctions formation and integrity are essential for 423 blastocyst development in mouse and pigs [75]. The reduction of the expression of ADAM19 in the TE 424 of embryos from old nulliparous mares, could therefore support the hypothesis that embryo integrity 425 is altered, leading to alteration of ion and nutrient exchanges in both TE and ICM. Alterations of 426 cytoskeleton, ECM and integrity of the TE in ON embryos, probably related to adaptation to the embryo
427 environment, could affect embryo-maternal exchanges and consequently embryonic development.

428 The lipid metabolism is particularly important for mammalian embryo development as it is an 429 important source of energy for growth (for review [79]). In horses, the embryo is particularly reliant 430 on its environment as it develops free inside the mare's uterus until around 35 days post ovulation, 431 when it finally starts to implant. Several genes related to lipid metabolism and transport were affected 432 in the ICM by maternal parity. Among them, carnitine palmitoyltransferase 2 (CPT2) is downregulated 433 in the ICM of ON embryos. The protein encoded by this gene catalyzes the oxidation of fatty acids in 434 the mitochondria. In mouse embryos, CTP2 transcription is essential and could be used as a marker for 435 future implantation [80]. Therefore, the reduced expression of this gene could indicate that embryo 436 development is reduced because of reduced lipid oxidation in old nulliparous mares. This could be the 437 result or at least related to reduced lipolysis via lipase or a defect in the transport of lipids. Both 438 functions are altered in the ICM of ON embryos. Indeed, lipase E (LIPE), also known as hormone-439 sensitive lipase, and apolipoprotein B (APOB), involved in the transport of lipids, are downregulated in 440 the ICM of ON embryos. APOB has been identified as very important for equine embryo development 441 as its expression is increased by a factor of 200 between day 8 and day 14 [81]. Furthermore, the 442 production of the protein apolipoprotein B by the endoderm is required for the development of 443 preimplantation mouse embryos [82]. In the equine species, APOB expression is dysregulated in Day 8 444 embryos when mares do not produce enough progesterone [83]. It has been observed that the risk of 445 embryo mortality is increased in mares producing less than 4 ng/ml of progesterone through the post 446 ovulation period [84]. Progesterone, indeed, is mandatory for embryo development by modulating its 447 environment to ensure pregnancy [85]. Reduced expression of APOB suggests that the expression of 448 this gene is particularly dependent on the embryo environment. Here, except one, close to the cut-off 449 value, all mares produced sufficient progesterone for normal gestation at 8 days post ovulation and 450 there was no difference in progesterone concentration according to mare parity. It could be

- 451 hypothesized, however, that the environment in the uterus of old nulliparous mares could lead to the452 difference of expression of this gene, independently of progesterone production.
- 453

454 **Conclusion**

455 Mare's parity, especially in mares older than 10 years old, has never been considered for the study of 456 embryos. Here, however, it has been shown that mare's parity in old mares affects the expression of 457 genes in ICM and TE of blastocysts. Although only the expression of a few genes is altered by mare's 458 parity, some of these genes are particularly important for embryo growth and development. Genes 459 related to nutrient exchanges and responses to environment signaling in both ICM and TE are 460 particularly affected by the nulliparity of mares, suggesting that the developing environment from 461 these mares are not optimal for embryonic growth. It is not possible to conclude, however, if 462 differences in oocyte quality also play a role in those observations. More work on the oocyte, oviductal 463 and uterine environment are needed to elucidate the exact mechanism leading to the present 464 observations in Day 8 embryos.

465 In the present experiment, the destruction of the embryos made it impossible to predict their 466 individual chance of implantation. The observed alterations suggest that implantation defects may be 467 present in the embryos of old nulliparous mares. It is often assumed that nulliparous mares have a 468 better uterine environment, more favorable to the development and implantation of an embryo than 469 multiparous mares. In this study, however, being nulliparous and old does not seem to be the perfect 470 match for embryonic development. If embryos succeed to implant, the apparent lower quality of 471 embryos in nulliparous mares may accentuate differences in placentation and development observed in nulliparous mares, exacerbating the observed phenotype of smaller and lighter foals at birth. 472

474 Acknowledgments

484	Conflict of interest
483	
482	provided the coordinates for the XIST gene.
481	Christophe Klopp for their advice on RNA-seq de novo analysis. Many thanks to Pablo Ross who kindly
480	Phenotyping-Endocrinology laboratory for the hormonal assays. Many thanks to Matthias Zytnicki and
479	de Bioinformatique (IFB) (ANR-11-INBS-0013). The authors thank A.L. Lainé and the staff of the
478	expertise. The bioinformatics analyses were performed thanks to Core Cluster of the Institut Français
477	We acknowledge the high-throughput sequencing facility of I2BC for its sequencing and bioinformatics
476	experimental farm (Jumenterie du Haras du Pin, Exmes, France) for care and management of animals.
475	The authors are grateful to the staff of the Institut Français du Cheval et de l'Equitation (IFCE)

485 The authors declare that they have no conflict of interests.

486

487 Author contributions

PCP obtained the funding. PCP and VD conceived the project. FDG, LB, VD and PCP supervised the study. ED, CG, AM, CA, ND, NP, FDG, VD and PCP adapted the methodology for the project. ED, CG, AM, CA and YJ performed the experiments. AM, CG, CA, ND, NP, MD, LB and FDG provided the resources. ED, LJ, YJ and RL performed data curation. ED and LJ analyzed the data. ED wrote the original draft. All authors read, revised, and approved the submitted manuscript.

493

494

496 List of abbreviations

- 497 DEG: differential expressed genes
- 498 DeMixT_ICM_cells: deconvoluted gene expression in ICM cells
- 499 DeMixT_TE_cells: deconvoluted gene expression in TE cells
- 500 ECM: Extracellular matrix
- 501 ERR: embryo collection rate
- 502 FDR: false discovery rate
- 503 GO BP: Gene Ontology biological process
- 504 GO: Gene Ontology
- 505 GSEA: gene set enrichment analyses
- 506 ICM: inner cell mass
- 507 ICMandTE: inner cell mass enriched hemi-embryo
- 508 ICSI: intracytoplasmic sperm injection
- 509 KEGG: Kyoto Encyclopedia of Genes and Genomes
- 510 Log2FC: log2 fold change
- 511 NES: normalized enrichment score
- 512 OM: old multiparous mares
- 513 ON: old nulliparous mares
- 514 TE: trophoblast
- 515 TE_part: pure trophoblast hemi-embryo

516 XIST: X inactive Specific Transcript

518 **References**

- Allen WR, Brown L, Wright M, Wilsher S. Reproductive efficiency of Flatrace and National Hunt
 Thoroughbred mares and stallions in England. Equine Veterinary Journal 2007;39:438–45.
 https://doi.org/10.2746/042516407X1737581.
- 522 [2] Katila T, Reilas T, Nivola K, Peltonen T, Virtala A-M. A 15-year survey of reproductive efficiency
 523 of Standardbred and Finnhorse trotters in Finland descriptive results. Acta Veterinaria
 524 Scandinavica 2010:11.
- 525 [3] Physick-Sheard PW. Demographic analysis of the Canadian Standardbred broodmare herd.
 526 Preventive Veterinary Medicine 1995;24:285–99. https://doi.org/10.1016/0167527 5877(95)00492-F.
- 528 [4] Derisoud E, Auclair-Ronzaud J, Palmer E, Robles M, Chavatte-Palmer P. Female age and parity in 529 horses: how and why does it matter? Reproduction, Fertility and Development In press.
- [5] Rizzo M, Stout TAE, Cristarella S, Quartuccio M, Kops GJPL, De Ruijter-Villani M. Compromised
 MPS1 Activity Induces Multipolar Spindle Formation in Oocytes From Aged Mares: Establishing
 the Horse as a Natural Animal Model to Study Age-Induced Oocyte Meiotic Spindle Instability.
 Front Cell Dev Biol 2021;9:657366. https://doi.org/10.3389/fcell.2021.657366.
- 534[6]Rizzo M, du Preez N, Ducheyne KD, Deelen C, Beitsma MM, Stout TAE, et al. The horse as a
natural model to study reproductive aging-induced aneuploidy and weakened centromeric
cohesion in oocytes. Aging 2020. https://doi.org/10.18632/aging.104159.
- 537 [7] Rizzo M, Ducheyne KD, Deelen C, Beitsma M, Cristarella S, Quartuccio M, et al. Advanced mare
 538 age impairs the ability of in vitro-matured oocytes to correctly align chromosomes on the
 539 metaphase plate. Equine Vet J 2019;51:252–7. https://doi.org/10.1111/evj.12995.
- [8] Campos-Chillon F, Farmerie TA, Bouma GJ, Clay CM, Carnevale EM. Effects of aging on gene expression and mitochondrial DNA in the equine oocyte and follicle cells. Reprod Fertil Dev 2015;27:925. https://doi.org/10.1071/RD14472.
- 543 [9] Cox L, Vanderwall DK, Parkinson KC, Sweat A, Isom SC. Expression profiles of select genes in
 544 cumulus–oocyte complexes from young and aged mares. Reprod Fertil Dev 2015;27:914.
 545 https://doi.org/10.1071/RD14446.
- [10] Rambags BPB, van Boxtel DCJ, Tharasanit T, Lenstra JA, Colenbrander B, Stout TAE. Advancing
 maternal age predisposes to mitochondrial damage and loss during maturation of equine
 oocytes in vitro. Theriogenology 2014;81:959–65.
- 549 https://doi.org/10.1016/j.theriogenology.2014.01.020.
- [11] Catandi GD, Obeidat YM, Broeckling CD, Chen TW, Chicco AJ, Carnevale EM. Equine maternal
 aging affects oocyte lipid content, metabolic function and developmental potential.
 Reproduction 2021;161:399–409. https://doi.org/10.1530/REP-20-0494.
- 553 [12] Hendriks WK, Colleoni S, Galli C, Paris DBBP, Colenbrander B, Roelen BAJ, et al. Maternal age
 554 and in vitro culture affect mitochondrial number and function in equine oocytes and embryos.
 555 Reprod Fertil Dev 2015;27:957. https://doi.org/10.1071/RD14450.
- [13] Woods GL, Baker CB, Hillman RB, Schlafer DH. Recent studies relating to early embryonic death
 in the mare. Equine Veterinary Journal 1985;17:104–7. https://doi.org/10.1111/j.20423306.1985.tb04609.x.
- [14] Carnevale EM, Griffin PG, Ginther OJ. Age-associated subfertility before entry of embryos into
 the uterus in mares. Equine Veterinary Journal 1993;25:31–5. https://doi.org/10.1111/j.2042 3306.1993.tb04820.x.
- [15] Brinsko SP, Ball BA, Miller PG, Thomas PGA, Ellington JE. In vitro development of day 2 embryos
 obtained from young, fertile mares and aged, subfertile mares. Reproduction 1994;102:371–8.
 https://doi.org/10.1530/jrf.0.1020371.
- [16] Willmann C, Schuler G, Hoffmann B, Parvizi N, Aurich C. Effects of age and altrenogest
 treatment on conceptus development and secretion of LH, progesterone and eCG in early-

567 pregnant mares. Theriogenology 2011;75:421-8. https://doi.org/10.1016/j.theriogenology.2010.05.009. 568 [17] Panzani D, Rota A, Marmorini P, Vannozzi I, Camillo F. Retrospective study of factors affecting 569 570 multiple ovulations, embryo recovery, quality, and diameter in a commercial equine embryo transfer program. Theriogenology 2014;82:807–14. 571 572 https://doi.org/10.1016/j.theriogenology.2014.06.020. [18] Camargo Ferreira J, Linhares Boakari Y, Sousa Rocha N, Saules Ignácio F, Barbosa da Costa G, de 573 Meira C. Luteal vascularity and embryo dynamics in mares during early gestation: Effect of age 574 575 and endometrial degeneration. Reprod Dom Anim 2019;54:571-9. 576 https://doi.org/10.1111/rda.13396. 577 [19] Carnevale EM, Ginther OJ. Relationships of age to uterine function and reproductive efficiency 578 in mares. Theriogenology 1992;37:1101–15. https://doi.org/10.1016/0093-691X(92)90108-4. 579 [20] Carnevale EM, Ginther OJ. Defective Oocytes as a Cause of Subfertility in Old Mares. Biology of 580 Reproduction 1995;52:209-14. 581 https://doi.org/10.1093/biolreprod/52.monograph_series1.209. [21] Morel MCGD, Newcombe JR, Swindlehurst JC. The effect of age on multiple ovulation rates, 582 583 multiple pregnancy rates and embryonic vesicle diameter in the mare. Theriogenology 584 2005;63:2482–93. https://doi.org/10.1016/j.theriogenology.2004.09.058. [22] Frank BL, Doddman CD, Stokes JE, Carnevale EM. Association of equine oocyte and cleavage 585 stage embryo morphology with maternal age and pregnancy after intracytoplasmic sperm 586 587 injection. Reprod Fertil Dev 2019;31:1812. https://doi.org/10.1071/RD19250. 588 [23] Derisoud E, Jouneau L, Dubois C, Archilla C, Jaszczyszyn Y, Legendre R, et al. Maternal age 589 affects equine Day 8 embryo gene expression both in trophoblast and inner cell mass. BioRxiv 590 2021:2021.04.07.438786. https://doi.org/10.1101/2021.04.07.438786. 591 [24] Aguilar JJ, Woods GL, Miragaya MH, Olsen LM. Living fibroblast cells in the oviductal masses of 592 mares. Equine Veterinary Journal 1997;S25:103-8. https://doi.org/10.1111/j.2042-593 3306.1997.tb05112.x. [25] Adams GP, Kastelic JP, Bergfelt DR, Ginther OJ. Effect of uterine inflammation and 594 595 ultrasonically-detected uterine pathology on fertility in the mare. J Reprod Fertil Suppl 596 1987;35:445-54. 597 [26] Chevalier-Clément F. Pregnancy loss in the mare. Animal Reproduction Science 1989;20:231-598 44. https://doi.org/10.1016/0378-4320(89)90088-2. 599 [27] Bracher V, Mathias S, Allen WR. Videoendoscopic evaluation of the mare's uterus: II. Findings in 600 subfertile mares. Equine Veterinary Journal 1992;24:279-84. https://doi.org/10.1111/j.2042-601 3306.1992.tb02835.x. [28] Eilts BE, Scholl DT, Paccamonti DL, Causey R, Klimczak JC, Corley JR. Prevalence of Endometrial 602 603 Cysts and Their Effect on Fertility. Biology of Reproduction 1995;52:527–32. 604 https://doi.org/10.1093/biolreprod/52.monograph series1.527. 605 [29] Tannus RJ, Thun R. Influence of Endometrial Cysts on Conception Rate of Mares. Journal of 606 Veterinary Medicine Series A 1995;42:275-83. https://doi.org/10.1111/j.1439-607 0442.1995.tb00378.x. 608 [30] Ferreira JC, Gastal EL, Ginther OJ. Uterine blood flow and perfusion in mares with uterine cysts: 609 effect of the size of the cystic area and age. Reproduction 2008;135:541–50. https://doi.org/10.1530/REP-07-0447. 610 611 [31] Kenney RM. Cyclic and pathologic changes of the mare endometrium as detected by biopsy, 612 with a note on early embryonic death. Journal of the American Veterinary Medical Association 613 1978;172:241-62. 614 [32] Doig PA, McKNIGHT JD, Miller RB. The Use of Endometrial Biopsy in the Infertile Mare. Can Vet J 615 1981;22:72-6. [33] Siemieniuch MJ, Gajos K, Kozdrowski R, Nowak M. Advanced age in mares affects endometrial 616 617 secretion of arachidonic acid metabolites during equine subclinical endometritis. 618 Theriogenology 2017;103:191–6. https://doi.org/10.1016/j.theriogenology.2017.07.043.

619 [34] Barbacini S, Marchi V, Zavaglia G. Equine frozen semen: results obtained in Italy during the 620 1994-1997 period. Equine Veterinary Education 1999;11:109–12. https://doi.org/10.1111/j.2042-3292.1999.tb00930.x. 621 622 [35] Woodward EM, Christoffersen M, Campos J, Squires EL, Troedsson MHT. Susceptibility to 623 persistent breeding-induced endometritis in the mare: Relationship to endometrial biopsy 624 score and age, and variations between seasons. Theriogenology 2012;78:495–501. 625 https://doi.org/10.1016/j.theriogenology.2012.02.028. [36] LeBlanc MM, Neuwirth L, Jones L, Cage C, Mauragis D. Differences in uterine position of 626 627 reproductively normal mares and those with delayed uterine clearance detected by 628 scintigraphy. Theriogenology 1998;50:49–54. https://doi.org/10.1016/S0093-691X(98)00112-5. 629 [37] Esteller-Vico A, Liu IK, Couto S. Uterine vascular degeneration is present throughout the uterine 630 wall of multiparous mares. Colinearity between elastosis, endometrial grade, age and parity. Theriogenology 2012;78:1078–84. https://doi.org/10.1016/j.theriogenology.2012.04.014. 631 632 [38] LeBlanc MM, Neuwirth L, Asbury AC, Tran T, Mauragis D, Klapstein E. Scintigraphic 633 measurement of uterine clearance in normal mares and mares with recurrent endometritis. 634 Equine Veterinary Journal 1994;26:109–13. https://doi.org/10.1111/j.2042-635 3306.1994.tb04346.x. [39] Duchamp G, Bour B, Combarnous Y, Palmer E. Alternative solutions to hCG induction of 636 637 ovulation in the mare. J Reprod Fertil Suppl 1987;35:221-8. [40] Bucca S, Carli A. Efficacy of human chorionic gonadotropin to induce ovulation in the mare, 638 639 when associated with a single dose of dexamethasone administered at breeding time: Efficacy 640 of human chorionic gonadotropin to induce ovulation when associated with dexamethasone. 641 Equine Veterinary Journal 2011;43:32–4. https://doi.org/10.1111/j.2042-3306.2011.00488.x. 642 [41] Canepa S, Laine A-L, Bluteau A, Fagu C, Flon C, Monniaux D. Validation d'une méthode 643 immunoenzymatique pour le dosage de la progestérone dans le plasma des ovins et des bovins. 644 Cahier des Techniques de l'INRA 2008;64:19. 645 [42] Panzani D, Di Vita M, Lainé A-L, Guillaume D, Rota A, Tesi M, et al. Corpus Luteum Vascularization and Progesterone Production in Autumn and Winter Cycles of the Mare: 646 647 Relationship Between Ultrasonographic Characteristics of Corpora Lutea and Plasma Progesterone Concentration in the Last Cycles Before Anestrus. Journal of Equine Veterinary 648 649 Science 2017;56:35–9. https://doi.org/10.1016/j.jevs.2017.05.001. 650 [43] Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet Journal 2011;17:10-2. 651 652 [44] Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal 653 RNA-seq aligner. Bioinformatics 2013;29:15–21. 654 https://doi.org/10.1093/bioinformatics/bts635. 655 [45] Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics 2014;30:923-30. 656 657 https://doi.org/10.1093/bioinformatics/btt656. 658 [46] R Core Team. A Language and Environment for Statistical Computing. Vienna, Austria: R foundation for Statistical Computing; 2020. 659 660 [47] Rstudio Team. RStudio: Integrated Development for R. RStudio. Boston, USA: PBC; 2020. [48] Pinheiro J, Bates D, Debroy S, Sarkar D, R Core Team. nlme: Linear and Nonlinear Mixed Effects 661 662 Models. 2020. [49] Giraudoux P. pgirmess: Spatial Analysis and Data Mining for Field Ecologists. 2018. 663 664 [50] Cao S, Wang JR, Ji S, Yang P, Chen J, Shen JP, et al. Differing total mRNA expression shapes the molecular and clinical phenotype of cancer. BioRxiv 2020:57. 665 https://doi.org/10.1101/2020.09.30.306795. 666 [51] Wang Z, Cao S, Morris JS, Ahn J, Liu R, Tyekucheva S, et al. Transcriptome Deconvolution of 667 668 Heterogeneous Tumor Samples with Immune Infiltration. IScience 2018;9:451-60. 669 https://doi.org/10.1016/j.isci.2018.10.028.

[52] Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq 670 data with DESeq2. Genome Biol 2014;15:550. https://doi.org/10.1186/s13059-014-0550-8. 671 [53] Iqbal K, Chitwood JL, Meyers-Brown GA, Roser JF, Ross PJ. RNA-Seq Transcriptome Profiling of 672 673 Equine Inner Cell Mass and Trophectoderm. Biology of Reproduction 2014;90. https://doi.org/10.1095/biolreprod.113.113928. 674 675 [54] Gómez-Rubio V. ggplot2 - Elegant Graphics for Data Analysis (2nd Edition). J Stat Soft 2017;77. https://doi.org/10.18637/jss.v077.b02. 676 [55] Kassambara A, Mundt F. factoextra: Extract and Visualize the Results of Multivariate Data 677 678 Analyses. 2020. 679 [56] Kolberg L, Raudvere U. gprofiler2: Interface to the "g:Profiler" Toolset. 2020. 680 [57] Yates AD, Achutan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, et al. Ensembl 2020 n.d. 681 [58] Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set 682 enrichment analysis: A knowledge-based approach for interpreting genome-wide expression 683 profiles. Proceedings of the National Academy of Sciences 2005;102:15545-50. 684 https://doi.org/10.1073/pnas.0506580102. [59] Mootha VK, Lindgren CM, Eriksson K-F, Subramanian A, Sihag S, Lehar J, et al. PGC-1α-685 686 responsive genes involved in oxidative phosphorylation are coordinately downregulated in 687 human diabetes. Nat Genet 2003;34:267–73. https://doi.org/10.1038/ng1180. [60] Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular 688 Signatures Database Hallmark Gene Set Collection. Cell Systems 2015;1:417–25. 689 690 https://doi.org/10.1016/j.cels.2015.12.004. 691 [61] McKinnon AO, Squires EL. Morphologic assessment of the equine embryo. Journal of the 692 American Veterinary Medical Association 1988;192:401-6. 693 [62] Griffin PG, Ginther OJ. Uterine morphology and function in postpartum mares. Journal of 694 Equine Veterinary Science 1991;11:330–9. https://doi.org/10.1016/S0737-0806(06)81262-1. 695 [63] Lemes KM, Silva LA, Alonso MA, Celeghini ECC, Pugliesi G, Carvalho HF, et al. Uterine Vascular 696 Perfusion and Involution During the Postpartum Period in Mares. Journal of Equine Veterinary 697 Science 2017;51:61–9. https://doi.org/10.1016/j.jevs.2016.10.010. 698 [64] Andäng M, Hjerling-Leffler J, Moliner A, Lundgren TK, Castelo-Branco G, Nanou E, et al. Histone H2AX-dependent GABAA receptor regulation of stem cell proliferation. Nature 2008;451:460–4. 699 700 https://doi.org/10.1038/nature06488. 701 [65] Li S, Shi Y, Dang Y, Luo L, Hu B, Wang S, et al. NOTCH signaling pathway is required for bovine 702 early embryonic development. Biology of Reproduction 2021;105:332-44. 703 https://doi.org/10.1093/biolre/ioab056. 704 [66] Russ AP, Wattler S, Colledge WH, Aparicio SAJR, Carlton MBL, Pearce JJ, et al. Eomesodermin is 705 required for mouse trophoblast development and mesoderm formation. Nature 2000;404:95-706 9. https://doi.org/10.1038/35003601. 707 [67] Boroviak T, Loos R, Lombard P, Okahara J, Behr R, Sasaki E, et al. Lineage-Specific Profiling 708 Delineates the Emergence and Progression of Naive Pluripotency in Mammalian 709 Embryogenesis. Developmental Cell 2015;35:366-82. 710 https://doi.org/10.1016/j.devcel.2015.10.011. [68] Valdez Magaña G, Rodríguez A, Zhang H, Webb R, Alberio R. Paracrine effects of embryo-711 derived FGF4 and BMP4 during pig trophoblast elongation. Developmental Biology 712 713 2014;387:15-27. https://doi.org/10.1016/j.ydbio.2014.01.008. [69] Degrelle SA, Campion E, Cabau C, Piumi F, Reinaud P, Richard C, et al. Molecular evidence for a 714 715 critical period in mural trophoblast development in bovine blastocysts. Developmental Biology 716 2005;288:448-60. https://doi.org/10.1016/j.ydbio.2005.09.043. [70] Reinholt BM, Bradley JS, Jacobs RD, Ealy AD, Johnson SE. Tissue organization alters gene 717 718 expression in equine induced trophectoderm cells. General and Comparative Endocrinology 719 2017;247:174-82. https://doi.org/10.1016/j.ygcen.2017.01.030.

- [71] Apodaca G. Endocytic Traffic in Polarized Epithelial Cells: Role of the Actin and Microtubule
 Cytoskeleton: Role of Cytoskeleton in Endocytic Traffic 2001;2:149–59.
 https://doi.org/10.1034/j.1600-0854.2001.020301.x.
- Tögel M, Wiche G, Propst F. Novel Features of the Light Chain of Microtubule-associated
 Protein MAP1B: Microtubule Stabilization, Self Interaction, Actin Filament Binding, and
 Regulation by the Heavy Chain. Journal of Cell Biology 1998;143:695–707.
 https://doi.org/10.1083/jcb.143.3.695.
- [73] Seipel K, O'Brien S, Iannotti E, Medley Q, Streuli M. Tara, a novel F-actin binding protein,
 associates with the Trio guanine nucleotide exchange factor and regulates actin cytoskeletal
 organization. Journal of Cell Science 2001;114:389–99.
- 730 [74] Zagris N. Extracellular matrix in development of the early embryo. Micron 2001;32:427–38.
 731 https://doi.org/10.1016/S0968-4328(00)00011-1.
- [75] Jeong Y, Choi I. Tight Junction Assembly Ensures Maintenance of Pregnancy during
 Embryogenesis in a Mouse Model. Journal of Animal Reproduction and Biotechnology
 2019;34:318–21. https://doi.org/10.12750/JARB.34.4.318.
- [76] Wang H-X, Zhao Y-G, Wang H-M, Yang Q, Lin H-Y, Sang Q-XA, et al. Expression of adamalysin
 19/ADAM19 in the endometrium and placenta of rhesus monkey (Macaca mulatta) during early
 pregnancy. Molecular Human Reproduction 2005;11:429–35.
- 738 https://doi.org/10.1093/molehr/gah183.
- 739 [77] Osipovich AB, Gangula R, Vianna PG, Magnuson MA. Setd5 is essential for mammalian
 740 development and the co-transcriptional regulation of histone acetylation. Development
 741 2016;143:4595–607. https://doi.org/10.1242/dev.141465.
- [78] Nakasato M, Shirakura Y, Ooga M, Iwatsuki M, Ito M, Kageyama S, et al. Involvement of the
 STAT5 Signaling Pathway in the Regulation of Mouse Preimplantation Development. Biology of
 Reproduction 2006;75:508–17. https://doi.org/10.1095/biolreprod.105.047860.
- [79] Bradley J, Swann K. Mitochondria and lipid metabolism in mammalian oocytes and early
 embryos. Int J Dev Biol 2019;63:93–103. https://doi.org/10.1387/ijdb.180355ks.
- [80] Gentile L, Monti M, Sebastiano V, Merico V, Nicolai R, Calvani M, et al. Single-cell quantitative
 RT-PCR analysis of *Cpt1b* and *Cpt2* gene expression in mouse antral oocytes and in
 preimplantation embryos. Cytogenet Genome Res 2004;105:215–21.
 https://doi.org/10.1159/000078191.
- [81] Klein C, Troedsson MHT. Transcriptional Profiling of Equine Conceptuses Reveals New Aspects
 of Embryo-Maternal Communication in the Horse1. Biology of Reproduction 2011;84:872–85.
 https://doi.org/10.1095/biolreprod.110.088732.
- [82] Farese RV, Cases S, Ruland SL, Kayden HJ, Wong JS, Young SG, et al. A novel function for
 apolipoprotein B: lipoprotein synthesis in the yolk sac is critical for maternal-fetal lipid
 transport in mice. Journal of Lipid Research 1996;37:347–60. https://doi.org/10.1016/S00222275(20)37621-5.
- [83] Leisinger CA, Klein C, Markle ML, Premanandan C, Sones JL, Pinto CRF, et al. Altered gene
 expression in embryos and endometrium collected on day 8 of induced aluteal cycles in mares.
 Theriogenology 2019;128:81–90. https://doi.org/10.1016/j.theriogenology.2019.01.028.
- [84] Ginther OJ. Embryonic loss in mares: Incidence, time of occurrence, and hormonal involvement.
 Theriogenology 1985;23:77–89. https://doi.org/10.1016/0093-691X(85)90074-3.
- 763 [85] Allen W. Luteal Deficiency and Embryo Mortality in the Mare. Reprod Domest Anim
- 764 2001;36:121–31. https://doi.org/10.1046/j.1439-0531.2001.d01-43.x.
- 765

766 Figure legends

767 Figure 1: Gene expression in ICM and TE before and after deconvolution using DeMixT

A) Venn diagrams of the differential gene expression in ICMandTE vs TE_part (before deconvolution),
DeMixT_ICM_cells vs DeMixT_TE_cells (after deconvolution) and DeMixT_ICM_cells vs TE_part (gene
expression of ICM after deconvolution vs gene expression in TE_part without deconvolution); B)
Principal Component Analysis of gene expression of DeMixT_ICM_cells, DeMixT_TE_cells, ICMandTE
and TE part datasets.
Deconvolution was used to isolate gene expression of ICM and TE cells in ICMandTE hemi-embryos.
ICMandTE: inner cell mass + trophoblast; TE_part: pure trophoblast. Here trophoblast represents
trophectoderm + endoderm.
Figure 2: Analysis of differentially expressed genes (DEG) in embryos according to maternal parity
A) representation of down- (blue) and upregulated (red) DEG in ICM (from DeMixT_ICM_cells data
obtained after deconvolution of ICMandTE using DeMixT R package [50,51]) and TE (from TE_part
dataset) of embryos from ON vs OM.
DEG: Differentially Expressed Genes (FDR < 0.05); TE: Trophoblast; ICM: Inner Cell Mass; ON: Old
nulliparous mares; OM: Old multiparous mares

- 784 Additional file 1:
- 785 Supp1.xlsx
- 786 Differential gene analysis using DeSeq2 in DeMixT_ICM_cells of equine embryo at Day 8 post-ovulation
- 787 according to mares parity
- 788 Equine ensemble ID, orthologue human Ensembl ID, Orthologue human Entrez Gene ID, gene
- description, normalized counts for each embryo and parameters obtained after Deseg2 analysis
- 790 (log2FoldChange, pvalue and padj (after FDR correction)) of genes expressed in ICM (after gene
- 791 expression deconvolution of ICMandTE using DeMixT) of ON and OM embryos
- 792 ICM: Inner cell mass; ON: Old nulliparous mares; OM: old multiparous mares

793

- 794 Additional file 2:
- 795 Supp2.xlsx
- Differential gene analysis using DeSeq2 in TE_part of equine embryo at Day 8 post-ovulation according
 to mares parity

. ,

Equine ensemble ID, orthologue human Ensembl ID, Orthologue human Entrez Gene ID, gene description, normalized counts for each embryo and parameters obtained after Deseq2 analysis (log2FoldChange, pvalue and padj (after FDR correction)) of genes expressed in TE_part of ON and OM embryos

802 ICM: Inner cell mass; ON: Old nulliparous mares; OM: old multiparous mares

803

804 Additional file 3:

805 Supp3.xlsx

806 Gene set enrichment analysis results on gene expression of DeMixT_ICM_cell of embryos from old

807 nulliparous and multiparous mares

Gene Set Enrichment Analysis results (database, pathway name, size, enrichment score without and
with normalization, p-value and FDR corrected q-value) for GO biological process, KEGG and
REACTOME databases on genes expressed in ICM (after gene expression deconvolution of ICMandTE
using DeMixT).

- 812 ICM: Inner cell mass
- 813
- 814 Additional file 4:
- 815 Supp4.xlsx
- 816 Gene set enrichment analysis results on gene expression of TE_part of embryos from old nulliparous
- 817 and multiparous mares
- 818 Gene Set Enrichment Analysis results (database, pathway name, size, enrichment score without and
- 819 with normalization, p-value and FDR corrected q-value) for GO biological process, KEGG and
- 820 REACTOME databases on genes expressed in TE_part
- 821 TE: trophoblast

822 Tables

	Nulliparo	ous (ON)	Multipar	ous (OM)
Characteristics	All mares (n = 11)	Embryo donors (n = 5)	All mares (n = 14)	Embryo donors (n = 6)
Breed	SF n = 9; ST n = 1; SB n = 1	SF n = 4; ST n = 1	SF n = 7; AA n = 4; ST n = 2	SF n = 5; AA n = 1
Age (in years)	12.7 ± 2.0	13.6 ± 2.2	13.4 ± 2.1	13.5 ± 2.4
Parity (number of foalings)	0	0	1.7 ± 1.3	1.8 ± 1.6
Weight (in kg)	663.3 ± 49.1	644.8 ± 53.3	641.0 ± 76.1	658.0 ± 81.7
Withers' height (in cm)	168.2 ± 5.8	165.8 ± 7.6	163.6 ± 4.1	164.0 ± 5.1
Progesterone concentration (<i>in ng/ml</i>)	-	12.7 ± 5.8	-	14.5 ± 7.3

823 Table 1: Mares' characteristics at embryo collection time.

All mares were at least 10 years old.

825 AA: Anglo Arab or Anglo-Arabian type; SF: Selle Français section A or B; ST: Standardbred; SB:

826 Saddlebred. Age, parity, weight, and height are presented as mean ± SD

828	Table 2: C	Comparison of sel	ected genes expression b	efore and after deconvolu	ution

			ICMandTE <i>vs</i> TE_part		_	CM_cells vs part
	Gene name	Ensembl ID	log2FC from DeSeq2	padj	log2FC from DeSeq2	padj
	SOX2	ENSECAG00000010653	3.82	0.069	5.08	1.02E-04
	NANOG	ENSECAG00000012614	3.01	0.017	4.26	2.50E-06
	SPP1	ENSECAG00000017191	4.02	8.39E-09	3.99	2.19E-05
	LIN28B	ENSECAG00000020994	2.75	0.045	3.64	5.14E-06
	SMARCA2	ENSECAG00000024187	0.10	1.00	0.64	0.398
CM	POU5F1 (OCT4)	ENSECAG0000008967	0.40	0.046	0.96	1.6E-07
2	ID2	ENSECAG0000008738	0.31	0.409	0.82	1.01E-04
	DNMT3B ENSECAG00000012102		0.37	0.034	0.87	2.10E-09
	DPPA4	ENSECAG00000013271	0.38	0.089	0.87	5.8E-06
	SALL4	ENSECAG00000018533	0.09	0.829	0.29	1.41E-03
	KLF4	ENSECAG00000010613	0.05	0.999	-0.13	0.848
	UTF1	ENSECAG0000039888	-0.18	0.973	-0.91	3.28E-04
	TFAP2A	ENSECAG00000017468	-0.14	0.700	-0.19	0.259
	CDX2	ENSECAG00000027754	-0.10	0.924	-0.24	0.032
	ELF3	ENSECAG00000014608	-0.19	0.830	-0.67	1.73E-04
TE	GATA2	ENSECAG00000016768	-0.04	0.998	-0.11	0.614
	GATA3	ENSECAG00000024574	-0.11	0.789	-0.08	0.756
	TEAD4	ENSECAG00000011303	-0.11	0.692	-0.22	8.81E-03
	FREM2	ENSECAG0000020410	-0.01	1.000	0.24	0.439

Gene expressions were obtained from RNA of 11 equine embryos bissected in two hemi-embryos: one part was composed only of trophoblast (TE), TE_part, while the other part was composed of TE and inner cell mass (ICM), ICMandTE. As it is impossible to estimate the proportion of each cell in ICMandTE, a deconvolution algorithm (package DeMixT) was used to estimate gene expression of these different kind of cells. DeMixT_ICM_cells dataset corresponds to the deconvoluted gene expression of ICM cells from ICMandTE. Log2 fold change (log2FC) and padj (adjusted p-value with Benjamini-Hochberg correction) were obtained with Deseq2 package. TE_part is the reference group

in both analyses: when log2 fold changes (log2FC) > 0, gene is more expressed in ICMandTE or DeMixT_ICM_cells while when log2FC < 0, gene is more expressed in TE_part. Green is used to represent gene differentially expressed in the present study. Orange is used to represent genes that have been previously identified as predominant in the ICM [53] but which are identified here as predominant in the TE.

841

Table 3: Up- and down-regulated genes coding for a protein in the inner cell mass of equine embryos according to old mare parity

Ensembl Name	Entrez Gene ID	Description	GO Molecular function	GO Biological process	log2 Fold Change	Adjusted p-value
ENSECAG00000002796	GABRR1	Gamma-aminobutyric acid type A receptor subunit rho1	Chloride channel Ion channel	Ion transport Chloride transmembrane transport Gamma-aminobutyric acid signaling pathway Regulation of membrane potential Signal transduction	5.19	0.004
ENSECAG00000007423	ZKSCAN2	zinc finger with KRAB and SCAN domains 2	DNA-binding transcription factor activity, RNA polymerase II-specific Metal ion binding	Regulation of transcription by RNA polymerase II	4.42	0.008
ENSECAG00000012957	RHAG	Rh associated glycoprotein	Ammonium transmembrane transporter activity Ankyrin binding Leak channel activity	Ammonia transport Cellular ion homeostasis Bicarbonate transport Carbon dioxide transport	4.25	0.024
ENSECAG00000017370	PPT1	palmitoyl-protein thioesterase 1	Hydrolase	Nervous system development Lipid catabolism process Protein transport Protein catabolic process	1.97	0.008
ENSECAG00000023194	XYLB	xylulokinase	ATP binding Kinase Transferase	Carbohydrate metabolism Xylose metabolism	0.63	0.009
ENSECAG00000036419	RPL36	ribosomal protein L36	Ribonucleoprotein Ribosomal protein	Translation Nuclear-transcribed mRNA catabolic process, nonsense-mediated decay rRNA processing	-0.36	0.001
ENSECAG00000012792	MIF	macrophage migration inhibitory factor	Cytokine Isomerase	Immunity Inflammatory response Innate immunity	-0.39	0.001
ENSECAG0000007113	ATXN10	ataxin 10	Enzyme binding	Nervous system development Cilium assembly Neuron projection development	-0.46	3.78E-08

ENSECAG00000020469	VPS51	VPS51 subunit of GARP complex		Lipid transport Protein transport Transport Autophagy Brain morphogenesis Golgi organization	-0.56	0.024
ENSECAG0000006539	EXOSC7	exosome component 7	RNA-binding	rRNA processing	-0.65	0.002
ENSECAG00000042897	H2AC20	H2A clustered histone 20	DNA-binding	Chromatin silencing	-0.76	0.007
ENSECAG00000020116	RABEP2	rabaptin, RAB GTPase binding effector protein 2	Growth factor activity GTPase activator activity	Cilium biogenesis/degradation Endocytosis Protein transport Transport	-0.76	0.009
ENSECAG00000018683	PCF11	PCF11 cleavage and polyadenylation factor subunit	mRNA processing	mRNA 3'-end processing mRNA cleavage mRNA polyadenylation mRNA splicing, via spliceosome termination of RNA polymerase II transcription	-0.78	0.010
ENSECAG00000012328	STAT5A	signal transducer and activator of transcription 5A	Activator DNA-binding	Transcription Transcription regulation	-0.89	0.001
ENSECAG00000014662	SLX4	SLX4 structure-specific endonuclease subunit	DNA damage DNA recombination DNA repair	DNA repair	-0.90	0.020
ENSECAG00000028573	TRIOBP	TRIO and F-actin binding protein	Actin-binding	Cell cycle Cell division Mitosis	-0.99	0.014
ENSECAG00000016843	SETD5	SET domain containing 5	Chromatin regulator Methyltransferase Transferase	Transcription Transcription regulation Histone H3-K36 trimethylation Regulation of histone acetylation Regulation of chromatin organization	-1.02	0.023

ENSECAG0000008726	NPY	neuropeptide Y	Neuropeptide Calcium channel regulator activity G protein-coupled receptor activity Hormone activity signaling receptor binding	Adult feeding behavior G protein-coupled receptor signaling pathway Innate immune response Central nervous system neuron development	-1.27	0.001
ENSECAG00000021312	ABHD14A	abhydrolase domain containing 14A	Hydrolase		-1.30	0.007
ENSECAG00000030141	PXMP2	peroxisomal membrane protein 2	Transporter	Protein import into peroxisome membrane	-1.37	0.019
ENSECAG00000014057	ASB1	ankyrin repeat and SOCS box containing 1	Developmental protein	Ubiquitine-like conjugation pathway	-1.42	0.006
ENSECAG00000022130	CPT2	carnitine palmitoyltransferase 2	Acyltransferase	Fatty acid metabolism Lipid metabolism Transport	-1.52	0.004
ENSECAG00000015074	LIPE	lipase E, hormone sensitive type	Hydrolase	Cholesterol metabolism Lipid degradation Lipid metabolism Steroid metabolism Sterol metabolism	-1.53	0.018
ENSECAG0000008600	АРОВ	apolipoprotein B	Heparin-binding	Cholesterol metabolism Lipid metabolism Lipid transport Steroid metabolism Sterol metabolism Transport	-1.99	0.041
ENSECAG00000011240	CNTN1	Contactin 1	Carbohydrate binding	Cell adhesion Notch signaling pathway	-2.36	0.001

Log2 Fold-Change > 0 indicates up-regulation of the gene in embryos from old nulliparous mares; Log2 Fold-Change < 0 indicates down-regulation of the gene in embryos

845 from old nulliparous mares

Table 3: Up- and down-regulated genes coding for a protein in the trophoblast part of equine embryos according to old mare parity

Entrez Ensembl Name Gene ID Description		GO Molecular function	GO Biological process	log2 Fold Change	Adjusted p-value	
ENSECAG00000022403	EFCAB2	EF-hand calcium binding domain 2	Calcium ion binding		2.70	0.029
ENSECAG00000012686	SNCAIP	synuclein alpha interacting protein	Binding	0.98	0.012	
ENSECAG00000001080	PLPP5	phospholipid phosphatase 5	Hydrolase	0.86	0.003	
ENSECAG00000012682	NFU1	NFU1 iron-sulfur cluster scaffold	Iron ion bindng	0.61	0.024	
ENSECAG00000004829	SYNJ1	synaptojanin 1	Hydrolase RNA-binding	Endocytosis Brain development Membrane organization Phosphatidylinositol metabolic process Inositol phosphate metabolic process	0.57	0.043
ENSECAG00000019000	CHPT1	choline phosphotransferase 1	Transferase Lipid metabolism Phospholipid metabolism		0.52	0.026
ENSECAG00000009962	MSH3	mutS homolog 3	DNA-binding	-0.41	0.030	
ENSECAG00000024071	MZT2B	mitotic spindle organizing protein 2B			-0.43	0.029

ENSECAG00000015935	ACTB	Actin beta	BindingStructural constituent of cytoskeleton	Apical protein localization ATP-dependent chromatin remodeling Axonogenesis Cell junction assembly Cell motility Establishment or maintenance of cell polarity Membrane organization Morphogenesis of a polarized epithelium Negative regulation of protein binding Positive regulation of gene expression, epigenetic Protein deubiquitination Regulation of cyclin-dependent protein Regulation of transepithelial transport	-0.44	0.001
ENSECAG0000003599	GK	glycerol kinase	Kinase Transferase	Glycerol metabolism Phosphorylation Triglyceride biosynthetic process	-0.48	0.040
ENSECAG00000009054	СКВ	creatine kinase B	Kinase Transferase	Cellular chloride ion homeostasis Cerebellum development Creatine metabolic process	-0.95	0.040
ENSECAG00000037199	H4C3	H4 clustered histone 3	DNA-binding	DNA-templated transcription, initiation Double-strand break repair Mitotic chromosome condensation Negative regulation of gene expression, epigenetic	-1.06	6.03E-09
ENSECAG00000020400	ENO3	enolase 3	Lyase	Glycolysis Aging Gluconeogenesis Skeletal muscle tissue regeneration	-1.31	0.030
ENSECAG00000039059	CD99	CD99 molecule (Xg blood group)		Cell adhesion Diapedesis Positive regulation of neutrophil extravasation Positive regulation of T cell extravasatiion	-1.33	0.001

ENSECAG00000012921	ALPK2	alpha kinase 2	Serine/threonine-protein kinaseTransferase	Establishment of cell polarity Heart morphogenesis Negative regulation of Wnt signaling pathway involved in heart development Regulation of apoptotic processRegulation of gene expression	-2.23	0.001
ENSECAG00000017922	ADAM19	ADAM metallopeptidase domain 19	Hydrolase Metalloprotease	Amyloid precursor protein catabolic process Extracellular matrix organization Membrane protein ectodomain proteolysis Placenta development Positive regulation of cell-cell adhesion mediated by cadherin Positive regulation of gene expression Protein processing	-2.65	0.047
ENSECAG00000021052	EOMES	eomesodermin	Activator Developmental protein DNA-binding	Adaptive immunity Differentiation Gastrulation Transcription Transcription regulation	-2.88	0.029
ENSECAG00000008777	MAP1B	microtubule associated protein 1B	Actin binding Phospholipid binding Microtubule binding	Cellular process Dendrite development Mitochondrion transport among microtubule	-4.61	0.009

Log2 Fold-Change > 0 indicates up-regulation of the gene in embryos from old nulliparous mares; Log2 Fold-Change < 0 indicates down-regulation of the gene in embryos

849 from old nulliparous mares

			Number		FDR	
	Database	Geneset name	of genes	NES	q-val	Subset of genes that contributes most to the enrichment result
ICM	GO BP	CYTOPLASMIC MICROTUBULE ORGANIZATION	45	-2.17	0.013	SLAIN2, CCDC88A, EZR, CHP1, DYNC1H1, IFT172, KATNA1, TUBGCP6, TUBGCP4, FIGN, HOOK3, CLASP1, DLG1, DST, TUBGCP5, NUMA1, WDR73, CCDC88B, SLK, CCDC88C
	KEGG	NEUROACTIVE LIGAND RECEPTOR INTERACTION	65	1.98	0.019	GABRR1 , GPR50, SCTR, GLP1R, CRHR2, P2RY11, GABRA5, PLG, MCHR2, ADORA2B, EDNRA, GABRA4, GHR, LPAR4, F2RL1, GRIA2, NPY1R, GABRG1, CHRNA5, GABRG2, GH2, F2R, CHRNA3, CGA, ADORA2A, CNR2, GRIN2C, CHRNA2, HCRTR2, NR3C1
	GO BP	REGULATION OF GLIAL CELL DIFFERENTIATION	35	-2.21	0.008	PTN, SLC45A3, NF1, DAG1, PRPF19, LIN28A, CXCR4, TNFRSF1B, LDLR, ID2, TGFB1, EPHA4, BIN1, BMP2, MDK, NOTCH1, TMEM98
	REACTOME	ECM PROTEOGLYCANS	46	-2.20	0.002	BCAN, ITGA7, COL6A6, HSPG2, COL4A5, COL5A2, ITGAV, NCAN, DAG1, TNXB, LAMB1, COL1A2, PTPRS, LAMA1, TGFB3, AGRN, MATN4, COL4A1, TGFB1, COL9A2, SPARC, VCAN, COL1A1, NCAM1, COL4A2, TNC
	GO BP	POSITIVE REGULATION OF GLIAL CELL DIFFERENTIATION	23	-2.12	0.018	PTN, DAG1, PRPF19, CXCR4, TNFRSF1B, ID2, TGFB1, BIN1, BMP2, MDK, NOTCH1
	REACTOME	INTEGRIN CELL SURFACE INTERACTIONS	50	-2.11	0.004	COL3A1, COL4A6, ITGA5, ITGA7, FBN1, ICAM3, COL6A6, BSG, HSPG2, CD47, COL4A5, COL5A2, ITGAV, CDH1, ICAM1, SPP1, COL1A2, AGRN, COL4A1, FGA, COL9A2, COL1A1, COL4A2, TNC, ITGA3, COL18A1, FGB
TE	REACTOME	EXTRACELLULAR MATRIX ORGANIZATION	178	-1.99	0.018	CAPN15, COL4A6, ADAMTS2, ITGA5, MMP9, MMP11, ITGB4, MFAP3, CASK, BCAN, CAPNS1, ITGA7, P3H3, ADAMTS1, FBN1, ICAM3, COL6A6, BSG, HSPG2, CD47, LTBP2, ADAMTS18, COL4A5, CAPN1, DMD, TIMP1, COL5A2, ITGAV, LAMB3, PLG, ACTN1, NCAN, DAG1, CDH1, CTSD, MMP2, TNXB, LAMB1, ICAM1, SPP1, COL1A2, SPOCK3, LAMC2, PTPRS, LAMA1, PXDN, TGFB3, EFEMP1, FBLN2, AGRN, COL14A1, MATN4, COL4A1, FURIN, TGFB1, FGA, ADAMTS5, COL9A2, LTBP3, SPARC, SERPINH1, DST, VCAN, COL1A1, PLOD3, ADAM19 , NCAM1, MMP14, BMP2, COL4A2, PLEC, TNC, ADAM8, ITGA3, COL18A1, FGB
	REACTOME	LAMININ INTERACTIONS	23	-1.98	0.015	NID2, COL4A6, ITGB4, ITGA7, HSPG2, COL4A5, ITGAV, LAMB3, LAMB1, LAMC2, LAMA1, COL4A1, COL4A2, ITGA3, COL18A1
	KEGG	HISTIDINE METABOLISM	18	-1.93	0.033	HNMT, ALDH3A1, ALDH7A1, HAL, AMDHD1, UROC1, HEMK1, FTCD, MAOB
	REACTOME	REGULATION OF BETA CELL DEVELOPMENT	22	-1.91	0.038	HNF4A, CREBBP, HNF1B, HNF1A, FOXA2, NR5A2, MAML1, AKT2, PKLR, PDX1, NOTCH1

REACT	OME NR1H3 NR1H2 REGULATE GEN EXPRESSION LINKED TO CHOLESTEROL TRANSPORT ANI EFFLUX	28	-1.90	0.034	EEPD1, ARL4C, TNRC6A, GPS2, NCOR2, APOC2, RXRB, KDM4A, MOV10, PLTP, NR1H2, APOE, TNRC6B, ABCA1, RXRA
REACT	OME NOTCH3 INTRACELLULAR OME DOMAIN REGULATES TRANSCRIPTION	16	-1.90	0.031	STAT1, KAT2A, CREBBP, WWC1, PTCRA, MAML1, PBX1, NOTCH1
REACT	OME DEGRADATION OF THE EXTRACELLULAR MATRIX	76	-1.89	0.028	FBN2, COL3A1, CAPN15, COL4A6, MMP9, MMP11, BCAN, CAPNS1, ADAMTS1, FBN1, COL6A6, BSG, HSPG2, ADAMTS18, COL4A5, CAPN1, TIMP1, COL5A2, LAMB3, PLG, CDH1, CTSD, MMP2, LAMB1, SPP1, COL1A2, SPOCK3, LAMC2, COL14A1, COL4A1, FURIN, ADAMTS5, COL9A2, COL1A1, MMP14, COL4A2, ADAM8, COL18A1
KEG	G ECM RECEPTOR INTERACTION	50	-1.86	0.037	COL3A1, COL4A6, ITGA5, ITGB4, ITGA7, COL6A6, HSPG2, CD47, COL5A2, ITGAV, LAMB3, DAG1, TNXB, LAMB1, SPP1, COL1A2, LAMC2, LAMA1, AGRN, COL4A1, COL1A1, COL4A2, TNC, ITGA3

852 ICM: Inner cell mass; TE: Trophoblast; Genes in bold have also been identified as differentially expressed. Gene sets in blue are related to the extracellular matrix organization

853 and function while ones in green represent gene sets related to NOTCH signaling pathway.





