1                 # A new protocol for speed vernalisation of winter cereals

3     Jin-Kyung Cha[1*], Kathryn O'Connor[2*], Samir Alahmad[3], Jong-Hee Lee[1], Eric Dinglasan [3],

4     Hyeonjin Park[1], So-Myeong Lee[1], Dominique Hirsz[2], Soon-Wook Kwon[4], Youngho Kwon[1],

5     Kyeong-Min Kim[1], Jong-Min Ko[1], Lee T. Hickey[3+], Dongjin Shin[1+], Laura E. Dixon[2+]

[1]National Institute of Crop Science, RDA, Miryang, 50424, Korea

[2]School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT, UK

[3]Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, St Lucia, Queensland, Australia

[4]Department of Plant Bioscience, Pusan National University, Miryang 60463, Korea

* These authors contributed equally to the work

E-mails for corresponding authors:

l.hickey@uq.edu.au, jacob1223@korea.kr, l.dixon2@leeds.ac.uk

***Keywords***

***Abstract***

There are many challenges facing the development of high-yielding, nutritious crops for
future environments. One limiting factor is generation time, which prolongs research and
plant breeding timelines. Recent advances in speed breeding protocols have dramatically
reduced generation time for many short-day and long-day species by optimising light and
temperature conditions during plant growth. However, winter crops with a vernalisation
requirement still require up to 6–10 weeks in low-temperature conditions before transition to
reproductive development. Here, we tested a suite of environmental conditions and protocols
to investigate if vernalisation can be satisfied more efficiently. We identified a vernalisation
method consisting of exposing seeds at the soil surface to an extended photoperiod of 22 h
day:2 h night at 10°C with transfer to speed breeding conditions that dramatically reduces
generation time in both winter wheat (*Triticum aestivum*) and winter barley (*Hordeum*
*vulgare*). Implementation of this protocol achieved up to five generations per year for winter
wheat or barley, instead of the two typically obtained under standard vernalisation and plant
growth conditions. The protocol has great potential to enhance training and to accelerate
research, pre-breeding, and breeding outcomes focussed on winter crop improvement.

***Main***

Vernalisation is the requirement for a prolonged period of cold before certain plants can
transition from vegetative to reproductive development. Vernalisation thus coordinates a
plant's development with its environment. In agriculture, vernalisation maximises the growing
season of a crop by enabling autumn sowing without the risk of plants transitioning to
reproductive development and becoming damaged by winter conditions[1]. The longer growth
duration improves crop productivity and is a common feature of many wheat (*Triticum*
*aestivum*) and barley (*Hordeum vulgare*) cultivars grown throughout the high-yielding regions
of Northern Europe, the United States, Asia, and the South Pacific. However, retaining
vernalisation in elite germplasm comes at a cost for generation turnover in breeding
programmes. To date, the agronomic and academic standard cereal vernalisation protocol
entails 6–10 weeks at a low temperature of 2–6°C[1]. Here, we optimised the environmental
conditions during vernalisation and obtained a substantial reduction in the generation time
for winter wheat and winter barley. This new soil-surface protocol uses an extended
photoperiod (22 h light:2 h dark) and warmer temperatures (8–10°C) than typical
vernalisation treatments and was effective across a genetically diverse set of germplasm.
This 'speed vernalisation' protocol will accelerate the pace of research, training, and pre-
breeding and breeding outcomes for winter cereal crops.

Due to vernalisation, crop improvement programmes that focus on developing winter
cultivars do not fully benefit from recent advances made via speed breeding, in which the
photoperiod is extended to a 22-h daylength and plants are grown at 22°C during this period
(Supplementary Table 1)[2,3]. Vernalisation is considered a winter response, and artificial
vernalisation conditions reflect that with a standard protocol of 6–10 weeks at 2–6°C under a
short-day, 8-h-light:16-h-dark, photoperiod[1,4]. However, recent and historic evidence showed
that vernalisation proceeds sometimes more efficiently at slightly warmer conditions[4-6], which
suggested to us that current vernalisation conditions may be sub-optimal. To investigate this
possibility, we tested vernalisation efficiency at warmer temperatures and under short-day
photoperiods (Extended Data Fig. 1a-f). As the vernalisation response is quantitative (i.e., up
to the point of vernalization saturation, increasing amounts of vernalisation will lead to an
acceleration in flowering time), the rate and point of completion of vernalisation can be
assessed by transferring vernalising plants to floral inductive conditions and scoring
flowering time. To challenge the protocols, we tested wheat cultivars with weak (e.g., cv.
Claire, one copy of *VERNALIZATION1* (*VRN-A1*)), moderate (e.g., cv. Buster, two *VRN-A1*
copies), and strong (e.g., cv. Hereward and Charger, three *VRN-A1* copies) vernalisation
requirements[7]. For all cultivars tested, vernalisation completed efficiently following 6 weeks
at 10°C or 14°C (Extended Data Fig. 1a-b), with warmer temperatures leading to the
development of large vegetative meristems (Extended Data Fig. 1c-f). Our results support
evidence from *Arabidopsis thaliana* in which vernalisation completes in the autumn[8],
suggesting that artificial vernalisation conditions using low temperatures of 4–6°C may not
be necessary. Here, raising the vernalisation temperature to 10°C for 6 weeks met all
vernalisation needs of all tested cultivars, even those with long vernalisation requirements.
Importantly, these new conditions of 10°C under an 8-h-light:16-h-dark photoperiod, which
we refer to as warm regular vernalisation (wRV), can be supported by most controlled
growth chambers.

Building on this, we were curious if there were opportunities for further optimisation. Short-
day photoperiods are typically used for artificial vernalisation of cereals as they repress the
expression of the long-day-activated flowering repressor *VRN2*[9]. However, there is
increasing evidence that pre-vernalisation repression of flowering in cereals is a multigenic
response[10,11], prompting us to hypothesise that *VRN2* may not be a limiting factor. To test
this possibility, we vernalised plants at 10°C under a 22-h-light:2-h-dark photoperiod (speed
vernalisation: SV) before transfer to speed breeding (SB) conditions[2] (SB: 22 h light:2 h
dark, 22°C:17°C cycles or constant 22°C) (Fig. 1a). As with wRV, meristems remained
vegetative during SV, and the plants went on to produce fully developed, fertile spikes
(Extended Data Fig. 1g-n). To challenge our protocol, we used genotypes with a range of

pedigrees, including Korean, European, and American wheat cultivars. We observed a
reduction in generation time for many cultivars subjected to SV compared to wRV also
transferred to SB (Fig. 1: Supplementary Table 2) and a substantial reduction in generation
time when compared to wRV followed by regular glasshouse conditions (RG 20°C; 16 h
light:8 h dark)(Extended Data Fig. 1). For example, following 4 weeks of SV, the generation
time for cv. Claire was accelerated by at least 10 weeks compared to standard practice (Fig.
1 and Extended Data Fig 1a). For cv. Charger, SV enabled an acceleration of flowering
following 4 weeks of vernalization from non-flowering under wRV to 125 days under SV, both
transferring into SB (Fig. 1c). However, this acceleration still meant that cv. Charger
generation time was longer than a cultivar with a lower vernalisation requirement (e.g., cv.
Claire). The variability in the response suggested that *VRN-A1* copy number may be
important, so we were interested in identifying ways to improve our protocol to mitigate this.

We thus used other parameters to increase the universal nature of the vernalisation protocol.
Under standard practice, plants are transferred to vernalisation following 1–2 weeks of
growth under regular glasshouse conditions (RG 20°C; 16 h light:8 h dark) to allow seedling
establishment. Our SV protocol uses germination and growth under 22-h-light:2-h-dark
photoperiods, so seedlings grow faster than with wRV. Therefore, to limit the extent of initial
seedling growth, we tested if seedlings responded efficiently without pre-growth in the
glasshouse. Accordingly, we subjected seeds to four treatments: T1) germination and growth
at 4°C in the dark, T2) germination in SB before transfer to SV, T3) SV with the seed buried
in soil, and T4) SV with the seed on the soil surface (Fig. 1f, Supplementary Table 3).
Unexpectedly, vernalisation was most efficient when seeds were placed on the soil surface
and exposed to light under SV conditions (Fig. 1g), which significantly reduced the time to
flowering by an additional 8 days compared to the other conditions (Duncan's multiple test, α
= 0.05) for cv. Keumgang (one *VRN-A1* copy). To optimise the protocol, we tested a suite of
durations and temperatures of seed-surface SV conditions using the Korean winter wheat
cv. Keumgang and the American winter wheat cv. Sturdy (one *VRN-A1* copy). These
experiments confirmed that 8–10°C is the most efficient and reliable vernalisation
temperature (Fig. 1j-k, Supplementary Tables 4 and 5). This result revealed a seed-based
aspect of the vernalisation response in cereals that is similar to dormancy in dicots[12]. The
same method was also successful in cultivars with a strong vernalisation requirement (e.g.,
cv. Hereward and Charger) and reduced generation time by at least 4 weeks compared to
SV or wRV and transfer to SB (Figure 1b-i). Importantly, the combination of seed-surface
vernalisation via SV followed by SB (hereafter SGV_SB) conditions reduced the duration of
vernalisation needed, with 3–4 weeks at 10°C being optimal; therefore, our protocol enables
a higher throughput of plants through vernalisation compared to RV and SV.

For use in academic and industrial breeding programmes, the SGV_SB protocol should
support usual plant development. Therefore, we measured multiple plant growth parameters,
obtained normal seed set, and observed standard plant development, although plant height
was slightly reduced (Extended Data Fig. 2). We also tested the effectiveness of the
SGV_SB protocol on a wheat diversity panel regularly used in wheat breeding programmes
in Korea. Of the 51 winter cultivars in the panel, 45 were fully responsive to SGV_SB
conditions (Supplementary Table 6). Overall, our data demonstrate that vernalisation on the
soil surface imposed by the SGV_SB protocol reduces the vernalisation requirement and
generation time for many winter wheat cultivars tested and is effective even on cultivars that
traditionally have a longer vernalisation requirement.

Given the similarity between the vernalisation response of wheat and barley, we
hypothesised that SGV_SB may accelerate generation time in winter barley as well.
Accordingly, we subjected 60 diverse winter barley cultivars originating from 34 growing
regions or countries to RV_RG, RV_SB, and SGV_SB (Extended Data Fig. 3a, b). We
recorded days to flowering in each experiment (Fig. 2a, b). Notably, all cultivars flowered and
produced viable seeds under all conditions. Under SGV_SB, the entire population flowered
substantially earlier, on average after 50 days compared to 92 days under RV_RG and 68
days under RV–SB (Fig. 2a). Therefore, in contrast to wheat, the collection of winter barley
cultivars was completely responsive to the SGV_SB protocol.

Crossing and the subsequent development of genetically stable or inbred lines are routine
practices in both breeding and research programmes. However, these techniques are
particularly time consuming for populations derived from winter × spring and winter × winter
crosses, as each generation must be vernalised to avoid unintended selection against the
vernalisation mechanism. To test the SGV_SB protocol for use in population development,
we crossed cv. Jokyung (spring) and cv. Joongmo2008 (winter) wheat cultivars in the field in
Korea during the 2018–2019 winter season. Throughout the subsequent 15 months, we
reached the F5 generation, with each generation taking an average of 82.4 days (Fig. 2c,
Supplementary Table 7, 8). The SGV_SB protocol considerably reduced the variation in
days to flowering for populations derived from winter × spring and winter × winter crosses
(Fig. 2d; Extended Data Fig. 3, Supplementary Table 9). Early and more synchronous
flowering across segregating or diverse germplasm can facilitate more efficient crossing and
rapid generation times. We harvested at least 25 seeds from each F5 plant, which is
sufficient to bulk seed in the field and subsequent evaluation. Using a projected timeline for
plant generations (Fig. 2e-f), we highlight the opportunity to increase the number of plant
generations within a 12–18-month period for both spring and winter wheat and barley.
Impressively, the SGV_SB protocol applied to some winter cultivars reached generation

turnover times similar to those of spring cultivars. Furthermore, the seed surface
vernalisation is extremely high-throughput, with a density of up to 1,709 seeds per $m^2$ being
treated when utilising 128-cell seed trays.

Our optimised vernalisation conditions (SV and SGV) that lead to a reduction in overall
generation time challenge our current understanding of the vernalisation mechanism itself,
where short days reduce *VRN2* expression and cold temperatures activate *VRN1*
expression[4]. Therefore, we investigated how vernalisation genes responded during SV_SB
and if they differed between the seed surface (SGV) versus buried seed (SV) protocol (Fig.
3). The expression patterns of vernalisation genes followed the expected profiles in SV_SB
buried seed, using cv. Claire, with *VRN1* and *VRN3* (also named *FLOWERING LOCUS T-*
*like1* (*FT1*)) transcript levels increasing during vernalisation and *VRN2* (a locus comprised of
*ZCCT1* [for *Zinc domain and CONSTANS, CONSTANS-LIKE, TOC1*] and *ZCCT2*) transcript
levels decreasing (Fig. 3a-d). Notably, *VRN1* and *VRN3* expression was lower under SV
compared to wRV conditions (Fig. 3a-b), indicating that these genes may not represent the
exclusive route promoting flowering under SV. We observed a sustained repression of
*ZCCT1* expression during and following SV when plants were transferred to SB, which was
unexpected given the extended photoperiod of SV, which would be anticipated to promote
*VRN2* expression (Fig. 3d). However, comparing the SV_SB conditions in cv. Hereward,
which vernalised more rapidly with the seed-surface protocol, *ZCCT1* steadily increased in
expression, opposite to that in the buried seed condition (Fig. 3j-k). This result indicated that
considering the *VRN2* locus as a single gene is potentially misleading for our understanding
of the vernalisation response and that an additional or alternative vernalisation response is
activated during SGV. Further examination of this response may identify new breeding
targets in the regulation of vernalisation and flowering time in cereals.

The time required for the introgression and stacking of novel alleles remains a bottleneck in
the development of improved cultivars. One of the limiting factors is the vernalisation
requirement, which imposes a biological constraint on generation time. Here, we identify a
method to reduce generation time for winter wheat and barley germplasm using alternative
vernalisation conditions. The protocols we developed identify the environmental parameters
that can be further modified to account for local or genotypic variation in vernalisation
efficiencies and therefore offer a framework to universally reduce generation times in winter
cereals. The approach could also be adapted to other winter crops (e.g., canola [*Brassica*
*napus*]) or vegetables with a vernalisation requirement to reduce generation time and
accelerate breeding outcomes.

**Materials and methods**

**Cultivars used:** Wheat and barley cultivars used in this study are in Supplementary Table 10.

**European winter wheat assessment and gene expression (assessed at The University of Leeds, UK)**

**Conditions used: Wheat (*Triticum aestivum*)**

*wRV_RG:* 8 h light:16 h dark 10°C into 16 h light:8 h dark 22°C

*wRV_SB*: 8 h light:16 h dark 10°C into 22 h light:2 h dark 22°C

*SV (and SGV) _SB*: 22 h light:2 h dark 10°C into 22 h light:2 h dark 22°C

Please note warmer conditions were used in wRV than are classically used in RV to enable direct photoperiod comparison for phenotype and gene expression analysis.

Seeds were germinated for 2 days in darkness at 4°C in 9-cm petri dishes that had a layer of filter paper saturated with 5 mL dH$_2$O.  The germinated seeds were transferred to 3 x 3-cm cell pots of JIC cereal mix[4] and placed under vernalisation conditions in Snijders MICROCLIMA MC1000 cabinets. Plants were watered when required, and no additional nutrients were added. At 1- or 2-week intervals, plants were sampled for gene expression and apex analysis (see below), and three plants were transferred to glasshouse growth conditions for RG (PhytoLux Plessey; model ATTIS-7) and SB (Heliospectra; model MITRA). Plants were placed in cereal mix (as above) in 9 x 9-cm pots. Plants were sampled 1 and 2 weeks after transfer to SB conditions. Flowering time was recorded as half-ear emergence (Zadok scale 55). Plants that did not flower after 170 days were recorded as "non-flowering." A control group using the same four genotypes ($n$ = 10, $s$ = 40) was grown under constant SB conditions, and a representative for each genotype was imaged once the plant had reached maturity.

**Speed-green vernalisation.** Seeds were placed in 9-cm petri dishes containing filter paper and 5 mL dH$_2$O and kept in complete darkness at 4°C for 48 hours. Seeds were then placed on the soil surface (John Innes cereal mix) in a P24 seedling tray. Care was taken to press the seed into the soil surface while ensuring the seed remained uncovered and exposed to the light. Plant trays were watered from the base, and a spray bottle of dH$_2$O was used to mist the soil surface; particular care was taken misting the soil during the first week of growth, when the roots were anchoring into the soil. At set weekly intervals, plants were moved into SB conditions and flowering was recorded.

**Gene expression samples.** Leaf samples from three plants per biological replicate and for three biological replicates ($n$ = 3) were taken at each sampling stage. Samples were taken 1 h after lights on. To investigate gene expression during SV, leaf tissue was sampled at 1, 2, 3, 4, 5, 6, 7, and 8 weeks of growth under SV conditions. Total RNA was extracted using the Spectrum™ Plant Total RNA Kit (Sigma-Aldrich) following the manufacturer's recommended protocol. RNA samples were treated using RQ1 RNase-Free Dnase (Promega), and first-strand cDNA synthesis primed with Oligo dT was processed using SuperScript™ III Reverse Transcriptase (Invitrogen) and RNaseOUT™ (Invitrogen). The cDNA was diluted (1:10), and quantitative reverse transcription PCR (RT-qPCR) was performed using the CFX96™ Thermal Cycler (Bio-Rad) and GoTaq® Master Mix (Promega). Primers used are provided in Supplementary Table 11. Expression levels of the genes of interest were calculated relative to TraesCS5A02G015600 following the $2^{\wedge -\Delta CT}$ format (where $\Delta CT$ = *GOI* CT – TraesCS5A02G015600 CT).

Apex samples: three plants were dissected for each apex sample ($n$ = 3) and imaged on a Keyence microscope, and apex length was measured using ImageJ[13].

**Methods for Wheat (*Triticum aestivum*) (assessed at RDA, S. Korea)**

**Conditions used: Wheat (*Triticum aestivum*)**

*wRV–SB*: 8 h light:16 h dark 10°C into 22 h light:2 h dark 22°C:17°C

SGV_RG: 22 h light:2 h dark 10°C into 16 h light:8 h dark 22°C:17°C

*SGV–SB*: 22 h light:2-h dark 10°C into 22 h light:2 h dark 22°C:17°C

**Optimization of vernalization treatment method.** Seeds were germinated in 9-cm petri dishes containing 10 mL of water and at 4°C for 3 days under dark conditions. Then, seeds were transferred to 25°C until they reached the growth stage 07 and about 0.2-mm coleoptile length[14]. For all experiments, each square pot (7(L) × 7(W) × (7H)) used was filled with 245 mL of soil, which is a mixture of paddy rice soil (Punong, Korea) and horticulture soil (Seoul-Bio, Korea) at a 2:1 ratio. To find an optimum vernalisation treatment method, four cold treatment methods were applied using the Korean winter variety Keumgang as follows: T1) seed vernalisation, T2) green vernalisation, T3) speed-green vernalisation with covering soil, and T4) speed-green vernalisation without covering soil about 1.5 cm in depth. Regarding the seed vernalisation method (T1), germinated seeds were placed in a 4°C refrigerator for 4 weeks without light and cultivated under speed breeding (SB) conditions. For green vernalisation (T2), plants initially placed under SB for 1 week were further grown under 22-h-light:2-h-dark photoperiod cycles at 8°C for 4 weeks. Concerning the speed-green vernalisation with (T3) and without (T4) covering soil, germinated seeds were sown

and grown at 8°C for 4 weeks under 22-h-light:2-h-dark cycles. LED lights (red 8: blue 3:

white 2, Estech LED, Korea, Extended Data Fig. 2b-c) in SGV condition were set to 410

µmol m$^{-2}$ s$^{-1}$ of illumination. All plants were then transferred to SB (2,500 µmol m$^{-2}$ s$^{-1}$ of

light) to evaluate the time to flowering after vernalisation treatment; $n$ = 8 plants for each

evaluation. Spectral measurement of light composition was performed using the MK350N

handheld spectrometer (UPRtek, Miaoli county, Taiwan).

**Optimization of vernalisation temperature and periods.** To optimize the cold treatment

protocol for speed vernalisation, temperatures ranging from 6–12°C were applied for 2–6

weeks under SGV without covering soil (SGV condition 4). Six plants were then transferred

to SB conditions to examine days from germination to heading. The growth stages (GS32

and GS59) were recorded as reported by[14], and plant height was measured from the ground

to the start point of the spike.

**Evaluation of heading time of genetic resources.** To investigate days from germination to

heading time of wheat genetic resources, germinated seeds were placed in the cell of a 72-

cell tray (34 mL/cell) and cultivated under the speed-green vernalisation without covering soil

condition (SGV condition 4). Plants were grown under SGV for 4 weeks, followed by transfer

to SB and observation until heading time (GS59), $n$ = 4.

**Plant density.** SGV was conducted under a range of planting densities and supported very-

high-throughput vernalisation. When testing seed trays, the 72-cell seed tray, providing a

total number was 466 plant/m$^2$, and 105-cell tray (680 plant/m$^2$) both supported normal plant

growth and development.

**Development of breeding materials.** To evaluate actual application of SGV in a breeding

programme, 10 F1 seeds were obtained from a cross derived between cv. Jokyung (spring)

and cv. Joongmo2008 (winter) wheat varieties in the field in the 2018–2019 winter season.

Germinated seeds from the F1 to F5 generation were placed in the cell of a 72-cell tray (34

mL/cell) and cultivated under the speed-green vernalisation without covering soil condition

(SGV condition 4). Plants were grown under SGV for 4 weeks, followed by transfer to SB

until harvest. After harvest, seeds were dried in the glasshouse for 7 days for the next

generation. For F6 generation, 25 seeds of each line were planted in the field at the National

Institute of Crop Science, Miryang Korea (35.3$^{o}$ N; 128.5$^{o}$ E).

**Barley collection (assessed at The University of Queensland, Australia)**

**Conditions used: Barley (*Hordeum vulgare*)**

*RV–RG*: 8 h light:16 h dark 6°C into 16 h light:8 h dark 22°C

*RV–SB*: 8 h light:16 h dark 6°C into 22 h light:2 h dark 22°C:17°C

*SGV–SB*: 22 h light:2 h dark 8°C into 22 h light:2 h dark 22°C:17°C

**Experimental design and treatments.** The barley accessions were evaluated in three
experiments: 1) regular vernalisation and regular glasshouse (RV–RG), 2) regular
vernalisation and speed breeding (RV–SB), and 3) speed vernalisation and speed breeding
(SGV–SB). Under regular vernalisation conditions, plants received a standard vernalisation
treatment at 6°C for 6 weeks under 8-h-light:16-h-dark photoperiod cycles. For speed
vernalisation (SGV), plants were exposed to 8°C for 4 weeks under 22-h-light:2-h-dark
cycles. Vernalisation was performed in a fully enclosed walk-in growth cabinet fitted with
LED growth lights (Heliospectra, model E602G). For the regular vernalisation (RV), seeds
were sown directly into 100-cell trays (18 mL per cell), covered with UQ23 potting mix[15],
watered, and moved into the vernalisation chamber. For speed vernalisation (SGV), seeds
were pre-germinated in petri plates and then placed onto the surface of the potting mix.
Three seeds per accession were transplanted into a single cell of the tray for vernalisation.
To retain moisture in the cells during vernalisation, the 100-cell trays were placed inside a
sealed bottom tray with two to three small drainage holes. Trays were lightly watered daily
during the vernalisation process.

After vernalisation, the bottom trays were removed and filled with UQ23 potting mix and
Osmocote® slow release fertiliser (at a rate of 2 g per litre) to provide developing plants with
sufficient media and resources. For the regular glasshouse conditions, plants were grown in
a temperature-controlled glasshouse (22/17°C, light/dark) under a natural 12-h diurnal
photoperiod. For the speed breeding treatment, trays of barley plants were moved to a
temperature-controlled glasshouse (22/17°C, light/dark) fitted with Heliospectra LEDs using
a 22-h-light:2-h-dark photoperiod[15]. The day of anthesis for each accession was recorded as
the first spike to reach awn-peep stage (GS49).

**Benchmarking diversity of the 60 winter barley accessions.** A diverse panel of barley
accessions acquired from Australian Grains Genebank Collection (AGG, *n* = 806) was
genotyped using the Illumina Infinium–40K XT SNP chip assay (InterGrain and AgriBio-
Victoria) and returned a total of 12,599 SNP markers. Polymorphic markers with known
chromosome positions were used to investigate the genetic diversity of the barley
accessions (9,221 high-quality SNP markers with <10% missing data and <10%
heterozygosity; and 737 barley lines with <10% missing values). To investigate the
population structure, we calculated the pairwise Roger's distances between the accessions
using       'SelectionTools'       (downloadable       at       http://population-genetics.uni-

giessen.de/~software/) implemented in R. Principal coordinate analysis based on the
Roger's genetic distance matrix and k-means clustering was performed and plotted using
ggplot2[16] in R.

**Statistical analyses.** Variation in DTA for the barley accessions in each vernalisation
treatment was visualised in the form of density plots, generated using ggjoy package in R.
To determine if DTA differed across the three treatments, one-way analysis of variance
(ANOVA) was performed. Tukey's multiple comparison test (Tukey's HSD test) was then
performed to evaluate the effect of each treatment on DTA. HSD test applies appropriate
adjustments to the mean for each treatment suitable to multiple testing[17]. The analysis was
performed using (Agricolae package) in R. To investigate the relationship between barley
flowering behaviour across vernalisation treatments, the Pearson's correlation coefficient ($r$)
was calculated for DTA. The degree of correlation was also tested for significance ($P$-value;
$\alpha = 0.05$). The analysis was performed using corrgram and corrplot packages in R.

**Acknowledgements** This research was supported by the Research Program for Agricultural
Science and Technology Development (Project No. PJ011202) Rural Development
Administration. L.T.H received funding from the Australian Research Council (ARC), project
codes DP190102185 and LP170100317. Genotyping of the winter barley accessions at The
University of Queensland was funded through the Grains Research and Development
Corporation (GRDC), project code UOQ2005-012RTX. S.A was supported a GRDC
Postdoctoral Fellowship, project code UOQ1903-007RTX. L. E. D. at the University of Leeds
received funding from UKRI FLF MR/S031677/1 and the Rank Prize Funds New Lecturer
Award.

**Author contributions** L.T.H., D.S. and L.E.D. conceived and supervised the project. J.H.L.,
L.T.H., D.S. and L.E.D. designed the experiments. J.K.C. K.O.C., D.H. and K.M.K.
investigated flowering time of wheat. S.A. and E.D. investigated flowering time of barley.
J.K.C, H.P., S.M.L., Y.K. and J.M.K. developed wheat breeding materials. J.K.C., K.O.C. and
S.W.K. analysed the data. L.T.H., D.S. and L.E.D. wrote the manuscript. All the authors
discussed the results and contributed to the manuscript.

**Competing interests** The authors declare that they have no competing interests.

***Figure 1. Identifying speed vernalisation conditions for wheat (Triticum aestivum)***

Speed (green) vernalisation (SV (SGV)) combined with speed breeding (SB) accelerates winter wheat life cycles. **a**. Scheme of SV–SB conditions. Comparison of flowering time for **b**. cv. Claire and **c**. cv. Charger between wRV, SV, and SGV treatments and transfer to SB conditions following the vernalisation duration indicated. Total time to flowering is shown (including vernalisation duration); $n$ = at least 6. Fastest generation time following wRV, SV, and SGV indicated by the time the first plant flowered following transfer to SB for **d**. cv. Claire and **e**. cv. Charger. **f**. Comparison of seed treatments with example images taken when the first plant flowered under T4 conditions for the winter wheat cv. Keumgang. **g**. Flowering time following the four seed treatments (T1-4 in f) in cv. Keumgang; $n$ = 7. **h**. Comparison of plant development between SV and SGV. **i**. Comparison of flowering time for cv. Hereward between wRV and SGV; $n$ = at least 6. Flowering times for different durations **j**. and temperatures **k**. for cv. Keumgang under SGV. Significance is shown according to Student's $t$-test $P$ < 0.05. SV (speed vernalisation), SGV (speed-green vernalisation), wRV (warm regular vernalisation), SB (speed breeding), and vernalisation duration are included in days to flowering.

**Figure 2. Speed vernalisation conditions for barley (Hordeum vulgare) and cereal population development**

Flowering time responses of diverse winter barley accessions following various vernalization conditions. **a**. Density plots displaying days to anthesis (DTA) for 60 diverse winter barley accessions evaluated using three treatments: 1) speed vernalisation (at 8°C) and speed

breeding conditions (SGV–SB), 2) regular vernalisation and speed breeding conditions (RV– SB), and 3) regular vernalisation and regular glasshouse conditions (RV–RG). **b**. Example image of barley population following SGV–SB. **c**. Population development timeline for winter x spring wheat population over six generations and **d**. flowering comparison between SB and SGV–SB for spring x winter population. **e.** An example breakdown of growth cycle under SGV–SB conditions for the same winter x spring cross in **c** and **d**. **f.** Projected generation times for different cereal types and growth conditions: A = SB Spring barley (e.g., cv. Commander, Golden Promise), B = SB Spring wheat cv. Suntop, Cadenza, C = SGV_SB Winter barley, D = SGV_SB Wheat winter x spring (using data from d), E = SB wheat winter x spring (using data from d), F = SGV–SB winter wheat cv. Claire, G = SV–SB winter wheat cv. Claire, H = SGV–SB winter wheat cv. Charger, I = SV–SB winter wheat cv. Charger, J = wRV–RG winter wheat cv. Charger, and K = wRV–RG cv. Claire. Spring generation times calculated from published data[2]. SV (speed vernalisation), SGV (speed-green vernalisation), wRV (warm regular vernalisation), SB (speed breeding).

**Figure 3. Vernalisation-related gene expression under SV and SGV conditions**

In cv. Claire, **a**. Comparison of expression of *VRN1* between wRV (green) and SV (pink) conditions. **b**. Comparison of expression of *FT1-B* between wRV (green) and SV (pink) conditions. **c**. Expression of *VRN1* (green) and *FT1-B* (red) between 6 weeks of SV and then 6 weeks of SV and 1 week in SB. **d**. Expression of *ZCCT1* and *ZCCT2* between 6 weeks of

SV and then 6 weeks of SV and 1 week in SB. Representative images following different
vernalisation treatments. For cv. Claire following SV_SB with 3 weeks **e**., 4 weeks **f**., and 5
weeks **g**. vernalisation treatment. Comparison between wRV and SGV conditions for cv.
Claire following 4 weeks vernalization and photographed at the same age **h**. and cv.
Hereward following 6 weeks vernalization and photographed at the same age **i**. All
conditions result in vernalised plants. Scale bar is 30 cm. In cv. Hereward, expression of
*VRN1*, *ZCCT1*, *ZCCT2*, and *FT-1B* for **j**. SV and **k**. SGV. All show three biological replicates
that each comprise at least three plants, with standard error of the mean. SGV (speed-green
vernalisation), wRV (warm regular vernalisation), duration in weeks refers to vernalisation
duration experienced.

414
415
416

light:16 h dark) photoperiod and transferred at the weeks indicated on the x-axis to long-day
(16 h light:8 h dark) glasshouse conditions with constant 20–22°C. Vernalisation duration is
included in the flowering time. $n$ = at least 3, NF = No flowering by date indicated. **c** - **f** Apex
lengths from the same conditions; $n$ = 3. Apex length following SV for cultivars **g**. Buster, **h**.
Charger, **i**. Hereward, and **j**. Claire; $n$ = 3 with representative images between weeks 2 and
7 of SV for each cultivar, scale bar represents 1000 µm, for cultivars **k**. Buster, **l**. Charger,
**m**. Hereward, and **n**. Claire. Spikelet counts from plants moved following 2, 4, or 6 weeks of
vernalisation for cv. Claire for **o**. wRV_SB, **p**. SV_SB, and **q**. SGV_SB. $n$ = at least 8.

**Extended data figure 2**: An example of SGV_SB conditions using LED lighting. **a.** Representative image of plants in SGV conditions with **b.** showing the spectral measurement of light composition in SGV. **c.** Representative image of plants in SB conditions and **d.** showing the spectral measurement of light composition SB. Spectral measurements were taken using the RS-3500 spectroradiometer from spectral Evolution Inc. X-axis values are wavelengths in nanometres, and y-axis represents proportion (1 unit = 0.1 proportion). Under SGV–SB conditions, wheat plants followed standard development shown over a 2- to 6-week vernalisation treatment for cv. Sturdy: **e.** Growth stage 59, **f.** days until GS32, **g.** plant height for cv. Keumgang, **h.** growth stage 59, **i.** days until GS32, **j.** plant height for cv. Sturdy, **k.** Spike length, **l.** tiller number per plant, and **m.** grain number per spike; and for cv. Keumgang: **n.** spike length, **o.** tiller number per plant, and **p.** grain number per spike, where $n = 6$.

**Extended data figure 3**: Accelerated generation times also achieved for barley and breeding populations. **a**. Principal component analysis (PCA) of barley cultivars used in Figure 2. **b**. Spectral measurement of light composition during SGV for barley and **c**. flowering distribution for three additional spring x winter or winter x winter populations following SGV–SB; crosses 1 and 2 are spring x winter populations, and cross 3 is a winter x winter population.

**References**

Kim, D. H., Doyle, M. R., Sung, S. & Amasino, R. M. Vernalization: winter and the timing of flowering in plants. *Annu Rev Cell Dev Biol* **25**, 277-299, doi:10.1146/annurev.cellbio.042308.113411 (2009).

Watson, A. *et al.* Speed breeding is a powerful tool to accelerate crop research and breeding. *Nat Plants* **4**, 23-29, doi:10.1038/s41477-017-0083-8 (2018).

Hickey, L. T. *et al.* Breeding crops to feed 10 billion. *Nat Biotechnol* **37**, 744-754, doi:10.1038/s41587-019-0152-9 (2019).

Dixon, L. E. *et al.* VERNALIZATION1 controls developmental responses of winter wheat under high ambient temperatures. *Development* **146**, doi:10.1242/dev.172684 (2019).

HUNT, W. Y. a. L. A. Reanalysis of Vernalization Data of Wheat and Carrot. *Annals of Botany* **84**, 615-619 (1999).

Duncan, S. *et al.* Seasonal shift in timing of vernalization as an adaptation to extreme winter. *Elife* **4**, doi:10.7554/eLife.06620 (2015).

7    Diaz, A., Zikhali, M., Turner, A. S., Isaac, P. & Laurie, D. A. Copy number variation affecting
the Photoperiod-B1 and Vernalization-A1 genes is associated with altered flowering time in
wheat (Triticum aestivum). *PLoS One* **7**, e33234, doi:10.1371/journal.pone.0033234 (2012).
8    Hepworth, J. *et al.* Absence of warmth permits epigenetic memory of winter in Arabidopsis.
*Nat Commun* **9**, 639, doi:10.1038/s41467-018-03065-7 (2018).
9    Yan, L. *et al.* The wheat VRN2 gene is a flowering repressor down-regulated by vernalization.
*Science* **303**, doi: 10.1126/science.1094305 (2004).
10   Greenup, A. G. *et al.* ODDSOC2 is a MADS box floral repressor that is down-regulated by
vernalization in temperate cereals. *Plant Physiol* **153**, 1062-1073,
doi:10.1104/pp.109.152488 (2010).
11   Xie, L. *et al.* TaVrt2, an SVP-like gene, cooperates with TaVrn1 to regulate vernalization-
induced flowering in wheat. *New Phytol* **231**, 834-848, doi:10.1111/nph.16339 (2021).
12   Chen, M. *et al.* Maternal temperature history activates Flowering Locus T in fruits to control
progeny dormancy according to time of year. *Proc Natl Acad Sci U S A* **111**, 18787-18792,
doi:10.1073/pnas.1412274111 (2014).
13   Schneider. NIH Image to ImageJ: 25 years of Image Analysis. *Nature Methods* **9**, 671-675
(2012).
14   TOTTMAN, D. R. The decimal code for the growth stages of cereals, with illustrations.
*Annuals Applied Biology* (1987).

15   Ghosh, S. *et al.* Speed breeding in growth chambers and glasshouses for crop breeding and
model plant research. *Nat Protoc* **13**, 2944-2963, doi:10.1038/s41596-018-0072-z (2018).
16   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis.*, (Springer-Verlag, 2016).
17   Rogan, J. C. a. K., H.J. . Is the ANOVA F-Test Robust to Variance Heterogeneity When Sample
Sizes Are Equal? An Investigation via a Coefficient of Variation. *American Educational
Research Journal* **14**, 493-498 (1977).