Community Composition of Bacteria Isolated from Swiss Banknotes Varies Depending

on Collection Environment

Authors: Anna M. Bischofberger[1†], Alex R. Hall[1]

Contact information: [1]Institute of Integrative Biology, ETH Zurich, Switzerland

[†]anna.bischofberger@env.ethz.ch

Abstract:

Humans interact constantly with surfaces and associated microbial communities in the

environment. The factors shaping the composition of these communities are poorly understood:

some proposed explanations emphasize the influence of local habitat conditions (niche-based

explanations), while others point to geographic structure and the distance among sampled

locations (dispersal-based explanations). However, the relative roles of these different drivers

for microbial community assembly on human-associated surfaces are not clear. Here, we used

a combination of sampling, sequencing (16S rRNA) and culturing to show that the composition

of banknote-associated bacterial communities varies depending on the local collection

environment. Using banknotes collected from various locations and types of shops across

Switzerland, we found taxonomic diversity dominated by families such as *Pseudomonadaceae*,

Staphylococcaceae and *Streptococcaceae*, but with banknote samples from particular types of

shops (especially butcher shops) having distinct community structure. By contrast, we found

no evidence of geographic structure: similarity of community composition did not decrease

with increasing distance among sampled locations. These results show that microbial

communities associated with banknotes, one of the most commonly encountered and

exchanged human-associated surfaces, can reflect the local environmental conditions (in this

case, the type of shop), and the signal for this type of variation was stronger than that for

geographic structure among the locations sampled here.

Keywords:

microbiome, banknote, community assembly, 16S rRNA

Introduction

Humans encounter microbial communities constantly and we now know that they contribute,

among other things, to food and drink quality (Wareing, 2016; Reese *et al.*, 2020), soil fertility

(Fierer, 2017) and human health (Libertucci and Young, 2019). Understanding variation of

microbial community structure (relative abundances of different taxa) is therefore an important

challenge. As for other organisms and in the wider field of community ecology, the

contributions of different drivers of microbisl community assembly remain unclear (Chesson,

2000; Ebach, 2015). Selection imposed by local habitat conditions and biotic interactions

(niche-based explanations invoking deterministic factors) (Macarthur and Levins, 1967;

Diamond, 1975) is expected to result in communities matched to their local environments.

Other possible drivers include more stochastic processes, ecological drift or dispersal limitation

(Hubbell, 2011), which may result in geographic structuring of communities independent from

the effects of local habitat conditions. There is evidence from microbes and other taxa

supporting both types of explanation (Webb; Finlay, 2002; Fargione *et al.*, 2003; Whitaker *et*

al., 2003a; Horner-Devine *et al.*, 2004; Zhou *et al.*, 2013), and it is increasingly recognised that

multiple drivers are in play (Chase, 2003; Chase and Myers, 2011; HilleRisLambers *et al.*,

2012). However, we still lack a clear picture of their relative roles (Chase, 2010; Hanson *et al.*,

2012; Stegen *et al.*, 2015; Zhou and Ning, 2017). This problem is especially acute for microbial

communities on surfaces that humans interact with in daily life. For example, a recent global

study of microbial communities on metro and transit systems identified city-specific

community composition linked to local conditions, albeit with a shared core microbiota across

locations (Danko *et al.* 2021). However, it is not yet clear whether such biogeographical

structuring is also evident on other human-associated surfaces, or how strongly such

microbiomes reflect local habitat conditions (selection) as opposed to other predictors such as

geographic distance among sites.

One of the most important human-associated surfaces is cash money: few everyday objects

are as frequently, widely and actively exchanged among humans as banknotes. Paper-based

money has been in use for hundreds of years and is now commonplace around the world.

Despite the advent of credit cards and other cashless payment instruments, banknotes and coins

are still the most common payment option for purchases at the physical point of sale and for

person-to-person payments in the EU and Switzerland (Gehring *et al.*; Central Bank, 2020).

The rough surface of most banknotes lends itself well to the attachment of microorganisms

(Jalali *et al.*, 2015; Vriesekoop *et al.*, 2016), something first investigated in the late 19th century

(Schaarschmidt, 1884). Since then, tools such as metagenomic sequencing have enabled more

detailed description of microorganisms on money, including characterization of communities

on dollar bills in New York City (Maritz *et al.*, 2017), identification of pathogens and antibiotic

resistance genes on Rupees in New Delhi (Jalali *et al.*, 2015), and monitoring survival of

influenza viruses on banknotes (Thomas *et al.*, 2008). However, to our knowledge it remains

poorly understood whether banknotes collected in different places harbour microbial

communities that reflect their local environment, which would indicate a role for selection

(Hanson *et al.*, 2012; Zhou and Ning, 2017), and/or show biogeographic structuring (e.g.,

increasing dissimilarity with geographic distance).

Here, we aimed to quantify variation of the bacterial microbiome associated with banknotes

originating from different locations across a country, and to test whether this variation can be

explained by geographic distances among locations, or other characteristics of the local

sampling environments. We did this by collecting banknotes from each of ten locations in

Switzerland (Fig. 1A), visiting five different types of shops in each location. We hypothesized

community composition would vary among banknotes collected from different types of shops

(if there is an effect of local habitat selection on the monetary microbiome), and/or that

banknotes collected from more distant locations would harbour more dissimilar communities

(if there is biogeographic structuring in general). We studied the samples isolated from the 50

banknotes by (1) 16S rRNA sequencing to determine community composition, including viable

and non-viable, culturable and non-culturable bacteria, and (2) plating on chromogenic agar to

detect viable, culturable bacteria. Our results show that the composition of bacterial

communities isolated from banknotes – determined by both sequencing and by plating – varies

depending on the local environment (shop type) in which banknotes are collected. By contrast,

we find no evidence for geographic structuring of bacterial community composition across

banknotes sampled in Switzerland.

<u>Materials and Methods</u>

Sample Collection

We collected banknotes from ten locations in Switzerland. In choosing locations, we adhered

to several criteria: (1) countrywide distribution of sampled locations (Fig. 1A; (2) fewer than

10'000 inhabitants per location (because we expected any signal for geographic structuring

would be stronger among smaller towns); (3) all of the targeted shop types (bakery, butcher,

florist, kiosk/newsagent, and pharmacy) present at each location; (4) no significant commuter

traffic between locations (according to *Erwerbstätige nach Wohn- und Arbeitsgemeinde, 2014*

und 2018 by the Federal Office of Statistics; again we expected geographic structuring of

microbial communities would be stronger among less connected towns). The chosen locations

were (Fig. 1A): Biasca (BIA), Brienz BE (BRI), Frick (FRI), Klosters-Serneus (KLO),

Langnau am Albis (LAN), Orbe (ORB), Porrentruy (POR), Rorschach (ROR), Stein am Rhein

(STE), Visp (VIS). We chose the five shop types in an attempt to represent different

environments, products and clientele. More details on locations and shops can be found in

supplementary files SuppFile_01 and SuppFile_02, respectivley.

On arrival in a location, we collected the banknotes from the different shops as follows:

donning sterile gloves before entering the shop; choosing an item for less than CHF 10; paying

the item with a CHF 20 banknote; folding the CHF 10 banknote received as change in half

twice and collecting it in a sterile Nasco Whirl-Pak® sample bag (product nr. 11701973; Fisher

Scientific AG, Reinach, Switzerland); storing the sample bag in a paper envelope. Once we

had the five banknotes of a given location, we transported the banknotes back to the laboratory

and did the initial processing as described below. All samples were collected by the same

person between March 12th and March 26th, 2021. Samples from different locations were

collected on different days. One sample (butcher-BIA) was classified as "butcher" even though

it came from a supermarket with a served meat counter (both butchers in Biasca (BIA) were

closed on collection day); this is discussed further below. We contacted all shop owners via

email to inform them about our project and gave them the option to have the banknote collected

in their shop removed from the study. No shop owner vetoed inclusion of the banknote

collected in their shop in the study.

Initial Sample Processing and Plating

We performed initial processing for all banknotes on the day of collection as follows: after

removing the banknote from the sample bag and placing it on a sterile surface, we swabbed the

front of the banknote (not exposed to the bag surface) along the width from top to bottom

(Maritz *et al.*, 2017), using a sterile cotton swab (product nr. 80.628; Sarstedt, Nürnberg,

Germany) moistened with phosphate-buffered saline (PBS; product nr. P4417-100TAB;

Sigma-Aldrich, Merck KgaA, Germany). Turning the banknote 90 degrees, we repeated the

procedure. With the same swab, we swabbed the back of the banknote according to the same

protocol. We cut the tip of the swab with sterile cissors and placed it in an 2mL Eppendorf tube

filled with 950μL PBS. We repeated the procedure with the other four banknotes, sterilizing

the surface between samples. As a negative control, we swabbed the interior of a sample bag

and followed the protocol just described. We vortexed all Eppendorf tubes for 5min at the

highest amplitude and removed the swab tips with sterile tweezers, squeezing out the remaining

liquid.

To screen for viable and culturable cells, we plated a 50μL aliquot of each sample on a

separate Chromatic™ MH agar plate (product nr. LF-611618; Apteq AG, Cham, Switzerland).

Chromatic™ MH agar is a chromogenic agar that allows for the identification of bacteria from

environmental and clinical isolates by differential colouring of colonies from different species.

We incubated plates at 37°C for 24h and stored the remaining samples at -20°C until further

processing as described below. Note that plating on chromogenic agar provides a relatively

coarse-grained identification and only for certain taxa, but we employed it here only to test (1)

for the presence of viable and culturable (i.e., colony-forming) cells and (2) whether the

putative species identities of resulting colonies were consistent with taxa identified by sequence

analysis. More details on colony morphology and colour can be found in supplementary file

SuppFile_03.

Genetic Analysis and Bioinformatics

We used the DNeasy® PowerLyzer® PowerSoil® Kit (product nr. 12855-100; Qiagen, Hilden,

Germany) to extract bacterial DNA from thawed samples, according to the manufacturer's

protocol with slight adaptations. In brief: For each sample, we added 700μL to a PowerBead

tube (positive control: 700μL PBS plus *Escherichia coli* colony from agar plate; negative

control: from empty sample bag, see above) and centrifuged all PowerBead tubes at 13'000rpm

for 10min at 4°C. Discarding the supernatant, we dissolved the cell pellet by adding 750μL

PowerBead solution and 60μL Solution C1, and inverting the tubes 10 times. After incubating

the tubes at 65°C for 10min followed by incubation at 95°C for 10min, we performed bead

beating with a Bead Ruptor 24 (Omni International, Inc, Kennesaw (Georgia), United States

of America (USA)) with the following settings: cycle nr = 2, strength = 5.5, time = 45s, interval

= 0. After transferring the supernatant to clean 2mL collection tubes, we added 250μL Solution

C2, vortexed the tubes for 5s at the highest amplitude and incubated them at 4°C for 5min.

After centrifugation at 13'000rpm for 1min at 4°C, we transferred 600μL supernatant into clean

2mL collection tubes and added 200μL Solution C3. We vortexed tubes briefly at the highest

amplitude and incubated them at 4°C for 5min, followed by centrifugation at 13'000 rpm for

1min at 4°C. Avoiding the pellet, we transferred 750μL supernatant into clean 2mL collection

tubes, added 1200μL well-mixed Solution C4 and vortexed tubes for 5s at the highest

amplitude. We loaded 675μL sample onto MB spin Columns, centrifuged them at 13'000rpm

for 1min at 4°C and discarded the flow-through. We repeated this until the entire sample was

processed. After adding 500μL Solution C5, we centrifuged samples at 13'000rpm for 1min at

4°C and discarded the flow-through, followed by centrifugation at 13'000rpm at 4°C for 30s.

We placed the MB Spin Columns into clean 2mL collection tubes and added 70μL Solution

C6 to the centre of the filter membrane. Following incubation at 65°C for 5min and

centrifugation at 13'000rpm at 4°C for 1min, we discarded the MB Spin Columns and stored

samples at -20°C. The 50 samples were processed in three batches (consisting of randomly

selected samples) on different days, with a positive and negative control for each batch. After

sequencing, we did not detect an effect of extraction batch on either read counts per sample

(ANOVA: $F_{2,47} = 2.521$, $p = 0.091$) or community composition (PERMANOVA: $F_{2,41} = 1.711$,

$p > 0.1$).

We quantified the obtained DNA using the Quant-iT™ dsDNA BR Assay Kit (product nr.

Q33130; Thermo Fisher Scientific, Waltham, USA) in the Spark® 10M multimode microplate

reader (Tecan Group Ltd, Männedorf, Switzerland). We sequenced at the Genetic Diversity

Centre, Zurich, Switzerland, using the MiSeq Reagent kit v3 (600 cycle PE) (Illumina, San

Diego, USA) after library preparation with the Nextera XT 96 Index kit v2, Set D (Illumina,

San Diego, USA). The sequences of the four sets of primers used during library preparation

(limited cycle PCR) are listed in supplementary table 1.

If not noted otherwise, raw sequence data were processed with USEARCH

(v11.0.667_i86linux64; Edgar (2010)). After end trimming reads (if needed) and merging them

into amplicons (commands: *filter_phix*, *filter_lowc* (threshold 25), *fastx_truncate* (trim R1/R2:

25/50), *fastx_syncpairs*, *fastq_mergepairs* (min. overlap: 30; min. identity: 60%; min. merged

length: 100; min. merged quality: 8)), we trimmed reads for primers (command: *search_pcr*

(amplicon size range 100-600; number of mismatches: 2; coverage: full-length (no end gaps)).

We used PRINSEQ-lite 0.20.4 to check quality and size filter (parameters: size range: 150-

550; GC range: 30-70; min. Q mean: 20; number of Ns: 1; low complexity: dust / 30). We

clustered operational taxonomic units (OTUs) at 97% using the UPARSE-OTU algorithm

(number of OTUs: 2378), and used UNOISE3 algorithm to perform denoising (error-

correction) of amplicon sequence variants (zero-radius OTUs, number of ZOTUs: 4464), and

to cluster ZOTUs at different identity levels (number of ZOTUs 99%: 3148; number of ZOTUs

98%: 2513; number of ZOTUs 97%: 2039). OTU annotation was carried out using SINTAX

(reference: SILVA_128_16S_utax_work.fa; tax filter: 0.75).

After removing positive and negative controls and mitochondrial and chloroplast sequences,

and before filtering, denoised sequencing data (ZOTUs) consisted of 50 samples with

14'900'848 raw reads in total, ranging from four to 6'752'341 raw reads per sample (mean

number of raw reads: 266'086.57; median number of raw reads 158'243). We filtered data by

removing samples with low number of raw reads (<1000) and ZOTUs with total abundance

lower than 0.1% of total depth, ending up with 42 samples with 10'136'186 filtered reads in

total, ranging from 138 filtered reads per sample for butcher-BRI to 6'167'513 filtered reads

per sample for florist-BRI (mean number of filtered reads = 241'337.76; median number of

filtered reads = 88'192; Fig. S1). We had at least three samples per location and at least six

samples per shop type.

The sequencing data have been deposited in the European Nucleotide Archive under the

study accession number PRJEB45390 (https://www.ebi.ac.uk/ena; data will be made

accessible upon publication of the study).

Statistics

We used denoised sequences, i.e. ZOTUs, and conducted all statistical analysis in R (version

4.1.2).

To quantify within-sample diversity, we used the alpha() function in the microbiome

package (version 1.16.0) on filtered data to calculate several different diversity indices

(Shannon diversity index; Gini-Simpson index; observed richness; Chao1 index). The Shannon

diversity index H takes into account both richness and evenness; the Gini-Simpson index gives

the probabilty that two random draws from the same sample (with replacement) are not from

the same type (here, the same family); the observed richness counts the number of unique

species (ZOTUs) present in the sample; Chao1 index is a non-parametric estimator of species

richness that assumes Poisson distribution of the data (Chao, 1984; Chao and Bunge, 2002).

We used analysis of variance (ANOVA) to test whether within-sample diversity varied among

locations or shop types (aov() function in the stats package (R version 4.1.3)).

Before analysing between-sample diversity and to account for different library sizes (Weiss

et al., 2017), we normalized filtered data with a regularized-logarithm (rlog() function in the

DESeq2 package (version 1.34.0)), after comparing the two available transformations offered

in DESeq2 package, regularized-logarithm transformation and variance-stabilizing

transformation. We chose the regularized-logarithm transformation because it (1) is

recommended for smaller datasets and for cases with large ranges of sequencing depths across

samples (Love *et al.*, 2014); (2) it performed better in the meanSdPlot generated.

As a first test of whether samples from different locations and shop types differed in terms

of bacterial community structure, we used permutational multivariate analysis of variance

(PERMANOVA; Anderson, 2001; Zapala and Schork, 2006) with the adonis2() function in the

vegan package (version 2.5-7), with filtered-and-normalized data and using Bray-Curtis

dissimilarity (vegdist() function) to measure differences among sampled communities. Next,

to visualise any clustering among samples in terms of community composition, we performed

ordination of filtered-and-normalized data with the ordinate() function in the phyloseq package

(version 1.38.0), with non-parametric multidmensional scaling (*method* = "NMDS") and Bray-

Curtis dissimilarity (*distance* = "bray") as input parameters. We used NMDS as the ordination

method because we obtained a stress value of 0.11 for our data with NMDS ordination; stress

values <0.2 generally indicate good fit (Clarke, 1993). We visualised ordination results with

the plot_ordination() function in the phyloseq package (version 1.38.0). We used a third type

of analysis as an additional test for geographic structuring of microbial community

composition, by conducting a Mantel test. This allowed us to account explicitly for the physical

distance between individual sampling locations (shops), rather than testing for average

differences among locations (as above in PERMANOVA). We did this using the mantel.test()

function in the ape package (version 5.5), with the vegdist() function in the vegan package

(version 2.5-7) for Bray-Curtis dissimilarity of filtered-and-normalized data. We used the

geodist() function in the geodist package (version 0.0.7) to calculate geographical distance (in

metres) between individual sampling locations. We used the same functions when repeating

the analysis for sample sets from individual shop types; here and elsewhere, we accounted for

multiple testing using the Holm–Bonferroni method (sequential Bonferroni).

The core microbiome is defined as the species present above a certain threshold and shared

among a set of samples (Salonen *et al.*, 2012; Shade and Handelsman, 2012). We determined

the core microbiome, based on ZOTUs, for each shop type as follows: Using filtered-and-

normalized relative abundance data (transform() function in the microbiome package (version

1.16.0; "compositional" as input for *transform* argument) and subsetting data by shop type, we

applied the core_members() function in the microbiome package (version 1.16.0; *detection* =

0.001 and *prevalence* = 0.5) to filter out core ZOTUs for each shop type. Then, as a fourth

analysis of how microbiome structure varies among shop types/locations, we drew a Venn

diagram with the venn() function in the eulerr package (version 6.1.0) and determined ZOTU

identity with the format_to_besthit() function in the microbiomeutilities package (version

1.00.16), followed by identification of shared ZOTUs with reduce() and intersect() functions

(base package, R version 4.1.2).

To test for an association between the abundance of viable-and-culturable bacteria

(presence/absence of colonies and number of colonies per plate, after plating samples on

chromatic agar) and shop type and location we conducted hurdle regression for count data via

maximum likelihood (hurdle() function in pscl package (version 1.5.5), with formula

number.colonies ~ location or *number.colonies ~ shop.type* and negative binomial regression

(with log link) as count model). As our data contained several samples with zero viable cells

we adjusted the count table with a dummy variable (dummy species present at count 1 in all

samples; Clarke *et al.*, 2006) to perform further analysis, being aware that we cannot be sure

that all empty samples had zero viable cells for the same reason. We used the adjusted count

table to perform PERMANOVA as described above.

The map of Switzerland with locations (Fig. 1A) was made with the mun.plot() function in

RSwissMaps package (version 0.1.0.1) with the following arguments (where relevant): *bfs_id*

= dt$bfs_nr, *year* = 2016, *boundaries* = "c". Other packages and functions used in data

processing and drawing figures: cowplot (version 1.1.1); dplyr (version 1.0.7); ggplot2 (version

3.3.5); ggpubr (version 0.4.0); microbial (version 0.0.20); ochRe (version 1.0.0);

phyloseq.extended (version 0.1.1.6); phylosmith (version 1.0.6); ranacapa (version 0.1.0); yarrr

(version 0.1.5); VennDiagram (version 1.7.0); vsn (version 3.62.0).

Results

Communities Isolated from Banknotes Include *Pseudomonadaceae*, *Streptococcaceae*,

and *Staphylococcaceae*

We detected some families of bacteria (Fig. 1B) in every banknote sample: *Pseudomonadaceae*

and *Staphylococcaceae* were present in all samples (42/42, 100%; Table S2). These are both

diverse families that include species closely associated with humans, e.g. as part of the human

skin, respiratory tract or gut microbiome (Cogen *et al.*, 2008; Fierer *et al.*, 2008; Murphy and

Parameswaran, 2009; Kim *et al.*, 2013; Fourquin-Gomez *et al.*, 2014; Robert-Pillot *et al.*, 2014;

Rock and Donnenberg, 2014; Delanghe *et al.*, 2021; Skowron *et al.*, 2021), but have also been

isolated from animal hosts and soil and aquatic environments (Cousin, 1999; Madhaiyan *et al.*,

2020). Other families were detected in significant abundance, but only in subsets of samples,

such as *Vibrionaceae* and *Lactobacillaceae* (in 21 and 30 samples out of 42 respectively). The

same taxa resurfaced when we calculated the core microbiome for each shop type (taxa found

in >50% of samples per shop type at abundance >0.1%). We identified seven core ZOTUs

shared between all shop types, attributed to the genera: *Corynebacterium*, *Propionibacterium*,

Pseudomonas, *Streptococcus* (n = 2) and *Staphylococcus* (n = 2) (Fig. S2; detection threshold

= 0.001, prevalence threshold = 0.5). Thus, our sequence data revealed diverse microbial

communities associated with Swiss banknotes, but with the relative abundances of individual

taxa varying among samples.

Within-sample diversity estimated by Shannon diversity index (Fig. 2) did not vary

significantly with either geographic location or shop type (ANOVA: effect of shop type – $F_{4,37}$

$= 0.844$, $p = 0.506$; effect of location – $F_{9,32} = 1.013$, $p = 0.45$). We obtained the same

qualitative result with an alternative diversity index (Gini-Simpson index), and the two

diversity indices were highly correlated with each other (linear regression Gini-Simpson ~

Shannon: adjusted $R^2 = 0.851$). Similarly, within-sample species richness (taken as observed

richness (Fig. 2B) or Chao1 index) did not vary significantly among locations or shop types

(ANOVA: $p > 0.1$ for shop type and location effects for both measures) and was correlated

with Shannon diversity (adjusted $R^2 = 0.629$ for observed richness and $R^2 = 0.478$ for Chao1).

Thus, within-sample diversity did not vary significantly among shop types or geographic

locations.

Community Composition Was Specific to Shop Type, but Not Location

Bacterial community composition was more similar among banknotes collected from the same

shop type than from different shop types (PERMANOVA: effect of shop type – $F_{4,41} = 1.321$,

$p = 0.03$). Visualising the similarities/differences among samples by ordination analysis (non-

metric multidimensional scaling NMDS) further supported variation of community

composition among shop types (Fig. 3). In particular, several samples from butcher shops

clustered together in the top-right corner of the ordination plot (Fig. 3). Note one sample

classified as "butcher" but separate from this cluster (butcher-BIA) was collected from a

supermarket with a served meat counter but separate cash desk (see Methods). Consistent with

the variation among shop types being primarily driven by the distinctiveness of butcher shop

samples, if we excluded butcher samples from the analysis we no longer found significant

variation among shop types (PERMANOVA: effect of shop type, excluding butcher samples

– $F_{3,32} = 0.949$, $p = 0.584$); this was not the case when re-running the analysis excluding sample

sets from all other shop types, except for florist samples which resulted in marginal non-

significance (PERMANOVA: effect of shop type, without florist samples – $F_{3,32} = 1.287$, $p =$

0.056).

Looking at the ZOTUs making up the core microbiome for each shop type again supported

butcher samples as being distinctive: the set of banknotes from butchers (n = 9) had the greatest

number of core ZOTUs, both in total and unique to a single shop type (31 core ZOTUs in total,

13 unique core ZOTUs; Fig. S2). The 13 core ZOTUs unique to butchers belonged to the

following genera: *Photobacterium* (n = 3; family *Vibrionaceae*), *Streptococcus* (n = 2; family

Streptococcaceae), *Brochothrix* (n = 1; family *Listeriaceae*), *Corynebacterium* (n = 1; family

Corynebacteriaceae), *Kocuria* (n = 1; family *Micrococcaceae*), *Lactobacillus* (n = 1; family

Lactobacillaceae), *Lactococcus* (n = 1; family *Staphylococcaceae*), *Novosphingobium* (n = 1;

family *Sphingomonadaaceae*), *Staphylococcus* (n = 1; family *Staphylococcaceae*), *Veillonella*

(n = 1; family *Veillonellaceae*). This is consistent with our family-level analysis (Fig. 1), where

some of the families to which these butcher-associated genera belong were over-represented

among butcher samples (e.g., *Vibrionaceae*, *Lactobacillaceae*).

A comparison between core ZOTUs unique to butcher samples and NMDS loadings (Fig.

3) provides further information about the taxa that drive the among-shop-type variation: the

Veillonella core ZOTU and one *Streptococcus* core ZOTU unique to butchers were among the

ten ZOTUs with the highest positive loadings for NMDS1; the *Lactococcus* and three

Photobacterium core ZOTUs unique to butchers were among the ten ZOTUs with the highest

positive loadings for NMDS2 (again consistent with families *Vibrionaceae* and

Lactobacillaceae being abundant and over-represented in butcher samples in Fig. 1). Note that

the observed variation in community composition among shop types was not explained by

variable numbers of reads among samples from different shop types (Fig. S1; effect of shop

type in ANOVA – $F_{4,45} = 0.792$, $p > 0.5$).

We found no evidence that samples collected in different locations (towns) harboured

different bacterial microbiomes (PERMANOVA: effect of location – $F_{9,41} = 1.173$, $p = 0.067$;

Fig. 3). In a second test for geographic structuring of the sampled communities (Mantel test),

we quantified the association between pairwise similarity among samples in terms of

community composition and the geographic distances between sampling locations. This

revealed no significant association between the two distance measures, neither in the full

dataset (n = 42; Mantel test – $z = 39'978'433$, $p = 0.751$), nor when performed separately for

each shop type (Mantel test, five tests total with p-values corrected for multiple testing – p_{bakery}

= 1, $p_{butcher}$ = 1; $p_{florist}$ = 0.315; p_{kiosk} = 1; $p_{pharmacy}$ = 1). As for shop type above, there was no

association between number of reads and location (ANOVA: effect of location – $F_{9,40} = 0.836$,

$p > 0.5$). Thus, unlike for the variation among shop types observed above, we found no

evidence that the differences in bacterial community composition among samples increased

with geographic distance.

Cultivation of Samples on Agar Supports Conclusions Drawn from Sequencing Data

When we plated aliquots of samples on Chromatic^{TM} MH agar plates, we detected a total of

157 viable colonies across 24 of 50 plates (48%). The number of colonies per plate varied

among samples (Fig. 4), and was highest on average for butcher samples (count model in hurdle

regression model for shop type: butcher: $z = 3.44$, $p < 0.001$; all other shop types: $p > 0.05$;

Fig. 4). There was no significant variation of the number of colonies per plate among

geographic locations (count model in hurdle regression model for location: $p > 0.05$ for all

locations). Several plates produced no colonies, but presence/absence of viable colonies was

not predicted by either shop type or location (zero hurdle model in hurdle regression model for

shop type and for location: $p > 0.05$ for all shop types and all locations).

The identity of colonies (as determined by colony morphology and colour on Chromatic™

MH agar plates, see Methods) corresponded to key taxa identified by sequencing: *E. coli* =

Enterobacteriaceae; *Staphylococcus aureus* = *Staphlyococcaceae*; *Pseudomonas spp* =

Pseudomonadaceae; *Klebsiella spp.*, *Enterobacter spp.*, *Serratia spp.* = *Enterobacteriaceae*

(Fig. 1B Fig. 4). The exception to this trend was *Enterococcus faecalis* (3/157 colonies or 2%),

which belongs to the *Enterococcaceae* family. This was not among the most abundant families

in our sequence data (Fig. 1B), and we only found ZOTUs categorized as *Enterococcaceae*

prior to filtering the data, indicating it was present but at lower abundances than suggested by

colony formation data. For most of the other key taxa identified by sequence data that we did

not find on agar plates, prior information about these taxa suggests the growth conditions

(Chromatic™ MH agar; 37°C; 24h aerobic incubation) are not conducive to colony formation.

For example, *Corynebacteriaceae, Lactobacillaceae*, (environmental) *Moraxellaceae*, and

Vibrionaceae have growth optima <37°C (De Angelis and Gobbetti, 2011; Vinet and

Zhedanov, 2011; Lonvaud-Funel, 2014; Tauch and Sandbote, 2014; Baron *et al.*, 2020). In

addition, several of the key taxa (e.g. *Brevibacteriaceae, Corynebacteriaceae*

Halomonadaceae, *Lactobacillaceae*, *Moraxellaceae*, *Vibrionaceae)* may have required

incubation times >24h (Vinet and Zhedanov, 2011; Argandoña *et al.*, 2012a; Forquin and

Weimer, 2014; Tauch and Sandbote, 2014; De Angelis and Gobbetti, 2016; Baron *et al.*, 2020),

or different media composition such as higher salt or nutrient content (*Halomonadaceae* and

Corynebacteriaceae, respectively, (Argandoña *et al.*, 2012b; Tauch and Sandbote, 2014)).

Therefore, the absence of these taxa from our cultured samples is probably because they do not

grow well in these conditions, and does not necessarily contradict our finding them in the

sequence data.

As above for our community-level sequence data, the relative abundances of different taxa

inferred from colony counts on chromatic agar varied among different shop types

(PERMANOVA with zero-adjusted Bray-Curtis dissimilarity (Clarke *et al.*, 2006): effect of

shop type – $F_{4,49} = 2.507$, $p = 0.003$) but not geographic locations (effect of location – $F_{9,41} =$

0.080, $p = 0.701$). Thus, banknotes yielded viable colonies and taxonomic identification on

chromatic agar, albeit relatively coarse-grained compared to our sequence data, supporting

similar conclusions (both in terms of which taxa were present and variation of community

composition with collection environment but not geographic location).

DISCUSSION

We found that Swiss banknotes taken directly from circulation harbour a variety of bacterial

DNA, with the key taxa (Fig. 1B) including families found before in humans and human-

associated environments, e.g. the human skin, gut or respiratory tract microbiome, or human

food production (Cogen *et al.*, 2008; Fierer *et al.*, 2008; Murphy and Parameswaran, 2009;

Oren, 2011; Fourquin-Gomez *et al.*, 2014; Delanghe *et al.*, 2021; Skowron *et al.*, 2021). Our

finding that community composition varied depending on the shop type in which each banknote

was collected is surprising: this suggests money has a microbiome reflecting its local

environment, even though we assume banknotes are only transiently exposed to each

habitat/shop type. By extension, we expect individual banknotes to have a dynamic

microbiome that changes during their "lifecycle" as they pass from one habitat to another. By

contrast, we found no evidence for geographic structuring, unlike some previous studies with

microbes in other contexts (Whitaker *et al.*, 2003b; Horner-Devine *et al.*, 2004), but consistent

with other observations of relatively weak biogeographic structuring in microbes (Hillebrand

et al., 2001; Finlay, 2002; Zhou *et al.*, 2008). Our plating results corresponded well with

conclusions from sequencing, indicating that at least some taxa are present as viable, culturable

cells. Together these observations suggest, on the scale investigated here, deterministic

assembly processes (e.g., selection imposed by habitat heterogeneity) are more important

drivers of banknote-associated bacterial community composition than other processes

associated with geographic distance among sites, such as dispersal limitation (Chase, 2010).

The taxa we identified as being relatively more abundant on banknotes collected from

butcher shops (*Brevibacteriaceae*, *Halomonadaceae*, *Lactobacillaceae*, *Vibrionaceae*) are

known to include bacteria previously identified in "meaty" contexts. *Halomonadaceae* and

Brevibacteriaceae have previously been found in high-salinity environments and salty foods

such as poultry skin or Chinese pork cured with salt (Oren, 2011; Fourquin-Gomez *et al.*,

2014). *Brevibacteriaceae* and *Vibrionaceae* are frequently found in samples from butchers,

slaughterhouses, abattoirs (Sierra *et al.*, 1995; Połka *et al.*, 2015; Fuertes-Perez *et al.*, 2019;

Dourou *et al.*, 2021). Representatives of the *Lactobacillaceae* family are key species used as

starter cultures in production of, and have been isolated from, cured and fermented meat

(Deibel *et al.*, 1961; Samelis *et al.*, 1994; Cocolin *et al.*, 2011). It is important to note that

members of the same families have also been isolated from a wide range of other environments

including humans, plants, animals and foods (Walter, 2008; Oren, 2011; Felis and Pot, 2014;

Forquin and Weimer, 2014; Franz *et al.*, 2014; Leisner and Pot, 2014; Pot *et al.*, 2014;

Takemura *et al.*, 2014)(Oren, 2011) . Therefore, while our results are consistent with past work

finding these taxa in "meaty" environments, they can also occur elsewhere. It is interesting in

this context that the sample butcher-BIA – classified as "butcher" but in fact originating from

a supermarket till due to the respective location's two butchers being closed on collection day

– was relatively impoverished for the taxa linked above to the other butcher samples (butcher-

BIA: 0.45% *Brevibacteriaceae*, 0.00% *Halomonadaceae*, 0.07% *Lactobacillaceae*, 0.00%

Vibrionaceae).

At a broader level, the taxa we identified across all shop types are in line with previous

research into monetary microbiomes. For example, in a metagenomic analysis of dollar bills

from New York City, Maritz *et al.* (2017) found representatives of several of the families that

were dominant in our dataset (including *Staphylococcaceae*, *Micrococcaceae*, *Streptococcaceae*, *Corynebacteriaceae*, and *Moraxellaceae*). Similarly, an analysis of Chinese RMB banknotes and US dollar bills (Lin *et al.*, 2021) revealed the most abundant taxa on the family level were *Moraxellaceae*, *Pseudomonadaceae* and *Enterobacteriaceae* (all abundant in our data set). Other studies of monetary microbiomes identified similarly overlapping sets of taxa, including analysis of Indian Rupee banknotes (*Actinobacteria*, *Bacteriodetes*, *Firmicutes*, and *Proteobacteria*; no lower-taxa resolution available) (Jalali *et al.*, 2015), banknotes from Hong Kong (*Moraxellaceae*, *Micrococcaceae*, *Enterobacteriaceae, Vibrionaceae*, *Corynebacteriaceae*, *Pseudomonadaceae*, among others) (Heshiki *et al.*, 2017) and from Brazil (*Enterobacteriaceae*, *Staphylococcaceae*, *Lactobacillaceae*, *Moraxellaceae*, and *Corynebacteriaceae*, among others) (da Fonseca *et al.*, 2015). Our results go beyond past work on monetary microbiomes by showing that the relative abundances of different taxa (families and ZOTUs) vary among collection environments (shop types), but not with geographic distance within a European country.

A key implication of our results is that even temporary exposure of banknotes to certain environments is reflected by the community composition of colonizing bacteria. This is surprising because we assume the residence time of individual banknotes in a given shop is short (hours, days or weeks). It also suggests high variability in temporal beta diversity in the monetary microbiome. We found several taxa (ZOTUs) that were unique to particular shop types, and at the family level it was also visible that some taxa were overrepresented in butcher samples. Despite this, analogous to the core microbiome Danko *et al.* (2021) found in urban mass transit systems, we found something akin to a core microbiome across our samples in that two families (*Pseudomonadaceae, Staphylococcaceae*) were present in every sample.

Another important implication of our results is that many of the key taxa of the monetary microbiome in Switzerland are also present as viable cells (culturable on agar). This raises an

interesting question about colonization dynamics in the monetary microbiome. If viable

microbes are present on banknotes, are they growing and interacting on the banknote itself?

We can speculate the rough surface of most banknotes not only allows attachment of bacteria,

but also human skin cells, secretions and remnants of food or products during handling.

Potentially, this could support bacterial growth after colonization. If the local

microenvironment influences the relative growth/death rates of different taxa, this could even

shape community composition and contribute to habitat-specific community composition as

we observed here. Alternatively, community composition may be determined primarily at the

colonization step, with different shop types harbouring different microbiomes in general (due

to local habitat selection), which are "sampled" by banknotes passing through the shop. One

avenue for future work aiming to tease apart these different potential drivers of the monetary

microbiome would be to experimentally colonize banknotes and track community abundance

and composition over time (longitudinal sampling of individual banknotes).

We emphasize strongly that our results do not address or make any predictions regarding

food hygiene or potential health hazards posed to humans. We were solely interested in the

microbial community composition on banknotes and how this relates to the sampling

environment. Finding distinct microbiomes in different shop types does not necessarily

translate to variable infection risk or indicate variable hygiene standards, only that the local

monetary microbiome reflects the local commercial environment. One limitation of our study

is that the SARS-CoV2 pandemic has led to changes in customers' payment behaviour

(Kraenzlin *et al.*, 2020). This might influence the transfer of microbiota to banknotes, because

of increased hygienic awareness in customers and shop staff. For example, we observed in one

case a banknote being disinfected by shop staff (the sample extracted from the banknote from

this shop was removed in the initial filtering step due to low read counts, see Methods).

However if anything we would expect any effect of pandemic-related hygiene measures to

make our conclusions conservative: when hygiene standards are less stringent we might expect

even more exchange of microbiota between the local environment and banknotes. A further

limitation of our study is the spatial scale and number of shop types. While our study design

revealed significant variation among shop types, the lack of geographic structure across

Switzerland may differ from that across larger spatial scales or across different currency types.

In summary, we found that the bacterial communities associated with Swiss banknotes can

vary depending on the environment in which the banknotes were collected and that, by contrast,

geographic distance between sampled locations does not predict variation of the associated

microbial communities. Our results also show that the most abundant taxa are not only present

in the form of cell debris and genetic material, but as viable cells. This raises the question of

banknotes potentially being involved in transmission of bacteria between people and the

environment, and/or of genes, such as antibiotic resistance genes, among bacteria co-occurring

on the same banknotes. Therefore, a key avenue for future work building on the link between

microbial community structure on banknotes and the local environment (shop type) identified

here is to ask whether this translates to variable infection risk or horizontal gene transfer.

<u>Acknowledgments</u>

AMB thanks Richard Allen, Elrina Bomben, Carole Imhof, Ricardo Leon Sampedro; Ethics

Commission ETHZ, GDC ETHZ, Legal Services ETHZ, Statistical Consulting Group ETHZ.

AMB and ARH thank Andrew Letten.

References

Anderson, M.J. (2001) A new method for non-parametric multivariate analysis of variance. *Austral Ecol* **26**: 32–46.

De Angelis, M. and Gobbetti, M. (2011) Lactobacillus SPP.: General Characteristics. In, Fuquay,J.W. (ed), *Encyclopedia of Dairy Sciences*. Academic Press; Copyright © 1999 Elsevier Ltd. All rights reserved., pp. 78–90.

De Angelis, M. and Gobbetti, M. (2016) Lactobacillus SPP.: General Characteristics ☆. *Ref Modul Food Sci*.

Argandoña, M., Vargas, C., Reina-Bueno, M., Rodríguez-Moya, J., Salvador, M., and Nieto, J.J. (2012a) An extended suite of genetic tools for use in bacteria of the halomonadaceae: An overview. In, Lorence,A. (ed), *Methods in Molecular Biology*. Heidelberg: Springer, pp. 167–201.

Argandoña, M., Vargas, C., Reina-Bueno, M., Rodríguez-Moya, J., Salvador, M., and Nieto, J.J. (2012b) An extended suite of genetic tools for use in bacteria of the halomonadaceae: An overview. *Methods Mol Biol* **824**: 167–201.

Baron, S., Ceccarelli, D., Dalsgaard, I., Granier, S.A., Haenen, O., Jansson, E., et al. (2020) Influence of incubation time on antimicrobial susceptibility testing of pathogenic Vibrio anguillarum and Vibrio vulnificus isolated from fish. *Aquaculture* **524**: 1–5.

Central Bank, E. (2020) Study on the payment attitudes of consumers in the euro area (SPACE).

Chao, A. (1984) Nonparametric estimation of the number of classes in a population. *Scand Sournal Stat* **11**: 265–270.

Chao, A. and Bunge, J. (2002) Estimating the number of species in a stochastic abundance model. *Biometrics* **58**: 531–539.

Chase, J.M. (2003) Community assembly: When should history matter? *Oecologia* **136**: 489–498.

Chase, J.M. (2010) Stochastic community assembly causes higher biodiversity in more productive environments. *Science (80-)* **328**: 1388–1391.

Chase, J.M. and Myers, J.A. (2011) Disentangling the importance of ecological niches from stochastic processes across scales. *Philos Trans R Soc B Biol Sci* **366**: 2351–2363.

Chesson, P. (2000) Mechanisms of Maintenance of Species Diversity. *Annu Rev Ecol Syst* **31**: 343–66.

Clarke, K.R. (1993) Non-parametric multivariate analyses of changes in community structure. *Aust J Ecol* **18**: 117–143.

Clarke, K.R., Somerfield, P.J., and Chapman, M.G. (2006) On resemblance measures for ecological studies, including taxonomic dissimilarities and a zero-adjusted Bray-Curtis coefficient for denuded assemblages. *J Exp Mar Bio Ecol* **330**: 55–80.

Cocolin, L., Dolci, P., and Rantsiou, K. (2011) Biodiversity and dynamics of meat fermentations: The contribution of molecular methods for a better comprehension of a complex ecosystem. *Meat Sci* **89**: 296–302.

Cogen, A.L., Nizet, V., and Gallo, R.L. (2008) Skin microbiota: A source of disease or defence? *Br J Dermatol* **158**: 442–455.

Cousin, M.A. (1999) PSEUDOMONAS | Introduction. In, Robinson,R.K.B.T.-E. of F.M. (ed), *Encyclopedia of Food Microbiology*. Oxford: Elsevier, pp. 1864–1867.

Danko, D., Bezdan, D., Afshin, E.E., Ahsanuddin, S., Bhattacharya, C., Butler, D.J., et al. (2021) A global metagenomic map of urban microbiomes and antimicrobial resistance. *Cell* **184**: 3376–3393.

Deibel, R.H., Wilson, G.D., and Niven, C.F. (1961) Microbiology of Meat Curing. *Appl Microbiol* **9**: 239–243.

Delanghe, L., Spacova, I., Van Malderen, J., Oerlemans, E., Claes, I., and Lebeer, S. (2021) The role of lactobacilli in inhibiting skin pathogens. *Biochem Soc Trans* **49**: 617–627.

Diamond, J.M. (1975) Assembly of species communities. *Ecol Evol communities* 342–444.

Dourou, D., Spyrelli, E.D., Doulgeraki, A.I., Argyri, A.A., Grounta, A., Nychas, G.J.E., et al. (2021) Microbiota of chicken breast and thigh fillets stored under different refrigeration temperatures assessed by next-

generation sequencing. *Foods* **10**: 1–17.

Ebach, M.C. (2015) Origins of Biogeography, Dordrecht: Springer Netherlands.

Edgar, R.C. (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**: 2460–2461.

Fargione, J., Brown, C.S., and Tilman, D. (2003) Community assembly and invasion: An experimental test of neutral versus niche processes. *Proc Natl Acad Sci U S A* **100**: 8916–8920.

Felis, G.E. and Pot, B. (2014) The family Lactobacillaceae. In, Holzapfel,W.H. and Wood,B.J. (eds), *Lactic Acid Bacteria: Biodiversity and Taxonomy*. John Wiley & Sons, Ltd., pp. 245–247.

Fierer, N. (2017) Embracing the unknown: Disentangling the complexities of the soil microbiome. *Nat Rev Microbiol* **15**: 579–590.

Fierer, N., Hamady, M., Lauber, C.L., and Knight, R. (2008) The influence of sex, handedness, and washing on the diversity of hand surface bacteria.

Finlay, B.J. (2002) Global dispersal of free-living microbial eukaryote species. *Science (80-)* **296**: 1061–1063.

da Fonseca, T.A.P., Pessôa, R., and Sanabani, S.S. (2015) Molecular analysis of bacterial microbiota on Brazilian currency note surfaces. *Int J Environ Res Public Health* **12**: 13276–13288.

Forquin, M.P. and Weimer, B.C. (2014) Brevibacterium. *Encycl Food Microbiol Second Ed* **1**: 324–330.

Fourquin-Gomez, M.-P., Weimer, B.C., Sorieul, L., Kalinowski, J., and Vallaeys, T. (2014) The Family Brevibacteriaceae. In, Rosenberg,E., DeLong,E.F., Lory,S., Stackebrandt,E., and Thompson,F. (eds), *The Prokaryotes: Actinobacteria*. Heidelberg: Springer, pp. 141–153.

Franz, C.M.A.P., Endo, A., Abriouel, H., Van Reenen, C.A., Gálvez, A., and Dicks, L.M.T. (2014) The genus Pediococcus.

Fuertes-Perez, S., Hauschild, P., Hilgarth, M., and Vogel, R.F. (2019) Biodiversity of Photobacterium spp. Isolated From Meats. *Front Microbiol* **10**:.

Gehring, B., Graf, S., and Trütsch, T. Swiss Payment Monitor 2020 Wie bezahlt die Schweiz?

Hanson, C.A., Fuhrman, J.A., Horner-Devine, M.C., and Martiny, J.B.H. (2012) Beyond biogeographic patterns: Processes shaping the microbial landscape. *Nat Rev Microbiol* **10**: 497–506.

Heshiki, Y., Dissanayake, T., Zheng, T., Kang, K., Yueqiong, N., Xu, Z., et al. (2017) Toward a metagenomic understanding on the bacterial composition and resistome in Hong Kong banknotes. *Front Microbiol* **8**:.

Hillebrand, H., Watermann, F., Karez, R., and Berninger, U.G. (2001) Differences in species richness patterns between unicellular and multicellular organisms. *Oecologia* **126**: 114–124.

HilleRisLambers, J., Adler, P.B., Harpole, W.S., Levine, J.M., and Mayfield, M.M. (2012) Rethinking community assembly through the lens of coexistence theory. *Annu Rev Ecol Evol Syst* **43**: 227–248.

Horner-Devine, M.C., Lage, M., Hughes, J.B., and Bohannan, B.J.M. (2004) A taxa-area relationship for bacteria. *Nature* **432**: 750–753.

Hubbell, S.P. (2011) The Unified Neutral Theory of Biodiversity and Biogeography (MPB-32), Princeton University Press.

Jalali, S., Kohli, S., Latka, C., Bhatia, S., Vellarikal, S.K., Sivasubbu, S., et al. (2015) Screening currency notes for microbial pathogens and antibiotic resistance genes using a shotgun metagenomic approach. *PLoS One* **10**:.

Kim, K.K., Lee, J.S., and Stevens, D.A. (2013) Microbiology and epidemiology of Halomonas species. *Future Microbiol* **8**: 1559–1573.

Kraenzlin, S., Meyer, C., and Nellen, T. (2020) COVID-19 and regional shifts in Swiss retail payments (Version of July 23, 2020).

Leisner, J.J. and Pot, B. (2014) The genus Paralactobacillus.

Libertucci, J. and Young, V.B. (2019) The role of the microbiota in infectious diseases. *Nat Microbiol* **4**: 35–45.

Lin, J., Jiang, W., Shi, Y., and Cai, W. (2021) Metagenomic Sequencing Revealed the Potential Pathogenic Threats of Banknotes. *ACS Omega* **6**: 3499–3507.

Lonvaud-Funel, A. (2014) Leuconostocaceae Family, Second Edi. Elsevier.

Love, M.I., Huber, W., and Anders, S. (2014) Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol* **15**:.

Macarthur, R. and Levins, R. (1967) The Limiting Similarity, Convergence, and Divergence of Coexisting
Species. *Am Nat* **101**: 377–385.

Madhaiyan, M., Wirth, J.S., and Saravanan, V.S. (2020) Phylogenomic analyses of the Staphylococcaceae
family suggest the reclassification of five species within the genus Staphylococcus as heterotypic
synonyms, the promotion of five subspecies to novel species, the taxonomic reassignment of five
Staphylococcu. *Int J Syst Evol Microbiol* **70**: 5926–5936.

Maritz, J.M., Sullivan, S.A., Prill, R.J., Aksoy, E., Scheid, P., and Carlton, J.M. (2017) Filthy lucre: A
metagenomic pilot study of microbes found on circulating currency in New York City. *PLoS One* **12**:
e0175527.

Murphy, T.F. and Parameswaran, G.I. (2009) Moraxella catarrhalis, a human respiratory tract pathogen. *Clin*
Infect Dis **49**: 124–131.

Oren, A. (2011) The Halophilic World of Lourens Baas Becking. In, Ventosa,A., Oren,A., and Ma,Y. (eds),
Halophiles and Hypersaline Environments. Heidelberg: Springer, pp. 9–25.

Połka, J., Rebecchi, A., Pisacane, V., Morelli, L., and Puglisi, E. (2015) Bacterial diversity in typical Italian
salami at different ripening stages as revealed by high-throughput sequencing of 16S rRNA amplicons.
Food Microbiol **46**: 342–356.

Pot, B., Felis, G.E., De Bruyne, K., Tsakalidou, E., Papadimitriou, K., Leisner, J., and Vandamme, P. (2014)
The genus Lactobacillus.

Reese, A.T., Madden, A.A., Joossens, M., Lacaze, G., and Dunn, R.R. (2020) Influences of Ingredients and
Bakers on the Bacteria and Fungi in Sourdough Starters and Bread. *mSphere* **5**: 1–15.

Robert-Pillot, A., Copin, S., Himber, C., Gay, M., and Quilici, M.L. (2014) Occurrence of the three major
Vibrio species pathogenic for human in seafood products consumed in France using real-time PCR. *Int J*
Food Microbiol **189**: 75–81.

Rock, C. and Donnenberg, M.S. (2014) Human Pathogenic Enterobacteriaceae. In, *Reference Module in*
Biomedical Sciences. Elsevier.

Salonen, A., Salojärvi, J., Lahti, L., and de Vos, W.M. (2012) The adult intestinal core microbiota is determined
by analysis depth and health status. *Clin Microbiol Infect* **18**: 16–20.

Samelis, J., Maurogenakis, F., and Metaxopoulos, J. (1994) Characterisation of lactic acid bacteria isolated from
naturally fermented Greek dry salami. *Int J Food Microbiol* **23**: 179–196.

Schaarschmidt, J. (1884) Upon the occurrence of bacteria and minute algæ on the surface of paper money [3].
Nature **30**: 360.

Shade, A. and Handelsman, J. (2012) Beyond the Venn diagram: The hunt for a core microbiome. *Environ*
Microbiol **14**: 4–12.

Sierra, M.L., Gonzalez-Fandos, E., García-López, M.L., Camino Garcia Fernandez, M., and Prieto, M. (1995)
Prevalence of salmonella, yersinia, aeromonas, campylobacter, and cold-growing escherichia coli on
freshly dressed lamb carcasses. *J Food Prot* **58**: 1183–1185.

Skowron, K., Bauza-Kaszewska, J., Kraszewska, Z., Wiktorczyk-Kapischke, N., Grudlewska-Buda, K.,
Kwiecińska-Piróg, J., et al. (2021) Human skin microbiome: Impact of intrinsic and extrinsic factors on
skin microbiota. *Microorganisms* **9**: 543.

Stegen, J.C., Lin, X., Fredrickson, J.K., and Konopka, A.E. (2015) Estimating and mapping ecological
processes influencing microbial community assembly. *Front Microbiol* **6**:.

Takemura, A.F., Chien, D.M., and Polz, M.F. (2014) Associations and dynamics of vibrionaceae in the
environment, from the genus to the population level. *Front Microbiol* **5**:.

Tauch, A. and Sandbote, J. (2014) The Family Corynebacteriaceae. In, Rosenberg,E., Delong,E.F., Lory,S.,
Stackebrandt,E., and Thompson,F. (eds), *The Prokaryotes: Actinobacteria*. Heidelberg: Springer, pp. 239–
278.

Thomas, Y., Vogel, G., Wunderli, W., Suter, P., Witschi, M., Koch, D., et al. (2008) Survival of influenza virus
on banknotes. *Appl Environ Microbiol* **74**: 3002–3007.

Vinet, L. and Zhedanov, A. (2011) A "missing" family of classical orthogonal polynomials. In, Rosenbert,E.,
DeLong,E.F., Lory,S., Stackebrandt,E., and Thompson,F. (eds), *Journal of Physics A: Mathematical and*
Theoretical. Heidelberg: Springer, pp. 443–476.

Vriesekoop, F., Chen, J., Oldaker, J., Besnard, F., Smith, R., Leversha, W., et al. (2016) Dirty money: A matter
of bacterial survival, adherence, and toxicity. *Microorganisms* **4**:.

Walter, J. (2008) Ecological role of Lactobacilli in the gastrointestinal tract: Implications for fundamental and
biomedical research. *Appl Environ Microbiol* **74**: 4985–4996.

Wareing, P. (2016) Microbiology of soft drinks and fruit juices. In, Ashurst,P.R. (ed), *Chemistry and*
Technology of Soft Drinks and Fruit Juices. Chichester: John Wiley & Sons, Ltd., pp. 290–309.

Webb, C.O. Exploring the Phylogenetic Structure of Ecological Communities: An Example for Rain Forest
Trees.

Weiss, S., Xu, Z.Z., Peddada, S., Amir, A., Bittinger, K., Gonzalez, A., et al. (2017) Normalization and
microbial differential abundance strategies depend upon data characteristics. *Microbiome* **5**: 1–18.

Whitaker, R.J., Grogan, D.W., and Taylor, J.W. (2003a) Geographic barriers isolate endemic populations of
hyperthermophilic archaea. *Science (80-)* **301**: 976–978.

Whitaker, R.J., Grogan, D.W., and Taylor, J.W. (2003b) Geographic barriers isolate endemic populations of
hyperthermophilic archaea. *Science (80-)* **301**: 976–978.

Zapala, M.A. and Schork, N.J. (2006) Multivariate regression analysis of distance matrices for testing
associations between gene expression patterns and related variables. *Proc Natl Acad Sci U S A* **103**:
19430–19435.

Zhou, J., Kang, S., Schadt, C.W., and Garten, C.T. (2008) Spatial scaling of functional gene diversity across
various microbial taxa. *Proc Natl Acad Sci U S A* **105**: 7768–7773.

Zhou, J., Liu, W., Deng, Y., Jiang, Y.H., Xue, K., He, Z., et al. (2013) Stochastic assembly leads to alternative
communities with distinct functions in a bioreactor microbial community. *MBio* **4**:.

Zhou, J. and Ning, D. (2017) Stochastic Community Assembly: Does It Matter in Microbial Ecology?

Data Accessibility and Benefit-Sharing Statement

The sequencing data have been deposited in the European Nucleotide Archive under the study

accession number PRJEB45390 (https://www.ebi.ac.uk/ena; data will be made accessible upon

publication of the study).

Other data presented in this study will be made available at the Dryad Digital Repository:

http://dx.doi.org/...: to be completed after manuscript is accepted for publication.

Benefits Generated: Benefits from this research accrue from the sharing of our data and

results on public databases as described above.

<u>Author Contributions</u>

AMB, ARH   designed research.

AMB   performed research.

AMB, ARH   analyzed data.

AMB, ARH   wrote paper.

Tables

Supplementary Table 1. Sequences of primers used during library preparation. These
primers were used to amplify the V3-V4 regions of the 16S rRNA gene during the first course
of limited cycle PCR.

forward primers	
340F_nex 0	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGGCCTACGGGNGGCWGCAG
340F_nex 1	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGNGCCTACGGGNGGCWGCAG
340F_nex 2	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGNNGCCTACGGGNGGCWGCAG
340F_nex 3	TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGNNNGCCTACGGGNGGCWGCAG
reverse primers	
805R_nex 0	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGTGACTACHVGGGTATCTAATCC
805R_nex 1	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGNTGACTACHVGGGTATCTAATCC
805R_nex 2	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGNNTGACTACHVGGGTATCTAATCC
805R_nex 3	GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGNNNTGACTACHVGGGTATCTAATCC

 <u>Figures</u>

A

B

Figure 1. Bacterial community composition across sampling locations and shop types.
Panel A: Distribution of sampling locations across Switzerland. The map shows the 10
municipalities, highlighted in dark green, in which banknotes were collected. Panel B: Relative
abundances of the main bacterial families identified by amplicon sequencing from banknotes
sampled in each shop (lower *x*-axis) and location (upper *x*-axis, indicated by three-letter
abbreviations). *Unknown* contains non-assigned ZOTUs; *Other* contains assigned ZOTUs not
belonging to the most abundant families. Abbreviations in both panels: BIA = Biasca, BRI =
Brienz BE, FRI = Frick, KLO = Klosters-Serneus, LAN = Langnau am Albis, ORB = Orbe,
POR = Porrentruy, ROR = Rorschach, STE = Stein am Rhein, VIS = Visp.

Figure 2. Within-sample taxonomic diversity of samples from banknotes collected in various shop types across Switzerland. Panel A: The Shannon diversity index *H* takes into account both richness (number of species present) and evenness (how evenly do present species occur). Panel B: Observed richness gives the number of unique ZOTUs present in a sample. Colours indicate locations (abbreviations: BIA = Biasca, BRI = Brienz BE, FRI = Frick, KLO = Klosters-Serneus, LAN = Langnau am Albis, ORB = Orbe, POR = Porrentruy, ROR = Rorschach, STE = Stein am Rhein, VIS = Visp).

729 **Figure 3. Ordination analysis.** NMDS ordination based on Bray-Curtis dissimilarity between
730 the bacterial communities isolated from banknotes (stress = 0.11). Symbols represent different
731 shop types and are coloured according to location. Abbreviations: BIA = Biasca, BRI = Brienz
732 BE, FRI = Frick, KLO = Klosters-Serneus, LAN = Langnau am Albis, ORB = Orbe, POR =
733 Porrentruy, ROR = Rorschach, STE = Stein am Rhein, VIS = Visp.

Figure 4. Number and identity of colonies isolated from banknotes collected in various shop types across Switzerland. Each composite bar gives the number and identity of colonies detected in samples isolated from banknotes. The bar colour indicates species identity (based on morphology and colour on Chromatic^{TM} MH agar, see Methods). Colonies classified as "unknown" had uncertain identity (regular-shaped, small size) or a morphology more commonly associated with yeasts/fungi (astral appearance, polychromatic, with spores/fruiting body-like structures). Abbreviations: BIA = Biasca, BRI = Brienz BE, FRI = Frick, KLO = Klosters-Serneus, LAN = Langnau am Albis, ORB = Orbe, POR = Porrentruy, ROR = Rorschach, STE = Stein am Rhein, VIS = Visp.

Supplementary Figures

Supplementary Figure 1. Number of reads per sample. The figure shows the samples and
number of reads for each sample retained after filtering. Total reads per sample varied greatly,
with 140 reads for the butcher in BRI and 7'152'829 reads for the florist in BRI (~51'092x
difference). Note: *y*-axis is log10-transformed. Abbreviations: BIA = Biasca, BRI = Brienz BE,
FRI = Frick, KLO = Klosters-Serneus, LAN = Langnau am Albis, ORB = Orbe, POR =
Porrentruy, ROR = Rorschach, STE = Stein am Rhein, VIS = Visp.

Supplementary Figure 2. Venn diagram of unique and shared core ZOTUs samples from banknotes collected in various shop types across Switzerland. Each ellipse gives the number of ZOTUs present at an abundance of >0.1% in >50% of a set of banknotes attributed to a certain shop type. Numbers in overlapping areas of ellipses indicate shared ZOTUs between sets of banknotes collected in different types of shops. The seven core microbiome ZOTUs present in >50% of every set of banknotes were attributed to the genera: *Escherichia-Shigella*, *Pseudomonas*, *Propionibacterium*, *Corynebacterium*, *Streptococcus*, *Streptococcus*, and *Staphylococcus*.