A stepwise route to domesticate rice by controlling seed
shattering and panicle shape

Ryo Ishikawa[a,1,*], Cristina Cobo Castillo[a,f,1,*], Than Myint Htun[a,h,1], Koji Numaguchi[a,i], Kazuya

Inoue[b], Yumi Oka[a], Miki Ogasawara[a], Shohei Sugiyama[a], Natsumi Takama[a], Chhourn Orn[a,j],

Chizuru Inoue[a], Ken-ichi Nonomura[c,d], Robin Allaby[e], Dorian Q Fuller[f,g] and Takashige Ishii[a]

[a]Laboratory of Plant Breeding, Graduate School of Agricultural Science, Kobe University; Kobe,

657-8501, Japan

[b]Laboratory of Hydraulic Structures and Geo-Environmental Engineering, Graduate School of

Agricultural Science, Kobe University; Kobe, 657-8501, Japan

[c]Laboratory of Plant Cytogenetics, National Institute of Genetics; Mishima, Shizuoka, 411-8540,

Japan

[d]Department of Genetics, School of Life Science, The Graduate University for Advanced Studies

(SOKENDAI); Mishima, Shizuoka, 411-8540, Japan

[e]School of Life Sciences, University of Warwick; Coventry, CV4 7AL, UK

[f]Institute of Archaeology, University College London; London, WC1H 0PY, UK

[g]School of Cultural Heritage, Northwest University, Shaanxi, 710069, China

[h]Present address: Department of New Genetics, Advanced Center of Agricultural Research and

Education, Yezin Agricultural University; Yezin, Nay Pyi Taw, 15013, Myanmar

[i]Present address: Japanese Apricot Laboratory, Wakayama Fruit Tree Experiment Station;

Wakayama, 645-0021, Japan

[j]Present address: Plant Breeding Division, Cambodian Agricultural Research and Development

Institute (CARDI); Phnom Penh, Cambodia

[1] These authors contributed equally to this work.

* To whom correspondence may be addressed.

Email: Ryo Ishikawa (r-ishika@port.kobe-u.ac.jp) or Cristina Cobo Castillo (criscastillo@me.com)

Author Contributions:

R.I. and C.C.C. designed the research; R.I., C.C.C., T.M.H, Y.O., M.O., S.S., N.T., C.O., and C.I.
conducted genetic analysis; K.N. and R.A. carried out population genetic analysis; C.C.C. carried
out seed-gathering experiment; C.I. conducted SEM analysis; K.I. and R.I. carried out structural
mechanic simulation; K-I.N. contributed wild rice materials and their evaluation; R.I., C.C.C., R.A.,
D.F., and T.I. wrote the manuscript.

Competing Interest Statement:

The authors declare no conflict of interest.

Classification:

Biological Sciences/Plant Biology or Biological Sciences/Anthropology

Keywords:

Oryza sativa, *Oryza rufipogon*, domestication, seed shattering, closed panicle

This PDF file includes:

Main Text
Figures 1 to 5.

Abstract

Rice (*Oryza sativa* L.) is consumed by more than half of the world's population, but despite

its global importance the mechanisms of domestication remain unclear. During domestication, wild

rice (*O. rufipogon* Griff.) was transformed by acquiring non-seed-shattering behaviour, an important

genetic change that allowed humans to increase grain yield. However, we show previously

identified loci, *sh4* and *qSH3*, are individually insufficient to explain loss of seed shattering nor

increases in harvest yield in wild rice. We identify the complementary interaction of key mutations

for abscission layer interruption and panicle architecture that were causal in the early domestication

of Asian rice. An interruption of abscission layer formation requires both *sh4* and *qSH3*, which

presents an apparent barrier to selection of shattering loss. We identified the causal single

nucleotide polymorphism at *qSH3* within a seed-shattering gene *OsSh1* conserved in *indica* and

japonica subspecies, but absent in the *circum*-aus group of rice. We demonstrate through harvest

experiments that seed-shattering alone does not significantly impact yield. Instead, we observed

yield increases under a *SPR3*-controlled closed panicle formation, which is augmented by the

integration of *sh4* and *qSH3* alleles causing a slight inhibition of abscission layer. Complementary

manipulation of seed shattering and panicle shape result in a panicle structure that is mechanically

stable. We propose a stepwise route in the earliest phase of rice domestication in which selection

for visible *SPR3*-controlled closed panicle morphology was instrumental in the sequential

recruitment of *sh4* and *qSH3* and leading to loss of shattering.

Significance Statement

Rice is one of the most important crops worldwide. Loss of seed shattering in domesticated

rice, previously attributed to single mutations such as in *sh4*, is considered the principal genetic

change which resulted in yield increases. However, we show that *sh4* is insufficient on its own to

cause abscission layer disruption and other genes, such as *qSH3* are required, making

mechanisms for the initial selection of non-shattering unclear. We show that shattering loss in wild

rice genetic backgrounds does not increase yields. We identify an interaction in which a second

trait, closed panicle formation controlled by *SPR3*, both increases yield and facilitates recruitment

of *sh4* and *qSH3* which synergistically augment yield, leading to a stepwise route for rice

domestication.

116   **Main Text**
117
118   **Introduction**
119
120   Selection of naturally occurring variations in wild plants that provided useful agronomic

traits was an essential step for crop domestication. These traits are often related to yield, such as

seed number, seed size and a loss of seed dispersal, as well as ease of cultivation, including plant

architecture, seed dormancy, and photoperiod sensitivity. Plants with these traits provided the

necessary impetus for humans to shift from hunting and gathering to cultivating. The visible

morphological changes were key for triggering domestication in many crop species. Among several

domestication-related traits, the suppression of seed shattering is considered the most important

genetic change that allowed humans to increase grain yield and differentiated domesticated from

wild plants (1, 2).

129   Rice (*Oryza sativa* L.) is consumed by almost half of the world's population and is

particularly important in Asia. During its domestication, wild rice (*O. rufipogon* Griff.) was

transformed by acquiring non-seed-shattering behaviour. Investigations on prehistoric rice spikelet

bases from archaeological sites demonstrate that suppression of seed shattering leaves a

phenotypic trace that can be observed through the examination of the abscission layer (3).

Archaeological rice spikelet bases with rough and deeply torn scars in the spikelet base (rachilla)

indicate that humans had to actively detach the spikelets from the pedicels by threshing. More than

a decade ago, two loci, *qSH1* and *sh4*, were identified as major loci that influence the reduction in

seed shattering based on quantitative trait locus (QTL) analyses (4, 5). Since *sh4* and *qSH1*

displayed large phenotypic variances, they were recognised as crucial to rice domestication.

139   To better understand how *qSH1* and *sh4* were involved in the loss of seed shattering in

wild rice, we developed introgression lines (ILs) named IL(*qSH1*-N) and IL(*sh4*-N), which contain

small chromosomal segments from *O. sativa* Nipponbare at the *qSH1* and *sh4* loci, respectively, in

the genetic background of wild rice *O. rufipogon* W630 (6). Complete seed shattering was still

observed in these lines, indicating that neither of the single mutations at *qSH1* and *sh4* could

account for non-shattering during rice domestication. As the *sh4* mutation is commonly observed

in cultivated rice, and the *qSH1* mutation is only found in some *japonica* cultivars (4), additional

mutation(s) together with *sh4* must have played a role in reducing seed shattering during the early

stages of rice domestication.

In this study, we conducted genetic analysis of the *qSH3*, an additional locus involved in

the loss of seed shattering together with *sh4*, and identified the causal mutation selected during

rice domestication. The frequency of the *qSH3* alleles in rice cultivars and their effects on the loss

of seed shattering were evaluated. In addition, we assessed the interaction between seed

shattering and panicle shape, and also investigated their role in seed shattering behaviour based

on structural mechanic analysis to better understand the selection process on a loss of seed

shattering in rice.

Results and discussion
157

Identification of the causal mutation at *qSH3*, a locus involved in the loss of seed shattering

during rice domestication

To identify additional genomic regions involved in reducing seed shattering, we produced

an IL with chromosomal segments of *O. sativa* Nipponbare at both *sh4* and *qSH1* in the genetic

background of wild rice, *O. rufipogon* W630 (7). IL(*qSH1*-N+*sh4*-N) was crossed with Nipponbare,

and their F_2 population was subjected to QTL analysis to determine the degree of seed shattering,

and the locus *qSH3* was identified (7, 8). The effect of *qSH3* was confirmed in the wild genetic

background as IL(*qSH1*-N+*sh4*-N+*qSH3*-N) showed reduced seed-shattering behaviour compared

with IL(*qSH1*-N+*sh4*-N) (7). To identify the gene for *qSH3*, we performed high-resolution linkage

analysis using a mapping population from a cross between IL(*qSH1*-N+*sh4*-N) and Nipponbare (9)

(Fig. S1). The genomic location of *qSH3* was delimited by the two recombinants to a 1,228 bp

region that harboured a part of a previously known seed-shattering gene, *OsSh1* (*Os03g0650000*)

(Figs. S2 and S3). This gene was identified as a homolog of *Sh1*, which controls abscission layer

formation in sorghum (10). The function of *OsSh1* in seed shattering was studied using artificially

induced rice materials showing that the null mutations caused a complete loss of seed shattering

(10, 11), but the causal mutation selected in rice domestication was not determined. A sequence

survey in the 1,228 bp region of 5′ UTR, exon 1, and intron 1 of *OsSh1* identified seven

polymorphisms between Nipponbare and W630 (Fig. 1A and Table S1). Among them, only a single

nucleotide polymorphism (SNP), namely SNP-70, was located in the coding region (exon1) of

OsSh1, causing an amino acid substitution from leucine in W630 to phenylalanine in Nipponbare

(Fig. S4). To test whether SNP-70 is associated with the degree of seed shattering, we performed

a transformation test in the Nipponbare genetic background of two constructs that differed only at

the SNP-70 position. In the preliminary experiment, we observed that IL(*qSH3*-W), carrying the

W630 allele at *qSH3* in Nipponbare genetic background (Fig. S5), showed a non-seed-shattering

behaviour similar to Nipponbare, whereas IL(*qSH1*-W+*qSH3*-W) displayed a significantly higher

seed-shattering degree than the IL(*qSH1*-W) (Fig. 1B and Fig. S6). These findings indicate that the

seed-shattering effect of *qSH3* was considerably enhanced by a functional allele at *qSH1*.

Therefore, we introduced two types of constructs carrying the *qSH3* cDNA sequence of W630

(*qSH3^W630^*) and Nipponbare (*qSH3^Npb^*) into IL(*qSH1*-W) (Fig. 1C). As a result, transformants with

*qSH3^W630^* showed enhanced seed shattering, compared with the control IL(*qSH1*-W), whereas

those with *qSH3^Npb^* did not (Fig. 1D), confirming that SNP-70 is the causal mutation for reduced

seed shattering.

Causal mutation at *qSH3* is conserved in both *japonica* and *indica* but not in *circum*-aus rice

cultivars

Next, we analysed the distribution of SNP-70 at *qSH3* in cultivated rice. First, genotyping

at the causal SNPs of *sh4*, *qSH1*, and *qSH3* was conducted for the three rice cultivars, Nipponbare,

IR36, and Kasalath, belonging to *japonica*, *indica*, and *circum*-aus, respectively. All cultivars

possessed the causal mutation at *sh4* and only Nipponbare had the causal mutation at *qSH1*. As

for *qSH3*, Nipponbare and IR36 carried the causal mutation, but it was absent in Kasalath (Fig. 2A).

We further found that the Kasalath haplotype is similar to that of W630 around the *qSH3* region

(Table S1). Using the diverse varieties of the World Rice Core Collection (WRC) (12), we found

that 14 lines, all belonging to *circum*-aus, carried the functional allele at *qSH3* (Table S2 and Figs.

S7 and S8). Furthermore, the Rice 3K genome project data (13) clearly showed that both *indica*

and *japonica* carried the causal mutation, but almost 90% of *circum*-aus rice carried a functional

allele at *qSH3* (Fig. 2B). To further understand the footprint of *qSH3* selection, we analysed

nucleotide diversity across the *qSH3* genomic region. Using sequences of the Rice 3K genome

collection (13), we detected a selective sweep at *qSH3* in both *indica* and *japonica*, but not in the

circum-aus linage (Fig. 2C, Figs. S9 and S10). These results suggest that *circum*-aus may have

experienced a separate trajectory to evolve reduced seed shattering (14). Thus, the reduction in

seed shattering is dependent on lineage-specific variations in the subspecies, which is key to

understanding the processes of rice domestication.

Role of *qSH3* causal mutation in an initial loss of seed shattering

We next aimed to understand how the causal SNP at *qSH3* contributed to initial rice

domestication by reducing seed shattering in *japonica* and *indica*. Since the *sh4* mutation is

conserved in all cultivated rice, we produced IL(*sh4*-N) and IL(*qSH3*-N) in the genetic background

of wild rice *O. rufipogon* W630. Complete formation of the abscission layer similar to W630 was

observed in both ILs (Fig. 3), suggesting that the single mutation at each locus was insufficient for

phenotypic change in the abscission layer formation. However, a slight inhibition of the abscission

layer around vascular bundles was observed in IL(*sh4*-N+*qSH3*-N) (Fig. 3). A slight abscission

layer inhibition was also observed in several wild rice accessions of *O. rufipogon* carrying mutations

at both *sh4* and *qSH3* (Figs. S11 and S12, and Table S3), although they may have gained these

domestic-type alleles through introgressions from cultivated rice (15). Even if the mutations in the

two loci displayed phenotypic difference in the abscission layer formation adequate to reduce

shattering under a greenhouse conditions without wind, but their effect on seed shattering was

much less than expected under field conditions.

Seed-shattering behaviour associated with a slight inhibition of abscission layer was mitigated by a closed panicle formation

Previously, we reported that a closed panicle trait controlled by *OsLG1* had a major effect in rice domestication by facilitating grain harvest (1). A closed panicle reduces seed shedding by retaining seeds that get stuck by long awns found in the lower sections of the panicle. Interestingly, the *SPR3* (a locus regulating *OsLG1*) region was under strong selection in *indica*, *japonica*, and *circum*-aus (Fig. S13), suggesting that the closed panicle trait was selected in the early phase of rice domestication and so might have been associated with seed-shattering changes, although this trait, unlike seed shattering, is not visible archaeologically. Therefore, we generated seven wild ILs with different combinations of the three loci (*sh4*, *qSH3*, and *SPR3*) (Fig. S14) and compared their morphology together with the wild rice *O. rufipogon* W630 (Fig. S15). The heading date of all ILs was similar to those of W630 (Fig. S16), but with distinct abscission layer formation (Fig. 3 and Fig. S17) and panicle structure (Figs. 4A and 4B). Small BTS values were observed only for IL(*sh4*-N+*qSH3*-N) and IL(*sh4*-N+*qSH3*-N+*SPR3*-N), while the rest were close to zero as wild rice (Fig. 4C). We were interested in assessing the differences in yields brought about by the causal mutations and which combination of alleles would have provided humans in prehistory with higher yields, including whether a closed panicle conferred additional value to humans when gathering wild rice. We therefore subjected the seven ILs and W630 to a seed-gathering experiment in the field (Fig. S18, Movies S1 and S2). The seed-gathering rate from IL(*sh4*-N) and IL(*qSH3*-N), and IL(*sh4*-N+*qSH3*-N) was not significantly different from that of W630 (Fig. 4D and Table S4). However, the three ILs with a closed panicle structure, IL(*SPR3*-N), IL(*sh4*-N+*SPR3*-N), and IL(*qSH3*-N+*SPR3*-N), presented a slightly increased yield. In contrast, IL(*sh4*-N+*qSH3*-N+*SPR3*-N), with a combination of closed panicle and abscission layer inhibition, showed a significant increase in gathering rate compared with W630 (Fig. 4D and Table S4). As closed panicles would be easily visible in the field, humans could have targeted this higher yielding rice (Fig. S19).

When plants were sown from harvests, closed panicles would have been selected for, and unintentionally would have selected also for reduced shattering, because of the enhanced

gathering rate that would occur in the presence of the domesticated alleles. As these alleles

increased in frequency and became fixed this would have increased yields, which would have

encouraged further investment in rice cultivation (16). The selection for closed panicle instigates

self-pollination behaviour owing to the long awns, which disturb the free exposure of anthers and

stigmas (stigmas and closed panicle) (17). Thus, a closed panicle may also be advantageous in

mitigating natural variations in seed-shattering loci by reducing outcrossing. Although awns in

modern cultivated rice are undesirable, gathering of rice in the early stages of rice domestication

might have benefitted from the presence of awns in plants with closed panicles, as these would

have increased yield and self-pollination rates.

Complementary interaction of a slight inhibition of abscission layer and closed panicle

formation synergistically contributed to structural stability of panicle to increase yield

To better understand how the closed panicle, caused by *SPR3*, and the inhibition of

abscission layer, caused by *sh4* and *qSH3*, contributed to the initial loss of seed shattering, we

analysed their roles by performing structural mechanics analysis. The awns of wild rice play a

pivotal role in seed dispersal in spreading panicles (Fig. 5A). We measured the lengths and weights

of awns and grains in wild rice *O. rufipogon* W630, as well as calculated the sectional force exerted

on the spikelet base depending on panicle angles (Fig. 5B). The axial and shear forces were slightly

increased in a closed panicle compared to an open panicle (Fig. S20). However, the bending

moment, which is the predominant factor affecting seed dispersal in an open panicle, was

considerably reduced in a closed panicle (Fig. S20). A slight inhibition of the abscission layer by

sh4 and *qSH3* led to an increase in the length of abscission layer inhibition (Fig. 3). Therefore, we

measured the length of the abscission layer and central vascular bundle in *O. rufipogon* W630 by

scanning microscopy (Figs. 5C and D). We also calculated the moment of inertia of the area, a

property that describes the torque required to break the abscission layer and so disarticulate the

grain, for the disrupted abscission layer (Fig. S20). The value increased exponentially with

increasing lengths of abscission layer inhibition. A reduction in the bending moment and an

increase in the moment of inertia of the area act synergistically to reduce bending stress (Fig. 4E),

contributing to the structural stability of spikelets without shattering. Thus, the interaction between

a closed panicle and abscission layer inhibition acted complementarily to increase yield.

Conclusions

In this study, we identified key functional genes that contributed to rice domestication by

increasing harvest yields. We identified the causal SNP at *qSH3* involved in reduced seed

shattering, whereas the previously proposed *sh4* mutation alone could not trigger the non-

shattering morphology in rice (6, 17). We also explained how the early selection of a closed panicle

and the resulting mechanics of spikelet retention would have increased yields and facilitated the

selection for non-shattering. Our results showed that the initial step of rice domestication is more

complex than previously thought. With new information from our studies, a change in the long-held

perspective from a single domestication allele model to one where synergistic effects of several

domestication genes is needed.

The origin and spread of rice subspecies based on population genetic analysis have been

the subject of discussions (14, 18-21). However, most of the studies have been conducted using

the genome information of modern cultivars and wild rice, without considering the importance of

visible phenotypic changes targeted by humans. Therefore, the lineage-specific variations

associated with quantitative traits are a key to further understand the process of rice domestication.

Based on our work, we propose a stepwise route for rice domestication. In wild rice, any of

the natural variations in the loci for seed shattering and closed panicle formation alone had little

effect on increasing yield (Fig. 4). Their combination, however, established an archaic rice which

could have been recognised by ancient gatherers as advantageous to increasing yields. The

change in harvesting efficiency along with the use of harvesting tools further promoted the

selections of natural variants in domestication-related traits in rice, a crop which now supports

billions of people worldwide.

306
307

Materials and Methods

Details regarding plant materials, QTL mapping, fine mapping, transformation, population genetic analysis, histological analysis, seed gathering experiment, and structural mechanics analysis are provided in the Supplementary Information.

Figure legends

Fig. 1. Identification of a causal SNP of *qSH3* associated with the degree of seed shattering.
(*A*) Fine mapping of the causal mutation at *qSH3* in a 1,228-bp region on chromosome 3. A causal SNP (SNP-70) identified in *OsSh1* (*Os03g0650000*) is shown in bold. (*B*) Comparison of seed-shattering degree by breaking tensile strength (BTS) values in three introgression lines, IL(*qSH3*-W), IL(*qSH1*-W), and IL(*qSH1*-W+*qSH3*-W) in the Nipponbare genetic background. Data are mean ± s.d. of four plants. n.s. and ** indicate not significant and significant at the 1% level based on unpaired Student's t-test, respectively. (*C*) Two types of constructs carrying *qSH3* cDNA sequences of W630 and Nipponbare (*qSH3^{W630}* and *qSH3^{Npb}*) driven by a 3-kb region of the promoter used for transgenic analysis. (*D*) The BTS values observed for the transformants with *qSH3^{W630}* and *qSH3^{Npb}*. Black triangles represent the average BTS value of IL(*qSH1*-W).

Fig. 2. Lineage-specific selection at *qSH3* locus in rice.
(*A*) Genotyping at *qSH3*, *sh4*, and *qSH1* for *O. rufipogon* W630, *O. sativa japonica* Nipponbare, *indica* IR36, and *circum*-aus Kasalath based on the causal SNPs by dCAPS markers. (*B*) Allele frequency of *qSH3* causal SNP (%) in cultivated rice based on the 3K rice project data. Nipponbare type ('T') and W630 type ('C') are shown in yellow and green, respectively. (*C*) Nucleotide diversity (π) observed for the domesticated rice in the physical position of 25.0-25.3 Mb on chromosome 3. A flanking region of the *qSH3* locus (around 25.2 Mb) was substantially decreased in *japonica* and *indica* cultivars. π was calculated in 10-kb windows using SNPs data of the 3K rice project (13).

Fig. 3. Abscission layer formation partially inhibited by *sh4* and *qSH3* in wild rice.

Abscission layer formation in *O. rufipogon* W630, IL(*sh4*-N), IL(*qSH3*-N), IL(*sh4*-N+*qSH3*-N), and

O. sativa Nipponbare. Each enlarged section indicated by the dotted square is shown in the lower

panel. A black triangle indicates an inhibited area of the abscission layer observed for IL(*sh4*-

N+*qSH3*-N). Scale bars = 100 μm.

Fig. 4. Role of the *sh4, qSH3,* and *SPR3* mutations on the initial loss of seed shattering in

rice domestication.

(*A*, *B*) Panicle shape of the seven ILs at flowering (*A*) and seed-maturation (*B*) stages. Scale bars

= 5 cm (*C*) Breaking tensile strength (BTS) values for the seven ILs. BTS values for *O. sativa*

japonica Nipponbare and *indica* IR36 are shown as control. n.s. indicates not significant by

unpaired Student's t-test. (*D*) Seed-gathering rates (mean ± s.d. of four plot replicates, with nine

plants in each plot) for *O. rufipogon* W630 and the seven ILs. Mean values labelled with different

letters are significantly different, whereas those with the same letters are not (Tukey's test with

arcsine transformed values, $P < 0.05$).

Fig. 5. Structural mechanics analysis of panicle shape and abscission layer inhibition for

the initial loss of seed shattering.

(*A*), Panicle shape of *O. rufipogon* W630. Bar = 5cm. (*B*) Schematic representation of awn and

grain with panicle angle. *θ* represents the panicle angle. l_1, l_2, m_1g, and m_2g indicate the length of

grain, awn, weight of grain and awn, respectively. (*C*) Detachment of grain and pedicel (left) and

scanning electron microscopy (SEM) analysis of the abscission layer of pedicel in *O. rufipogon*

W630 (right). AL and VB indicate abscission layer and vascular bundle, respectively. Bars = 1 mm

(left) and 100 μm (right). (*D*) Schematic representation of the abscission layer. *D* and *d* indicate the

diameter of abscission layer and vascular bundles, respectively. Dotted circle indicates the area of

disrupted abscission layer. (*E*) Simulation of the bending stress exerted on the spikelet base

depending on panicle shape and abscission layer inhibition, with higher stress experience in open

panicle with less inhibited abscission.

Acknowledgments

The wild rice accessions of *O. rufipogon* were provided by the National Institute of Genetics supported by the National Bioresource Project, MEXT, Japan. We thank R. Morita and H. Fukayama, Kobe University, for their support on producing transgenic plants, and H. Furuumi, National Institute of Genetics, for growing wild rice accessions. This work was partly supported by Grants-in-Aid from the Japan Society for the Promotion of Science, 15KK0280 and 18K05594 (R.I.), JSPS overseas research program (C.C.C.), JSPS Bilateral Open Partnership Joint Research Project, No. JPJSBP120189948 (R.I., D.F.), Nikki Saneyoshi and Kinoshita foundations (R.I.), NIG-Joint research (K-I.N, T.I.).

References

1. J. R. Harlan, J. M. de Wet, E. G. Price. Comparative evolution of cereals. *Evolution* **27**, 311–325 (1973). doi.org/10.1111/j.1558-5646.1973.tb00676.x

2. J. F. Doebley, B. S. Gaut, B. D. Smith, The molecular genetics of crop domestication. *Cell* **127**, 1309–1321 (2006). doi.org/10.1016/j.cell.2006.12.006

3. D. Q. Fuller, L. Qin Y. Zheng, Z. Zhao, X. Chen, L. A. Hosoya, G-P. Sun. The Domestication process and domestication rate in rice: spikelet bases from the lower Yangtze. *Science* **323**, 1607–1610 (2009). doi.org/10.1126/science.1166605

4. S. Konishi, T. Izawa, S. Y. Lin, K. Ebana, Y. Fukuta, T. Sasaki, M. Yano. An SNP caused loss of seed shattering during rice domestication. *Science* **312**, 1392–1396 (2006). doi.org/10.1126/science.1126410

5. C. Li, A. Zhou, T. Sang. Rice domestication by reducing shattering. *Science* **311**, 1936–1939 (2006). doi.org/10.1126/science.1123604

6. R. Ishikawa, P. T. Thanh, N. Nimura, T. M. Htun, M. Yamasaki, T. Ishii. Allelic interaction at seed-shattering loci in the genetic backgrounds of wild and cultivated rice species. *Genes Genet Syst* **85**, 265–271 (2010). doi.org/10.1266/ggs.85.265

7. T. M. Htun, C. Inoue, O. Chhourn, T. Ishii, R. Ishikawa. Effect of quantitative trait loci for seed shattering on abscission layer formation in Asian wild rice *Oryza rufipogon*. *Breed Sci* **64**, 199–205 (2014). doi.org/10.1270/jsbbs.64.199

8. K. Onishi, K. Takagi, M. Kontani, T. Tanaka, Y. Sano. Different patterns of genealogical relationships found in the two major QTLs causing reduction of seed shattering during rice domestication. *Genome* **50**, 757–766 (2007). doi.org/10.1139/G07-051

9. C. Inoue, T. M. Htun, K. Inoue, K. Ikeda, T. Ishii, R. Ishikawa. Inhibition of abscission layer formation by an interaction of two seed-shattering loci, *sh4* and *qSH3*, in rice. *Genes Genet Syst* **90**, 1–9 (2015). doi.org/10.1266/ggs.90.1

10. Z. Lin, X. Li, L. M. Shannon, C-T. Yeh, M. L. Wang, G. Bai, Z. Peng, J. Li, H. N. Trick, T. E. Clemente, J. Doebley, P. S. Schnable, M. R. Tuinstra, T. T. Tesso, F. White, J. Yu. Parallel domestication of the *Shattering1* genes in cereals. *Nat Genet* **44**, 720–724 (2012). doi.org/10.1038/ng.2281

11. F. Li, A. Komatsu, M. Ohtake, H. Eun, A. Shimizu, H. Kato. Direct identification of a
mutation in *OsSh1* causing non-shattering in a rice (*Oryza sativa* L.) mutant cultivar using
whole-genome resequencing. *Sci Rep* **10**, 14936 (2020). doi.org/10.1038/s41598-020-
71972-1

12. N. Tanaka, M. Shenton, Y. Kawahara, M. Kumagai, H. Sakai, H. Kanamori, J. Yonemaru,
S. Fukuoka, K. Sugimoto, M. Ishimoto, J. Wu, K. Ebana. Whole-genome sequencing of the
NARO World Rice Core Collection (WRC) as the basis for diversity and association studies.
Plant Cell Physiol **61**, 922–932 (2020). doi.org/10.1093/pcp/pcaa019

13. W. Wang, R. Mauleon, Z. Hu, D. Chebotarov, S. Tai, Z. Wu, M. Li, T. Zheng, R. R. Fuentes,
F. Zhang, L. Mansueto, D. Copetti, M. Sanciangco, K. C. Palis, J. Xu, C. Sun, B. Fu, H.
Zhang, Y.Gao, X. Zhao, F. Shen, X. Cui, H. Yu, Z. Li, M. Chen, J. Detras, Y. Zhou, X.
Zhang, Y. Zhao, D. Kudrna, C. Wang, R. Li, B. Jia, J. Lu, X. He, Z. Dong, J. Xu, Y. Li, M.
Wang, J. Shi, J. Li, D. Zhang, S. Lee, W. Hu, A. Poliakov, I. Dubchak, V. J. Ulat, F. N. Borja,
J. R. Mendoza, J. Ali, J. Li, Q. Gao, Y. Niu, Z. Yue, M. E. B. Naredo, J. Talag, X. Wang, J.
Li, X. Fang, Y. Yin, J-C. Glaszmann, J. Zhang, J. Li, R. S. Hamilton, R. A. Wing, J. Ruan,
G. Zhang, C. Wei, N. Alexandrov, K. L. McNally, Z. Li, H. Leung. Genomic variation in
3,010 diverse accessions of Asian cultivated rice. *Nature* **557**, 43–49 (2018).
doi.org/10.1038/s41586-018-0063-9

14. P. Civáň, H. Craig, C. J. Cox, T. A. Brown. Three geographically separate domestications
of Asian rice. *Nat Plants* **1**, 15164 (2015). doi.org/10.1038/nplants.2015.164

15. X. Jin, Y. Chen, P. Liu, C. Li, X. Cai, J. Rong, B-R. Lu. Introgression from cultivated rice
alters genetic structures of wild relative populations: implications for *in situ* conservation.
AoB Plants **10**, plx055 (2017). doi.org/10.1093/aobpla/plx055

16. D. Q. Fuller. Transitions in productivity: rice intensification from domestication to
urbanisation. *Archaeol Int* **23**, 88–103 (2020). doi.org/10.14324/111.444.ai.2020.08

17. D. Q. Fuller. Transitions in productivity: rice intensification from domestication to
urbanisation. *Archaeol Int* **23**, 88–103 (2020). doi.org/10.14324/111.444.ai.2020.08

18. X. Huang, N. Kurata, X. Wei, Z-X. Wang, A. Wang, Q. Zhao, Y. Zhao, K. Liu, H. Lu, W. Li,
Y. Guo, Y. Lu, C. Zhou, D. Fan, Q. Weng, C. Zhu, T. Huang, L. Zhang, Y. Wang, L. Feng,
H. Furuumi, T. Kubo, T. Miyabayashi, X. Yuan, Q. Xu, G. Dong, Q. Zhan, C. Li, A. Fujiyama,
A. Toyoda, T. Lu, Q. Feng, Q. Qian, J. Li, B. Han. A map of rice genome variation reveals
the origin of cultivated rice. *Nature* **490**, 497–501 (2012). doi.org/10.1038/nature11532

19. J. Y. Choi, A. E. Platts, D. Q. Fuller, Y-I. Hsing, R. A. Wing, M. D. Purugganan. The rice
paradox: multiple origins but single domestication in Asian rice. *Mol Biol Evol* **34**, 969-979
(2017). doi.org/10.1093/molbev/msx049

20. R. M. Gutaker, S. C. Groen, E. S. Bellis, J. Y. Choi, I. S. Pires, R. K. Bocinsky, E. R. Slayton,
O. Wilkins, C. C. Castillo, S. Negrão, M. M. Oliveira, D. Q. Fuller, J. A. d'A. Guedes, J. R.
Lasky, M. D. Purugganan. Genomic history and ecology of the geographic spread of rice.
Nat Plants **6**, 492-502 (2020). doi.org/10.1038/s41477-020-0659-6

21. X. Wei, J. Qiu, K. Yong, J. Fan, Q. Zhang, H. Hua, J. Liu, Q. Wang, K. M. Olsen, B. Han,
X. Huang. A quantitative genomics map of rice provides genetic insights and guides
breeding. *Nat Genet* **53**, 243–253 (2021). doi.org/10.1038/s41588-020-00769-9

22. P. T. Thanh, P. D. T. Phan, N. Mori, R. Ishikawa, T. Ishii. Development of backcross
recombinant inbred lines between *Oryza sativa* Nipponbare and *O. rufipogon* and QTL
detection on drought tolerance and submergence avoidance *Breed Sci* **61**, 76–79 (2011).
doi.org/10.1270/jsbbs.61.76

23. M. Kumagai, J. Kim, R. Itoh, T. Itoh. TASUKE: a web-based visualization program for large-
scale   resequencing   data.   *Bioinformatics*   **29**,   1806–1808.
doi.org/10.1093/bioinformatics/btt295

24. L. Mansueto, R. R. Fuentes, F. N. Borja, J. Detras, J. M. Abriol-Santos, D. Chebotarov, M.
Sanciangco, K. Palis, D. Copetti, A. Poliakov, I. Dubchak, V. Solovyev, R. A. Wing, R. S.
Hamilton, R. Mauleon, K. L. McNally, N. Alexandrov. Rice SNP-seek database update:
new SNPs, indels, and queries. *Nucl Acids Res* **45**, D1075–D1081 (2017).
doi.org/10.1093/nar/gkw1135

25. S. Purcell, B. N., K. Todd-Brown, L. Thomas, M. A. R Ferreira, D. Bender, J. Maller, P.
Sklar, P. I. W. de Bakker, M. J. Daly, P. C. Sham. PLINK: a tool set for whole-genome
association and population-based linkage analyses. *Am J Hum Genet* **81**, 559–575 (2007).
doi.org/10.1086/519795

26. B. L. Browning, Y. Zhou, S. R. Browning. A one-penny imputed genome from next-
generation reference panels. *Am J Hum Genet* **103**, 338-348 (2018).
doi.org/10.1016/j.ajhg.2018.07.015

27. Z. A. Szpiech, R. D. Hernandez. selscan: an efficient multithreaded program to perform
EHH-based scans for positive selection. *Mol Biol Evol* 31, 2824–2827 (2014).
doi.org/10.1093/molbev/msu211

28. A. Ferrer-Admetlla, M. Liang, T. Korneliussen, R. Nielsen. On detecting incomplete soft or
hard selective sweeps using haplotype structure. *Mol Biol Evol* **31**, 1275–1291 (2014).
doi.org/10.1093/molbev/msu077
29. P. C. Sabeti, P. Varilly, B. Fry, J. Lohmueller, E. Hostetter, C. Cotsapas, X. Xie, E. H. Byrne,
S. A. McCarroll, R. Gaudet, S. F. Schaffner, E. S. Lander, International HapMap
Consortium. Genome-wide detection and characterization of positive selection in human
populations. *Nature* **449**, 913–918 (2007). doi.org/10.1038/nature06250

Fig. 1

Fig. 1. Identification of a causal SNP of *qSH3* associated with the degree of seed shattering.
(*A*) Fine mapping of the causal mutation at *qSH3* in a 1,228-bp region on chromosome 3. A causal SNP (SNP-70) identified in *OsSh1* (*Os03g0650000*) is shown in bold. (*B*) Comparison of seed-shattering degree by breaking tensile strength (BTS) values in three introgression lines, IL(*qSH3*-W), IL(*qSH1*-W), and IL(*qSH1*-W+*qSH3*-W) in the Nipponbare genetic background. Data are mean ± s.d. of four plants. n.s. and ** indicate not significant and significant at the 1% level based on unpaired Student's t-test, respectively. (*C*) Two types of constructs carrying *qSH3* cDNA sequences of W630 and Nipponbare (*qSH3^{W630}* and *qSH3^{Npb}*) driven by a 3-kb region of the promoter used for transgenic analysis. (*D*) The BTS values observed for the transformants with *qSH3^{W630}* and *qSH3^{Npb}*. Black triangles represent the average BTS value of IL(*qSH1*-W).

Fig. 2

Fig. 2. Lineage-specific selection at *qSH3* locus in rice.
(*A*) Genotyping at *qSH3*, *sh4*, and *qSH1* for *O. rufipogon* W630, *O. sativa japonica* Nipponbare, *indica* IR36, and *circum*-aus 'Kasalath' based on the causal SNPs by dCAPS markers. (*B*) Allele frequency of *qSH3* causal SNP (%) in cultivated rice based on the 3K rice project data. Nipponbare type ('T') and W630 type ('C') are shown in yellow and green, respectively. (*C*) Nucleotide diversity (π) observed for the domesticated rice in the physical position of 25.0-25.3 Mb on chromosome 3. A flanking region of the *qSH3* locus (around 25.2 Mb) was substantially decreased in *japonica* and *indica* cultivars. π was calculated in 10-kb windows using SNPs data of the 3K rice project (13).

Fig. 3

Fig. 3. Abscission layer formation partially inhibited by *sh4* and *qSH3* in wild rice.
Abscission layer formation in *O. rufipogon* W630, IL(*sh4*-N), IL(*qSH3*-N), IL(*sh4*-N+*qSH3*-N), and *O. sativa* Nipponbare. Each enlarged section indicated by the dotted square is shown in the lower panel. A black triangle indicates an inhibited area of the abscission layer observed for IL(*sh4*-N+*qSH3*-N). Scale bars = 100 μm.

Fig. 4

Fig. 4. Role of the *sh4, qSH3*, and *SPR3* mutations on the initial loss of seed shattering in rice domestication.
(*A*,*B*) Panicle shape of the seven ILs at flowering (*A*) and seed-maturation (*B*) stages. Scale bars = 5 cm (*C*) Breaking tensile strength (BTS) values for the seven ILs. BTS values for *O. sativa japonica* Nipponbare and *indica* IR36 are shown as control. n.s. indicates not significant by unpaired Student's t-test. (*D*) Seed-gathering rates (mean ± s.d. of four plot replicates, with nine plants in each plot) for *O. rufipogon* W630 and the seven ILs. Mean values labelled with different letters are significantly different, whereas those with the same letters are not (Tukey's test with arcsine transformed values, *P* < 0.05).

Fig. 5

Fig. 5. Structural mechanics analysis of panicle shape and abscission layer inhibition for the initial loss of seed shattering.
(*A*), Panicle shape of *O. rufipogon* W630. Bar = 5cm. (*B*) Schematic representation of awn and grain with panicle angle. θ represents the panicle angle. l_1, l_2, m_1g, and m_2g indicate the length of grain, awn, weight of grain and awn, respectively. (*C*) Detachment of grain and pedicel (left) and scanning electron microscopy (SEM) analysis of the abscission layer of pedicel in *O. rufipogon* W630 (right). AL and VB indicate abscission layer and vascular bundle, respectively. Bars = 1 mm (left) and 100 μm (right). (*D*) Schematic representation of the abscission layer. *D* and *d* indicate the diameter of abscission layer and vascular bundles, respectively. Dotted circle indicates the area of disrupted abscission layer. (*E*) Simulation of the bending stress exerted on the spikelet base depending on panicle shape and abscission layer inhibition, with higher stress experience in open panicle with less inhibited abscission.