1   Title page

Title:

Obesity affects mitochondrial metabolism and proliferative potential of equine endometrial progenitor

cells.

Running head:

Obesity-induced deterioration of metabolism in equine endometrial progenitor cells.

Keywords:

obesity; endometrium; horse progenitor cells; metabolism; proliferation

Authors:

Agnieszka Smieszek[1,*], Klaudia Marcinkowska[1], Ariadna Pielok[1], Mateusz Sikora[1], Lukas Valihrach[2],
Elaine Carnevale[3] and Krzysztof Marycz[1,4]

[1]Department of Experimental Biology, The Faculty of Biology and Animal Science, University of
Environmental and Life Sciences, 50-375 Wroclaw, Poland;

[2]Laboratory of Gene Expression, Institute of Biotechnology CAS, Biocev, 25250 Vestec, Czech
Republic;

[3] Equine Reproduction Laboratory, Department of Biomedical Sciences, Colorado State University, Fort
Collins, CO 80523-1693, United States of America;

[4]International Institute of Translational Medicine, Jesionowa 11 St, 55-124 Malin, Poland

*Correspondence: agnieszka.smieszek@upwr.edu.pl,

Abstract:

The study aimed to investigate the influence of obesity on cellular features of equine endometrial progenitor cells (Eca EPCs), including viability, proliferation capacity, mitochondrial metabolism, and oxidative homeostasis. Eca EPCs derived from non-obese (Non-OB) and obese (OB) mares were characterised by cellular phenotype and multipotency.

Obesity-induced changes in the activity of Eca EPCs include the decline of their proliferative activity, clonogenic potential, mitochondrial metabolism and enhanced oxidative stress. Eca EPCs isolated from obese mares were characterised by an increased occurrence of early apoptosis, loss of mitochondrial dynamics, and senescence-associated phenotype. Attenuated metabolism of Eca EPCs OB was related to increased expression of pro-apoptotic markers (*CASP9, BAX, P53, P21*), enhanced expression of OPN, PI3K and AKT, simultaneously with decreased signalling stabilising cellular homeostasis (including mitofusin, SIRT1, FOXP3).

Obesity alters functional features and the self-renewal potential of endometrial progenitor cells. The impaired cytophysiology of progenitor cells from obese endometrium predict lower regenerative capacity if used as autologous transplants.

1. Introduction

As is widely known, contributes to an increased risk of cardiovascular disease, type 2 diabetes, and stroke. However, there is also growing body of evidence supporting the negative influence of obesity on reproduction, including infertility, low quality of oocytes, increased early pregnancy loss[1]. Additionally, obesity reduces the success of attaining pregnancy, even when accompanied by assisted reproductive technology. It was also shown that obesity is strongly associated with the development of endometrial cancer. The endometrial environment has an essential role in obesity-related reproductive impairments, but molecular mechanisms by which obesity contribute to the degeneration of the endometrium niche are still investigated. However, mitochondrial dysfunction is highlighted as a significant link between cellular metabolic imbalance and reproductive problems[2].

In this study, we have incorporated the equine model to study the effect of obesity on the cytophysiology of endometrial progenitor cells.

From studies conducted in Europe, North America, Australia and New Zealand, 2 to 72% of horses are considered overweight, and 1 to 19% of animals are obese[3,4]. Obesity contributes to developing metabolic disorders, such as insulin resistance, equine metabolic syndrome (EMS), and laminitis - all conditions accompanied by inflammation. Infertility and altered reproductive function of mares are also consequences of obesity [4,5]. Maternal obesity also influences foal metabolism and increases the occurrence of low-grade inflammation and osteochondrosis in yearlings[3]. The effect of obesity on the equine uterus has not been well defined, although gene expression in the endometrium was altered even after short-term obesity, showing a proinflammatory profile with increased expression of TNFα and IL1β [6].

Excessive accumulation and expansion of adipose tissue affect the endogenous self-renewal capacity
of the organism[7]. Our previous studies have demonstrated that equine metabolic syndrome affects the
regenerative potential of adipose-derived stromal/stem cells (ASCs)[8–11]. Progenitor cells, isolated
from horses with metabolic syndrome, have limited clinical potential compared to healthy subjects
because of reduced proliferation capacity, disturbed cellular homeostasis, and hampered multipotency.
The physiological status and age of the horse affect the metabolism of progenitor cells. Ageing and
equine metabolic syndrome reduce the self-renewal potential of progenitor cells, significantly impairing
mitochondrial function while increasing autophagy, oxidative stress, and endoplasmic stress[8–10]. The
lowered regenerative features of progenitor cells, resulting from obesity-induced cellular damage,
notably limits therapies based on autologous transplants[12].

The biology of equine multipotent stromal/stem cells (MSCs) isolated from adipose tissue and bone
marrow is well described. These cells are commonly used to treat equine locomotive disorders[12,13].
However, since MSCs keep an epigenetic memory specific to the tissue of origin, other adult tissues are
being explored as potential sources for progenitor cells[14]. Recently, attention has been given to the
cytophysiology of progenitor cells originating from the endometrium [14–16]. The endometrium has
increased regenerative potential, which is needed for remodelling during the oestrous cycle. Progenitor
cells from the uterus can be harvested by a biopsy, which is a nonsurgical procedure without the need
for anaesthesia in mares. The endometrium provides an attractive source of easily obtained and highly
proliferating cells [14,15,17].

Cells isolated from equine endometrial tissue were previously characterised as a multipotent population
with enhanced self-renewal properties [15]. Uterine pathologies, such as post-breeding endometritis and
endometriosis lead to a substantial economic burden for the equine industry [18]. Thus, therapies based
on the application of endometrial progenitor cells are appealing.

The main objective of this study was to evaluate the effects of obesity on basic cytophysiological features
of equine endometrial progenitor cells (Eca EPCs), including viability and self-renewal potential, as well
as mitochondrial metabolism. We compared proliferation capacity, mitochondrial dynamics, and
molecular phenotypes of EPCs obtained from non-obese (Non-OB) and obese mares (OB).

103
2. Materials and methods

2.1. Harvesting and isolation of equine endometrial progenitor cells (Eca EPCs) from obese and
non-obese mares

Tissue samples were obtained post-mortem from non-obese (n=4) and obese (n=4) mares during
anestrus. Uteri were collected from a local slaughterhouse at Rawicz in November and December of
2019 and 2020, and samples were isolated from the uterine body. The mean age of non-obese mares
was 9.5 ± 2, while obese mares were 9± 1. The obesity of mares was determined based on body
condition score (BCS), using the protocol established previously [10,19] and a system developed by
Henneke *et al*. [20]. BSC is expressed by numerical value for fat deposition ranging from 1 (poor) to 9
(extremely obese). The horses in the non-obese group had BSC = 6.5 ± 0.289 (mean+SEM), while
obese mares were characterised by BCS = 8.75 ±0.250 (mean+SEM).

The tissue fragments (~ 4g) were washed three times using Hank's balanced salt solution (HBSS)
supplemented with 1% of penicillin/streptomycin mix (P/S, Sigma Aldrich/Merck, Poznan, Poland). The

endometrium was dissociated from myometrium, cut into small fragments, and placed into an enzymatic
solution. The solution consisted of collagenase type I (1 mg/ml) prepared in Dulbecco's modified Eagle's
medium/F12 (DMEM/F12, Sigma Aldrich/Merck, Poznan, Poland). The tissue fragments were digested
for 40 minutes in CO_2 incubator at 37 °C with periodic vortexing. After incubation, the homogenates
were filtered using a sterile 70-μm cell strainer (Greiner Bio-One, Biokom, Janki, Poland) to discard the
undigested tissue fragments. Further, the suspension of cells was centrifuged at 300×g for 10 minutes.
Obtained pellets were washed in HBSS and then resuspended in DMEM/F12 medium supplemented
with 15% of foetal bovine serum (FBS, Sigma Aldrich/Merck, Poznan, Poland) and 1% P/S solution,
hereinafter referred to as a complete growth medium for endometrial stromal cells (CGM$_{EPC}$). The cells
were transferred into a T-75 culture flask (Nunc, Biokom, Janki, Poland). Both primary and subsequently
passaged cultures were cultured at constant conditions in CO_2 incubator at 37°C, and maintaining 95%
humidity. The CGM$_{EPC}$ was changed every 2 days. The culture was passaged when reaching 90%
confluence using a trypsin solution (StableCell Trypsin, Sigma Aldrich/Merck, Poznan, Poland). Cells
used for the experiment were pulled at passage 3 (p3).

2.2. Phenotype and multipotency of Eca EPCs

The phenotype of cells was determined based on mRNA expression of (i) the mesenchymal markers,
including CD29, CD44, CD90, CD105; (ii) hematopoietic markers, i.e. CD34 and CD45; (iii) perivascular
markers, including CD146 and NG2; (iv) an epithelial marker, i.e. mucin-1 and (v) smooth muscle
markers, including actin alpha 1 (ACTA2), calponin 1 (CNN1) and myosin heavy chain 11 (MHY11).
The expression of selected genes was determined using the RT-qPCR protocol described in paragraph
2.6. The sequences of oligonucleotides were published by Rink et al. [15] and were fully characterised
in Table 1S. Amplified PCR products were separated in 2% agarose gel and stained with SYBR™ Safe
DNA Gel Stain Thermo Fisher Scientific, Warsaw, Poland. The molecular weight of the obtained
products was compared to DNA Ladder (M50pz, Blirt DNA, Gdansk, Poland).

Immunocytochemical staining aimed at detecting CD44, CD45, CD73, CD90 and CD105 surface
markers is described in paragraph 2.8. Moreover, expression of CD34, CD45, CD90 and CD105 was
detected using flow cytometry. The analysis was performed using FACSLyric™ Clinical Flow Cytometry
System (Becton-Dickinson and Company, Franklin Lakes, NJ, USA). The detailed staining protocol was
described elsewhere [21], while antibodies used for analysis are listed in Table 4S.

To verify the multipotency, Eca EPCs were cultured under osteogenic, chondrogenic and adipogenic
conditions. For this purpose, cells (p3) were seeded into 6-well plates at inoculum $1x10^5$ cells per well.
The differentiation was induced using commercially available kits (StemPro®, Thermo Fisher Scientific,
Warsaw, Poland) after 48 hours of culture when cells reached 90% confluency. The effectiveness of
tissue-specific differentiation was evaluated after 21 days in osteogenic cultures and after 16 days in
chondrogenic and adipogenic cultures. Following differentiation, the extracellular matrix was stained
accordingly to well-established protocols[22]. Osteogenesis in vitro was verified with Alizarin Red
staining. The chondrogenic nodules were detected with Safranin O. In contrast, lipid droplets in
adipogenic cultures were detected after Oil Red O staining. Documentation of cultures was performed
using an inverted microscope (Axio Observer A.1, Zeiss, Oberkochen, Germany) with Power Shot digital

camera (Canon, Woodhatch, UK). Images were processed and analysed, as was previously described
[21].

2.3. Evaluation of proliferation and migratory capacity of cells

Proliferation activity of cells was evaluated based on multiple parameters established in a clonogenic
and wound healing assay, MTS assay, and cell cycle analysis. The protocols were used previously for
progenitor cells of different origins and were published in detail [11,23,24]. In this experiment, the
efficiency of colony-forming unit (CFU-E) occurrence was evaluated after 12 days of culture using the
protocol described by Rink et $al.$[15]. Cells were inoculated in 6-well plates at a density equal 10
cells/cm^2 in 2 mL of CGM$_{EPC}$. After the test, cultures were fixed in 4% of paraformaldehyde (PFA) for 30
minutes, subsequently washed using HBSS, and stained with 2 % pararosaniline solution (Sigma
Aldrich/Merck, Poznan, Poland) for 5 minutes, before being rewashed with HBSS. Colonies were
defined as clusters formed by more than 50 cells.

For the wound healing assay, cultures were maintained at a high confluence, reaching 90%. The scratch
was made with a 200-μl pipette tip, and cultures were propagated for an additional 24 hours. After that,
cultures were fixed with 4% PFA and stained with the pararosaniline solution. Cultures were documented
using a Canon PowerShot digital camera (Woodhatch, UK).

The population doubling time (PDT) was determined based on the growth curve of Eca EPCs,
established by a routine counting of cells during the passage. The cells were counted using the Muse$^®$
Count & Viability Kit (Luminex/Merck, Poznan, Poland) according to the manufacturer's protocol and
analysed using Muse Cell Analyser (Sigma-Aldrich/Merck, Poznan, Poland). Obtained data were used
to determine PDT with an algorithm published by Heuer et al.[25] and Cell Calculator [26].

The proliferation determined based on metabolic activity was measured using MTS (Cell Proliferation,
Colorimetric, Abcam, Cambridge, UK). For this purpose, Eca EPCs were seeded in 96-well plates at a
density equal to 5x10^3 per well in 200 µl CGM$_{EPC}$. After 24 h of cells inoculation, 20 µl of MTS solution
was added per well. The test included blank control. Cultures were incubated with the dye for 2 h at
37°C in a CO$_2$ incubator. Following incubation, absorbance was measured spectrophotometrically with
a plate reader (Epoch, Biotek, Germany) at a wavelength of 490 nm. MTS assay was performed after
24, 48 and 72 h of Eca EPCs culturing.

The cell cycle was determined using Muse Cell Cycle Kit (Luminex/Merck, Poznan, Poland) using the
manufacturer's protocol. The analysis was performed using Muse Cell Analyser (Sigma-Aldrich/Merck,
Poznan, Poland), and for each assay, 0.5×10^6 of Eca EPCs were tested.

The proliferation activity of Eca EPCs from non-obese and obese mares was also evaluated based on
immunocytochemical staining of Ki67 marker and miRNA levels. A transwell system with a pore size of
8 µm (Corning, Biokom, Warsaw, Poland) was used to test the motility capabilities of Eca EPCs. Cells
were inoculated into the upper chamber at a density equal to 0.2 ×10^5 in a 300 μl of serum-free culture
medium (DMEM/F12 supplemented 1% P/S). The CGM$_{EPC}$ was added to the lower compartment of the
transwell system. The cells were stained with pararoseline solution, as described above. Chambers
were evaluated under an inverted microscope (Leica DMi1), with an attached Digital Camera MC170
and Leica Application Suite (LAS) software (all from Leica Microsystems, KAWA.SKA Sp. z o.o., Zalesie
Gorne, Poland).

2.4. The evaluation of Eca EPC growth pattern, morphology and ultrastructure

The growth pattern and morphology of cultures were evaluated using an inverted microscope (Leica DMi1) equipped with a camera MC170 (Leica Microsystems, KAWA.SKA Sp. z o.o., Zalesie Gorne, Poland). Cell condition was also determined from images obtained after senescence-associated beta-galactosidase staining. The ultrastructure of Eca EPCs was analysed using a confocal microscope (Leica TCS SPE, Leica Microsystems, KAWA.SKA Sp. z o.o., Zalesie Gorne, Poland) based on localisation and distribution of nuclei, cytoskeleton and mitochondrial network. The mitochondria were stained in vital cultures with MitoRed dye (Sigma-Aldrich/Merck, Poznan, Poland) prepared in CGM_{EPC} (1:1000). Subsequently, cultures were incubated with the dye in CO_2 incubator at 37°C for 30 minutes. Next, cells were fixed with 4% paraformaldehyde (PFA) for 30 minutes at room temperature and rinsed three times with HBSS. Following fixation, the actin cytoskeleton was imaged using Phalloidin-Atto 488 (Sigma-Aldrich/Merck, Poznan, Poland). For this purpose, cells were permeabilised with 0.2% Tween 20 and prepared in HBSS for 20 minutes at room temperature. After permeabilisation, cells were stained with phalloidin solution dissolved in HBSS at a concentration of 1:800. The staining lasted 30 minutes and was performed at 37°C in the dark. Cells nuclei were counterstained using 4′,6-Diamidine-2′-phenylindole dihydrochloride (DAPI) mounting medium (ProLong™ Diamond Antifade Mountant with DAPI, Thermo Fisher Scientific, Warsaw, Poland). Preparations were observed using the confocal microscope under magnification 630× and processed with Fiji software (ImageJ 1.52n, Wayne Rasband, National Institute of Health, USA). Additionally, to evaluate the mitochondrial dynamics, the cells were visualised under 1000× and analysed using MicroP software [27].

To determine mitochondrion gene expression, mitochondria were isolated using the Mitochondria Isolation Kit for Cultured Cells (Thermo Fisher Scientific, Warsaw, Poland). All steps of the procedure were performed following the manufacturer's instructions. The mitochondria were isolated from 2×10^7 cells. The obtained mitochondria were homogenised with 1 ml of Extrazol® (Blirt DNA, Gdansk, Poland).

2.5. Evaluation of Eca EPCs viability, metabolism and oxidative status

The cellular health of Eca EPCs Non-OB and Eca EPCs OB was evaluated using Guava® Muse® Cell Analyser (Sigma-Aldrich/Merck, Poznan, Poland). The metabolic activity of cells was determined based on mitochondrial membrane potential using Muse™ Mitopotential Assay Kit. Apoptosis profile was established in the test with Muse® Annexin V & Dead Cell Kit. In addition, oxidative status was evaluated based on the intracellular accumulation of reactive oxygen species (Muse® Oxidative Stress Kit) and nitric oxide (Muse® Nitric Oxide Kit). All assays were performed according to protocols provided by the manufacturer (Luminex/Merck, Poznan, Poland) and were described in detail previously [28–30].

2.6. Determination of transcripts levels

The experimental cultures of Eca EPCs and isolated mitochondria were homogenised using Extrazol® (Blirt DNA, Gdansk, Poland). Total RNA and mitochondrial RNA was isolated using the phenol-chloroform method. Ther genomic DNA (gDNA) residuals were removed from samples with DNase I from PrecisionDNAse kit (Primerdesign, Blirt DNA, Gdansk, Poland). Purified RNA (500 ng) was transcribed into cDNA with Tetro cDNA Synthesis Kit (Bioline Reagents Limited, London, UK) to analyse genes expression. Both digestion of gDNA and reverse transcription (RT) was performed using T100 Thermal Cycler (Bio-Rad, Hercules, CA, USA). Quantitative PCR (qPCR) was performed using 1 μl of

cDNA and specific primers (sequences listed in Table 1S; Supporting Information). Two-tailed RT-qPCR

was used for specific detection of miRNA levels. The exact conditions of reactions were described

elsewhere [11], while primers are listed in Table 2S (Supporting Information). All qPCR measurements

were carried out using CFX Connect™ Real-Time PCR Detection System (Bio-Rad, Hercules, CA,

USA). The obtained gene expression results were calculated using RQ_{MAX} algorithm and converted into

the log2 scale as described previously [11,24,30].

2.7. Detection of protein expression with Western blot

To determine the intracellular accumulation of selected proteins, cells were lysed using ice-cold RIPA

buffer containing 1% of protease and phosphatase inhibitor cocktail (Sigma Aldrich/Merck, Poznan,

Polska). The total concentration of proteins in samples was determined using the Bicinchoninic Acid

Assay Kit (Sigma Aldrich/Merck, Poznan, Polska). The samples containing 20 μg of protein were mixed

with 4× Laemmli loading buffer (Bio-Rad, Hercules, CA, USA). Subsequently, samples were incubated

for 5 minutes at 95 °C in T100 Thermal Cycler (Bio-Rad, Hercules, CA, USA). Samples were separated

in 12.5 % sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE; 100V for 90

minutes) in Mini-PROTEAN Tetra Vertical Electrophoresis Cell (Bio-Rad, Hercules, CA, USA) and

transferred into polyvinylidene difluoride (PVDF) membrane in 1× Transfer buffer (Tris-

base/Glycine/Methanol, Sigma-Aldrich/Merck, Poznan, Poland) using the Mini Trans-Blot® system (Bio-

Rad, Hercules, CA, USA) at 100 V for 60 minutes. The membranes were blocked using 5% bovine

serum albumin (BSA) in TBS-T for 60 minutes. Detection of protein was performed by overnight

incubation at 4°C with primary antibodies. The incubation of membranes with secondary antibodies was

performed for 60 minutes at 4°C. After each incubation with the antibody, the membranes were washed

five times for 5 minutes with TBST buffer.

Details of antibodies used for the reaction are presented in Table 3S. Chemiluminescent signal was

detected using Bio-Rad ChemiDoc™ XRS system (Bio-Rad, Hercules, CA, USA) using DuoLuX®

Chemiluminescent and Fluorescent Peroxidase (HRP) Substrate (Vector Laboratories, Biokom, Janki,

Poland;). The Image Lab™ Software (Bio-Rad, Hercules, CA, USA) was used in order to analyse the

molecular weight and intensity of signals.

2.8. Immunocytochemical detection

Immunocytochemical(ICC) staining was performed using previously described protocols[24,31]. Here,

Eca EPCs cultures were performed in 24-well dishes covered by glass cover slides. Cultures were fixed

with 4% PFA after 72 hours of propagation. The incubation was performed 30 minutes at room

temperature. Subsequently, specimens were washed three times using HBSS and prepared for

permeabilisation with 0.2% PBS-Tween solution supplemented with 10% goat serum. Specimens were

permeabilized for 1h. The incubation of samples with the primary antibody in HBSS was performed

overnight at 4 °C. The secondary antibody was used at a concentration equal to 1:1000 in HBSS. Details

of antibodies used for the reaction are presented in Table 4S. Each step of the protocol was preceded

by gentle washing of specimens with HBSS. After staining, slides were fixed with a mounting medium

with DAPI (ProLong™ Diamond Antifade Mountant with DAPI, Thermo Fisher Scientific, Warsaw,

Poland). Observations were made using a confocal microscope (Leica TCS SPE, Leica Microsystems,

KAWA.SKA Sp. z o.o., Zalesie Gorne, Poland). Images were processed and analysed, as was
previously described [24].

2.8. Statistical analysis

The results obtained in the study are presented as the mean with standard deviation (±SD). The values
derived from at least three technical repetitions. The normality of the population data was determined
using Shapiro–Wilk test, while equality of variances was assessed by Levene's test. The measurements
were conducted with STATISTICA 10.0 software (StatSoft, Inc., Statistica for Windows, Tulsa, OK,
USA). Comparative analysis was performed using t-Student test or One-way analysis of variance with
Dunnett's post hoc test. These calculations were performed using GraphPad Software (Prism 8.20, San
Diego, CA, USA). Differences with a probability of $p< 0.05$ were considered significant.

3. Results

The animals used for the experiment were classified using a system proposed by Henneke *et al.*[20].
Significant differences in terms of BSC between non-obese (Non-OB) and obese (OB) groups were
noted, warranting proper classification of animals (**Fig.1S**).

Furthermore, we have evaluated the phenotype of the cells used for the study and tested their
multipotent character. The mesenchymal stem cells markers were assessed using RT-qPCR and with
immunocytochemistry staining (**Fig. 1**). The population of progenitor cells isolated from equine
endometrium (Eca EPCs) consisted of cells with features characteristic for cells of mesenchymal origin,
i.e. expression of markers CD44, CD90 and CD105 and low expression of hematopoietic markers, i.e.
CD34 and CD45 (**Fig. 1 b-c**). Progenitor cells from the endometrium of non-obese mares and obese
mares (Eca EPCs Non-OB and Eca EPCs OB) differed in CD105 expression (**Fig. 1**). The Eca EPCs
OB showed decreased levels of mRNA for CD105. Moreover, the reduced transcript levels for CD105
noted in Eca EPCs OB correlated with a decreased mRNA expression for CD90 (**Fig. 1 a-b**). Both Eca
EPCs Non-OB and OB showed mRNA expression for perivascular markers CD146 and NG2
proteoglycan (neural/glial antigen 2), an epithelial marker, i.e. mucin-1 and smooth muscle markers,
including actin alpha 1 (ACTA2), calponin 1 (CNN1) and myosin heavy chain 11 (MHY11). Notably, Eca
EPCs OB were characterized by increased accumulation of transcripts for ACTA2 (**Fig. 1 a-b**).

(a)

(b)

(c)

Fig.1. Phenotype of Eca EPCs. The RT-qPCR analysis aimed to determine relative transcript levels of
cell surface markers (a). The mRNA expression was established for MSCs markers (CD29, CD44,
CD90, CD105), perivascular markers (NG2, CD146), epithelial marker (MUC-1) and smooth muscle
markers (ACTA2, CNN1, MHY11). The mRNA expression for hematopoietic markers (CD34, CD45) was
not included on the graphs due to the lack of a specific signal. The PCR products were analysed using
electrophoresis (b). The cellular phenotype was tested using immunocytochemistry (c). All results are
shown as mean ± SD. Columns with bars represent means ± SD. * p-value < 0.05, **** p-value < 0.0001.
317

319 Additionally, Eca EPCs used in the study underwent specific differentiation under osteogenic,

320 chondrogenic and adipogenic conditions, indicating their multipotency (**Fig. 2**). However, the analysis

321 of the extracellular matrix showed that Eca EPCs OB had lowered osteogenic and chondrogenic

322 potential compared to Eca EPCs Non-OB (**Fig.2 a-c**). The adipogenic potential of Eca EPCs OB was

323 also decreased (**Fig.2 d**).

(a)

(b) **(c)** **(d)**

Fig. 2. The analysis of Eca EPCs multipotency. Tissue-specific differentiation extracellular matrix
features were documented (a). Cultures were maintained under osteogenic, chondrogenic and
adipogenic conditions. Calcium deposits formed under osteogenic conditions were detected using
Alizarin Red staining, while chondrogenic nodules were stained with Safranin-O dye. The lipid-rich
vacuoles after adipogenic stimulation were detected with Oil Red O. The staining efficiency was
measured to compare the differentiation potential of Eca EPCs Non-OB with Eca EPCs OB (b-d).
Columns with bars represent means ± SD. **** p-value < 0.0001.

Furthermore, the progenitor cells isolated from the endometrium of obese mares (Eca EPCs OB) were

characterised by decreased proliferative potential. In contrast, endometrial progenitor cells isolated from

non-obese mares (Eca EPCs Non-OB) showed faster initial growth and enhanced proliferation capacity

(**Fig. 3**). The lowered self-renewal potential of Eca EPCs OB was demonstrated by a significantly

reduced ability to form colony-forming units (CFUs) (**Fig 3 a and b**). The migratory and wound healing

assay revealed that Eca EPCs from OB were characterised by lowered expansion potency (**Fig. 3 c-f**),

confirming lower CFU efficiency. Reduced proliferation of Eca EPCs OB in culture was reflected by the

prolonged time needed for cell population doubling (PDT; **Fig. 3g**). The metabolic activity of EPCs OB

was also disturbed compared to Eca EPCs from non-obese mares. The significant difference between

the metabolism of Eca EPCs from non-obese and obese mares was noted after 48 hours of culture (**Fig.**

3h).

Cell cycle analysis confirmed that Eca EPCs OB had decreased potential for division (Fig 2).

The intracellular accumulation of Ki67, a marker of actively proliferating cells, was significantly

decreased in EPCs Ob cultures (**Fig. 4 a-c**). Furthermore, the percentage of actively proliferating cells

in Eca EPCs OB cultures, evidenced by a decreased S-phase ratio, shifted toward G0/G1-phase (**Fig.**
4 d and e). Reduced Ki67 protein expression and proliferation were correlated with the lower levels of
microRNAs from the let7 family, known regulators of genes associated with proliferation of progenitor
cells, i.e. let-7b and let-7c. However, let-7a expression was increased in Eca EPCs OB (**Fig. 4 f –h**).
352

353
354
355 **Fig. 3:** The proliferative capacity of equine endometrial progenitor cells (Eca EPCs) isolated from non-
356 obese mares (Non-OB) and obese mares (OB), expressed by colony formation capability (a, b),
357 migratory efficiency (c-f), population doubling time (g), metabolic activity (h). Columns with bars
358 represent mean ± SD. * p-value < 0.05, ** p-value < 0.01 and *** p-value < 0.001.
359

Fig. 4: The proliferative capacity of equine endometrial progenitor cells (Eca EPCs) isolated from non-obese mares (Non-OB) and obese mares (OB) evaluated based on Ki67 expression (a-c), distribution of cells within the cell cycle (d-e), and levels of particular let7 family members. Columns with bars represent means ± SD. * p-value < 0.05, ** p-value < 0.01 and *** p-value < 0.001.

Analysis of cell morphology and ultrastructure showed that Eca EPCs isolated from non-obese and obese mares showed fibroblast-like, spindle-shaped morphology (**Fig. 5 a and b**). Eca EPCs isolated from non-obese mares formed dense colonies in which well-developed intracellular connections were observed. In contrast, Eca EPCs OB formed less abundant aggregations, and intracellular connections between those cells were less marked. The actin cytoskeleton, as well as mitochondrial network, were more developed in EPCs from non-obese mares. The identification of senescence cells revealed that Eca EPCs from OB mares show significantly increased activity of β-galactosidase (**Fig. 5 f**). Additionally, mRNA levels for vimentin (*VIM*), type III intermediate filament (IF) were decreased in Eca EPCs OB (**Fig. 5 c**). The lowered transcript levels of *VIM* noted for Eca EPCs OB, were correlated with decreased levels of miRNA-29a-3p and miR-181-5p, which function as regulators of cytoskeleton assembly and self-renewal of stem cells (**Fig. 5 d and e**).

Fig. 5: Morphology, ultrastructure, the growth pattern of equine endometrial progenitor cells (Eca EPCs)
isolated from non-obese mares (Non-OB) and obese mares (OB), and analysis of transcripts associated
with a cytoskeletal network. The cultures were imaged with a confocal microscope to determine
ultrastructure. Growth pattern of cell cultures was monitored under a phase-contrast microscope. Scale
bars are indicated in the representative photographs (a and b). Additionally, RT-qPCR was performed
to determine the mRNA expression of vimentin (*VIM*; c) and levels of miR-29a-3p and miR-181-5p (d
and e). Results from RT-qPCR are presented as a column with bars representing means ± SD. ** p-
value < 0.01 and *** p-value < 0.001.

A decreased number of mitochondria were observed in Eca EPCs derived from obese mares. Moreover,
these cells were distinguished by the presence of respiratory deficient, globular mitochondria (**Fig. 6**).
Further, analysis of mitochondrial membrane potential (MMP) confirmed the lowered metabolism of Eca
EPCs from obese mares (**Fig 7**). The significantly decreased mitochondrial metabolism was also
correlated with declining viability of cells. Gene expression patterns confirmed that metabolism of
mitochondria in EPCs Ob was disturbed, with downregulation of transcripts essential for mtDNA function
and the mitochondrial respiratory chain stability (**Fig. 8**).
Analysis of parkin RBR E3 ubiquitin-protein ligase (PARKIN), PTEN-induced kinase 1 (PINK1)
and mitofusin 1 (MFN1) expression was performed at mRNA and protein level, both in Eca EPCs Non-
OB and OB. The mRNA level of *PARKIN* was decreased, while PARKIN protein was accumulated within
Eca EPCs OB. In contrast, PINK mRNA levels were elevated, but protein expression was lower in Eca
EPCs OB. However, mitofusin expression was decreased in EPCs OB, both at mRNA and protein levels.
Transcript levels determined for fission (*FIS*) were also reduced in Eca EPCs OB compared to Eca
EPCs derived from non-obese mares (**Fig. 9**).

403
404
405 **Fig. 6:** Mitochondrial network, numbers, and morphology of equine endometrial progenitor cells (Eca
406 EPCs) isolated from non-obese mares (Non-OB) and obese mares (OB). Imaging of Eca EPCs with
407 high magnification allowed assessment of the mitochondrial net and its dynamics (a-b), number of
408 mitochondria per cell (c) and classification of mitochondrial morphology (d-h). Means ± SD are presented
409 as columns and bars. *** p-value < 0.001.
410

(a)
(b)
(c)
(d)

411
412 **Fig.7:** The mitochondrial membrane potential (MMP) determined in equine endometrial progenitor cells
413 (Eca EPCs) isolated from non-obese mares (Non-OB) and obese mares (OB) The representative graphs
414 show the distribution of cells, taking into account their viability and mitochondrial membrane
415 depolarisation (a and b). Comparative analysis was performed to determine differences between EPCs
416 Non-Ob and Ob in terms of viability (c) and mitochondrial activity (d). Means ± SD. *** p-value < 0.001.
417

418
Fig. 8: Transcript levels for genes associated with mitochondria homeostasis determined for equine
endometrial progenitor cells (Eca EPCs) isolated from non-obese mares (Non-OB) and obese mares
(OB). The analysis was performed using RT-qPCR technologies. The following genes were measured:
ubiquinol-cytochrome c reductase core protein 2 (UQCRC2), transcription factor A, mitochondrial
(TFAM), pseudouridylate synthase-like 1 (PUSL1), NADH: ubiquinone oxidoreductase subunit A9
(NDUFA9), cytochrome c oxidase subunit 4I1 (COX4I1), PIGB opposite strand 1 (PIGBOS1),
mitochondrial ribosomal protein L24 (MRPL24), mitochondrial transcription termination factor 4
(MTERF4), mitochondrial inner membrane protein (OXA1L) and PPARG coactivator 1 beta
(PPARGC1β). Obtained data were normalised to the expression of the reference gene and expressed
as using RQ(max) algorithm. Results are presented as columns with bars representing means ± SD. ***
p-value < 0.001.

430
431
Fig. 9: The expression profile of molecular markers associated with mitochondrial dynamics determined
for equine endometrial progenitor cells (Eca EPCs) isolated from non-obese mares (Non-OB) and obese
mares (OB). Parkin RBR E3 ubiquitin-protein ligase (PARKIN), PTEN-induced kinase 1 (PINK1) and
mitofusin 1 (MFN1) were determined based on mRNA (b, d, f) and protein levels (a- representative blots,
c, e, g), additionally transcript levels for fission (FIS) were determined (h). Means ± SD. *** p-value <
0.001.

Apoptosis profile analysis supported the result indicating lower viability of Eca EPCs derived
from obese than non-obese mares (**Fig. 10**). The Eca EPCs OB showed an increased percentage of
early apoptotic cells compared to Eca EPCs Non-OB (**Fig. 10 a, b, e**). Although we observed increased
early apoptosis in EPC cultures obtained from obese mares, the number of total apoptotic cells were
not different for OB and Non-OB (**Fig. 10 a, b, d**). EPCs OB showed a pro-apoptotic gene expression
pattern, with an elevated *BAX/BCL-2* ratio and increased mRNA expression for caspase-9 (*CASP-9*),
P53 and *P21 (***Fig. 11***)*.

446
447
Fig. 10: The apoptosis profile determined in equine endometrial progenitor cells (Eca EPCs) isolated
from non-obese mares (Non-OB) and obese mares (OB). The representative graphs show the
distribution of cells, taking into account their viability (a and b). Comparative analysis was performed to
determine differences between Eca EPCs Non-OB and OB for viability (c) total cell apoptosis (d) early
apoptosis (e) and late apoptosis (f). Means ± SD. *** p-value < 0.001.

454
Fig. 11: The expression profile of genes associated with apoptosis determined in equine endometrial
progenitor cells (Eca EPCs) isolated from non-obese mares (Non-OB) and obese mares (OB). The
transcript levels were determined using RT-qPCR and normalised to the expression of *GAPDH*
(glyceraldehyde-3-phosphate dehydrogenase) and spike control. The tested apoptotic markers were
Bcl-2-associated X protein (*BAX*, B-cell lymphoma 2 (*BCL-2*), caspase 9 (*CASP9*), cellular tumour
antigen p53 (*P53*) and cyclin-dependent kinase inhibitor 1A (*P21*). The relative values of gene
expression were established using RQ_{MAX} algorithm. Means ± SD. *** p-value < 0.001.
462
463   The decreased viability of EPCs Ob was associated with oxidative stress and accumulation of
464 nitric oxygen (NO) and reactive oxygen species (ROS). Dysregulated oxidative stress balance noted in
465 EPCs OB was correlated with decreased levels of miR-20a-5p and miR-133b-3p (**Fig 12**).
466

467
468
Fig.12: The analysis of the oxidative status in equine endometrial progenitor cells (Eca EPCs) isolated
from non-obese mares (Non-OB) and obese mares (OB). Representative dot plots show the distribution
of cells based on intracellular accumulation of nitric oxide (a and b) while column graphs show results
of statistical analyses (d and e). The increased occurrence of NO positive cells (NO+) noted in EPCs
Ob corresponds with the lowered expression of miR-20a (c) and miR-133b (f). Distribution of cells based
on ROS accumulation visible on histograms (g and h) and comparative analysis confirmed oxidative
stress in Eca EPCs OB. Means ± SD. * p-value < 0.05 and ** p-value < 0.01.
476

Expression pattern of FOXP3 and SIRT1 confirmed the disturbance of oxidative status in EPCs
OB. The mRNA level of FOXP3 noted in EPCs was not affected by the physiological status of mares.
However, we observed significantly decreased intracellular accumulation of FOXP3 protein in EPCs OB.
This phenotype may be related to lower regulatory potential and accumulation of oxidative stress
markers of EPC Ob. Lower expression of SIRT1 in EPCs OB supports lower metabolic activity. Elevated
mRNA levels for osteopontin (*OPN)* and accumulated intracellular protein for EPCs OB confirms the
molecular phenotype associated with obesity. Significantly increased expression of OPN was noted for
glycosylated protein, with a molecular weight equal to 66 kDa, and fragments with molecular weight
equal to 30 kDa (**Fig 13**).

486
Fig. 13: Expression of forkhead box P3 (FOXP3), sirtuin 1 (SIRT1) and osteopontin (OPN) determined
in equine endometrial progenitor cells (Eca EPCs) isolated from non-obese mares (Non-OB) and obese
mares (OB).   Protein expression was detected using Western blot technique (a), while RT-qPCR (b-d)
was used to establish the mRNA level for genes of interest. The densitometric measurements of
membranes were performed to compare the expression of proteins between EPCs Non-Ob and Ob (e-
j). Significant differences were calculated for normalised values and shown as means ± SD. * p-value <
0.05 and ** p-value < 0.01.
494
495   We noted an increased constitutive expression of AKT/PI3K in Eca EPCs derived from obese
496 mares. The analysis of mRNA levels determined for *AKT* and *PI3K* indicated the increased accumulation
497 of those transcripts in Eca EPCs OB. However, a statistically significant difference was noted only in
498 *AKT* mRNA expression. In turn, intracellular detection of AKT and PI3K revealed a significant increase
499 of those proteins in Eca EPCs from obese mares (**Fig. 14**).
500

(a)

(b)

(c)

(d)

(e)

(f)

(g)

(h)

(i)

Fig. 14: Expression of Pi3K/AKT determined and compared between EPCs Non-OB and OB. The intracellular accumulation of AKT and PI3K was visualised using immunocytochemistry (ICC, a-b) and determined with Western blot (c). RT-qPCR (d-i) was used to establish mRNA levels for genes of interest. Densitometric measurements of membranes were performed to compare expression of proteins between EPCs Non-Ob and Ob (e-j). Significant differences were calculated for normalised values and shown as means ± SD. * p-value < 0.05 and ** p-value < 0.01.

4. Discussion

The endometrium has gained attention as a source of progenitor cells with high proliferative potential. The endometrial progenitor cells (EPCs) are described as a multipotent population, i.e. able to differentiate into adipogenic, osteogenic, chondrogenic, and even myogenic cell lineages or neurons[32]. The pro-regenerative features of EPCs, with their self-renewal properties, make them a treatment option for endometrial healing. Their potential therapeutic application has been strictly associated with treating endometrial diseases in women, such as Asherman's syndrome characterised by endometrial scarring, fibrosis, and dense adhesions with occlusion of the uterine cavity and endometriosis[32,33]. However, because of the high cellular plasticity and multipotent properties of EPCs, their therapeutic effectiveness in regenerative medicine could be much broader. For instance, endometrial stem cells were proposed as an alternative therapeutic tool for Parkinson's disease [34]. Also, in veterinary regenerative medicine, EPCs-based therapies could be a possible treatment for subfertility caused by endometrial dysfunction, which is a severe problem in the horse breeding industry[35]. Endometrial diseases occur commonly in mares without a specific aetiology.

Obesity and insulin resistance can affect reproduction in mares, impacting the embryo and uterus, as well as the oocyte [2, 30]. We have previously shown that the regenerative potential of multipotent stromal cells isolated from equine adipose tissue depends on the animal's physiological status [8,9,11,36]. Disturbed cellular metabolism affects the pro-regenerative potential of progenitor cells, decreasing their utility in autologous transplantations. For instance, in this study, we have shown that obesity affects the molecular phenotype of equine endometrial progenitor cells (Eca EPCs) and may

reduce their multipotency. We observe, inter alia, that Eca EPCs from obese mares had significantly
decreased mRNA expression for CD105 surface marker (endoglin) compared to Eca EPCs from non-
obese mares. The CD105 is an essential factor associated with the differentiation potential of adult stem
cells originated from mesenchymal tissues, i.e. multipotent stromal/stem cells (MSCs)[37]. The
expression of this marker varies between MSCs of different origins and is related to their heterogeneity
[38]. Indeed, the population of Eca EPCs OB used in this study had lowered capacity for osteogenic and
chondrogenic differentiation[39]. The functional role of CD105 in MSC biology is still deliberated,
especially its effect on chondrogenesis in vitro. Thus, future studies related to multipotency of Eca EPCs
should also include an analysis of CD105 association with their plasticity and potential for tissue-specific
differentiation. Notably, the increased mRNA expression for α-smooth muscle actin (ACTA/α-SMA) was
also a characteristic of Eca EPCs OB. Studies by Talele et al. show that α-SMA is a feature of cells with
lowered clonogenic potential and limited cellular plasticity, confirming results obtained in this study[40].
Essentially, the current study demonstrated that obesity reduces the proliferation and mitochondrial
metabolism of equine endometrial progenitor cells (Eca EPCs). The regenerative potential of progenitor
cells is reflected by high proliferative (self-renewal) capacity and viability. We have demonstrated that
Eca EPCs isolated from mares that are obese have decreased clonogenic and proliferative potential, as
well as migratory activity. Lower mitochondrial metabolism in Eca EPCs was associated with apoptosis
and oxidative stress. Our results correspond with the study of Allesio et al. [41], who showed that obesity
in mice not only affects the proliferation of progenitor cells isolated from subcutaneous and visceral
adipose tissue but also affects cells derived from bone marrow.
In our study, Eca EPCs from obese mares had a reduced percentage of progenitor cells in the S-phase
of the cell cycle, associated with a lower potential for cell division and a prolonged time for population
doubling. The clonogenic potential of EPCs from obese mares was also decreased compared to EPCs
from non-obese mares, which agrees with the finding that obesity negatively impacts the clonogenicity
of endometrial mesenchymal stem cells of women with reproductive failure [42]. In previous studies, we
observed that the proliferative potential of multipotent adipose-derived stromal cells (ASC) could be
affected by increased age and the occurrence of equine metabolic syndrome. An increased percentage
of G1/G0-arrested cells was characteristic for cells isolated from middle-aged horses (5–15), old
(> 15 years old), or had metabolic syndrome [43].
The decreased self-renewal potential of EPCs isolated from obese mares was correlated with
cell-cycle arrest and reduced expression of Ki67, a marker of actively proliferating cells. Obesity
increases Ki67 expression in human progenitor cells from adipose tissue and smaller adipocytes as well
as leads to their cellular death [44]. A hallmark of the inflammation accompanying obesity is a reduction
in Ki67 expression and inhibited proliferation of progenitor cells in the taste buds[45]. Thus, obesity-
induced changes in Ki67 levels can depend on tissue niche and cell type. For instance, functional
endometrium is characterised by the presence of stromal cells showing increased Ki67 expression in
addition to increased levels of the anti-apoptotic protein, BCL-2[46].
We obtained a similar expression profile in our study, as EPCs isolated from obese mares
exhibited a phenotype of early apoptotic cells, with increased mRNA levels for pro-apoptotic *BAX*,
caspase 9, *P53* and *P21* and decreased expression of *BCL-2*. The balance between apoptosis and
proliferation in endometrial cells is essential to regulate regeneration of the endometrium during the
reproductive cycle. The decreased viability of Eca EPCs OB can also be related to the lowered
expression of mRNA for vimentin (*VIM*). Generally, vimentin is recognised as a type III intermediate
filament protein abundant in tissues of mesenchymal origin. This cytoskeleton protein plays a pivotal
role in maintaining the shape and integrity of cells. It is also notable that vimentin is crucial for tissue
homeostasis, protecting cells from protein misfolding stress[47]. Given the above, the Eca EPCs OB
show features of early apoptotic cells with decreased cellular function, including self-renewal potential.
Our study showed, for the first time, that progenitor cells isolated from the endometrium of
obese mares during anestrus have a pro-apoptotic profile and lower mitochondrial potential than cells
from non-obese mares. The molecular phenotype of obese endometrium was also consistent with an
increased expression of osteopontin (OPN). During obesity, high expression of OPN could correlate
with the development of insulin resistance. OPN is depicted as a biomarker for obesity-associated
inflammation and deteriorating adipose tissue metabolism related to enhanced macrophage
infiltration[48]. Knockout of OPN in mice endometrial stromal cells resulted in lower adhesion and
invasion of the blastocyst, as well as decreased expression of matrix metalloproteinase-9 (MMP-9)

associated with impaired invasive competency of the trophoblast[49]. Increased levels of OPN are also
correlated with oxidative stress and decreased mitochondrial membrane potential, which was observed
using the model of mouse cardiac myocytes [50]. Collectively, our results showed that OPN may be
associated with the deterioration of Eca EPCs metabolism induced by obesity. Thus OPN may serve as
a potential target in terms of developing novel therapeutic strategies in endometrial diseases.
Furthermore, analysis of mitochondrial dynamics revealed the increased occurrence of fragmented
globular mitochondria in EPCs derived from the obese endometrium compared to EPCs from non-obese
mares. Obesity-induced morphological and functional changes of mitochondria have been noted in
skeletal muscle and linked with impaired tissue metabolism [51]. Metabolic imbalance related to
decreased mitochondrial activity and oxidative stress is linked with disturbed decidualisation, the
pathogenesis of endometriosis, and endometrial cancer development [52]. The cellular stress noted in
endometrial progenitor cells is related to the pro-inflammatory uterine environment. In the current study,
expression of FOXP3 and SIRT1 was lower in progenitor cells isolated from obese than non-obese
endometrium. A characteristic of regulatory T cells, FOXP3 is responsible for the immunosuppressive
capacity of multipotent stromal cells [53] and may participate in regulating immune responses in
endometriosis. Endometrial FOXP3 in women with endometriosis decreases linearly from early to late
proliferative phase[54]. In turn, SIRT1 suppresses nuclear factor-κB (NF-κB), a transcription factor
regulating the expression of various pro-inflammatory cytokines and chemokines [55,56]. Moreover,
SIRT1 may be an essential molecule for regulating embryo implantation into the endometrium through
stimulation of E-cadherin [57].
Compared to normal-weight mares, embryos from obese mares have higher expression of transcripts
associated with inflammation, endoplasmic reticulum, oxidative and mitochondrial stress [6]. Defects of
mitochondrial metabolism occur in endometriosis with a decrease in electron transport chain complex I-
and II-mediated mitochondrial respiration [58], which correlates with our study. Comparative analysis
showed that EPCs OB compared to EPCs Non-OB had decreased expression of transcripts essential
for mtDNA function and mitochondrial respiratory chain stability.
Defects in mitochondrial metabolism, observed in EPCs OB, could originate from accumulation of
reactive oxygen species (ROS) and nitric oxide, which is also accompanied by decreased mRNA and
protein expression of PTEN-induced kinase 1 (PINK1) and mitofusin 1 (MNF1). PINK1 is an essential
regulator of mitochondrial homeostasis and energy metabolism. Studies performed using the PINK1-
knockout mice model showed that mammalian PINK1 is a key factor protecting against oxidative stress
in the mitochondrion[59]. Decreased levels of PINK1 are noted in the endometrium with minimal and
mild endometriosis [60]. Mitofusins, including mitofusin 1 and 2 (MFN1 and MFN2, respectively), are
crucial mediators of mitochondrial fusion, which compensate for the functional defects of the
mitochondrion. Moreover, MFN1 is particularly important in terms of female fertility. The absence of
MFN1 is associated with oocyte apoptosis and cell-cycle arrest, depletion of the ovarian follicular
reserve, and characteristic of female reproductive ageing [61]. Moreover, increased expression of
PARKIN noted in Eca EPCs OB can be associated with clearance of dysfunctional mitochondria and
activation of mechanisms that protect cells against apoptosis[62].
We also noted that obesity-induced changes in endometrial progenitor cells were associated with
increased constitutive expression of PI3K/AKT and increased β−galactosidase activity. The PI3K/AKT
pathway controls various critical cellular processes, including glucose homeostasis, lipid metabolism,
and protein synthesis; it also affects cell proliferative activity and viability. Increased AKT expression is
related to increased adipogenesis and the obese phenotype [63]. An overactive AKT pathway is a
characteristic of endometrial cancer and endometriosis, thus regulation of PI3K/AKT signalling also
became a consideration in the matter of therapeutic strategies for endometrial disorders [64].
Obesity induces cellular senescence in mesenchymal stromal/stem cells, related to a chronic
inflammatory condition[7]. Consistent with our results, Conley et al. showed that obesity elicits an early
senescence program in progenitor cells, manifested by an increased expression of such markers as p53
and p21. However, in the study, Coloney et al. indicated that senescence-associated beta-galactosidase
activity (SA-β-gal) is not significantly increased in adipose-derived MSCs of obese subjects. In contrast,
we found that EPCs from obese mare endometrium accumulated significantly more SA-β-gal in
comparison to EPCs isolated from non-obese endometrium.
Using two-tailed RT-qPCR, we established equine non-coding RNA levels. We have tested miRNAs
regulating oxidative status and counteracting the senescent cell burden of progenitor cells. The obtained

miRNA profile confirmed that obesity attenuates the molecular phenotype of EPCs, resulting in decreased proliferative activity and oxidative stress. Eca EPCs from obese endometrium were characterised by decreased levels of let-7b and let-7c, as well as miR-20a-5p, -29a-3p, -133b-3p and -181a-5p, but significantly increased levels of let-7a. Loss of let-7b expression is connected with endometriosis pathophysiology and inflammation [65]. In human adipose-derived stem cells, let-7c is recognised as a molecule promoting ectopic bone formation and suppressing adipogenesis via high-mobility group AT-hook 2 [66]. Thus, decreased expression of let-7b and let-7c in EPCs from obese endometrium could be related to the deterioration of self-renewal potential and decreased cellular plasticity. Moreover, increased levels of let-7a may reflect the lowered immunomodulatory potential of Eca EPCs OB, as linked to decreased expression of FOXP3 [67]. Lowered proliferative potential of Eca EPCs OB, was also confirmed by decreased levels of miR-29a and miR-181a-5p. Those molecules are not only essential regulators of the self-renewal potential of MSCs, but they also promote the wound healing capacity of tissues[68]. Furthermore, miR-29a promotes angiogenesis and protects against fibrosis post-injury, regulating biological activity of MMPs and vascular endothelial growth factors (VEGFs) [69]. Down-regulation of miR-20a-5p and miR-133b-3p are observed in mesenchymal stem cells as a result of oxidative stress damage [70,71].

While not the focus of this manuscript, our results marked differences in the endometrium of mares associated with obesity. These changes suggest that the obese mare's uterus could be less able to adapt to cyclic changes and provide an optimal environment during pregnancy. While further studies are needed in this area, body condition is a potential consideration in broodmare management.

In summary, we demonstrated that obesity in the mare alters functional features of endometrial progenitor cells, such as molecular phenotype associated with their potential for specific differentiation, self-renewal activity and mitochondrial metabolism and dynamics. Obesity attenuated cellular homeostasis of endometrial progenitor cells, resulting in increased oxidative stress and hampered pro-regenerative signalling. Early apoptosis and increased senescence characterised progenitor cells when derived from obese endometrium. The impaired cytophysiology of progenitor cells from obese endometrium predict lower regenerative capacity if used as autologous transplants. In our opinion, further studies are required to evaluate the cellular plasticity and immunomodulatory properties of endometrial progenitor cells. Future studies should also focus on developing methods to change the trajectory of obesity-induced metabolic deterioration to improve their functional properties.

Declarations

Authors' declaration of interests

Authors declare no financial or other potential conflict of interest.

Source of funding

The project was financed from an individual research project funded by Wrocław University of

Environmental and Life Sciences grant no. N090/0003/21. Financial support from the National Science

Centre over the course of the realization of the project OPUS 18: Exploring the role and therapeutic

potential of sex hormone binding globulin (SHBG) in the course of insulin resistance, inflammation,

lipotoxicity in adipose stem progenitor cells and adipocytes in equine metabolic syndrome (EMS)

mares (UMO-2019/35/B/NZ7/03651). The studies aimed at the evaluation of equine miRNA profile

also were supported by the following grants: Czech Science Foundation GACR 18-21942S, RVO

86652036 and BIOCEV CZ.1.05/1.1.00/02.0109.

Authorship

A.Ś. and K.Marycz conceived and designed the study; A.Ś. and KMarycz participated in

experiment supervision; A.P. collected and categorised the specimens; AŚ and KMarcinkowska

isolated cells; K.Marcinkowska performed the in vitro cultures and molecular biology experiments;

M.S. performed the immunocytochemical studies and analysis; A.Ś. KMarcinkowska and M.S.

prepared figures; KMarcinkowska, M.S. and A.Ś wrote the material and method section; A.Ś.wrote the

manuscript; KMarycz, L.V. and E.C. participated in critical reading of the manuscript.

All of the authors read and approved the final manuscript.

The tissues used for the isolation of progenitor cells were derived from commercial abattoirs.

The Abattoir owner permits the publication of the abattoir identity indicated in the Materials and

Method section.

The data that support the findings of this study are available from the corresponding author upon

request.

List of legends for Supplementary items:

Figure 1S. The results of body condition score (BCS) examination performed on live mares for proper

classification of animals. Columns with bars represent mean ± SD. * p-value < 0.05, ** p-value < 0.01

Table 1S. Sequences and characteristics of primers used for the mRNA detection with RT-qPCR.

Table 2S. List of primers used for establishing miRNA levels with the Two-tailed RT-qPCR approach.

Table 3S. List of antibodies used for protein detection using Western blot.

Table 4S. List of antibodies used for immunocytochemistry.

References:

1. Rhee, J.S.; Saben, J.L.; Mayer, A.L.; Schulte, M.B.; Asghar, Z.; Stephens, C.; Chi, M.M.-Y.; Moley, K.H. Diet-Induced Obesity Impairs Endometrial Stromal Cell Decidualization: A Potential Role for Impaired Autophagy. *Hum Reprod* **2016**, *31*, 1315–1326, doi:10.1093/humrep/dew048.

2. Das, M.; Sauceda, C.; Webster, N.J.G. Mitochondrial Dysfunction in Obesity and Reproduction. *Endocrinology* **2021**, *162*, doi:10.1210/endocr/bqaa158.

3. Robles, M.; Nouveau, E.; Gautier, C.; Mendoza, L.; Dubois, C.; Dahirel, M.; Lagofun, B.; Aubrière, M.-C.; Lejeune, J.-P.; Caudron, I.; et al. Maternal Obesity Increases Insulin Resistance, Low-Grade Inflammation and Osteochondrosis Lesions in Foals and Yearlings until 18 Months of Age. *PLoS One* **2018**, *13*, doi:10.1371/journal.pone.0190309.

4. Johnson, P.J.; Wiedmeyer, C.E.; Messer, N.T.; Ganjam, V.K. Medical Implications of Obesity in Horses—Lessons for Human Obesity. *J Diabetes Sci Technol* **2009**, *3*, 163–174.

5. Kosolofski, H.R.; Gow, S.P.; Robinson, K.A. Prevalence of Obesity in the Equine Population of Saskatoon and Surrounding Area. *Can Vet J* **2017**, *58*, 967–970.

6. Sessions-Bresnahan, D.R.; Heuberger, A.L.; Carnevale, E.M. Obesity in Mares Promotes Uterine Inflammation and Alters Embryo Lipid Fingerprints and Homeostasis†. *Biology of Reproduction* **2018**, *99*, 761–772, doi:10.1093/biolre/ioy107.

7. Conley, S.M.; Hickson, L.J.; Kellogg, T.A.; McKenzie, T.; Heimbach, J.K.; Taner, T.; Tang, H.; Jordan, K.L.; Saadiq, I.M.; Woollard, J.R.; et al. Human Obesity Induces Dysfunction and Early Senescence in Adipose Tissue-Derived Mesenchymal Stromal/Stem Cells. *Front. Cell Dev. Biol.* **2020**, *8*, doi:10.3389/fcell.2020.00197.

8. Marycz, K.; Kornicka, K.; Marędziak, M.; Golonka, P.; Nicpoń, J. Equine Metabolic Syndrome Impairs Adipose Stem Cells Osteogenic Differentiation by Predominance of Autophagy over Selective Mitophagy. *Journal of Cellular and Molecular Medicine* **2016**, *20*, 2384–2404, doi:https://doi.org/10.1111/jcmm.12932.

9. Marycz, K.; Weiss, C.; Śmieszek, A.; Kornicka, K. Evaluation of Oxidative Stress and Mitophagy during Adipogenic Differentiation of Adipose-Derived Stem Cells Isolated from Equine Metabolic Syndrome (EMS) Horses. *Stem Cells Int* **2018**, *2018*, 5340756, doi:10.1155/2018/5340756.

10. Marycz, K.; Kornicka, K.; Basinska, K.; Czyrek, A. Equine Metabolic Syndrome Affects Viability, Senescence, and Stress Factors of Equine Adipose-Derived Mesenchymal Stromal Stem Cells: New Insight into EqASCs Isolated from EMS Horses in the Context of Their Aging Available online: https://www.hindawi.com/journals/omcl/2016/4710326/ (accessed on 19 February 2021).

11. Smieszek, A.; Kornicka, K.; Szłapka-Kosarzewska, J.; Androvic, P.; Valihrach, L.; Langerova, L.; Rohlova, E.; Kubista, M.; Marycz, K. Metformin Increases Proliferative Activity and Viability of Multipotent Stromal Stem Cells Isolated from Adipose Tissue Derived from Horses with Equine Metabolic Syndrome. *Cells* **2019**, *8*, doi:10.3390/cells8020080.

12. Al Naem, M.; Bourebaba, L.; Kucharczyk, K.; Röcken, M.; Marycz, K. Therapeutic Mesenchymal Stromal Stem Cells: Isolation, Characterization and Role in Equine Regenerative Medicine and Metabolic Disorders. *Stem Cell Rev and Rep* **2020**, *16*, 301–322, doi:10.1007/s12015-019-09932-0.

13. Zahedi, M.; Parham, A.; Dehghani, H.; Kazemi Mehrjerdi, H. Equine Bone Marrow-Derived Mesenchymal Stem Cells: Optimization of Cell Density in Primary Culture. *Stem Cell Investig* **2018**, *5*, doi:10.21037/sci.2018.09.01.

14. Lara, E.; Rivera, N.; Cabezas, J.; Navarrete, F.; Saravia, F.; Rodríguez-Alvarez, L.; Castro, F.O. Endometrial Stem Cells in Farm Animals: Potential Role in Uterine Physiology and Pathology. *Bioengineering (Basel)* **2018**, *5*, doi:10.3390/bioengineering5030075.

15. Rink, B.E.; Amilon, K.R.; Esteves, C.L.; French, H.M.; Watson, E.; Aurich, C.; Donadeu, F.X. Isolation and Characterization of Equine Endometrial Mesenchymal Stromal Cells. *Stem Cell Res Ther* **2017**, *8*, doi:10.1186/s13287-017-0616-0.

16. Rink, B.E.; Beyer, T.; French, H.M.; Watson, E.; Aurich, C.; Donadeu, F.X. The Fate of Autologous
Endometrial Mesenchymal Stromal Cells After Application in the Healthy Equine Uterus. *Stem*
Cells Dev **2018**, *27*, 1046–1052, doi:10.1089/scd.2018.0056.
17. Serrato López, A.G.; Montesinos Montesinos, J.J.; Anzaldúa Arce, S.R.; Serrato López, A.G.;
Montesinos Montesinos, J.J.; Anzaldúa Arce, S.R. The Endometrium as a Source of Mesenchymal
Stem Cells in Domestic Animals and Possible Applications in Veterinary Medicine. *Veterinaria*
México OA **2017**, *4*, 41–58, doi:10.21753/vmoa.4.3.441.
18. Canisso, I.F.; Segabinazzi, L.G.T.M.; Fedorka, C.E. Persistent Breeding-Induced Endometritis in
Mares—A Multifaceted Challenge: From Clinical Aspects to Immunopathogenesis and
Pathobiology. *International Journal of Molecular Sciences* **2020**, *21*, 1432,
doi:10.3390/ijms21041432.
19. Basinska, K.; Marycz, K.; Śieszek, A.; Nicpoń, J. The Production and Distribution of IL-6 and TNF-a
in Subcutaneous Adipose Tissue and Their Correlation with Serum Concentrations in Welsh
Ponies with Equine Metabolic Syndrome. *J Vet Sci* **2015**, *16*, 113–120,
doi:10.4142/jvs.2015.16.1.113.
20. Henneke, D.R.; Potter, G.D.; Kreider, J.L.; Yeates, B.F. Relationship between Condition Score,
Physical Measurements and Body Fat Percentage in Mares. *Equine Vet J* **1983**, *15*, 371–372,
doi:10.1111/j.2042-3306.1983.tb01826.x.
21. Marycz, K.; Pielok, A.; Kornicka-Garbowska, K. Equine Hoof Stem Progenitor Cells (HPC)
CD29 + /Nestin + /K15 + - a Novel Dermal/Epidermal Stem Cell Population With a Potential
Critical Role for Laminitis Treatment. *Stem Cell Rev Rep* **2021**, *17*, 1478–1485,
doi:10.1007/s12015-021-10187-x.
22. Smieszek, A.; Tomaszewski, K.A.; Kornicka, K.; Marycz, K. Metformin Promotes Osteogenic
Differentiation of Adipose-Derived Stromal Cells and Exerts Pro-Osteogenic Effect Stimulating
Bone Regeneration. *J Clin Med* **2018**, *7*, E482, doi:10.3390/jcm7120482.
23. Śmieszek, A.; Stręk, Z.; Kornicka, K.; Grzesiak, J.; Weiss, C.; Marycz, K. Antioxidant and Anti-
Senescence Effect of Metformin on Mouse Olfactory Ensheathing Cells (MOECs) May Be
Associated with Increased Brain-Derived Neurotrophic Factor Levels-An Ex Vivo Study. *Int J Mol*
Sci **2017**, *18*, doi:10.3390/ijms18040872.
24. Smieszek, A.; Marcinkowska, K.; Pielok, A.; Sikora, M.; Valihrach, L.; Marycz, K. The Role of MiR-
21 in Osteoblasts-Osteoclasts Coupling In Vitro. *Cells* **2020**, *9*, doi:10.3390/cells9020479.
25. Heuer, G.G.; Skorupa, A.F.; Prasad Alur, R.K.; Jiang, K.; Wolfe, J.H. Accumulation of Abnormal
Amounts of Glycosaminoglycans in Murine Mucopolysaccharidosis Type VII Neural Progenitor
Cells Does Not Alter the Growth Rate or Efficiency of Differentiation into Neurons. *Mol Cell*
Neurosci **2001**, *17*, 167–178, doi:10.1006/mcne.2000.0917.
26. Doubling Time - Online Computing with 2 Points Available online: https://www.doubling-
time.com/compute.php (accessed on 28 January 2021).
27. Peng, J.-Y.; Lin, C.-C.; Chen, Y.-J.; Kao, L.-S.; Liu, Y.-C.; Chou, C.-C.; Huang, Y.-H.; Chang, F.-R.; Wu,
Y.-C.; Tsai, Y.-S.; et al. Automatic Morphological Subtyping Reveals New Roles of Caspases in
Mitochondrial Dynamics. *PLoS Comput Biol* **2011**, *7*, e1002212,
doi:10.1371/journal.pcbi.1002212.
28. Sikora, M.; Marcinkowska, K.; Marycz, K.; Wiglusz, R.J.; Śmieszek, A. The Potential Selective
Cytotoxicity of Poly (L- Lactic Acid)-Based Scaffolds Functionalized with Nanohydroxyapatite and
Europium (III) Ions toward Osteosarcoma Cells. *Materials (Basel)* **2019**, *12*,
doi:10.3390/ma12223779.
29. Marycz, K.; Smieszek, A.; Targonska, S.; Walsh, S.A.; Szustakiewicz, K.; Wiglusz, R.J. Three
Dimensional (3D) Printed Polylactic Acid with Nano-Hydroxyapatite Doped with Europium(III)
Ions (NHAp/PLLA@Eu3+) Composite for Osteochondral Defect Regeneration and Theranostics.
Mater Sci Eng C Mater Biol Appl **2020**, *110*, 110634, doi:10.1016/j.msec.2020.110634.
30. Targonska, S.; Sikora, M.; Marycz, K.; Smieszek, A.; Wiglusz, R.J. Theranostic Applications of
Nanostructured Silicate-Substituted Hydroxyapatite Codoped with Eu3+ and Bi3+ Ions-A Novel

Strategy for Bone Regeneration. *ACS Biomater Sci Eng* **2020**, *6*, 6148–6160, doi:10.1021/acsbiomaterials.0c00824.

31. Sikora, M.; Śmieszek, A.; Marycz, K. Bone Marrow Stromal Cells (BMSCs CD45- /CD44+ /CD73+ /CD90+) Isolated from Osteoporotic Mice SAM/P6 as a Novel Model for Osteoporosis Investigation. *J Cell Mol Med* **2021**, *25*, 6634–6651, doi:10.1111/jcmm.16667.

32. Figueira, P.G.M.; Abrão, M.S.; Krikun, G.; Taylor, H. STEM CELLS IN ENDOMETRIUM AND THEIR ROLE IN THE PATHOGENESIS OF ENDOMETRIOSIS. *Ann N Y Acad Sci* **2011**, *1221*, 10–17, doi:10.1111/j.1749-6632.2011.05969.x.

33. Queckbörner, S.; Syk Lundberg, E.; Gemzell-Danielsson, K.; Davies, L.C. Endometrial Stromal Cells Exhibit a Distinct Phenotypic and Immunomodulatory Profile. *Stem Cell Research & Therapy* **2020**, *11*, 15, doi:10.1186/s13287-019-1496-2.

34. Wolff, E.F.; Mutlu, L.; Massasa, E.E.; Elsworth, J.D.; Eugene Redmond, D.; Taylor, H.S. Endometrial Stem Cell Transplantation in MPTP- Exposed Primates: An Alternative Cell Source for Treatment of Parkinson's Disease. *J Cell Mol Med* **2015**, *19*, 249–256, doi:10.1111/jcmm.12433.

35. Schöniger, S.; Schoon, H.-A. The Healthy and Diseased Equine Endometrium: A Review of Morphological Features and Molecular Analyses. *Animals (Basel)* **2020**, *10*, doi:10.3390/ani10040625.

36. Kornicka, K.; Houston, J.; Marycz, K. Dysfunction of Mesenchymal Stem Cells Isolated from Metabolic Syndrome and Type 2 Diabetic Patients as Result of Oxidative Stress and Autophagy May Limit Their Potential Therapeutic Use. *Stem Cell Rev Rep* **2018**, *14*, 337–345, doi:10.1007/s12015-018-9809-x.

37. Schmelzer, E.; McKeel, D.T.; Gerlach, J.C. Characterization of Human Mesenchymal Stem Cells from Different Tissues and Their Membrane Encasement for Prospective Transplantation Therapies. *Biomed Res Int* **2019**, *2019*, 6376271, doi:10.1155/2019/6376271.

38. Anderson, P.; Carrillo-Gálvez, A.B.; García-Pérez, A.; Cobo, M.; Martín, F. CD105 (Endoglin)- Negative Murine Mesenchymal Stromal Cells Define a New Multipotent Subpopulation with Distinct Differentiation and Immunomodulatory Capacities. *PLoS One* **2013**, *8*, e76979, doi:10.1371/journal.pone.0076979.

39. Cleary, M.A.; Narcisi, R.; Focke, K.; van der Linden, R.; Brama, P.A.J.; van Osch, G.J.V.M. Expression of CD105 on Expanded Mesenchymal Stem Cells Does Not Predict Their Chondrogenic Potential. *Osteoarthritis and Cartilage* **2016**, *24*, 868–872, doi:10.1016/j.joca.2015.11.018.

40. Talele, N.P.; Fradette, J.; Davies, J.E.; Kapus, A.; Hinz, B. Expression of α-Smooth Muscle Actin Determines the Fate of Mesenchymal Stromal Cells. *Stem Cell Reports* **2015**, *4*, 1016–1030, doi:10.1016/j.stemcr.2015.05.004.

41. Alessio, N.; Acar, M.B.; Demirsoy, I.H.; Squillaro, T.; Siniscalco, D.; Bernardo, G.D.; Peluso, G.; Özcan, S.; Galderisi, U. Obesity Is Associated with Senescence of Mesenchymal Stromal Cells Derived from Bone Marrow, Subcutaneous and Visceral Fat of Young Mice. *Aging (Albany NY)* **2020**, *12*, 12609–12621, doi:10.18632/aging.103606.

42. Murakami, K.; Bhandari, H.; Lucas, E.S.; Takeda, S.; Gargett, C.E.; Quenby, S.; Brosens, J.J.; Tan, B.K. Deficiency in Clonogenic Endometrial Mesenchymal Stem Cells in Obese Women with Reproductive Failure – a Pilot Study. *PLoS One* **2013**, *8*, doi:10.1371/journal.pone.0082582.

43. Age-Dependent Impairment of Adipose-Derived Stem Cells Isolated from Horses | Stem Cell Research & Therapy | Full Text Available online: https://stemcellres.biomedcentral.com/articles/10.1186/s13287-019-1512-6 (accessed on 26 March 2021).

44. Maumus, M.; Sengenès, C.; Decaunes, P.; Zakaroff-Girard, A.; Bourlier, V.; Lafontan, M.; Galitzky, J.; Bouloumié, A. Evidence of in Situ Proliferation of Adult Adipose Tissue-Derived Progenitor Cells: Influence of Fat Mass Microenvironment and Growth. *The Journal of Clinical Endocrinology & Metabolism* **2008**, *93*, 4098–4106, doi:10.1210/jc.2008-0044.

45. Kaufman, A.; Choo, E.; Koh, A.; Dando, R. Inflammation Arising from Obesity Reduces Taste Bud Abundance and Inhibits Renewal. *PLoS Biol* **2018**, *16*, doi:10.1371/journal.pbio.2001959.

46. Mertens, H.J.M.M.; Heineman, M.J.; Evers, J.L.H. The Expression of Apoptosis-Related Proteins Bcl-2 and Ki67 in Endometrium of Ovulatory Menstrual Cycles. *Gynecol Obstet Invest* **2002**, *53*, 224–230, doi:10.1159/000064569.

47. Pattabiraman, S.; Azad, G.K.; Amen, T.; Brielle, S.; Park, J.E.; Sze, S.K.; Meshorer, E.; Kaganovich, D. Vimentin Protects Differentiating Stem Cells from Stress. *Scientific Reports* **2020**, *10*, 19525, doi:10.1038/s41598-020-76076-4.

48. Nomiyama, T.; Perez-Tilve, D.; Ogawa, D.; Gizard, F.; Zhao, Y.; Heywood, E.B.; Jones, K.L.; Kawamori, R.; Cassis, L.A.; Tschöp, M.H.; et al. Osteopontin Mediates Obesity-Induced Adipose Tissue Macrophage Infiltration and Insulin Resistance in Mice. *J Clin Invest* **2007**, *117*, 2877–2888, doi:10.1172/JCI31986.

49. Qi, Q.-R.; Xie, Q.-Z.; Liu, X.-L.; Zhou, Y. Osteopontin Is Expressed in the Mouse Uterus during Early Pregnancy and Promotes Mouse Blastocyst Attachment and Invasion In Vitro. *PLOS ONE* **2014**, *9*, e104955, doi:10.1371/journal.pone.0104955.

50. Dalal, S.; Zha, Q.; Singh, M.; Singh, K. Osteopontin-Stimulated Apoptosis in Cardiac Myocytes Involves Oxidative Stress and Mitochondrial Death Pathway: Role of a pro-Apoptotic Protein BIK. *Mol Cell Biochem* **2016**, *418*, 1–11, doi:10.1007/s11010-016-2725-y.

51. Heo, J.-W.; No, M.-H.; Park, D.-H.; Kang, J.-H.; Seo, D.Y.; Han, J.; Neufer, P.D.; Kwak, H.-B. Effects of Exercise on Obesity-Induced Mitochondrial Dysfunction in Skeletal Muscle. *Korean J Physiol Pharmacol* **2017**, *21*, 567–577, doi:10.4196/kjpp.2017.21.6.567.

52. Yang, X.; Wang, J. The Role of Metabolic Syndrome in Endometrial Cancer: A Review. *Front. Oncol.* **2019**, *9*, doi:10.3389/fonc.2019.00744.

53. Sundin, M.; D'Arcy, P.; Johansson, C.C.; Barrett, A.J.; Lönnies, H.; Sundberg, B.; Nava, S.; Kiessling, R.; Mougiakakos, D.; Le Blanc, K. Multipotent Mesenchymal Stromal Cells Express FoxP3: A Marker for the Immunosuppressive Capacity? *J Immunother* **2011**, *34*, 336–342, doi:10.1097/CJI.0b013e318217007c.

54. Berbic, M.; Hey-Cunningham, A.J.; Ng, C.; Tokushige, N.; Ganewatta, S.; Markham, R.; Russell, P.; Fraser, I.S. The Role of Foxp3+ Regulatory T-Cells in Endometriosis: A Potential Controlling Mechanism for a Complex, Chronic Immunological Condition. *Human Reproduction* **2010**, *25*, 900–907, doi:10.1093/humrep/deq020.

55. Liu, T.; Zhang, L.; Joo, D.; Sun, S.-C. NF-KB Signaling in Inflammation. *Signal Transduct Target Ther* **2017**, *2*, 17023, doi:10.1038/sigtrans.2017.23.

56. Yeung, F.; Hoberg, J.E.; Ramsey, C.S.; Keller, M.D.; Jones, D.R.; Frye, R.A.; Mayo, M.W. Modulation of NF-KappaB-Dependent Transcription and Cell Survival by the SIRT1 Deacetylase. *EMBO J* **2004**, *23*, 2369–2380, doi:10.1038/sj.emboj.7600244.

57. Shirane, A.; Wada-Hiraike, O.; Tanikawa, M.; Seiki, T.; Hiraike, H.; Miyamoto, Y.; Sone, K.; Hirano, M.; Oishi, H.; Oda, K.; et al. Regulation of SIRT1 Determines Initial Step of Endometrial Receptivity by Controlling E-Cadherin Expression. *Biochem Biophys Res Commun* **2012**, *424*, 604–610, doi:10.1016/j.bbrc.2012.06.160.

58. Atkins, H.M.; Bharadwaj, M.S.; O'Brien Cox, A.; Furdui, C.M.; Appt, S.E.; Caudell, D.L. Endometrium and Endometriosis Tissue Mitochondrial Energy Metabolism in a Nonhuman Primate Model. *Reprod Biol Endocrinol* **2019**, *17*, doi:10.1186/s12958-019-0513-8.

59. Gautier, C.A.; Kitada, T.; Shen, J. Loss of PINK1 Causes Mitochondrial Functional Defects and Increased Sensitivity to Oxidative Stress. *PNAS* **2008**, *105*, 11364–11369, doi:10.1073/pnas.0802076105.

60. Holzer, I.; Machado Weber, A.; Marshall, A.; Freis, A.; Jauckus, J.; Strowitzki, T.; Germeyer, A. GRN, NOTCH3, FN1, and PINK1 Expression in Eutopic Endometrium – Potential Biomarkers in the Detection of Endometriosis – a Pilot Study. *J Assist Reprod Genet* **2020**, *37*, 2723–2732, doi:10.1007/s10815-020-01905-4.

61. Zhang, M.; Bener, M.B.; Jiang, Z.; Wang, T.; Esencan, E.; Scott Iii, R.; Horvath, T.; Seli, E. Mitofusin 1 Is Required for Female Fertility and to Maintain Ovarian Follicular Reserve. *Cell Death & Disease* **2019**, *10*, 1–15, doi:10.1038/s41419-019-1799-3.

62. Bonilla-Porras, A.R.; Arevalo-Arbelaez, A.; Alzate-Restrepo, J.F.; Velez-Pardo, C.; Jimenez-Del-Rio,
M. PARKIN Overexpression in Human Mesenchymal Stromal Cells from Wharton's Jelly
Suppresses 6-Hydroxydopamine-Induced Apoptosis: Potential Therapeutic Strategy in
Parkinson's Disease. *Cytotherapy* **2018**, *20*, 45–61, doi:10.1016/j.jcyt.2017.09.011.

63. Chu, C.-Y.; Chen, C.-F.; Rajendran, R.S.; Shen, C.-N.; Chen, T.-H.; Yen, C.-C.; Chuang, C.-K.; Lin, D.-
S.; Hsiao, C.-D. Overexpression of Akt1 Enhances Adipogenesis and Leads to Lipoma Formation
in Zebrafish. *PLOS ONE* **2012**, *7*, e36474, doi:10.1371/journal.pone.0036474.

64. Lee, I.I.; Kim, J.J. Influence of AKT on Progesterone Action in Endometrial Diseases. *Biol Reprod*
2014, *91*, doi:10.1095/biolreprod.114.119255.

65. Sahin, C.; Mamillapalli, R.; Yi, K.W.; Taylor, H.S. MicroRNA Let-7b: A Novel Treatment for
Endometriosis. *J Cell Mol Med* **2018**, *22*, 5346–5353, doi:10.1111/jcmm.13807.

66. Liu, G.-X.; Ma, S.; Li, Y.; Yu, Y.; Zhou, Y.-X.; Lu, Y.-D.; Jin, L.; Wang, Z.-L.; Yu, J.-H. Hsa-Let-7c
Controls the Committed Differentiation of IGF-1-Treated Mesenchymal Stem Cells Derived from
Dental Pulps by Targeting IGF-1R via the MAPK Pathways. *Experimental & Molecular Medicine*
2018, *50*, 1–14, doi:10.1038/s12276-018-0048-7.

67. Yu, Y.; Liao, L.; Shao, B.; Su, X.; Shuai, Y.; Wang, H.; Shang, F.; Zhou, Z.; Yang, D.; Jin, Y. Knockdown
of MicroRNA Let-7a Improves the Functionality of Bone Marrow-Derived Mesenchymal Stem
Cells in Immunotherapy. *Mol Ther* **2017**, *25*, 480–493, doi:10.1016/j.ymthe.2016.11.015.

68. Guo, L.; Zhao, R.C.H.; Wu, Y. The Role of MicroRNAs in Self-Renewal and Differentiation of
Mesenchymal Stem Cells. *Exp Hematol* **2011**, *39*, 608–616, doi:10.1016/j.exphem.2011.01.011.

69. Su, W.-H.; Wang, C.-J.; Hung, Y.-Y.; Lu, C.-W.; Ou, C.-Y.; Tseng, S.-H.; Tsai, C.-C.; Kao, Y.-T.; Chuang,
P.-C. MicroRNA-29a Exhibited Pro-Angiogenic and Anti-Fibrotic Features to Intensify Human
Umbilical Cord Mesenchymal Stem Cells—Renovated Perfusion Recovery and Preventing against
Fibrosis from Skeletal Muscle Ischemic Injury. *Int J Mol Sci* **2019**, *20*, doi:10.3390/ijms20235859.

70. Tai, L.; Huang, C.-J.; Choo, K.B.; Cheong, S.K.; Kamarul, T. Oxidative Stress Down-Regulates MiR-
20b-5p, MiR-106a-5p and E2F1 Expression to Suppress the G1/S Transition of the Cell Cycle in
Multipotent Stromal Cells. *Int J Med Sci* **2020**, *17*, 457–470, doi:10.7150/ijms.38832.

71. Yildirim, S.S.; Akman, D.; Catalucci, D.; Turan, B. Relationship Between Downregulation of
MiRNAs and Increase of Oxidative Stress in the Development of Diabetic Cardiac Dysfunction:
Junctin as a Target Protein of MiR-1. *Cell Biochem Biophys* **2013**, *67*, 1397–1408,
doi:10.1007/s12013-013-9672-y.

946