Title: Rapid genomic evolution in *Brassica rapa* with bumblebee selection in experimental evolution

Authors: Léa Frachon[1, *], Florian P. Schiestl[1]

[1]Department of Systematic and Evolutionary Botany, University of Zürich, Switzerland.

Corresponding author:

Léa Frachon

lea.frachon@systbot.uzh.ch

Key words: fast cycling *Brassica rapa*, experimental evolution, genomic variance, rapid genomic evolution, bumblebees' selection

Abstract

Insect pollinators shape rapid phenotypic evolution of traits related to floral attractiveness and plant reproductive success. However, the underlying genomic changes remain largely unknown despite their importance in predicting adaptive responses to natural or to artificial selection. Here, based on a previous nine-generation experimental evolution study with fast cycling *Brassica rapa* plants adapting to bumblebees, we documented genomic evolution associated with the adaptive process. We performed a genomic scan of the allele frequency changes along the genome and estimated the genomic variance changes. We detected signature of selection associated with rapid changes in allelic frequencies on multiple loci, and an increase of overall genomic variance. The genomic evolution observed in nine generations was different between replicates, suggesting non-parallel evolution. Our study highlights the involvement of multiple loci in short-term pollinator adaptation, and the maintenance of genomic variance during strong natural selection by biotic factors.

Introduction

Pollinator insects are important selective agents for wild- and crop plant species due to their essential role in the reproduction of most flowering plants (Van der Niet et al. 2014). While a decline of insect pollinators has been detected in different geographical regions and insect families (Biesmeijer et al. 2006, Hallmann et al. 2017, Wagner et al. 2020), the understanding of the adaptive potential of plants to such changes remains in its infancy. Plant adaptation to pollinators typically involves traits associated with flower attractiveness such as (1) flower morphology (Anderson et al. 2016, Wessinger et al. 2016, Minaar et al. 2019), flower colour (Bradshaw et al. 2003, Sobral et al. 2015), flower scent (Schiestl 2005, Xu et al. 2012, Gervasi and Schiestl 2017), and (2) traits associated with mating system like herkogamy (Moeller and Geber 2005, Luo and Widmer 2013) or selfing (Bodbyl Roles and Kelly 2011, Ramos and Schiestl 2019). While most studies assessed the result of long-term evolutionary adaptation to pollinators, tracking the adaptive processes across generations remains rarely done. Both the resurrection approach in natural populations, growing seeds from different generations together, or experimental evolution studies, applying the same selective pressure for multiple generations, can bridge this gap. For instance, using a resurrection approach, Thomann et al. (2015) observed phenological and reproductive trait changes over 18 years in *Adonis annua* plants in response to the loss of wild bees. While this approach benefits from ecological realism in natural populations, it makes it difficult to differentiate the effect of the factor of interest from other factors such as climate, also shaping plant evolution. Gervasi and Schiestl (2017) performed experimental evolution with fast-cycling *Brassica rapa* plants evolving with different pollinators and under controlled conditions, to identify the evolutionary response to pollinator-mediated selection. They showed, within nine generations of experimental evolution, rapid plant adaptation to bumblebee pollination in phenotypic traits, such as floral volatiles, UV reflection and plant height. However, while the evolved traits have also been shown to carry substantial heritability (Zu et al. 2016, Zu and Schiestl 2017), the genomic changes underlying these rapid plant phenotypic changes are still unknown.

In the current context of pollinator decline and the associated changes in pollinator

communities, analyzing the genetic architecture of adaptation to pollinators and its association to

standing genetic variation is essential to understand the adaptative potential of plants in changing

environments (Hansen et al. 2006, Bergelson et al. 2010, Bay et al. 2017). Molecular genetic studies

have uncovered the molecular and genetic bases of several traits involved in pollination and pollinator

attractiveness such as selfing (Sicard and Lenhard 2011), pollination syndromes (Bradshaw et al. 1995,

Bradshaw and Schemske 2003, Streisfeld and Rausher 2009, Yuan et al. 2013, Yarahmadov et al. 2020),

nectar (Lin et al. 2015, Solhaug et al. 2019) and volatiles (Dudareva et al. 2006, Klahre et al. 2011, Cai

et al. 2016, Fattorini and Glover 2020, Lo et al. 2020). However, insects use a combination of signals

(shape, colour, scent) and rewards for identifying suitable flowers leading to plant adaptation based

on multiple traits (Schiestl and Johnson 2013). For instance, honest signals (signals associated with

reward) and pollination syndromes (convergent evolution of specific signal combinations selected by

pollinators) are good examples of evolution of multiple traits. In a context of rapid changes, genetic

correlation among traits may allow the synchronous response of different phenotypic traits to varying

patterns of selection (Hermann et al. 2013, Smith 2015, Zu and Schiestl 2017). However, we are still in

the beginning of understanding the genetic basis involved in the rapid evolutionary response of plants

to pollinator changes. Identification of genomic regions involved in plant adaptation to pollinators is

essential to predict the adaptive potential of plants to pollinator changes and enable breeding of more

attractive crop plants. In addition, an important aspect in conservation is to understand the

mechanisms maintaining genetic diversity within populations. In fact, standing genetic variation is an

important resource for rapid response to environmental changes (Barrett and Schluter 2008, Chevin

et al. 2010), and strong directional selection is considered to lead to the loss of genetic diversity

resulting in a loss of adaptive potential in populations (Barton and Keightley 2002).

Here, based on previous experimental evolution performed by Gervasi and Schiestl (2017) with

outcrossing fast-cycling *Brassica rapa* plants, we tracked the genomic changes involved in the

adaptative response of plants to bumblebee selection compared to hand pollinated control plants

(**Figure 1**), and documented the changes in genetic diversity observed in the context of strong selection

by estimating the genomic variance before and after experimental evolution.

Methods

Plant material and experimental design. *Brassica rapa* (Brassicacea) is an outcrossing plant with

genetic self-incompatibility, pollinated by diverse insects such as bumblebees, flies or butterflies (El-

Esawi 2017). Our study used rapid-cycling *Brassica rapa* plants (Wisconsin Fast Plants™, purchased

from Carolina Biological Supply Company, Burlington, USA), selected for its short life cycle of

approximately two months from seed to seed. We used seeds produced by the study of Gervasi and

Schiestl (2017), performing experimental evolution with bumblebees and control hand pollination.

Briefly, the pollination was carried out over nine generations either by bumblebees within cages or by

random hand cross-pollination. A subset of seeds produced by individual plants, proportional to the

total seed set of each individual within the replicates was used to generate each subsequent

generation. For this study, we used one seed per individual from 64 plants (half of replicate A, and half

of replicate B) of (1) the starting generation (generation 1; here called A1 and B1); (2) the ninth

generation selected by bumblebees (bumblebee treatment; here called B9A and B9B); (3) the ninth

generation of control hand pollination plants (control treatment; here called C9A and C9B; **Figure 1**).

Finally, we performed crossings between replicates A and B within each treatment, (generation 10)

yielding 32 individuals from the bumblebee treatment (inter-replicate crossing in bumblebee

treatment; here called B10) and 32 individuals from the control treatment (inter-replicate crossing in

control treatment; here called C10). These manual crossing are commonly used for reducing the effect

of potential inbreed depression on trait changes. Pollen donors and receivers were randomly assigned

in these crossing. Each combination of generation*treatment*replicate is called a population (*e.g.*

ninth generation, treatment bumblebees, replicate A called B9A is a population). A total of 256 seeds

from these 8 populations (first, ninth and tenth generation) were sown out in a phytotron (first

generation in 2017 and ninth generation as well as the inter-replicate crossing in 2019) and the leaf

tissue of each plant was collected for DNA extraction and whole genomic sequencing.

DNA extraction and genomic characterization. Because leaf tissue was collected in 2017 for the first

generation and 2019 for the last generations, we adapted the collection storage (drying vs freezing).

Leaf material from the first generation was dried in vacuum at 40 °C for 20 hours, and leaf material

from the ninth and tenth generation was stored in -80°C. A high molecular weight DNA extraction

(average DNA concentration of 48 ng/µL, LGC extraction protocol) and library preparation for

genotyping-by-sequencing (restriction enzyme MsII, insert size mean range ~215bp) was performed

by the LGC Genomics group Berlin. Samples were sequenced with Illumina NextSeq 500 V2 sequencer

using 150 paired-end reads; the alignment of our samples was performed with BWA version 0.7.12

against the reference genome sequence of *Brassica rapa* FPsc v1.3, Phytozome release 12

(https://phytozome.jgi.doe.gov/pz/portal.html) by the LGC Genomics group Berlin. The variant

discovery and the genotyping were realized using Freebayes v1.0.2-16 with the following parameters

by the LGC Genomic Group Berlin: --min-base-quality 10 –min-supporting-allele-qsum 10 –read-

mismatch-limit 3 –min-coverage 5 –no-indels –min-alternate-count 4 –exclude-unobserved-genotypes

–genotype-qualities –ploidy 2 or 3 –no-mnps –no-complex –mismatch-base-quality-threshold 10. We

then performed a quality trimming on the 10 chromosomes *i.e.* by discarded unassigned scaffolds from

the chromosomes using vcftools (Danecek et al. 2011), by removing SNPs with missing data in more

than 5% of the individuals (function –max-missing 0.95, *i.e.* genotype calls had to be present for at

least 243 samples out of 256 for a SNP to be included in the downstream analysis), and retaining only

bi-allelic SNPs with a minimum average Phred quality score of 15 (function –minGQ 15, **Figure S1de**).

We removed SNPs with a maximum average read depth of 100 (function –max-meanDP 100, **Figure**

S1bc), and discarded SNPs with a minor allele frequency (MAF) lower than 0.1 (function –maf 0.1,

Figure S1a). The final dataset contained 4'713 SNPs in ~ 283Mb genome size (https://phytozome.jgi.doe.gov/pz/portal.html).

Allele frequency changes and genetic drift simulation. The allele frequencies of the reference allele for the 4'713 SNPs were estimated within each 8 populations using vcftools (function –freq, Danecek et al. 2011). To control for potential genetic drift during the nine generations of evolution, we simulated random final allele frequencies 10'000-fold for different ranges of initial allele frequencies (from 0 to 1 by an interval window of 0.01). The simulations were performed using the R environment package "learnPopGen" (function "drift.selection", Revell 2019) over eight transitions between generations (*i.e.* from the first generation to the ninth generation) considering 32 individuals within each population for an effective size (N_e) of 16 (*i.e.* individuals contributing to the next generation, see details of experimental evolution in Gervasi and Schiestl 2017), and considering an equal fitness for each individual. For each SNP along the genome, we assigned their initial allele frequency value ($AF_{initial}$) to the range estimated by the final allele frequency simulations (10'000 values). Finally, the observed final allele frequency (AF_{final}) was compared to 10'000 simulated allele frequency values ($AF_{simulated}$) to estimate a p-value for each SNP using the following equations:

(1) For a decrease of reference allelic frequency *i.e.* ($AF_{initial} - AF_{final}$) > 0, *pvalue* = (number of simulations with $AF_{simulated} \geq AF_{final}$)/10'000

(2) For an increase of reference allelic frequency *i.e.* ($AF_{initial} - AF_{final}$) < 0, *pvalue* = (number of simulations with $AF_{simulated} \leq AF_{final}$)/10'000

(3) For ($AF_{initial} - AF_{final}$) = 0, *pvalue* = 1

With $AF_{simulated}$ = simulated final allele frequency, $AF_{initial}$ = initial allele frequency from the reference allele (first generation), and AF_{final} = observed final allele frequency for the reference allele (ninth generation). The pvalues were controlled for the False Discovery Rate (fdr) of 5% using Benjamini-Hochberg method implemented in R environment (Benjamini and Hochberg 1995).

155   Finally, we estimated the allele frequency changes (Δh) from the reference allele according to

156   the equation (1) for both bumblebee and control treatments:

157                                $$\Delta h = \mathrm{AF}_{\mathrm{final}} - \mathrm{AF}_{\mathrm{initial}} \qquad (1)$$

Where Δh is the allelic frequency change between the first and the ninth generation, $\mathrm{AF}_{\mathrm{initial}}$ is the initial

observed allele frequency, and $\mathrm{AF}_{\mathrm{final}}$ is the observed final allele frequency at the ninth generation.

Linkage disequilibrium (LD) and haplotype block structure evolution. During selective process, an

increase of the linkage disequilibrium *i.e.* the non-random association of alleles between loci, is

expected in genomic regions strongly under selection. First, we calculated pairwise linkage

disequilibrium (LD) among all set of SNPs measured by the correlation coefficient between SNP pair r^2

within each chromosomes using VCFtools (function –geno-r2) for 256 samples within each population.

The associated median LD was then estimated and plotted. Finally, to understand whether changes in

median LD are due to random allelic associations along the genome, or aggregated in genomic regions

under selection, we calculated the haplotype blocks in each population using plink1.9 with the

following parameters: --blocks no-pheno-req –maf 0.07 –blocks-max-kb 200. This method estimates

the length of theses blocks with "strong LD" considering the allelic association D' metrics between 0.7

and 0.98 according to (Gabriel et al. 2002).

Candidate genes. We identified 171 candidate genes associated with 187 SNPs with significant allele

frequency changes (fdr < 0.05) during bumblebee selection. Because the median linkage disequilibrium

in bumblebee treatment is ~4kb, we retrieved the annotated genes around 4kb (2 kb upstream and 2

kb downstream) for 187 SNPs and extracted the gene description using phytozome.jgi.doe.gov.

Genome-wide variance changes. The genomic variance among individuals within each population was

estimated using principal component analysis (PCA) on scaled and centered genotype data (pcadapt

package in R environment, function pcadapt, Privé et al. 2020). In order to unravel the changes in

genomic variance over nine generations, we performed the PCA (1) on the total number of SNPs (4713

SNPs), and (2) on 187 SNPs showing significant allele frequency changes under bumblebee selection in

on or another replicate (fdr < 0.05). To compare genomic variance among populations, we estimated

the area of the polygon formed by scatter plots of axis 1 and 2 of each population for the full dataset

(4713 SNPs) and the subset of 187 SNPs using R package tidyverse and splancs (Wickhman et al. 2019,

Rowlingson and Diggle 2022). The relative area is the ratio between the polygon area within each

population and the polygon area circumventing all the points of the PCA.

Results

Genomic changes during experimental evolution. In our study, we observed allele frequency changes

(Δh) over nine generations in both bumblebee and control treatments (**Figure 2**). However, the

strongest ($\Delta h > 0.5$) and most significant (frd < 0.05) changes were observed in the bumblebee

treatment for the replicate A on 32 SNPs (4 SNPs for the replicate B in the bumblebee treatment, 0

SNP and 3 SNPs for replicate A and B respectively in the control treatment). Overall, the significant

changes observed in replicate A for the bumblebee treatment (fdr < 0.05), regardless of its strength,

involved 42-fold more SNPS than in the control treatment (4 SNPs with significant changes in the

control treatment, and 171 SNPs in the bumblebee treatment, **Figure 2**). In replicate B, we did not

observe difference in allelic frequency changes between the two treatments (21 SNPs with significant

changes in the control, and 19 in the bumblebee treatment considering fdr < 0.05, **Figure 2**).

As expected, the evolutionary process is associated with an increase of linkage disequilibrium

in the bumblebee treatment, but also in the control treatment. The decay of the median linkage

disequilibrium was slower in both treatments in the ninth generation than in the first one (**Figure S2A**).

Within the bumblebee treatment, we observed a slower decay of LD in replicate A than in B as

expected with the observed stronger allele frequency changes (**Figure S2A**). The decays of LD in inter-

replicate crossing were slower in both treatments compared to the ninth generation (**Figure S2A**).

Moreover, we observed a decrease of the number of haplotype blocks, *i.e.*, genomic regions with

strongly linked alleles, over nine generations (from 786 and 780 haplotype blocks for replicate A and B

respectively in the first generation, to 563 and 648 haplotype blocks for replicate A and B respectively

in the ninth generation of the control treatment, and 478 and 600 for replicate A and B respectively in

the ninth generation of the bumblebee treatment) associated with an increase of their length (**Figure**

S2BC, **Table S1**). We observed 743 and 696 haplotype blocks in control and bumblebee treatments

respectively in the inter-replicate crossing (**Figure S2BC, Table S1**).

Overall, the SNPs involved in the evolutive responses are independent between replicates, and

between treatments (**Figure 3**). For instance, among the 171 SNPs with significant allele frequency

changes in replicate A in bumblebee treatment, 164 SNPs are unique of this population, whereas 5

SNPs are shared with the control treatment in replicate B, and 3 SNPs are shared with the bumblebee

treatment in replicate B (**Figure 3**).

Identity of candidate genes underlying genomic evolution to bumblebees. After retrieving the

annotated genes around 4kb (2kb upstream, 2kb downstream) for 187 SNPs with significant allele

frequency changes in the bumblebee treatment (fdr < 0.05), we obtained a list of 171 candidate genes

(**Dataset1**). Briefly, we found genes with biological functions that may be involved in the emission of

volatiles (ABC transporter G), in the regulation of flowering time (B3), or in shoot architecture (UCH).

In addition, some genes are involved in the response to different biotic and abiotic stresses

(METACASPASE-2, PUB, Pum, PAT8, BON). Finally, some candidate genes do not have a known

biological function.

Genomic variance changes. To assess the variation of genomic variance across generation, we

performed a genomic principal component analysis (PCA) on the full set of SNPs (4'713 SNPs). We

observed a structuration of our samples determined by populations *i.e*. generations, treatments, and

replicates (**Figure 4A**). Along the two first principal components (PCs) axis, explaining 24.5% of the total

genomic variance, all individuals from the two replicates of generation one were grouped together,

and individuals from the ninth generations were clearly separated from the first generation, but they

were also separated between both treatments and replicates (**Figure 4A**). In genomic space, individuals

from the inter-replicate crossing fell between those from the scatterplot formed by replicates A and B

of generation nine for both treatments (**Figure 4A**). We estimated the genomic variance of our

populations by calculating the area of polygons in the genomic space formed by the first two PCs. We

observed a 2-fold and 4-fold increase in genomic variance (polygon area) between the first and the

ninth generation in replicates A and B respectively in bumblebee treatment, and a 0.5-fold and 2-fold

increase in replicates A and B respectively in control treatment (**Figure 4C**). In the inter-replicate

crossing for both treatments, we observed a 0.8-fold and 7.7-fold (3.9-fold considering only samples

without outliers) increase of the polygon area compared to replicate A in first generation, in control

and bumblebee treatments respectively (**Figure 4C**).

In the reduced dataset including significant allele frequency changes in bumblebee treatment

(both replicate A and B, *i.e.* 187 SNPs), we observed a clear distinction between individuals from the

bumblebee treatment and other populations (**Figure 4B**). Indeed, individuals from the first generation

and the control treatment are clustered closely together, and the individuals from bumblebee

treatment were clearly separated in the genomic space (**Figure 4B**). For the first generation and control

treatment, we observed a higher genomic variance (polygon relative area) for PCA performed on the

sub-dataset than the one performed with the full dataset (**Figure 4C**). In the bumblebee treatment, we

observed different pattern regarding the replicate; a small increase of the genomic variance (polygon

relative area) in replicate A, and a decrease in replicate B and inter-crossing (**Figure 3C**). These genomic

regions drove the evolutionary process observed in bumblebee selection, while maintaining genetic

diversity in the treatment under selection.

Discussion

Understanding pollinator-mediated evolutionary genomics of flowering plants and its impact

on standing genetic variation within the genome remains an important challenge in conservation.

Here, we screened for the genomic consequences of biotic selection in an experimental evolution

experiment by sequencing genome-wide SNP markers in *Brassica rapa* plant individuals before and

after nine generations of selection by bumblebees, and under random hand-pollination. As shown

previously at the level of the phenotype, this primarily outcrossing plant rapidly adapt to specific

pollinators (Gervasi and Schiestl 2017). We documented the signature of directional selection driven

by bumblebee pollinators with significant allele frequency changes at several loci. Interestingly, we

documented a maintenance of standing genetic variation across the genome, a finding challenging the

classical selective sweep model.

In agreement with the previously demonstrated phenotypic evolution in the fast-cycling *Brassica*

rapa experimental system (Gervasi and Schiestl 2017), we have shown genomic evolution across nine

generations. The here documented changes in allele frequencies, the increase of linkage disequilibrium

and the number of haplotype blocks, underlined the importance of pollinators in shaping plant rapid

genomic evolution. Some significant candidate genes identified in our study have biological functions

in line with the phenotypic evolution of traits observed by Gervasi and Schiestl (2017). For instance,

while this previous study highlighted increase in scent emission and plant height driven by bumblebee

pollination, we identified candidate genes who, or at least their family, are known to be involved in

emission of volatile organic compounds, like ABC transporter G family (ABCG35, ABCG38, ABCG1,

ABCB19, Adebesin et al. 2017, Hao et al. 2021), in shoot architecture like Ubiquitin C-terminal

hydrolase (UCH, Yang et al. 2007), or in growth and development like AUXIN-RESPONSIVE PROTEIN-

RELATED, or TREHALOSE-PHOSPHATE PHOSPHATASE E-RELATED (Figueroa and Lunn 2016). The U-box

protein is also an interesting candidate gene due to its involvement in different developmental and

physiological processes such as self-incompatibility, defence, or abiotic stress response (Wang et al.

2015). Moreover, we found significant allele frequency changes associated with flowering time like a

B3 transcription regulator (Swaminathan et al. 2008) or UCH (Hayama et al 2021). While Gervasi &

Schiestl (2017) did not observe any difference in flowering time between bumblebee and control

treatment in the last generation, flowering time is an important phenological trait involved in plant-

pollinator matching (Forrest 2005, Hegland et al. 2009). It is therefore likely that pollinators induce

flower phenological shift. Overall, our study highlighted the potential involvement of multiple loci in

rapid adaption to bumblebees, which agrees with studies highlighting such a genetic architecture

underlying floral evolution (Fishman et al. 2002, Yuan et al. 2013, Fishman et al. 2015, Smith 2015,

Campitelli et al. 2017). However, further analysis using higher quality of sequencing and the full

characterization of associated phenotypic and phenological traits (allowing a genome-wide association

approach) and functional validation of the genes are required to draw more solid conclusions about

the genetic basis of evolutionary changes induced by bumblebee selection. Moreover, among loci

under-selection, the lack of annotation and the unknown functions of some genes highlighted the need

to deepen our knowledge on genes involved in plant-pollinator interactions.

We found that the extent of genomic changes observed during evolutionary processes was

different between replicates suggesting non-parallel evolution (Bolnick et al. 2018). This result is in line

with the observed phenotypic evolution in Gervasi and Schiestl (2017) with significant differences

among replicates in the bumblebee treatment for different phenotypic traits. Such evolution was

observed in natural population of *Senecio lautus* with similar phenotypic variation regulated by

variation in genomic space (James et al. 2021). However, we observed significant changes in allele

frequencies only in one replicate, indicating that it is difficult to clearly distinguish potential non-

parallel evolution from a "two-speed" parallel evolution (*i.e.* the second replicate will follow the same

evolutionary path in the upcoming generations). Longer experimental evolution, with more replicates

and individuals are needed to understand the evolutionary pattern mediated by biotic factors such as

bumblebees.

Finally, we observed an increase of the overall genomic variance despite the selection imposed by

bumblebees during experimental evolution (2017), and the small effective size of populations. This

increase in overall genomic variance was observed among individuals in both bumblebee-pollinated

replicates (B9A and B9B), as well as in the inter-replicate crossing (B10). This pattern might be

explained by the polygenic model with a weaker selection acting on multiple standing variants (soft

sweep) and by multiple loci underlying individual phenotypic trait evolutionary changes. An increased

number of studies demonstrate the importance of polygenic adaptation (Pritchard and di Rienzo 2010,

Csilléry et al. 2018, Barghi et al. 2020) related to the infinitesimal model (reviewed in Barton et al.

2017), where local adaptation is driven by small allele frequency changes in multiple loci. This outcome

is supported by recent work observing the maintenance of standing genetic variation during long-term

artificial selection on chicken weight, mainly explained by selection acting on highly polygenic

architecture (Sheng et al. 2015). However, deepened analyses are needed using newly developed

models to validate such hypothesis (Berg et al. 2014, Boyle et al. 2017, Wieters et al. 2021). Multiple

genes underlying phenotypic variation are widely emphasized in plants with the advances of GWAs

(Barghi et al. 2020), however the interplay of evolutionary forces on these genes and their involvement

in the potential maintenance of standing genetic variation is still poorly understood and deserve

further studies.

Conclusion

We revealed important genomic changes on multiple loci during bumblebee selection during only

nine generations. Our study is a first step into the understanding of the complex genomic mechanisms

involved in rapid evolutionary adaptation to biotic factors, and we advocate further analyses to

understand (1) the genetic architecture underlying phenotypic and phenological variation, (2)

pleiotropic effects of quantitative-trait locus in rapid adaptation and (3) the mechanisms behind a

maintenance of genetic variance. We also underline the importance of better characterizing the gene

functions involved in plant-pollinator interactions. Overall, pollinators constitute complex patterns of

selection which deserve more attention for predicting the adaptive responses of wild and crop plant

species to their decline.

Acknowledgements

We thank Jörg Vogt for the first-generation DNA extraction, and the LGC Genomic group Berlin for the

DNA extraction of other samples, the sequencing and bioinformatic treatment. We thank Cyril Zipfel

for discussion regarding candidate genes. We thank Loren Rieseberg and Marius Rösti for their useful

comments on a previous version of the draft. Finally, we thank the University of Zürich for their

financial support.

Author Contributions

L.F. and F.S. designed the research. L.F. performed the DNA extraction, statistical analyses and wrote

the manuscript. L.F. and F.S. reviewed and edit the manuscript.

References
Adebesin, F. *et al.* Emission of volatile organic compounds from petunia flowers is facilitated by an ABC
transporter. *Science*. **356**, 1386-1388 (2017).
Anderson, B., Pauw, A., Cole, W. W. & Barrett S. C. H. Pollination, mating and reproductive fitness in a
plant population. *Journal of Evolutionary Biology*. **29**, 1631–1642 (2016).
Barghi, N., Hermisson, J. & Schlötterer, C. Polygenic adaptation: a unifying. *Nature Reviews Genetics*.
21, 769-781 (2020).
Barrett, R. D. H. & Schluter, D. Adaptation from standing genetic. *TRENDS in Ecology and Evolution*.
23(1), 38-44 (2008).
Barton, N. H. & Keightley, P.D. Understanding quantitative genetic variation. *Nature Reviews Genetics*.
3, 11-21 (2002).
Barton, N. H., Etheridge, A.M. & Véber, A. The infinitesimal model: Definition, derivation, and
implications. *Theoretical Population Biology*. 118, 50-73 (2017).
Bay, R. A. *et al*. Predicting responses to contemporary environmental change using evolutionary
response architectures. *The American naturalist*. **189**(5), 463-473 (2017).
Benjamini, Y. & Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to
multiple testing. *Journal of Royal Statistical Society: Series B*. **57**(1), 289-300 (1995).

Berg, J. J & Coop, G. A population genetic signal of polygenic adaptation. *PloS Genet.* **10**(8), e1004412
(2014).
Bergelson, J. & Roux, F. Towards identifying genes underlying ecologically relevant traits in Arabidopsis
thaliana. *Nature Reviews Genetics.* **11**, 867-879 (2010).
Biesmeijer, J. C. *et al*. Parallel declines in pollinators and insect-pollinated plants in Britain and the
Netherlands. *Science.* **313**, 351-354 (2006).
Bodbyl Roles, S. A. & Kelly, J. K. Rapid evolution caused by pollinator loss in *Mimulus guttatus.*
Evolution. **65**(9), 2541–2552 (2011).
Bolnick D. I., Barrett R. D. H., Oke K. B., Rennison D. J., Stuart Y. E. (Non)Parallel Evolution. *Annual*
Review of Ecology, Evolution, and Systematics. 19, 303-330 (2018).
Boyle, E. A., Li, Y. I. & Pritchard J. K. An expanded view of complex traits: from polygenic to
omnigenic. *Cell.* 169, 1177-1186 (2017).
Bradshaw, H. D. & Schemske, D. W. Allele substitution at a flower colour locus produces a pollinator
shift in monkeyflowers. *Nature.* **426**, 176-178 (2003).
Bradshaw, H. D., Wilbert, S. M., Otto, K. G. & Schemsket, D. W. Genetic mapping of floral traits
associated with reproductive isolation in monkeyflowers (*Mimulus*). *Nature.* **376**, 762-765
(1995).
Cai, J., Zu, P. & Schiestl F. P. The molecular bases of floral scent evolution under artificial selection:
insights from a transcriptome analysis in Brassica rapa. *Scientific Reports.* **6**, 36966 (2016).
Campitelli, B. E., Kenney, A. M., Hopkins, R., Soule, J., Lovell, J. T. & Juenger, T.E. Genetic mapping
reveals an anthocyanin biosynthesis pathway gene potentially influencing evolutionary
divergence between two subspecies of scarlet gilia (*Ipomopsis aggregata*). *Molecular Biology*
and Evolution. **35**(4), 807–822 (2017).
Chevin, L. M., Lande, R. & Mace, G. M. Adaptation, plasticity, and extinction in a changing environment:
towards a predictive theory. *PLOS biology.* **8**, e1000357 (2010).
Csilléry, K., Rodriguez-Verdugo, K., Rellstab, C. & Guillaume, F. Detecting the genomic signal of
polygenic adaptation and the role of epistasis in evolution. *Molecular Ecology.* **27**, 606-612
(2018).
Danecek, P., Auton A., Abecasis, G., Albers C., A., Banks, E., DePristo, M. A., Handsaker, R., Lunter, G.,
Marth, G., Sherry, S. T., McVean, G., Durbin, R. and 1000 Genomes Project Analysis Group. The
variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).
Dudareva, N., Negre, F., Nagegowda, D. A. & Orlova, I. Plant Volatiles: Recent Advances and Future
Perspectives. *Critical Reviews in Plant Sciences.* **25**, 417–440 (2006).
El-Esawi, M. A. Genetic diversity and evolution of Brassica genetic resources: from morphology to novel
genomic technologies – a review. *Plant Genetic Resources: Characterization and Utilization.*
15(5), 388–399 (2017).
Fattorini, R. & Glover, B. J. Molecular Mechanisms of Pollination Biology. *Annual Review of Plant*
Biology. **71**, 487-515 (2020).
Figueroa C. M., Lunn J. E. A Tale of Two Sugars: Trehalose 6-Phosphate and Sucrose. *Plant Physiology.*
172, 7-27 (2016).
Fishman, L., Beardsley, P. M., Stathos, A., Williams, C. F. & Hill, J. P. The genetic architecture of traits
associated with the evolution of self-pollination in Mimulus. *New Phytologist.* **205**, 907–917
(2015).
Fishman, L., Kelly, A. J. & Willis, J. H. Minor quantitative trait loci underlie floral traits associated with
mating system divergence in mimulus. *Evolution.* **56**(11), 2138–2155 (2002).
Forrest, J. R. K. Plant – pollinator interactions and phenological change: what can we learn about
climate impacts from experiments and observations? *Oikos.* **124**, 4-13 (2005).
Gabriel, S. B. et al. The structure of haplotype blocks in the Human genome. *Science.* **296**, 2225-2229
(2002).

Gervasi, D. D. L. & Schiestl, F. P. Real-time divergent evolution in plants driven by pollinators. *Nature*
communications. **8**, 14691 (2017).
Hallmann, C. *et al*. More than 75 percent decline over 27 years in total flying insect biomass in
protected areas. *PloS ONE*. **12**(10), e0185809 (2017).
Hansen, T. F. The evolution of genetic architecture. *Annual Review of Ecology, Evolution, and*
Systematics. **37**, 123–57 (2006).
Hao, R., Yang, R., Zhang, Z., Zhang, Y., Chang, J. & Qiu, C. Identification and specific expression patterns
in flower organs of ABCG genes related to floral scent from *Prunus mume*. *Scientia Horticulturae*.
288, 110218 (2021).
Hayama R., Yang P., Valverde F., Mizoguchi T., Furutani-Hayama I., Vierstra R. D., Coupland G.
Ubiquitin carboxyl-terminal hydrolases are required for period maintenance of the circadian
clock at high temperature in Arabidopsis. *Scientific Reports*. **9**, 17030 (2021).
Hegland, S. J., Nielsen, A., Lázaro, A., Bjerknes, A.-L. & Totland, Ø. How does climate warming affect
plant-pollinator interactions? *Ecology Letters*. **12**, 184-195 (2009).
Hermann, K., Klahre, U., Moser, M., Sheehan, H., Mandel, T. & Kuhlemeier, C. Tight genetic linkage of
prezygotic barrier loci creates a multifunctional speciation island in *Petunia*. *Current Biology*. **23**,
873–877 (2013).
James, M. E., Wilkinson, M. J., Bernal, D. M., Liu, H., North, H. L., Engelstädter, J., Ortiz-Barrientos, D.
Phenotypic and genotypic parallel evolution in parapatric ecotypes of *Senecio*. *Evolution*.
10.1111/evo.14387 (2021).
Klahre, U. *et al*. Pollinator choice in Petunia depends on two major genetic loci for floral scent
production. *Current Biology*. **21**, 730-739 (2011).
Lin, W. *et al*. Nectar secretion requires sucrose phosphate synthases and the sugar transporter
SWEET9. *Nature*. **508**, 546-549 (2015).
Lo, S., Fatokun, C., Boukar, O., Gepts, P., Close, T. J., & Muñoz-Amatriain, M. Identification of QTL for
perenniality and floral scent in cowpea (*Vigna unguiculata* [L.] Walp.). *PloS ONE*. **15**(4),
e0229167 (2020).
Luo, Y. & Widmer, A. Herkogamy and its effects on mating patterns in *Arabidopsis thaliana. PloS ONE*.
8(2), e57902 (2013).
Minnaar, C., Anderson, V., de Jager, M. L., & Karron, J. D. Plant–pollinator interactions along the
pathway to paternity. *Annals of Botany*. **123**, 225–245 (2019).
Moeller, D. A. & Geber, M. A. Ecological context of the evolution of self-pollination in *Clarkia xantiana*:
population size, plant communities, and reproductive assurance. *Evolution*. **59**(4), 786–799
(2005).
Pritchard, J. K. & Di Rienzo, A. Adaptation – not by sweeps alone. *Nature Reviews Genetics*. **11**(10),
665–667 (2010).
Privé, F., Luu, K., Vilhjálmsson, B. J. & Blum, M. G. B. Performing highly efficient genome scans for local
adaptation with R package pcadapt version 4. *Molecular Biology and Evolution*. **37**(7), 2153–
2154 (2020).
Ramos, S. E. & Schiestl, F. P. Rapid plant evolution driven by the interaction of pollination and
herbivory. *Science*. **364**, 193–196 (2019).
Revell, L. J. learnPopGen: An R package for population genetic simulation and numerical analysis.
Ecology and Evolution. **9**, 7896–7902 (2019).
Rowlingson, B., Diggle, P. splancs: Spatial and Space-Time Point Pattern Analysis. *R package version*
2.01-43. https://CRAN.R-project.org/package=splancs (2022).
Schiestl, F. P. On the success of a swindle: pollination by deception in orchids. *Naturwissenschaften*.
92, 255–264 (2005).
Schiestl, F.P. & Johnson, S. D. Pollinator-mediated evolution of floral signals. *Trends in Ecology &*
Evolution. **28**, 307-315 (2013).

Sheng, Z., Pettersson, M. E., Honaker, C. F., Siegel, P. B. & Carlborg, O. Standing genetic variation as a
major contributor to adaptation in the Virginia chicken lines selection experiment. *Genome*
Biology. **16**, 219 (2015).
Sicard, A. & Lenhard, M. The selfing syndrome: a model for studying the genetic and evolutionary basis
of morphological adaptation in plants. *Annals of Botany*. **107**, 433–1443 (2011).
Smith, S. D. Pleiotropy and the evolution of floral integration. *New Phytologist*. **209**, 80–85 (2015).
Sobral, M., Veiga, T., Domínguez, P., Guitián, J.A., Guitián, P., & Guitián, J. M. Selective pressures
explain differences in flower color among Gentiana lutea populations. *PloS ONE*. **10**(7),
e0132522 (2015).
Solhaug, E. M. *et al*. An integrated transcriptomics and metabolomics analysis of the Cucurbita pepo
nectary implicates key modules of primary metabolism involved in nectar synthesis and
secretion. *Plant Direct*. **3**, e00120 (2019).
Streisfeld, M. A. & Rausher, M. D. Genetic changes contributing to the parallel evolution of red floral
pigmentation among Ipomoea species. *New Phytologist*. **183**, 751–763 (2009).
Swaminathan, K., Peterson, K. & Jack, T. The plant B3 superfamily. *Trends in Plant Science*. **13**, 647-655
(2008).
Thomann, M., Imbert, E. & Cheptou, P.O. Is rapid evolution of reproductive traits in Adonis annua
consistent with pollinator decline? *Acta Oecologica*. **69**, 161-166 (2015).
Van der Niet, T., Peakall, R. & Johnson, S.D. Pollinator-driven ecological speciation in plants: new
evidence and future perspectives. *Annals of Botany*. **113**, 199–211 (2014).
Wagner, D. L., Grames, E. M., Forister, M. L., Berenbaum, M. R. & Stopak, D. Insect decline in the
Anthropocene: Death by a thousand cuts. *PNAS*. **118**(2), e2023989118 (2020).
Wang C., Duan W., RamuliMaquina Riquicho A., Jing Z., Liu T., Hou X., Li Y. Genome-wide survey and
expression analysis of the *PUB* family in Chinese cabbage (*Brassica rapa* ssp*. pekinesis*).
Molecular genetics and genomics. **290**, 2241-2260 (2015).
Wessinger, C. A. & Hileman, L. C. Accessibility, constraint, and repetition in adaptive floral evolution.
Developmental Biology. **419**, 175–183 (2016).
Wickham, H. *et al*. Welcome to the tidyverse. Journal of Open. *Source Software*. **4**(43), 1686 (2019).
Wieters, B., Steige, K. A., He, F., Koch, E. M., Ramos-Onsins, S. E., Gu, H., Guo, Y. L., Sunyaev, S. & de
Meaux, J. Polygenic adaptation of rosette growth in *Arabidopsis thaliana. PloS Genet*. **17**(1),
e1008748 (2021).
Xu, S., Schlüter, P. M., Grossniklaus, U. & Schiestl, F. P. The genetic basis of pollinator adaptation in a
sexually deceptive Orchid. *PloS Genet*. **8**(8), e1002889 (2012).
Yang P., Smalle J., Lee S., Yan N., Emborg T. J., Vierstra R. D. Ubiquitin C-terminal hydrolases 1 and 2
affect shoot architecture in Arabidopsis. *The Plant Journal*. **51**, 441-457 (2007).
Yarahmadov, T., Robinson, S., Hanemian, M., Pulver, V. & Kuhlemeier, C. Identification of transcription
factors controlling floral morphology in wild Petunia species with contrasting pollination
syndromes. *The Plant Journal*. **104**(2), 289–301 (2020).
Yuan, Y. W., Byers, K. J. R. P. & Bradshaw, H. D. J. The genetic control of flower–pollinator specificity.
Current Opinion in Plant Biology. **16**, 422–428 (2013).
Zu, P. & Schiestl, F. P. The effects of becoming taller: direct and pleiotropic effects of artificial selection
on plant height in Brassica rapa. *The Plant Journal*. **89**, 1009–1019 (2017).
Zu, P., Blanckenhorn, W. U. & Schiestl, F. P. Heritability of floral volatiles and pleiotropic responses to
artificial selection in *Brassica rapa*. *New Phytologist*. **209**, 1208–1219 (2016).
498
499
500

 Legends

Figure 1. Experimental evolution design. Partial experimental evolution design from Gervasi and
Schiestl (2017) from the first generation (G1) to the ninth generation in control (C9) and bumblebee
treatment (B9). The tenth generation was obtained by inter-replicate crossing (C10 and B10).
Population of *Brassica rapa* fast cycling used for seedling sequencing are coloured. The abbreviation
and the colours assigned to the populations are unchanged along the manuscript. The number of
seedlings (n) sequenced per population is indicated.

Figure 2. Allele frequency changes during experimental evolution. (**A**) Comparison of the allele
frequency changes (Δh) between the bumblebee treatment (*x-axis*) and the control treatment (*y-axis*).
The grey dots represent the 4'713 SNPs. Replicate A is in light grey, replicate B in darker grey. The
significant changes are highlighted in blue and green (see the legend in the figure for details).
Comparison of initial (first generation) and final (ninth generation) allele frequencies in the control
treatment for both replicate A **(B)** and replicate B **(D)**, and in the bumblebee treatment for both
replicate A **(C)** and replicate B **(E)**. The grey dots represent the non-significant changes in allele
frequencies between generations. The grey solid lines indicate the maximum (upper line) or minimum
(lower line) of final simulated allele frequencies obtained by 10'000 simulations of random genetic
drift (over nine generations, N_e=16). The color dots represent important changes with *fdr* < 0.05.

Figure 3. Intersection of genomic variants under selection between treatment and replication. The
UpSet plot illustrates the genomic variants involved in the evolution of *Brassica rapa* in control and
bumblebee treatment for both replicates A and B. On the left (Set Size), the number of significant
genomic variants involved in evolutionary response of *B. rapa* in different populations is indicated. The
dots indicate that the genomic variants are involved in the evolutionary response only in a particular
population, while the vertical lines indicate that the genomic variants are involved in the evolutionary
processes in several populations. The number of genomic variants involved in unique population or in
different populations is indicated in the upper part of the figure.

Figure 4. Genomic variance within populations. (A-B) Position of the 256 individuals in the genomic
space from the principal component analysis (PCA) performed on their genotypes (GT). The PCAs were
performed **(A)** on the total set of SNPs (4'713 SNPs), and **(B)** on the 187 SNPs with significant changes
in allele frequency during bumblebee selection (fdr < 0.05). The label of the population is shown on

their centroid. Polygons linking outer points of the scatter plot are displayed for all populations. For

the generation 10 in bumblebee treatment, the polygon linking points without outlier is represented

in grey (**C**) Bar plots presenting area values of polygons for the different populations. The filled bars

are the values from PCA performed on the full dataset (4'713 SNPs), and the dashed ones on the

reduced dataset (187 SNPs). The legend colours are indicated in the bottom right of the figure. For the

generation 10 in bumblebee treatment, the polygon areas based on all samples are represented in

blue, and the one based on samples without outliers are in grey.

560
561

562   **Figure 1**

564   **Figure 2**

Figure 3

Figure 4

Supplementary information

Figure S1. Distribution of filtered genomic data for our final dataset of 4'713 SNPs. (a) Distribution
of the minor allele frequency, (b) distribution of average read depth (DP) per SNPs, (c) distribution
of the average read depth (DP) per individual, (d) distribution of the average genotype quality
(GQ), (e) different thresholds of the minimum average GQ as a function of the number of SNPs in
the final dataset. The red line indicates the chosen value in our study (--minGQ = 15).

Figure S2. Linkage disequilibrium and haplotype block structure. (A) Distribution of the median
pairwise linkage disequilibrium (r^2) for each population by distance between two SNPs (kb). **(B)**
Number of haplotype block calculated within each population **(C)** Average length (kb) of haplotype
blocks per population (more details Table S1).

Table S1. Haplotype blocks. For each population, the number of haplotype blocks, the mean (+- sd)
of the number of SNPs per haplotype block, and the length (in kb) of haplotype blocks.

	Nber blocks	mean length/block	median length/block	mean nber of SNP/block	median nber of SNP/block
A1	786	11.75	0.11	3.13	3
B1	780	10.14	0.1	3.06	3
C9A	563	47.81	0.69	4.07	3
C9B	648	39.37	0.2	3.91	3
B9A	478	65.33	20	4.33	4
B9B	600	50.71	1.13	4.12	3
C10	743	24.87	0.15	3.53	3
B10	696	38.94	0.17	3.87	3