Title: Rapid genomic evolution in *Brassica rapa* with bumblebee selection in experimental evolution

Authors: Léa Frachon[1, *], Florian P. Schiestl[1]

[1]Department of Systematic and Evolutionary Botany, University of Zürich, Zürich, Switzerland.

[*]*Corresponding author*:

Léa Frachon

lea.frachon@systbot.uzh.ch

Abstract

Background. Insect pollinators shape rapid phenotypic evolution of traits related to floral attractiveness and plant reproductive success. However, the underlying genomic changes remain largely unknown despite their importance in predicting adaptive responses to natural or to artificial selection. Based on a nine-generation experimental evolution study with fast cycling *Brassica rapa* plants adapting to bumblebees, we investigate the genomic evolution associated with the previously observed parallel phenotypic evolution. In this current evolve and resequencing (E&R) study, we conduct a genomic scan of the allele frequency changes along the genome in bumblebee-pollinated and hand-pollinated plants and perform a genomic principal component analysis (PCA).

Results. We highlight rapid genomic evolution associated with the observed phenotypic evolution mediated by bumblebees. Controlling for genetic drift, we observe significant changes in allelic frequencies at multiple loci. However, this pattern differs according to the replicate of bumblebee-pollinated plants, suggesting putative non-parallel genomic evolution. Finally, our study underlines an increase in genomic differentiation implying the putative involvement of multiple loci in short-term pollinator adaptation.

Conclusions. Overall, our study enhances our understanding of the complex interactions between pollinator and plants, providing a steppingstone towards unravelling the genetic basis of plant genomic adaptation to biotic factors in the environment.

Key words: fast cycling *Brassica rapa*, experimental evolution, evolve and resequencing study, non-parallel genomic evolution, rapid genomic evolution, bumblebees' selection.

Background

Pollinator insects are important selective agents for wild- and crop plant species due to their

essential role in the reproduction of most flowering plants [1]. While a decline of pollinator insects has

been detected in different geographical regions and insect families [2, 3, 4], the understanding of the

adaptive potential of plants to such changes remains in its infancy. Plant adaptation to pollinators

typically involves traits associated with flower attractiveness such as (1) flower morphology [5, 6, 7],

flower colour [8, 9], flower scent [10, 11, 12], and (2) traits associated with mating system like

herkogamy [13, 14] or selfing [15, 16]. While most studies assessed the result of long-term evolutionary

adaptation to pollinators, tracking the adaptive processes across generations remains scarce. Both the

resurrection approach in natural populations, growing seeds from different generations together, or

experimental evolution studies, applying the same selective pressure for multiple generations, can

bridge this gap. For instance, using a resurrection approach, Thomann et al. (17) observed phenological

and reproductive trait changes over 18 years in *Adonis annua* plants in response to the loss of wild

bees. While this approach benefits from ecological realism in natural populations, it makes it difficult

to differentiate the effect of the factor of interest from other factors such as climate, also shaping plant

evolution. Gervasi and Schiestl [12] performed experimental evolution with fast-cycling *Brassica rapa*

plants evolving with different pollinators and under controlled conditions, to identify the evolutionary

response to pollinator-mediated selection. They showed, within nine generations of experimental

evolution, rapid plant adaptation to bumblebee pollination in phenotypic traits, such as floral volatiles,

UV reflection and plant height. However, while these evolved traits have also been shown to carry

substantial heritability in this study system [18, 19], the genomic changes underlying these rapid plant

phenotypic changes are still unknown.

In the current context of pollinator decline and the associated changes in pollinator

communities, understanding the genetic architecture involved in plant response to pollinators is

essential to understand their adaptative potential in changing environments [20, 21, 22]. Molecular genetic studies have uncovered the molecular and genetic bases of several traits involved in pollination and pollinator attractiveness such as selfing [23], pollination syndromes [24, 8, 25, 26, 27], nectar [28, 29] and volatiles [30, 31, 32, 33, 34]. However, insects use a combination of signals (shape, colour, scent) and rewards for identifying suitable flowers leading to plant adaptation based on multiple traits [35]. For instance, honest signals (signals associated with reward) and pollination syndromes (convergent evolution of specific signal combinations selected by pollinators) are good examples of evolution of multiple traits. In a context of rapid environmental changes, genetic correlation among traits may allow the synchronous response of different phenotypic traits to varying patterns of selection [36, 37, 19]. While essential to predict the adaptive potential of plants to pollinators and enable breeding of crop that are more attractive to pollinators, we are still in the beginning of understanding the genetic basis involved in the adaptive response to pollinators . By combining experiment evolution and next generation sequencing (NGS), evolve and resequencing (E&R) studies have proven their ability to unravel the genomic basis involved in rapid evolution [38, 39, 40].

Here, based on previous experimental evolution performed by Gervasi and Schiestl [12] with outcrossing fast-cycling *Brassica rapa* plants, we tracked the genomic changes involved in the adaptative response of plants to bumblebee selection compared to hand pollinated control plants in an E&R study (**Figure 1**). Gervasi and Schiestl [12] initiated their experimental evolution with 108 full sib seed families in the greenhouse. Over nine generations of selection, they grew plants in three isolated replicates (36 plants per replicate). The selection was mediated by three pollination treatments during the experiment: bumblebee (*Bombus terrestris*) pollination, hoverfly- (*Episyrphus balteatus*), and hand pollination. Overall, they observed parallel phenotypic evolution with bumblebee-pollinated plants evolving taller, more fragrant flowers, and being more attractive to bumblebees. In our E&R study, we re-sequenced plants from two replicates in bumblebee and hand-

pollination treatments. We performed a genome scan of allele frequency changes and a genomic

principal component analysis to observe putative genomic evolution shaped by bumblebees.

Figure 1

Results

Genomic changes during experimental evolution. In our study, we observed allele frequency changes

(Δh) over nine generations in both bumblebee and control treatments and between two replicates

within treatments (**Figure 2**). Based on 10'000 genetic drift simulations, we defined significant changes

(colored dots in **Figure 2**) as observed changes in allele frequency outside the drift simulation ranges

(upper and lower grey lines in **Figure 2**). The allele frequency changes varied substantially between

treatments and between replicates within treatments. For instance, we found 32 SNPs with strong

($\Delta h > 0.5$) and significant (fdr < 0.05) changes in one of the replicates of the bumblebee treatment

(replicate A). In contrast, we observed only 4 SNPs with significant allele frequency changes in the

other replicate in bumblebee treatment (replicate B). In the control (hand-pollinated) treatment,

there were 3 SNPs with significant allele frequency changes in one replicate (B) and zero SNPs with

allele frequency changes in the other replicate (A). In replicate A, we found a 42-fold increase in the

number of SNPs displaying significant allele frequency changes (fdr > 0.05) with the bumblebee

treatment compared to the control treatment, regardless of its strength i.e. considering all the value

of Δ*h* (4 SNPs with significant changes in the control treatment, and 171 SNPs in the bumblebee

treatment, **Figure 2**). However, no difference in the number of SNPs involved in significant allele

frequency changes between bumblebee and control treatment were observed in replicate B (21 SNPs

with significant changes in the control, and 19 SNPs in the bumblebee treatment considering fdr < 0.05,

considering all value of Δ*h*, **Figure 2**).

Figure 2

As expected, the evolutionary process is associated with an increase of linkage disequilibrium

in the bumblebee treatment, but also in the control treatment. In both treatments, we observed a

slower decay of the median linkage disequilibrium in the ninth generation compared to the first

generation (**Figure S3A**). Within the bumblebee treatment, we observed a slower decay of LD in

replicate A than in B as expected with the observed stronger allele frequency changes (**Figure S3A**). The decays of LD in inter-replicate crossing were slower in both treatments compared to the ninth generation (**Figure S3A**). Moreover, we observed a decrease of the number of haplotype blocks, *i.e.*, genomic regions with strongly linked alleles, over nine generations (from 786 and 780 haplotype blocks for replicate A and B respectively in the first generation, to 563 and 648 haplotype blocks for replicate A and B respectively in the ninth generation of the control treatment, and 478 and 600 for replicate A and B respectively in the ninth generation of the bumblebee treatment) associated with an increase of their length (**Figure S3BC**, **Table S1**). We observed 743 and 696 haplotype blocks in control and bumblebee treatments respectively in the inter-replicate crossing (**Figure S3BC, Table S1**). It is worth noting that the moderate density of genetic markers (18 SNPs per Mb in mean among the 10 chromosomes, **Figure S1**) suggests that the estimate of LD and haplotype length may be slightly underestimated, although this has no effect on the trends observed.

Overall, the SNPs involved in the evolutive responses are independent between replicates, and between treatments (**Figure 3**). For instance, among the 171 SNPs with significant allele frequency changes in replicate A in bumblebee treatment, 164 SNPs are unique of this population, whereas 5 SNPs are shared with the control treatment in replicate B, and 3 SNPs are shared with the bumblebee treatment in replicate B (**Figure 3**).

Figure 3

Identity of candidate genes underlying genomic evolution in bumblebee treatment. After retrieving

the annotated genes around 4kb (2kb upstream, 2kb downstream) for 187 SNPs with significant allele

frequency changes in the bumblebee treatment (fdr < 0.05), we obtained a list of 171 candidate genes

(**Dataset1**). Briefly, we found genes with biological functions that may be involved in the emission of

volatiles (ABC transporter G), in the regulation of flowering time (B3), or in shoot architecture (UCH).

In addition, some genes are involved in the response to different biotic and abiotic stresses

(METACASPASE-2, PUB, Pum, PAT8, BON). Finally, some candidate genes do not have a known

biological function. However, due to a putative underestimate of LD and haplotype length (see

previous section), we may not have been able to correctly identify all the candidate genes involved in

this evolutionary process. **Genome-wide population structure.** To assess the relatedness among

individuals across generation and treatments, we performed a genomic principal component analysis

(PCA) on the full set of SNPs (4'713 SNPs). We observed a structuration of our samples determined by

populations *i.e.* generations, treatments, and replicates (**Figure 4**). Along the six first principal

components (PCs) axis, explaining 56.07% of the total genomic variance (**Figure 4A**), all individuals

from the two replicates of generation one were grouped together, and individuals from the ninth

generations were clearly separated from the first generation (**Figure 4**). In this genomic space,

individuals from the inter-replicate crossing fell between those from the scatterplot formed by

replicates A and B of generation nine for both treatments (**Figure 4**). Interestingly, in the genomic space

created by the first 6 principal components explaining the most genetic variance, the individuals

resulting from selection by bumblebees were always more scattered than those from the control

treatment or the first generation.

169 **Figure 4**

Discussion

Understanding pollinator-mediated evolutionary genomics of flowering plants remains an

important challenge in biodiversity conservation and crop improvement. Here, we screened for the

genomic consequences of biotic selection in an E&R study by sequencing genome-wide SNP markers

in *Brassica rapa* plant individuals before and after nine generations of selection by bumblebees, and

under random hand-pollination. The previous experimental evolution has shown, at the phenotypic

level, that this primarily outcrossing plant rapidly adapts to specific pollinators [12]. For instance,

Gervasi and Schiestl [12] observed taller plants with an increase in total scent emission per flower in

bumblebee-pollinated plants. The pattern of phenotypic evolution was similar between replicates

from the same pollination treatment. In this E&R study, we documented the putative signature of

directional selection driven by bumblebee pollinators with significant allele frequency changes at

several loci. Moreover, we have shown different pattern of genomic evolution between replicates, and

an increase of genomic differentiation among individuals.

In agreement with the demonstrated phenotypic evolution in the fast-cycling *Brassica rapa*

experimental system [12], we have shown genomic evolution across nine generations. The here

documented changes in allele frequencies, the increase of linkage disequilibrium and the number of

haplotype blocks, underscore the importance of pollinators in shaping plant rapid genomic evolution.

The limited number of individuals (36 per replicate and per treatment) and replicates (2 replicates)

could impact the evolutionary processes and the identification of the underlying genomic bases.

Indeed, small population size can lead to an increase in the effect of genetic drift, and a low number

of replicates to a poor estimate of loci under selection [41]. However, at the phenotypic level, the

observed differences among the three replicates were smaller than those observed among treatments,

indicating that pollinator selection played a larger role than random drift [12]. Moreover, in our study,

we accounted for genetic drift bias and identified changes in allele frequency that were greater than

those expected from 10'000 genetic drift simulations. While we cannot completely exclude some

effects of genetic drift, this alone cannot explain the patterns observed. Moreover, some significant

candidate genes identified in our E&R study have biological functions in line with the phenotypic

evolution of traits observed by Gervasi and Schiestl [12]. For instance, while the previous experimental

evolution study highlighted an increase in scent emission and plant height driven by bumblebee

pollination, we identified candidate genes who, or at least their family, are known to be involved in

emission of volatile organic compounds, like ABC transporter G family (ABCG35, ABCG38, ABCG1,

ABCB19, [42, 43]), in shoot architecture like Ubiquitin C-terminal hydrolase (UCH [44]), or in growth

and development like AUXIN-RESPONSIVE PROTEIN-RELATED, or TREHALOSE-PHOSPHATE

PHOSPHATASE E-RELATED [45]. The U-box protein is also an interesting candidate gene due to its

involvement in different developmental and physiological processes such as self-incompatibility,

defence, or abiotic stress response [46]. Moreover, we found significant allele frequency changes

associated with flowering time like a B3 transcription regulator [47] or UCH [48]. While Gervasi &

Schiestl [12] did not observe any difference in flowering time between bumblebee and control

treatment in the last generation, flowering time is an important phenological trait involved in plant-

pollinator matching [49, 50]. It is therefore possible that pollinators induce flower phenological shift,

which was not detected in the phenotypic data, but picked up in the genomic changes as changes

influencing flowering time. Overall, our study highlighted the potential involvement of multiple loci in

rapid adaptation to bumblebees, which agrees with studies highlighting such a genetic architecture

underlying floral evolution [51, 26,52, 37,53]. However, further analysis using higher quality of

sequencing and the characterization of associated phenotypic and phenological traits (allowing a

genome-wide association approach) and functional validation of the genes are required to draw more

solid conclusions about the genetic basis of evolutionary changes induced by bumblebee selection.

In addition, we found that the extent of genomic changes observed during the evolutionary

processes was different between replicates suggesting non-parallel genomic evolution [54]. While

Gervasi and Schiestl [12] have shown convergent (or parallel) phenotypic evolution for the bumblebee

treatment, our results indicated that only one replicate exhibited several loci with significant changes

in allele frequency (i.e. outside the range of drift simulations). Such a convergent phenotypic evolution

associated with non-parallel genomic evolution has been observed in artificial selection of crops [55,

56] and in natural populations. For instance, a recent study highlighted different genomic regions

underpinning the evolutionary convergence of herbicide resistance in blackgrass among different

populations [57]. Likewise, in natural populations of *Senecio lautus,* similar phenotypic variation was

reported to be regulated by variation in genomic space across populations in dune-headland coastal

habitats [58]. However, the low number of replicates in our study and the significant changes in allele

frequencies observed in only one replicate do not provide a clear distinction between potential non-

parallel evolution and a "two-speed" parallel genomic evolution (*i.e.* the second replicate will follow

the same evolutionary path in the upcoming generations). Longer experimental evolution, with more

replicates and individuals are needed to clearly understand the evolutionary pattern mediated by

biotic factors such as bumblebees.

Finally, we observed an increase of genomic variance, within the genomic space created by the

main differentiation axes, during experimental evolution mediated by bumblebees [12]. This increase

in overall genomic variance was observed among individuals in both bumblebee-pollinated replicates

(B9A and B9B), as well as in the inter-replicate crossing (B10). This pattern might be explained by the

polygenic model with a weaker selection acting on multiple standing variants (soft sweep) and by

multiple loci underlying individual phenotypic trait evolutionary changes. An increased number of

studies demonstrate the importance of polygenic adaptation [56, 60, 61, 62] related to the

infinitesimal model (reviewed in Barton et al. [63]), where local adaptation is driven by small allele

frequency changes in multiple loci. Interestingly, highly polygenic architecture involved in phenotypic

evolution could contribute to the maintenance of standing genetic variation, as recently demonstrated

in long-term artificial selection on chicken weight [64]. However, deepened analyses are needed using

newly developed models to validate the involvement of polygenic architecture in rapid phenotypic

evolution [65, 66, 67]. Multiple genes underlying phenotypic variation are widely emphasized in plants

with the advances of GWAs [59, 62], however the interplay of evolutionary forces on these genes is

still poorly understood and deserves further studies.

Conclusion

We revealed important genomic changes on multiple loci associated with convergent phenotypic evolution resulting from bumblebee selection in only nine generations. Our study is a first step into the understanding of the complex genomic mechanisms involved in rapid evolutionary adaptation to biotic factors, and we advocate further analyses to understand (1) the genetic architecture underlying phenotypic and phenological variation, (2) pleiotropic effects of quantitative-trait locus in rapid adaptation and (3) the mechanisms behind a maintenance of genetic variance. We also underline the importance of better characterizing the gene functions involved in plant-pollinator interactions. Overall, pollinators constitute complex patterns of selection which deserve more attention for predicting the adaptive responses of wild and crop plant species to their decline.

Methods

Plant material and experimental design. *Brassica rapa* (Brassicacea) is an outcrossing plant with genetic self-incompatibility, pollinated by diverse insects such as bumblebees, flies or butterflies [68]. Our study used rapid-cycling *Brassica rapa* plants (Wisconsin Fast Plants™, purchased from Carolina Biological Supply Company, Burlington, USA), selected for its short life cycle of approximately two months from seed to seed in our greenhouse conditions. We used seeds produced by the study of Gervasi and Schiestl [12], performing experimental evolution with bumblebees and control hand pollination. Briefly, starting from 108 full sib seed families, the pollination was carried out over nine generations either by bumblebees (*Bombus terrestris*), hoverflies (*Episyrphus balteatus*) or by random hand cross-pollination (control treatment). This experiment was conducted with 3 isolated replicates (one replicate includes 36 plants) for each treatment. A representative subset of seeds from all pollinated flowers was used for the next generation; the contribution of seeds to the next generation being calculated as 36 divided by the sum of seeds per replicate over all individual seeds. In this study,

we focused on bumblebee and hand-pollination treatment, using two most distinct replicates (replicate A and B). We used the offspring of the starting populations and the ninth generation for both treatments. On seed per individual plant were used. In total, we used 32 32 plants from the starting generation (generation 1) in replicate A (called A1), and 32 plants from replicate B (called B1), 32 plants from the ninth generation selected by bumblebees (bumblebee treatment) in replicate A (called B9A) and 32 plants in replicate B (called B9B)and ; 32 plants from the ninth generation of control hand pollination plants (control treatment) in replicate A (called C9A) and replicate B (C9B; **Figure 1**). Finally, we performed crossings between replicates A and B within each treatment (generation 10) yielding 32 individuals from the bumblebee treatment (inter-replicate crossing in bumblebee treatment; here called B10) and 32 individuals from the control treatment (inter-replicate crossing in control treatment; here called C10). This manual crossing is commonly used for reducing the effect of potential inbreed depression on trait changes. Pollen donors and receivers were randomly assigned. Each combination of generation*treatment*replicate is called a population (*e.g.* ninth generation, treatment bumblebees, replicate A called B9A is a population). A total of 256 seeds from these 8 populations (first, ninth and tenth generation) were sown out in a phytotron (first generation in 2017 and ninth generation as well as the inter-replicate crossing in 2019) and the leaf tissue of each plant was collected for DNA extraction and whole genomic sequencing. The study conforms to the institutional, national, and international regulations.

DNA extraction and genomic characterization. Because leaf tissue was collected in 2017 for the first generation and 2019 for the last generations, we adapted the collection storage (drying vs freezing). Leaf material from the first generation was dried in vacuum at 40 °C for 20 hours, and leaf material from the ninth and tenth generation was stored in -80°C. A high molecular weight DNA extraction (average DNA concentration of 48 ng/µL, LGC extraction protocol) and library preparation for

genotyping-by-sequencing (restriction enzyme MsII, insert size mean range ~215bp) was performed

by the LGC Genomics group Berlin. Samples were sequenced with Illumina NextSeq 500 V2 sequencer

using 150 paired-end reads; the alignment of our samples was performed with BWA version 0.7.12

against the reference genome sequence of *Brassica rapa* FPsc v1.3, Phytozome release 12

(https://phytozome.jgi.doe.gov/pz/portal.html) by the LGC Genomics group Berlin. The variant

discovery and the genotyping were realized using Freebayes v1.0.2-16 with the following parameters

by the LGC Genomic Group Berlin: --min-base-quality 10 –min-supporting-allele-qsum 10 –read-

mismatch-limit 3 –min-coverage 5 –no-indels –min-alternate-count 4 –exclude-unobserved-genotypes

–genotype-qualities –ploidy 2 or 3 –no-mnps –no-complex –mismatch-base-quality-threshold 10. We

then performed a quality trimming on the 10 chromosomes *i.e.* by discarded unassigned scaffolds from

the chromosomes using vcftools [69], by removing SNPs with missing data in more than 5% of the

individuals (function –max-missing 0.95, *i.e.* genotype calls had to be present for at least 243 samples

out of 256 for a SNP to be included in the downstream analysis), and retaining only bi-allelic SNPs with

a minimum average Phred quality score of 15 (function –minGQ 15, **Figure S2DE**). We removed SNPs

with a maximum average read depth of 100 (function –max-meanDP 100, **Figure S2BC**), and discarded

SNPs with a minor allele frequency (MAF) lower than 0.1 (function –maf 0.1, **Figure S2A**). The final

dataset   contained   4'713   SNPs   in   ~   283Mb   genome   size

(https://phytozome.jgi.doe.gov/pz/portal.html).

Allele frequency changes and genetic drift simulation. The allele frequencies of the reference allele

for the 4'713 SNPs were estimated within each 8 populations using vcftools (function –freq, [64]). To

control potential genetic drift during the nine generations of evolution, we simulated random final

allele frequencies 10'000-fold for different ranges of initial allele frequencies (from 0 to 1 by an interval

window of 0.01). The simulations were performed using the R environment package "learnPopGen"

(function "drift.selection", [70]) over eight transitions between generations (*i.e.* from the first

generation to the ninth generation) considering 32 individuals within each population for an effective

size (N_e) of 16 (*i.e.* individuals contributing to the next generation, see details of experimental evolution

in Gervasi and Schiestl [12]), and considering an equal fitness for each individual. For each SNP along

the genome, we assigned their initial allele frequency value ($AF_{initial}$) to the range estimated by the final

allele frequency simulations (10'000 values). Finally, the observed final allele frequency (AF_{final}) was

compared to 10'000 simulated allele frequency values ($AF_{simulated}$) to estimate a p-value for each SNP

using the following equations:

(1) For a decrease of reference allelic frequency *i.e.* ($AF_{initial} - AF_{final}$) > 0, *pvalue* = (number of

simulations with $AF_{simulated} \geq AF_{final}$)/10'000

(2) For an increase of reference allelic frequency *i.e.* ($AF_{initial} - AF_{final}$) < 0, *pvalue* = (number of

simulations with $AF_{simulated} \leq AF_{final}$)/10'000

(3) For ($AF_{initial} - AF_{final}$) = 0, *pvalue* = 1

With $AF_{simulated}$ = simulated final allele frequency, $AF_{initial}$ = initial allele frequency from the

reference allele (first generation), and AF_{final} = observed final allele frequency for the reference allele

(ninth generation). The pvalues were controlled for the False Discovery Rate (fdr) of 5% using

Benjamini-Hochberg method implemented in R environment [71].

Finally, we estimated the allele frequency changes (Δh) from the reference allele according to

the equation (1) for both bumblebee and control treatments:

$$\Delta h = AF_{final} - AF_{initial} \qquad (1)$$

Where Δh is the allelic frequency change between the first and the ninth generation, $AF_{initial}$ is the initial

observed allele frequency, and AF_{final} is the observed final allele frequency at the ninth generation.

Linkage disequilibrium (LD) and haplotype block structure evolution. During the selective process, an

increase of the linkage disequilibrium *i.e.* the non-random association of alleles between loci, is

expected in genomic regions strongly under selection. First, we calculated pairwise linkage

disequilibrium (LD) among all set of SNPs measured by the correlation coefficient between SNP pair r^2

within each chromosomes using VCFtools (function –geno-r2) for each 8 populations containing 32

individuals. The associated median LD was then estimated and plotted. Finally, to understand whether

changes in median LD are due to random allelic associations along the genome, or aggregated in

genomic regions under selection, we calculated the haplotype blocks in each population using plink1.9

with the following parameters: --blocks no-pheno-req –maf 0.07 –blocks-max-kb 200. This method

estimates the length of these blocks with "strong LD" considering the allelic association D' metrics

between 0.7 and 0.98 according to Gabriel et al. [72].

Candidate genes. We identified 171 candidate genes associated with 187 SNPs with significant allele

frequency changes (fdr < 0.05) during bumblebee selection. Because the median linkage disequilibrium

in bumblebee treatment is ~4kb, we retrieved the annotated genes around 4kb (2 kb upstream and 2

kb downstream) for 187 SNPs and extracted the gene description using phytozome.jgi.doe.gov.

Genome-wide variance changes. The genomic variance among individuals within each population was

estimated using principal component analysis (PCA) on scaled and centered genotype data (pcadapt

package in R environment, function pcadapt, [73]). In order to unravel the changes in genomic variance

over nine generations, we performed the PCA (1) on the total number of SNPs (4713 SNPs), and (2) on

187 SNPs showing significant allele frequency changes under bumblebee selection in on or another

replicate (fdr < 0.05). To compare genomic variance among populations, we estimated the area of the

polygon formed by scatter plots of axis 1 and 2 of each population for the full dataset (4713 SNPs) and

the subset of 187 SNPs using R package tidyverse and splancs [74, 75]. The relative area is the ratio

between the polygon area within each population and the polygon area circumventing all the points

of the PCA.

Declarations

Ethics approval and consent to participate. The study conforms to the institutional, national, and international regulations.

Consent for publication. Not applicable.

Availability of data and material. The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request. The DNA sequences of all samples will be available at National Library for Biotechnology Information (NCBI) database (BioProject PRJNA931117) after acceptance of the manuscript. Reviewers can access to the data using the following link:

https://dataview.ncbi.nlm.nih.gov/object/PRJNA931117?reviewer=hocdpu4je8n22ueldu7potnjqn

Competing interests. The authors declare that they have no competing interests.

Funding. University of Zürich provided funding.

Authors' contributions. L.F. and F.S. designed the research. L.F. performed the DNA extraction and statistical analyses and wrote the manuscript. L.F. and F.S. reviewed and edited the manuscript.

Acknowledgements

We thank Jörg Vogt for the first-generation DNA extraction, and the LGC Genomic group Berlin for the DNA extraction of other samples, the sequencing and bioinformatic treatment. We thank Cyril Zipfel for discussion regarding candidate genes. We thank Loren Rieseberg and Marius Rösti for their useful comments on a previous version of the draft. Finally, we thank the University of Zürich for their financial support.

References

1. Van der Niet T, Peakall R, Johnson SD. Pollinator-driven ecological speciation in plants: new evidence and future perspectives. *Ann Bot*. 2014;113:199–211.

2. Biesmeijer JC, et al. Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science*. 2006;313:351-354.

3. Hallmann C, et al. More than 75 percent decline over 27 years in total flying insect biomass in protected areas. *PloS ONE*. 2017;12:e0185809.

4. Wagner DL, Grames EM, Forister ML, Berenbaum MR, Stopak D. Insect decline in the Anthropocene: Death by a thousand cuts. *Proc. Natl. Acad. Sci. U.S.A*. 2020;118:e2023989118.

5. Anderson B, Pauw A, Cole WW. Barrett SCH. Pollination, mating and reproductive fitness in a plant population. *J. Evol. Biol*. 2016;29:1631–1642.

6. Wessinger CA, Hileman LC. Accessibility, constraint, and repetition in adaptive floral evolution. *Dev. Biol*. 2016;419:175–183.

7. Minnaar C, Anderson V, de Jager ML, Karron JD. Plant–pollinator interactions along the pathway to paternity. *Ann. Bot*. 2019;123:225–245.

8. Bradshaw HD, Schemske DW. Allele substitution at a flower colour locus produces a pollinator shift in monkeyflowers. *Nature*. 2003;426:176-178.

9. Sobral M, Veiga T, Domínguez P, Guitián JA, Guitián P, Guitián JM. Selective pressures explain differences in flower color among Gentiana lutea populations. *PloS ONE*. 2015;10:e0132522.

10. Schiestl FP. On the success of a swindle: pollination by deception in orchids. Sci. Nat. 2005;92:255–264.

11. Xu S, Schlüter PM, Grossniklaus U, Schiestl FP. The genetic basis of pollinator adaptation in a sexually deceptive Orchid. *PloS Genet*. 2012;8:e1002889.

12. Gervasi DDL, Schiestl FP. Real-time divergent evolution in plants driven by pollinators. *Nat. comm*. 2017;8:14691.

13. Moeller DA, Geber MA. Ecological context of the evolution of self-pollination in *Clarkia xantiana*: population size, plant communities, and reproductive assurance. *Evolution*. 2005;59:786–799.

14. Luo Y, Widmer A. Herkogamy and its effects on mating patterns in *Arabidopsis thaliana. PloS ONE*. 2013;8: e57902.

15. Bodbyl Roles SA, Kelly JK. Rapid evolution caused by pollinator loss in *Mimulus guttatus. Evolution*. 2011;65:2541–2552.

16. Ramos SE, Schiestl FP. Rapid plant evolution driven by the interaction of pollination and herbivory. *Science*. 2019;364:193–196.

17. Thomann M, Imbert E, Cheptou PO. Is rapid evolution of reproductive traits in Adonis annua consistent with pollinator decline? *Acta Oecol*. 2015;69:161-166.

18. Zu P, Blanckenhorn WU, Schiestl FP. Heritability of floral volatiles and pleiotropic responses to artificial selection in *Brassica rapa*. *New Phytol*. 2016;209:1208–1219.

19. Zu P, Schiestl FP. The effects of becoming taller: direct and pleiotropic effects of artificial selection on plant height in Brassica rapa. *Plant J*. 2017;89:1009–1019.

20. Hansen TF. The evolution of genetic architecture. *Annu. Rev. Ecol. Evol. Syst*. 2006;37:123–57.

21. Bergelson J, Roux F. Towards identifying genes underlying ecologically relevant traits in Arabidopsis thaliana. *Nat. Rev. Genet.* 2010;11:867-879.

22. Bay RA, et al. Predicting responses to contemporary environmental change using evolutionary response architectures. *Am. nat*. 2017;189:463-473.

23. Sicard A, Lenhard M. The selfing syndrome: a model for studying the genetic and evolutionary basis of morphological adaptation in plants. *Ann. Bot*. 2011;107:433–1443.

24. Bradshaw HD, Wilbert SM, Otto KG, Schemsket DW. Genetic mapping of floral traits associated with reproductive isolation in monkeyflowers (*Mimulus*). *Nature*. 1995;376:762-765.

25. Streisfeld MA, Rausher MD. Genetic changes contributing to the parallel evolution of red floral
pigmentation among Ipomoea species. *New Phytol*. 2009;183:751–763.
26. Yuan YW, Byers KJRP, Bradshaw HDJ. The genetic control of flower–pollinator specificity. *Curr.*
Opin. Plant Biol. 2013;16:422–428.
27. Yarahmadov T, Robinson S, Hanemian M, Pulver V, Kuhlemeier C. Identification of transcription
factors controlling floral morphology in wild Petunia species with contrasting pollination
syndromes. *Plant J*. 2020;104:289–301.
28. Lin W, *et al*. Nectar secretion requires sucrose phosphate synthases and the sugar transporter
SWEET9. *Nature*. 2015;508:546-549.
29. Solhaug EM, et al. An integrated transcriptomics and metabolomics analysis of the Cucurbita pepo
nectary implicates key modules of primary metabolism involved in nectar synthesis and
secretion. *Plant Direct*. 2019;3:e00120.
30. Dudareva N, Negre F, Nagegowda DA, Orlova I. Plant Volatiles: Recent Advances and Future
Perspectives. *Crit. Rev. Plant Sci*. 2006;25:417–440.
31. Klahre U, *et al*. Pollinator choice in Petunia depends on two major genetic loci for floral scent
production. *Curr. Biol*. 2011;21:730-739.
32. Cai J, Zu P, Schiestl FP. The molecular bases of floral scent evolution under artificial selection:
insights from a transcriptome analysis in Brassica rapa. *Scientific Reports*. 2016;6:36966.
33. Fattorini R, Glover BJ. Molecular Mechanisms of Pollination Biology. *Ann. Rev. Plant Biol*.
2020;71:487-515.
34. Lo S, Fatokun C, Boukar O, Gepts P, Close TJ, Muñoz-Amatriain M. Identification of QTL for
perenniality and floral scent in cowpea (*Vigna unguiculata* [L.] Walp.). *PloS ONE*.
2020;15:e0229167.
35. Schiestl FP, Johnson SD. Pollinator-mediated evolution of floral signals. *Trends in Ecology &*
Evolution. 2013;28:307-315.
36. Hermann K, Klahre U, Moser M, Sheehan H, Mandel T, Kuhlemeier C. Tight genetic linkage of
prezygotic barrier loci creates a multifunctional speciation island in *Petunia*. *Curr. Biol*.
2013;23:873–877.
37. Smith SD. Pleiotropy and the evolution of floral integration. *New Phytologist*. 2015;209:80–85.
38. Turner TL, Stewart AD, Fields AT, Rice WR, Tarone AM. Population-based resequencing of
experimentally evolved populations reveals the genetic basis of body size variation in
Drosophila melanogaster. PLoS Genet. 2011; 7:e1001336.
39. Schlötterer C, Kofler R, Versace E, Tobler R, Franssen SU. Combining experimental evolution with
next-generation sequencing: a powerful tool to study adaptation from standing genetic
variation. *Hered*. 2015;114:431-440.
40. Long A, Liti G, Luptak A, Tenaillon O. Elucidating the molecular architecture of adaptation via
evolve and resequence experiments. *Nat. Rev. Genet*. 2015;16:567-582.
41. Kofler R, Schötterer C. A guide for the design of evolve and resequencing studies. *Mol. Biol. Evol.*
2013;31:474-483.
42. Adebesin F, et al. Emission of volatile organic compounds from petunia flowers is facilitated by an
ABC transporter. *Science*. 2017; 356:1386-1388.
43. Hao R, Yang R, Zhang Z, Zhang Y, Chang J, Qiu C. Identification and specific expression patterns in
flower organs of ABCG genes related to floral scent from *Prunus mume*. *Sci. Hortic*.
2021;288:110218.
44. Yang P, Smalle J, Le7]e S, Yan N, Emborg TJ, Vierstra RD. Ubiquitin C-terminal hydrolases 1 and 2
affect shoot architecture in Arabidopsis. *Plant J*. 2007;51:441-457.
45. Figueroa CM, Lunn JE. A Tale of Two Sugars: Trehalose 6-Phosphate and Sucrose. *Plant Physiol*.
2016;172:7-27.

46. Wang C, Duan W, Ramuli Maquina Riquicho A, Jing Z, Liu T, Hou X, Li Y. Genome-wide survey and
expression analysis of the *PUB* family in Chinese cabbage (*Brassica rapa* ssp*. pekinesis*). *Mol.*
Genet. genom. 2015;290:2241-2260.
47. Swaminathan K, Peterson K, Jack T. The plant B3 superfamily. *Trends Plant Sci*. 2008;13:647-655.
48. Hayama R, Yang P, Valverde F, Mizoguchi T, Furutani-Hayama I, Vierstra RD, Coupland G.
Ubiquitin carboxyl-terminal hydrolases are required for period maintenance of the circadian
clock at high temperature in Arabidopsis. *Sci. Rep*. 2021;9:17030.
49. Forrest JRK. Plant – pollinator interactions and phenological change: what can we learn about
climate impacts from experiments and observations? *Oikos*. 2005;124:4-13.
50. Hegland SJ, Nielsen A, Lázaro A, Bjerknes AL, Totland Ø. How does climate warming affect plant-
pollinator interactions? *Ecol. Lett*. 2009;12:184-195.
51. Fishman L, Kelly AJ, Willis JH. Minor quantitative trait loci underlie floral traits associated with
mating system divergence in mimulus. *Evolution*. 2002;56:2138–2155.
52. Fishman L, Beardsley PM, Stathos A, Williams CF, Hill JP. The genetic architecture of traits
associated with the evolution of self-pollination in Mimulus. *New Phytol*. 2015;205:907–917.
53. Campitelli BE, Kenney AM, Hopkins R, Soule J, Lovell JT, Juenger TE. Genetic mapping reveals an
anthocyanin biosynthesis pathway gene potentially influencing evolutionary divergence
between two subspecies of scarlet gilia (*Ipomopsis aggregata*). *Mol. Biol. Evol*. 2017;35:807–
822.
54. Bolnick DI, Barrett RDH, Oke KB, Rennison DJ, Stuart YE. (Non)Parallel Evolution. *Ann. Rev. Ecol.*
Evol. Syst. 2018;19:303-330.
55. Schmutz J, et al. A reference genome for common bean and genome-wide analysis of dual
domestications. *Nat. genet*. 2014;46:707-713.
56. Takuno S, Ralph P, Swarts K, Elshire RJ, Glaubitz JC, Buckler ES, Hufford MB, Ross-Ibarra J.
Independent molecular basis of convergent highland adaptation in maize. *Genetics*.
2015;200:1297-1312.
57. Cai L, Comont D, MacGregor D, Lowe C, Beffa R, Neve P, Saski C. The blackgrass genome reveals
patterns of non-parallel evolution of polygenic herbicide resistance. *New Phytol*.
2023;237:1891-1907.
58. James ME, Wilkinson MJ, Bernal DM, Liu H, North HL, Engelstädter J, et al. Phenotypic and
genotypic parallel evolution in parapatric ecotypes of *Senecio*. *Evolution*. 2021;10:1111-
evo.14387.
59. Laurie CC, Chasalow SD, LeDeaux JR, McCarroll R, Bush D, Hauge B, Lai C, Clark D, Rocheford TR,
Dudley JW. The genetic architecture of response to long-term artificial selection: for oil
concentration in the maize kernel. *Genetics*;168:2141-2155.
60. Pritchard JK, Di Rienzo A. Adaptation – not by sweeps alone. *Nat. Rev. Genet*. 2010;11:665–667.
61. Csilléry K, Rodriguez-Verdugo K, Rellstab C, Guillaume F. Detecting the genomic signal of polygenic
adaptation and the role of epistasis in evolution. *Mol. Ecol*. 2018;27:606-612.
62. Barghi N, Hermisson J, Schlötterer C. Polygenic adaptation: a unifying. *Nat. Rev. Genet*.
2020;21:769-781.
63. Barton NH, Etheridge AM, Véber A. The infinitesimal model: Definition, derivation, and
implications. *Theor. Popul. Biol*. 2017;118:50-73.
64. Sheng Z, Pettersson ME, Honaker CF, Siegel PB, Carlborg O. Standing genetic variation as a major
contributor to adaptation in the Virginia chicken lines selection experiment. *Genome Biol*.
2015;16:219.
65. Berg JJ, Coop G. A population genetic signal of polygenic adaptation. *PloS Genet.*
2014;10:e1004412.
66. Boyle EA, Li YI, Pritchard J K. An expanded view of complex traits: from polygenic to omnigenic.
Cell. 2017;169:1177-1186.

67. Wieters B, Steige KA, He F, Koch EM, Ramos-Onsins SE, Gu H, et al. Polygenic adaptation of
rosette growth in *Arabidopsis thaliana. PloS Genet*. 2021;17:e1008748.
68. El-Esawi MA. Genetic diversity and evolution of Brassica genetic resources: from morphology to
novel genomic technologies – a review. *Plant Genet. Resour*. 2017;15:388–399.
69. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo, et al. The variant call format and
VCFtools. J. *Bioinform.* 2011;27:2156–2158.
70. Revell LJ. learnPopGen: An R package for population genetic simulation and numerical analysis.
Ecol. Evol. 2019;97896–7902.
71. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to
multiple testing. *J. R. Stat. Soc. Series B*. 1995;57:289-300.
72. Gabriel SB, et al. The structure of haplotype blocks in the Human genome. *Science*. 2002;
296:2225-2229.
73. Privé F, Luu K, Vilhjálmsson BJ, Blum MGB. Performing highly efficient genome scans for local
adaptation with R package pcadapt version 4. *Mol. Biol. Evol*. 2020;37:2153–2154.
74. Wickham H, et al. Welcome to the tidyverse. *J. Open Source Softw*. 2019;4:1686.
75. Rowlingson B, Diggle P. splancs: Spatial and Space-Time Point Pattern Analysis. *R package version
2.01-43*. 2022;https://CRAN.R-project.org/package=splancs

Figure titles and legends

Figure 1. Experimental evolution design. Partial experimental evolution design from Gervasi and

Schiestl [12] from the first generation (G1) to the ninth generation in control (C9) and bumblebee

treatment (B9). The tenth generation was obtained by inter-replicate crossing (C10 and B10).

Population of *Brassica rapa* fast cycling used for seedling sequencing are coloured. The abbreviation

and the colours assigned to the populations are unchanged along the manuscript. The number of

seedlings (n) sequenced per population is indicated. Images: courtesy of F. Schiestl and L. Frachon.

Figure 2. Allele frequency changes during experimental evolution. (**A**) Comparison of the allele

frequency changes (Δh) between the bumblebee treatment (*x-axis*) and the control treatment (*y-axis*).

The grey dots represent the 4'713 SNPs. Replicate A is in light grey, replicate B in darker grey. The

significant changes are highlighted in blue and green (see the legend in the figure for details).

Comparison of initial (first generation) and final (ninth generation) allele frequencies in the control

treatment for both replicate A **(B)** and replicate B **(D)**, and in the bumblebee treatment for both

replicate A **(C)** and replicate B **(E)**. The grey dots represent the non-significant changes in allele

frequencies between generations. The grey solid lines indicate the maximum (upper line) or minimum

(lower line) of final simulated allele frequencies obtained by 10'000 simulations of random genetic

drift (over nine generations, N_e=16). The color dots represent important changes with *fdr* < 0.05.

Figure 3. Intersection of genomic variants under selection between treatment and replication. The

UpSet plot illustrates the genomic variants involved in the evolution of *Brassica rapa* in control and

bumblebee treatment for both replicates A and B. On the left (Set Size), the number of significant

genomic variants involved in evolutionary response of *B. rapa* in different populations is indicated. The

dots indicate that the genomic variants are involved in the evolutionary response only in a particular

population, while the vertical lines indicate that the genomic variants are involved in the evolutionary

processes in several populations. The number of genomic variants involved in unique population or in

different populations is indicated in the upper part of the figure.

Figure 4. Genome-wide population structure. (A) Eigenvalues of the 15 principal components of the

genomic PCA based on 4713 SNPs. (B-D) Position of the 256 individuals in the genomic space from the

principal component analysis (PCA) performed on their genotypes (GT). The label of the population is

shown on their centroid. Polygons linking outer points of the scatter plot are displayed for all

populations. The legend colours are indicated in the bottom of the figure.

602
603
604