Long term analysis of social structure: evidence of age-based consistent

associations in male Alpine ibex

Alice Brambilla[a,b], Achaz von Hardenberg[c], Claudia Canedoli[d], Francesca Brivio[e], Cédric

Sueur[f,g], Christina R Stanley[c]

[a] Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Switzerland

[b] Alpine Wildlife Research Center, Gran Paradiso National Park, Italy

[c] Conservation Biology Research Group, Department of Biological Sciences, University of

Chester, UK

[d] Department of Earth and Environmental Sciences, University of Milano Bicocca, Italy

[e] Department of Veterinary Medicine, University of Sassari, Italy

[f] Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

[g] Institut Universitaire de France, Saint-Michel 103, F-75005 Paris, France

Corresponding Author:

Alice Brambilla alicebrambilla1@gmail.com, +39 3289230658

Address: Y13-J-34, Department of Evolutionary Biology and Environmental Studies,

University of Zurich. Winterthurerstrasse 190, CH-8057 Zurich (CH).

ORCID ID 0000-0003-3061-9544

ABSTRACT

Despite its recognized importance for understanding the evolution of animal sociality as well as for conservation, long term analysis of social networks of animal populations is still relatively uncommon. We investigated social network dynamics in males of a gregarious mountain ungulate (Alpine ibex, *Capra ibex*) over ten years focusing on groups, sub-groups and individuals, exploring the dynamics of sociality over different scales. Despite the social structure changing between seasons, the Alpine ibex population was highly cohesive: fission-fusion dynamics lead almost every male in the population to associate with each other male at least once. Nevertheless, we found that male Alpine ibex showed preferential associations that were maintained across seasons and years. Age seemed to be the most important factor driving preferential associations while other characteristics, such as social status, appeared less crucial. We also found that centrality measures were influenced by age and were also related to individual physical condition. The multi-scale and long-term frame of our study helped us show that ecological constrains, such as resource availability, may play a role in shaping associations in a gregarious species, but they cannot solely explain sociality and preferential association that are likely also to be driven by life-history linked physiological and social needs. Our results highlight the importance of long-term studies based on individually recognizable subjects to help us build on our understanding of the evolution of animal sociality.

Keywords: Age-based preference, Alpine ibex, *Capra ibex*, fission-fusion dynamic, mountain ungulates, Social Network Analysis

INTRODUCTION

Understanding the drivers of spatial and temporal interactions between animals is of great importance to some of the most pressing questions in biology, such as understanding the evolution of animal sociality (Couzin and Laidre 2009), determining how genetic and cultural information spread within and among inter-connected populations (Van de Waal and Bshary 2011), tracking disease transmission (MacIntosh et al. 2012; Marchand et al. 2017; Silk et al. 2019), and predicting the invasion dynamics of introduced species (Fogarty et al. 2011).

In many social species, the process of group formation is highly dynamic, with frequent changes in group size and composition (Grueter et al. 2020). Groups can merge (fusion) or split (fission) over time and space, making group composition a dynamic property (Sueur et al. 2011). Social systems characterized by fission-fusion dynamics are widespread among vertebrates as for example fish (*Poecilia reticulata*, Wilson et al. 2014), birds (Silk et al. 2014; Papageorgiou and Farine 2020) and several mammals including primates (Smuts et al. 1987), Bechstein's bats (*Myotis bechsteinii*, Kerth and König 1999), bottlenose dolphins (*Tursiops spp.,* Connor et al. 2000), spotted hyenas (*Crocuta crocuta*, Holekamp et al. 1997), African elephants (*Loxodonta africana,* Wittemyer et al. 2005), giraffes (*Giraffa camelopardalis*, Bond et al. 2019), feral goats (*Capra hircus*, Stanley and Dunbar 2013) and sheep (*Ovis aries*, Jewell et al. 1974). The timescale over which fission-fusion dynamics occur may differ as environmental conditions or individual requirements change (Aureli et al. 2008; Sueur et al. 2011). Indeed, there is pronounced variation in the degree of fission-fusion dynamics both across and within species which can vary from a highly cohesive society with stable group membership to a highly fluid society with either relatively stable or flexible subgroup membership (Aureli et al. 2008). The flexibility of fission-fusion dynamics is likely to have evolved because it allows individuals to optimize the costs and benefits of group-living (Moscovice et al. 2020). Fission-fusion dynamics are predicted to be most frequent in

environments with spatial variability and to increase with temporal uncertainty and

unpredictability of the environment (Sueur et al. 2011). Other factors known to influence the

structure and cohesion of groups are demographic processes such as deaths, births, dispersal

and immigration (Ilany and Akçay 2016) that may influence social structure through the loss

of some social connections and the formation of new ones (Shizuka et al. 2020) as well as by

driving changes in patterns of association between remaining individuals (e.g., Flack et al.

2006).

Studies on social systems often focus on the association between individuals, as members of

a group can show a preference to associate with specific individuals and maintain certain

social links despite the frequent splitting and merging of the overall group. In many social

mammals there is a tendency towards homophily, i.e., for individuals to associate with others

that share similar characteristics such as age or sex (Carranza 1984, Le Pendu et al. 1995). A

possible explanation for these preferences is that the cost of living in groups with members of

the same age class or sex is lower due to similar physiological needs and a reduction in

harmful inter or intra-sexual interactions (Conradt and Roper 2000; Ruckstuhl and Neuhaus

2000). Likewise, social relationships between individuals may be affected by kinship

(Cassinello and Calabuig 2008; Wittenmyer et al. 2009; Podgórski et al. 2014) as well as by

social rank as observed in horses (*Equus caballus,* Kimura 1998) and red deer (*Cervus*

elaphus, Appleby 1983), where individuals spent more time close to other of similar rank.

Associations can be stable or can change during the life on an individual, varying from long-

term or even life-long associations (as shown, e.g., in birds, Teitelbaum et al. 2017; wild

boars, *Sus scrofa,* Podgórski et al. 2014; feral horses and ponies, Cameron et al. 2009 and

Stanley et al. 2018) to transitory or short-term associations (e.g., in spotted hyenas, Smith et

al. 2011). Also the levels of gregariousness can differ between individuals of the same

species or population and can be influenced by various factors such as age (Carter et al. 2013;

Machanda et al. 2020) or reproductive status (Fischhoff et al. 2009; Vander Wal et al. 2015;

Machanda et al. 2020). This can, however, also be a consistent individual characteristic that is

maintained throughout life as recent studies on personality (i.e., the presence of behavioural

differences that are repeatable over time and across situations) have shown (e.g., Krause et al.

2010; Firth et al. 2018). Sociability is indeed one of the five personality traits described in

literature and it can affect reproductive success and survival and hence fitness (Réale et al.

2007).

Social network analysis (SNA) is a powerful tool that can be applied to the analysis of several

aspects of social behaviour (Sueur et al 2011; Sosa et al. 2021b). One of the most attractive

features of SNA is that it allows to study the social organisation of animals at all levels

(individual, dyad, group, population) and for all types of interaction (e.g., aggressive,

cooperative, sexual, Krause et al. 2009), allowing a plethora of novel insights into the

evolution and maintenance of sociality to be elucidated (Wey et al. 2008; Pinter-Wollman et

al. 2014; Krause et al. 2015; Webber and Vander Wal 2019; Sosa et al. 2021b). As social

structure can also affect population growth rates, dispersal and gene flow (Strier 1997),

network analysis also has the potential to be an important tool in the management of wild

populations (Tarlow and Blumstein 2007; Schakner et al. 2017, Snijders et al. 2017, Welch et

al. 2020).

To date, few studies have investigated dynamic changes in social network structures of wild

populations in the long term, especially where the individuals composing the network change

due to demographic processes (e.g., death, emigration or immigration of individuals, Cantor

et al. 2012; Borgeaud et al. 2017). Shizuka et al. (2020) called for the need to integrate

demographic processes and consequent social processes into social network analyses.

However, the scarcity of such studies is probably due to the intensity of consistent data

collection required in order to incorporate environmental and demographic changes or

stochastic events, in addition to limitations in the available analytical tools (Pinter-Wollman

et al. 2014). Instead, many studies making use of SNA in free ranging populations rely on

relatively short-term datasets and, therefore, are only able to describe snapshots of the social

systems. Since long term monitoring of wild species has provided important contributions to

the study of ecology and evolution as well as to conservation (Festa-Bianchet et al. 2017,

Clutton-Brock 2021), long term analysis of social network dynamics could bring rich rewards

in terms of a better understanding of population-level processes (Pinter-Wollman et al. 2014).

Ungulate species are widespread in all continents except Antarctica and show high diversity

both in terms of size, habitat as well as behaviour (Wilson and Mittermeier 2011). Their

social organization has been extensively studied and provided crucial elements for

understanding the evolution of vertebrates' social systems (Jarman, 1974, Krause et al. 2002).

However, many of the studies on social networks of ungulates have focused on female

associations (Vander Wal et al. 2016; Ramos et al. 2019) meaning knowledge on male

sociality in ungulates remains scarce. To fill those gaps, we performed a long-term analysis

of male social structure in a gregarious ungulate: the Alpine ibex (*Capra ibex*). We took

advantage of a detailed long-term dataset resulting from ten years of behavioural observations

of individually identifiable male Alpine ibex to explore the male social network of a

gregarious ungulate and to investigate population-level changes in social structure over time.

The Alpine ibex, a mountain ungulate of conservation concern (Brambilla et al. 2015; 2018;

2020), is a gregarious species that lives in open-membership groups and shows fission-fusion

dynamics. Alpine ibex are highly sexually dimorphic and exhibit strong sexual segregation

all year round except for during the mating season, that occurs between December and

January (Villaret and Bon 1995; Ruckstuhl and Neuhaus 2001). As the degree of sexual

segregation increases with male age (Ruckstuhl and Neuhaus 2001), outside the rut, adult

males join groups mainly composed of males, while yearlings stay in groups mainly

composed of females, and young males (≤ 2-3 years old) may move between male and female

groups.

This study was conducted during spring and summer months, when sexual segregation is at

its peak, and focused on the social network of male Alpine ibex. Specifically, we aimed to

determine: a) whether males showed consistent associations and which factors influenced the

choice of preferred companions; b) which individual characteristics (age, life stage)

influenced network metrics at the individual level and if these metrics were consistent over

time; c) whether social structure was stable across seasons (i.e., across periods with different

ecological conditions); d) whether any changes in the social network structure over time

could be explained by demographic factors.

Similarly to observations in other bovids (Vander Wal et al. 2015; Ramos et al. 2019), we

expected to find a highly connected and cohesive social network, with all the individuals

connected either directly or indirectly. Despite the aforementioned studies having analysed

social structure in females, based on field observations we expected similar results also for

males. Age was expected to predict certain network node-based measures (Turner et al. 2018;

Sosa et al. 2021b). Particularly, we expected adult males to occupy the more central positions

in the network due to their reproductive status and generally higher rank. In addition, we

expected males to become less central once they had passed their reproductive peak due to

their decreased competitive ability. Age often underpins social preferences (e.g., Wey and

Blumstein 2010; Welch et al. 2019); we therefore expected stronger associations between

males of the same age class due to their similarity in body size, social motivation, and

behaviours. At the same time, and as rank-based homophily was observed in other species

(Sosa 2016), we also expected stronger associations between males with similar social rank.

At the population level, we wanted to verify if the structure of the social network remained

stable between seasons and between years over a period of ten years. Environmental

conditions during the study period were reasonably stable within seasons so we did not

expect variations in global network measures. However, as more experienced, older

individuals are expected to help maintain group cohesion (Allen et al. 2020), we expected to

find a correlation between global centrality measures and the proportion of old males in the

population which could change due to stochastic events (e.g., harsh winters).

METHODS

Study area and population

The study was conducted on a free-ranging population of Alpine ibex in the Levionaz basin

(Valsavarenche valley, AO), within the Gran Paradiso National Park (North-western Italian

Alps - 45°26′ N, 7°08′ E).

In the study site, ibex are captured and individually marked with coloured plastic ear tags

(Allflex®: Allflex Europe (UK), 77 Greenchurch Street, London) in the framework of a long-

term study on the life-history and conservation of the species (Bassano et al. 2003; Bergeron

et al. 2010; Brambilla and Brivio 2018). The capture and marking protocol used in this study

has been authorised by the Italian Ministry of Environment (authorisation nr. 25114 of

21/09/2004) after the positive review by the Italian National Institute for Environmental

Protection and Research (ISPRA) and was developed trying to minimize the effects on the

welfare of the animals (Brambilla et al. 2013; Brivio et al. 2015).

The mean number ± SD of individuals counted in the Levionaz population during the 10

years of this study was 180 ± 41 of which 63.9 ± 10.8 were males (table 6). Censuses were

conducted every year in July block-counts by the personnel of Gran Paradiso National Park.

The proportion of marked males varied from a minimum of 50% in 2017 to a maximum of

88% in 2011 and 2012 for a total of 111 marked individuals observed during the study.

Data collection

Data were collected for male Alpine ibex during a period of ten years from 2008 to 2017 in spring (May-June) and summer (July-Aug-Sept). As Alpine ibex perform yearly altitudinal migrations following the green up of the vegetation (Parrini et al. 2003), exact dates of the seasonal ranges varied between years and were selected according to the altitudinal movement of the animals on the study site.

Individual attributes

Age. The age of the marked individuals was determined at capture by counting the number of annual horn segments (Brambilla and Canedoli 2014). The age class of the unmarked individuals was estimated by using binoculars to count the number of annual horn segments. Unmarked individuals were divided in age classes, defined as follows based on clearly visible body mass differences (Couturier 1962): 2-5 years old (young); 6-8 years old (sub-adults); 9-11 years old (fully grown adults); >11 years old (old individuals).

Season preceding death. Male Alpine ibex adult survival is very high and most of them reach senescence (Toïgo et al. 2007). However, some individuals die earlier and, particularly in case of chronic diseases, their behaviour could change in the months preceding death. We therefore recorded all deaths of identified individuals to obtain a binary variable "season preceding death"; this indicated whether the animal died within that year of data collection or if it survived until the following year. This was possible as carcasses of dead individuals were often found in the field within a few days of death. When possible, the cause of death was also ascertained. Some individuals' carcasses, however, were not found. This happened especially when animals died during winter or in an inaccessible part of the study area. If an animal was not observed in the study site nor in the surrounding areas (monitored daily by park rangers) for a whole year, we considered it as dead. We could not exclude that some of

those individuals migrated; however, as all the animals included in our study were observed

several times every year and their movements were rather predictable, we were confident in

considering them as dead if they were not observed again. Indeed, during the time of our

study, we had no cases of animals that were observed again in our or in other study areas

after one whole year when they were not observed.

Male associations

Association patterns between individuals were defined via the 'gambit of the group' method

(Whitehead 2008), i.e., assuming that each animal in a group is associating with every other

individual in that group (Croft et al. 2008; Franks et al. 2010). Association data were

collected daily by means of surveys conducted when the animals were active/feeding. When

possible, associations were recorded twice per day, after dawn and before dusk, considering

each survey as independent due to the strongly bimodal daily activity pattern typical of the

species, but time and number of daily observations varied based on seasonal and weather

conditions (details on the number of surveys per season are provided in supplementary

material S0). Data were collected by an observer scanning the study area while walking along

transects consisting of the GPNP paths used during block counts. Transect routes changed

during the season according to the altitudinal movement of the animals in the study site

(Parrini et al. 2003). Each session of data collection lasted around one to three hours

depending on the location of the animals. Every year, data were collected by two to four

observers. New observers were trained and tested by the same person that was present for the

whole duration of the study. Before new observers could start collecting data, blind

contemporary data collection with the expert observer was conducted for up to one week,

until consensus in group identification was reached. For each group, the total number of

individuals, identity of the marked individuals and age class of the unmarked individuals

were recorded. Two groups were considered as distinct if their closest members were more

than 50 metres apart. This threshold was set after field observations considered this distance

sufficient to avoid social interactions during random movements while foraging. Any animal

observed alone was considered to be a separate group. Data on association were used to build

association matrices for each season of each year (see the data analysis paragraph for details).

Male dominance

All observed agonistic interactions when an individual was clearly dominant over another, as

defined by Bergeron et al. (2009), were recorded using all occurrence sampling (Altmann

1974). Agonistic interactions were collected opportunistically during the group composition

data collection as well as during focal samples conducted for other studies. The identity of the

winner and loser in each interaction was recorded, allowing dominance matrices to be built

based on the frequency of dominance interactions per dyad. If the identity of one or both of

the individuals involved in the interaction was unknown, this interaction was not included in

the matrix.

The outcome of agonistic interactions was also used to calculate hierarchical rank via the Elo-

rating method (Elo 1978, Albers and De Vries 2001, Neumann et al. 2011). This method is

based on the sequence in which interactions occur and continuously updates ratings by

looking at interactions sequentially (Neumann et al. 2011). As previous analysis showed that

hierarchical ranks are established early in spring and remain rather stable during summer, we

calculated the Elo score (R package EloRating, Neumann et al. 2011) for each individual as

of the 30[th] of June of each year in which the individual was observed.

Network analysis

Association networks

The association network was built based on the half weight index (HWI; Whitehead 2008).

HWI is defined as:

$$HWI = \frac{X}{X + 0.5(Y_a + Y_b) + Y_{ab}} \qquad (1)$$

where X is the number of sampling periods during which individuals a and b were observed

together, Y_a is the number of sampling periods when *a* was observed without *b*, Y_b is the

number of sampling periods in which b was observed without a, and Y_{ab} is the number of

sampling periods when *a* and *b* were both observed separately (Whitehead, 2008). The HWI

is considered to be less biased when not all associating individuals can be identified

(Whitehead 2008) and it was chosen as association index to account for the unequal detection

probability typical of gregarious mountain ungulates (Vander Wal et al. 2015) and for

potential uneven sampling between individuals. As ecological conditions (e.g., resource

quality and distribution, temperatures, precipitations) change during the year and Alpine ibex

perform seasonal altitudinal migrations (Parrini et al. 2003), the HWI for each pair of

individuals was calculated separately for spring and summer of each year (2008-2017)

resulting in 20 time-aggregated association matrices (for details on the methods and tools

used to build the networks, see the data analysis section).

Dominance networks

Dominance networks were built using the absolute frequency of dominance interactions

between each pair of individuals within each season and year (as we assume that our method

allowed random sampling of dominance interactions), resulting in 20 dominance networks.

Dominance networks were non-symmetrical as the directionality of the dominance resulting

from the interaction was preserved.

Network metrics

To analyse the network structure of male Alpine ibex, we used the metrics described below.

Individual-level metrics: strength centrality and *eigenvector centrality* were calculated for

each individual in each network and averaged among individuals for each network. *Strength*

centrality is calculated as the sum of the weights of the edges in a weighted network

(Wasserman and Faust 1994). This measure represents the sociality of an individual as it

estimates the frequency of its interactions (Sosa et al. 2021a) and it was also used to compute

gregariousness at the group level (see *Global metrics* section). *Eigenvector centrality* is

defined as the first non-negative eigenvector value obtained by transforming an adjacency

matrix linearly (Wasserman and Faust 1994). It measures the centrality of a node by

examining its connections as well as that of its alters (with alters being the other individuals

connected to the node). Eigenvector centrality can therefore be interpreted as the social

resources available to an individual (Brent et al. 2011).

Global metrics: as measures to describe the global structure of the network we calculated

network density, *gregariousness* and *Typical Group Size (TGS)*. *Network density* calculates

the ratio between existing links and all potential links of a network, and it assesses the

connection of the network as a whole (Sosa et al. 2021a). *Gregariousness* represents the

tendency of individuals to associate with few or many individuals (Godde et al. 2013). At the

individual level, it is represented by *strength centrality* and it is calculated as the sum of the

values of the association indices involving that individual. The overall gregariousness of a

population is calculated as the average gregariousness of all individuals in the population.

TGS quantifies group size as experienced by an average individual of the population and it

emphasises the extent to which members of the population tend to associate (Jarman 1974). It

is an animal-centred measure defined as the sum of the squares of the number of individuals

in each group, divided by the total number of animals sampled (Jarman 1974).

Data analysis

Association matrices were built in SOCPROG 2.9 (Whitehead 2009). Network analysis was

performed in R (R Core Team 2020) using the packages tnet (Opsahl 2015), network and sna

(Butts 2020), included in the statnet suite (Statnet Development Team, 2003-2020; Handcock

et al. 2008) and with the package ANTs (Sosa et al. 2020a). Generalised Linear Mixed

Models (GLMMs) were conducted using the R packages ANTs and lme4 (Bates et al. 2015).

Details of which tool was used for each analysis are provided below.

Association patterns

To investigate association patterns, we used a Quadratic Assignment Procedure (QAP)

approach to compare pairs of association networks. The QAP test is a specialised version of

the Mantel test that uses random permutations of node labelling to determine whether a

correlation between two matrices is significantly higher than expected (Krackhardt 1988;

Krause et al. 2015). QAP tests were carried out with 10 000 permutations using the sna

package.

To test for consistency in associations between seasons, a QAP test was carried out between

the two seasonal association networks within each year. For this analysis, the networks were

built excluding individuals that were not observed across both seasons within a year.

Furthermore, to test for consistency in associations between years, we compared the networks

built for the summer seasons of the years 2012-2016. The network used for this analysis only

included a subset of individuals that were observed in all those five years (hereafter referred

to as subset networks). This time period was selected as being optimal in terms of having the

largest number of individuals present across the entire period, hence allowing the greatest

power for testing for consistency in associations; the summer seasonal network was selected

as spring associations may partly be driven by resource availability. Moreover, spring and

summer networks within each year were correlated (see Results section). QAP tests were

carried out between each subset network and the subset network of the following year.

To test whether the age difference between pairs of individuals had an impact on whether

they were likely to associate or to be dominant over each other, we built matrices of

difference in age (in years) within each dyad for each year. We built both a symmetrical

matrix with absolute age difference as well as an asymmetrical matrix with the exact age

difference between dyads. We then used QAP tests to determine whether there was

significant consistency in structure between the summer association matrix and the absolute

age difference matrix, then between the dominance matrix and the age difference matrix.

Finally, to test whether dominance rank differences between individuals predicted their

likelihood of association, we built matrices of absolute difference in their Elo scores (used as

a proxy for hierarchical rank) and performed a QAP test to test for structural consistency

between the summer association matrix and the absolute Elo score difference matrix. As this

resulted in multiple hypothesis testing, for each of the sets of QAP tests described before, we

applied a sequential Bonferroni correction (Holm 1979) for the assessment of significance

levels. A combined *p*-value for all years was calculated using Fisher's method with the

package poolr (Cinar and Viechtbauer 2020).

Factors affecting node measures and seasonal network structure

To determine whether association network structure changed between seasons (i.e., between

periods with different ecological conditions) and which individual characteristics predicted

node-based measures, we carried out Generalized Linear Mixed Models (GLMMs) on

permuted association matrices with the package ANTs: time-aggregated networks for each

season of each year were built through data stream permutations resulting in a list of 20

symmetrical association matrices (each with 10 000 permutations) used to calculate the

centrality measures of interest (strength and eigenvector centrality) and to run permuted

GLMMs.

The fixed and random structures of the permuted GLMMs were selected using non-permuted

GLMMs (lme4 package): each GLMM included as the dependent variable either strength

centrality (modelled with a gaussian distribution) or eigenvector centrality (modelled with a

binomial distribution) and, as fixed effects, the season and individual characteristics: age as a

quadratic term (Bergeron et al. 2008) and the season preceding death (as a binary variable

that indicated whether the animal died within that year of data collection or if it survived until

the following year). The models also included year and individual identity (ID) as random

effects. The choice of the fixed effects as well as the assessment of the importance of

individual ID as a random effect was made based on the Akaike Information Criterion (AIC,

Akaike 1973; Burnham et al. 2011), with Δ AIC>2 used as a threshold for the selection of the

best fitting models. Graphical analysis of the residuals and the coefficient of determination R^2

were used to check model fit. Both marginal (R^2m, that describes the proportion of variance

explained by the fixed factors alone) and conditional coefficient of determination (R^2c, which

describes the proportion of variance explained by both the fixed and random factors) were

calculated with the R package MuMIn (Barton 2009).

Seasonal and annual changes in the social structure

To investigate seasonal differences in the social structure, we run linear models to compare

the spring and summer values of the global measures (mean gregariousness and TGS,

calculated in R with own-built functions). To account for the sampling effort, the number of

surveys for each season was added as an explanatory variable to the model.

To detect possible changes in the network structure during the ten years of the study, we

calculated the global measures of density (calculated within the network R package),

gregariousness and TGS within each year (across seasons). As we hypothesised that the age

structure of the population could affect the cohesion of the social structure and hence the

above mentioned global network measures, we calculated the proportion of adult individuals

(i.e., of individuals of 9 years and more) and the total number of males in the population

(counting the maximum total number of males, including both marked and unmarked

individuals, observed during the daily observations). We then ran two separate generalised

linear models (GLM) with binomial distribution to test the effect of the proportion of old

individuals and of the total number of males on density. In addition, we ran three GLMs to

test the effect of the proportion of old individuals and of the total number of males on TGS

and to test the effect of the total number of males on gregariousness.

RESULTS

The association networks of Alpine ibex showed a high level of connectivity (e.g., figure 1)

with the network density being close to one in all the years of the study.

a) 2010 Spring network   b) 2010 Summer network

Figure 1. An example of a weighted association network of male Alpine ibex in a) spring and

b) summer of the same year (2010). Graphs were created using Gephi 0.9.2 (Bastian et al.

2009) with the "ForceAtlas" spatialization tool. The colours and width of the edges are

proportional to the strength of association (with darker and thicker edges representing

stronger associations). The size of the nodes represents the age of the individuals. The colours

of the nodes represent the value of the binary variable "season preceding death" of the

individuals during the year. Individuals that die in the months following data collection are

represented in darker colour. The graphical visualization of all networks (all season of all

years) can be found in Supplementary Materials S1.

Association patterns

QAP tests performed to test for consistent associations across seasons within each year

showed significant correlations between association matrices within all the ten years of the

study (table 1) indicating that male Alpine ibex maintain consistent bonds throughout the

year, despite changes in environment and location. Consistent associations between adjacent

years were also observed over a five year period as shown by the results of the pairwise QAP

tests performed on the subset networks between summer association networks in adjacent

years (table 2).

Table 1. Results of QAP tests to investigate consistency in association network structure

across the spring and summer seasons within each year. Significant correlations after a

Bonferroni sequential correction are indicated by bold *p*-values. Combined *p*-value < 0.001.

Seasons	Correlation coefficient	*p*-value
Spring vs summer 2008	0.250	**0.004**
Spring vs summer 2009	0.511	**<0.001**
Spring vs summer 2010	0.526	**<0.001**
Spring vs summer 2011	0.490	**<0.001**
Spring vs summer 2012	0.407	**<0.001**
Spring vs summer 2013	0.315	**<0.001**
Spring vs summer 2014	0.236	**0.016**
Spring vs summer 2015	0.460	**<0.001**
Spring vs summer 2016	0.452	**0.007**
Spring vs summer 2017	0.493	**<0.001**

432 **Table 2.** Results of QAP tests to investigate consistency in association network structure

433 between summer seasons of adjacent years. Tests were performed on subset networks

434 including N=21 individuals that were observed in all the years 2012-2016. Significant

435 correlations after a Bonferroni sequential correction are indicated by bold *p*-values.

436 Combined *p*-value < 0.001.

Years	Correlation coefficient	*p*-value
2012 vs 2013	0.236	**0.014**
2013 vs 2014	0.192	**0.028**
2014 vs 2015	0.265	**0.016**
2015 vs 2016	0.400	**0.008**

We also found that dominance matrices mostly correlated positively with age difference

matrices within the summer season for each year (table 3a), indicating that dominance

interactions were more likely to occur between individuals the greater their difference in age,

with the older individual of the dyad being more likely to be dominant over the younger.

Association matrices, instead, mostly correlated negatively with their associated absolute age

difference matrices, showing that individuals were more likely to associate with those closest

in age (table 3b). Finally, the tests between association matrices and absolute difference in

Elo score matrices showed no evidence of a correlation between association choice and

dominance rank differences or similarities (table 3c).

Table 3. Results of QAP tests for correlations between a) dominance matrices and their

associated exact age difference matrices (combined p-value <0.001); b) association matrices

and their associated and absolute age difference matrices (combined p-value <0.001); c)

association matrices and their associated absolute Elo score difference matrices. QAP tests

were performed only on the summer season matrices. Significant correlations after a

Bonferroni sequential correction are indicated by bold P values.

	a) dominance vs age-diff.		b) association vs abs. age-diff.		c) association vs abs. Elo score-diff.	
Year	Correlation coefficient	p-value	Correlation coefficient	p-value	Correlation coefficient	p-value
2008	0.078	0.039	-0.049	0.304	0.148	0.131
2009	0.233	0.033	-0.239	**0.009**	0.106	0.166
2010	0.063	0.119	-0.274	**<0.001**	0.158	0.038
2011	0.068	0.028	-0.285	**<0.001**	-0.239	0.010
2012	0.108	**0.009**	-0.320	**<0.001**	0.004	0.544
2013	0.160	**<0.001**	-0.302	**<0.001**	-0.119	0.053
2014	0.138	**0.004**	-0.070	0.230	0.043	0.342
2015	0.247	**<0.001**	-0.261	**<0.001**	-0.036	0.334
2016	0.188	**<0.001**	-0.158	**0.007**	-0.103	0.114
2017	0.108	**0.007**	-0.295	**<0.001**	0.059	0.287

Factors affecting node measures and seasonal network structure

The results of the GLMMs, performed to test whether the association network structure

changed between seasons and which individual characteristics predicted node-level centrality

measures, are presented in tables 4 and 5. Model selection results are presented in the

supplementary materials S2.

Strength centrality was significantly lower in the summer season compared to spring and was

correlated with age following a quadratic curve, i.e., it increased with age, until around 9-10

years, but then decreased when animals became older (table 4, figure 2). Strength centrality

was also negatively correlated with the season preceding death (i.e., the occurrence of death

in the months following the observations). In the best fitting model, ID was retained as a

random factor (the ΔAIC between the selected model with and without ID as random factor

was 134.1) and accounted for almost 14% of the variance explained by the model ($13.88 \pm$

3.72). The coefficients of determination of the selected model were: $R^2m=0.30$, $R^2c=0.72$.

Table 4. Output of the permuted GLMM performed to explain the variance of strength

centrality. Perm *p*-value represents the permuted *p*-values obtained after 10 000 permutations

in ANTs. Model specification: strength ~ age + age^2 + season + season preceding death +

(1|year) + (1|ID), family=gaussian.

	Estimate	Std. Error	*t*-value	Perm *p*-value
(Intercept)	10.69	2.42	4.41	<0.001
age	**2.93**	**0.41**	**7.08**	**<0.001**
age^2	**-0.18**	**0.02**	**-7.47**	**<0.001**
season	**-9.63**	**0.36**	**-26.87**	**0.022**
season prec. death	**-2.72**	**0.65**	**-4.18**	**0.001**

Consistently with results for strength centrality, eigenvector centrality was significantly lower

in the summer season and was correlated with age, increasing until around 8-9 years and then

decreasing when animals became older (table 5, figure 2). Eigenvector centrality was also

negatively correlated with the occurrence of death in the months following the observations.

ID was retained as a random factor in the selected model (ΔAIC between the selected model

with and without ID as random factor was 13.5). The coefficients of determination of the

selected model were: $R^2m=0.12$, $R^2c=0.55$.

Table 5. Output of the permuted GLMM performed to explain the variance of eigenvector

centrality. Perm p-value represents the permuted p values obtained after 10 000 permutations

in ANTs. Model specification: eigenvector ~ age + age^2 + season + season preceding death +

(1|year) + (1|ID), family= binomial.

	Estimate	Std. Error	z-value	Perm p-value
(Intercept)	-0.97	0.71	-1.37	0.137
age	**0.76**	**0.17**	**4.51**	0.060
age^2	**-0.05**	**0.01**	**-4.89**	**0.009**
season	**-0.54**	**0.18**	**-2.93**	**0.003**
season prec. death	**-0.83**	**0.25**	**-3.29**	**0.030**

Figure 2. Partial regression plots representing the relationship between a) strength and b) eigenvector centrality respectively with age. Plots were built with the package visreg (Breheny and Burchett 2017). The Y axis is scaled around the mean value of the response variable.

Seasonal and annual changes in the social structure

Annual and seasonal measures calculated to describe the global structure of the network are presented in table 6. The density of the network was close to one in all the years of the study with little variation (min.= 0.82, max.=0.99). TGS and gregariousness showed more variation over the years and between spring and summer seasons with both TGS (β=-6.66, Std. Err=2.05, p=0.005) and gregariousness (β=-5.09, Std. Err=1.43, p=0.002) being significantly lower during summer than during spring seasons (with no effect of sampling effort in either model). Gregariousness also showed differences in the coefficient of variation between years (min=0.18, max=0.40). Neither the proportion of old individuals nor the total number of

males in the population had a significant effect on density or gregariousness. The proportion

of old individuals also had no effect on TGS.

Table 6. Global measures calculated for the 2008-2017 networks. Density was calculated for

the summer networks. Gregariousness (with coefficient of variation, CV, calculated as the

ratio of the standard deviation to the mean) and TGS were calculated for both spring and

summer networks.

year	N tot	Proportion ind > 9 y	Density	Gregariousness spring mean (CV)	Gregariousness summer mean (CV)	TGS spring	TGS summer
2008	59	0.32	0.89	6.01 (0.20)	3.09 (0.40)	12.03	8.17
2009	55	0.31	0.82	5.68 (0.34)	2.74 (0.32)	14.72	11.65
2010	60	0.29	0.95	9.36 (0.26)	6.85 (0.25)	15.65	13.16
2011	57	0.26	0.92	7.70 (0.28)	5.58 (0.36)	11.90	10.08
2012	58	0.33	0.95	12.66 (0.22)	7.59 (0.28)	18.67	12.93
2013	61	0.43	0.89	12.26 (0.21)	6.55 (0.31)	20.50	9.90
2014	58	0.45	0.99	9.20 (0.20)	6.76 (0.23)	17.38	13.42
2015	63	0.40	0.98	16.51 (0.19)	6.19 (0.27)	27.07	11.27
2016	84	0.25	0.96	18.07 (0.18)	9.20 (0.27)	27.83	18.84
2017	84	0.19	0.99	11.56 (0.26)	6.84 (0.29)	22.65	16.71

DISCUSSION

We analysed the social network structure of male Alpine ibex using association and

behavioural data collected over ten years. Male and female Alpine ibex are strongly

segregated all year round, except for during the rutting season that occurs in December-

January (Ruckstuhl and Neuhaus 2001). As they do not associate during spring and summer,

our study did not consider interactions between males and females and focused on the social

network structure of males alone.

Association patterns

The association network for male Alpine ibex formed one discrete component, with a high

density of connections. Fission-fusion dynamics lead almost every male in the population to

associate with each other male at least once by being members of the same sub-group; field

observations showed frequent splitting and merging events and that all males occasionally

joined together in a single group. For example, during bad weather (i.e., mainly

thunderstorms), male ibex tend to group at specific places used as shelters, usually at a lower

altitude (A. Brambilla, *pers. obs.*). The Alpine ibex has a relatively small home range (Parrini

et al. (2003) reported an average yearly home range of around 450-500 ha for males in the

same study area) and lives in open high-altitude alpine habitat. During summer, males feed in

large alpine pasturelands where resources are not concentrated in specific sites making

resource attraction unlikely to explain the high network density observed in this species. The

high density of connections observed in male Alpine ibex was therefore probably due to the

general gregariousness of the species (Villaret and Bon 1998) that is also observed in other

mountain ungulates with a similar social structure and mating system (e.g., Bighorn sheep).

The Alpine ibex is a polygynous species with males actively tending oestrus females and

preventing subordinate males from approaching them (Willisch and Neuhaus 2009;

Apollonio et al. 2013). As the rutting season of most mountain ungulates occurs in winter,

when environmental conditions can be harsh and moving in the snow is energetically

demanding (Signer et al. 2011), the establishment of the hierarchies often begins earlier.

Indeed, agonistic interactions in Alpine ibex occur all year round except immediately after

the rutting season (Brivio et al. 2010) and are more frequent during spring and summer

months. Being part of a highly connected network may allow males of polygynous species

living in highly seasonal environments to interact with many other males reducing the need of

agonistic contests during winter, when it is more energetically costly. This advantage may

compensate for the costs of group-living that, in the case of gregarious ungulates, are

probably mostly related to infection risk (Brambilla et al. 2013; Marchand et al. 2017) as

highly connected networks allows for potentially fast spreading of diseases (Wey et al. 2008;

Marchand et al. 2017).

We found significant correlations between association matrices for spring and summer

seasons within all the ten years of the study and also between adjacent years. The correlation

coefficients were rather low, requiring caution in the interpretation of these results. However,

the fact that all correlations between spring and summer association matrices of the same year

and between summer association matrices of adjacent years were significant, seems to

indicate that there are consistent associations within the population, both across seasons and

years. These stable associations must convey some advantage as they are maintained despite

a negligible risk of predation in this population and despite seasonal changes to the global

association network structure. Our analyses carried out across a ten year period revealed that

the factors leading to assortativity in Alpine ibex appeared constant throughout the study

period. Males tended to associate with individuals of a similar age; dominance rank did not

seem to predict associations, but difference in age did predict the likelihood of engaging in

dominance interactions. The finding that age predicts association confirmed our expectation

and supports the results of previous studies (*Capra ibex,* Villaret and Bon 1995; *Capra*

pyrenaica, Alados 1986). However, we were not able to disentangle whether preferential

associations were also present within individuals in the same age class and which factors

drove these associations (e.g., kinship as observed by Cassinello and Calabuig (2008) in

aoudades). A possible explanation of the fact that age predicts association is that individuals

with similar characteristics (e.g., age and hence body size, Bergeron et al. 2010) also have

similar energetic requirements and this leads them to associate and synchronize their activity

so as to maintain group cohesion (Ruckstuhl and Neuhaus 2000; Conradt and Roper 2000).

The large differences in body size of male Alpine ibex of different ages (Bergeron et al.

2010) further strengthen this explanation and in addition suggests that life history as well can

contribute to the evolution of social structures. Furthermore, individuals may associate with

others of similar age or size because they share similar social motivations and behaviours that

enhance social cohesion and increase association (Cransac et al. 1998; Bon et al. 2001). In a

polygynous species with male-male competition that lives in a strongly seasonal environment

such as the Alps, adult males may share the need to interact to establish hierarchies before the

reproductive season (Brivio et al. 2010; Willisch and Neuhaus 2010). At the same time,

young males may share the need to interact with other males through sparring to increase

fighting skills, as observed for example in African elephants (Chiyo et al. 2011).

In this Alpine ibex population, groups of males of similar age appear to merge and split with

other groups regardless of their age. An implication of this could be that there is not a

constant transfer of information between old and young males. In this species, the transfer of

information between old and young individuals, if present, is likely to be more important for

females that have to learn where to find safe and suitable places for parturition and the

weaning of calves. In the case of males, since resources are homogeneously distributed

during the summer season and predation risk is negligible, young individuals may avoid the

costs of being constantly associated with mature males (with different time activity rhythm

and physiological needs) but yet join them on some occasions.

As expected, and not surprisingly, age difference was also positively correlated with an

increased likelihood of dominance interactions. Adult individuals were more frequently

dominant, as has also been observed in other studies on the same species and on other

polygynous mountain ungulates (Pelletier and Festa-Bianchet, 2006; Bergeron et al. 2010;

Willisch et al. 2012; Apollonio et al. 2013). The Alpine ibex is a capital breeder with a slow

life history strategy and reaches social sexual maturity at around 9-10 years of age (Willisch

et al. 2012). Young individuals are therefore often in the lower positions of the hierarchy

(Bergeron et al. 2010; Willisch et al. 2012). However, we did not find a correlation between

likelihood of association and dominance rank differences, indicating that dominance rank is

not a factor that seems to predict social preferences. Only a few of the agonistic interactions

among males escalate into physical contact. When an individual's dominance is clear, most

of the interactions consists of threat displays. Therefore, associating with individuals of

higher rank may not be costly, in terms of a risk of being injured, for low-rank males of

species that evolved threat displays as a mean to establish hierarchies.

Factors affecting node measures and seasonal network structure

Analysing the network at the node level, we also found that age predicted the position of the

individuals in the network; as we were expecting, both strength centrality and eigenvector

centrality showed a significant quadratic relationship with age. We found lower centrality

values for young and old individuals and maximum centrality values for adult individuals at

the peak of fitness (9-10 years old). Young individuals may join groups of adults but then

leave and form separate groups, thus explaining the lower centrality values observed in this

age class. Moreover, in accordance with the known life history of the species, old and

senescent individuals also showed lower strength centrality and eigenvector centrality values.

Because of their conservative life history strategy, adult survival is very high in Alpine ibex

(Toïgo et al. 2007). Many males therefore reach senescence with progressively deteriorating

body condition (Bergeron et al. 2008; Brambilla et al. 2020) and changes in activity rhythm

and spatial behaviour, meaning they are less likely to remain central in the network. Our

results suggests that social behaviour changes in Alpine ibex may occur as a result of

senescence (Siracusa et al. 2022). Similar patterns were recently observed in red deer where

Albery et al. (2021, preprint) found that older animals changed their spatial behaviour and

became less socially connected. In accordance with our interpretation, we also found that

strength and eigenvector centrality were lower if an animal died within subsequent months.

Further evidence for this explanation came from field observations as well as from the

structure of the network graphs (fig.1), where it appears that some individuals occupied a

more peripheral position in the months preceding their death (fig.1, fig. S1). In particular, this

seemed to happen for individuals that died from medium to long term diseases (e.g.,

pneumonia). In the months preceding death, those individuals were often observed alone,

away from the other males. This likely happened because the progressive deterioration of

their physical condition did not allow them to follow the daily movements and rhythm of the

other individuals. It has been observed that Alpine ibex significantly decrease investment in

horn growth in the years preceding death (von Hardenberg et al. 2004) suggesting that

energetic investment in non-vital traits is reduced when the animal is in poor physical

condition. Our observations suggest that the same is also true for behavioral traits. The

observation of changes in individual behavior due to physical conditions and health status

may have important implications for understanding the spreading dynamic of infectious

diseases (Craft 2015; Silk et al. 2019). However, we were not able to establish the cause of

death for all the individuals and therefore our interpretation of this result remains speculative;

a more detailed collection of data on the presence of infectious disease could help shed light

on this important issue.

We also found preliminary evidence that inter-individual differences other than age are likely

to play a role in determining social network positions in male ibex. Interestingly, an

individual's identity concurred to explain their association network position in terms of their

strength centrality, suggesting that sociability varies consistently between individuals

independent of their age. The presence of behavioural differences that are repeatable over

time and across situations (i.e., personality or temperament) in animals has become widely

recognized in the last few decades (Réale et al. 2007) and, since then, it has been observed in

several species, including mountain ungulates (e.g., in bighorn sheep, Réale et al. 2000; Réale

and Festa-Bianchet 2003; Poissant et al. 2013). Sociability is one of the five temperament

traits described in literature (Réale et al. 2007) and our ten-year dataset provided a unique

opportunity to follow the associations of several individuals for many years, highlighting the

potential impact of personality on network position. However, we have only found very

preliminary evidence of repeatable individual differences in sociability (i.e., retention of the

individual identity as a random effect in the selected model) while other measurements of

sociability as well as its heritability should be assessed in order to reliably quantify

personality in this species (Réale et al. 2007).

Seasonal and annual changes in the social structure

Analysing seasonal differences in the social structure, we found that mean strength and mean

eigenvector centrality in the association network, gregariousness and Typical Group Size

(TGS) were all significantly lower in the summer compared with the spring. The overall

higher values of sociality observed during spring compared to summer can partly be

explained by seasonal differences in resource availability (Peignier et al. 2019). During

spring, resources are more concentrated as there are only a few pastures available at low

altitude, where snow melts earlier. Hence, all male Alpine ibex group together there

following the winter. In summer, as snow melts and more pastures become available, male

Alpine ibex migrate to a higher altitude (Parrini et al. 2003) and therefore have fewer

constraints on where to feed and with whom to associate. Accordingly, overall gregariousness

and TGS decrease in addition to the mean strength centrality. Similar results have been found

in other ungulates, for example in reindeer (*Rangifer tarandus*): Peignier et al. (2019)

observed low home range overlap and social associations in summer, when resources were

distributed homogeneously, but higher home range overlap and social associations in winter,

when resources were distributed heterogeneously. Our results hence strengthen the

hypothesis that ecological constrains may influence social behaviour in ungulates. On the

other side, in horses, strength centrality was found to be lower outside of the mating season;

this highlighted a similar increased freedom in the choice of social partners and feeding

locations when constraints were reduced, but in this species, the constraints appeared to be

social rather than ecological (Stanley et al. 2018).

Finally, we aimed to verify if the overall structure of the social network remained consistent

over a period of ten years and whether possible changes could be explained by demographic

characteristics of the population. Due to their experience and knowledge, older individuals

are expected to help maintain group cohesion, e.g., leading group movement (McComb et al.

2001). Despite most of the study on this topic being conducted on females in matrilineal

societies (McComb et al. 2001; Brent et al. 2015), it has recently been shown that the same

may apply also to males (Allen et al. 2020). We therefore expected to find the highest values

for global centrality measures in years with a higher number of mature individuals. However,

the proportion of males over 9 years of age seemed not to affect any of the global measures

considered (TGS, gregariousness, density). Also the total number of individuals in the

population did not affect either gregariousness or density. The absence of a relationship

between global network measures and the demographic structure of the population could be

due to the seasonal movement patterns apparent in this species. Unlike animals such as

African elephants, the home range of the Alpine ibex in our study area is rather small, around

450-500 ha (Parrini et al. 2003). The seasonal movement of most male Alpine ibex occurs

within the same valley, and they show a high site fidelity, possibly because they live in a

rather predictable environment (Morrison et al. 2021). This likely makes the role of old,

experienced individuals less crucial. Similar conditions may apply to other mountain

ungulates with low dispersal rate and high site fidelity (Festa-Bianchet 1986). If this

hypothesis is confirmed, it could mean that the social cohesion of male mountain ungulates

group is stable and is not dependent on the presence of old and experienced individuals, i.e.,

it shows resilience to changes in group membership. However, we have to acknowledge that,

during the years of the study, the demographic structure of the population was not subject to

drastic changes as the environmental conditions remained relatively stable and some old

males were alive in the population across all the years of the study, even after harsh winters.

This could mean that the presence of a few older individuals may be enough to maintain

social cohesion; however, it could also signify that we did not capture enough variation to

draw biologically meaningful conclusions. As this aspect could have important conservation

implications, further data should be collected to determine the effect of demographic

structure on the social structure of mountain ungulates. Particularly, building networks on a

finer scale (e.g., daily networks) coupled with more detailed data collection on associations

(e.g., using the nearest neighbour method or conducting focal observations), as well as

accounting for spatial behaviour (Albery et al. 2021), could help in better understanding the

factors shaping associations and the global network structure.

In conclusion, this study investigated social network dynamics of male of a mountain

ungulate over a relatively long time period focusing at the level of group and individual,

allowing us to explore ungulate sociality over different scales. As most previous studies on

ungulate social networks have focused on female associations, this is one of the few to

quantify male social structure in a gregarious ungulate with high levels of sexual segregation.

We found that the male Alpine ibex social network is highly cohesive with all individuals

being connected; this has important implications for the management of disease outbreaks.

Social structure changed during seasons, suggesting that ecological constrains such as

resource availability may play a role in shaping associations in gregarious ungulates.

However, we also demonstrated that male Alpine ibex did not associate randomly but showed

stable associations, and that the strength of these associations varies across individuals, with

age similarity being a factor driving these associations. This, in turn, suggests that ecological

constrains are not enough to explain gregariousness and preferential associations in
ungulates; these are likely to also be driven by physiological and social needs, ultimately
varying according to individual characteristics and species life history. Finally, our results
highlight the importance of long-term studies based on individually recognizable subjects
(Festa-Bianchet et al. 2017).

ACKNOWLEDGEMENTS

We deeply thank Bruno Bassano from the Gran Paradiso National Park biodiversity and
scientific research service and the Park Rangers of the surveillance service for captures and
marking of Alpine ibex and for providing logistic support during field work. We are grateful
to all the students and researchers that helped with data collection in the ten years of the study
and to Edward Monovich for the artwork. We also thank David Laniado and Cristian
Pasquaretta for helpful discussion and suggestion on data analysis, and Brenda McCowan and
Sandra Smith Aguilar for constructive reviews of the manuscript.

CONFLICT OF INTEREST

736 The authors declare they have no conflict of interest relating to the content of this article.
737 Cédric Sueur is a recommender for PCI Network Science.

DATA AND SCRIPTS AVAILABILITY

739 Data and code used to produce this manuscript are deposited on the Dryad Digital Repository
740 with the following DOI: https://doi.org/10.5061/dryad.w0vt4b8st

SUPPLEMENTARY INFORMATION CAPTION

742 Supplementary information S0, S1 and S2 are deposited on bioRxiv

REFERENCES

Akaike, H. (1973). Information theory and the maximum likelihood principle in *2nd International Symposium on Information Theory* (B.N. Petrov and F. Cs ä ki, eds.). Akademiai Kiàdo, Budapest.

Alados, C. L. (1986). Spatial structure in groups of Spanish ibex (*Capra pyrenaica*). *Biol Behav*, *11*, 176-185.

Albers, P. C., & Vries, H. D. (2001). Elo-rating as a tool in the sequential estimation of dominance strengths. *Animal Behaviour,* 489-495.

Albery, G. F., Clutton-Brock, T. H., Morris, A., Morris, S., Pemberton, J. M., Nussey, D. H., & Firth, J. A. (2021). Ageing red deer alter their spatial behaviour and become less social. *bioRxiv*.

Albery, G. F., Morris, A., Morris, S., Pemberton, J. M., Clutton‐Brock, T. H., Nussey, D. H., & Firth, J. A. (2021). Multiple spatial behaviours govern social network positions in a wild ungulate. *Ecology Letters*, *24*(4), 676-686.

Allen, C. R., Brent, L. J., Motsentwa, T., Weiss, M. N., & Croft, D. P. (2020). Importance of old bulls: leaders and followers in collective movements of all-male groups in African savannah elephants (*Loxodonta africana*). *Scientific reports*, *10*(1), 1-9.

Altmann, J. (1974). Observational study of behavior: sampling methods. *Behaviour*, *49*(3-4), 227-266.

Apollonio, M., Brivio, F., Rossi, I., Bassano, B., & Grignolio, S. (2013). Consequences of snowy winters on male mating strategies and reproduction in a mountain ungulate. *Behavioural processes*, *98*, 44-50.

Appleby M. C. (1983). Competition in a red deer stag social group: rank, age and relatedness of opponents. *Animal Behaviour,* 31, 913–918.

Aureli F. et al. 2008. Fission–fusion dynamics: new research frameworks. *Current Anthropology*, 49: 627–654.

Bassano, B., von Hardenberg, A., Pelletier, F., & Gobbi, G. (2003). A method to weigh free-ranging ungulates without handling. *Wildlife Society Bulletin*, 1205-1209.

Barton, K. (2009) Mu-MIn: Multi-model inference. R Package Version 1.43.17 https://CRAN.R-project.org/package=MuMIn

Bates D, Mächler M, Bolker B & Walker S (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, **67**(1), 1–48.

Bergeron, P., Grignolio, S., Apollonio, M., Shipley, B., & Festa-Bianchet, M. (2010). Secondary sexual characters signal fighting ability and determine social rank in Alpine ibex (*Capra ibex*). *Behavioral Ecology and Sociobiology*, 64(8), 1299-1307.

793   Bond, M. L., Lee, D. E., Ozgul, A., & König, B. (2019). Fission–fusion dynamics of a
794   megaherbivore are driven by ecological, anthropogenic, temporal, and social factors.
795   *Oecologia*, *191*(2), 335-347.
796
797   Borgeaud, C., Sosa, S., Sueur, C., & Bshary, R. (2017). The influence of demographic
798   variation on social network stability in wild vervet monkeys. *Animal Behaviour*, 134, 155-
799   165.
800
801   Brambilla, A., von Hardenberg, A., Kristo, O., Bassano, B., & Bogliani, G. (2013). Don't spit
802   in the soup: faecal avoidance in foraging wild Alpine ibex, Capra ibex. *Animal Behaviour,*
803   86(1), 153-158.
804
805   Brambilla, A., & Brivio, F. (2018). Assessing the effects of helicopter disturbance in a
806   mountain ungulate on different time scales. *Mammalian Biology*, 90(1), 30-37.
807
808   Brambilla, A., & Canedoli, C. (2014). How to continue measuring horn growth after capture
809   in Alpine ibex. *Journal of Mountain Ecology,* 9.
810
811   Brambilla, A., Biebach, I., Bassano, B., Bogliani, G., & von Hardenberg, A. (2015). Direct
812   and indirect causal effects of heterozygosity on fitness-related traits in Alpine ibex.
813   *Proceedings of the Royal Society B: Biological Sciences*, 282(1798), 20141873.
814
815   Brambilla, A., Keller, L., Bassano, B., & Grossen, C. (2018). Heterozygosity–fitness
816   correlation at the major histocompatibility complex despite low variation in Alpine ibex
817   (*Capra ibex*). *Evolutionary applications*, 11(5), 631-644.
818
819   Brambilla, A., Von Hardenberg, A., Nelli, L., & Bassano, B. (2020). Distribution, status, and
820   recent population dynamics of Alpine ibex Capra ibex in Europe. *Mammal Review*, *50*(3),
821   267-277.
822
823   Breheny, P and Burchett, W (2017). Visualization of Regression Models Using visreg. *The R
824   Journal*, 9: 56-71.
825
826   Brent, L. J., Lehmann, J., & Ramos-Fernández, G. (2011). Social network analysis in the
827   study of nonhuman primates: A historical perspective. *American Journal of Primatology,*
828   73(8), 720-730.
829
830   Brent, L. J., Franks, D. W., Foster, E. A., Balcomb, K. C., Cant, M. A., & Croft, D. P. (2015).
831   Ecological knowledge, leadership, and the evolution of menopause in killer whales. *Current
832   Biology*, *25*(6), 746-750.
833
834   Brivio, F., Grignolio, S., & Apollonio, M. (2010). To feed or not to feed? Testing different
835   hypotheses on rut-induced hypophagia in a mountain ungulate. *Ethology*, *116*(5), 406-415.
836
837   Brivio, F., Grignolio, S., Sica, N., Cerise, S., & Bassano, B. (2015). Assessing the impact of
838   capture on wild animals: the case study of chemical immobilisation on alpine ibex. *PLoS
839   One*, 10(6), e0130957.
840

Burnham, K. P., Anderson, D. R., & Huyvaert, K. P. (2011). AIC model selection and
multimodel inference in behavioral ecology: some background, observations, and
comparisons. *Behavioral ecology and sociobiology*, 65(1), 23-35.

Butts, C., T. (2020). sna: Tools for Social Network Analysis. R package version 2.6.

Cameron, E. Z., Setsaas, T. H., & Linklater, W. L. (2009). Social bonds between unrelated
females increase reproductive success in feral horses. *Proceedings of the National Academy*
of Sciences, 106(33), 13850-13853.

Cantor, M., Wedekin, L. L., Guimaraes, P. R., Daura-Jorge, F. G., Rossi-Santos, M. R., &
Simoes-Lopes, P. C. (2012). Disentangling social networks from spatiotemporal dynamics:
the temporal structure of a dolphin society. *Animal Behaviour*, 84(3), 641-651.

Carter, K. D., Brand, R., Carter, J. K., Shorrocks, B., & Goldizen, A. W. (2013). Social
networks, long-term associations and age-related sociability of wild giraffes. *Animal*
Behaviour, *86*(5), 901-910.

Cassinello, J. and Calabuig, G. (2008). Spatial Association in a Highly Inbred Ungulate
Population: Evidence of Fine-Scale Kin Recognition. *Ethology*, 114: 124–132.

Chiyo, P. I., Archie, E. A., Hollister-Smith, J. A., Lee, P. C., Poole, J. H., Moss, C. J., &
Alberts, S. C. (2011). Association patterns of African elephants in all-male groups: the role of
age and genetic relatedness. *Animal Behaviour*, *81*(6), 1093-1099.

Cinar, O., & Viechtbauer, W. (2020). Poolr: methods for pooling P-values from (dependent)
tests. *R Package Version*, 9-8.

Clutton-Brock, T. (2021). Social evolution in mammals. Science, 373(6561), eabc9699.

Connor, R. C., & Wells, R. S. (2000). The bottlenose dolphin: social relationships in a
fission–fusion society. In 'Cetacean Societies: Field Studies of Dolphins and Whales'.(Eds J.
Mann, RC Connor, PL Tyack and H. Whitehead.) pp. 91–126.

Conradt, L., & Roper, T. J. (2000). Activity synchrony and social cohesion: a fission-fusion
model. *Proceedings of the Royal Society of London. Series B: Biological Sciences*,
267(1458), 2213-2218.

Couturier M.A.J. (1962) Le Bouquetin des Alps: Capra aegagrus ibex ibex. L. Arthaud,
Grenoble France

Couzin, I. D., & Laidre, M. E. (2009). Fission–fusion populations. *Current biology*, *19*(15),
R633-R635.

Cransac, N., Gerard, J. F., Maublanc, M. L., & Pépin, D. (1998). An example of segregation
between age and sex classes only weakly related to habitat use in mouflon sheep (*Ovis*
gmelini). *Journal of Zoology*, *244*(3), 371-378.
888

Craft, M. E. (2015). Infectious disease transmission and contact networks in wildlife and livestock. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *370*(1669), 20140107.

Croft D. P., James R., Krause J. (2008). Exploring Animal Social Networks. Princeton University Press.

Elo A. E. (1978). The Rating of Chess Players, Past and Present. New York: Arco.

Festa-Bianchet, M. (1986). Site fidelity and seasonal range use by bighorn rams. *Canadian Journal of Zoology*, *64*(10), 2126-2132.

Festa-Bianchet, M., Douhard, M., Gaillard, J. M., & Pelletier, F. (2017). Successes and challenges of long-term field studies of marked ungulates. *Journal of Mammalogy*, 98(3), 612-620.

Firth, J. A., Cole, E. F., Ioannou, C. C., Quinn, J. L., Aplin, L. M., Culina, A., ... & Sheldon, B. C. (2018). Personality shapes pair bonding in a wild bird social system. *Nature ecology & evolution*, *2*(11), 1696-1699.

Fischhoff, I. R., Dushoff, J., Sundaresan, S. R., Cordingley, J. E., & Rubenstein, D. I. (2009). Reproductive status influences group size and persistence of bonds in male plains zebra (Equus burchelli). *Behavioral Ecology and Sociobiology*, *63*(7), 1035-1043.

Flack, J. C., Girvan, M., De Waal, F. B., & Krakauer, D. C. (2006). Policing stabilizes construction of social niches in primates. *Nature*, 439(7075), 426-429.

Fogarty S., Cote J., Sih A. 2011. Social personality polymorphism and the spread of invasive species: a model. *The American Naturalist*, 177(3), 273-287.

Franks, D. W., Ruxton, G. D., & James, R. (2010). Sampling animal association networks with the gambit of the group. *Behavioral ecology and sociobiology*, *64*(3), 493-503.

Godde, S., Humbert, L., Côté, S. D., Réale, D., & Whitehead, H. (2013). Correcting for the impact of gregariousness in social network analyses. *Animal Behaviour*, 85(3), 553-558.

Grueter, C. C., Qi, X., Zinner, D., Bergman, T., Li, M., Xiang, Z., ... & Swedell, L. (2020). Multilevel organisation of animal sociality. *Trends in ecology & evolution*. 35(9), 834-847.

Handcock, M. S., Hunter, D. R., Butts, C. T., Goodreau, S. M., & Morris, M. (2008). statnet: Software tools for the representation, visualization, analysis and simulation of network data. *Journal of Statistical Software,* 24(1), 1548.

Holekamp, K. E., Cooper, S. M., Katona, C. I., Berry, N. A., Frank, L. G., & Smale, L. (1997). Patterns of association among female spotted hyenas (Crocuta crocuta). *Journal of Mammalogy,* 78(1), 55-64.

Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian journal of statistics*, 65-70.

939 Ilany, A., & Akcay, E. (2016). Social inheritance can explain the structure of animal social
940 networks. *Nature communications,* 7(1), 1-10.
941
942 Jarman, P. (1974). The social organisation of antelope in relation to their ecology. *Behaviour*,
943 *48*(1-4), 215-267.
944
945 Jewell P. A., Milner C., Boyd J. M. (1974). Island survivors. The ecology of the Soay sheep
946 of St. Kilda. The Athlone Press, University of London, London
947
948 Kimura, R. (1998). Mutual grooming and preferred associate relationships in a band of free-
949 ranging horses. *Applied Animal Behaviour Science*, *59*(4), 265-276.
950
951 Kerth, G., & Konig, B. (1999). Fission, fusion and nonrandom associations in female
952 Bechstein's bats (*Myotis bechsteinii*). *Behaviour*, *136*(9), 1187-1202.
953
954 Krackhardt, D. (1988). Predicting with networks: Nonparametric multiple regression analysis
955 of dyadic data. *Social networks*, *10*(4), 359-381.
956
957 Krause, J., Ruxton, G. D., Ruxton, G., & Ruxton, I. G. (2002). *Living in groups*. Oxford
958 University Press.
959
960 Krause, J., Lusseau, D., & James, R. (2009). Animal social networks: an introduction.
961 *Behavioral Ecology and Sociobiology*, *63*(7), 967-973.
962
963 Krause, J., James, R., & Croft, D. P. (2010). Personality in the context of social networks.
964 *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1560), 4099-
965 4106.
966
967 Krause, J., James, R., Franks, D. W., & Croft, D. P. (Eds.). (2015). *Animal social networks*.
968 Oxford University Press, USA.
969
970 Le Pendu, Y., Briedermann, L., Gerard, J. F., & Maublanc, M. L. (1995). Inter-individual
971 associations and social structure of a mouflon population (*Ovis orientalis musimon*).
972 *Behavioural processes*, *34*(1), 67-80.
973
974 Machanda, Z. P., & Rosati, A. G. (2020). Shifting sociality during primate ageing.
975 *Philosophical Transactions of the Royal Society B*, *375*(1811), 20190620.
976
977 MacIntosh, A. J., Jacobs, A., Garcia, C., Shimizu, K., Mouri, K., Huffman, M. A., &
978 Hernandez, A. D. (2012). Monkeys in the middle: parasite transmission through the social
979 network of a wild primate. *PloS one*, *7*(12), e51144.
980
981 Marchand, P., Freycon, P., Herbaux, J. P., Game, Y., Toïgo, C., Gilot-Fromont, E., ... & Hars,
982 J. (2017). Sociospatial structure explains marked variation in brucellosis seroprevalence in an
983 Alpine ibex population. *Scientific reports*, *7*(1), 1-12.
984
985 McComb, K., Moss, C., Durant, S. M., Baker, L., & Sayialel, S. (2001). Matriarchs as
986 repositories of social knowledge in African elephants. *Science*, *292*(5516), 491-494.
987

Morrison, T. A., Merkle, J. A., Hopcraft, J. G. C., Aikens, E. O., Beck, J. L., Boone, R. B., ...
& Kauffman, M. J. (2021). Drivers of site fidelity in ungulates. *Journal of animal ecology*,
90(4), 955-966.

Moscovice, L. R., Sueur, C., & Aureli, F. (2020). How socio-ecological factors influence the
differentiation of social relationships: an integrated conceptual framework. *Biology Letters*,
16(9), 20200384.

Neumann, C., Duboscq, J., Dubuc, C., Ginting, A., Irwan, A. M., Agil, M., ... & Engelhardt,
A. (2011). Assessing dominance hierarchies: validation and advantages of progressive
evaluation with Elo-rating. *Animal Behaviour, 82*(4), 911-921.

Opsahl, T., & Opsahl, M. T. (2015). Package 'tnet'.

Papageorgiou, D., & Farine, D. R. (2020). Multilevel societies in birds. *Trends in Ecology &*
Evolution, 36.

Parrini, F., Grignolio, S., Luccarini, S., Bassano, B., & Apollonio, M. (2003). Spatial
behaviour of adult male Alpine ibex Capra ibex ibex in the Gran Paradiso National Park,
Italy. *Acta Theriologica*, *48*(3), 411-423.

Peignier, M., Webber, Q. M., Koen, E. L., Laforge, M. P., Robitaille, A. L., & Vander Wal,
E. (2019). Space use and social association in a gregarious ungulate: Testing the conspecific
attraction and resource dispersion hypotheses. *Ecology and evolution*, *9*(9), 5133-5145.

Pelletier, F., & Festa-Bianchet, M. (2006). Sexual selection and social rank in bighorn rams.
Animal Behaviour, *71*(3), 649-655.

Pinter-Wollman, N., Hobson, E. A., Smith, J. E., Edelman, A. J., Shizuka, D., De Silva, S., ...
& Fewell, J. (2014). The dynamics of animal social networks: analytical, conceptual, and
theoretical advances. *Behavioral Ecology*, *25*(2), 242-255.

Poissant, J., Réale, D., Martin, J. G. A., Festa-Bianchet, M., & Coltman, D. W. (2013). A
quantitative trait locus analysis of personality in wild bighorn sheep. *Ecology and evolution*,
3(3), 474-481.

Podgórski, T., Lusseau, D., Scandura, M., Sönnichsen, L., & Jędrzejewska, B. (2014). Long-
lasting, kin-directed female interactions in a spatially structured wild boar social network.
PLoS One, *9*(6), e99875.

R Core Team (2020). R: A language and environment for statistical computing. Vienna,
Austria: R Computing Foundation for Science. Retrieved from http://www.R-project.org

Ramos, A., Manizan, L., Rodriguez, E., Kemp, Y. J., & Sueur, C. (2019). The social network
structure of a semi-free roaming European bison herd (Bison bonasus). *Behavioural*
processes, *158*, 97-105.

Réale, D., Dingemanse, N. J., Kazem, A. J., & Wright, J. (2010). Evolutionary and ecological
approaches to the study of personality. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, *365*(1560), 3937-3946.

Réale, D., & Festa-Bianchet, M. (2003). Predator-induced natural selection on temperament in bighorn ewes. *Animal behaviour*, *65*(3), 463-470.

Réale, D., Gallant, B. Y., Leblanc, M., & Festa-Bianchet, M. (2000). Consistency of temperament in bighorn ewes and correlates with behaviour and life history. *Animal behaviour*, *60*(5), 589-597.

Réale, D., Reader, S. M., Sol, D., McDougall, P. T., & Dingemanse, N. J. (2007). Integrating animal temperament within ecology and evolution. *Biological reviews*, *82*(2), 291-318.

Ruckstuhl, K., & Neuhaus, P. (2001). Behavioral synchrony in ibex groups: effects of age, sex and habitat. Behaviour, 138(8), 1033-1046.

Ruckstuhl, K., & Neuhaus, P. (2000). Sexual segregation in ungulates: a new approach. *Behaviour*, *137*(3), 361-377.

Schakner, Z. A., Petelle, M. B., Tennis, M. J., Van der Leeuw, B. K., Stansell, R. T., & Blumstein, D. T. (2017). Social associations between California sea lions influence the use of a novel foraging ground. *Royal Society open science*, *4*(5), 160820.

Siracusa, E. R., Higham, J. P., Snyder-Mackler, N., & Brent, L. J. (2022). Social ageing: exploring the drivers of late-life changes in social behaviour in mammals. *Biology Letters*, *18*(3), 20210643.

Shizuka, D., & Johnson, A. E. (2020). How demographic processes shape animal social networks. *Behavioral Ecology*, *31*(1), 1-11.

Signer, C., Ruf, T., & Arnold, W. (2011). Hypometabolism and basking: the strategies of Alpine ibex to endure harsh over‐wintering conditions. *Functional Ecology*, *25*(3), 537-547.

Silk, M. J., Croft, D. P., Tregenza, T., & Bearhop, S. (2014). The importance of fission–fusion social group dynamics in birds. *Ibis*, *156*(4), 701-715.

Silk, M. J., Hodgson, D. J., Rozins, C., Croft, D. P., Delahay, R. J., Boots, M., & McDonald, R. A. (2019). Integrating social behaviour, demography and disease dynamics in network models: applications to disease management in declining wildlife populations. *Philosophical Transactions of the Royal Society B*, *374*(1781), 20180211.

Smith, J. E., Powning, K. S., Dawes, S. E., Estrada, J. R., Hopper, A. L., Piotrowski, S. L., & Holekamp, K. E. (2011). Greetings promote cooperation and reinforce social bonds among spotted hyaenas. *Animal Behaviour*, *81*(2), 401-415.

Smuts B. B., Cheney D. L., Seyfarth R. M., Wrangham R. W., Struhsaker T. T. (1987). Primate societies. Chicago: The University of Chicago Press.

Snijders, L., Blumstein, D. T., Stanley, C. R., & Franks, D. W. (2017). Animal social network theory can help wildlife conservation. *Trends in ecology & evolution*, *32*(8), 567-577.

1087 Sosa, S. (2016). The influence of gender, age, matriline and hierarchical rank on individual
1088 social position, role and interactional patterns in Macaca sylvanus at 'La Forêt des singes': A
1089 multilevel social network approach. *Frontiers in psychology*, 7, 529.
1090
1091 Sosa, S., Puga-Gonzalez, I., Hu, F., Pansanel, J., Xie, X., & Sueur, C. (2020a). A multilevel
1092 statistical toolkit to study animal social networks: the Animal network toolkit Software (Ants)
1093 R package. *Scientific reports*, 10(1), 1-8.
1094
1095 Sosa, S., Sueur, C., & Puga-Gonzalez, I. (2021a). Network measures in animal social network
1096 analysis: Their strengths, limits, interpretations and uses. *Methods in Ecology and Evolution*,
1097 *12*(1), 10-21.
1098
1099 Sosa, S., Jacoby, D. M., Lihoreau, M., & Sueur, C. (2021b). Animal social networks:
1100 Towards an integrative framework embedding social interactions, space and time. *Methods in*
1101 *Ecology and Evolution*, 12,4-9.
1102
1103 Stanley, C. R., & Dunbar, R. I. M. (2013). Consistent social structure and optimal clique size
1104 revealed by social network analysis of feral goats, Capra hircus. *Animal Behaviour*, *85*(4),
1105 771-779.
1106
1107 Stanley, C. R., Mettke-Hofmann, C., Hager, R., & Shultz, S. (2018). Social stability in
1108 semiferal ponies: networks show interannual stability alongside seasonal flexibility. *Animal*
1109 *Behaviour*, *136*, 175-184.
1110
1111 Statnet Development Team (2003-2020). Statnet: Software tools for the Statistical Modeling
1112 of Network Data. http://statnet.org.
1113
1114 Sueur, C., King, A. J., Conradt, L., Kerth, G., Lusseau, D., Mettke-Hofmann, C., ... & Aureli,
1115 F. (2011). Collective decision-making and fission–fusion dynamics: a conceptual framework.
1116 *Oikos*, *120*(11), 1608-1617.
1117
1118 Tarlow, E. M., & Blumstein, D. T. (2007). Evaluating methods to quantify anthropogenic
1119 stressors on wild animals. *Applied Animal Behaviour Science*, *102*(3-4), 429-451.
1120
1121 Teitelbaum, C. S., Converse, S. J., & Mueller, T. (2017). Birds choose long-term partners
1122 years before breeding. *Animal Behaviour*, *134*, 147-154.
1123
1124 Toïgo, C., Gaillard, J. M., Festa-Bianchet, M.., Largo, E., Michallet, J., & Maillard, D.
1125 (2007). Sex-and age-specific survival of the highly dimorphic Alpine ibex: evidence for a
1126 conservative life-history tactic. *Journal of Animal Ecology*, *76*(4), 679-686.
1127
1128 Turner, J. W., Bills, P. S., & Holekamp, K. E. (2018). Ontogenetic change in determinants of
1129 social network position in the spotted hyena. *Behavioral ecology and sociobiology*, *72*(1), 10.
1130
1131 van de Waal, E., & Bshary, R. (2011). Social-learning abilities of wild vervet monkeys in a
1132 two-step task artificial fruit experiment. *Animal Behaviour*, *81*(2), 433-438.
1133
1134 Vander Wal, E., Festa-Bianchet, M., Réale, D., Coltman, D. W., & Pelletier, F. (2015).
1135 Sex-based differences in the adaptive value of social behavior contrasted against morphology
1136 and environment. *Ecology*, *96*(3), 631-641.

1137
1138 Vander Wal, E., Gagné-Delorme, A., Festa-Bianchet, M., & Pelletier, F. (2016). Dyadic
1139 associations and individual sociality in bighorn ewes. *Behavioral Ecology*, *27*(2), 560-566.
1140
1141 von Hardenberg, A., Bassano, B., Arranz, M. D. P. Z., & Bogliani, G. (2004). Horn growth
1142 but not asymmetry heralds the onset of senescence in male Alpine ibex (*Capra ibex*). *Journal*
1143 *of Zoology*, *263*(4), 425-432.
1144
1145 Wasserman, S., & Faust, K. (1994). *Social network analysis: Methods and applications* (Vol.
1146 8). Cambridge university press.
1147
1148 Welch, M. J., Smith, T., Hosie, C., Wormell, D., Price, E., & Stanley, C. R. (2020). Social
1149 Experience of Captive Livingstone's Fruit Bats (*Pteropus livingstonii*). *Animals*, *10*(8), 1321.
1150
1151 Wey, T., Blumstein, D. T., Shen, W., & Jordán, F. (2008). Social network analysis of animal
1152 behaviour: a promising tool for the study of sociality. *Animal behaviour*, *75*(2), 333-344.
1153
1154 Wey, T. W., & Blumstein, D. T. (2010). Social cohesion in yellow-bellied marmots is
1155 established through age and kin structuring. *Animal Behaviour*, *79*(6), 1343-1352.
1156
1157 Wilson, A. D., Krause, S., James, R., Croft, D. P., Ramnarine, I. W., Borner, K. K., ... &
1158 Krause, J. (2014). Dynamic social networks in guppies (Poecilia reticulata). *Behavioral*
1159 *Ecology and Sociobiology*, *68*(6), 915-925.
1160
1161 Wilson, D. E., & Mittermeier, R. A. (2011). Handbook of the mammals of the word. Vol. 2:
1162 Hoofed mammals. Lynx Editions.
1163
1164 Villaret, J. C., & Bon, R. (1995). Social and spatial segregation in Alpine ibex (Capra ibex) in
1165 Bargy, French Alps. *Ethology*, *101*(4), 291-300.
1166
1167 Villaret J. C., Bon, R. (1998). Sociality and relationships in Alpine ibex (Capra ibex). *Revue*
1168 *d'écologie*.
1169
1170 Whitehead, H. (2008). Analysing animal societies: Quantitative Methods for Vertebrate
1171 Social Analysis. Chicago: University of Chicago Press.
1172
1173 Whitehead, H. (2009). SOCPROG programs: analysing animal social structures. *Behavioral*
1174 *Ecology and Sociobiology*, *63*(5), 765-778.
1175
1176 Williams, R., & Lusseau, D. (2006). A killer whale social network is vulnerable to targeted
1177 removals. *Biology letters*, *2*(4), 497-500.
1178
1179 Willisch, C. S., & Neuhaus, P. (2009). Alternative mating tactics and their impact on survival
1180 in adult male Alpine ibex (*Capra ibex ibex*). *Journal of Mammalogy*, *90*(6), 1421-1430.
1181
1182 Willisch, C. S., & Neuhaus, P. (2010). Social dominance and conflict reduction in rutting
1183 male Alpine ibex, *Capra ibex*. *Behavioral Ecology*, *21*(2), 372-380.
1184
1185 Willisch, C. S., Biebach, I., Koller, U., Bucher, T., Marreros, N., Ryser-Degiorgis, M. P., ...
1186 & Neuhaus, P. (2012). Male reproductive pattern in a polygynous ungulate with a slow life-

1187   history: the role of age, social status and alternative mating tactics. *Evolutionary Ecology*,
1188   *26*(1), 187-206.
1189
1190   Wittemyer, G., Okello, J. B., Rasmussen, H. B., Arctander, P., Nyakaana, S., Douglas-
1191   Hamilton, I., & Siegismund, H. R. (2009). Where sociality and relatedness diverge: the
1192   genetic basis for hierarchical social organization in African elephants. *Proceedings of the*
1193   *Royal Society B: Biological Sciences*, *276*(1672), 3513-3521.