Secreted ORF8 is a pathogenic cause of severe COVID-19 and is potentially targetable with select NLRP3 inhibitors

Xiaosheng Wu[1]*, Michelle K. Manske[1], Gordon J. Ruan[1], Taylor L. Witter[2], Kevin E. Nowakowski[1], Jithma P. Abeykoon[1], Xinyi Tang[1], Yue Yu[3], Kimberly A. Gwin[1], Annie Wu[4], Vanessa Taupin[6], Vaishali Bhardwaj[1], Jonas Paludo[1], Surendra Dasari[3], Haidong Dong[4], Stephen M. Ansell[1], Andrew D. Badley[5], Matthew J. Schellenberg[2], Thomas E. Witzig[1]*

[1] Division of Hematology, Department of Medicine, [2] Department of Biochemistry and Molecular Biology, [3] Department of Quantitative Health Sciences, [4] Department of Immunology, [5] Division of infectious diseases, Department of Medicine, Mayo Clinic, Rochester, MN, USA; [6] Electron Microscopy Core, University of California San Diego, La Jolla, CA, USA

Correspondence:

        Dr. Xiaosheng Wu (507-293-0058, wu.xiaosheng@mayo.edu,) or
        Dr. Thomas E. Witzig (507-284-6134, Thomas.witzig@mayo.edu)
        Division of Hematology, Department of Medicine
        Mayo Clinic
        Rochester, MN 55905, USA
        Fax: (507) 266-9277

Abstract:	153 words
Main text:	3972 words
Main Figure:	7
Suppl Figure:	7
Suppl Table:	4
Reference:	35

Scientific Category: inflammation, monocytes, cytokines/chemokines.

Key points:

- Secreted glycoprotein ORF8 induces monocytic pro-inflammatory cytokines involving the activation of the NLPR3 inflammasome pathway.

- ORF8 is prognostically present in the blood of symptomatic patients with covid-19 and is targetable with NLRP3 inhibitor MCC-950.

41   **ABSTRACT**

43   COVID-19 is a significant cause of morbidity and mortality in blood cancer

patients, especially those on immunosuppressive therapy. Despite extensive research,

the specific factor associated with SARS-CoV-2 infection that mediates the life-

threatening inflammatory cytokine response in patients with severe COVID-19 remains

unidentified. Herein we demonstrate that the virus-encoded Open Reading Frame 8

(ORF8) protein is abundantly secreted as a glycoprotein *in vitro* and in symptomatic

patients with COVID-19. ORF8 specifically binds to the NOD-like receptor family pyrin

domain-containing 3 (NLRP3) in CD14$^+$ monocytes to induce a non-canonical

inflammasomal response, and a canonical response when the second activation signal is

present. Levels of ORF8 protein in the blood correlate with severity and disease-specific

mortality in patients with acute SARS-CoV-2 infection. Furthermore, the ORF8-induced

inflammasome response was readily inhibited by the NLRP3 inhibitor MCC950 *in vitro*.

Our study identifies a dominant cause of pathogenesis, its underlying mechanism, and

a potential new treatment for severe COVID-19.

INTRODUCTION

COVID-19, the global pandemic caused by infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has infected more than half a billion people and caused over six million deaths worldwide.[1] Numerous studies have shown that the production of pro-inflammatory cytokines/chemokines including IL1β, IL6, IL8, and CCL2 is responsible for life-threatening symptoms.[2,3] It is also known that the viral load, cytokine levels, and disease severity are tightly associated[4-6], and the virus-neutralizing antibodies and IL1β pathway antagonists could readily mitigate symptoms and improve clinical outcomes.[7,8] However, the intermediate viral factor that directly causes the inflammatory cytokine responses remains unidentified. It has been demonstrated that SARS-CoV-2 infection localizes to nasal and pulmonary epithelial cells,[9,10] while the cytokine response is more systemic. It seems irreconcilable how this cytokine response is initiated given that no live virus has been reported in the blood based on transfusion medicine studies.[11-14] We hypothesized that an inflammatory byproduct of SARS-CoV-2 replication is released into the bloodstream resulting in a systematic cytokine response in severe COVID-19 patients.

Upon infection of human cells, the SARS-CoV-2 virus replicates its 29.9kb RNA genome and produces up to 29 possible viral proteins, including 16 non-structural proteins (NSP1-16), four structural proteins Spike (SPK), Membrane (MEM), Envelope (ENV) and Nucleocapsid (NUC) and nine accessory proteins (ORF3A, 3B, 6, 7A, 7B, 8, 9b, 9c, and 10). While only four structural proteins along with the RNA genome are assembled into new viral particles, the other viral proteins are thought to be left behind[15] which may disrupt host cell functions.[16,17] Herein, we demonstrate that the SARS-CoV-2 encoded ORF8 is abundantly secreted as a glycoprotein into culture supernatant *in vitro* and into the bloodstream in patients with COVID-19. Glycosylated

84   ORF8 stimulates CD14$^+$ monocytes to produce a group of pro-inflammatory cytokines

85   including IL1β through an NLRP3-mediated inflammasome response, and the

86   ORF8/NLRP3 axis is targetable by NLRP3 inhibitor MCC-950.

MATERIALS AND METHODS

Study Samples

Human serum samples were collected from Mayo Clinic patients with a documented diagnosis of COVID-19 infection and consented to COVID-19 Research Task Force Specimen Biobank. Samples for these studies were requested and approved by the Task Force Review Committee and the Mayo Clinic Institutional Review Board. Deidentified fresh leukocyte cones of healthy donors were obtained from Mayo Clinic Blood Bank. Use of mouse splenic B cells from C57/B6 mice was approved by the Institutional Animal Care and Use Committee of the Mayo Clinic.

SARS-CoV-2 protein constructs and their expression in human cells

Lentiviral constructs expressing SARS-CoV-2 proteins originally described by Gordon, et al [16] were purchased from Addgene (Supplemental Table 1), and used to make stable expression cell lines in HEK293 cells by lentiviral transduction, and transient expression in HEK293F cells with HyCell TransFx media (Cytiva) method for ORF8 protein purification. During the study, we first used the conditioned media (CM) (made by dialyzing the culture supernatants of ORF8-expressing HEK293 cells) for PBMC stimulation, it was later replaced with purified glycosylated ORF8.

Stimulation PBMCs or THP-1 and cytokine detection

Fresh PBMCs (3×10^6) isolated from healthy donors were stimulated with either 20% (v/v) of CM-ORF8 for 72 hours, or 200ng/ml pure ORF8 for 24 hours in a total volume of 1.5ml RPMI/10% FCS based on the titration results (Supplemental Figure 6). The cytokines expression was determined using qPCR (SYBR-green method, primers are listed in Supplemental Table 3), Luminex using a custom procarta Luminex 6-plex-cytokines (IL1β, IL6, IL8, IL18, CCL2, and TNFα) detection kit (ThermoFisher)

(Supplemental Fig. 5), intracellular flowcytometry, or Western blotting. For qPCR,

housekeeping gene β-Actin or HPRT1 was used for normalization.

Detection of ORF8 protein in sera of patients newly infected with SARS-CoV-2

For detection of ORF8 protein in serum samples from patients newly infected

with SARS-CoV-2, typically four microliters of serum were directly mixed with 36

microliters of 2x Laemmli buffer with 5% of BME, boiled for 3 min before load to 10-

20% SDS-PAGE gels, the blots were probed with 1:5000 diluted rabbit anti-ORF8

antibody (MyBioSource, Cat# MBS3014575) at 4°C overnight. The images were

developed using an ECL reagent kit.

Analysis of ORF8 expression and survival in patients with COVID-19

Overall survival was calculated from the time of COVID-19 diagnosis, using the

Kaplan-Meir method on JMP 14.0 software (SAS Institute, Cary, NC).

Additional methods are described in supplemental Materials and methods.
131

RESULTS

Secretion of SARS-CoV-2 proteins from human cells

To determine the secretion potential of SARS-CoV-2 encoded proteins from human cells, we transduced HEK293 cells with one of 22 lentiviral constructs available to us each expressing a SARS-CoV-2 protein tagged Strep II (Supplemental Table 1). The secreted viral proteins in the culture supernatants were enriched and analyzed by Western blotting, and their levels were compared to their total cellular expression in whole cell lysates. As shown in Fig. 1a, most viral proteins were robustly expressed with the expected molecular sizes except for NSP13, ORF10, MEM, and NSP4 due to their size or solubility issues. While NSP2, 5, 7, 9, 10, 12, 14, 15, ORF8, and NUC proteins were all detected as secreted proteins, ORF8 was the single most robustly secreted protein (Fig. 1b). This is consistent with *in silico* analysis (Supplemental Fig. 2) showing that all secreted NSPs possess an unconventional protein secretion signal (UPS) while ORF8 carries a classical protein signal sequence (SS) and an N-link glycosylation sites (^{78}NYTV). These results suggest that ORF8 is efficiently secreted through the classical ER/Golgi protein secretion pathway, and its glycosylation explains the up-shifting and smearing of the ORF8 protein band (Fig. 1b).

Secreted ORF8 protein induces the production of pro-inflammatory cytokines

We then queried if any of the secreted SARS-CoV-2 proteins could induce the expression of pro-inflammatory cytokines seen in patients with COVID-19. We treated human peripheral blood mononuclear cells (PBMCs) from healthy donors with conditioned media (CM) containing major secreted NSPs or ORF8 proteins followed by an assessment of cytokine expression. We found that none of the secreted NSP proteins induced cytokine expression (Fig. 2a) while ORF8-containing CM (ORF8-CM) induced the expression of IL1β, IL6, IL8, and CCL2 up to 5-fold in PBMCs from select donors

(Fig. 2b). Since IL1β, IL6, IL8, and CCL2 are among the key cytokines elevated in

patients with severe COVID-19,[18] it is conceivable that secreted ORF8 is the viral factor

responsible for inflammatory cytokine response in patients with severe COVID-19.

To evaluate whether the glycosylation of ORF8 affected cytokine induction, we

purified ORF8 from the HEK293 culture supernatant and from ORF8-expressing *E. coli*

and designated them as ORF8-glyco^{hi} and ORF8-glyco^{null}, respectively. As shown in Fig.

2c that IL1β, IL-6, and IL-8 but not IL-2 were robustly induced by the pure ORF8-glyco^{hi}

in a dose-dependent manner confirming that secreted ORF8 is capable of inducing

proinflammatory cytokines seen in COVID-19 patients. Interestingly, *E. coli* expressed

ORF8-glyco^{null}, had no cytokine induction activity even at higher doses, suggesting that

proper ORF8 glycosylation is necessary for cytokine induction.

We then tested the cytokine induction activities of ORF8 from HEK293 cells

treated with the Golgi inhibitor cocktail Brefeldin-A and Monensin (BFA/M) and found

that while the glycosylation level was reduced (Supplemental Fig. 3) the cytokine

induction activity of BFA/M-treated ORF8 was indeed altered (actually increased),

suggesting that the cytokine induction activity of ORF8 is governed by the Golgi

complex. Our protein localization data (Supplemental Fig 1, 2) indicate that several

SARS-CoV-2 proteins are localized to Golgi membranes and may potentially modify the

glycosylation and the cytokine induction activity of ORF8. Indeed, co-expressing MEM,

ENV, ORF3A, or ORF7A was able to alter the molecule weight and the cytokine-

inducing activities of ORF8 (Supplemental Fig 4).

ORF8 stimulates CD14^{+} monocytes to produce pro-inflammatory cytokines/chemokines.

Next, we asked which specific cell subsets in PBMCs are the primary targets of

ORF8, and what is the resulting transcriptional signatures in those cells. We performed

single-cell RNA sequencing (scRNA-Seq) analysis on three PBMC samples treated with

ORF8-CM or control-CM. Single-cell transcriptome-based cell clustering showed that

cells in cluster 5 (C5) on the t-SNE plot (Fig. 3a) were induced to express cytokines

including IL1β, IL8, and CCL2 and inflammasome pathway components upon ORF8

treatment (Fig. 3b) (Supplemental Table 2). The same cytokines are known to be

elevated in COVID-19 patients.[19] The expression of CD14, CD16, CD68, and HLA-DR

identifies those cytokine secreting cells as activated monocytes.[20,21]

To identify which monocyte subsets are the targets of ORF8, we analyzed

intracellular cytokines in ORF8 stimulated PBMCs from 15 donors by flow cytometry.

Significant IL1β expression was detected in classical (CD14$^+$/CD16$^-$) and intermediate

(CD14$^+$/CD16$^+$) subsets but not non-classical (CD14^{low}/CD16$^+$) monocytes nor cells of

other lineages (B or T-lymphocytes, dendritic, NK cells) in all 15 donors (Fig. 3c).

Similar results were observed for IL8 and CCL2 expression as well (Fig. 5a). Our data

demonstrate that the CD14$^+$ monocyte subsets are the producer of the pro-inflammatory

cytokines upon ORF8 stimulation.

203
Secreted ORF8 directly binds to and activates the NLRP3-mediated inflammasome response

To decipher the mechanism by which ORF8 induces pro-inflammatory cytokines

in CD14+ monocytes, using the Gene Set Enrichment Analysis (GSEA) and Kyoto

Encyclopedia of Gene and Genome (KEGG) tools, we found that the SARS-CoV-2

infection pathway, NOD-like receptor signaling pathway, and NFκB pathway were

among the top enriched, especially, mRNAs of lysosomal enzymes (LYZ, CTSD, CTSB,

CTSS), inflammasomal protein NLPR3, and cytokines IL1β, IL8, CCL2 were among the

top expressed (Fig. 3b).

We then examined the role of the NLRP3-mediated inflammasome pathway in

ORF8-mediated cytokine production. Fig. 3d shows that both NLPR3 and IL1β proteins

were induced in PBMCs upon ORF8 treatment, and such induction was diminished in

NLPR3 knockdown cells (THP1-defNLRP3, InvivoGen) compared to the parental THP-

1 cells, demonstrating the NLRP3 dependency of ORF8 mediated cytokine response in

human monocytes (Fig. 3e).

To determine how ORF8 molecules enter CD14⁺ monocytes, we first incubated

PBMCs with pure ORF8 protein on ice for one hour to saturate any potential ORF8

binding receptors on the monocyte surface. After extensive washes with ice-cold PBS to

remove unbound ORF8 protein, the cells were then incubated at 37°C to initiate ORF8-

mediated cell activation in the presence or absence of additional ORF8. Fig. 4a shows

robust IL1β mRNA expression was observed only in cells exposed to additional ORF8

but not those only pre-incubated with ORF8, suggesting that ORF8 likely enters

monocytes through a non-receptor-mediated process, such as phagocytosis.

Complementarily, we asked if ORF8 would bind to TLR2, TLR4, CD14, or NLRP3

known to be involved in inflammasome activation. We incubated ORF8 protein-coated

beads with monocyte lysates from two ORF8 responding and two ORF8 nonresponding

healthy donors. Fig. 4b shows that neither TLR2, TLR4, nor CD14 were co-precipitated

with ORF8 (CD14 data not shown). However, NLRP3 was readily detected in one of the

ORF8-responders, suggesting that indeed ORF8 binds to NLRP3 in primary human

monocytes.

To further dissect how NLRP3 binds to ORF8, we transiently transfected ORF8-

expressing HEK293 cells with Flag-tagged NLRP3 constructs Flag-N3-FL (full-length

NLRP3), or one of the three deletion mutants Flag-N3-N, Flag-N3-M, or Flag-N3-C (Fig.

4c).[22] Cell lysates were then incubated with Strep-Tactin™ beads to pulldown Strep-

tagged ORF8 and its binding proteins. In addition to NLRP3-FL (Fig. 4d), both Flag-N3-

M, and Flag-N3-C but not Flag-N3-N also strongly co-precipitated with ORF8

suggesting both the middle NACHT domain and C-terminal LRR domain of NLRP3

each can independently bind to ORF8. These results clearly demonstrate inflammasome

protein NLRP3 directly interacts with ORF8.

ORF8 activates the NLRP3-mediated inflammasome pathways.

247   The effect of LPS in inflammasomal activation has been well established;

therefore, it was important to demonstrate that our purified ORF8 protein was free of

LPS contamination. ORF8 and LPS were compared for their effects on IL1β, IL8, and

CCL2 induction in three monocyte subsets (classical, intermediate, and non-classical)

from 15 healthy donors. Fig 5a shows that all three subsets responded to LPS while only

CD14$^+$ (classical and intermediate) responded to ORF8 with variable induction of IL1β,

IL8, and CCL2. These results show that ORF8 and LPS target monocytes with different

cell type specificities. We then examined the responsiveness of CD14$^+$ monocytes to LPS

or ORF8 treatment using IL1β, IL8, and CCL2 expression as the readout, and we found

that intracellular IL1β was detected in the majority (90.1%) of LPS-treated CD14$^+$

monocytes but only in a subpopulation (mean=38%) of ORF8 treated CD14$^+$ monocytes.

In contrast, CCL2 expression was detected only in a minor population (mean=22.1%) of

LPS-treated CD14$^+$ monocytes but in a larger population of ORF8-treated CD14+

monocytes (mean=63.7%) (Fig. 5b). These data clearly demonstrate that ORF8 and LPS

also have different cytokine induction specificities. In addition, we also examined the

cell proliferation and the IgM induction of mouse splenic B cells upon ORF8 and LPS

treatment, and we found that the mouse splenic B cells did not proliferate, nor

produced any polyclonal IgM 96 hours after ORF8 treatment, while LPS treated cells

showed robust proliferation morphology and massive production of IgM (Fig. 5c). In

the aggregate, our data clearly demonstrated that the cytokine inducing activity of

ORF8 is not due to LPS contamination.

To examine the activation status of the canonical NFκB pathway necessary for

the induction of inflammasomal proteins including NLPR3, Casp-1, and IL1β in ORF8

treated monocytes, we examined the GSEA analysis results and found that NFκB

pathway components were transcriptionally enriched (NES=1.539662, FDR q-value=0)

in ORF8 treated monocytes in cluster 5 (Fig. 6a, left panel). To validate these at the

protein level, we treated THP-1 cells with ORF8 for 4 and 24 hours and determined

their NFκB pathway protein expression. Fig. 6a (right panels) shows that the increased

level of phospho-p65 and the conversion of p100 to p52 were readily detected 4 hours

after ORF8 treatment. Similar results were also observed in LPS-treated cells. These data

demonstrate that ORF8 stimulation activates the NFκB pathway triggering the

activation of the inflammasome pathway in human monocytes.

Similarly, the GSEA analysis data revealed the mRNAs of inflammasome

pathway components were also significantly enriched (NES=1.812154, FDR q-value

=0.015483) (Fig. 6b, left panel). We then treated THP-1 cells with ORF8 or LPS alone or

in combination with nigericin, a potent agent that provides a second signal for the

canonical inflammasomal pathway activation, for 4 or 24 hours. The cell lysates and

culture supernatants were analyzed by Western blotting for the expression of NLPR3,

Caspase-1, Caspase-5, IL1β, and the pore-forming fragment of gasdermin D (GSDMD)

proteins. We found that ORF8 induces the inflammasome pathway by 1) activating the

production of IL1β without requiring a separate priming step; 2) signaling through the

non-canonical pathway using Casp-1 (early hours) followed by switching to Casp-4/5;

3) maintaining cell viability without triggering pyroptosis unless the second signal such

as nigericin is present which rapidly initiates pyroptosis (Fig. 6b, right panel). Our data

suggest that ORF8 and LPS may represent a family of pyrogens by triggering a non-

canonical inflammasomal pathway leading to the production and release of IL1β

without activating GSDMD to the level necessary for pyroptosis, a process called

hyperactivation for LPS.[23,24] However, ORF8 can also serve as a priming agent for the

canonical inflammasome pathway to evoke a prompt pyroptosis in the presence of

second signal molecules such as extracellular ATP from damaged cells.

To validate the dual roles of ORF8 in the activation of canonical and non-

canonical inflammasome pathways, we examined ASC speck formation in ORF8- or

LPS-treated THP-1 cells by immunofluorescence. As shown in Fig. 6c, ORF8 alone was

insufficient to induce ASC speck formation. However, distinct pre-nuclear ASC specks

were readily detected when nigericin or extracellular ATP was also present. In addition,

the NLPR3 protein was also co-localized with ASC protein to the specks (Fig. 6c), ASC

speck formation always precedes cell pyroptosis in our system, and ORF8 is slightly

more robust than LPS in mediating ASC formation in the presence of nigericin or

extracellular ATP. The results confirm the dual roles of ORF8 in the activation of the

non-canonical inflammasome pathway when the second signal is absent, and the

canonical inflammasome pathway by priming the cells when the second signal is

present. These data further suggest that the mode of action of ORF8 is dependent on the

presence or absence of a second signal such as extracellular ATP released from

damaged cells/tissues nearby.

Blood ORF8 protein levels correlate with the mortality and disease course of severe patients with

COVID-19

Having demonstrated that ORF8 is secreted as a glycoprotein and capable of

inducing inflammatory cytokine responses _in vitro_, we then asked if ORF8 is also

secreted and glycosylated in blood from patients infected with SARS-CoV-2. By

analyzing serum samples collected from patients within 4-7 days of COVID-19

diagnosis using Western blotting, we found that ORF8 protein was readily detected at

various levels in 92% (23/25) of newly infected patients. Interestingly, two other major

secreted proteins NSP9 and NSP10 were not detected in the same samples suggesting

not all the secreted SARS-CoV-2 proteins are present at detectable levels in blood

circulation. To determine the glycosylation status of the serum ORF8, we compared the

sizes of ORF8 from patients and HEK293 lysate. Our results (Fig. 7A) show that patient

serum ORF8 is indeed glycosylated when compared to the ORF8 from HEK293 lysate

which has an extra Strep-II tag (28 aa) and possibly an uncleaved signal sequence (16

aa). Due to the lack of a quantitative assay, we were unable to determine the absolute

quantity of ORF8 in patient samples; however, our semi-quantitative Western blot

could readily detect ORF8 in as little as four microliters of patient serum suggesting the

protein is present at significant levels in patient blood at the onset of COVID-19.

Given its activity in inducing inflammatory cytokines _in vitro_, and its significant

presence in the blood of newly infected patients, we then asked if the levels of ORF8

protein would correlate with disease severity and outcome. To that end, we examined

the correlation between blood ORF8 levels and patients' survival outcomes in our

cohort of 25 hospitalized patients. Fig. 7b shows that after 120 days of follow-up, all

seven fatalities were exclusively associated with the ORF8-high group (scored 2+ or 3+

on Western blot) while all patients in the ORF8-low group (scored 0 or 1+) had mild

symptoms and quick recovery without any events of death. These results demonstrate a

clear correlation between the ORF8 load and disease severity in newly infected patients.

We then monitored the ORF8 levels at various time points during the disease course in

two patients with prolonged COVID-19, and one patient with long covid lasting more

than 17 months. As demonstrated in Fig. 7c, two patients with high blood ORF8 levels

had persistent severe symptoms requiring treatment in the intensive care unit (ICU),

and their blood ORF8 levels diminished by the time they were recovered and

discharged from the hospitals. The long covid patient with lingering symptoms

throughout remained ORF8 positive in all three serial blood samples collected over 16-

month period (Fig. 7c). These results suggest that ORF8 level is prognostic for COVID-

19 outcome, further supporting our hypothesis that the ORF8 protein is a pathogenic

cause of severe COVID-19 in patients.

ORF8-mediated cytokine induction is targetable by select NLRP3 inhibitors

Given its causative role in COVID-19 pathogenesis, we then tested the

targetability of ORF8 mediated cytokine response using three investigational NLRP3

inhibitors - MCC950, Tranilast, and OLT117 (Dapansutrile). As shown in Fig. 7d, the

NLRP3 inhibitor MCC950 showed effective inhibition on the production of IL1β in both

THP-1 cells and PBMCs (blue bars), while Tranilast and OLT117 (hatched bars)

exhibited moderate to marginal effect in PBMCs but no effect in THP-1 cells. Our results

suggest that ORF8 mediated IL1β production can be effectively inhibited by the NLRP3

inhibitor MCC950 at a nanomolar dose range, and possibly by other classes of inhibitors

as well, demonstrating that targeting the ORF8/NLRP3 axis is a promising strategy for

treating patients with symptomatic COVID-19.

DISCUSSION

Despite the success in the development and implementation of effective vaccines to prevent SARS-CoV-2 infection, COVID-19 remains a major challenge for many reasons. These include 1) a significant population remains unvaccinated, 2) new variants keep emerging and evading current vaccine protection, 3) patients with compromised immunity including those with blood cancers on immunosuppressive medications are poorly protected, [25,26] and more importantly, 4) many previously infected patients have developed lingering symptoms or long Covid. Therefore, better therapies and clinical management tools for COVID-19 are urgently needed.

SARS-CoV-2 accessory protein ORF8, a 121 amino acid protein, is the least conserved protein in the beta-coronavirus family.[27,28] Here we report that ORF8 is a major secreted viral glycoprotein *in vitro* and in COVID-19 patients. Our data, summarized in Supplemental Fig. 7, support the concept that the ORF8 protein secreted from locally infected cells in the lung and released to the bloodstream then stimulates circulating CD14$^+$ monocytes to initiate systematic cytokine responses. This is consistent with findings by others that monocytes are the single most affected WBC subset in the blood by SARS-CoV-2 infection.[29] It has been reported that about 6% of monocytes were infected with the virus in COVID-19 patients through CD16-mediated uptake of antibody-opsonized SARS-CoV-2 virus. Whether these cells were truly infected or simply engulfed with infected cells/debris remains to be seen since no live virus was detected.[30] Nevertheless, these monocytes may represent a parallel and complementary process to the ORF8-mediated inflammatory pathway described herein. Our intracellular IL1β data showed a mean of 43% (range, 7.2% - 90.7%) of monocytes that are ORF8 responsive, suggesting that the ORF8-mediated process dominates. We

conclude that secreted ORF8 is a key disease-causing viral factor and can be targeted by

NLRP3 inhibitors, offering a potential new treatment option.

NLRP3-mediated inflammasome activation is a critical part of the innate immune

response. However, the precise mechanisms for different pathogens have not been fully

delineated. Our study indicated that ORF8 may enter monocytes through a non-

receptor mediated ORF8 internalization, followed by lysosomal action involving

lysozyme (LYZ) and cathepsin proteases (CTSB, CTSD). ORF8 then may bind to the

NACHT and/or LRR domain to activate NLRP3 and recruit Casp-1 (early) and Casp-4/5

(later). It is not known what the roles of these two different bindings play, and if they

are important in pathway choosing (canonical vs. non-canonical). It is also interesting

that, like LPS, ORF8 induces the expression/activation of Caspase-1 early (hour 4) but

switches to Casp-4/5 later (at hour 24) even when nigericin is present (data not shown).

Our data shows that the mode of action of ORF8 is readily changed by the presence of

the second signal leading to the switch from non-canonical to canonical inflammasomal

response. Therefore, it is conceivable that, in patients with COVID-19, the inflammation

may be aggravated by the additional tissue/cell damages that release ATP, leading to

the activation of an ORF8-mediated canonical inflammasome pathway and causing

pyroptosis.

Due to the lack of effective COVID-19 animal models, it is challenging to

genetically test the pathogenic role and targetability of the ORF8 protein *in vivo*.

However, a SARS-CoV-2 variant (Δ382) found in Taiwan and Singapore with a

complete loss of the ORF8 gene serves as a tailor-made natural genetic model that

validates our findings.[31,32] Young et al. analyzed a cohort of 92 patients infected with the

wildtype virus and 29 patients infected with the Δ382 variant and found that patients

infected with the wildtype SARS-CoV-2 exhibited much higher levels of pro-

inflammatory cytokines than those infected with the Δ382 variant. Furthermore, no

patients with the Δ382 variant required supplemental oxygen whereas it was required

for 26 (28%) patients infected with wild-type virus. These observations provide yet

another line of evidence supporting the link between ORF8 and COVID-19 disease

severity, and an *in vivo* genetic proof-of-principle for targeting ORF8 to prevent severe

COVID-19 infection. Additionally, various clinical studies and trials have shown that

the IL1 receptor antagonist anakinra significantly reduced the mortality risk in

hospitalized patients with moderate to severe inflammation symptoms.[33] [34,35] The data

from our present study suggests that targeting NLRP3 or ORF8 upstream of the IL1

receptor would provide greater therapeutic benefit to patients with severe COVID-19

symptoms, and could also provide prophylactic benefit to patients at risk of developing

severe COVID-19 infection.

ACKNOWLEDGEMENT

We thank The Mayo Clinic Genome Analysis Core, Bioinformatics Core, Microscopy and Cell Analysis Core, and the CMM Electron Microscopy Facility of the University of California San Diego (supported by NIH equipment grant 1S10OD023527-01) for their expert service, Mayo Clinical Blood Bank for providing fresh leukocyte cones of healthy donor blood, and the Mayo Clinic Center for Individualized Medicine for providing access to their biobank data. We also thank the University of Iowa/Mayo Clinic Lymphoma SPORE (NIH/NCI grant P50 CA97274) and the Predolin Foundation Biobank for sharing some of their resources for this project.

AUTHOR CONTRIBUTION

XW and TEW designed the experiments, analyzed, and interpreted results, and wrote the manuscript; XW, MKM, GJR, KEN, KAG, XT, TLW, VT, AW, and MJS conducted experiments and analyzed data; XW, JPA, YY, XT, SD, and MJS analyzed data and generated figures; ADB, JPA, GR, JP, SMA, VB, HD, and TEW provided experiment samples; JPA, GJR, JP, TLW, SMA, MJS discussed and interpreted results; all authors read and approved the manuscript.

CONFLICT-OF-INTEREST DISCLOSURE

The authors declare no competing financial interests.

 FIGURE LEGEND

Figure 1. Secretion property of SARS-CoV-2 proteins. a. Western blot image of cell

lysates for validating SARS-CoV-2 protein expression in HEK293 cells. The blots were

probed with an anti-Strep II tag antibody and anti-HA (for SPK protein). **b.** Western

image of SARS-CoV-2 protein pulldown using StrepTactin™ beads and anti-HA-beads

(for SPK) from the culture supernatants for validating SARS-CoV-2 protein secretion

from HEK293 cells. The blots were also probed with an anti-Strep II tag antibody and

anti-HA (for SPK protein).

Figure 2. Secreted ORF8 specifically induces the expression of pro-inflammatory

cytokines and is glycosylation dependent. a. Conditioned media containing major

secreted NSP proteins did not induce pro-inflammatory cytokines in PBMCs of healthy

donors; **b**. Conditioned media containing secreted ORF8 specifically induced pro-

inflammatory cytokines (highlighted in yellow) IL1β, IL6, IL8, and CCL2 in PBMCs

from unselected healthy donors. **c.** Purified highly glycosylated ORF8 (ORF8-glycol^{hi})

from HEK293 but not unglycosylated ORF8 (ORF8-glycol^{null}) from *E. coli* stimulates

PBMCs to produce pro-inflammatory cytokines IL1β, IL6, IL8 but not T cell cytokine

IL2.

Figure 3. Single-cell RNA sequencing of ORF8 treated PBMCs. a. tSNE maps of cell

clusters using concatenated scRNA-Seq data from two ORF8-responsive PBMC

samples. Cells in cluster 5 were responsive to ORF8 treatment. **b.** Violin plots show the

monocyte markers CD14, CD16, CD68, and HLA-DRB1, key inflammasome pathway

components LYZ, CTSB, NLRP3, and IL1β, and pro-inflammatory cytokines IL1β, IL8,

and CCL2 mRNA expression in cells of cluster 5. **c.** Flow cytometry analysis of

intracellular IL1β protein expression in different PBMC subsets in response to ORF8

treatment. The plots show that IL1β was exclusively produced by CD14$^+$CD16$^-$

(classical) and CD14$^+$CD16$^+$ (intermediate) monocyte subsets. **d.** Western blot images

validate that the induction of NLRP3 and IL1β in PBMCs upon stimulation with ORF8

indeed occurs at the protein level. **e.** Western images showing in response to ORF8

treatment, the IL1β induction in parental THP-1 cells became diminished in NLRP3

knockdown (NLRP3^{kd}) THP-1 cells. The data shown are representative of at least two

independent experiments except for the scRNA-Seq.

Figure 4. ORF8 induces proinflammatory cytokines through direct binding to NLRP3.

a. ORF8 induces IL1β production through a non-surface-receptor-mediated process.

PBMCs were pre-incubated on ice with ORF8 (500 ng/ml) for 2 hours to allow ORF8 to

bind to its "surface receptors" followed by washing and incubating at 37ºC to activate

the cytokine production in the presence or absence of additional ORF8 (200 ng/ml). IL1β

expression was measured by qPCR. **b.** ORF8 directly binds to NLRP3 but not TLR2 or

TRL4 in primary monocytes shown by affinity pulldown assay. **c.** Schematic drawing

of NLRP3 deletion constructs for mapping ORF8 binding domains in NLPR3. **d.**

Mapping NLRP3 domains that bind to ORF8 by affinity pulldown assay using ORF8-

Strep-Tacin® beads. Western images of total cell lysates (left half) were used as

expression controls and Western images of precipitated proteins (right half) showing

ORF8 efficiently binds the full-length as well as two NLRP3 deletion mutant proteins.

Figure 5. ORF8 is functionally different from LPS. a. Representative flow data

showing LPS could activate multiple cytokine responses in all three monocyte subsets

of PMBCs from all 15 healthy donors examined while ORF8 only activates CD14$^+$

Monocytes but not CD14^{low}/CD16^{++} non-classical monocytes (red boxed) suggesting LPS

and ORF8 may target monocytes through different mechanisms. **b.** Intracellular

cytokine flow cytometry data summary on 15 healthy donors upon treatment with

ORF8 or LPS, suggesting that ORF8 and LPS are different in cytokine induction

specificities. **c.** Mouse B cells responded differently to the treatment of ORF8 and LPS as

measured by IgM production and the morphological features of cell proliferation status,

demonstrating the lack of any detectable contaminating LPS activity in purified ORF8

protein.

Figure 6. ORF8 induces proinflammatory cytokines through activation of NLRP3-

mediated inflammasome pathways. a. Gene enrichment analysis of scRNA-Seq (left

panel) and Western confirmation of key NFkB pathway molecules (right panels) upon

stimulation of ORF8 for 4 or 24 hours. **b**. Gene enrichment analysis of scRNA-Seq (left

panel) and Western confirmation of NLRP3, IL1β, Casp-1, and Casp-5 in THP-1 cell

lysates and IL1β and GSDMD in the culture supernatants (right panels) upon

stimulation of ORF8 for 4 or 24 hours. **c.** Immunofluorescence images show that, LPS

and ORF8 activate ASC speck (arrowhead pointed) formation only when a second

signal (nigericin or extracellular ATP) was present. Cells were first treated with 200

ng/ml ORF8 or 100 ng/ml LPS for 2.5 hours followed by 90 min treatment with 5 μM

Nigericin or 5 μM ATP before cell harvest and staining for immunofluorescence.

Figure 7. The levels of ORF8 protein in sera of COVID-19 patients correlate with the

outcome and disease course, and the ORF8 pathway is targetable with NLRP3

inhibitors. a. Representative Western blot images showing ORF8 protein but not NSP9

or NSP10 were detectable in as little as 4.0 μl of serum samples from newly infected

COVID-19 patients. ORF8 protein in patient sera shown as glycosylated by size

estimation; **b.** Kaplan-Meier curve showing fatality in hospital patients is associated

with higher serum level of ORF8; **c.** Western images showing ORF8 levels at different

time points correlate with disease course in two patients with prolonged clinical

courses, and in one patient with long-covid. **d.** Targetability of ORF8 mediated IL1β

expression with various NLRP3 inhibitors in THP1 cells and human primary PBMCs.

The data shown are representative of two or three independent experiments.

Abbreviations: Dx = diagnosis; ED = emergency department; ICU = intensive care unit

REFERENCE

1.   Johns Hopkins University Coronavirus Resource Center (2021) Retrived (June 20, 2021) from: https://coronavirus.jhu.edu/map.html.

2.   Arunachalam PS, Wimmers F, Mok CKP, et al. Systems biological assessment of immunity to mild versus severe COVID-19 infection in humans. *Science*. 2020;369(6508):1210-1220.

3.   Schultze JL, Aschenbrenner AC. COVID-19 and the human innate immune system. *Cell*. 2021;184(7):1671-1692.

4.   Chen X, Zhao B, Qu Y, et al. Detectable Serum Severe Acute Respiratory Syndrome Coronavirus 2 Viral Load (RNAemia) Is Closely Correlated With Drastically Elevated Interleukin 6 Level in Critically Ill Patients With Coronavirus Disease 2019. *Clin Infect Dis*. 2020;71(8):1937-1942.

5.   Fajnzylber J, Regan J, Coxen K, et al. SARS-CoV-2 viral load is associated with increased disease severity and mortality. *Nat Commun*. 2020;11(1):5493.

6.   Lucas C, Wong P, Klein J, et al. Longitudinal analyses reveal immunological misfiring in severe COVID-19. *Nature*. 2020;584(7821):463-469.

7.   Chen P, Nirula A, Heller B, et al. SARS-CoV-2 Neutralizing Antibody LY-CoV555 in Outpatients with Covid-19. *N Engl J Med*. 2021;384(3):229-237.

8.   Weinreich DM, Sivapalasingam S, Norton T, et al. REGN-COV2, a Neutralizing Antibody Cocktail, in Outpatients with Covid-19. *N Engl J Med*. 2021;384(3):238-251.

9.   Qi F, Qian S, Zhang S, Zhang Z. Single cell RNA sequencing of 13 human tissues identify cell types and receptors of human coronaviruses. *Biochem Biophys Res Commun*. 2020;526(1):135-140.

10.   Zou X, Chen K, Zou J, Han P, Hao J, Han Z. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med*. 2020;14(2):185-192.

11.   Rubin R. COVID-19 and Blood Donation. *JAMA*. 2022;327(7):615.

12.   Andersson MI, Arancibia-Carcamo CV, Auckland K, et al. SARS-CoV-2 RNA detected in blood products from patients with COVID-19 is not associated with infectious virus. *Wellcome Open Res*. 2020;5:181.

13.   Cappy P, Candotti D, Sauvage V, et al. No evidence of SARS-CoV-2 transfusion transmission despite RNA detection in blood donors showing symptoms after donation. *Blood*. 2020;136(16):1888-1891.

14.   Owusu M, Sylverken AA, El-Duah P, et al. Low risk of SARS-CoV-2 in blood
transfusion. *PLoS One*. 2021;16(4):e0249069.

15.   Mariano G, Farthing RJ, Lale-Farjat SLM, Bergeron JRC. Structural Characterization of
SARS-CoV-2: Where We Are, and Where We Need to Be. *Front Mol Biosci*. 2020;7:605236.

16.   Gordon DE, Jang GM, Bouhaddou M, et al. A SARS-CoV-2 protein interaction map
reveals targets for drug repurposing. *Nature*. 2020;583(7816):459-468.

17.   Gordon DE, Hiatt J, Bouhaddou M, et al. Comparative host-coronavirus protein
interaction networks reveal pan-viral disease mechanisms. *Science*. 2020;370(6521).

18.   Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel
coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506.

19.   Jin K, Bardes EE, Mitelpunkt A, et al. Implicating Gene and Cell Networks Responsible
for Differential COVID-19 Host Responses via an Interactive Single Cell Web Portal. *bioRxiv*.
2021.

20.   Lambert C, Preijers F, Yanikkaya Demirel G, Sack U. Monocytes and macrophages in
flow: an ESCCA initiative on advanced analyses of monocyte lineage using flow cytometry.
Cytometry B Clin Cytom. 2017;92(3):180-188.

21.   Kapellos TS, Bonaguro L, Gemund I, et al. Human Monocyte Subsets and Phenotypes in
Major Chronic Inflammatory Diseases. *Front Immunol*. 2019;10:2035.

22.   Shi H, Wang Y, Li X, et al. NLRP3 activation and mitosis are mutually exclusive events
coordinated by NEK7, a new inflammasome component. *Nat Immunol*. 2016;17(3):250-258.

23.   Zanoni I, Tan Y, Di Gioia M, Springstead JR, Kagan JC. By Capturing Inflammatory
Lipids Released from Dying Cells, the Receptor CD14 Induces Inflammasome-Dependent
Phagocyte Hyperactivation. *Immunity*. 2017;47(4):697-709 e693.

24.   Evavold CL, Ruan J, Tan Y, Xia S, Wu H, Kagan JC. The Pore-Forming Protein
Gasdermin D Regulates Interleukin-1 Secretion from Living Macrophages. *Immunity*.
2018;48(1):35-44 e36.

25.   Roeker LE, Knorr DA, Thompson MC, et al. COVID-19 vaccine efficacy in patients with
chronic lymphocytic leukemia. *Leukemia*. 2021.

26.   Greenberger LM, Saltzman LA, Senefeld JW, Johnson PW, DeGennaro LJ, Nichols GL.
Antibody response to SARS-CoV-2 vaccines in patients with hematologic malignancies. *Cancer
Cell*. 2021;39(8):1031-1033.

27.   Ceraolo C, Giorgi FM. Genomic variance of the 2019-nCoV coronavirus. *J Med Virol*.
2020;92(5):522-528.

28.   Tan Y, Schneider T, Leong M, Aravind L, Zhang D. Novel Immunoglobulin Domain
Proteins Provide Insights into Evolution and Pathogenesis of SARS-CoV-2-Related Viruses.
mBio. 2020;11(3).

29.   Notarbartolo S, Ranzani V, Bandera A, et al. Integrated longitudinal immunophenotypic,
transcriptional and repertoire analyses delineate immune responses in COVID-19 patients. *Sci
Immunol*. 2021;6(62).

30.   Junqueira C, Crespo A, Ranjbar S, et al. FcgammaR-mediated SARS-CoV-2 infection of
monocytes activates inflammation. *Nature*. 2022.

31.   Gong YN, Tsao KC, Hsiao MJ, et al. SARS-CoV-2 genomic surveillance in Taiwan
revealed novel ORF8-deletion mutant and clade possibly associated with infections in Middle
East. *Emerg Microbes Infect*. 2020;9(1):1457-1466.

32.   Young BE, Fong SW, Chan YH, et al. Effects of a major deletion in the SARS-CoV-2
genome on the severity of infection and the inflammatory response: an observational cohort
study. *Lancet*. 2020;396(10251):603-611.

33.   Caricchio R, Abbate A, Gordeev I, et al. Effect of Canakinumab vs Placebo on Survival
Without Invasive Mechanical Ventilation in Patients Hospitalized With Severe COVID-19: A
Randomized Clinical Trial. *JAMA*. 2021;326(3):230-239.

34.   Kyriazopoulou E, Poulakou G, Milionis H, et al. Early treatment of COVID-19 with
anakinra guided by soluble urokinase plasminogen receptor plasma levels: a double-blind,
randomized controlled phase 3 trial. *Nat Med*. 2021;27(10):1752-1760.

35.   Kyriazopoulou E, Huet T, Cavalli G, et al. Effect of anakinra on mortality in patients
with COVID-19: a systematic review and patient-level meta-analysis. *Lancet Rheumatol*.
2021;3(10):e690-e697.
626

Figure 1. Secretion property of SARS-CoV-2 proteins.

Figure 2. Secreted ORF8 specifically induces the expression of pro-inflammatory cytokines is glycosylation dependent.

Figure 3. Single cell RNA sequencing of ORF8 treated PBMCs.

Figure 4. ORF8 induces proinflammatory cytokines through direct binding to NLRP3.

Figure 5. ORF8 is functionally different from LPS.

Figure 6. ORF8 induces proinflammatory cytokines through activation of NLRP3 mediated inflammasome pathways.

Figure 7. The levels of ORF8 protein in sera of Covid-19 patients correlates with the outcome and disease course, and the ORF8 pathway is targetable with NLRP3 inhibitors.