A chromosomal inversion may facilitate adaptation despite periodic gene flow in a

freshwater fish

Matt J. Thorstensen[1], Peter T. Euclide[2], Jennifer D. Jeffrey[1,3], Yue Shi[2,4], Jason R. Treberg[1],

Douglas A. Watkinson[5], Eva C. Enders[5], Wesley A. Larson[2,6], Yasuhiro Kobayashi[7,8], and Ken

M. Jeffries[1]

[1] Department of Biological Sciences, University of Manitoba, Winnipeg, MB R3T 2N2, Canada

[2] U.S. Geological Survey, Wisconsin Cooperative Fishery Research Unit, College of Natural

Resources, University of Wisconsin-Stevens Point, Stevens Point, WI 54481-3897, USA

[3] Department of Biology, University of Winnipeg, Richardson College, 2RC047, 515 Portage

Avenue, Winnipeg, Manitoba, R3B 2E9, Canada

[4] College of Fisheries and Ocean Sciences, University of Alaska Fairbanks, 17101 Point Lena

Loop Road, Juneau, AK 99801, USA

[5] Freshwater Institute, Fisheries and Oceans Canada, Winnipeg, MB R3T 2N6, Canada

[6] National Oceanographic and Atmospheric Administration, National Marine Fisheries Service,

Alaska Fisheries Science Center, Auke Bay Laboratories, 17109 Point Lena Loop Road, Juneau,

AK 99801, USA

[7] Department of Biological Sciences, Fort Hays State University, 600 Park Street, Hays, KS

67601, USA

[8] Department of Biology, The College of St. Scholastica, 1200 Kenwood Avenue, Duluth, MN

55811, USA

Abstract

Differences in genomic architecture between populations, such as chromosomal inversions, may play an important role in facilitating adaptation despite opportunities for gene flow. One system where chromosomal inversions may be important for eco-evolutionary dynamics are in freshwater fishes, which often live in heterogenous environments characterized by varying levels of connectivity and varying opportunities for gene flow. In the present study, reduced representation sequencing was used to study possible adaptation in $n = 345$ walleye (*Sander vitreus*) from three North American waterbodies: Cedar Bluff Reservoir (Kansas, USA), Lake Manitoba (Manitoba, Canada), and Lake Winnipeg (Manitoba, Canada). Haplotype and outlier-based tests revealed a putative chromosomal inversion that contained three expressed genes and was nearly fixed for alternate genotypes in each Canadian lake. These patterns exist despite several opportunities for gene flow between these proximate Canadian lakes, suggesting that the inversion may be important for facilitating adaptive divergence between the two lakes despite gene flow. Our study illuminates the importance of genomic architecture for facilitating local adaptation in freshwater fishes. Furthermore, our results provide additional evidence that inversions may facilitate local adaptation in many organisms that inhabit connected but heterogenous environments.

Keywords

selection, haplotype, fishery, genomic architecture, adaptive variation, walleye, *Sander vitreus*

Introduction

Differences in genomic architecture genomic architecture such as structural variants may help to facilitate local adaptation, and are therefore key mechanisms of eco-evolutionary processes (Dorant et al., 2020; Shi et al., 2021; Tigano & Friesen, 2016; Wellenreuther & Bernatchez, 2018). One type of structural variant, chromosomal inversions, are taxonomically widespread reversed regions of DNA that have been linked to both adaptation and speciation (reviewed in Wellenreuther & Bernatchez, 2018). Inversions can effectively protect large chromosomal regions containing hundreds of genes from recombination, facilitating local adaptation even in the face of high gene flow (Wellenreuther & Bernatchez, 2018). Most inversions found in previous studies have been relatively large because commonly used reduced representation sequencing methods can more easily detect large inversions (Wellenreuther & Bernatchez, 2018). However, inversions of any size may be important. According to a simulation study, intermediate-to-large inversions have been linked to local adaptation, but small inversions were preferentially fixed in directly beneficial mutation models and contributed the most to genome evolution in neutral and underdominant scenarios (Connallon & Olito, 2021). Among fishes, chromosomal inversions have largely been characterized in well studied groups of species such as representative salmonids (Arostegui et al., 2019; Leitwein et al., 2017; McKinney et al., 2020; Pearse et al., 2014) and gadids (Berg et al., 2017; Berg et al., 2016; Kirubakaran et al., 2016; Puncher et al., 2019; Sinclair-Waters et al., 2018; Sodeland et al., 2016), but little information exists on the importance of inversions in most taxa especially those inhabiting fresh water (Shi et al., 2021).

Freshwater fish often live in heterogenous environments characterized by varying levels of connectivity and therefore, varying opportunities for gene flow (Griffiths, 2015; Mushet et al.,

2019). Gene flow tends to work in contrast to local adaptation (Lenormand, 2002) but

nevertheless, evidence for local adaptation has been observed among different fish that use

freshwater habitats (see Fraser et al., 2011 for a review of salmonids; Shi et al., 2021). While

genomic architecture has been proposed to facilitate local adaptation (Tigano & Friesen, 2016),

few studies test for this possibility in freshwater fishes (Shi et al., 2021). Freshwater habitats

contribute a disproportionate richness in species diversity to global species diversity, given that

only 0.01% of the planet's total surface is fresh water (Balian et al., 2008). Fishes are a

significant proportion of overall freshwater species diversity (approx. 10% of overall freshwater

species and 70% of vertebrate freshwater species are fish; Balian et al., 2008). Therefore, an

understanding of the mechanisms of local adaptation in freshwater fishes can contribute

substantially to our overall understanding of local adaptation in heterogenous environments.

Walleye (*Sander vitreus*, but see Bruner, 2021) is a freshwater fish with a native range

from the Northwest Territories (Canada) to Alabama (USA) (Hartman, 2009) and support

important commercial and recreational fisheries (Fisheries and Oceans Canada, 2019, 2021). The

economic importance of walleye underscores the need for research supporting the sustainable

management of walleye fisheries, while its wide native distribution provides an opportunity to

study the genomic basis of adaptation to a variety of freshwater environments. In addition,

northern populations of walleye (such as those in Manitoba, Canada) were suspected to have

been recently isolated into smaller lakes following the drainage of glacial Lake Agassiz

approximately 7,000 years ago (Rempel & Smith, 1998; Stepien et al., 2009). This event may

have resulted in walleye populations that remain closely related at neutral markers but show

evidence of divergence at adaptive markers, such as in chromosomal inversions.

We sought to study how heterogenous environments with varying connectivity may have

shaped contemporary walleye populations, and the potential for genomic architecture to facilitate

adaptation despite opportunities for gene flow. Walleye from Lake Manitoba and Lake Winnipeg

in Manitoba (Canada) were analyzed and compared with an outgroup from Cedar Bluff

Reservoir in Kansas (USA) (Fig 1). Walleye in Lake Manitoba and Lake Winnipeg live in

largely separated waterbodies with a shared river—Lake Manitoba drains eastward into Lake

Winnipeg via the Fairford River, Lake St. Martin, and the Dauphin River (Fig 1). As a mobile

species that undergoes broad seasonal migrations, historic gene flow may have been extensive

between these Manitoba waterbodies (Backhouse-James & Docker, 2012; Munaweera

Arachchilage et al., 2021; Thorstensen et al., 2020; Turner et al., 2021). In addition, historical

flooding in 1882, 1902, 1904, 2011, and 2014 brought significantly increased waterflow

eastward from Lake Manitoba toward Lake Winnipeg (Ahmari et al., 2016), and such flooding

could have facilitated increased fish movement between the waterbodies. Last, walleye fry were

opportunistically stocked in Lake Winnipeg from Lake Manitoba (Manitoba, Canada) between

1917-2002 (Manitoba Government, 2020). While intensive stocking complicated analyses of

walleye ancestry in walleye of Minnesota and Wisconsin (USA) (Bootsma et al., 2020), stocking

of walleye fry, which likely experience significant mortality, had little to no effect in South

Dakota and Missouri (USA) (Fielder, 1992; Koppelman et al., 1992). Therefore, the genetic

impacts of fry stocking in this system are likely minimal.

Here, we employed Rapture (Ali et al., 2016), a reduced representation approach, to

genotype $n = 345$ individuals from the three waterbodies described above. Population structure

and demographic history were explored to provide context for signatures of selection.

Simulations with EASYPOP were used to estimate historical migration rates between Lake

Manitoba and Lake Winnipeg (Balloux, 2001). Candidate regions of adaptive variation were

identified with XP-EHH, and outlier tests were used to confirm haplotype-based results. In

addition, mRNA transcript expression in one candidate chromosomal inversion was assessed for

Lake Winnipeg walleye to show the potential for the functional significance of genomic

architecture in this system. The present study illuminates the importance of genomic architecture

for facilitating local adaptation in freshwater fishes, and provides additional evidence that

inversions may facilitate local adaptation in many organisms that inhabit connected but

heterogenous environments.

Methods

Sample Collection

Walleye anal fin clips from Lake Winnipeg individuals were collected by boat

electrofishing in Spring 2017 and 2018 from spawning sites in the Red River and possible

spawning sites near Matheson Island and in the Dauphin River, representing the south basin,

channel, and north basin, respectively (n = 129, 51, and 55 respectively). Prior to sampling, a

Portable Electroanesthesia System (Smith Root) was used to anesthetize fish following approved

animal use protocols of Fisheries and Oceans Canada (FWI-ACC-2017-001, FWI-ACC-2018-

001), the University of Manitoba (F2018-019), and the University of Nebraska-Lincoln (Project

ID: 1208). Fin clips were stored in 95% ethanol. For Lake Manitoba fish, n = 50 fry from the

2019-year class were sampled at Swan Creek Hatchery and stored in 95% ethanol. From Cedar

Bluff Reservoir, anal fin clips were taken from n = 60 sexually mature walleye (n = 40 females

and n = 20 males). Fin clips were collected after gametes were harvested. Procedures for sample

collection at Cedar Bluff Reservoir were approved by the Fort Hays State University Institutional

Animal Care and Use Committee (Protocol #19-0023). Fish were captured using seven 25 mm-

mesh trap nets and four 76 mm mesh gill nets (91.44 X 1.83 m panel per net). In total, $n = 345$

walleye were genotyped for the present study. See Figure 1 for information on sampling

locations.

DNA Extraction, Sequencing, and Single Nucleotide Polymorphism Calling

DNA was extracted using the Qiagen DNeasy Blood & Tissue Kit (QIAGEN, Venlo,

Netherlands) following manufacturer protocols. RAD-seq library preparation was performed

with the NEBNext Ultra DNA Library Prep Kit for Illumina (New England Biolabs, Ipswich,

Massachusetts, USA) with the *PstI* restriction enzyme. BestRAD libraries were baited using

protocols outlined in Euclide et al., (2021) using a RAD-Capture panel for a different walleye

research project in the Laurentian Great Lakes that was still ongoing at the time of publication

(Ali et al., 2016). The panel includes a single bait for 99,636 SNP loci identified from a

preliminary PSTI RAD-sequencing survey of 48 walleye collected in the Great Lakes ($n = 9 – 10$

from each of Lakes Superior, Huron, Michigan, Erie, and Ontario; USA) designed by Arbor

Biosciences (Ann Arbor, Michigan, USA). Target loci were selected based on minor allele

frequency (>0.01) and alignment position along a draft walleye genome to retain one SNP

approximately every 5000 bp along every contig greater than 0.1 megabases. Bait-captured

libraries were then sequenced on a S4 NovaSeq 6000 (Illumina, San Diego, California, USA) at

NovoGene (Sacramento, California, USA).

Raw reads were processed with STACKS v2.3 (Catchen et al., 2013; Catchen et al.,

2011), and aligned to the yellow perch (*Perca flavescens*) genome with bwa v0.7.17 (Feron et

al., 2020; Li & Durbin, 2009). Single nucleotide polymorphisms (SNPs) were called following

best practices with RAD data (Linck & Battey, 2019), and filtered for paralogy with HDPlot

(McKinney et al., 2017). Data for population structure and demographic history were pruned for

linkage with PLINK v1.9 (Purcell et al., 2007). Because relatedness can introduce bias in

association studies, pairwise relatedness was checked using the method of moments as

implemented by PLINK in SNPRelate v1.22 (Zheng et al., 2012). Because XP-EHH requires

complete data, imputation of a set of SNPs with 10% missing data was performed with Beagle

v5.1 (Browning et al., 2018; Browning & Browning, 2007; Weng et al., 2013; Yang et al., 2014).

VCFtools v0.1.16, PGDSpider v2.1.5, the statistical computing environment R v4.0.3, along with

the packages Tidyverse v1.3.0 and vcfR v1.12.0, were used throughout these analyses (Danecek

et al., 2011; Knaus & Grünwald, 2017; Lischer & Excoffier, 2012; R Core Team, 2021;

Wickham et al., 2019). Full details of SNP calling and processing are provided in the

supplemental materials.

To further investigate predicted synteny between the walleye and yellow perch genomes,

synteny was analyzed using progressiveMauve and i-ADHoRe following a pipeline published in

Doerr & Moret (2018), with details provided in the supplementary materials (yellow perch

genome, PLFA_1.0 at NCBI SRA BioProject PRJNA514308; walleye genome, ASM919308v1

at NCBI SRA BioProject PRJNA528354) (Darling et al., 2010; Ghiurcuta & Moret, 2014;

Krzywinski et al., 2009; Proost et al., 2012).

Population Genetics

Admixture v1.3.0 was run over K values of one through six, with 10-fold cross validation

and 1000 bootstrap replicates for parameter standard errors (Alexander et al., 2009). Pophelper

v2.3.0 in R was used to visualize and organize Admixture results (Francis, 2017). For population

assignments, individuals were considered assigned to a cluster when their Q-values were > 0.85

for that cluster. Only assigned individuals were used for population differentiation and

demographic reconstruction.

Hierfstat v0.5-7 was used to find β_{WT} as a measure of population-specific differentiation

from the entire pool and Weir & Cockerham's pairwise F_{ST} between populations (Goudet, 2005;

Weir & Cockerham, 1984; Weir & Goudet, 2017). Ninety five percent confidence intervals were

generated for F_{ST} and β_{WT} over 1,000 bootstrapped iterations each. In addition, observed

heterozygosity (H_O), gene diversity (H_S), and inbreeding coefficients (F_{IS}, 95% confidence

intervals generated over 1,000 bootstrapped iterations) for each population were calculated.

NeEstimator v2.1 was used to estimate effective population size in 95% confidence

intervals for each population using the linkage disequilibrium method and only comparing SNPs

on different chromosomes (Do et al., 2014). Here, datasets were filtered for 10% missing data

per population using population assignments with vcftools, pulled from the pruned SNP dataset.

EASYPOP v2.0.1 (Balloux, 2001) was used to test different scenarios of historic gene

flow that may have led to contemporary F_{ST} between Lake Winnipeg and Lake Manitoba.

Migration rates varied between 0, 0.0001, 0.001, 0.002, 0.003, 0.004, 0.005, 0.01, 0.02, and 0.03

individuals per generation, with migration rates held constant within each simulation run.

Population sizes of 4,500 individuals in Lake Winnipeg and 350 individuals in Lake Manitoba

were assumed constant with equal males and females, based on estimates of N_e (see results).

1,000 simulated loci with two allelic states were used with free recombination allowed, using a

KAM mutation model with mutation rate μ of 3.28×10^{-9} based on divergence between the

blackfin icefish (*Chaenocephalus aceratus*) and dragonfish (*Parachaenichthys charcoti*) (Kim et

al., 2019). Maximal variability (i.e., randomly assigned alleles) was chosen for the initial

population. Each simulation was run over 930 generations, based on the 4.3 year generation time

estimated in Franckowiak et al. (2009) and an approximate lower bound of 4,000 radiocarbon

years since Lake Winnipeg had a reduced northern lakebed (Lewis et al., 2002). This reduced

northern lakebed would have prevented walleye migration between populations via the Dauphin

River and Lake St. Martin in the north basin (Fig 1). 100 replicate runs of EASYPOP were used

for each migration rate tested, with 1000 simulation runs total. For each run, Weir &

Cockerham's pairwise F_{ST} was measured with Hierfsat. These F_{ST} values were compared to the

F_{ST} observed between Lake Winnipeg and Lake Manitoba walleye populations using linkage

disequilibrium-pruned SNPs.

Signatures of Selection

Signatures of selection, where one genomic region has approached or reached fixation in

one population compared to another, were explored with XP-EHH using the R package rehh

v3.2.1, with the markers previously described as phased and imputed after filtering for 90%

present data (Gautier et al., 2017; Sabeti et al., 2007). Data were unpolarized for analyzing XP-

EHH because ancestral and derived alleles were unknown. Individuals with phased and imputed

SNPs were split into the three assigned populations using Admixture results (Q>0.85 per

individual), and extended haplotype homozygosity (EHH) was analyzed for each population

separately (scan_hh). Three pairwise comparisons between each population were used to identify

XP-EHH using false discovery rate corrected *p*-values (*q*-values). Significant candidate regions

of selection were identified by analyzing 100 kilobase (kb) windows overlapping by 10 kb, in

which at least three significant SNPs ($q < 0.05$) showing XP-EHH were found. For visualization,

only significant SNPs within candidate regions were highlighted (thus a significant SNP outside

a candidate region would be de-emphasized), using the R package ggman v0.99.0 with relative

SNP positions to reflect distances between points in the reduced representation data

(https://rdrr.io/github/veera-dr/ggman/). Results for XP-EHH were divided into three pairwise

tests between each assigned population (Cedar Bluff Reservoir, Lake Manitoba, and Lake

Winnipeg), and significant results for each pairwise test were further categorized into which

population showed elevated XP-EHH scores (e.g., in a pairwise test between Lake Manitoba and

Lake Winnipeg, certain SNPs were significant among haplotypes in Lake Manitoba versus others

significant among haplotypes in Lake Winnipeg).

The program pcadapt v4.3.3 was used to explore potential signatures of local adaptation

with outlier tests (Luu et al., 2017; Privé et al., 2020). Here, a scree plot showed that the optimal

choice for principal components (PC) was $K = 2$ based on Cattell's rule, an observation

confirmed by a lack of discernible populations structure in $K > 2$ principal components (Cattell,

1966) (Fig S1). Significance for individual SNPs was determined using *q* values implemented in

the R package qvalue v2.20.0, where a SNP was accepted as significant at $q < 0.05$ (Storey et al.,

2020). The PC associated with a significant SNP was retrieved with *get.pc*, which enabled us to

link significant SNPs with different axes of population structure. pcadapt results were separated

into datasets of those significant along PC1 or PC2, corresponding to latitude and longitude,

respectively.

Hierfstat v0.5-7 was used to provide supporting evidence for selection at individual SNPs

using F'_{ST} with *basic.stats* on the same set of SNPs used with pcadapt. Here, F'_{ST} was used as a

sample size corrected F_{ST} and measured in pairwise comparisons between each of three

populations defined by admixture. In addition, absolute allele frequency differences were

analyzed between populations. VCFtools was used to calculate allele frequencies for each

population. Minor alleles in Lake Winnipeg-assigned individuals were used for comparisons

across populations.

Putative Inversion Analysis

To assess evidence that a region of interest that we identified (chromosome 8, position

15,260,000 - 15,900,000; see results for more detail) was a putative inversion, we conducted the

following analyses as suggested by Huang et al. (2020) and Shi et al. (2021). First, to test

whether the region showed elevated linkage disequilibrium or LD (r^2), we calculated r^2 using

PLINK v1.9 for SNPs on chromosome 8 (--r2 --ld-window-r2 0.05 --ld-window 999999999 --ld-

window-kb 30000) and generated a LD heatmap (Chang et al., 2015; Purcell et al., 2007).

Second, we conducted regional PCA using SNPs within the region to identify whether

individuals were grouped into three clusters along PC1, which is characteristic of chromosomal

inversions. The discreteness of the clustering was calculated as the proportion of the between-

cluster sum of squares over the total using the R function *kmeans* in *adegenet* (Jombart, 2008).

Third, we compared heterozygosity (the proportion of heterozygotes) among the three identified

PCA clusters using Wilcoxon tests ($\alpha = 0.05$) to further confirm that individuals in the middle

PCA cluster was heterozygous for the putative inversion. Lastly, we calculated genotype

frequencies of the putative inversion in all populations and assumed that the more derived

inversion arrangement would have had lower heterozygosity given its relatively recent origin

compared to the ancestral state (Knief et al., 2016; Laayouni et al., 2003; Twyford & Friedman,

2015). Individual assignments for the three identified PCA clusters for the putative inversion

were then visualized with respect to population assignments and site collected, to assess the

frequency of the inversion across the different waterbodies studied.

To provide evidence of potential functional significance for the identified putative

inversion on chromosome 8, gene expression data was collected for genes within the inversion.

RNA-seq data was used from n=48 walleye in three sites in Lake Winnipeg, using previously

published reads (NCBI SRA database accession #PRJNA596986) (Thorstensen et al., 2020).

Three genes in the yellow perch genome were within the putative inversion on chromosome 8:

PDHX, *EHF*, and *LRRC4C*. Therefore, expression of these genes was assessed with counts per

million (CPM) values from the walleye transcriptome-aligned data (Jeffrey et al., 2020; Patro et

al., 2017; Soneson et al., 2015). Differential gene expression was not tested for because the

putative inversion was nearly fixed in the Lake Winnipeg-assigned walleye; differential

expression would thus be expected between populations and not within populations. Details for

RNA-seq are provided in supplementary materials.

Results

DNA Extraction, Sequencing, and SNP Calling

96,955 SNPs were called in STACKS, of which 63,882 were accepted as non-paralogous

SNPs after using HDPlot. Pruning non-paralogous SNPs for linkage with PLINK left 46,342

SNPs for use for population structure and demographic reconstruction. The dataset of 63,882

SNPs was filtered for a maximum of 10% missing data and imputed and phased into a dataset of

18,601 SNPs for use with XP-EHH. Synteny between the walleye and yellow perch genomes

was high, indicating a large degree of concordance between the two genomes (mean weighted

synteny score = 0.999, 95% CI [0.997, 1.0]; mean relaxed synteny sore = 0.999, 95% CI [0.997,

1.0]) (Fig S2).

Population Genetics

Admixture identified K = 2 populations based on the lowest cross validation error

between K = 1 through 6, separating Canadian and Kansas populations of walleye (Fig S3).

However, cross validation error was similar at K = 3 and population structure delineated the

three different water bodies used in the present study; individuals were thus assigned to

populations based on K = 3. Using ancestry coefficients Q > 0.85 for population assignments, 60

individuals were assigned to the Kansas population, 67 individuals to the Lake Manitoba

population, 189 individuals to a Lake Winnipeg population, and 29 to no population. Notably, 27

out of the 29 unassigned individuals were found in Lake Winnipeg (the 2 remaining in Lake

Manitoba), and of those, 18 were caught in the northern Dauphin River, 6 in the central

Matheson Channel, and 2 in the southern Red River. In addition, out of the 67 individuals

assigned to the Lake Manitoba population, 20 were sampled from the Dauphin River in Lake

Winnipeg.

Pairwise F_{ST} showed the greatest differentiation between each Canadian lake and Cedar

Bluff Reservoir, and moderate differentiation between Lake Manitoba and Lake Winnipeg (Fig

2). Population-specific differentiation from the overall pool, β_{WT}, was lowest for Cedar Bluff

Reservoir (Fig S4). Filtering for 90% present data for NeEstimator v2 from the pruned SNP

dataset left 11,784 SNPs available in with the Cedar Bluff Reservoir population, 16,436 SNPs in

the Lake Manitoba population, and 13,213 SNPs in the Lake Winnipeg population. Linkage

disequilibrium-based N_e was highest for the Lake Winnipeg population, and approximately 20-

35x lower in each the Lake Manitoba and Cedar Bluff Reservoir populations (Lake Winnipeg N_e

= 12,084, 95% CI [9,986, 15,282]; Lake Manitoba N_e=336, 95% CI [331, 342]; Cedar Bluff

Reservoir N_e = 587, 95% CI [568, 608]) (Fig 2). Observed heterozygosity, gene diversity, and

inbreeding coefficient (H_O, H_S, and F_{IS}, respectively) were each highest for the Cedar Bluff

Reservoir population, with F_{IS} approximately two times higher for Cedar Bluff walleye than the

walleye in the northern lakes (Fig 2, Fig S4).

Simulations with EASYPOP showed observed F_{ST} between Lake Manitoba and Lake

Winnipeg (between 0.026 and 0.029, Fig 2) most consistent with continuous migration of 0.001

individuals per generation starting 930 generations prior to the present (Fig S5).

Signatures of Selection

The XP-EHH scores between Lake Winnipeg- and Lake Manitoba-assigned individuals

identified 45 SNPs with elevated XP-EHH scores (15 SNPs higher in Lake Winnipeg, 30 higher

in Lake Manitoba), of which 33 SNPs were in candidate regions under selection (11 with higher

XP-EHH scores in Lake Winnipeg, 22 with higher XP-EHH scores in Lake Manitoba) (Fig 3).

The XP-EHH scores between Lake Winnipeg- and Cedar Bluff Reservoir-assigned individuals

identified 26 SNPs with elevated XP-EHH scores (24 higher in Lake Winnipeg, 2 higher in

Cedar Bluff Reservoir), of which 18 were in candidate regions under selection (17 with higher

XP-EHH scores in Lake Winnipeg, 1 with higher XP-EHH scores in Cedar Bluff Reservoir) (Fig

3). Between Lake Manitoba- and Cedar Bluff Reservoir-assigned individuals, 7 SNPs showed

elevated XP-EHH scores (7 with higher XP-EHH scores in Lake Manitoba, 10 with higher XP-

EHH scores in Cedar Bluff Reservoir), of which 5 were in candidate regions under selection (5

with higher XP-EHH scores in Lake Manitoba, 0 with higher XP-EHH scores in Cedar Bluff

Reservoir) (Fig 3). There were 10 candidate regions under selection in XP-EHH between Lake

Winnipeg and Lake Manitoba, 5 candidate regions between Lake Winnipeg and Cedar Bluff

Reservoir, and 0 candidate regions between Lake Manitoba and Cedar Bluff Reservoir (Table

S1). Prominent among candidate regions was a section of chromosome 8 between approximately

15.26 Mb and 15.90 Mb in Lake Winnipeg with high differentiation compared to both Lake

Manitoba and Cedar Bluff Reservoir.

Two principal components (PCs) showed population structure in pcadapt, with PC1

showing 5.76% variance explained by latitudinal differences between sites and PC2 showing

2.28% variance explained by longitudinal differences (Fig 4). 345 SNPs were identified as

significant outliers by pcadapt, of which 226 were significant along PC1 and 119 were

significant along PC2 (Fig 4). Along PC2, between approximately 15.35 Mb and 15.91 Mb,

chromosome 8 showed a region of strong differentiation between Lake Manitoba and Lake

Winnipeg.

The region between 15.26 and 15.90 Mb along chromosome 8 was notable for its

candidate regions identified in XP-EHH that were unique to Lake Winnipeg (Table S1), outlier

SNPs in pcadapt (Fig 4), F'_{ST} outlier SNPs (Fig S6), and SNPs with absolute allele frequency

differences (Fig 3, Fig S7) between Lake Manitoba and Lake Winnipeg. Many of the outlier

SNPs in the chromosome 8 peak for F'_{ST} and absolute allele frequency differences overlapped

with PC2 outlier SNPs from pcadapt at $q < 0.05$ (Fig 3, Fig 4, Fig S6, Fig S7).

Putative Inversion Analysis

We found strong evidence of the genomic region 15.26 - 15.90 Mb on chromosome 8 as a

putative chromosomal inversion (Fig 5). This region demonstrated elevated LD with a mean r^2 of

0.30 and values up to 0.96 (Fig 5B). Regional PCA showed that individuals were clustered into

three distinct groups along PC1 with a discreteness of 0.9626 (Fig 5C) and the middle PCA

cluster displaying significantly higher heterozygosity than the other two clusters (Fig 5D, Fig

S8). There were also significant differences in heterozygosity between the two homokaryotypes

(Fig 5D) and the arrangement with lower heterozygosity (cluster 2) was assumed to be the

derived inverted type. This putative inversion (cluster 2) was nearly fixed in the Red River and

Matheson Island sites of Lake Winnipeg (toward the channel and south basin; Fig 1), was at

intermediate frequency in the Dauphin River (north basin) of Lake Winnipeg, and was nearly

fixed or heterozygous for the opposite genotype near Swan Creek Hatchery in Lake Manitoba

and in Cedar Bluff (Fig 5E). Frequency of the putative inversion visualized in relation to

individual assignments and site collected revealed the inversion was nearly fixed in Lake

Winnipeg-assigned walleye for one genotype, nearly fixed for the other genotype among Lake

Manitoba-assigned walleye, and among unassigned, likely admixed fish in Lake Winnipeg,

nearly fixed for the Lake Winnipeg genotype (Fig S9).

Three genes were identified within the putative inversion: *pyruvate dehydrogenase*

protein X component (*PDHX*), *ETS homologous factor* (*EHF*), and *leucine-rich repeat-*

containing protein 4C (*LRRC4C*). Each of these genes had transcripts expressed in Lake

Winnipeg walleye, with counts per million between 0.4 and 20 (Fig S10).

Discussion

We used a reduced representation approach to genotype *n* = 345 walleye from three

different waterbodies in North America at 46,342 genetic markers. Walleye of the Canadian

lakes diverged from each other despite three scenarios for gene flow: a connecting river, stocking

of fry throughout the 20^th century, and flooding in 1882, 1902, 1904, 2011, and 2014 (and

possibly periodically over evolutionary timescales). Historic gene flow likely occurred in

sporadic pulses, as we observed evidence of limited but present contemporary gene flow possibly

facilitated by flooding in 2011 and 2014. These potential pulses of gene flow were consistent

with population differentiation (F_{ST}) between the Canadian lakes that was lower in magnitude

than each Canadian lake compared to a waterbody in Kansas, USA. Lake Winnipeg walleye

showed unexpectedly strong signatures of selection on chromosome 8, with concurrent evidence

for a chromosomal inversion between 15.26 and 15.90 Mb. An inversion may have played a role

in adaptive divergence between Lake Winnipeg and Lake Manitoba walleye (Wellenreuther &

Bernatchez, 2018; Shi et al., 2021), and three expressed genes (i.e., mRNA transcripts) within it

indicate at least the potential for functional significance of the putative inversion.

Population Genetics

Despite moderate signals of differentiation between the Canadian lakes, admixture

analysis and population assignment identified 11% of individuals sampled from primarily the

north basin of Lake Winnipeg as originating from Lake Manitoba. This gene flow into Lake

Winnipeg may have been associated with stocking programs from Lake Manitoba to Lake

Winnipeg in the 1900s and early 2000s, floods may have facilitated movement in 2011 and 2014

(Ahmari et al., 2016). However, individuals possibly admixed between Lake Manitoba and Lake

Winnipeg were likely the direct descendants (i.e., F_1 crosses) from individuals originating from

each lake, based on admixture analyses (Bray et al., 2010; Reynolds & Fitzpatrick, 2013; Wilson

& Goldstein, 2000). Therefore, we believe flooding in 2011 and 2014 is the more likely route for

movement than stocking, because stocking ended in 2002, approximately 3.5 generations before

sampling in 2017 and 2018, given an estimated 4.3 year generation time (Franckowiak et al.,

2009). Simulations confirmed low gene flow between each lake, assuming continuous gene flow

post-glaciation. Taken together, we suggest that gene flow between Lakes Manitoba and

Winnipeg prior to stocking programs and recent flooding has been sporadic, and that several

differentiated genomic regions suggestive of local adaptation remain in each population despite

recent gene flow.

We observed moderate population differentiation (F_{ST}) between two large lakes in

Canada (Lake Winnipeg and Lake Manitoba), and comparatively stronger differentiation

between the two lakes in Canada and Cedar Bluff Reservoir, Kansas, USA. This relatively

stronger differentiation between the Canadian lakes used here and the waterbody in Kansas was

similar in magnitude to differentiation observed between walleye in Minnesota and Wisconsin

(USA), differences which were attributed to Pleistocene lineages (Bootsma et al., 2020). Walleye

population differentiation between Lake Winnipeg and Lake Manitoba was consistent with

certain other walleye populations within watersheds (e.g., Grand River, Ontario, Canada vs other

Lake Erie stocks in Euclide et al., 2020; Sarah Lake vs Lake Koronis, Minnesota, USA & in

Bootsma et al., 2020). Retreating glacial refugia dispersed walleye into different watersheds

approximately 7,000 - 10,000 years ago (Bootsma et al., 2020; Stepien et al., 2009). Therefore,

the population structure identified in the present study represents differentiation following

geologically recent changes to the overall landscape. This differentiation is consistent with high

spawning site fidelity previously observed in walleye (Horrall, 1981; Jennings et al., 1996;

Stepien et al., 2009; Stepien & Faber, 1998), despite the species' otherwise broad movements in

their waterbodies (Munaweera Arachchilage et al., 2021; Raby et al., 2018; Turner et al., 2021).

Given widespread stocking and potamodromous movements in walleye, it may be

surprising that the species showed as much population differentiation as has been observed

between waterbodies in multiple watersheds (Bootsma et al., 2020; Euclide et al., 2021;

Munaweera Arachchilage et al., 2021; Raby et al., 2018; Turner et al., 2021). While stocking has

led to extensive homogenization in some walleye populations, the lack of gene flow from

stocking in the present data and among other waterbodies implicates potential mechanisms by

which gene flow is prevented (Bootsma et al., 2020). One possible mechanism is environmental.

Here, environmental factors impede gene flow because local adaptation may select against

migrants (Hendry, 2004; Sexton et al., 2014). In a comparison between environmental

mechanisms of isolation and isolation by distance, isolation by environment was more common

in all taxa, including vertebrates (Sexton et al., 2014). As such, environmental differences

between source and stocked waterbodies may have precluded gene flow in different walleye

systems. Genomic architecture is another possible mechanism that can maintain differentiation

despite stocking. Here, physical linkage or chromosomal rearrangements have been hypothesized

to play an important role in local adaptation in the face of gene flow (Tigano & Friesen, 2016).

Walleye in waterbodies with historical pulses of gene flow or recent stocking may thus maintain

differentiation because of genomic architecture facilitating local adaptation (Tigano & Friesen,

2016). Moreover, the hypotheses that environment or genomic architecture may have maintained

differentiation despite opportunities for gene flow are not mutually exclusive. Because genomic

architecture has played a role in local adaptation in several freshwater fishes (Shi et al., 2021),

and environment is a key variable in local adaptation (Coop et al., 2010; Forester et al., 2018),

these two mechanisms may often work in tandem.

Putative Chromosomal Inversion

A putative chromosomal inversion was identified on chromosome 8. Based on PCA,

absolute allele frequency differences, and F'_{ST}, this 640 Kb region on chromosome 8 showed

evidence as an outlier region between Lake Manitoba and Lake Winnipeg. For example, the top

10 outlier SNPs (by $-\log_{10}$ q-value) out of 119 significant outliers ($q < 0.05$) along PC 2 in a

pcadapt analysis are all from this small outlier region (Luu et al., 2017; Privé et al., 2020).

Similarly, the top 10 SNPs by absolute allele frequency difference between Lake Manitoba and

Lake Winnipeg are also from this small outlier region. As a haplotype-based test, XP-EHH

provided evidence that this region was selected for not in Lake Manitoba, but in Lake Winnipeg

instead (Sabeti et al., 2007). This chromosome 8 region was also a consistent outlier with F_{ST}

and BAYESCAN analyses (Foll & Gaggiotti, 2008) in a different study on walleye in Wisconsin

and Minnesota (USA) (Bootsma et al., 2020). However, SNP data were too sparse to interrogate

the genomic architecture of this outlier region. In the present study, a dense panel of SNPs

provided sufficient resolution to investigate the possibility that this outlier region was a

chromosomal inversion.

Linkage disequilibrium, regional PCA, discreteness of clustering, and heterozygosity

provided evidence for a putative chromosomal inversion at higher frequency in Lake Winnipeg

relative to both other waterbodies. This putative inversion was nearly fixed in Lake Winnipeg-

assigned walleye, and at intermediate frequencies in walleye assigned to the other populations in

this study. If the putative inversion had a neutral effect on walleye biology, an intermediate

frequency would have been expected among admixed fish given the parental populations were

almost fixed for opposite genotypes. Instead, among Lake Manitoba-assigned fish caught in Lake

Winnipeg that we hypothesize migrated into Lake Winnipeg (the majority of which were found

in the northern Dauphin River), the putative inversion was nearly fixed for the Lake Winnipeg

genotype. Therefore, this putative inversion may play a role in walleye adaptation to Lake

Winnipeg.

Three genes were within the putative inversion: *PDHX*, *EHF*, and *LRRC4C*. All three

were expressed in Lake Winnipeg walleye, indicating at least some gene transcription activity

within the putative inversion. However, the specific functional significance of these genes in

Lake Winnipeg walleye is unknown. Several other characteristics of Lake Winnipeg walleye are

consistent with signatures of selection unique to the waterbody. Observations of recently-evolved

dwarf walleye morphotypes possibly due to fishing pressure suggest ongoing selection within

Lake Winnipeg (Moles et al., 2010; Sheppard et al., 2018). Additionally, there are patterns of

low gonadal investment despite high lipid concentrations in Lake Winnipeg walleye relative to

walleye in other lakes, indicating a possibly unusual phenotype in walleye of Lake Winnipeg

relative to others (Moles et al., 2008). Walleye habitats are different between Lake Manitoba and

Lake Winnipeg, as well. Lake Winnipeg is one of the largest lakes in the world based on surface

area, but is relatively shallow compared with other large freshwater lakes (Brunskill et al., 1980;

ECCC & MARD, 2020). However, Lake Manitoba is shallower still, with a 4.9 m mean depth

(compared with a mean depth for Lake Winnipeg of 13.3 m and 9 m in the north and south

basins, respectively) (ECCC & MARD, 2020; Rawson, 1952). Drainage from the more saline

Lake Manitoba increases salinity in Lake Winnipeg's north basin, but not in Lake Winnipeg's

south basin (Brunskill et al., 1980; ECCC & MARD, 2020; ECCC, 2011). Consequently,

possible local adaptation in Lake Winnipeg may be an evolutionary response to environmental

differences in the waterbody.

Conclusion

The identification of a putative inversion contributes to a broad literature on

chromosomal inversions in diverse taxa (see Wellenreuther & Bernatchez, 2018), although

information on inversions is relatively scarce for freshwater fishes (Shi et al., 2021). In addition,

many observed inversions have been > 1 Mb in length, possibly because reduced representation

sequencing methods may be biased for detecting large inversions (Wellenreuther & Bernatchez,

2018). The present data show a small putative inversion in an obligate freshwater fish discovered

via reduced representation sequencing (i.e., Rapture). That this putative inversion may have

played a role in recent divergence between fish of two connected lakes, and that its frequency

among likely admixed fish is consistent with a possible adaptive role in Lake Winnipeg walleye,

indicates a potential importance for chromosomal inversions in freshwater fishes more generally.

Inversions have been previously related to environmental adaptation, such as in freshwater

emerald shiner (*Notropis atherinoides*) with high gene flow (Wellenreuther & Bernatchez, 2018;

Shi et al., 2021). Heterogeneous habitats with opportunities for gene flow are common for

freshwater fishes (Griffiths, 2015; Mushet et al., 2019), and chromosomal inversions, along with

other genomic architectures, may facilitate local adaptation in many organisms that live in such

connected environments.

Acknowledgements

We thank Dr. Jeff Long for contributing walleye fry from Swan Creek Hatchery and for valuable discussions of Manitoba walleye ecology, diversity, and stocking. We thank Colin Charles, Colin Kovachik, Doug Leroux, Nicole Turner, Mike Gaudry, Sarah Glowa, and Emily Barker, who provided fin clips used for walleye DNA from Lake Winnipeg. We thank Evelien de Greef for guidance on synteny analyses and the map figure, and Dr. Kristen Gruenthal for sharing her expertise in sequencing library preparation. We thank David Splausbury and his staff from the Kansas Department of Wildlife, Parks, and Tourism for their assistance in the collection of samples at Cedar Bluff Reservoir. We thank Dr. Colin Garroway for discussions about β_{WT}. Many analyses in this manuscript were enabled using computing resources provided by WestGrid (www.westgrid.ca) and Compute Canada (www.computecanada.ca). This work was supported by a Fisheries and Oceans Canada Ocean and Freshwater Science Contribution Program Partnership Fund grant awarded to KMJ, JRT, and Dr. Darren Gillis, and a Natural Sciences and Engineering Research Council of Canada Discovery Grants awarded to KMJ (#05479) and JRT (#06052). Work by JRT is also supported by the Canada Research Chairs program (#223744) and the Faculty of Science, University of Manitoba (#319254).

References

Ahmari, H., Blais, E. L., & Greshuk, J. (2016). The 2014 flood event in the Assiniboine River
Basin: Causes, assessment and damage. *Canadian Water Resources Journal*, *41*(1–2), 85–
93. https://doi.org/10.1080/07011784.2015.1070695

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in
unrelated individuals. *Genome Research*, *19*(9), 1655–1664.
https://doi.org/10.1101/gr.094052.109

Ali, O. A., O'Rourke, S. M., Amish, S. J., Meek, M. H., Luikart, G., Jeffres, C., et al. (2016).
Rad capture (Rapture): Flexible and efficient sequence-based genotyping. *Genetics* 202,
389–400. doi:10.1534/genetics.115.183665.

Arostegui, M. C., Quinn, T. P., Seeb, L. W., Seeb, J. E., & McKinney, G. J. (2019). Retention of
a chromosomal inversion from an anadromous ancestor provides the genetic basis for
alternative freshwater ecotypes in rainbow trout. *Molecular Ecology*, *28*(6), 1412–1427.
https://doi.org/10.1111/mec.15037

Backhouse-James, S. M., & Docker, M. F. (2012). Microsatellite and mitochondrial DNA
markers show no evidence of population structure in walleye (Sander vitreus) in Lake
Winnipeg. *Journal of Great Lakes Research*, *38*(SUPPL. 3), 47–57.
https://doi.org/10.1016/j.jglr.2011.05.005

Balian, E. V., Segers, H., Lévèque, C., & Martens, K. (2008). The Freshwater Animal Diversity
Assessment: an overview of the results. *Hydrobiologia*, *595*(1), 627–637.
https://doi.org/10.1007/s10750-007-9246-3

Balloux, F. (2001). EASYPOP (version 1.7): A computer program for population genetics
simulations. *Journal of Heredity*, *92*(3), 301–302. https://doi.org/10.1093/jhered/92.3.301

Berg, P. R., Star, B., Pampoulie, C., Bradbury, I. R., Bentzen, P., Hutchings, J. A., Jentoft, S., &
Jakobsen, K. S. (2017). Trans-oceanic genomic divergence of Atlantic cod ecotypes is
associated with large inversions. *Heredity*, *119*(6), 418–428.
https://doi.org/10.1038/hdy.2017.54

Berg, Paul R., Star, B., Pampoulie, C., Sodeland, M., Barth, J. M. I., Knutsen, H., Jakobsen, K.
S., & Jentoft, S. (2016). Three chromosomal rearrangements promote genomic divergence
between migratory and stationary ecotypes of Atlantic cod. *Scientific Reports*, *6*(1), 23246.
https://doi.org/10.1038/srep23246

Bootsma, M. L., Miller, L., Sass, G. G., Euclide, P. T., & Larson, W. A. (2020). The ghosts of
propagation past: haplotype information clarifies the relative influence of stocking history
and phylogeographic processes on contemporary population structure of walleye (Sander
vitreus). *Evolutionary Applications*, *June 2020*, 1–21. https://doi.org/10.1111/eva.13186

Bray, S. M., Mulle, J. G., Dodd, A. F., Pulver, A. E., Wooding, S., & Warren, S. T. (2010).
Signatures of founder effects, admixture, and selection in the Ashkenazi Jewish population.
Proceedings of the National Academy of Sciences of the United States of America, *107*(37),
16222–16227. https://doi.org/10.1073/pnas.1004381107

Browning, B. L., Zhou, Y., & Browning, S. R. (2018). A One-Penny Imputed Genome from Next-Generation Reference Panels. *American Journal of Human Genetics*, *103*(3), 338–348. https://doi.org/10.1016/j.ajhg.2018.07.015

Browning, S. R., & Browning, B. L. (2007). Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *American Journal of Human Genetics*, *81*(5), 1084–1097. https://doi.org/10.1086/521987

Bruner, J. C. (2021). Stizostedion Rafinesque, 1820 (Percidae) is the Valid Generic Name for Walleye, Sauger, and Eurasian Pikeperch. *Fisheries*, *1820*(1816), 1–5. https://doi.org/10.1002/fsh.10582

Brunskill, G. J., Elliott, S. E. M., & Campbell, P. (1980). Watershed data pertinent to the limnology of Lake Winnipeg: Canadian manuscript report of fisheries & aquatic sciences. In *Canadian Manuscript Report of Fisheries and Aquatic Sciences* (Issue 1556). http://hdl.handle.net/1993/23608

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: An analysis tool set for population genomics. *Molecular Ecology*, *22*(11), 3124–3140. https://doi.org/10.1111/mec.12354

Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W., & Postlethwait, J. H. (2011). Stacks: Building and genotyping loci de novo from short-read sequences. *G3: Genes, Genomes, Genetics*, *1*(3), 171–182. https://doi.org/10.1534/g3.111.000240

Cattell, R. B. (1966). The Scree Test For The Number Of Factors. *Multivariate Behavioral Research*, *1*(2), 245–276. https://doi.org/10.1207/s15327906mbr0102_10

Chang, C. C., Chow, C. C., Tellier, L. C. A. M., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-generation PLINK: Rising to the challenge of larger and richer datasets. *GigaScience*, *4*(1), 1–16. https://doi.org/10.1186/s13742-015-0047-8

Connallon, T., & Olito, C. (2021). Natural selection and the distribution of chromosomal inversion lengths. *Molecular Ecology*, *July*, 1–15. https://doi.org/10.1111/mec.16091

Coop, G., Witonsky, D., Di Rienzo, A., & Pritchard, J. K. (2010). Using environmental correlations to identify loci underlying local adaptation. *Genetics*, *185*(4), 1411–1423. https://doi.org/10.1534/genetics.110.114819

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., & Durbin, R. (2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

Darling, A. E., Mau, B., & Perna, N. T. (2010). Progressivemauve: Multiple genome alignment with gene gain, loss and rearrangement. *PLoS ONE*, *5*(6). https://doi.org/10.1371/journal.pone.0011147

Do, C., Waples, R. S., Peel, D., Macbeth, G. M., Tillett, B. J., & Ovenden, J. R. (2014). NeEstimator v2: Re-implementation of software for the estimation of contemporary effective population size (Ne) from genetic data. *Molecular Ecology Resources*, *14*(1), 209–

214. https://doi.org/10.1111/1755-0998.12157

Doerr, D., & Moret, B. M. E. (2018). Comparative RNA Genomics. In J. C. Setubal, J. Stoye, & P. F. Stadler (Eds.), *Methods in Molecular Biology* (Vol. 1704, pp. 317–329). Humana Press. https://doi.org/10.1007/978-1-4939-7463-4_14

Dorant, Y., Cayuela, H., Wellband, K., Laporte, M., Rougemont, Q., Mérot, C., Normandeau, E., Rochette, R., & Bernatchez, L. (2020). Copy number variants outperform SNPs to reveal genotype-temperature association in a marine species. *Molecular Ecology*, *July*, 1–18. https://doi.org/10.1101/2020.01.28.923490

ECCC. (2011). *State of Lake Winnipeg: 1999–2007*. http://hdl.handle.net/1993/23915

ECCC & MARD. (2020). *State of Lake Winnipeg, 2nd Edition*.

Euclide, P. T., MacDougall, T., Robinson, J. M., Faust, M. D., Wilson, C. C., Chen, K., Marschall, E. A., Larson, W., & Ludsin, S. (2021). Mixed□stock analysis using Rapture genotyping to evaluate stock□specific exploitation of a walleye population despite weak genetic structure. *Evolutionary Applications*, *14*(5), 1403–1420. https://doi.org/10.1111/eva.13209

Feron, R., Zahm, M., Cabau, C., Klopp, C., Roques, C., Bouchez, O., Eché, C., Valière, S., Donnadieu, C., Haffray, P., Bestin, A., Morvezen, R., Acloque, H., Euclide, P. T., Wen, M., Jouano, E., Schartl, M., Postlethwait, J. H., Schraidt, C., … Guiguen, Y. (2020). Characterization of a Y-specific duplication/insertion of the anti-Mullerian hormone type II receptor gene based on a chromosome-scale genome assembly of yellow perch, Perca flavescens. *Molecular Ecology Resources*, *20*(2), 531–543. https://doi.org/10.1111/1755-0998.13133

Fielder, D. G. (1992). Evaluation of Stocking Walleye Fry and Fingerlings and Factors Affecting Their Success in Lower Lake Oahe, South Dakota. *North American Journal of Fisheries Management*, *12*(2), 336–345. https://doi.org/10.1577/1548-8675(1992)012<0336:eoswfa>2.3.co;2

Fisheries and Oceans Canada. (2019). *Survey of recreational fishing in Canada, 2015*. 21.

Fisheries and Oceans Canada. (2021). *Freshwater Fisheries - Catches and Landed Values by Species, By Province/Territory, 2019*. https://www.dfo-mpo.gc.ca/stats/commercial/land-debarq/freshwater-eaudouce/2019-eng.htm

Foll, M., & Gaggiotti, O. (2008). A Genome-Scan Method to Identify Selected Loci Appropriate for Both Dominant and Codominant Markers: A Bayesian Perspective. *Genetics*, *180*(2), 977–993. https://doi.org/10.1534/genetics.108.092221

Forester, B. R., Lasky, J. R., Wagner, H. H., & Urban, D. L. (2018). Comparing methods for detecting multilocus adaptation with multivariate genotype–environment associations. *Molecular Ecology*, *27*(9), 2215–2233. https://doi.org/10.1111/mec.14584

Francis, R. M. (2017). pophelper: an R package and web app to analyse and visualize population structure. *Molecular Ecology Resources*, *17*(1), 27–32. https://doi.org/10.1111/1755-0998.12509

Franckowiak, R. P., Sloss, B. L., Bozek, M. A., & Newman, S. P. (2009). Temporal effective
size estimates of a managed walleye Sander vitreus population and implications for genetic-
based management. *Journal of Fish Biology*, *74*(5), 1086–1103.
https://doi.org/10.1111/j.1095-8649.2008.02170.x

Fraser, D. J., Weir, L. K., Bernatchez, L., Hansen, M. M., & Taylor, E. B. (2011). Extent and
scale of local adaptation in salmonid fishes: Review and meta-analysis. *Heredity*, *106*(3),
404–420. https://doi.org/10.1038/hdy.2010.167

Gautier, M., Klassmann, A., & Vitalis, R. (2017). rehh 2.0: a reimplementation of the R package
rehh to detect positive selection from haplotype structure. *Molecular Ecology Resources*,
17(1), 78–90. https://doi.org/10.1111/1755-0998.12634

Ghiurcuta, C. G., & Moret, B. M. E. (2014). Evaluating synteny for improved comparative
studies. *Bioinformatics*, *30*(12), 9–18. https://doi.org/10.1093/bioinformatics/btu259

Goudet, J. (2005). HIERFSTAT, a package for R to compute and test hierarchical F-statistics.
Molecular Ecology Notes, *5*(1), 184–186. https://doi.org/10.1111/j.1471-8286.2004.00828.x

Griffiths, D. (2015). Connectivity and vagility determine spatial richness gradients and
diversification of freshwater fish in North America and Europe. *Biological Journal of the
Linnean Society*, *116*(4), 773–786. https://doi.org/10.1111/bij.12638

Hartman, G. F. (2009). A Biological Synopsis of Walleye (Sander vitreus). *Canadian
Manuscript Report of Fisheries and Aquatic Sciences*, *2888*, v–48. http://www.dfo-
mpo.gc.ca/Library/337847.pdf

Hendry, A. P. (2004). Selection against migrants contributes to the rapid evolution of
ecologically dependent reproductive isolation. *Evolutionary Ecology Research*, *6*(8), 1219–
1236.

Horrall, R. M. (1981). Behavioral Stock-Isolating Mechanisms in Great Lakes Fishes with
Special Reference to Homing and Site Imprinting. *Canadian Journal of Fisheries and
Aquatic Sciences*, *38*(12), 1481–1496. https://doi.org/10.1139/f81-201

Huang, K., Andrew, R. L., Owens, G. L., Ostevik, K. L., & Rieseberg, L. H. (2020). Multiple
chromosomal inversions contribute to adaptive divergence of a dune sunflower ecotype.
Molecular Ecology, *29*(14), 2535–2549. https://doi.org/10.1111/mec.15428

Jeffrey, J. D., Carlson, H., Wrubleski, D., Enders, E. C., Treberg, J. R., & Jeffries, K. M. (2020).
Applying a gene-suite approach to examine the physiological status of wild-caught walleye
(Sander vitreus). *Conservation Physiology*, *8*(1), 1–12.
https://doi.org/10.1093/conphys/coaa099

Jennings, M. J., Claussen, J. E., & Philipp, D. P. (1996). Evidence for Heritable Preferences for
Spawning Habitat between Two Walleye Populations. *Transactions of the American
Fisheries Society*, *125*(6), 978–982. https://doi.org/10.1577/1548-
8659(1996)125<0978:EFHPFS>2.3.CO;2

Jombart, T. (2008). Adegenet: A R package for the multivariate analysis of genetic markers.
Bioinformatics, *24*(11), 1403–1405. https://doi.org/10.1093/bioinformatics/btn129

Kim, B.-M., Amores, A., Kang, S., Ahn, D.-H., Kim, J.-H., Kim, I.-C., Lee, J. H., Lee, S. G.,
Lee, H., Lee, J., Kim, H.-W., Desvignes, T., Batzel, P., Sydes, J., Titus, T., Wilson, C. A.,
Catchen, J. M., Warren, W. C., Schartl, M., … Park, H. (2019). Antarctic blackfin icefish
genome reveals adaptations to extreme environments. *Nature Ecology & Evolution*, *3*(3),
469–478. https://doi.org/10.1038/s41559-019-0812-7

Kirubakaran, T. G., Grove, H., Kent, M. P., Sandve, S. R., Baranski, M., Nome, T., De Rosa, M.
C., Righino, B., Johansen, T., Otterå, H., Sonesson, A., Lien, S., & Andersen, Ø. (2016).
Two adjacent inversions maintain genomic differentiation between migratory and stationary
ecotypes of Atlantic cod. *Molecular Ecology*, *25*(10), 2130–2143.
https://doi.org/10.1111/mec.13592

Knaus, B. J., & Grünwald, N. J. (2017). vcfr: a package to manipulate and visualize variant call
format data in R. *Molecular Ecology Resources*, *17*(1), 44–53.
https://doi.org/10.1111/1755-0998.12549

Knief, U., Hemmrich-Stanisak, G., Wittig, M., Franke, A., Griffith, S. C., Kempenaers, B., &
Forstmeier, W. (2016). Fitness consequences of polymorphic inversions in the zebra finch
genome. *Genome Biology*, *17*(1), 1–22. https://doi.org/10.1186/s13059-016-1056-3

Koppelman, J. B., Sullivan, K. P., & Jeffries, P. J. (1992). Survival of Three Sizes of Genetically
Marked Walleyes Stocked into Two Missouri Impoundments. *North American Journal of*
Fisheries Management, *12*(2), 291–298. https://doi.org/10.1577/1548-
8675(1992)012<0291:SOTSOG>2.3.CO;2

Krzywinski, M., Schein, J., Birol, I., Connors, J., Gascoyne, R., Horsman, D., Jones, S. J., &
Marra, M. A. (2009). Circos: An information aesthetic for comparative genomics. *Genome*
Research, *19*(9), 1639–1645. https://doi.org/10.1101/gr.092759.109

Laayouni, H., Hasson, E., Santos, M., & Fontdevila, A. (2003). The evolutionary history of
Drosophila buzzatii. XXXV. Inversion polymorphism and nucleotide variability in different
regions of the second chromosome. *Molecular Biology and Evolution*, *20*(6), 931–944.
https://doi.org/10.1093/molbev/msg099

Leitwein, M., Garza, J. C., & Pearse, D. E. (2017). Ancestry and adaptive evolution of
anadromous, resident, and adfluvial rainbow trout (Oncorhynchus mykiss) in the San
Francisco bay area: application of adaptive genomic variation to conservation in a highly
impacted landscape. *Evolutionary Applications*, *10*(1), 56–67.
https://doi.org/10.1111/eva.12416

Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology &*
Evolution, *17*(4), 183–189. https://doi.org/10.1016/S0169-5347(02)02497-7

Lewis, C. F. M., Forbes, D. L., Todd, B. J., Nielsen, E., Thorleifson, L. H., Henderson, P. J.,
McMartin, I., Anderson, T. W., Betcher, R. N., Buhay, W. M., Burbidge, S. M., Schröder-
Adams, C. J., King, J. W., Moran, K., Gibson, C., Jarrett, C. A., Kling, H. J., Lockhart, W.
L., Last, W. M., … Vance, R. E. (2002). Uplift-driven expansion delayed by middle
Holocene desiccation in Lake Winnipeg, Manitoba, Canada. *Geology*, *29*(8), 743–746.
https://doi.org/10.1130/0091-7613(2001)029<0743:UDEDBM>2.0.CO;2

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics*, *25*(14), 1754–1760.
https://doi.org/10.1093/bioinformatics/btp324

Linck, E., & Battey, C. J. (2019). Minor allele frequency thresholds strongly affect population
structure inference with genomic data sets. *Molecular Ecology Resources*, *19*(3), 639–647.
https://doi.org/10.1111/1755-0998.12995

Lischer, H. E. L., & Excoffier, L. (2012). PGDSpider: An automated data conversion tool for
connecting population genetics and genomics programs. *Bioinformatics*, *28*(2), 298–299.
https://doi.org/10.1093/bioinformatics/btr642

Luu, K., Bazin, E., & Blum, M. G. B. (2017). pcadapt: an R package to perform genome scans
for selection based on principal component analysis. *Molecular Ecology Resources*, *17*(1),
67–77. https://doi.org/10.1111/1755-0998.12592

Manitoba Government. (2020). *Lake Information for Anglers*. Lake Information for Anglers.
https://experience.arcgis.com/experience/2557cda82dcc4a348fbb71304cedcf6d/page/page_
2/

McKinney, G. J., Waples, R. K., Seeb, L. W., & Seeb, J. E. (2017). Paralogs are revealed by
proportion of heterozygotes and deviations in read ratios in genotyping-by-sequencing data
from natural populations. *Molecular Ecology Resources*, *17*(4), 656–669.
https://doi.org/10.1111/1755-0998.12613

McKinney, G., McPhee, M. V., Pascal, C., Seeb, J. E., & Seeb, L. W. (2020). Network analysis
of linkage disequilibrium reveals genome architecture in chum salmon. *G3: Genes,*
Genomes, Genetics, *10*(4), 1553–1561. https://doi.org/10.1534/g3.119.400972

Moles, M. D., Robinson, B. W., Johnston, T. A., Cunjak, R. A., Jardine, T. D., Casselman, J. M.,
& Leggett, W. C. (2010). Morphological and trophic differentiation of growth morphotypes
of walleye (Sander vitreus) from Lake Winnipeg, Canada. *Canadian Journal of Zoology*,
88(10), 950–960. https://doi.org/10.1139/Z10-062

Moles, Matthew D., Johnston, T. A., Robinson, B. W., Leggett, W. C., & Casselman, J. M.
(2008). Is gonadal investment in walleye (Sander vitreus) dependent on body lipid reserves?
A multipopulation comparative analysis. *Canadian Journal of Fisheries and Aquatic*
Sciences, *65*(4), 600–614. https://doi.org/10.1139/F07-186

Munaweera Arachchilage, I. P., Muthukumarana, S., Gillis, D. M., Watkinson, D. A., Charles,
C., & Enders, E. C. (2021). Assessing Movement Patterns using Bayesian State-Space
Models on Lake Winnipeg Walleye. *Canadian Journal of Fisheries and Aquatic Sciences*,
1–48. https://doi.org/10.1139/cjfas-2020-0262

Mushet, D. M., Alexander, L. C., Bennett, M., Schofield, K., Christensen, J. R., Ali, G., Pollard,
A., Fritz, K., & Lang, M. W. (2019). Differing Modes of Biotic Connectivity within
Freshwater Ecosystem Mosaics. *Journal of the American Water Resources Association*,
55(2), 307–317. https://doi.org/10.1111/1752-1688.12683

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast
and bias-aware quantification of transcript expression. *Nature Methods*, *14*(4), 417–419.
https://doi.org/10.1038/nmeth.4197

Pearse, D. E., Miller, M. R., Abadía-Cardoso, A., & Garza, J. C. (2014). Rapid parallel evolution
of standing variation in a single, complex, genomic region is associated with life history in
steelhead/rainbow trout. *Proceedings of the Royal Society B: Biological Sciences*,
281(1783). https://doi.org/10.1098/rspb.2014.0012

Privé, F., Luu, K., Vilhjálmsson, B. J., Blum, M. G. B., & Rosenberg, M. (2020). Performing
Highly Efficient Genome Scans for Local Adaptation with R Package pcadapt Version 4.
Molecular Biology and Evolution, *37*(7), 2153–2154.
https://doi.org/10.1093/molbev/msaa053

Proost, S., Fostier, J., De Witte, D., Dhoedt, B., Demeester, P., Van De Peer, Y., & Vandepoele,
K. (2012). i-ADHoRe 3.0-fast and sensitive detection of genomic homology in extremely
large data sets. *Nucleic Acids Research*, *40*(2). https://doi.org/10.1093/nar/gkr955

Puncher, G. N., Rowe, S., Rose, G. A., Leblanc, N. M., Parent, G. J., Wang, Y., & Pavey, S. A.
(2019). Chromosomal inversions in the Atlantic cod genome: Implications for management
of Canada's Northern cod stock. *Fisheries Research*, *216*(March), 29–40.
https://doi.org/10.1016/j.fishres.2019.03.020

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J.,
Sklar, P., De Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: A tool set for
whole-genome association and population-based linkage analyses. *American Journal of
Human Genetics*, *81*(3), 559–575. https://doi.org/10.1086/519795

R Core Team. (2021). *R: A language and environment for statistical computing. R Foundation
for Statistical Computing*. https://www.r-project.org/

Raby, G. D., Vandergoot, C. S., Hayden, T. A., Faust, M. D., Kraus, R. T., Dettmers, J. M.,
Cooke, S. J., Zhao, Y., Fisk, A. T., & Krueger, C. C. (2018). Does behavioural
thermoregulation underlie seasonal movements in Lake Erie walleye? *Canadian Journal of
Fisheries and Aquatic Sciences*, *75*(3), 488–496. https://doi.org/10.1139/cjfas-2017-0145

Rawson, D. S. (1952). Mean Depth and the Fish Production of Large Lakes. *Ecology*, *33*(4),
513–521. https://doi.org/10.2307/1931525

Rempel, L. L., & Smith, D. G. (1998). Postglacial fish dispersal from the Mississippi refuge to
the Mackenzie River basin. *Canadian Journal of Fisheries and Aquatic Sciences*, *55*(4),
893–899. https://doi.org/10.1139/f97-257

Reynolds, R. G., & Fitzpatrick, B. M. (2013). Tests of two methods for identifying founder
effects in metapopulations reveal substantial type II error. *Genetica*, *141*(1–3), 119–131.
https://doi.org/10.1007/s10709-013-9711-z

Sabeti, P. C., Varilly, P., Fry, B., Lohmueller, J., Hostetter, E., Cotsapas, C., Xie, X., Byrne, E.
H., McCarroll, S. A., Gaudet, R., Schaffner, S. F., Lander, E. S., Frazer, K. A., Ballinger, D.
G., Cox, D. R., Hinds, D. A., Stuve, L. L., Gibbs, R. A., Belmont, J. W., … Stewart, J.
(2007). Genome-wide detection and characterization of positive selection in human
populations. *Nature*, *449*(7164), 913–918. https://doi.org/10.1038/nature06250

Sexton, J. P., Hangartner, S. B., & Hoffmann, A. A. (2014). Genetic isolation by environment or
distance: Which pattern of gene flow is most common? *Evolution*, *68*(1), 1–15.

https://doi.org/10.1111/evo.12258

Sheppard, K. T., Hann, B. J., & Davoren, G. K. (2018). Growth rate and condition of walleye
(Sander vitreus), sauger (sander canadensis), and dwarf walleye in a large Canadian lake.
Canadian Journal of Zoology, *96*(7), 739–747. https://doi.org/10.1139/cjz-2017-0276

Shi, Y., Bouska, K. L., McKinney, G. J., Dokai, W., Bartels, A., McPhee, M. V, & Larson, W.
A. (2021). Gene flow influences the genomic architecture of local adaptation in six riverine
fish species. *BioRxiv*. https://doi.org/10.1101/2021.05.18.444736

Sinclair-Waters, M., Bradbury, I. R., Morris, C. J., Lien, S., Kent, M. P., & Bentzen, P. (2018).
Ancient chromosomal rearrangement associated with local adaptation of a postglacially
colonized population of Atlantic Cod in the northwest Atlantic. *Molecular Ecology*, *27*(2),
339–351. https://doi.org/10.1111/mec.14442

Sodeland, M., Jorde, P. E., Lien, S., Jentoft, S., Berg, P. R., Grove, H., Kent, M. P., Arnyasi, M.,
Olsen, E. M., & Knutsen, H. (2016). "Islands of Divergence" in the atlantic cod genome
represent polymorphic chromosomal rearrangements. *Genome Biology and Evolution*, *8*(4),
1012–1022. https://doi.org/10.1093/gbe/evw057

Soneson, C., Love, M. I., & Robinson, M. D. (2015). Differential analyses for RNA-seq:
transcript-level estimates improve gene-level inferences. *F1000Research*, *4*(2), 1521.
https://doi.org/10.12688/f1000research.7563.1

Stepien, C. A., Murphy, D. J., Lohner, R. N., Sepulveda-Villet, O. J., & Haponski, A. E. (2009).
Signatures of vicariance, postglacial dispersal and spawning philopatry: Population genetics
of the walleye Sander vitreus. *Molecular Ecology*, *18*(16), 3411–3428.
https://doi.org/10.1111/j.1365-294X.2009.04291.x

Stepien, Carol A., & Faber, J. E. (1998). Population genetic structure, phylogeography and
spawning philopatry in walleye (Stizostedion vitreum) from mitochondrial DNA control
region sequences. *Molecular Ecology*, *7*(12), 1757–1769. https://doi.org/10.1046/j.1365-
294x.1998.00512.x

Storey A. J.; Dabney, A.; Robinson, D., J. D. . B. (2020). qvalue: Q-value estimation for false
discovery rate control. In *R package version 2.22.0: Vol. null* (Issue null, p. null-null).
http://github.com/jdstorey/qvalue

Thorstensen, M. J., Jeffrey, J. D., Treberg, J. R., Watkinson, D. A., Enders, E. C., & Jeffries, K.
M. (2020). Genomic signals found using RNA sequencing show signatures of selection and
subtle population differentiation in walleye (Sander vitreus) in a large freshwater
ecosystem. *Ecology and Evolution*, *10*(14), 7173–7188. https://doi.org/10.1002/ece3.6418

Tigano, A., & Friesen, V. L. (2016). Genomics of local adaptation with gene flow. *Molecular*
Ecology, *25*(10), 2144–2164. https://doi.org/10.1111/mec.13606

Turner, N. A., Charles, C., Watkinson, D. A., Enders, E. C., Klein, G., & Rennie, M. D. (2021).
Historical and contemporary movement and survival rates of walleye (Sander vitreus) in
Lake Winnipeg, Canada. *Journal of Great Lakes Research*, *xxxx*.
https://doi.org/10.1016/j.jglr.2021.01.012

Twyford, A. D., & Friedman, J. (2015). Adaptive divergence in the monkey flower Mimulus

guttatus is maintained by a chromosomal inversion. *Evolution*, *69*(6), 1476–1486. https://doi.org/10.1111/evo.12663

Weir, B. S., & Cockerham, C. C. (1984). ESTIMATING F -STATISTICS FOR THE ANALYSIS OF POPULATION STRUCTURE. *Evolution*, *38*(6), 1358–1370. https://doi.org/10.1111/j.1558-5646.1984.tb05657.x

Weir, Bruce S., & Goudet, J. (2017). A unified characterization of population structure and relatedness. *Genetics*, *206*(4), 2085–2103. https://doi.org/10.1534/genetics.116.198424

Wellenreuther, M., & Bernatchez, L. (2018). Eco-Evolutionary Genomics of Chromosomal Inversions. *Trends in Ecology and Evolution*, *33*(6), 427–440. https://doi.org/10.1016/j.tree.2018.04.002

Weng, Z., Zhang, Z., Zhang, Q., Fu, W., He, S., & Ding, X. (2013). Comparison of different imputation methods from low- to high-density panels using Chinese Holstein cattle. *Animal*, *7*(5), 729–735. https://doi.org/10.1017/S1751731112002224

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., … Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Wilson, J. F., & Goldstein, D. B. (2000). Consistent long-range linkage disequilibrium generated by admixture in a bantu-semitic hybrid population. *American Journal of Human Genetics*, *67*(4), 926–935. https://doi.org/10.1086/303083

Yang, Y., Wang, Q., Chen, Q., Liao, R., Zhang, X., Yang, H., Zheng, Y., Zhang, Z., & Pan, Y. (2014). A new genotype imputation method with tolerance to high missing rate and rare variants. *PLoS ONE*, *9*(6). https://doi.org/10.1371/journal.pone.0101025

Zheng, X., Levine, D., Shen, J., Gogarten, S. M., Laurie, C., & Weir, B. S. (2012). A high-performance computing toolset for relatedness and principal component analysis of SNP data. *Bioinformatics*, *28*(24), 3326–3328. https://doi.org/10.1093/bioinformatics/bts606

Data Accessibility

All metadata and scripts used for analyses in this manuscript are available at: https://github.com/BioMatt/walleyeDNA. Raw sequence reads are available at the National Center for Biotechnology Information Sequence Read Archive (accession #PRJNA800092).

Author Contributions

Matt J. Thorstensen conducted the analyses and wrote the original manuscript. Matt J.

Thorstensen, Jennifer D. Jeffrey, Ken M. Jeffries, Jason R. Treberg, Douglas A. Watkinson, and

893 Eva C. Enders conceived of the study. Eva C. Enders, Douglas A. Watkinson, and Yasuhiro

894 Kobayashi contributed to obtaining samples. Peter T. Euclide and Wesley A. Larson developed

895 the sequencing panel used. Yue Shi analyzed the putative inversion. Ken M. Jeffries and Jason

896 R. Treberg acquired funding. All authors participated in editing the manuscript and contributed

897 to interpretation and analyses.

Tables and Figures

Figure 1. Map of waterbodies included in the present study. Cedar Bluff Reservoir, Kansas, USA, represents walleye (*Sander vitreus*) in an entirely stocked population of unknown origin. Lake Manitoba and Lake Winnipeg (Manitoba, Canada) walleye represent native populations, although Lake Winnipeg has been stocked by Lake Manitoba walleye throughout the 20[th] century. In addition, floods in 1882, 1902, 1904, 2011, and 2014 increased waterflow from Lake Manitoba to Lake Winnipeg, and may have facilitated walleye gene flow.

Figure 2. Population differentiation, effective population size, and observed heterozygosity for walleye (*Sander vitreus*) in three North American waterbodies. Cedar Bluff Reservoir (Kansas, USA) represents an entirely stocked population of walleye, while Lake Manitoba and Lake Winnipeg (Manitoba, Canada) walleye represent populations with possible gene flow. F_{ST} represents Weir & Cockerham's pairwise F_{ST}, while H_O observed heterozygosity, each found using Hierfstat. Effective population size for each population is represented by N_e, found using NeEstimator v2. For F_{ST} and N_e, 95% confidence intervals are provided in parentheses.

Figure 3. Signatures of selection inferred from cross-population extended haplotype homozygosity (XP-EHH) between each population of walleye (*Sander vitreus*) analyzed in the present study, along with absolute allele frequency differences between two waterbodies. In A), Cedar Bluff Reservoir (Kansas, USA) represents an entirely stocked population, while Lake Manitoba and Lake Winnipeg (Manitoba, Canada) each represent native populations with possible gene flow. Walleye DNA was aligned to the yellow perch (*Perca flavescens*) reference genome, and chromosome numbers represent yellow perch chromosomes. Unplaced scaffolds are not visualized. Single nucleotide polymorphisms (SNPs) were considered significant with $q<0.05$ and only if the SNPs were in a candidate region defined by 100 kilobase regions of the genome overlapping by 10 kilobases, in which ≥ 3 SNPs total were significant. Because XP-EHH focuses on haplotypes of unusual length and high frequency in one population compared to another, the population in which haplotypes are long and are at or approaching fixation can be identified. Here, the population in which XP-EHH was significant is identified with colours, where blue represents Lake Winnipeg, red represents Lake Manitoba, and yellow represents Cedar Bluff Reservoir. In B), absolute allele frequency differences were calculated between Lake Manitoba and Lake Winnipeg-assigned walleye. PC1 and PC2 refer to significant SNPs from principal component 1 or 2 from pcadapt (see Fig. 4). The genes *pyruvate dehydrogenase protein X component (PDHX), ETS homologous factor (EHF), and leucine-rich repeat-containing protein 4C (LRRC4C)* were within the candidate region for selection in Lake Winnipeg on chromosome 8. The candidate regions for selection from XP-EHH are emphasized with gray background, which together were identified as the putative chromosomal inversion.

Figure 4. Principal components analysis of walleye (*Sander vitreus*) from three waterbodies across North America. Cedar Bluff Reservoir (Kansas, USA) represent an entirely stocked population of walleye of unknown origin. Lake Winnipeg and Lake Manitoba (Manitoba, Canada) represent native populations of walleye with possible gene flow. Principal components analysis was done using pcadapt, while population assignments were done using Admixture with K=3 groups. Individuals were assigned to a population with $Q > 0.85$ for a population, and individuals with maximum $Q \leq 0.85$ were considered unassigned. Significant SNPs were distinguished according to the first two principal components in which they most strongly contributed variation, where principal component 1 (PC1) corresponded to latitudinal differences among populations, while principal component 2 (PC2) was most strongly characterized by variation between Lake Winnipeg and Cedar Bluff Reservoir relative to Lake Manitoba. Only SNPs with *q*-values < 0.05 were highlighted for visualization and retained for downstream analyses. Walleye DNA was aligned to the yellow perch (*Perca flavescens*) reference genome. Unplaced scaffolds were not visualized, and chromosome numbers refer to chromosomes in the yellow perch genome.

Figure 5. Characterization of the putative inversion on chromosome 8 (15.26 - 15.90 Mb). A) PCA based on SNPs within the putative inversion, coloured by site. B) Linkage disequilibrium heatmap for SNPs within the putative inversion, with boundary of the putative inversion emphasized by dashed lines. C) Clusters identified using k-means clustering represent two homozygous groups, red and blue, and a heterozygous purple group. D) Observed individual heterozygosity within each cluster identified. E) Genotype frequency distribution by collection site and cluster. Bars represent the proportion of individuals from each k-means cluster.

Lake Manitoba

Lake Winnipeg

F_{ST} = 0.027
(0.026 - 0.029)

H_O = 0.082

N_e = 336
(331 - 342)

H_O = 0.082
N_e = 12,084
(9,986 - 15,282)

F_{ST} = 0.076
(0.074 - 0.078)

F_{ST} = 0.072
(0.070 - 0.074)

H_O = 0.085
N_e = 587
(568 - 608)

Cedar Bluff Reservoir

Principal Components Analysis

Site Collected

- Cedar Bluff Reservoir
- Swan Creek Hatchery
- Dauphin River
- Matheson Island
- Red River

Population Assignments

- ☆ Admixed
- ☐ Kansas
- ○ Manitoba
- △ Winnipeg

PC2 (2.28% variance)

PC1 (5.76% variance)

Significant PC1 SNPs

Significant PC2 SNPs

Chromosome 8 (Position 15260000 – 15900000)