Main Manuscript for

Reduction of free-roaming cat population requires high-intensity neutering
in spatial contiguity to mitigate compensatory effects

Idit Gunther[a], Hadas Hawlena[b], Lior Azriel[a], Dan Gibor[a], Olaf Berke[c], and Eyal
Klement[a*]

[a] Koret School of Veterinary Medicine, The Robert H. Smith Faculty of Agriculture,
Food & Environment, The Hebrew University of Jerusalem, Rehovot 76100, Israel

[b] Mitrani Department of Desert Ecology, Swiss Institute for Dryland Environmental and
Energy Research, The Jacob Blaustein Institutes for Desert Research, Ben-Gurion
University of the Negev, Sde-Boker 8499000, Israel

[c] Department of Population Medicine, Ontario Veterinary College, University of Guelph,
Ontario N1G 2W1, Canada

* Corresponding author:
Eyal Klement
Tel: 972-54-6766004, email: eyal.klement@mail.huji.ac.il

Author Contributions:

Conception and design: Gunther I., Klement E.
Acquisition of data: Gunther I., Azriel L., Gibor D.
Interpretation of data and analyzes: Gunther I., Hawlena H., Berke O., Klement E.
Drafting the manuscript: Gunther I., Hawlena H., Berke O., Klement E.
Review and confirmation of manuscript: all authors.

Competing Interest Statement: All the authors declare that they do not have competing
interests in this manuscript.

Classification: Biological sciences/ecology

Keywords: Population management, TNR, Free-Roaming Cats, Fertility control,
Population compensatory mechanisms.

Abstract
When free-roaming in natural areas, the domestic cat (*Felis silvestris catus*) is ranked
high among the most destructive alien species. Near human dwellings, it might risk
humans, impair sanitation, and suffer from poor welfare. Cats' popularity as companion
animals complicates their population control. Thus, culling is often replaced by a fertility
control method called "Trap-Neuter-Return/Release (TNR)," which is considered more
humane. Despite the extensive application of TNR, a long-term controlled study was
never performed to test its effectiveness. We present a uniquely designed controlled field

experiment for examining TNR effectiveness. The study was performed over a twelve-
year period, divided into pre-intervention, mixed- and full-intervention phases, and
spanned a 20Km^2 urban area. Trends of cat, intact-female, and kitten counts, cat
reproduction, and carcass reports were compared among study phases and areas with
different neutering intensities. The cat population increased during the first two study
phases and did not decline in highly neutered populations, presumably due to cat
immigration. Expansion of high-intensity neutering to the entire city in the full-
intervention phase (>70% neutering percentage) reversed cat population growth, reaching
an annual ca. 7% reduction. This population reduction was limited by a rebound increase
in cat reproduction and longevity. We conclude that cat population management by TNR
should be performed in high-intensity, continuously, and in geographic contiguity to
enable population reduction. To enhance management effectiveness and mitigate
compensatory effects, we recommend further evaluating an integrated strategy that
combines TNR with complementary methods (e.g., vital resource regulation, ill cat
euthanasia, and adoption).

Significance Statement

Though popular companion animals, domestic cats pose numerous problems when free-
roaming, including predation of wildlife, hazards to humans, impaired sanitation, and a
decrease in their welfare. Thus, managing their populations is essential. The Trap-Neuter-
Return method (TNR, capturing, sterilizing, returning/releasing) is widely employed for
managing cat populations. However, there is a lack of long-term controlled evidence for
its effectiveness. We examined the outcomes of high-intensity TNR by performing a
twelve-year controlled field experiment, the largest to date. Neutering over 70% of the
cats caused population decline when applied over contiguous areas. However, it was
limited by a rebound increase in reproduction and survival. These findings provide a
robust quantification of the limitations and the long-term effectiveness of TNR.

Main Text

Introduction

As a generalist predator, the domestic cat (*Felis silvestris catus*) has been listed among the world's 100 worst non-native species (1). The domestic cat is a fast life-history species (i.e., early maturation, small body size, rapid reproduction) (2-4), which has been distributed around the globe mainly as a pet (5). Cats have been known to form non-domiciliary and often human-independent populations known as free-roaming cats, which tend to cause adverse environmental effects (1, 6-11). The most prominent ecological negative impact of cats is on islands. They are considered responsible for at least 14% of global bird, mammal, and reptile extinctions and are the principal threat to almost 8% of critically endangered birds, mammals, and reptiles (11). Moreover, cats also have a significant ecological effect on the mainland due to direct predation, the transmission of diseases to other species, fear-related effects, and the alteration of demographic processes such as source-sink dynamics (6). As a result of their potential to transmit certain zoonotic diseases and injure humans due to aggressive behavior, free-roaming cats constitute a hazard to public health (7, 8). They might also cause a nuisance to humans, mainly by impairing sanitation (9, 10). These conflicts are intensified in the urban setting where cats form large populations, increasing human-cat interaction (12, 13).

The adverse effects caused by free-roaming cats have increased the motivation to artificially manage their populations, aiming either to diminish their related nuisances or to preserve natural ecosystems (14-19). Animal population management is based on two strategies: resource limitation in the habitat or actions applied to individuals. The latter strategy can be further divided into culling and fertility control methods. While culling aims to increase mortality above the natural rate, fertility control aims to decrease reproduction below the natural rate (20). Culling has been successful in eradicating cat populations on certain islands (14, 16, 17). However, there are several examples of the failure of culling to accomplish population control in fast life-history species, such as voles (21), mice, rats, jirds (22), rabbits (23), cats (24, 25), and foxes (26). In contrast, information on the efficacy of fertility control on vertebrate populations is scarce (27). Specifically for free-roaming cats, theoretical studies predicted that culling performs better than Trap-Neuter-Return (TNR), a common method for controlling cat fertility (20, 28-31). However, despite these predictions, the TNR method has been progressively implemented in cat populations over widespread areas instead of culling, mainly due to moral considerations and public opinion (18, 32-36).

A growing collection of studies has examined the effects of TNR on cat population dynamics, however ours is the first controlled study conducted to observe the long-term effectiveness of TNR. Past studies have yielded inconsistent finding with several finding decreases in population growth indicators (37-52), and others finding stabilization or even increases (53-58). These inconsistent results might stem from differences in management duration and efforts, the examined populations (e.g., closed versus open populations, small versus large-scale populations), or the study environment. The inference of the overall long-term consequences of TNR is further limited due to: absence

of control (37-41, 43-45, 48-52, 54, 56, 57), combining TNR with other control tactics
(i.e., adoption and euthanasia of clinically ill or retrovirus-positive cats) (37-46, 49-52),
short-term follow-up (37, 38, 42, 45, 46, 51, 53-56, 58), small sample size (37, 38, 40, 43,
45, 47, 49, 50, 52-54, 58), relying on indirect indices of population growth (38, 39, 42,
46, 49, 51, 52, 57), and examining populations in secluded areas (41, 44).

Boon et al. (59) called for a long-term and large-scale study in cat populations to close
the knowledge gap and overcome shortcomings of previous studies. Here we present the
results of a 12-year longitudinal large-scale experiment. We assessed the effect of
sterilization (i.e., spaying/neutering) on the long-term temporal and spatial dynamics of
free-roaming cat populations. While adjusting for environmental factors, we compared
cat population dynamics before and at partial and full implementation of a TNR program.
We found that population compensatory mechanisms may limit the magnitude of the
neutering effect. Thus, TNR should be performed continuously, at high intensity, and in
spatial contiguity to enable long-term cat population regulation.

120
121 **Results**
122
Neutering intensity across statistical areas and study phases

The study was set to assess long-term TNR effectiveness over time and space (**Figures 1,**
S1). In the 1st (pre-intervention) phase, neutering was not applied. In the 2nd (mixed-
intervention) phase, high-intensity neutering was applied in about half of the city, and in
the 3rd (full-intervention) phase, TNR was applied in the entire city. We confirmed the
neutering status in 96% of the surveyed cats by ear-mark detection, counting both
unneutered (un-marked) females (queens) and males and neutered (marked) cats, along
with kittens. Repeated annual surveys were performed during September and October of
2012, 2013, 2014, and 2018. Overall, in the 50 surveyed statistical areas, 13,718 cat-
observations were documented during the study period, of which 1,486 were kittens. The
phase- and group-specific neutering percentages corroborated the respective TNR
intensity (**Table 1**, **Table S1**). For a visual perspective of the municipal TNR intensity
across statistical areas and study phases, see **Figure S2**.

At the end of the mixed-intervention phase (end of 2014), the overall neutering
percentages in the entire surveyed area, the low TNR (n=26) and high TNR (n=24)
statistical areas, were 51, 33, and 71%, respectively (**Table 1**). To compare the effect of
the highest-intensity neutering to the lowest-intensity, we further divided the statistical
areas into quartiles according to the observed neutering percentage in that year (2014).
Group-1 (n=12) represents the lowest neutering intensity and consists of the statistical
areas with neutering percentage at the lowest quartile. Group-2 (n=15) represents the
highest neutering intensity and consists of the statistical areas with neutering percentage
at the highest quartile (see **Figure S1** for the detailed spatial study design). At the end of
the mixed-intervention phase, the overall neutering percentages in Group-1 and Group-2
were 19 and 80%, respectively. At the end of the full-intervention phase (end of 2018),
the neutering percentage in the entire area was 72%, and in the low TNR, high TNR,

Group-1, and Group-2, the overall percentages were 75, 69, 69, and 77%, respectively
(**Table 1, Figure 2A**).

Environmental human-related factors affecting cat population dynamics

We modeled year-based cross-sectional data (adjusting for neutering percentage) to
screen for potential human-related factors that might influence cat population dynamics.
As expected, we found a consistent negative association between neutering percentage
and kitten counts (**Table S3**). Interestingly, in some years, the cat counts were positively
associated with the neutering percentage (i.e., 2012 and 2018), indicating possibly higher
TNR efforts in areas with higher, rather than lower, cat densities. In addition, cat
population size parameters, cat counts, and resident reports of cat carcasses and
reproduction were positively correlated with the human population density (**Table S3**).
Though less consistent, the volume of accessed-waste-bins was also positively correlated
with some cat population parameters. Therefore, we adjusted for human population
density and accessed-waste-bins by including them in the analysis of neutering
association with cat population trends. The other tested human-related factors were not
consistently associated with the cat population parameters. We did not find a consistent
spatial correlation in these models, and therefore it was not further included in the trend
models.

Survey results: Cat and kitten counts across statistical areas and study phases

During the mixed-intervention phase (2012-2014), we found an overall increase of 26.5%
in the cat counts in the surveyed area. Therein, we found an average annual growth of
20.7% in Group-1, which was exposed to the lowest neutering intensity (**Table 1, Figure
2B**). At the same time, the cat counts remained stable in Group-2, which was exposed to
the highest neutering intensity (**Table 2**). In contrast, during the full-intervention phase,
we found an overall reversal of population growth, with a 23.1% reduction in the cat
counts in the entire surveyed area. During this phase, the annual decrease in the two
treatment groups was similar (ca. -7% annually; **Figure 2B)**.

As opposed to cats, we did not find a significant reduction in the kitten counts in the
entire surveyed area (**Table 1, Figure 2C**). This overall stability was a result of opposite
trends in the two treatment groups. As expected, during the mixed-intervention phase,
kitten counts increased annually by 20.2% in Group-1 but decreased by 42.4% in Group-
2. Whereas, during the full-intervention phase, these trends reversed, with a decrease of
14.1% in Group-1 and an increase of 21.4% in Group-2 (**Table 2**). The latter increase in
kitten counts in Group-2, during the full-intervention phase, was unexpected. A possible
explanation for this increase is a reduction in neutering intensity in some statistical areas
in this group during the full-intervention phase. To examine this possibility, we
reanalyzed the data of both groups, excluding statistical areas in which the neutering
percentages at the end of the full-intervention phase were below 70%.

Here, too, we found the same trend reversal in kitten counts (**Table S4**). Therefore, a
decrease in neutering intensity in some statistical areas is unlikely to explain the observed
increase. An alternative explanation is that reduced agonistic behavior in neutered cats

enhanced compensatory mechanisms such as increased kitten survival, a rise in the
frequency of pregnancies per queen, or increased litter size due to lower competition.
Assuming this hypothesis was correct, we predicted to find an overall increase in the
kitten-to-queen ratio in the full-intervention study phase. We indeed found a 2.25-fold
increase during this phase (**Table 1, Figure 2E**). This increase was significant in both
treatment groups (**Table 2**). In Group-2, it occurred due to a significant increase in the
kitten number and stabilization of the queen number. In Group-1, the queen number
decreased annually by 38% shortly after introducing high-intensity neutering, while a
14% insignificant decrease in the kitten number was observed (**Table 2, Figure 2C,D,E**).
No significant reduction in the kitten-to-queen ratio was observed in Group-2 during the
mixed-intervention period, possibly due to parallel immigration of intact females.

Resident report results: Cat carcasses and reproduction events across statistical areas
and study phases

We complemented the survey data analysis with those of resident reports on cat
reproduction events and cat carcasses. This dataset increased the study resolution in both
space (covering more statistical areas) and time (daily reports from 2007 to 2018). To
examine the construct validity of these reports, we tested the correlation of reproduction
reports with the surveyed kitten counts and found a strong correlation between the two
datasets (Pearson correlation coefficient=0.55, p<0.001, **Figure S6**).

In addition, to ensure that these reports represent trends unique to cats, we compared
them with the number of all available reports to the call center (after excluding cat-related
reports) and with the reports of all carcasses (excluding cats and including canines,
poultry, rodents, hyraxes, hedgehogs, equines, reptiles, boar and tortoises) . The trends of
these two latter report types showed a consistent increase during the study period (**Figure**
S4). Overall, this increased trend suggests that people's awareness or tendency to use the
call center increased with time. Unlike the general trends, the reports of cat carcasses and
reproduction were linked to the application of TNR. As with the general reports, both
showed a mild positive trend during the pre-intervention phase. However, this was
followed by a negative trend, from the onset of the mixed-intervention phase until one
year after its end (from 2011 to 2015) (**Figure 3**). The magnitude of this negative trend
was ca. 50%, starting at 325 and 120 and reaching a low of 200 and 60 of carcasses and
reproduction reports per month, respectively. Notably, during the mixed-intervention
phase, the number of carcass and reproduction reports decreased while the cat counts
increased (**Table 1**, **Figure 2**), indicating a reduction in cat reproduction and mortality.
One year after the onset of the full-intervention phase (2016-2018), the number of carcass
reports stabilized, but the number of reproduction reports showed an unexpected increase
(**Figure 3**). This increase suggests a possible rebound effect of TNR on reproduction.

Both report types exhibited a prominent seasonal pattern: carcasses peaked twice, in
May-June and in October, whereas reproduction reports peaked only once in April-May
(**Figure 3**). Unlike these reports, the general reports to the call center peaked in July-
August (**Figure S4**), showing that the seasonal pattern of carcass and reproduction
reports is specific for cats and did not represent human behavior tendency to refer to the

call center. While the seasonal pattern of cat reproduction reports was maintained throughout the study, the seasonality of carcass reports gradually diminished from the onset of TNR until it disappeared during the full-intervention phase (see 'crude data' in **Figure 3b**). Thus, it implies that TNR resulted in a decrease in mortality and a decrease in the seasonal (reproduction-linked) mortality pattern.

We next compared the carcass and reproduction report trends between the treatment groups. Both trends coincided with TNR efforts (**Figure 4**, **Table 2**). In the pre-intervention phase, the numbers of carcass and reproduction reports were stable. During the mixed-intervention phase, both carcass and reproduction reports declined considerably in Group-2 (-12% annually for carcass and -24.5% for reproduction reports), while their numbers were stable in Group-1 (0.9% for carcass and 3.9% for reproduction reports). The number of carcass reports decreased considerably in Group-1 (-13.6%) in the full-intervention phase, indicating a decrease in cat number and/or mortality. Similar to kitten counts, in Group-2, a considerable rebound increase of reproduction reports occurred during the full-intervention phase (22.6%). We found similar trends when the analysis was repeated after excluding statistical areas in which the neutering percentages at the end of the full-intervention phase were below 70%. (**Figure S5, Table S4**).

Discussion

The current study is the first long-term controlled field experiment examining TNR effectiveness in free-roaming cat populations. Using a unique experimental design in which we manipulated neutering intensity across time and space, we distinguished between short-term and long-term population consequences of TNR. To achieve high-resolution data we used two independent data-collection methods-a repeated cross-sectional annual survey and daily resident reports. Parameters of population dynamics derived by both methods showed the same trends, suggesting that TNR can reduce population size, but only when applied at high rates in spatial contiguity. Below, we discuss the evidence suggesting processes that might challenge TNR effectiveness and thus should be considered when applying it. These include compensatory effects, such as increased cat survival and/or litter size, as well as possible influx of immigrant cats into vacant niches. We then compare our results to those generated by theoretical simulations, discussing the TNR option compared to its alternatives, and end by offering directions for further simulations and planning strategies.

Maintaining high neutering intensity in spatial contiguity is required to mitigate counteracting effects

During the mixed-intervention phase, the cat population growth in Group-2 was considerably lower than in Group-1. However, the cat population did not decline in Group-2 despite maintaining a neutering percentage of 80% (**Figures 2A, B**), a number previously suggested as sufficient for reducing the population size of free-roaming cats (28, 60). We propose that during the mixed-intervention phase, the TNR effect on cat number was counteracted by the immigration of intact cats from unneutered surrounding

populations. It was previously shown that sterilization-enhanced behavioral changes
resulted in increased immigration of free-roaming cats from low- into high-TNR regions
(53). This important population compensatory process was also documented in other
managed species (reviewed in Ransom et al. 2014 (27)). The study area boundaries may
have enabled long-distance cat immigration mainly from the neighboring northern cities,
where municipal TNR campaigns were not implemented . However, we believe it was
negligible due to the small home range of urban cats (60, 61).

During the full-intervention phase, an average annual reduction of more than 7% was
observed in groups 1 & 2 . Our surveys further indicated that during the four years of this
phase, the cat population in the entire city declined by ca. 25%. It is possible that the cat
counts decreased in the entire city due to temporal effects unrelated to TNR. For
example, a change in vital resource availability (e.g., decreased feeding or change in
sanitary conditions), decreased abandonment, or increased adoption. However, this
possibility is unlikely for three reasons: First, the finding of cat counts reduction was
adjusted for changes in the environmental factors influencing cat population size (i.e.,
human population density and the per human accessible waste bins). Second, the survey
periods following each intervention change were not longer than 4–5 years. A marked
change in human behavior to opposite directions both spatially and temporally in such
short periods is unlikely. Third, though abandonment and adoption of free roaming cats
do occur in Israel, they constitute only a negligible factor relative to the remarkable
density of the free-roaming cats' population (751-2300/Km^2 (61, 62)). The most probable
alternative explanation for the cat population trend changes between the phases is the
expansion of high neutering intensity (>70%) to the entire city during the full-
intervention phase. This expansion may have reduced the effect of the immigration of
cats from low TNR to high TNR areas. We conclude that maintaining high-intensity
neutering in spatial contiguity is a prerequisite for reducing cat population size.

Limiting processes should be considered in TNR programs

Along with the encouraging cat population decline, our findings support the occurrence
of several compensatory effects, limiting TNR effectiveness: **Reduced mortality**; Cat
carcass trend reversed from an increase during pre-intervention to a constant decrease
during the mixed-intervention phase. This decrease opposed growth in cat population size
at this phase. These opposite trends indicate that the reduced carcass reports are due to
reduced mortality following TNR. **Reduced reproduction-related mortality**; The
diminished seasonal pattern of carcass reports following increased neutering intensity
(see 'crude data' in Fig 3B and Fig 4b) indicates a decreased reproduction-related
mortality. It was previously shown that in the Northern hemisphere, kittens present a
seasonal appearance (4, 10, 52, 64) and experience higher morbidity and mortality than
adults (53, 63, 64). Moreover, the mortality of intact cats was higher than neutered ones
(65, 66), probably due to associated reproductive behavior. **Increased fertility and**
reduced kitten mortality; While the neutering percentage exceeded 72% in the entire
city during the full-intervention phase, a considerable increase in the kitten-to-queen ratio
was documented in both study groups. This increased ratio may stem from several
plausible explanations: an elevation in kitten survival, a rise in the frequency of

pregnancies per queen, and a rise in litter size. This conclusion is supported by a rise in
cat reproduction reports in Group-2 during the full-intervention phase. Increased fertility
and decreased mortality of juveniles and adults were previously reported to follow
fertility control in domestic cats (34, 53, 63, 64) and other vertebrates (27, 67-71). Both
compensatory processes could result from higher food availability and decreased resource
competition following population decline (64, 72). In addition, they could be caused by
diminished agonistic behavior in neutered male cats (73). These mechanisms, together
with cat immigration, might limit the TNR-derived decrease in cat population size.

Comparison of our findings to the results of cat population simulation models

Several theoretical simulations aimed to predict the long-term consequences of TNR and
provide specific management insights (20, 28, 30, 31). Of these, only two studies
examined the effect of TNR in an open population (20, 30). In these studies, population
reduction was achieved by implementing TNR, though the immigration of non-neutered
cats limited its effectiveness. This situation resembles the mixed-intervention phase in the
current study when untreated areas surrounded treated areas. Interestingly, in our study,
during this phase, the cat population did not decline. There are at least two possible
explanations for this discrepancy. First, we assume that similar to other population
dynamics, immigration is a density-dependent process. Thus, the immigration rate may
depend on density differences of both total cats and intact cats between the managed and
surrounding populations (53). By modeling immigration as a density-independent
process, the theoretical models might have underestimated this process's magnitude.
Second, these models did not include other density-dependent population compensatory
mechanisms such as increased survival and reproduction, which probably occurred in our
study.

Comparison of TNR to culling

When culling or other means of cat removal were applied intensively, they eradicated cat
populations on certain islands (14, 16, 17). Although culling may be efficient, concurrent
similar compensatory mechanisms to those observed in our study might also limit its
effectiveness (24, 25). In some environmental contexts, TNR is considered as an
alternative to culling. Since the public acceptance of TNR is better than culling (9, 35, 36,
74), it improves public cooperation with policy-makers, which is an important factor for
population reduction success. The merits of TNR go beyond those achieved by
population reduction and include improved welfare (64) and a reduction in cat-related
nuisances, which are mainly associated with reproduction (38, 75). On the other hand, the
costs of TNR implementation should be taken into account. Especially challenging is the
maintenance of high-intensity neutering for long periods, considering the significant
decrease in trapping efficiency when the percentage of neutered cats is high. More than
one million dollars ($US) were invested in the current project over the nine years of TNR
implementation. This cost should be compared to alternative population management
programs, considering the differences between countries, economies, jurisdictions, and
ethical perceptions.

Complementary approaches that can further improve TNR benefits

Combining TNR with removing a significant portion of the cat population (i.e.,
euthanasia of ill cats and adoption) can improve its effectiveness (40, 42-44). However,
the removal of cats might not be applicable everywhere. Our study's analysis of the
environmental factors influencing cat populations highlights an alternative approach to
complement TNR. The results show an association between vital anthropogenic resources
(i.e. accessed-waste-bins and human density, which may be associated with higher food
availability to cats) and cat population parameters. As suggested by Boone (2015),
limiting such resources might be combined with TNR to reduce carrying capacity along
with the reduction in population size (76). Despite population reduction, neutered cats
might continue to hunt (77, 78). Therefore, TNR is not an effective means by itself for
preventing predation. It was shown that manipulating food content (e.g. providing high-
meat protein and grain-free diet) and applying environmental enrichment (e.g. object
play) reduced cat tendency to hunt (79), and placing collar-mounted devices interfered
with hunting (80). Such measures can be therefore combined with TNR to diminish this
adverse effect further.

Strengths and limitations

The current experiment is unique not only in the controlled and long-term design but also
in the large number and depth of data of neutered and observed cats, the large covered
urban area which is systematically stratified (i.e., statistical areas), and the high-
resolution data of real-time events (resident report data) and environmental factors. With
this, it should be mentioned that the compensatory processes we observed are presumably
density-dependent. Therefore, generalization of our results to areas with low cat densities
should be taken cautiously. In addition, such a large study might also suffer from other
limitations. For example, some processes might affect study results, such as changes in
human behavior, changes in the urban environment (e.g., constructions), local changes in
cat numbers, and even stochastic changes during the study period. We excluded the effect
of these processes by controlling TNR both in space and time, incorporating possible
confounders in the analysis, performing comparisons of the two datasets (resident data
and annual counts data performed by experts), and comparing our data to datasets
representing mortality of other animals and human behavior.

One limitation is a possible overestimation of the number of neutered cats due to a higher
probability of detecting sociable cats than shy cats. These sociable cats are also more
likely to be captured and neutered before capturing shy cats. Using friendly calling to
increase the counted cat number might have enhanced this bias. However, a recent study
showed that while friendly calling significantly increased the number of detected cats by
79%, its effect on the estimated neutering percentage was insignificant. Moreover, the
mean insignificant increase by 18% (from 46% to 56%) of neutering percentage
diminished almost completely when the neutering percentage was high (above 74%).
This observation was probably due to the growing proportion of neutered shy cats as the
neutering percentage increased (81). Therefore, despite the inherent uncertainty in the
number of neutered cats not seen by the observers, we assume that our estimations of

neutering intensity are reliable and valid and thus enable robust evidence for TNR effects
and limitations.

Conclusion

Our results indicate that fertility control can be applied to manage and even decrease the
size of open free-roaming cat populations. However, the enhancement of compensatory
mechanisms and the influx of immigrant cats might limit this effect. Therefore, we
conclude that maintaining a high neutering rate for a prolonged period and in spatial
contiguity is necessary to reduce cat population size. We recommend integrating these
with complementary methods such as vital resource regulation, ill cat euthanasia, and
adoption. In a broader view, our study suggests that the success of TNR should be
measured by considering its environmental context, aims, and public acceptance. While
simulation studies may aid in assessing such considerations, for enabling realistic
interpretation, we recommend including parameters of possible compensatory effects and
density-dependent immigration in these models. In particular, we call for a future robust
evaluation of the effectiveness of integrated approaches.

Materials and Methods

Study site

We conducted the study from 2007 to 2018 in the city of Rishon-LeZion, located within
the greater Tel-Aviv metropolitan area in Israel. The jurisdictional area of Rishon-LeZion
is 58.7 km^2, divided according to the Israeli Central Bureau of Statistics into 64
commercial and residential statistical areas. Three statistical areas spanning 33.8 Km^2
include an isolated research institute, industrial and non-inhabited areas, and, therefore,
not part of the study area. The city's human population comprised 240,666 residents
living in a 25 km^2 area at the end of 2014 (Central Bureau of Statistics, Israel), divided
into 58 residential areas (each consists of approximately 4000 residents) and three
commercial areas (**Figure S1**). These 61 statistical areas constituted the study site.

Study design

The study period was divided into three consecutive phases (**Figure 1, Table S1**): (1) the
pre-intervention phase, (2) the mixed-intervention phase, and (3) the full-intervention
phase. The pre-intervention phase preceded the initiation of the TNR program from
January 2007 to the end of 2009. In the mixed-intervention phase, we allocated the 61
statistical areas into high- and low- intervention. In 31 statistical areas, the municipality
implemented a high-intensity multi-annual TNR program from October 2009 until
October 2014. The municipal TNR actions were limited in the other 30 statistical areas.
We attempted to allocate TNR so that the human-related characteristics will be
comparable between the high and low TNR regions but still maintain spatial contiguity
among the un-neutered and neutered areas (see **Table S2** for demographic comparison
between the high and low intervention areas). During this phase, 10,925 free-roaming
cats were neutered with a male to female ratio of 1:1.07. In the full-intervention phase
(November 2014 until December 2018), high-TNR efforts were expanded to the entire

city. During this phase, 11,219 free-roaming cats were neutered with a male to female
ratio of 1:1.06. The specific cat trapping and neutering procedures are detailed in the
supplementary information.

Data collection

To achieve high-resolution data on cat population dynamics, we used two independent
data collection methods—a repeated annual survey and daily resident reports.

Annual surveys:

We counted free-roaming cats via repeated annual surveys performed during September
and October of 2012, 2013, 2014, and 2018. We conducted these surveys in 50 statistical
areas (n=26 in the low-TNR statistical areas and n=24 in the high-TNR statistical areas)
along fixed walking transects. In each statistical area, we chose one or two transects using
a stratified random sampling design. Each statistical area was sampled by two observers,
with each transect sampled twice each year. To increase the number of the observed cats
the observer used friendly calling. We documented the individual characteristics of each
observed cat, including sex, presumptive age (kitten<6 months vs. adults > 6 months),
and sterilization status (according to the presence/absence of ear-marks). We summarized
the annual counts of cats (adults (neutered and intact), intact-females (queens)), kittens
(<6 months of age), and kitten-to-queen ratio per statistical area and year. We calculated
the annual neutering percentage by dividing the total counts of neutered cats observed in
each statistical area and year by the total counts of observed cats with identified neutering
status. The transect walks were recorded using a cellular GPS recording application
("Endomondo™—Running & Walking, Android application," Under Armour Inc.,
Maryland, USA), to ensure walking on precisely the same path of each transect and to
measure the transect length. The total length of the transects was 100.87 km (**Figure S3**).
A comprehensive description of this sampling method, including its intra-observer, inter-
observer, inter-transect agreement, and validity, is detailed in Gunther et al. 2020 (81).

Resident reports:

We retrieved data on cat carcasses and reproduction during 2007-2018 from the
continuously available municipal emergency call center. The center receives voice
complaints and reports from concerned residents on real-time events in the city
jurisdiction area. The following data were recorded for each reported event: time and date
of the call, location of the event, personal details about the calling resident, and a
synopsis of the reported event. To confirm that the observed trends were unique to cat-
related reports, we compared them to other municipal-based reports. The 2007-2018 cat
carcass reports were compared to reports of carcasses of other animals (n=7,535), and cat
reproduction reports were compared to the general municipal reports (apart from cat-
related reports, n= 3,725,873) **(Figure S4)**. Further information on the call center data
and its retrieval methods are detailed in the **supplementary information.**

Environmental human-related data:

The following environmental human-related factors were collected for Rishon-LeZion
statistical areas during the study period: *human population density*, *socioeconomic status,*
volume and type of *waste bins, cat feeding locations, educational institutions, food*
marketing businesses, built-up area and *park area, age of neighborhood (years) and TNR*
actions performed by the municipality. We provide further details on these factors in the
supplementary information.

Statistical analysis

Data were aggregated for each combination of statistical area and year. These values
constituted the sampling units in all analyses. The following outcome variables were
analyzed: cat counts, kitten counts, queen counts, kitten-to-queen ratio, neutering
percentage, cat carcass reports, and cat reproduction reports. Except for the time-series
analyses, all statistical analyses were performed by fitting generalized linear models
(GLM) or generalized linear mixed models (GLMM) to these data (where the statistical
area was used as a random effect). In all of these models, a negative binomial distribution
was used. Since we analyzed rates, we used offsets as denominators (an offset is a
variable that is forced to have a coefficient of 1 in the model. It is used in Poisson (or
negative binomial) models to model rates. see **supplementary information for details).**
To account for potential spatial autocorrelation, we corrected the analyses by using an
exponential variogram.

We performed the statistical analysis in two stages. The trends in the entire city were
explored using all available data (i.e., survey data from 50 statistical areas and resident
reports from 61 statistical areas). To discern the effect of TNR intensity on the cat
population dynamics, we compared the cat population trends between the statistical areas
with the highest to the lowest neutering intensity at the end of the mixed-intervention
phase (designated as group-level analysis). To accomplish this, we divided the surveyed
statistical areas into quartiles according to the observed neutering percentage at the end of
the mixed-intervention phase. We compared two groups: Group-1 (n=12) consisted of the
statistical areas with the lowest quartile of neutering percentage at this stage (≤30%), and
Group-2 (n=15) consisted of the highest quartile of neutering percentage at this stage
(≥75% with four statistical areas roughly tied at this threshold).

First, the effects of human-related factors (predictor variables) on the various cat
population parameters in the entire city and their pairwise interactions with neutering
were tested by GLM (outcome variables: surveyed cat and kitten counts, and cat carcass
and reproduction reports). As we found that human population density and the volume of
accessed waste bins per resident were the main human-related predictors of most cat
population parameters, these were used in the following group-level analyses as
covariates. To test the difference of cat population outcome variables between the annual
surveys in the entire city, we performed GLMM. The group-level trend analyses were
performed separately for the survey and resident report data. In both analyses, the effects
of the treatment group, study phase, and their interaction (predictor variables) on the cat
population parameters (outcome variables, as detailed above, plus queen counts and
kitten-to queen ratio) were explored by GLMM. However, in the analysis of the resident

report data, the year was added as a covariate in order to realize the high resolution of this
data type. The specific modeling approach and statistical software used in each analysis
are detailed in the **supplementary information**.

To visualize the temporal trends of the high-resolution resident report data (cat carcass
and reproduction reports) collected both over the entire city or at the group level, we used
time-series analysis (Seasonal and Trend decomposition using Loess (STL) method (82)).
Through this method, we decomposed the monthly reports of the entire city (61statistical
areas) into three compartments: the general trend, cyclic changes (designated as
seasonality), and residual variability (remainder).

To confirm the construct validity of resident report data, we examined the correlation
between reproduction reports and the survey-derived kitten counts. We aggregated the
data to the neighborhood level and calculated the Pearson correlation coefficient of the
log-transformed kitten counts + 0.5 per transect length and the log-transformed
reproduction reports per the overall street length in each neighborhood in each year.
(**Figure S6**).

Acknowledgments

The authors are thankful to Y. Even-Zor, head of the municipal veterinary services, E.
Gian, head of the municipal GIS department, S. Keidar, principal at the maintenance
department, I. Ashkenazy from the municipal call center and E. Levi, head of the
municipal information and research center at the city of Rishon-LeZion.

References

1.   Lowe S, Browne M, Boudjelas S, & De Poorter M (2000) *100 of the world's worst*
invasive alien species: a selection from the global invasive species database (Invasive
Species Specialist Group Auckland, New Zealand).
2.   Feldman EC & Nelson RW (1996) Canine and feline endocrinology and reproduction. in
Ch. 31 Feline reproduction (W. B. Saunders Co., Philadelphia, PA 19106; USA), pp 741-
766.
3.   England GC & Heimendahl Av (2010) *BSAVA manual of canine and feline reproduction*
and neonatology (British Small Animal Veterinary Association).
4.   Scott KC, Levy JK, & Crawford PC (2002) Characteristics of free-roaming cats evaluated
in a trap-neuter-return program. *Journal of the American Veterinary Medical Association*
221(8):1136-1138.
5.   Serpell JA (2000) Domestication and history of the cat. *The domestic cat: The biology of*
its behaviour 179.
6.   Loss SR & Marra PP (2017) Population impacts of free☐ranging domestic cats on
mainland vertebrates. *Frontiers in Ecology and the Environment* 15(9):502-509.
7.   Gerhold RW & Jessup DA (2013) Zoonotic diseases associated with free-roaming cats.
Zoonoses and Public Health 60(3):189-195.
8.   Dabritz HA & Conrad PA (2010) Cats and Toxoplasma: implications for public health.
Zoonoses and Public Health 57(1):34-52.

9.   Slater MR*, et al.* (2008) Free-roaming dogs and cats in central Italy: Public perceptions of the problem. *Preventive Veterinary Medicine* 84(1–2):27-47.

10.   Gunther I, Raz T, Berke O, & Klement E (2015) Nuisances and welfare of free-roaming cats in urban settings and their association with cat reproduction. *Preventive Veterinary Medicine* 119(3–4):203-210.

11.   Medina FM*, et al.* (2011) A global review of the impacts of invasive cats on island endangered vertebrates. *Global Change Biology* 17(11):3503-3510.

12.   Rowan AN, Kartal T, & Hadidian J (2019) Cat demographics & impact on wildlife in the USA, the UK, Australia and New Zealand: facts and values. *Journal of Applied Animal Ethics Research* 2(1):7-37.

13.   Lepczyk CA, Lohr CA, & Duffy DC (2015) A review of cat behavior in relation to disease risk and management options. *Applied Animal Behaviour Science* 173:29-39.

14.   Bester M*, et al.* (2002) A review of the successful eradication of feral cats from sub-Antarctic Marion Island, Southern Indian Ocean. *South African Journal of Wildlife Research-24-month delayed open access* 32(1):65-73.

15.   Parkes J & Murphy E (2003) Management of introduced mammals in New Zealand. *New Zealand Journal of Zoology* 30(4):335-359.

16.   Ramsey DSL, Parkes JP, Will D, Hanson CC, & Campbell KJ (2011) Quantifying the success of feral cat eradication, San Nicolas Island, California. *New Zealand Journal of Ecology* 35(2):163-173.

17.   Nogales M*, et al.* (2004) A review of feral cat eradication on islands. *Conservation Biology* 18(2):310-319.

18.   Kennedy B, Cumming B, & Brown WY (2020) Global strategies for population management of domestic cats (Felis catus): a systematic review to inform best practice management for remote Indigenous communities in Australia. *Animals* 10(4):663.

19.   Calver MC, Crawford HM, & Fleming PA (2020) Response to Wolf et al.: Furthering debate over the suitability of trap-neuter-return for stray cat management. *Animals* 10(2):362.

20.   Schmidt PM, Swannack TM, Lopez RR, & Slater MR (2009) Evaluation of euthanasia and trap-neuter-return (TNR) programs in managing free-roaming cat populations. *Wildlife Research* 36(2):117-125.

21.   Hein S & Jacob J (2019) Population recovery of a common vole population (Microtus arvalis) after population collapse. *Pest Management Science* 75(4):908-914.

22.   Shilova S & Tchabovsky A (2009) Population response of rodents to control with rodenticides. *Current Zoology* 55(2):81-91.

23.   Williams CK, Parer I, Coman B, Burley J, & Braysher M (1995) Managing vertebrate pests: rabbits.

24.   Lazenby BT, Mooney NJ, & Dickman CR (2015) Effects of low-level culling of feral cats in open populations: a case study from the forests of southern Tasmania. *Wildlife Research* 41(5):407-420.

25.   Palmas P*, et al.* (2020) Rapid recolonisation of feral cats following intensive culling in a semi-isolated context. *NeoBiota* 63:177-200.

26.   Baker PJ & Harris S (2006) Does culling reduce fox (Vulpes vulpes) density in commercial forests in Wales, UK? *European Journal of Wildlife Research* 52(2):99-108.

27.   Ransom JI, Powers JG, Hobbs NT, & Baker DL (2014) Ecological feedbacks can reduce population□level efficacy of wildlife fertility control. *Journal of Applied Ecology* 51(1):259-269.

610   28.   McCarthy RJ, Levine SH, & Reed JM (2013) Estimation of effectiveness of three
611   methods of feral cat population control by use of a simulation model. *Journal of the*
612   *American Veterinary Medical Association* 243(4):502-511.
613   29.   Loyd KAT & DeVore JL (2010) An evaluation of feral cat management options using a
614   decision analysis network. *Ecology and Society* 15(4):art 10-art 10.
615   30.   Miller PS*, et al.* (2014) Simulating free-roaming cat population management options in
616   open demographic environments.
617   31.   Andersen MC, Martin BJ, & Roemer GW (2004) Use of matrix population models to
618   estimate the efficacy of euthanasia versus trap-neuter-return for management of free-
619   roaming cats. *Journal of the American Veterinary Medical Association* 225(12):1871-
620   1876.
621   32.   Denny EA & Dickman C (2010) Review of cat ecology and management strategies in
622   Australia. *Invasive Animals Cooperative Research Centre, Canberra*.
623   33.   Longcore T, Rich C, & Sullivan LM (2009) Critical assessment of claims regarding
624   management of feral cats by trap-neuter-return. *Conservation Biology* 23(4):887-894.
625   34.   Boone JD*, et al.* (2019) A long-term lens: Cumulative impacts of free-roaming cat
626   management strategy and intensity on preventable cat mortalities. *Frontiers in Veterinary*
627   *Science* 6:238.
628   35.   Wolf PJ & Schaffner JE (2019) The road to TNR: Examining Trap-Neuter-Return
629   through the lens of our evolving ethics. *Frontiers in Veterinary Science* 5:341.
630   36.   Wolf PJ & Hamilton F (2020) Managing free-roaming cats in US cities: An object lesson
631   in public policy and citizen action. *Journal of Urban Affairs*:1-22.
632   37.   Neville PF & Remfry J (1984) Effect of neutering on two groups of feral cats. *Veterinary*
633   *Record* 114(18):447-450.
634   38.   Hughes KL & Slater MR (2002) Implementation of a feral cat management program on a
635   university campus. *Journal of Applied Animal Welfare Science* 5(1):15-28.
636   39.   Hughes KL, Slater MR, & Haller L (2002) The effects of implementing a feral cat
637   spay/neuter program in a Florida county animal control service. *Journal of Applied*
638   *Animal Welfare Science* 5(4):285-298.
639   40.   Levy JK, Gale DW, & Gale LA (2003) Evaluation of the effect of a long-term trap-
640   neuter-return and adoption program on a free-roaming cat population. *Journal of the*
641   *American Veterinary Medical Association* 222(1):42-46.
642   41.   Algar D, Hilmer S, Nickels D, & Nickels A (2011) Successful domestic cat neutering:
643   first step towards eradicating cats on Christmas Island for wildlife protection. *Ecological*
644   *Management & Restoration* 12(2):93-101.
645   42.   Levy J, Isaza N, & Scott K (2014) Effect of high-impact targeted trap-neuter-return and
646   adoption of community cats on cat intake to a shelter. *The Veterinary Journal*
647   201(3):269-274.
648   43.   Swarbrick H & Rand J (2018) Application of a protocol based on Trap-Neuter-Return
649   (TNR) to manage unowned urban cats on an Australian University Campus. *Animals*
650   8(5):77.
651   44.   Kreisler RE, Cornell HN, & Levy JK (2019) Decrease in Population and Increase in
652   Welfare of Community Cats in a Twenty-Three Year Trap-Neuter-Return Program in
653   Key Largo, FL: the ORCAT Program. *Frontiers in Veterinary Science* 6:1-14.
654   45.   Jones AL & Downs CT (2011) Managing feral cats on a university's campuses: how
655   many are there and is sterilization having an effect? *Journal of Applied Animal Welfare*
656   *Science* 14(4):304-320.
657   46.   Zito S, Aguilar G, Vigeant S, & Dale A (2018) Assessment of a targeted trap-neuter-
658   return pilot study in Auckland, New Zealand. *Animals* 8(5):73.

47.   Nutter F (2005) Evaluation of a trap-neuter-return management program for feral cat
colonies: population dynamics, home ranges, and potentially zoonotic diseases. . *PhD*
thesis in Comparative Biomedical Science, North Carolina State University, Raleigh, NC.
48.   Natoli E*, et al.* (2006) Management of feral domestic cats in the urban environment of
Rome (Italy). *Preventive Veterinary Medicine* 77(3/4):180-185.
49.   Spehar DD & Wolf PJ (2017) An examination of an iconic trap-neuter-return program:
The Newburyport, Massachusetts case study. *Animals* 7(11):81.
50.   Spehar DD & Wolf PJ (2019) Back to school: An updated evaluation of the effectiveness
of a Long-Term Trap-Neuter-Return program on a university's free-roaming cat
population. *Animals* 9(10):768.
51.   Tan K, Rand J, & Morton J (2017) Trap-Neuter-Return Activities in Urban Stray Cat
Colonies in Australia. *Animals* 7(6):46.
52.   Spehar DD & Wolf PJ (2018) A case study in citizen science: The effectiveness of a trap-
neuter-return program in a Chicago neighborhood. *Animals* 8(1):14.
53.   Gunther I, Finkler H, & Terkel J (2011) Demographic differences between urban feeding
groups of neutered and sexually intact free-roaming cats following a trap-neuter-return
procedure. *Journal of the American Veterinary Medical Association* 238(9):1134-1140.
54.   Castillo D & Clarke AL (2003) Trap/neuter/release methods ineffective in controlling
domestic cat "colonies" on public lands. *Natural Areas Journal* 23(3):247-253.
55.   Kilgour RJ*, et al.* (2017) Estimating free-roaming cat populations and the effects of one
year Trap-Neuter-Return management effort in a highly urban area. *Urban Ecosystems*
20(1):207-216.
56.   Mannhart T (2007) A catch-neuter-release project for free-roaming dogs and cats in
Rhodes, Greece: Problem analysis and effectiveness of the strategy. PhD thesis
(University of Bern.).
57.   Natoli E*, et al.* (2019) Evaluation of Unowned Domestic Cat Management in the Urban
Environment of Rome After 30 Years of Implementation of the No-Kill Policy (National
and Regional Laws). *Frontiers in veterinary science* 6:31.
58.   Bissonnette V, Lussier B, Doizé B, & Arsenault J (2018) Impact of a trap-neuter-return
event on the size of free-roaming cat colonies around barns and stables in Quebec: A
randomized controlled trial. *Canadian Journal of Veterinary Research* 82(3):192-197.
59.   Boone JD*, et al.* (2014) Improving and evaluating trap-neuter-return (TNR) management
for outdoor cats on the human landscape. *Proceedings of the Vertebrate Pest Conference*.
60.   Foley P, Foley JE, Levy JK, & Paik T (2005) Analysis of the impact of trap-neuter-return
programs on populations of feral cats. *Journal of the American Veterinary Medical*
Association 227(11):1775-1781.
61.   Mirmovitch V (1995) Spatial organisation of urban feral cats (Felis catus) in Jerusalem.
Wildlife Research 22(3):299-310.
62.   Finkler H, Hatna E, & Terkel J (2011) The influence of neighbourhood socio-
demographic factors on densities of free-roaming cat populations in an urban ecosystem
in Israel. *Wildlife Research* 38(3):235-243.
63.   Nutter FB, Levine JF, & Stoskopf MK (2004) Reproductive capacity of free-roaming
domestic cats and kitten survival rate. *Journal of the American Veterinary Medical*
Association 225(9):1399-1402.
64.   Gunther I, Raz T, & Klement E (2018) Association of neutering with health and welfare
of urban free-roaming cat population in Israel, during 2012-2014. *Preventive Veterinary*
Medicine.

65.   Murray JK, Skillings E, & Gruffydd-Jones TJ (2008) A study of risk factors for cat
707   mortality in adoption centres of a UK cat charity. *Journal of Feline Medicine & Surgery*
708   10(4):338-345.
709   66.   Childs JE & Ross L (1986) Urban cats: characteristics and estimation of mortality due to
710   motor vehicles. *American Journal of Veterinary Research* 47(7):1643-1648.
711   67.   Chambers LK, Singleton GR, & Hinds L (1999) Fertility control of wild mouse
712   populations: the effects of hormonal competence and an imposed level of sterility.
713   *Wildlife Research* 26(5):579-591.
714   68.   Twigg LE*, et al.* (2000) Effects of surgically imposed sterility on free☐ranging rabbit
715   populations. *Journal of Applied Ecology* 37(1):16-39.
716   69.   Turner A & Kirkpatrick JF (2002) Effects of immunocontraception on population,
717   longevity and body condition in wild mares (Equus caballus). *Reprod Suppl* 60:187-195.
718   70.   Williams CK*, et al.* (2007) Population responses to sterility imposed on female European
719   rabbits. *Journal of Applied Ecology*:291-301.
720   71.   Smith LM*, et al.* (2019) The Effectiveness of Dog Population Management: A Systematic
721   Review. *Animals* 9(12):1020.
722   72.   Scott KC, Levy JK, Gorman SP, & Newell SM (2002) Body condition of feral cats and
723   the effect of neutering. *Journal of Applied Animal Welfare Science* 5(3):203-213.
724   73.   Finkler H, Gunther I, & Terkel J (2011) Behavioral differences between urban feeding
725   groups of neutered and sexually intact free-roaming cats following a trap-neuter-return
726   procedure. *Journal of the American Veterinary Medical Association* 238(9):1141-1149.
727   74.   Tennent JK, Downs CT, Wald DM, & Watson HK (2010) Public perceptions of feral cats
728   within an urban conservancy on a campus of the University of KwaZulu-Natal. *South
729   African Journal of Wildlife Research* 40(1):16-26.
730   75.   Ireland T & Neilan RM (2016) A spatial agent-based model of feral cats and analysis of
731   population and nuisance controls. *Ecological Modelling* 337:123-136.
732   76.   Boone JD (2015) Better trap–neuter–return for free-roaming cats Using models and
733   monitoring to improve population management. *Journal of Feline Medicine and Surgery*
734   17(9):800-807.
735   77.   Bruce SJ*, et al.* (2019) Predation and risk behaviors of free-roaming owned cats in
736   Auckland, New Zealand via the use of animal-borne cameras. *Frontiers in Veterinary
737   Science* 6.
738   78.   Loyd KAT, Hernandez SM, Carroll JP, Abernathy KJ, & Marshall GJ (2013) Quantifying
739   free-roaming domestic cat predation using animal-borne video cameras. *Biological
740   Conservation* 160:183-189.
741   79.   Cecchetti M, Crowley SL, Goodwin CE, & McDonald RA (2021) Provision of High
742   Meat Content Food and Object Play Reduce Predation of Wild Animals by Domestic
743   Cats Felis catus. *Current Biology* 31(5):1107-1111. e1105.
744   80.   Nelson S, Evans A, & Bradbury R (2005) The efficacy of collar-mounted devices in
745   reducing the rate of predation of wildlife by domestic cats. *Applied Animal Behaviour
746   Science* 94(3-4):273-285.
747   81.   Gunther I, Azriel L, Wolf H, Raz T, & Klement E (2020) An accessible scheme for
748   monitoring free☐roaming cat population trends. *Ecology and Evolution* 10:1288-1298.
749   82.   Cleveland RB, Cleveland WS, McRae JE, & Terpenning I (1990) STL: a seasonal-trend
750   decomposition. *Journal of Official Statistics* 6(1):3-73.

751
752

Figures and Tables

Figure 1: The experimental design was set to assess long-term TNR effectiveness over time and space. The study period was divided into three phases: 1st phase – pre-intervention period - between 2007 to the end of 2009; 2nd phase -mixed-intervention - allocating TNR to half of the city neighborhoods between the end of 2009 to end of 2014; 3rd phase, full-intervention period, allocating TNR to the entire city between the end of 2014 and the end of 2018. Dots present locations of TNR in the city area. The study area was composed of 50 statistical areas (presented as polygons in the upper map; see also Fig. S1), 24 of which are termed high-TNR, as they were treated in both TNR phases, and the other 26, termed low-TNR, as they were mainly treated during the full-intervention phase. From these high- and low-TNR statistical areas, 15 and 12 areas with the highest and lowest neutering percentage at the 2014 survey, respectively, were served as the main study units (Group-1 and -2). Survey years are marked by an asterisk.

Figure 2: Annual trends (geometric mean±$CI_{95\%}$) of (**A**) neutering percentage, (**B**) free-roaming cat counts, (**C**) kitten counts, (**D**) queen counts, and (**E**) kitten-to-queen ratio in the city of Rishon-LeZion (black, n = 50 statistical areas), and in Group-1 (turquoise, lowest neutering intensity, n=12 statistical areas) and Group-2 (purple, highest neutering intensity, n=15 statistical areas). Observations were performed in 2012–2014 and 2018. Superscript letters represent years, where observations differ.

A

B

Figure 3: Municipal TNR efforts and time series decomposition plots of the monthly carcass
and cat-reproduction citizen reports in the entire city of Rishon-Lezion, between 2007-2018 (the
crude data is divided into three compartments: trend, seasonality and remainder) . (**A**) Monthly-
based time series plots of municipal TNR actions (n = 22,144), cat carcass reports (n = 36,544)
and reproduction reports (n = 12,217) from January 2007 to December 2018 across the entire city
(61 statistical areas). (**B**) Time series decomposition plots of carcass and reproduction reports
linked with cumulative TNR percentage. The 1st phase refers to the pre-intervention period
(before the initiation of the TNR program), the 2nd phase refers to the mixed-intervention period
(applying TNR in half of the city), and the 3rd phase refers to the full-intervention period
(applying TNR in the entire city).

Figure 4: Municipal TNR efforts and time series decomposition plots of the monthly carcass and cat-reproduction citizen reports in group 1 and group 2 statistical areas in the city of Rishon-Lezion, between 2007-2018 (the crude data is divided into three compartments: trend, seasonality and remainder) of (**A**) cat carcass reports and (**B**) cat reproduction reports in Group-1 (lowest neutering intensity, n = 12) compared to Group-2 (highest neutering intensity, n = 15). The 1st phase refers to the pre-intervention period (before the initiation of the TNR program), the 2nd phase refers to the mixed-intervention period (applying TNR primarily in Group-2), and the 3rd phase refers to the full-intervention period (applying TNR in both groups).

795 **Table 1:** Annual counts of observed cats, neutered cats, neutering percentage, kittens,
796 queens, and kitten-to-queen ratio in the low-TNR (n=26), high-TNR statistical areas
797 (n=24), Group-1 (lowest neutering intensity, n=12) and Group-2 (highest neutering
798 intensity, n=15).

Variable	Year	Low-TNR	High-TNR	Group-1	Group-2	Overall
Cats	2012	1726	1485	675	938	3211
	2013	1880	1443	770	906	3233
	2014	2219	1843	1000	1082	4062
	2018	1705	1417	688	765	3122
Neutered	2012	599	918	145	620	1517
	2013	644	1003	136	684	1647
	2014	742	1312	189	866	2054
	2018	1274	982	477	590	2256
Neutering percentage	2012	34.7	61.8	21.5	66.1	47.2
	2013	34.3	69.5	17.7	75.5	50.9
	2014	33.4	71.2	18.9	80.0	50.6
	2018	74.7	69.3	69.3	77.1	72.3
Kittens	2012	291	144	111	96	435
	2013	228	90	108	57	318
	2014	292	104	162	35	396
	2018	165	172	72	79	337
Queens	2012	273	114	123	61	387
	2013	308	103	158	49	411
	2014	400	113	226	50	513
	2018	91	103	43	36	194
Kitten-to-queen ratio	2012	1.07	1.26	0.90	1.57	1.12
	2013	0.74	0.87	0.68	1.16	0.77
	2014	0.73	0.92	0.72	0.70	0.77
	2018	1.81	1.67	1.67	2.19	1.74

799
800

Table 2. Annual trends of the cat and kitten counts, kitten-to-queen ratio (surveyed
during the mixed- and full-intervention phases), and cat carcass and reproduction data
(received from resident reports during all the three study phases). The trends are depicted
for Group-1, (lowest neutering intensity, n=12), and Group-2, (highest neutering
intensity, n=15). All the models are adjusted for human population density and accessed
waste bins.

Population parameter	Phase	Measurement period	Annual trend for Group-1 ($CI_{95\%}$) [%]	Annual trend for Group-2 ($CI_{95\%}$) [%]	P-value
Cat counts	2nd phase	2012-2014	20.72 (8.19 to 34.71) [a1*]	4.02 (-5.62 to 14.65) [a2]	0.046
	3rd phase	2014-2018	-7.95 (-13.09 to -2.50) [b1]	-7.83 (-12.29 to -3.15) [b2]	0.973
Kitten counts	2nd phase	2012-2014	20.20 (-11.31 to 62.91) [a1]	-42.39 (-58.19 to -20.60) [a2]	0.001
	3rd phase	2014-2018	-14.14 (-27.00 to 1.00) [b1]	21.37 (3.20 to 42.74) [b2]	0.003
Queen counts	2nd phase	2012-2014	40.40 (13.31 to 73.97) [a1]	-11.21 (-30.92 to 14.12) [a2]	0.007
	3rd phase	2014-2018	-38.34 (-46.47 to -28.98) [b1]	-8.76 (-20.50 to 4.71) [a2]	<0.001
Kitten-to-queen ratio	2nd phase	2012-2014	-7.72 (-36.10 to 33.28) [a1]	-28.48 (-50.60 to 3.55) [a2]	0.338
	3rd phase	2014-2018	29.88 (7.40 to 57.08) [a1]	23.35 (2.26 to 48.28) [b2]	0.706
Carcass reports	1st phase	2007-2009	7.96 (-1.44 to 18.26) [a1]	6.01 (-1.62 to 14.23) [a2]	0.761
	2nd phase	2010-2014	0.88 (-3.04 to 4.95) [a1]	-12.02 (-15.17 to -8.75) [b2]	<0.001
	3rd phase	2015-2018	-13.59 (-19.02 to -7.79) [b1]	0.57 (-4.82 to 6.27) [a2]	<0.001
Reproduction reports	1st phase	2007-2009	8.02 (-8.03 to 26.87) [a1]	-5.70 (-18.44 to 9.04) [a2]	0.219
	2nd phase	2010-2014	3.89 (-2.95 to 11.21) [a1]	-24.50 (-29.94 to -18.63) [b2]	<0.001
	3rd phase	2015-2018	-7.23 (-17.18 to 3.91) [a1]	22.56 (9.96 to 36.59) [c2]	<0.001

807
808   * The superscript letters represent comparisons between pairs of phases <u>within each group</u> of
809   statistical areas (pairs with insufficient evidence for a difference are labeled with the same
810   superscript letter).