1

## Main Manuscript for

Reduction of free-roaming cat population requires high-intensity neutering in spatial contiguity to mitigate compensatory effects

Idit Gunther[a], Hadas Hawlena[b], Lior Azriel[a], Dan Gibor[a], Olaf Berke[c], and Eyal Klement[a*]

[a] Koret School of Veterinary Medicine, The Robert H. Smith Faculty of Agriculture, Food & Environment, The Hebrew University of Jerusalem, Rehovot 76100, Israel

[b] Mitrani Department of Desert Ecology, Swiss Institute for Dryland Environmental and Energy Research, The Jacob Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Sde-Boker 8499000, Israel

[c] Department of Population Medicine, Ontario Veterinary College, University of Guelph, Ontario N1G 2W1, Canada

* Corresponding author:
Eyal Klement
Tel: 972-54-6766004, email: eyal.klement@mail.huji.ac.il

**Author Contributions:**

Conception and design: Gunther I., Klement E.
Acquisition of data: Gunther I., Azriel L., Gibor D.
Interpretation of data and analyzes: Gunther I., Hawlena H., Berke O., Klement E.
Drafting the manuscript: Gunther I., Hawlena H., Berke O., Klement E.
Review and confirmation of manuscript: all authors.

**Competing Interest Statement:** All the authors declare that they do not have competing interests in this manuscript.

**Classification:** Biological sciences/ecology

**Keywords:** Population management, TNR, Free-Roaming Cats, Fertility control, Population compensatory mechanisms.

**Abstract**

When free-roaming in natural areas, the domestic cat (*Felis silvestris catus*) is ranked high among the most destructive alien species. Near human vacancy, it might risk humans, impair sanitation, and suffer from poor welfare. Cats' popularity as companion animals complicates their population control. Thus, culling is often replaced by a fertility control method called "Trap-Neuter-Return/Release (TNR)," which is considered more humane. Despite the extensive application of TNR, a long-term controlled study was never performed to test its effectiveness. We present a uniquely designed controlled field experiment for examining TNR effectiveness. The study was performed over a twelve-year period, divided into pre-intervention, mixed- and full-intervention phases, and spanned a 20Km$^2$ urban area. Trends of cat, queen, and kitten counts, cat reproduction, and carcass reports were compared among study phases and areas with different neutering intensities. The cat population increased during the first two study phases and did not decline in highly neutered populations, presumably due to cat immigration. Expansion of high-intensity neutering to the entire city in the full-intervention phase (>70% neutering percentage) reversed cat population growth, reaching an annual ca. 7% reduction. This population reduction was limited by a rebound increase in cat reproduction and longevity. We conclude that cat population management by TNR should be performed in high-intensity, continuously, and in geographic contiguity to enable population reduction. To enhance management effectiveness and mitigate compensatory effects, we recommend further evaluating an integrated strategy that combines TNR with complementary methods (e.g., vital resource regulation, ill cat euthanasia, and adoption).

**Significance Statement**

Though popular companion animals, domestic cats pose numerous problems when free-roaming, including predation of wildlife, hazards to humans, impaired sanitation, and a decrease in their welfare. Thus, managing their populations is essential. The Trap-Neuter-Return method (TNR, capturing, sterilizing, returning/releasing) is widely employed for managing cat populations. However, its use is under continuous debate due to the lack of long-term controlled evidence. We examined the outcomes of high-intensity TNR by performing a twelve-year controlled field experiment, the largest to date. Neutering over 70% of the cats caused population decline when applied over contiguous areas. However, it was limited by a rebound increase in reproduction and survival. These findings provide a robust quantification of the limitations and the long-term effectiveness of TNR.

**Main Text**

**Introduction**

As a generalist predator, the domestic cat (*Felis silvestris catus*) has been listed among the world's 100 worst non-native species (1). The domestic cat is a fast life-history species (i.e., early maturation, small body size, rapid reproduction) (2-4), which has been distributed around the globe mainly as a pet (5). Cats have been known to form non-domiciliary and often human-independent populations known as free-roaming cats, which tend to cause adverse environmental effects (1, 6-11). The most prominent ecological negative impact of cats is on islands. They are considered responsible for at least 14% of global bird, mammal, and reptile extinctions and are the principal threat to almost 8% of critically endangered birds, mammals, and reptiles (11). Moreover, cats also have a significant ecological effect on the mainland due to direct predation, the transmission of diseases to other species, fear-related effects, and the alteration of demographic processes such as source-sink dynamics (6). As a result of their potential to transmit certain zoonotic diseases and injure humans due to aggressive behavior, free-roaming cats constitute a hazard to public health (7, 8). They might also cause a nuisance to humans, mainly by impairing sanitation (9, 10). Over the past decades, increased urbanization has further enhanced the formation of large cat populations and, hence, increased human-cat interactions (12).

The adverse effects caused by free-roaming cats have increased the motivation to artificially manage their populations, aiming either to diminish their related nuisances or to preserve natural ecosystems (13-18). Animal population management is based on two strategies: resource limitation in the habitat or actions applied to individuals. The latter strategy can be further divided into culling and fertility control methods. While culling aims to increase mortality above the natural rate, fertility control aims to decrease reproduction below the natural rate (19). Culling was successful in eradicating cat populations on islands (13, 15, 16). However, there are several examples of the failure of culling to accomplish population control in fast life-history species, such as voles (20), mice, rats, jirds (21), rabbits (22), cats (23, 24), and foxes (25). In contrast, information on the efficacy of fertility control on vertebrate populations is scarce (26). Specifically for free-roaming cats, theoretical studies predicted that culling performs better than Trap-Neuter-Return (TNR), a common method for controlling cat fertility (19, 27-30). However, despite these predictions, the TNR method has been progressively implemented in cat populations over widespread areas instead of culling, mainly due to moral considerations and public opinion (17, 31-35).

Several field studies aimed at determining the association of TNR management programs with cat population dynamics demonstrated a decrease in population growth indicators (36-51). Others showed stabilization or even an increase in population growth (52-57). These inconsistent results might stem from differences in management duration and efforts, the examined populations (e.g., closed versus open populations, small versus large-scale populations), or the study environment. The inference of the overall long-term consequences of TNR is further limited due to: absence of control (36-40, 42-44, 47-51,

53, 55, 56), combining TNR with other control tactics (i.e., adoption and euthanasia of
clinically ill or retrovirus-positive cats) (36-45, 48-51), short-term follow-up (36, 37, 41,
44, 45, 50, 52-55, 57), small sample size (36, 37, 39, 42, 44, 46, 48, 49, 51-53, 57),
relying on indirect indices of population growth (37, 38, 41, 45, 48, 50, 51, 56), and
examining populations in secluded areas (40, 43). Most importantly, to date, a controlled
study was never performed to test the long-term effectiveness of TNR.

Boon et al. (58) called for a long-term and large-scale study in a cat metapopulation to
close the knowledge gap and overcome shortcomings of previous studies. Here we
present the results of a 12-year longitudinal large-scale experiment. We assessed the
effect of sterilization (i.e., spaying/neutering) on the long-term temporal and spatial
dynamics of free-roaming cat metapopulations. While adjusting for environmental
factors, we compared cat population dynamics before and at partial and full
implementation of a TNR program. We found that population compensatory mechanisms
may limit the magnitude of the neutering effect. Thus, TNR should be performed
continuously, at high intensity, and in spatial contiguity to enable long-term cat
population regulation.

124
125 **Results**
126
*Neutering intensity across statistical areas and study phases*

The study was set to assess long-term TNR effectiveness over time and space (**Figures 1,**
**S1**). In the 1st (pre-intervention) phase, neutering was not applied. In the 2nd (mixed-
intervention) phase, high-intensity neutering was applied in about half of the city, and in
the 3rd (full-intervention) phase, TNR was applied in the entire city. We confirmed the
neutering status in 96% of the surveyed cats by ear-mark detection, counting both
unneutered (un-marked) and neutered (marked) cats, along with kittens. Repeated annual
surveys were performed during September and October of 2012, 2013, 2014, and 2018.
Overall, in the 50 surveyed statistical areas, 13,718 cat-observations were documented
during the study period, of which 1,486 were kittens. The phase- and group-specific
neutering percentages corroborated the respective TNR intensity (**Table 1**, **Table S1**).
For a visual perspective of the municipal TNR intensity across statistical areas and study
phases, see **Figure S2**.

At the end of the mixed-intervention phase (end of 2014), the overall neutering
percentages in the entire surveyed area, the low TNR (n=26) and high TNR (n=24)
statistical areas, were 51, 33, and 71%, respectively (**Table 1**). To compare the effect of
the highest-intensity neutering to the lowest-intensity, we further divided the statistical
areas into quartiles according to the observed neutering percentage in that year (2014).
Group-1 (n=12) represents the lowest neutering intensity and consists of the statistical
areas with neutering percentage at the lowest quartile. Group-2 (n=15) represents the
highest neutering intensity and consists of the statistical areas with neutering percentage
at the highest quartile (see **Figure S1** for the detailed spatial study design). At the end of
the mixed-intervention phase, the overall neutering percentages in Group-1 and Group-2

were 19 and 80%, respectively. At the end of the full-intervention phase (end of 2018), the neutering percentage in the entire area was 72%, and in the low TNR, high TNR, Group-1, and Group-2, the overall percentages were 75, 69, 69, and 77%, respectively (**Table 1, Figure 2A**).

*Environmental human-related factors affecting cat population dynamics*

We modeled year-based cross-sectional data (adjusting for neutering percentage) to screen for potential human-related factors that might influence cat population dynamics. As expected, we found a consistent negative association between neutering percentage and kitten counts (**Table S3**). Interestingly, in some years, the cat counts were positively associated with the neutering percentage (i.e., 2012 and 2018), indicating possibly higher TNR efforts in areas with higher, rather than lower, cat densities. In addition, cat population size parameters, cat counts, and resident reports of cat carcasses and reproduction were positively correlated with the human population density (**Table S3**). Though less consistent, the volume of accessed-waste-bins was also positively correlated with some cat population parameters. Therefore, we adjusted for human population density and accessed-waste-bins by including them in the analysis of neutering association with cat population trends. The other tested human-related factors were not consistently associated with the cat population parameters. We did not find a consistent spatial correlation in these models, and therefore it was not further included in the trend models.

*Survey results: Cat and kitten counts across statistical areas and study phases*

During the mixed-intervention phase (2012-2014), we found an overall increase of 26.5% in the cat counts in the surveyed area. Therein, we found an average annual growth of 20.7% in Group-1, which was exposed to the lowest neutering intensity (**Table 1, Figure 2B**). At the same time, the cat counts remained stable in Group-2, which was exposed to the highest neutering intensity (**Table 2**). In contrast, during the full-intervention phase, we found an overall reversal of population growth, with a 23.1% reduction in the cat counts in the entire surveyed area. During this phase, the annual decrease in the two treatment groups was similar (ca. -7% annually; **Figure 2B**).

As opposed to cats, we did not find a significant reduction in the kitten counts in the entire surveyed area (**Table 1, Figure 2C**). This overall stability was a result of opposite trends in the two treatment groups. As expected, during the mixed-intervention phase, kitten counts increased annually by 20.2% in Group-1 but decreased by 42.4% in Group-2. Whereas, during the full-intervention phase, these trends reversed, with a decrease of 14.1% in Group-1 and an increase of 21.4% in Group-2 (**Table 2**). The latter increase in kitten counts in Group-2, during the full-intervention phase, was unexpected. A possible explanation for this increase is a reduction in neutering intensity in some statistical areas in this group during the full-intervention phase. To examine this possibility, we reanalyzed the data of both groups, excluding statistical areas in which the neutering percentages at the end of the full-intervention phase were below 70%.

Here, too, we found the same trend reversal in kitten counts (**Table S4**). Therefore, a
decrease in neutering intensity in some statistical areas is unlikely to explain the observed
increase. An alternative explanation is that reduced agonistic behavior in neutered cats
enhanced compensatory mechanisms such as increased kitten survival or increased litter
size due to lower competition. Assuming this hypothesis was correct, we predicted to find
an overall increase in the kitten-to-queen ratio in the full-intervention study phase. We
indeed found a 2.25-fold increase during this phase (**Table 1, Figure 2E**). This increase
was significant in both treatment groups (**Table 2**). In Group-2, it occurred due to a
significant increase in the kitten number and stabilization of the queen number. In Group-
1, the queen number decreased annually by 38% shortly after introducing high-intensity
neutering, while a 14% insignificant decrease in the kitten number was observed (**Table
2**). No significant reduction in the kitten-to-queen ratio was observed in Group-2 during
the mixed-intervention period, possibly due to parallel immigration of intact females.

*Resident report results: Cat carcasses and reproduction events across statistical areas*
*and study phases*

We complemented the survey data analysis with those of resident reports on cat
reproduction events and cat carcasses. This dataset increased the study resolution in both
space (covering more statistical areas) and time (daily reports from 2007 to 2018). To
examine the construct validity of these reports, we tested the correlation of reproduction
reports with the surveyed kitten counts and found a strong correlation between the two
datasets (Pearson correlation coefficient=0.55, p<0.001, **Figure S6**).

In addition, to ensure that these reports represent trends unique to cats, we compared
them with the number of all available reports to the call center (after excluding cat-related
reports) and with the reports of all carcasses (excluding cats and including canines,
poultry, rodents, hyraxes, hedgehogs, equines, reptiles, boar and tortoises) . The trends of
these two latter report types showed a consistent increase during the study period (**Figure
S4**). Overall, this increased trend suggests that people's awareness or tendency to use the
call center increased with time. Unlike the general trends, the reports of cat carcasses and
reproduction were linked to the application of TNR. As with the general reports, both
showed a mild positive trend during the pre-intervention phase. However, this was
followed by a negative trend, from the onset of the mixed-intervention phase until one
year after its end (from 2011 to 2015) (**Figure 3**). The magnitude of this negative trend
was ca. 50%, starting at 325 and 120 and reaching a low of 200 and 60 of carcasses and
reproduction reports per month, respectively. Notably, during the mixed-intervention
phase, the number of carcass and reproduction reports decreased while the cat counts
increased (**Table 1**, **Figure 2**), indicating a reduction in cat reproduction and mortality.
One year after the onset of the full-intervention phase (2016-2018), the number of carcass
reports stabilized, but the number of reproduction reports showed an unexpected increase
(**Figure 3**). This increase suggests a possible rebound effect of TNR on reproduction.

Both report types exhibited a prominent seasonal pattern: carcasses peaked twice, in
May-June and in October, whereas reproduction reports peaked only once in April-May
(**Figure 3**). Unlike these reports, the general reports to the call center peaked in July-

August (**Figure S4**), showing that the seasonal pattern of carcass and reproduction
reports is specific for cats and did not represent human behavior tendency to refer to the
call center. While the seasonal pattern of cat reproduction reports was maintained
throughout the study, the seasonality of carcass reports gradually diminished from the
onset of TNR until it disappeared during the full-intervention phase (see 'crude data' in
**Figure 3b**). Thus, it implies that TNR resulted in a decrease in mortality and a decrease
in the seasonal (reproduction-linked) mortality pattern.

We next compared the carcass and reproduction report trends between the treatment
groups. Both trends coincided with TNR efforts (**Figure 4**, **Table 2**). In the pre-
intervention phase, the numbers of carcass and reproduction reports were stable. During
the mixed-intervention phase, both carcass and reproduction reports declined
considerably in Group-2 (-12% annually for carcass and -24.5% for reproduction
reports), while their numbers were stable in Group-1 (0.9% for carcass and 3.9% for
reproduction reports). The number of carcass reports decreased considerably in Group-1
(-13.6%) in the full-intervention phase, indicating a decrease in cat number and/or
mortality. Similar to kitten counts, in Group-2, a considerable rebound increase of
reproduction reports occurred during the full-intervention phase (22.6%). We found
similar trends when the analysis was repeated after excluding statistical areas in which
the neutering percentages at the end of the full-intervention phase were below 70%.
(**Figure S5, Table S4**).

**Discussion**
253
The current study is the first long-term controlled field experiment examining TNR
effectiveness in free-roaming cat populations. Using a unique experimental design in
which we manipulated neutering intensity across time and space, we distinguished
between short-term and long-term population consequences of TNR. We used two
independent data-collection methods to achieve high-resolution data—a repeated cross-
sectional annual survey and daily resident reports. Parameters of population dynamics
derived by both methods showed the same trends, suggesting that TNR can reduce
population size, but only when applied at high rates in spatial contiguity. Below, we
discuss the evidence suggesting processes that might challenge TNR effectiveness and
thus should be considered when applying it. These include compensatory effects, such as
increased cat survival and/or litter size, as well as possible occupancy of vacant niches by
immigrant cats. We then compare our results to those gathered by theoretical simulations,
discussing the TNR option compared to its alternatives, and end by offering directions for
further simulations and planning strategies.

*Maintaining high neutering intensity in spatial contiguity is required to mitigate*
*counteracting effects*

During the mixed-intervention phase, the cat population growth in Group-2 was
considerably lower than in Group-1. However, the cat population did not decline in
Group-2 despite maintaining a neutering percentage of 80% (**Figures 2A, B**), a number
previously suggested as sufficient for reducing the population size of free-roaming cats

(27, 59). We propose that during the mixed-intervention phase, the TNR effect was
counteracted by the immigration of intact cats from unneutered surrounding populations.
It was previously shown that sterilization-enhanced behavioral changes resulted in
increased immigration of free-roaming cats from low- into high-TNR regions (52). This
important population compensatory process was also documented in other managed
species (reviewed in Ransom et al. 2014 (26)). Theoretically, long-distance cat
immigration could have also occurred from adjacent cities. However, it was probably
negligible due to the small home range of urban cats (60, 61) and the study area
boundaries, enabling migration mainly from the north.

During the full-intervention phase, an average annual reduction of more than 7% was
observed in both groups. Our surveys further indicated that during the four years of this
phase, the cat population in the entire city declined by ca. 25%. It is possible that the cat
counts decreased in the entire city due to temporal effects unrelated to TNR. For
example, a change in vital resource availability (e.g., decreased feeding or change in
sanitary conditions), decreased abandonment, or increased adoption. However, this
possibility is unlikely for three reasons: First, the finding of cat counts reduction was
adjusted for changes in the environmental factors influencing cat population size (i.e.,
human population density and the per human accessible waste bins). Second, the survey
periods following each intervention change were not longer than 4–5 years. A marked
change in human behavior to opposite directions both spatially and temporally in such
short periods is unlikely. Third, abandonment and adoption in Israel constitute only a
negligible factor relative to the remarkable density of the free-roaming cats' population
(60, 61). The most probable alternative explanation for the cat population trend changes
between the phases is the expansion of high neutering intensity (>70%) to the entire city
during the full-intervention phase. This expansion may have reduced the effect of the
immigration of cats from low TNR to high TNR areas. We conclude that maintaining
high-intensity neutering in spatial contiguity is a prerequisite for reducing cat population
size.

*Limiting processes should be considered in TNR programs*

Along with the encouraging cat population decline, our findings support the occurrence
of several compensatory effects, limiting TNR effectiveness: **Reduced mortality**; Cat
carcass trend reversed from an increase during pre-intervention to a constant decrease
during the mixed-intervention phase. This decrease occurred despite growth in cat
population size at this phase. These opposite trends indicate that the reduced carcass
reports are due to reduced mortality following TNR. **Reduced reproduction-related**
**mortality**; The diminished seasonal pattern of carcass reports following increased
neutering intensity (see 'crude data' in Fig 3B and Fig 4b) indicates a decreased
reproduction-related mortality. It was previously shown that in the Northern hemisphere,
kittens present a seasonal appearance (4, 10, 52, 64) and experience higher morbidity and
mortality than adults (52, 62, 63). Moreover, the mortality of intact cats was higher than
neutered ones (64, 65), probably due to associated reproductive behavior. **Increased**
**fertility and reduced kitten mortality**; While the neutering percentage exceeded 72% in
the entire city during the full-intervention phase, a considerable increase in the kitten-to-

queen ratio was documented in both study groups. This increased ratio may indicate
either an elevation in kitten survival and/or a rise in litter size. This conclusion is
supported by a rise in cat reproduction reports in Group-2 during the full-intervention
phase. Increased fertility and decreased mortality of juveniles and adults were previously
reported to follow fertility control in domestic cats (33, 52, 62, 63) and other vertebrates
(26, 66-70). Both compensatory processes could result from higher food availability and
decreased resource competition following population decline (63, 71). In addition, they
could be caused by diminished agonistic behavior in neutered male cats (72). These
mechanisms, together with cat immigration, might limit the TNR-derived decrease in cat
population size.

*Comparison of our findings to the results of cat population simulation models*

Several theoretical simulations aimed to predict the long-term consequences of TNR and
provide specific management insights (19, 27, 29, 30). Of these, only two studies
examined the effect of TNR in an open population (19, 29). In these studies, population
reduction was achieved by implementing TNR, though the immigration of non-neutered
cats limited its effectiveness. This situation resembles the mixed-intervention phase in the
current study when untreated areas surrounded treated areas. Interestingly, in our study,
during this phase, the cat population did not decline. There are at least two possible
explanations for this discrepancy. First, we assume that similar to other population
dynamics, immigration is a density-dependent process. Thus, the immigration rate may
depend on density differences of both total cats and intact cats between the managed and
surrounding populations (52). By modeling immigration as a density-independent
process, the theoretical models might have underestimated this process's magnitude.
Second, these models did not include other density-dependent population compensatory
mechanisms such as increased survival and reproduction, which probably occurred in our
study.

*Comparison of TNR to culling*

When culling is applied intensively, it may eradicate cat populations, mainly on islands
(13, 15, 16). Although culling may be efficient, concurrent similar compensatory
mechanisms to those observed in our study might also limit its effectiveness (23, 24). In
some environmental contexts, TNR is considered as an alternative to culling. Since the
public acceptance of TNR is better than culling (9, 34, 35, 73), it improves public
cooperation with policy-makers, which is an important factor for population reduction
success. The merits of TNR go beyond those achieved by population reduction and
include improved welfare (63) and a reduction in cat-related nuisances, which are mainly
associated with reproduction (37, 74). On the other hand, the costs of TNR
implementation should be taken into account. Especially challenging is the maintenance
of high-intensity neutering for long periods, considering the significant decrease in
trapping efficiency when the percentage of neutered cats is high. More than one million
dollars ($US) were invested in the current project over the nine years of TNR
implementation. This cost should be compared to alternative population management

programs, considering the differences between countries, economies, jurisdictions, and
ethical perceptions.

*Complementary approaches that can further improve TNR benefits*

Combining TNR with removing a significant portion of the cat population (i.e.,
euthanasia of ill cats and adoption) can improve its effectiveness (39, 41-43). However,
the removal of cats might not be applicable everywhere. Our study's analysis of the
environmental factors influencing cat populations highlights an alternative approach to
complement TNR. The results show an association between vital anthropogenic resources
and cat population parameters. Limiting such resources might be combined with TNR to
enhance cat population reduction. Despite population reduction, neutered cats might
continue to hunt (75, 76). Therefore, TNR is not an effective means by itself for
preventing predation. It was shown that manipulating food content and applying
environmental enrichment reduced predatory cat behavior (77), and placing collar-
mounted devices interfered with hunting (78). Such measures can be therefore combined
with TNR to diminish this adverse effect further.

*Strengths and limitations*

The current experiment is unique not only in the controlled and long-term design but also
in the large number and depth of data of neutered and observed cats, the large covered
urban area which is systematically stratified (i.e., statistical areas), and the high-
resolution data of real-time events (resident report data) and environmental factors.
However, such a large study might also suffer from limitations. For example, some
processes might affect study results, such as changes in human behavior, changes in the
urban environment (e.g., constructions), local changes in cat numbers, and even
stochastic changes during the study period. We excluded the effect of these processes by
controlling TNR both in space and time, incorporating possible confounders in the
analysis, performing comparisons of the two datasets (resident data and annual counts
data performed by experts), and comparing our data to datasets representing mortality of
other animals and human behavior.

One limitation is a possible overestimation of the number of neutered cats due to a higher
probability of detecting sociable cats than shy cats. These sociable cats are also more
likely to be captured and neutered before capturing shy cats. Using friendly calling to
increase the counted cat number might have enhanced this bias. However, a recent study
showed that while friendly calling significantly increased the number of detected cats by
79%, its effect on the estimated neutering percentage was insignificant. Moreover, the
mean insignificant increase by 18% (from 46% to 56%) of neutering percentage
diminished almost completely when the neutering percentage was high (above 74%).
This observation was probably due to the growing proportion of neutered shy cats as the
neutering percentage increased (79). Hence, we assume that our estimations of neutering
intensity are reliable and valid and thus enable robust evidence for TNR effects and
limitations.

*Conclusion*

Our results indicate that fertility control can be applied to manage and even decrease the
size of open free-roaming cat populations. However, the enhancement of compensatory
mechanisms and the occupancy of immigrant cats might limit this effect. Therefore, we
conclude that maintaining a high neutering rate for a prolonged period and in spatial
contiguity is necessary to reduce cat population size. We recommend integrating these
with complementary methods such as vital resource regulation, ill cat euthanasia, and
adoption. In a broader view, our study suggests that the success of TNR should be
measured by considering its environmental context, aims, and public acceptance. While
simulation studies may aid in assessing such considerations, for enabling realistic
interpretation, we recommend including parameters of possible compensatory effects and
density-dependent immigration in these models. In particular, we call for a future robust
evaluation of the yield of integrated approaches.

**Materials and Methods**

**Study site**

We conducted the study from 2007 to 2018 in the city of Rishon-LeZion, located within
the greater Tel-Aviv metropolitan area in Israel. The jurisdictional area of Rishon-LeZion
is 58.7 km$^2$, divided according to the Israeli Central Bureau of Statistics into 64
commercial and residential statistical areas. Three statistical areas spanning 33.8 Km$^2$
include an isolated research institute, industrial and non-inhabited areas, and, therefore,
not part of the study area. The city's human population comprised 240,666 residents
living in a 25 km$^2$ area at the end of 2014 (Central Bureau of Statistics, Israel), divided
into 58 residential areas (each consists of approximately 4000 residents) and three
commercial areas (**Figure S1**). These 61 statistical areas constituted the study site.

**Study design**

The study period was divided into three consecutive phases (**Figure 1, Table S1**): (1) the
pre-intervention phase, (2) the mixed-intervention phase, and (3) the full-intervention
phase. The pre-intervention phase preceded the initiation of the TNR program from
January 2007 to the end of 2009. In the mixed-intervention phase, we allocated the 61
statistical areas into high- and low- intervention. In 31 statistical areas, the municipality
implemented a high-intensity multi-annual TNR program from October 2009 until
October 2014. The municipal TNR actions were limited in the other 30 statistical areas
(see **Table S2** for demographic comparison between the high and low intervention areas).
During this phase, 10,925 free-roaming cats were neutered with a male to female ratio of
1:1.07. In the full-intervention phase (November 2014 until December 2018), high-TNR
efforts were expanded to the entire city. During this phase, 11,219 free-roaming cats were
neutered with a male to female ratio of 1:1.06. The specific cat trapping and neutering
procedures are detailed in the **supplementary information**.

**Data collection**

To achieve high-resolution data on cat population dynamics, we used two independent
data collection methods—a repeated annual survey and daily resident reports.

*Annual surveys:*

We counted free-roaming cats via repeated annual surveys performed during September and October of 2012, 2013, 2014, and 2018. We conducted these surveys in 50 statistical areas (n=26 in the low-TNR statistical areas and n=24 in the high-TNR statistical areas) along fixed walking transects. In each statistical area, we chose one or two transects using a stratified random sampling design. Each statistical area was sampled by two observers, with each transect sampled twice each year. We documented the individual characteristics of each observed cat, including sex, presumptive age (kitten<6 months vs. adults > 6 months), and sterilization status (according to the presence/absence of ear-marks). We summarized the annual counts of cats (adults, queens, and kittens), kittens, and kitten-to-queen ratio per statistical area and year. We calculated the annual neutering percentage by dividing the total counts of neutered cats observed in each statistical area and year by the total counts of observed cats with identified neutering status. The transect walks were recorded using a cellular GPS recording application ("Endomondo™— Running & Walking, Android application," Under Armour Inc., Maryland, USA), to ensure walking on precisely the same path of each transect and to measure the transect length. The total length of the transects was 100.87 km (**Figure S3**). A comprehensive description of this sampling method, including its intra-observer, inter-observer, inter-transect agreement, and validity, is detailed in Gunther et al. 2020 (79).

*Resident reports:*

We retrieved data on cat carcasses and reproduction during 2007-2018 from the continuously available municipal emergency call center. The center receives voice complaints and reports from concerned residents on real-time events in the city jurisdiction area. The following data were recorded for each reported event: time and date of the call, location of the event, personal details about the calling resident, and a synopsis of the reported event. To confirm that the observed trends were unique to cat-related reports, we compared them to other municipal-based reports. The 2007-2018 cat carcass reports were compared to reports of carcasses of other animals (n=7,535), and cat reproduction reports were compared to the general municipal reports (apart from cat-related reports, n= 3,725,873) **(Figure S4)**. Further information on the call center data and its retrieval methods are detailed in the **supplementary information.**

*Environmental human-related data:*

The following environmental human-related factors were collected for Rishon-LeZion statistical areas during the study period: *human population density*, *socioeconomic status,* volume and type of *waste bins, cat feeding locations, educational institutions, food marketing businesses, built-up area* and *park area, age of neighborhood (years) and TNR actions performed by the municipality*. We provide further details on these factors in the **supplementary information**.

**Statistical analysis**

Data were aggregated for each combination of statistical area and year. These values
constituted the sampling units in all analyses. The following outcome variables were
analyzed: cat counts, kitten counts, queen counts, kitten-to-queen ratio, neutering
percentage, cat carcass reports, and cat reproduction reports. Except for the time-series
analyses, all statistical analyses were performed by fitting generalized linear models
(GLM) or generalized linear mixed models (GLMM) to these data (where the statistical
area was used as a random effect). In all of these models, a negative binomial distribution
was used. Since we analyzed rates, we used offsets as denominators (see **supplementary**
**information for details).** To account for potential spatial autocorrelation, we corrected
the analyses by using an exponential variogram.

We performed the statistical analysis in two stages. The trends in the entire city were
explored using all available data (i.e., survey data from 50 statistical areas and resident
reports from 61 statistical areas). To discern the effect of TNR intensity on the cat
population dynamics, we compared the cat population trends between the statistical areas
with the highest to the lowest neutering intensity at the end of the mixed-intervention
phase (designated as group-level analysis). To accomplish this, we divided the surveyed
statistical areas into quartiles according to the observed neutering percentage at the end of
the mixed-intervention phase. We compared two groups: Group-1 consisted of the
statistical areas with the lowest quartile of neutering percentage at this stage (n=12), and
Group-2 consisted of the highest quartile of neutering percentage at this stage (n=15).

First, the effects of human-related factors (predictor variables) on the various cat
population parameters in the entire city and their pairwise interactions with neutering
were tested by GLM (outcome variables: surveyed cat and kitten counts, and cat carcass
and reproduction reports). As we found that human population density and the volume of
accessed waste bins per resident were the main human-related predictors of most cat
population parameters, these were used in the following group-level analyses as
covariates. To test the difference of cat population outcome variables between the annual
surveys in the entire city, we performed GLMM. The group-level trend analyses were
performed separately for the survey and resident report data. In both analyses, the effects
of the treatment group, study phase, and their interaction (predictor variables) on the cat
population parameters (outcome variables, as detailed above, plus queen counts and
kitten-to queen ratio) were explored by GLMM. However, in the analysis of the resident
report data, the year was added as a covariate in order to realize the high resolution of this
data type. The specific modeling approach and statistical software used in each analysis
are detailed in the **supplementary information**.

To visualize the temporal trends of the high-resolution resident report data (cat carcass
and reproduction reports) collected both over the entire city or at the group level, we used
time-series analysis (Seasonal and Trend decomposition using Loess (STL) method (80)).
Through this method, we decomposed the monthly reports of the entire city (61statistical
areas) into three compartments: the general trend, cyclic changes (designated as
seasonality), and residual variability (remainder).

To confirm the construct validity of resident report data, we examined the correlation
between reproduction reports and the survey-derived kitten counts. We aggregated the
data to the neighborhood level and calculated the Pearson correlation coefficient of the
log-transformed kitten counts + 0.5 per transect length and the log-transformed
reproduction reports per the overall street length in each neighborhood in each year.
(**Figure S6**).

**Acknowledgments**

The authors are thankful to Y. Even-Zor, head of the municipal veterinary services, E.
Gian, head of the municipal GIS department, S. Keidar, principal at the maintenance
department, I. Ashkenazy from the municipal call center and E. Levi, head of the
municipal information and research center at the city of Rishon-LeZion.

**References**

1. Lowe S, Browne M, Boudjelas S, & De Poorter M (2000) *100 of the world's worst invasive*
*alien species: a selection from the global invasive species database* (Invasive Species
Specialist Group Auckland, New Zealand).
2. Feldman EC & Nelson RW (1996) Canine and feline endocrinology and reproduction. in
*Ch. 31 Feline reproduction* (W. B. Saunders Co., Philadelphia, PA 19106; USA), pp 741-
766.
3. England GC & Heimendahl Av (2010) *BSAVA manual of canine and feline reproduction*
*and neonatology* (British Small Animal Veterinary Association).
4. Scott KC, Levy JK, & Crawford PC (2002) Characteristics of free-roaming cats evaluated in
a trap-neuter-return program. *Journal of the American Veterinary Medical Association*
221(8):1136-1138.
5. Serpell JA (2000) Domestication and history of the cat. *The domestic cat: The biology of*
*its behaviour* 179.
6. Loss SR & Marra PP (2017) Population impacts of free-ranging domestic cats on
mainland vertebrates. *Frontiers in Ecology and the Environment* 15(9):502-509.
7. Gerhold RW & Jessup DA (2013) Zoonotic diseases associated with free-roaming cats.
*Zoonoses and Public Health* 60(3):189-195.
8. Dabritz HA & Conrad PA (2010) Cats and Toxoplasma: implications for public health.
*Zoonoses and Public Health* 57(1):34-52.
9. Slater MR*, et al.* (2008) Free-roaming dogs and cats in central Italy: Public perceptions of
the problem. *Preventive Veterinary Medicine* 84(1–2):27-47.
10. Gunther I, Raz T, Berke O, & Klement E (2015) Nuisances and welfare of free-roaming
cats in urban settings and their association with cat reproduction. *Preventive Veterinary*
*Medicine* 119(3–4):203-210.
11. Medina FM*, et al.* (2011) A global review of the impacts of invasive cats on island
endangered vertebrates. *Global Change Biology* 17(11):3503-3510.
12. Boone JD (2015) Better trap–neuter–return for free-roaming cats Using models and
monitoring to improve population management. *Journal of Feline Medicine and Surgery*
17(9):800-807.

13. Bester M*, et al.* (2002) A review of the successful eradication of feral cats from sub-
Antarctic Marion Island, Southern Indian Ocean. *South African Journal of Wildlife*
*Research-24-month delayed open access* 32(1):65-73.
14. Parkes J & Murphy E (2003) Management of introduced mammals in New Zealand. *New*
*Zealand Journal of Zoology* 30(4):335-359.
15. Ramsey DSL, Parkes JP, Will D, Hanson CC, & Campbell KJ (2011) Quantifying the success
of feral cat eradication, San Nicolas Island, California. *New Zealand Journal of Ecology*
35(2):163-173.
16. Nogales M*, et al.* (2004) A review of feral cat eradication on islands. *Conservation*
*Biology* 18(2):310-319.
17. Kennedy B, Cumming B, & Brown WY (2020) Global strategies for population
management of domestic cats (Felis catus): a systematic review to inform best practice
management for remote Indigenous communities in Australia. *Animals* 10(4):663.
18. Calver MC, Crawford HM, & Fleming PA (2020) Response to Wolf et al.: Furthering
debate over the suitability of trap-neuter-return for stray cat management. *Animals*
10(2):362.
19. Schmidt PM, Swannack TM, Lopez RR, & Slater MR (2009) Evaluation of euthanasia and
trap-neuter-return (TNR) programs in managing free-roaming cat populations. *Wildlife*
*Research* 36(2):117-125.
20. Hein S & Jacob J (2019) Population recovery of a common vole population (Microtus
arvalis) after population collapse. *Pest Management Science* 75(4):908-914.
21. Shilova S & Tchabovsky A (2009) Population response of rodents to control with
rodenticides. *Current Zoology* 55(2):81-91.
22. Williams CK, Parer I, Coman B, Burley J, & Braysher M (1995) Managing vertebrate pests:
rabbits.
23. Lazenby BT, Mooney NJ, & Dickman CR (2015) Effects of low-level culling of feral cats in
open populations: a case study from the forests of southern Tasmania. *Wildlife Research*
41(5):407-420.
24. Palmas P*, et al.* (2020) Rapid recolonisation of feral cats following intensive culling in a
semi-isolated context. *NeoBiota* 63:177-200.
25. Baker PJ & Harris S (2006) Does culling reduce fox (Vulpes vulpes) density in commercial
forests in Wales, UK? *European Journal of Wildlife Research* 52(2):99-108.
26. Ransom JI, Powers JG, Hobbs NT, & Baker DL (2014) Ecological feedbacks can reduce
population-level efficacy of wildlife fertility control. *Journal of Applied Ecology*
51(1):259-269.
27. McCarthy RJ, Levine SH, & Reed JM (2013) Estimation of effectiveness of three methods
of feral cat population control by use of a simulation model. *Journal of the American*
*Veterinary Medical Association* 243(4):502-511.
28. Loyd KAT & DeVore JL (2010) An evaluation of feral cat management options using a
decision analysis network. *Ecology and Society* 15(4):art 10-art 10.
29. Miller PS*, et al.* (2014) Simulating free-roaming cat population management options in
open demographic environments.
30. Andersen MC, Martin BJ, & Roemer GW (2004) Use of matrix population models to
estimate the efficacy of euthanasia versus trap-neuter-return for management of free-
roaming cats. *Journal of the American Veterinary Medical Association* 225(12):1871-
1876.

31.    Denny EA & Dickman C (2010) Review of cat ecology and management strategies in
Australia. *Invasive Animals Cooperative Research Centre, Canberra*.

32.    Longcore T, Rich C, & Sullivan LM (2009) Critical assessment of claims regarding
management of feral cats by trap-neuter-return. *Conservation Biology* 23(4):887-894.

33.    Boone JD*, et al.* (2019) A long-term lens: Cumulative impacts of free-roaming cat
management strategy and intensity on preventable cat mortalities. *Frontiers in*
*Veterinary Science* 6:238.

34.    Wolf PJ & Schaffner JE (2019) The road to TNR: Examining Trap-Neuter-Return through
the lens of our evolving ethics. *Frontiers in Veterinary Science* 5:341.

35.    Wolf PJ & Hamilton F (2020) Managing free-roaming cats in US cities: An object lesson in
public policy and citizen action. *Journal of Urban Affairs*:1-22.

36.    Neville PF & Remfry J (1984) Effect of neutering on two groups of feral cats. *Veterinary*
*Record* 114(18):447-450.

37.    Hughes KL & Slater MR (2002) Implementation of a feral cat management program on a
university campus. *Journal of Applied Animal Welfare Science* 5(1):15-28.

38.    Hughes KL, Slater MR, & Haller L (2002) The effects of implementing a feral cat
spay/neuter program in a Florida county animal control service. *Journal of Applied*
*Animal Welfare Science* 5(4):285-298.

39.    Levy JK, Gale DW, & Gale LA (2003) Evaluation of the effect of a long-term trap-neuter-
return and adoption program on a free-roaming cat population. *Journal of the American*
*Veterinary Medical Association* 222(1):42-46.

40.    Algar D, Hilmer S, Nickels D, & Nickels A (2011) Successful domestic cat neutering: first
step towards eradicating cats on Christmas Island for wildlife protection. *Ecological*
*Management & Restoration* 12(2):93-101.

41.    Levy J, Isaza N, & Scott K (2014) Effect of high-impact targeted trap-neuter-return and
adoption of community cats on cat intake to a shelter. *The Veterinary Journal*
201(3):269-274.

42.    Swarbrick H & Rand J (2018) Application of a protocol based on Trap-Neuter-Return
(TNR) to manage unowned urban cats on an Australian University Campus. *Animals*
8(5):77.

43.    Kreisler RE, Cornell HN, & Levy JK (2019) Decrease in Population and Increase in Welfare
of Community Cats in a Twenty-Three Year Trap-Neuter-Return Program in Key Largo,
FL: the ORCAT Program. *Frontiers in Veterinary Science* 6:1-14.

44.    Jones AL & Downs CT (2011) Managing feral cats on a university's campuses: how many
are there and is sterilization having an effect? *Journal of Applied Animal Welfare Science*
14(4):304-320.

45.    Zito S, Aguilar G, Vigeant S, & Dale A (2018) Assessment of a targeted trap-neuter-return
pilot study in Auckland, New Zealand. *Animals* 8(5):73.

46.    Nutter F (2005) Evaluation of a trap-neuter-return management program for feral cat
colonies: population dynamics, home ranges, and potentially zoonotic diseases. . *PhD*
*thesis in Comparative Biomedical Science, North Carolina State University, Raleigh, NC.*

47.    Natoli E*, et al.* (2006) Management of feral domestic cats in the urban environment of
Rome (Italy). *Preventive Veterinary Medicine* 77(3/4):180-185.

48.    Spehar DD & Wolf PJ (2017) An examination of an iconic trap-neuter-return program:
The Newburyport, Massachusetts case study. *Animals* 7(11):81.

49. Spehar DD & Wolf PJ (2019) Back to school: An updated evaluation of the effectiveness
of a Long-Term Trap-Neuter-Return program on a university's free-roaming cat
population. *Animals* 9(10):768.
50. Tan K, Rand J, & Morton J (2017) Trap-Neuter-Return Activities in Urban Stray Cat
Colonies in Australia. *Animals* 7(6):46.
51. Spehar DD & Wolf PJ (2018) A case study in citizen science: The effectiveness of a trap-
neuter-return program in a Chicago neighborhood. *Animals* 8(1):14.
52. Gunther I, Finkler H, & Terkel J (2011) Demographic differences between urban feeding
groups of neutered and sexually intact free-roaming cats following a trap-neuter-return
procedure. *Journal of the American Veterinary Medical Association* 238(9):1134-1140.
53. Castillo D & Clarke AL (2003) Trap/neuter/release methods ineffective in controlling
domestic cat "colonies" on public lands. *Natural Areas Journal* 23(3):247-253.
54. Kilgour RJ*, et al.* (2017) Estimating free-roaming cat populations and the effects of one
year Trap-Neuter-Return management effort in a highly urban area. *Urban Ecosystems*
20(1):207-216.
55. Mannhart T (2007) A catch-neuter-release project for free-roaming dogs and cats in
Rhodes, Greece: Problem analysis and effectiveness of the strategy. PhD thesis
(University of Bern. ).
56. Natoli E*, et al.* (2019) Evaluation of Unowned Domestic Cat Management in the Urban
Environment of Rome After 30 Years of Implementation of the No-Kill Policy (National
and Regional Laws). *Frontiers in veterinary science* 6:31.
57. Bissonnette V, Lussier B, Doizé B, & Arsenault J (2018) Impact of a trap-neuter-return
event on the size of free-roaming cat colonies around barns and stables in Quebec: A
randomized controlled trial. *Canadian Journal of Veterinary Research* 82(3):192-197.
58. Boone JD*, et al.* (2014) Improving and evaluating trap-neuter-return (TNR) management
for outdoor cats on the human landscape. *Proceedings of the Vertebrate Pest
Conference*.
59. Foley P, Foley JE, Levy JK, & Paik T (2005) Analysis of the impact of trap-neuter-return
programs on populations of feral cats. *Journal of the American Veterinary Medical
Association* 227(11):1775-1781.
60. Mirmovitch V (1995) Spatial organisation of urban feral cats (Felis catus) in Jerusalem.
*Wildlife Research* 22(3):299-310.
61. Finkler H, Hatna E, & Terkel J (2011) The influence of neighbourhood socio-demographic
factors on densities of free-roaming cat populations in an urban ecosystem in Israel.
*Wildlife Research* 38(3):235-243.
62. Nutter FB, Levine JF, & Stoskopf MK (2004) Reproductive capacity of free-roaming
domestic cats and kitten survival rate. *Journal of the American Veterinary Medical
Association* 225(9):1399-1402.
63. Gunther I, Raz T, & Klement E (2018) Association of neutering with health and welfare of
urban free-roaming cat population in Israel, during 2012-2014. *Preventive Veterinary
Medicine*.
64. Murray JK, Skillings E, & Gruffydd-Jones TJ (2008) A study of risk factors for cat mortality
in adoption centres of a UK cat charity. *Journal of Feline Medicine & Surgery* 10(4):338-
345.
65. Childs JE & Ross L (1986) Urban cats: characteristics and estimation of mortality due to
motor vehicles. *American Journal of Veterinary Research* 47(7):1643-1648.

66.    Chambers LK, Singleton GR, & Hinds L (1999) Fertility control of wild mouse populations:
the effects of hormonal competence and an imposed level of sterility. *Wildlife Research*
26(5):579-591.
67.    Twigg LE*, et al.* (2000) Effects of surgically imposed sterility on free-ranging rabbit
populations. *Journal of Applied Ecology* 37(1):16-39.
68.    Turner A & Kirkpatrick JF (2002) Effects of immunocontraception on population,
longevity and body condition in wild mares (Equus caballus). *Reprod Suppl* 60:187-195.
69.    Williams CK*, et al.* (2007) Population responses to sterility imposed on female European
rabbits. *Journal of Applied Ecology*:291-301.
70.    Smith LM*, et al.* (2019) The Effectiveness of Dog Population Management: A Systematic
Review. *Animals* 9(12):1020.
71.    Scott KC, Levy JK, Gorman SP, & Newell SM (2002) Body condition of feral cats and the
effect of neutering. *Journal of Applied Animal Welfare Science* 5(3):203-213.
72.    Finkler H, Gunther I, & Terkel J (2011) Behavioral differences between urban feeding
groups of neutered and sexually intact free-roaming cats following a trap-neuter-return
procedure. *Journal of the American Veterinary Medical Association* 238(9):1141-1149.
73.    Tennent JK, Downs CT, Wald DM, & Watson HK (2010) Public perceptions of feral cats
within an urban conservancy on a campus of the University of KwaZulu-Natal. *South
African Journal of Wildlife Research* 40(1):16-26.
74.    Ireland T & Neilan RM (2016) A spatial agent-based model of feral cats and analysis of
population and nuisance controls. *Ecological Modelling* 337:123-136.
75.    Bruce SJ*, et al.* (2019) Predation and risk behaviors of free-roaming owned cats in
Auckland, New Zealand via the use of animal-borne cameras. *Frontiers in Veterinary
Science* 6.
76.    Loyd KAT, Hernandez SM, Carroll JP, Abernathy KJ, & Marshall GJ (2013) Quantifying
free-roaming domestic cat predation using animal-borne video cameras. *Biological
Conservation* 160:183-189.
77.    Cecchetti M, Crowley SL, Goodwin CE, & McDonald RA (2021) Provision of High Meat
Content Food and Object Play Reduce Predation of Wild Animals by Domestic Cats Felis
catus. *Current Biology* 31(5):1107-1111. e1105.
78.    Nelson S, Evans A, & Bradbury R (2005) The efficacy of collar-mounted devices in
reducing the rate of predation of wildlife by domestic cats. *Applied Animal Behaviour
Science* 94(3-4):273-285.
79.    Gunther I, Azriel L, Wolf H, Raz T, & Klement E (2020) An accessible scheme for
monitoring free-roaming cat population trends. *Ecology and Evolution* 10:1288-1298.
80.    Cleveland RB, Cleveland WS, McRae JE, & Terpenning I (1990) STL: a seasonal-trend
decomposition. *Journal of Official Statistics* 6(1):3-73.

737
738

739 **Figures and Tables**

740
741
**Figure 1:** The experimental design was set to assess long-term TNR effectiveness over time and
space. The study period was divided into three phases: 1ˢᵗ phase – pre-intervention period -
between 2007 to the end of 2009; 2ⁿᵈ phase -mixed-intervention - allocating TNR to half of the
city neighborhoods between the end of 2009 to end of 2014; 3ʳᵈ phase, full-intervention period,
allocating TNR to the entire city between the end of 2014 and the end of 2018. Dots present
locations of TNR in the city area. The study area was composed of 50 statistical areas (presented
as polygons in the upper map; see also Fig. S1), 24 of which are termed high-TNR, as they were
treated in both TNR phases, and the other 26, termed low-TNR, as they were mainly treated
during the full-intervention phase. From these high- and low-TNR statistical areas, 15 and 12
areas with the highest and lowest neutering percentage at the 2014 survey, respectively, were
served as the main study units (Group-1 and -2). Survey years are marked by an asterisk.

**Figure 2**: Annual trends (geometric mean±CI$_{95\%}$) of (**A**) neutering percentage, (**B**) free-roaming
cat counts, (**C**) kitten counts, (**D**) queen counts, and (**E**) kitten-to-queen ratio in the city of
Rishon-LeZion (black, n = 50 statistical areas), and in Group-1 (turquoise, lowest neutering
intensity, n=12 statistical areas) and Group-2 (purple, highest neutering intensity, n=15 statistical
areas). Observations were performed in 2012–2014 and 2018. Superscript letters represent years,
where observations differ.

**Figure 3:** Municipal TNR efforts monthly-based Time series decomposition plots (between 2007-2018; the crude data is divided into three compartments: trend, seasonality and remainder). (**A**) Monthly-based time series plots of municipal TNR actions (n = 22,144), cat carcass reports (n = 36,544) and reproduction reports (n = 12,217) from January 2007 to December 2018 across the entire city (61 statistical areas). (**B**) Time series decomposition plots of carcass and reproduction reports linked with cumulative TNR percentage. The 1st phase refers to the pre-intervention period (before the initiation of the TNR program), the 2nd phase refers to the mixed-intervention

period (applying TNR in half of the city), and the 3rd phase refers to the full-intervention period (applying TNR in the entire city).

**Figure 4**: Cumulative TNR percentage and monthly-based time series decomposition plots (between 2007-2018; (the crude data is divided into three compartments: trend, seasonality and remainder) of (**A**) cat carcass reports and (**B**) cat reproduction reports in Group-1 (lowest neutering intensity, n = 12) compared to Group-2 (highest neutering intensity, n = 15). The 1st phase refers to the pre-intervention period (before the initiation of the TNR program), the 2nd phase refers to the mixed-intervention period (applying TNR primarily in Group-2), and the 3rd phase refers to the full-intervention period (applying TNR in both groups).

**Table 1:** Annual observation counts of cats, neutered cats, neutering percentage, kittens, queens, and kitten-to-queen ratio in the low-TNR (n=26), high-TNR statistical areas (n=24), Group-1 (lowest neutering intensity, n=12) and Group-2 (highest neutering intensity, n=15).

| Variable | Year | Low-TNR | High-TNR | Group-1 | Group-2 | Overall |
|---|---|---|---|---|---|---|
| Cats | 2012 | 1726 | 1485 | 675 | 938 | 3211 |
| | 2013 | 1880 | 1443 | 770 | 906 | 3233 |
| | 2014 | 2219 | 1843 | 1000 | 1082 | 4062 |
| | 2018 | 1705 | 1417 | 688 | 765 | 3122 |
| Neutered | 2012 | 599 | 918 | 145 | 620 | 1517 |
| | 2013 | 644 | 1003 | 136 | 684 | 1647 |
| | 2014 | 742 | 1312 | 189 | 866 | 2054 |
| | 2018 | 1274 | 982 | 477 | 590 | 2256 |
| Neutering percentage | 2012 | 34.7 | 61.8 | 21.5 | 66.1 | 47.2 |
| | 2013 | 34.3 | 69.5 | 17.7 | 75.5 | 50.9 |
| | 2014 | 33.4 | 71.2 | 18.9 | 80.0 | 50.6 |
| | 2018 | 74.7 | 69.3 | 69.3 | 77.1 | 72.3 |
| Kittens | 2012 | 291 | 144 | 111 | 96 | 435 |
| | 2013 | 228 | 90 | 108 | 57 | 318 |
| | 2014 | 292 | 104 | 162 | 35 | 396 |
| | 2018 | 165 | 172 | 72 | 79 | 337 |
| Queens | 2012 | 273 | 114 | 123 | 61 | 387 |
| | 2013 | 308 | 103 | 158 | 49 | 411 |
| | 2014 | 400 | 113 | 226 | 50 | 513 |
| | 2018 | 91 | 103 | 43 | 36 | 194 |
| Kitten-to-queen ratio | 2012 | 1.07 | 1.26 | 0.90 | 1.57 | 1.12 |
| | 2013 | 0.74 | 0.87 | 0.68 | 1.16 | 0.77 |
| | 2014 | 0.73 | 0.92 | 0.72 | 0.70 | 0.77 |
| | 2018 | 1.81 | 1.67 | 1.67 | 2.19 | 1.74 |

**Table 2.** Annual trends of the cat and kitten counts, kitten-to-queen ratio (surveyed
during the mixed- and full-intervention phases), and cat carcass and reproduction data
(received from resident reports during all the three study phases). The trends are depicted
for Group-1, (lowest neutering intensity, n=12), and Group-2, (highest neutering
intensity, n=15). All the models are adjusted for human population density and accessed
waste bins.

| Population parameter | Phase | Measurement period | Annual trend for Group-1 ($CI_{95\%}$) [%] | Annual trend for Group-2 ($CI_{95\%}$) [%] | P-value |
|---|---|---|---|---|---|
| Cat counts | 2nd phase | 2012-2014 | 20.72 (8.19 to 34.71) [a1]* | 4.02 (-5.62 to 14.65) [a2] | 0.046 |
| | 3rd phase | 2014-2018 | -7.95 (-13.09 to -2.50) [b1] | -7.83 (-12.29 to -3.15) [b2] | 0.973 |
| Kitten counts | 2nd phase | 2012-2014 | 20.20 (-11.31 to 62.91) [a1] | -42.39 (-58.19 to -20.60) [a2] | 0.001 |
| | 3rd phase | 2014-2018 | -14.14 (-27.00 to 1.00) [b1] | 21.37 (3.20 to 42.74) [b2] | 0.003 |
| Queen counts | 2nd phase | 2012-2014 | 40.40 (13.31 to 73.97) [a1] | -11.21 (-30.92 to 14.12) [a2] | 0.007 |
| | 3rd phase | 2014-2018 | -38.34 (-46.47 to -28.98) [b1] | -8.76 (-20.50 to 4.71) [a2] | <0.001 |
| Kitten-to-queen ratio | 2nd phase | 2012-2014 | -7.72 (-36.10 to 33.28) [a1] | -28.48 (-50.60 to 3.55) [a2] | 0.338 |
| | 3rd phase | 2014-2018 | 29.88 (7.40 to 57.08) [a1] | 23.35 (2.26 to 48.28) [b2] | 0.706 |
| Carcass reports | 1st phase | 2007-2009 | 7.96 (-1.44 to 18.26) [a1] | 6.01 (-1.62 to 14.23) [a2] | 0.761 |
| | 2nd phase | 2010-2014 | 0.88 (-3.04 to 4.95) [a1] | -12.02 (-15.17 to -8.75) [b2] | <0.001 |
| | 3rd phase | 2015-2018 | -13.59 (-19.02 to -7.79) [b1] | 0.57 (-4.82 to 6.27) [a2] | <0.001 |
| Reproduction reports | 1st phase | 2007-2009 | 8.02 (-8.03 to 26.87) [a1] | -5.70 (-18.44 to 9.04) [a2] | 0.219 |
| | 2nd phase | 2010-2014 | 3.89 (-2.95 to 11.21) [a1] | -24.50 (-29.94 to -18.63) [b2] | <0.001 |
| | 3rd phase | 2015-2018 | -7.23 (-17.18 to 3.91) [a1] | 22.56 (9.96 to 36.59) [c2] | <0.001 |

794
795 * The superscript letters represent comparisons between pairs of phases within each group of
796 statistical areas (pairs with insufficient evidence for a difference are labeled with the same
797 superscript letter).