The interkingdom horizontal gene transfer in 44 early diverging fungi is driven by bursts of gene exchange and duplication, resulting in boosted adaptive capabilities of selected lineages

Michał Ciach*[1,3] ,Julia Pawłowska[2], Paweł Górecki[3] and Anna Muszewska*[1]

[1] Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Pawinskiego 5A, 02-106 Warsaw, Poland

[2] Department of Molecular Phylogenetics and Evolution, Faculty of Biology, Biological and Chemical Research Centre, University of Warsaw, ul. Żwirki i Wigury 101, 02-089 Warsaw, Poland;

[3] Faculty of Mathematics, Informatics and Mechanics, University of Warsaw, Stefana Banacha 2, 02-097 Warsaw, Poland

*corresponding authors

Running title: HGT in 44 early diverging fungi

Keywords: Early Diverging Fungi, Horizontal Gene Transfer, Fungal immunity, Ecological adaptation, Unrooted Gene Trees, HGT patterns, associated bacteria

Abstract

Numerous studies have been devoted to individual cases of horizontally acquired genes in fungi. It has been shown that such genes expand the hosts' metabolic capabilities and contribute to their adaptations as parasites or symbionts. Some studies have provided an extensive characterization of the horizontal gene transfer (HGT) in Dikarya. However, in the early diverging fungi (EDF), a similar characterization is still missing. In order to fill this gap, we have designed a computational pipeline to obtain a statistical sample of reliable HGT events with a possibly minimal number of false detections. We have analyzed 44 EDF genomes and identified 185 phylogenetic trees with 1077 xenologs ranging from *Chytridiomycota* and *Blastocladiomycota* to *Mucoromycota*. We have identified several patterns and statistical properties of EDF HGT. Ancestrally aquatic fungi are generally more likely to acquire foreign genetic material than terrestrial ones. A close ecological relationship with another organism is a predisposing condition for HGT, as exemplified by a transfer of NOD-like receptors between Mortierellomycotina and their symbiotic bacteria, but does not always result in an extensive gene exchange. Closely related fungi have similar rates of intronization of xenologs. The evolution of the number of xenologs after transfer can be explained by the Yule model of speciation. Post-transfer gene fusions complicate the landscape of HGT.

Importance

Although it is now recognized that horizontal gene exchange is a factor influencing the adaptation and evolution of eukaryotic organisms, the so far described cases in early

diverging fungi (EDF) are fragmentary, and a large-scale comprehensive study is lacking. We have designed a methodology to obtain a reliable, statistical sample of inter-kingdom xenologs across the tree of life of EDF to give a preliminary characterization of their general properties and patterns. We study how different fungal lineages vary in terms of the number of xenologs, what are their ecological associations, and the molecular properties of proteins encoded by the acquired genes. Our results help to better understand to what extent and in what way the incorporation of foreign genetic material shaped the metabolic, adaptive, and immune capabilities of fungi.

Introduction

Horizontal gene transfer (HGT) is a process of exchange of genetic material between distantly related species. It has been extensively studied in prokaryotes and proved to be an important mechanism that increases genomic diversity, contributes to niche adaptation, antibiotic resistance, new metabolic pathways, microbiome stability (Coyte et al. 2022) and numerous other traits. In eukaryotes, the significance and frequency of HGT remain mostly unclear, and the mechanisms of gene transfer beyond *Agrobacterium*-mediated gene transformation are still a matter of debate (Naranjo-Ortiz and Gabaldón 2020). It has been shown, however, that HGT does occur both within eukaryotes (including animals) and between eukaryotes and prokaryotes (Gabaldón 2020; Keeling and Palmer 2008).

Fungi represent very diverse life strategies, making them a unique model to study the connections between HGT and ecological adaptations. Genomes of fungi are protected against the incorporation of foreign DNA at multiple levels: starting from the the physical separation of chromatin from the rest of the cell by the nuclear envelope, through the use of alternative genetic codes (e.g. in CTG *Candida albicans* and *Pachysolen tannophilus*), to foreign DNA elimination mechanisms RNAi and RIP, which are ubiquitous in fungi (Krassowski et al. 2018; Clutterbuck and John Clutterbuck 2011; Torres-Martínez and Ruiz-Vázquez 2017). On the other hand, the intake of foreign DNA may be facilitated by the osmotrophic lifestyle and hyphal fusion (Priest, Yadav, and Heitman 2020; Feurtey and Stukenbrock 2018; Richards and Talbot 2013). The complex interplay of factors facilitating and preventing the incorporation of foreign DNA places fungi somewhere between bacteria and animals in terms of the rate of HGT and contributes to large differences in the numbers of HGT between different fungal lineages.

Several horizontally transferred genes have documented adaptive roles for fungal fitness. A fungal pathogen of insects, *Metarhizium robertsii*, has acquired sterol carrier and peptidase genes that facilitate the penetration of the victim's body (Zhao et al. 2014; Zhang et al. 2019). Multiple transfers of toxins in plant-infecting fungi have improved their ability to infect new hosts (Mehrabi et al. 2011; Ambrose, Koppenhöfer, and Belanger 2014). Bacteria have been described as a source of extensive HGT in plant-associated *Colletotrichum* sp. (Jaramillo et al. 2015), *Verticillium dahliae* (Shi-Kunne et al. 2019) and animal skin pathogens *Malassezia* (Ianiri et al. 2020). Spectacular examples of transfers of clusters of functionally-linked genes (Wisecaver and Rokas 2015; Wisecaver, Slot, and Rokas 2014), genes facilitating metabolism expansions (Gao et al. 2019) or even conditionally dispensable chromosomes in pathogenic *Fusarium* species (Ma et al. 2010) have influenced the

perception of genomic evolution in fungi. They have shown that HGT can provide a faster way to gain the ability to colonize new hosts and degrade new compounds than vertical inheritance.

Existing studies of fungal HGT have mostly focused on the Dikarya fungi (Richards et al. 2011; Marcet-Houben and Gabaldón 2010), while HGT in the early diverging fungi (EDF, non-Dikarya fungi) is relatively understudied. Although no comprehensive characterization of HGT in EDF has been presented so far, it is already known that it does occur and may have a profound impact on the recipient organism. Recent reports show that EDF experienced several documented waves of transfers, like the acquisition of secondary metabolite clusters in *Basidiobolus* (Tabima et al. 2020), *Mortierella* (Wurlitzer et al. 2021) and *Rhizophagus* (Venice et al. 2020), or number of enzymes transferred to *Neocallimastigomycota* involved in the metabolism of nucleotides (Alexander et al. 2016) and carbohydrates (Murphy et al. 2019). A more complex scenario was reported for *Rhizophagus irregularis* that acquired genes both from the host plant and its endosymbiotic bacterium (Li et al. 2018). Catalase genes of bacterial origin, which presumably increase ROS resistance, were found in the amphibian pathogen *Batrachochytrium dendrobatidis* (Hansberg, Salas-Lizana, and Domínguez 2012).

In this work, we study the patterns of gene exchange between fungi and non-fungal organisms on a representative set of 44 EDFs ranging from *Chytridiomycota* to *Mucoromycota*. We have designed a custom phylogenetic-based pipeline to obtain a reliable sample of xenologs, focusing on limiting the number of false positive results that distort the observed signal (Irwin et al. 2022) while retaining a sufficient sample size to infer statistical trends. As potential sources of false positives, we have focused on genome contaminations, spurious homology and promiscuous domains, the discrepancy between closest blast hits and phylogenetically nearest neighbors, unreliable alignments, long branches, weak bootstrap supports, uncertain taxonomy and sequence homoplasy. We then analyze the resulting data set of HGTs from a taxonomic, ecological, and molecular perspective. Our results provide a comprehensive characterization of the patterns of HGT across the EDF tree of life.

Methods

Dataset

Proteomes of 44 early diverging fungi, referred to as the *target proteomes* (Supplementary Table S1), were downloaded from NCBI in April 2018 (NCBI Resource Coordinators 2017). Fungal ecological tagging was obtained from Fungal Traits (Põlme et al. 2020). Taxonomical annotations were obtained from the NCBI taxonomy database, with manual fine-tuning of taxonomic names and ranks when needed according to (Voigt et al. 2021). Since the NCBI Taxonomy tree contains many polytomies, we take a conservative approach and report HGTs only if they are supported by all possible binarizations. Genomic features, including intron count, were obtained from the GFF files of the genome assemblies.

HGT identification

Low-complexity regions in all target proteins were masked using ncbi-seg with relaxed settings (window length 12, trigger complexity 1.8, extension complexity 2.0) (John C. Wootton and Federhen 1993). Target proteins with less than 30 non-masked amino acids, more than 2000 characters, or more than 50% masking were discarded as prone to produce false-positive hits. Target protein homology within the 44 target proteomes was evaluated using the blastp suite with an e-value threshold 1e-05 (NCBI Resource Coordinators 2017). Contigs without any detectable homology to the other 44 target fungi were discarded as likely genome contaminants (first-stage contaminant filtering). Target proteins with homologs in multiple fungal families were discarded as likely ancestrally fungal.

Homologous sequences for the remaining proteins were found in the Non-Redundant NCBI database (last updated December 01, 2020) using blastp suite with an e-value threshold 1e-04 and maximum of 500 target sequences. Detected homologs were masked using ncbi-seg (default settings). Proteins with less than 30 non-masked amino acids, more than 2000 characters, or more than 75% masking were discarded. The taxonomic lineage of each homolog was inspected using the NCBI Taxonomy database. Only homologs from bacteria, archaea and eukarya were retained in order to discard viral and artificial sequences due to their uncertain location in the species tree. Proteins with 'unclassified', 'uncultured' or 'environmental' keywords anywhere in their lineage, as well as non-fungal proteins with the '*incertae sedis*' keyword, were discarded for the same reason (note that some of the fungi analyzed in this paper belong to *incertae sedis* families; their homologs were not discarded).

All remaining proteins were clustered with MCL. First, an all-vs-all BlastP was performed, with an increased word size to increase specificity and speed up computations (-word_size 6 -threshold 21 -evalue 1e-06). Only HSP matches with identity at least 30% and covering at least 50% of both the query and the subject proteins were retained. This step was essential to avoid cluster merging caused by long, multi-domain proteins and by small but highly conserved regions. The resulting protein homology graph was clustered with MCL (inflation parameter = 2). The results were filtered by discarding clusters without target proteins, with less than 4 different species, with more than 60% fungal species, and clusters with fungal sequences from more than a single phylum (as either ancestral or exhibiting evolutionary patterns too complex for an automated analysis).

For each remaining target protein, up to 10 additional proteins from the same contig were selected for the second stage contaminant filtering. The selected additional proteins were blasted against the NR database. Contigs with no detected protein with a fungal first blast hit were discarded as likely contaminants. The remaining target proteins and their homologs from the remaining clusters were clustered again with MCL using the same strategy. This step was performed because removing large numbers of nodes from a homology graph can alter the structure of clusters.

The resulting clusters were filtered again using the same strategy. The remaining clusters were aligned using mafft 3.7 (Nakamura et al. 2018) with E-INS-i strategy, recommended for the alignment of general sequences due to having fewer assumptions than other strategies

(--genafpair --maxiterate 5000). Alignments were then filtered by discarding constant columns and columns with more than 80% gaps, and next by discarding sequences with less than 20 non-gap characters or with more than 75% gaps.

Maximum likelihood (ML) trees were inferred using IqTree (Minh et al. 2020) with automated model selection. Branch supports were assessed with UFBoot, 2000 replicates. Leaves were discarded if no fungal leaf was closer than 3.33 substitutions per site (corresponding to an average of 15% sequence identity (Grishin 1995)). This step was performed in order to reduce the complexity of the trees and focus on the neighborhoods of fungal clades. Next, long branches (both terminal and internal ones) were cut iteratively until all branches were no longer than 1.598 substitutions per site (corresponding to an average of 30% sequence identity). This step was performed in order to limit the influence of long branches on tree topology. Resulting trees were retained if they contained proteins from at least one target and three non-fungal species. See the Supplementary Material for a detailed description of the branch-cutting algorithm.

For all remaining trees, their sequences were re-aligned and the alignments were filtered using the same strategy. Next, ML trees were recomputed using the same strategy. This step was performed because removing sequences can alter the topologies, branch lengths, and branch supports of ML trees.

For each tree, a corresponding species tree was downloaded from a local copy of the NCBI Taxonomy database. For each gene tree/species tree pair, all fungal clades were identified. Next, the species composition of the neighboring non-fungal clades was compared between the trees. Fungal clades which neighbored different clades in the gene tree and the species tree (for all possible binarizations of the latter) were considered displaced and therefore indicative of HGT. Otherwise, if the location of a fungal clade agreed in both trees, it was deemed as a possible vertical inheritance or sequence homoplasy and therefore not sufficient evidence to infer a horizontal gene transfer. See **Fig. 1** for a visual description of the approach and the Supplementary Material for a detailed description of the rationale and algorithms used for the detection of fungal clades and comparison of their locations between trees.

To further limit the number of false positive results, we discarded gene trees in which fungal clades could not be repositioned to their correct locations, i.e. gene trees with no branch giving the same species neighborhood as in the species tree (see **Fig. 1**). In gene trees without such a branch, any location of a fungal clade would be artifactually indicative of an HGT. This approach allows for additional transfers and evolutionary events within the neighboring clades but discards trees with transfers between the neighboring clades or gene tree inference errors as too complex for an automated analysis and prone to return false positive results.

For each fungal clade, an average of its UFBoot support value and the support of the two neighboring clades was computed (see **Fig. 1**). Clades for which the average was lower than 90% were discarded as weakly supported HGTs. For the remaining clades, we have estimated the origin of the transfer by inspecting the last common ancestors (LCAs) of

species in its two neighboring clades and the joint LCA of both neighboring clades, ignoring fungal species. If one of the neighbors had the same LCA as the whole neighborhood, we assumed that the other neighbor was the donor (see **Fig. 1**). This effectively selects the taxonomically smaller neighbor as the donor. Note that this procedure only approximates the donor group in case of additional evolutionary events or gene tree inference errors influencing the neighboring clades' LCAs. In case when both neighbors had different LCAs than their joint LCA, we assumed that the clade has an undetermined origin and discarded it from the analysis. Such situations can arise in complex evolutionary scenarios with multiple transfers. See the Supplementary Material for a more detailed description of the rationale.

Fig. 1. An example of a displaced fungal clade indicative of a horizontal gene transfer. The fungal clade branches with a {C1} and a {C2, C3} clade in the species tree, and with a {C3} and a {C1, C2} clade in the gene tree, indicating a different location of the clade in both trees. The correct location of the fungal clade in the gene tree is given by the branch adjacent to the {C1} clade. The clade {C3} can be inferred as the donor group because the species LCA of the {C1, C2} clade is the same as the species LCA of the whole neighborhood {C1, C2, C3}. In this work, we define clades only by their species compositions, as clades may have different topologies in both trees due to gene tree inference errors and additional evolutionary events. Trees with an average of the three Ultrafast Bootstrap supports UFB_1, UFB_2 and UFB_3 below 90% are discarded in this study.

Analysis of xenolog features

Xenolog protein products were mapped against PFAM (El-Gebali et al. 2019) using pfamscan.pl with default parameters (only hits with significance=1 taken), mapped against KEGG using kofamscan (Aramaki et al. 2020), scanned for signal peptides with TargetP (Armenteros et al. 2019) and cellular localization with WolfPsort II (Horton et al. 2007). For the latter two methods, the result for a given fungal clade was assumed to be consistent if it was the same for at least 75% of the clade's proteins. Low-complexity regions were predicted with seg (J. C. Wootton and Federhen 1996) and intrinsically disordered regions with IUPred3 (Erdős, Pajkos, and Dosztányi 2021). A species tree of the 44 target fungi was inferred from whole proteomes with OMA2 based on OMA groups of orthologs (Altenhoff et

al. 2019). To assess the evolutionary history of xenologs after transfer, we have reconciled the fungal clades (rooted at the transfer acceptor node) with the OMA2 tree in the Duplication-Loss (DL) model using the ete3 Python package (Page and Charleston 1997). Statistical and phylogenetic analyses were performed with the R statistical package (https://www.R-project.org/) and the Python 3 programming language using the Jupyter Notebook (Beg et al. 2021) and numpy (Harris et al. 2020), scipy (Virtanen et al. 2020) and ete3 (Huerta-Cepas, Serra, and Bork 2016) libraries. Plots were created using the matplotlib (Bisong 2019) library.

Results

We have detected 829 xenologs in the 44 target fungi with well-supported phylogenetic evidence for a horizontal gene transfer. The xenologs were nested within 226 clades in 206 phylogenetic trees. Including other fungal species, the 226 clades contained 1208 fungal xenologs. For over 50% of trees, iqTree selected the LG model as the best-fitting one, followed by WAG, VT and JTT models, which jointly accounted for 90% of clusters (**Supplementary Fig. S1**). Nuclear models accounted for 95% of clusters, followed by chloroplast (3%), mitochondrial (1.3%), viral (0.4%).

Additional 283 trees with 326 fungal clades containing 254 target xenologs and 1857 xenologs from other fungal species had weakly supported or incomplete phylogenetic evidence for HGT and were not used for the statistical analysis of patterns of HGT. The causes for lack of support are summarized per clade in **Supplementary Fig. S1.** Finally, 109 trees contained 123 fungal clades with 282 target proteins which had a congruent location on the gene tree, indicative of a simple vertical inheritance, sequence homoplasy or HGT from fungi rather than HGT into fungi. Note that these scenarios are not mutually exclusive in gene trees, as a single tree may contain multiple clades with strong and weak support for HGT as well as clades in congruent locations.

The list of all identified xenologs, including ones discarded from analysis due to insufficient support, is available in the Supplementary Material. In the remaining part of this work, unless explicitly stated otherwise, by fungal clades we understand clades of fungal proteins in gene trees with well-supported phylogenetic evidence for HGT, and by xenologs we understand proteins from target fungi in those clades. In this work, we refrain from using the term "transfer event" commonly used in bioinformatic literature, because it suggests one biological event of gene exchange which may result in transferring multiple genes, and its connection with protein families and gene tree clades is difficult to establish.

Genome contamination is an order of magnitude more common than HGT. In this work, we have adopted a two-stage blastp-based contamination filtering. In the first step, we used a coarse filter in which we discarded contigs without any detectable homology to the other 44 target fungi. Since this filter is insufficient to detect contaminants with distant fungal homologs, we then refined the filtering by discarding contigs in which we did not find a protein with a fungal top blast hit (excluding protein's host).

The first step removed 19578 proteins (**Supplementary Fig. S2**). This constitutes 3% of the 44 proteomes jointly, and 2% of each proteome on average. The number of removed proteins varied highly between proteomes, from 1 in *Absidia repens* NRRL 1336 and *Spizellomyces punctatus* DAOM BR117 through 203 in *Basidiobolus meristosporus* CBS 931.73 up to 5426 in *Rhizophagus irregularis* A1 and *Rhizophagus irregularis* DAOM 197198w. The extensive contamination of the *Rhizophagus irregularis* A1 proteome is in part likely caused by an associated *Lysinibacillus* bacterium.

The second, refined step removed additional 292 proteins. Here, the removed numbers varied to a lesser extent, with the largest being 107 for *Batrachochytrium dendrobatidis* JAM81; notably, the first stage discarded only 36 proteins in this fungus. This was followed by 57 and 53 proteins further removed from *Rhizophagus irregularis* DAOM 197198w and *Rhizophagus irregularis* A1. In all the remaining fungi, the second stage removed less than 10 proteins, suggesting that the initial coarse filter was mostly sufficient to discard contaminants in most fungi. The numbers of sequences discarded in both stages were highly correlated across the 44 proteomes (ρ=0.95), indicating that genome quality is the deciding factor.

The evolution of the number of xenologs after transfer can be explained with a simple but unexpected stochastic model. Over half of the fungal clades contained a single fungal protein (**Fig. 2**). If the numbers of xenologs evolved under a neutral birth-and-death model of gene duplications and losses, the distribution of clade size should follow a geometric distribution (Nee, May, and Harvey 1994). However, this seems not to be the case, and the chi-square test of goodness of fit performed on clades with 1 to 12 proteins rejects the hypothesis of a geometric distribution (p<1e-06; ML estimated geometric distribution parameter 0.47). Instead, the distribution of clade sizes appears to closely follow the Yule-Simon distribution with parameter alpha=1 (chi-square test p=0.40; **Fig. 2**), originally derived to describe the number of species in a randomly selected genus (E., F., and Udny Yule 1925; Steel and McKenzie 2001). This distribution differs qualitatively from the geometric one by having a heavy tail, i.e. noticeable occurrences of extremely large values. Those values can in turn be explained by a Yule-Simon distribution of the number of post-transfer duplications (chi-square p=0.33 for distribution parameter alpha=2; **Fig. 2**). This suggests a *rich get richer* dynamics of the evolution of the number of xenologs, where a xenolog in an already large clade is more likely to undergo a gene duplication than a xenolog in a smaller clade.

Characteristically for heavy-tailed distributions, while 80% of clades in our data set contained up to three fungal proteins, 1.3% contained more than 50, with the largest one containing 219 *Mucoromycota* xenologs (26 in the target fungi). The target proteins from this clade contain only an uncharacterized domain (DUF4379); other proteins from this cluster are involved in DNA repair and recombination (K03746). The distribution of the number of post-transfer duplications within the target fungi also had a heavy tail, explaining the occurrence of large clusters. For 86% of the fungal clades, there was up to 1 duplication per speciation event within the target fungi, while for 1.3% there were more than 20.

The number of xenologs is highly divergent across fungi. The number of xenologs in different proteomes was also heavily tailed, with 5 fungal species accounting for over 80% of all detected xenologs (**Supplementary Fig. S3**). These included the gut-dwelling *Neocallimastigomycota* (with xenologs constituting more than 1% of their proteomes), the chytrid *Rhizoclosmatium globosum* (0.5%), and the soil saprotrophic *Mortierella elongata* (0.5%). These groups were followed by the plant-symbiotic *Glomeromycota* and parasitic *Chytridiomycota* (with xenologs forming less than 0.2% of their proteomes). We did not observe any statistically significant correlation between the proteome size and the number of xenologs (ρ=0.19, p=0.21).

The heavy-tailed distribution of post-transfer duplications only partially explains these numbers. Each of the aforementioned fungi seems to have its own evolutionary and ecological mechanism for acquiring large numbers of xenologs. In chytrid species, each strain was the recipient of multiple transfers, while the xenologs in *Neocallimastigomycota* were mostly transferred to the ancestor of the whole group. Both groups were involved in multiple HGT events, while the large number of xenologs in *Mortierella elongata* was caused by only a few transfers and extensive post-transfer duplications.

The average proportion of xenologs in proteomes was 0.12%. This figure is within the range of 0.00% to 0.38% (with an identical per-genome average) reported previously for Dikarya (Marcet-Houben and Gabaldón 2010). We remark, however, that describing heavily tailed distributions with averages is highly inaccurate and may be misleading. The average proportion of xenologs may indeed be 0.12%, but, according to our data, a typical fungus is expected to have none or few at most. Furthermore, this proportion is expected to vary highly between reports due to sensitivity to dataset selection.

Fig. 2. The rates of HGT and post-transfer duplications among the 44 EDF. A) The number of post-transfer duplications among the target fungi. The red line on the log-log plot corresponds to the Yule-Simons distribution with parameter alpha=2. B) The OMA2 phylogenetic tree of fungi annotated with the numbers of clades of xenologs. *Method threshold:* HGTs older than the dashed line are rarely detected with our method. C) The number of clades of xenologs per acceptor-donor pair. Taxonomic names in B) and C) are based on NCBI Taxonomy with manual fine-tuning according to Voigt et al. (Voigt et al. 2021).

Transfer rates are highly divergent across the fungal tree of life and do not always reflect the numbers of xenologs in the proteome. To analyze the distribution of HGTs across the fungi, we have generated an OMA2 phylogenomic tree of the 44 EDFs and labelled it with the numbers of transferred protein clades (**Fig. 2**). Note that this procedure only approximates the depth of the transfer in the tree due to possible post-transfer gene losses or incomplete taxon sampling (making the transfer appear more recent), or inter-fungal transfers after gene acquisition (making the transfer appear more ancient).

The resulting distribution of HGTs across the fungi was highly uneven. There was a clear pattern dividing ancestrally aquatic taxa grouped in *Chytridiomycota* and *Neocallimastigomycota* (living in the digestive tract of terrestrial herbivores) with higher HGT rates compared to terrestrial fungi. The mainly saprotrophic *Mucoromycota*, despite the highest sampling in our dataset, shows the lowest number of HGT events per genome, just

as the animal-related *Kickxellomycota* fungi. *Mortierellomycota*, despite *Mortierella elongata* having one of the highest numbers of xenologs in our data set, appears to have rather low transfer rates, just as the closely related *Glomeromycota*. As opposed to ancestrally aquatic fungi, post-transfer duplication seems to be the major force driving the number of xenologs in these fungi.

Transfer rates reflect physical proximity. Diverse bacteria were the most common donors of xenologs, including *Firmicutes* (most prominently *Clostridiaceae, Lachnospiraceae, Eubacteriaceae, Oscillospiraceae* and *Bacilli*), *Proteobacteria* (*Gammaproteobacteria, Burkholderiales*) and *Actinobacteria*. These were followed by eukaryotic Oomycota, Metazoa and Viridiplantae, and only a few transfers from Archaea (**Fig. 2**). In agreement with previous results (Alexandre et al. 2021), in *Neocallimastigomycota* we have detected genes from a myriad of diverse bacteria, including *Bacteroidetes, Bacilli (Firmicutes), Slackia (Actinomycetota)* and *Aeromonas (Proteobacteria)*. The *Firmicutes* bacteria were a particularly characteristic donor for these fungi*,* with numerous genes obtained from the anaerobic *Clostridia*, which, according to previous reports, may have enabled them to adapt to the gut environment (Murphy et al. 2019).

For *Mortierellomycota* fungi, where *Burkholderia*-related endobacteria (BRE) are common, *Burkholderiales* stand out as one of the main sources of xenologs. Gene donors include *Mycoavidus cysteinexigens* (OAQ22128.1)*,* an endohyphal symbiont of *Linnemannia elongata* (J. Uehling et al. 2017), and *Mycetohabitans rhizoxinica* species (OAQ29846.1), which were previously described as endohyphal symbionts of *Rhizopus microsporus*.
The second main source of transfers to *Mortierellomycota* were the soil-inhabiting *Actinomycetia*. This bacterial group was recently reported as the most prevalently associated with fungal isolates (Robinson et al. 2021).

In the water-living *Chytridiomycota*, most xenologs come from loosely associated bacteria such as *Alphaproteobacteria, Gammaproteobacteria* and *Actinomycetia*. In addition, we have identified several transfers from Oomycota, including a tRNA (N6-threonylcarbamoyladenosine(37)-N6)-methyltransferase TrmO (ORY41869.1) and tandem tetratricopeptide repeat proteins (e.g. ORY27572.1). TrmOs are widely present in most domains of life but were apparently lost in Fungi. The transfer of the Oomycota gene seems to have compensated for this loss. The repeat proteins are predicted to form a protein-binding surface, but their function remains unknown.

***Neocallimastigomycota* appear unique in multiple independent transfers of homologous proteins.** Out of 206 trees in our data set, 187 contained a single fungal clade 18 contained two, and one contained three clades. With the single exception of one, overall rather weakly supported, tree with two *Conidiobolus* clades, all the trees with more than one fungal clade contained only *Neocallimastigomycota* species. In 7 trees with two fungal clades and in the single tree with three, all the clades contained different sets of species, indicating independent transfers of similar proteins after the divergence of *Neocallimastigomycota*. An example is an independent acquisition of proteins with a solute binding domain *SBP_bac_3* by *Neocallimastix californiae* G1 from *Oscillospiraceae* (ORY20588.1) and by *Anaeromyces robustus* from *Methanobacteriaceae* (ORX65467.1). On

the other hand, in four trees, the two clades contained identical species, indicating two independent transfers to the ancestral lineage of *Neocallimastigomycota*. An example is a repeated acquisition of a bacterial cellulase by the ancestor of *Neocallimastigomycota*, once from *Bacilli* (e.g. *Neocallimastix californiae* G1 ORY56492.1) and once from *Clostridia* (e.g. *Neocallimastix californiae* G1 ORY21818.1). This result stands in agreement with a recently reported major HGT event in the ancestor of *Neocallimastigomycota* that might have driven its evolution towards a gut-dwelling fungus of herbivores (Murphy et al. 2019). In the remaining 7 trees, the species composition overlapped partially between the two clades, indicating a repeated ancestral transfer followed by gene losses.

Low-complexity sequences are rarely transferred. The distributions of the proportions of protein sequences covered by low-complexity regions, as well as intrinsically unstructured regions, have lighter tails for xenologs than for the proteome background (**Fig. 3**). A comparison of the average low-complexity proportions of fungal xenologs to the donor group showed no visible trend (**Supplementary Fig. S4**). This suggests that extensive low-complexity or unstructured regions tend to inhibit transfers, rather than being caused by them. Note, however, that it may be an artifact caused by impared homology detection between low-complexity regions (Jarnot et al. 2022).

The distribution of xenolog sequence length follows the background up to around 1800 amino acids and drops sharply afterwards. This seems to be an artifact of our method, which puts a threshold on xenolog sequence length at 2000 aa. Individual cases of transfers of much longer sequences are documented, for instance, secondary metabolic clusters in *Basidiobolus* and multiple Dikarya (Tabima et al. 2020; Wisecaver, Slot, and Rokas 2014). As a consequence, we cannot conclude whether the distribution of sequence length of xenologs follows the protein background or not and whether sequence length poses a barrier for HGT in fungi. A comparison of the average sequence length of xenologs to the donor group showed no visible trend of shortening or lengthening of sequences post-transfer, but revealed differences of up to 400 amino acids in both ways (**Supplementary Fig. S4**).

Xenologs are secreted more often than typical fungal proteins, but determining their precise subcellular location may be challenging. TargetP 2.0 detected a signal peptide in 33% of identified xenologs, compared to 7.7% of the proteome background. This stands in agreement with the osmotrophic lifestyle of fungi, their massive secretome, and the accessory function of many xenologs. Consistently with the bacterial origin of many xenologs, only 9 xenologs had mitochondrial transfer peptides. We did not observe biologically meaningful differences in the numbers of transmembrane helices predicted by TMHMM, although a chi-square test weakly rejects the hypothesis of identical distributions in xenologs and the background (p=0.03, ddof=0; assuming xenolog observations as observed, background as expected).

The subcellular localization predicted by WolfPsort was consistent for 77% of target protein clades, compared to 95% for TargetP (**Fig. 3**, **Supplementary Fig. S5**). Compared to the proteome background, xenologs are predicted to localize more often in the extracellular space, the endoplasmic reticulum, and the cytoplasm, and less often in the nucleus and the mitochondria. Most xenologs localize in the extracellular space, closely followed by the

nucleus. In contrast to TargetP, WolfPSort predicts 90 proteins to localize in the mitochondria. The unexpectedly high number of mitochondrial and nuclear predictions may suggest that WolfPsort is biased towards these locations. To verify this hypothesis, we have generated a decoy dataset by randomly permuting each sequence from the 44 proteomes. The predicted distribution of locations is very similar for the proteome background and the decoy dataset, with the nucleus as the most common location, and the decoy proteins attain similar prediction scores (**Supplementary Fig. S5**). This is likely caused by the fact that WolfPSort is based on the *k*-Nearest Neighbours classification algorithm, which tends to reflect the distribution of the training dataset, in this case, fungal proteomes. The prediction may be further complicated by the fact that xenolog sequences have a non-fungal origin.

Fig. 3: Molecular characteristics of xenolog protein products (orange) and background proteins (blue). A) Quantile plots comparing the xenolog and background sequence low-complexity fraction, intrinsically unstructured fraction, and sequence length. B) Subcellular localizations. C) The distribution of the intron count in xenologs for selected fungi. D) The overall distribution of the number of introns in xenologs and proteome

background. E) The number of introns and their dependence on the kingdom of the donor. F) The number of PFAM domains.

Xenologs typically contain few introns but can acquire many. Over 50% of xenologs did not contain any intron, and almost 90% contained at most two, compared to respectively 25% and 59% in the background (**Fig. 3**). This result is in line with recommended HGT verification measures (Jaramillo et al. 2015). However, we have also detected 16 xenologs, mostly of eukaryotic origin, with 6 or more introns (**Fig. 3**). In particular, the *Glomeromycota* fungus *Rhizophagus irregularis DAOM 197198w* and the chytrid fungi *Gonapodya prolifera*, *Batrachochytrium dendrobatidis JAM81* and, to some extent, *Rhizoclosmatium globosum* are prone to have intron-rich xenologs from both unicellular eukaryotic and prokaryotic donors. Although there are chytrid species referred to as *intron-rich chytrids* (van de Vossenberg et al. 2019), neither *G. prolifera* nor *R. globosum* have been described as such in their genomic papers (Y. Chang et al. 2015; Mondo et al. 2017). The preference for the intron structure of xenologs seems to some extent correlated taxonomically: while the xenologs in *Chytridiomycota* tend to contain multiple introns, the ones in *Neocallimastigomycota* tend to contain few if any.

The number of introns closely follows a geometric distribution in the fungal proteome background, but not in the set of xenologs. In particular, the distribution in xenologs is zero-inflated, consistent with the bacterial origin of many xenologs. However, many bacteria-derived xenologs contain multiple introns, suggesting that they acquire introns after the transfer, consistent with previous reports (Lage et al. 2013). This seems to be the case in an *R. globosum* xenolog with twelve introns (ORY15655.1), derived from a deltaproteobacterium *Labilithrix luteola* magnesium transporter. This protein contains three post-transfer paralogs, out of which one contains only one intron, one contains five, and one contains six (respectively ORY29294.1, ORY29292.1 and ORY15656.1). A similar scenario seems to have shaped the story of arginine decarboxylase biosynthesis gene, which was transferred from an intronless gene of an anaerobic deltaproteobacterium *Anaeromyxobacter dehalogenans* (locus_tag A2cp1_2068) and acquired nine introns after transfer to *R. globosum* (KXS13513.1). In addition to the intronization of bacteria-derived xenologs, we have also observed an apparent intron gain in an E3 ubiquitin-protein ligase gene with 8 introns, transferred from some early diverged metazoans to *Glomeromycota* (EXX59914.1). The closest metazoan homologs from *Acropora millepora* (LOC114972538) and *Orbicella faveolata* (LOC110047255) have only three or four introns.

To test the hypothesis of intron acquisition in bacteria-derived xenologs, we have compared the number of introns to the branch distance to the root of the corresponding fungal clade (i.e. the post-transfer number of substitutions per site). A linear model fitted without an intercept estimated 0.6 introns per unit branch length (p < 0.001). However, the overall picture is more complicated, as some xenologs quickly acquire multiple introns, while many others have none even after a long time.

Gene transfer seems to frequently involve acquiring and losing protein domains. Compared to the proteome background, fungal sequences obtained *via* HGT are more likely to map to one or more distinct PFAM domains (**Fig. 3**). However, a considerable number of

those domains seems to be acquired after a transfer, since 100 fungal xenologs had domains which had no clan-level siblings in non-fungal sequences from the same gene tree. Most xenologs acquired a single domain, while 10 acquired two distinct domains. Except for one xenolog in *Mortierella elongata* AG 77 and five xenologs in chytrids, all the post-transfer domain acquisitions happened in *Neocallimastigomycota*. As therefore expected, the by far most commonly added domain was the carbohydrate-binding module CBM_10 (PF02013), a part of the dockerin cellulosome typical of anaerobic fungi, acquired in 86 xenologs. Consistent with the function of CBM_10, this domain was fused preferentially to xenologs containing domains associated with carbohydrate metabolism, such as the bacterial glycosyl hydrolase (PF02011) in 31 xenologs. Some exceptions to this rule involved 13 xenologs with a bacterial CotH kinase domain, the main invasin involved in mucormycosis which binds to glucose-regulated protein 78 (GRP78) on nasal epithelial cells (Nguyen et al. 2016; Alqarihi et al. 2020). The cellulose-binding domain together with a CotH kinase may play a role in the interaction with the animal host. The next most commonly acquired domain was the ricin B lectin domain (PF14200) acquired in nine *Neocallimastigomycota* xenologs.

Some domains were seemingly also lost during transfer, with 89 xenologs not containing domains which were present in at least 20% of sequences in both neighboring gene tree clades. An example is a bacteria-derived cellulase precursor ORX60185.1 in *Piromyces finnis*, which is missing an N-terminal fragment of its bacterial homologs with a cellulose binding domain CBM_2 and a Bacterial Ig domain Big_7. On the other hand, this protein contains a bacterial glycoside hydrolase domain Glyco_hydro_48 and is post-transfer fused with a cellulose binding domain CBM_10. Effectively, a bacterial carbohydrate-binding domain was replaced with a fungal one in this protein.

Overall, out of 829 xenologs detected in our target fungi, 170 had a novel or a lost domain, and 48 had both (such as the aforementioned cellulase precursor in *Piromyces finnis*). In some cases, this resulted in conflicting phylogenetic signals. The *Rhizoclosmatium globosum* ORY15655.1 (one of the aforementioned intron-rich xenologs) contains a xenologous N-terminal region of 530 aa of bacterial origin with a glycoside hydrolase domain (InterPro *Glycoside_hydrolase_SF*) and an apparently ancestrally fungal C-terminal region of 180 aa with a mainly eukaryotic magnesium transporter NIPA domain Mg_trans_NIPA. We note that this apparent gene fusion may simply be a gene calling error merging two neighboring genes (frequently encountered in genomes without transcriptomic evidence and submitted as drafts), and requires further proof for confirmation. Regardless of the actual source of the apparently ancestrally fungal region, the presence of conflicting signals in this protein's sequence severely impacts the topology of the phylogenetic tree (**Supplementary Fig. S6**).

Notably, the 12 introns of this xenolog are located in both the ancestrally fungal and the bacteria-derived region. The three putative post-transfer paralogs of this protein, with fewer introns, lack the apparently ancestral C-terminal region with the magnesium transporter domain (**Fig. 4**). It is unknown whether the presence of an ancestrally fungal genetic material has facilitated the intronization of ORY15655.1.

Fig. 4: A post-transfer fusion of a fungal domain with a xenolog. Plots visualize the taxonomic composition of the first 250 BlastP hits per amino acid of the query protein. The ORY15655.1 protein contains a distinct C-terminal domain which is likely ancestrally fungal. The post-transfer paralogs of this protein have a similar taxonomic distribution of their BlastP hits at each amino acid but lack the domain.

Metabolic domains constitute the majority of xenologs. To analyze HGT-derived protein domains of fungal xenologs, we have selected domains which were present in at least half of fungal proteins and in at least 20% of proteins in one of the neighboring clades. Domains of unknown function (DUFs/UPFs) constituted a numerous functional class with 16 diverse Pfam families found in 60 xenologs. Among domains with known functions we have found numerous transfers of genes involved in metabolism, in particular coding abhydrolases, acetyltransferases and esterases. Many carbohydrate active enzymes, such as glycohydrolases and glycosyltransferases, were transferred to *Neocallimastigomycota*. On the other hand, peptidases were underrepresented in the dataset (peptidase xenologs n=24) compared to the carbohydrate active enzymes and carbohydrate binding modules (n=63), but were transferred to diverse recipients. They include trypsins transferred from *Phytohabitans flavus* to *Spizellomyces punctatus* (KNC97026.1) and from *Araneus ventricosus* to *Conidiobolus coronatus* (KXN68345.1) and a membrane-bound dipeptidase, metalloproteinase M19, transferred from *Micromonosporaceae* to *G. prolifera* JEL478 (KXS08983.1). Trypsin expansions can be linked to enhanced degradation of protein-rich substrates such as insects, while a M19 peptidase is part of an *Aspergillus* spp. gliotoxin biosynthesis cluster. However, the *Aspergillus* spp. M19 peptidase is likely ancestrally fungal and distant to the *G. prolifera* M19 peptidase*.

The dominance of carbohydrate active enzyme encoding genes over protein metabolism genes in our data set can be linked to the high number of transfers to *Neocallimastigomycota*. These fungi also acquired genes with a myriad of specificities connected to secondary metabolism and transport, extending their nutritional capabilities. Condensation domain (PF00668) containing candidate NRPS secondary metabolite clusters were transferred from *Eubacteriales* to several *Neocallimastigomycota* (e.g. *Anaeromyces robustus* ORX81431.1). General transporters from MFS_1 and ABC families were acquired by *Neocallimastigomycota* (eg. ORY49205.1).

We have detected a few transfers (n=7) of mobile elements, including a transfer of a PiggyBac element from *Phytophthora* to *Smittium culicis* (OMJ27075.1) and of a retrotransposon from *Globisporangium splendens* to *Rhizoclosmatium globosum* (ORY50945.1). However, most of the potential transposon xenologs were likely excluded from our data set because of the usage of protein sequences instead of genomic sequences for HGT inference, as well as low support values and elevated branch lengths. We have also identified several phage proteins without clear function in fungi, eg. a phage tail lysozyme, PF18013, in *Piromyces finnis* protein ORX49608.1.

We have detected transfers of several genes involved in resistance, including a bacterial putative betalactamase resistance gene transferred from to *G. prolifera* JEL478 (KXS12969.1). *Mortierellomycota* acquired NOD-like receptors with NACHT-PPR-WD40 architecture from *Mycoavidus cysteinexigens* endosymbionts. NOD-like receptors are suspected to play a role in Dikarya immunity (Jessie Uehling, Deveau, and Paoletti 2017; Dyrka et al. 2014). Interestingly, all of the *Mortierellomycota* immunity proteins in our dataset seem to be of a bacterial origin or at least shared with the endosymbiont. *Mycoavidus* homologs have ~50-45% of sequence identity and are the best reciprocal blast hits. It is an open question whether these proteins have a role in the symbiotic interaction.

On the other side, we have detected transfers of several putative toxins. Several *Neocallimastigomycota* representatives gained ricin-type sugar binding modules (18 xenologs in a single gene tree clade; e.g. ORX83988.1). Similar proteins were recently documented in *Mucor* acting as toxins (mucoricin) (Soliman et al. 2021). Most of those domains were associated with bacterial glycoside hydrolase domains and nearly all were post-transfer fused with the carbohydrate binding CBM_10 domain. Additional nine xenologs in another gene tree appear to have gained the ricin-type sugar binding module post-transfer, suggesting a proliferation of a xenologous domain. Among other examples, a sea anemone pore-forming toxin (PF06369) was transferred from *Selaginella moellendorffii* to *R. irregularis* and is present in all three strains analyzed in this study (e.g. PKC66727.1).

Putative sequence homoplasy has a similar taxonomic distribution to HGTs. Annotation of the OMA2 tree of target fungi with the numbers of non-displaced fungal clades in trees with at least 60% non-fungal species and at least 60% sequences from non-fungal species (indicative of vertical inheritance or sequence homoplasy) reveals a striking similarity to taxonomic patterns of HGTs (**Supplementary Fig. S7**). The correlation between the numbers of displaced and non-displaced clades mapped to the nodes of the OMA2 tree was 0.80 (p < 1e-12). This suggests that, for fungal proteins with a predominantly non-fungal homology, even in the absence of tree incongruence a HGT may be a more likely hypothesis than vertical inheritance with multiple gene losses or sequence homoplasy. Although homoplasy has been reported to occur frequently on the scale of amino acids in Eukaryota (Rokas and Carroll 2008), it seems unlikely on the scale of proteins. Therefore, the 282 target proteins discarded in this work solely due to lack of tree incongruence are likely to represent additional cases of HGT. They include e.g. *Mortierellomycota* fatty acid desaturases similar to delta-12 fatty acid desaturases of bacterial origin, likely acquired from

Deltaproteobacteria (KFH67875.1). The representatives of this fungal group are known to produce long polyunsaturated fatty acids (L. Chang et al. 2021).

Discussion

Our aim in this work was to provide a statistical characterization of the patterns of interkingdom HGT across diverse EDF. Multiple factors can inflate the false positive rate of HGT detection in a high-throughput setting (see the **Supplementary Methods** for a detailed discussion), including, but not limited to:

- Relying on sequence similarity instead of phylogenetic trees (Koski and Brian Golding 2001)
- Excessively long, multidomain proteins, sometimes originating from ORF prediction errors (Poptsova and Gogarten 2010),
- Sequences without a well-defined location in the species tree (e.g. viral or environmental),
- Low-quality alignments and a wrong choice of the trimming software (Tan et al. 2015),
- Automatic rooting of gene trees (Lamarca, Mello, and Schrago 2022; Wade et al. 2020),
- Low-quality gene trees with low support values and long branches (Rodríguez-Ezpeleta et al. 2007),
- Using tree reconciliation algorithms with improper parameters (Libeskind-Hadas et al. 2014).

In order to avoid those pitfalls, we have applied a series of filters and developed a custom-made algorithm for the detection of gene trees with displaced fungal clades. Notably, our method allows us to use unrooted gene trees to detect transfers and approximately identify their donors and does not require the specification of any parameters. Our data and results should provide a reliable source of fungal xenologs and their properties which can further be used to benchmark new HGT identification methods (**Supplementary Table S1**). In particular, statistical properties such as the Yule-Simons distribution of the numbers of xenologs per transfer event could be used to validate future results in a rigorous, statistical framework. This would be a particularly valuable addition to other criteria if a mechanistic explanation of this distribution was provided and verified.

Our results suggest that the false positive risk factors listed above have a greatly varying impact on the quality of the results. Moreover, while there has been much research devoted to some, others seem to be underappreciated in the literature. Phylogenetic tree incongruence is widely considered the gold standard of HGT detection and has been the basis of our approach. However, the exceptionally high correlation between the numbers of congruent and incongruent trees per fungal taxon, combined with the lack of correlation between them and the recipient's proteome size, suggests that taxonomy-based methods such as the Alien Index may be equally reliable. The added benefit of phylogeny-based methods, however, is a more precise determination of the gene donors.

While tree-based methods seem to offer only a marginal improvement over taxonomy-based

heuristics, the filtering of contaminant sequences is of tremendous importance. We have detected nearly 20 000 likely contaminating sequences, 20 times more than the number of detected xenologs. As a consequence, a randomly selected gene with an atypical taxonomy of homologs or incongruent phylogeny is much more likely to be a contaminant than a xenolog. In EDF, genome contamination may occur due to sample contamination or the presence of endohyphal bacteria. The ultimate determination of whether a gene is incorporated into the genome and expressed requires experimental evidence; in bioinformatic analysis, the minimum requirement should be the presence of ancestral genes in the same contig. Heuristic approaches based on sequence identity and GC content may be insufficient to separate xenologs from contaminants, especially in the case of genes transferred from endosymbionts. The picture is further complicated by the fact that fungal endosymbionts are still understudied, and most likely there still are many unknown endohyphal organisms (Robinson et al. 2021).

One of the most predictable features of EDF HGT appears to be its unpredictability. The number of xenologs in a given proteome is governed by at least two burst-like events: bursts of gene exchange like in *Neocallimastigomycota* and bursts of gene duplications like in *Mortierella* sp. This agrees with unexpectedly high numbers of xenologs in some metazoan proteomes, like *Rotifera* sp. Nevertheless, thanks to separating HGTs from contaminants, our results allowed us to detect some clear patterns in the properties of protein products of horizontally transferred genes. Xenologs are characterized by relatively short low-complexity and intrinsically unstructured regions. The depletion of low-complexity regions can be related to the so-called complexity hypothesis, which states that xenologs tend to be far from central hubs of interaction networks (Cohen, Gophna, and Pupko 2011), and instead participate in secondary metabolism, drug resistance etc. On the other hand, regions with a skewed amino acid abundance, which constitute 20% of eukaryotic proteins (Mier et al. 2020), are involved in diverse molecular functions usually linked to regulation (Wright and Dyson 2015) and protein-protein interactions (Coletta et al. 2010).

In agreement with the current notions, the xenologs in our dataset tend to contain few introns. We have, however, detected notable exceptions to this rule in chytrids and some *Glomeromycota*. This may result from the acquisition of intron-rich genes or the intronization of xenologs. The distribution of introns seems to be taxonomy-dependent, with certain fungal groups showing a preference for intronless xenologs, while others for intron-rich ones. Bacteria-derived xenologs acquire introns over time, but some are particularly resistant to intronization or not yet ameliorated to the fungal genome. Together with the different distributions of the numbers of introns in xenologs and the proteome background, this suggests that the intronization dynamics is more complex in xenologs than in fungal proteins.

The lack of correlation between the number of xenologs and proteome size implies that HGT has little or no influence on the proteome sizes of EDF. Moreover, large proteomes do not predispose to HGT and the overall amount of genetic material does not seem to be a limiting factor for the incorporation of transferred genes (except, perhaps, for species with a particularly high pressure for reduced genomes).

The receptiveness for foreign genes varies greatly between fungal lineages, with bacteria being the most common source of xenologs. In agreement with previous reports, we have found that fungi belonging to ancestrally aquatic groups experienced more HGT events than terrestrial lineages. This can be explained by the easier exchange of genetic material in the aquatic ecosystem, the ubiquity of unicellular aquatic life forms, and the possibility of forming biofilms with bacteria (McDaniel et al. 2010; Grüll, Mulligan, and Lang 2018; Abe, Nomura, and Suzuki 2020). Another possible explanation is that terrestrial fungi may have lost some of the HGT predisposing traits or gained defensive mechanisms preventing foreign gene incorporation. Despite the impact of the aquatic environment, the relationship with the host and the necessity to adapt to a new environment seem to have a higher influence on the rate of HGT. Within fungi with at least 10 xenologs, most have a documented long-term association with another organism. On the other hand, many endomycorrhizal fungi did not experience extensive transfers (or the transfers remained undetected, for instance due to the diversity of nuclei in Glomeromycota cells). These observations suggest that, while symbiosis or parasitism does not always result in gene exchange, it is a predisposing condition.

Some of the early diverging fungal lineages have associated bacteria, either endohyphal or loosely related to their hyphae, some of which are co-evolving with the fungal host and transmitted vertically (Pawlowska et al. 2018). Those bacteria have been previously pointed to as a likely source of xenologs (Naito, Morton, and Pawlowska 2015), because such close endosymbiotic relations pose a natural opportunity for a large-scale gene exchange, as exemplified by the massive gene transfer from the mitochondrion to the nucleus of eukaryotic organisms (Anselmetti et al. 2021). This pattern was confirmed in this study, with a notable example of *Podila verticillata* acquiring fatty acid desaturases probably from *Deltaproteobacteria* and NOD-like receptors from endohyphal *Burkholderia*-related endosymbiont *Mycoavidus* sp. The latter suggests a parallel evolution of NOD-like receptors and perhaps a NOD-like receptor-based immunity in Dikarya and *Mortierellomycotina*. Moreover, NOD-like receptors may be involved in the host-bacteria interaction. Our results also confirm the reports on a massive enzyme encoding gene acquisition by *Neocallimastigomycota* (Murphy et al. 2019) which has likely played a role in their transition to a gut-dwelling mode of life.

Out of 829 xenologs detected in our target fungi, 100 had apparently post-transfer fused domains. This may point to an unexpectedly high frequency of gene fusions occurring during or after the integration of foreign DNA. Cross-kingdom HGT has been reported previously as an important factor in gene fusion (Yanai, Wolf, and Koonin 2002). As the foreign genetic material is incorporated into the host's genome, it can be located within an existing gene. Alternatively, homologous or non-homologous recombination can occur after the new gene is incorporated, especially if it contains a fragment similar to the host's genes. While it is unclear which mechanism was responsible for the swapping of the bacterial cellulose binding domain with a fungal one in ORX60185.1, a replacement of a domain which preserves the protein's function agrees with the current notions about the evolution of domain architectures.

Note that the 100 fusions were detected only based on identified domains; since many sequences lack identified domains, the full extent of post-transfer gene fusion is unknown.

Frequent fusions may significantly complicate the studies of HGT. Recombination of transferred and ancestral genetic material may, on one hand, confound the impact of HGT on the host's ecology and evolution, and on the other hand, the proteins' phylogenetic signals. While most research to date, including this one, has focused on detecting "whole-protein" xenologs, our results indicate the need to develop domain-level methods of xenology detection and investigate their results. This agrees with the recent ideas that although terms like orthology, paralogy or xenology are traditionally applied on a level of whole genes or proteins, they should perhaps also be considered on a domain-level basis (Linard et al. 2021; Persson et al. 2019; Sjölander et al. 2011; Forslund et al. 2018). Domain-level HGT detection methods may also be more robust against gene calling errors merging neighboring genes or missing their parts. We also note that sequence regions with independent evolutionary histories are another reason why the first blast hit may not be the phylogenetically closest neighbor, as a small but highly conserved region may have a smaller E-value than a large but less conserved one.

In this work, we have focused on xenologs which homology is confined to a single taxonomic family of fungi. This was done in order to obtain clear-cut cases of HGTs in simple gene trees, and therefore to mitigate the false positive rates. However, discarding proteins with homology to more than one fungal family means that our methodology mostly detects relatively recent transfers from distantly related donors. In particular, it does not detect transfers between fungal species, such as the massive transfer from a fungal parasite *Parasitella parasitica* to its host *Absidia glauca (Kellner et al. 1993)*, and may also have a limited sensitivity towards transfers of animal genes, many of which have a fungal homolog. As a consequence, while our results confirm the occurrence of cross-kingdom gene transfers between eukaryotes (Schönknecht, Weber, and Lercher 2014; Van Etten and Bhattacharya 2020; Richards et al. 2009), they may significantly underestimate their prevalence.

The list of xenologs reported in this work is far from exhaustive. We have found several probable cases of HGT among gene trees discarded due to different reasons in different steps of our pipeline. The ORY16048.1 protein in *Rhizoclosmatium globosum* is a plausible transfer from mosquitoes, which occupy a similar ecological niche as the recipient fungus in their larval states; however, the protein has a low identity to its closest homolog and therefore has been discarded in the long branch cutting stage. One of the possible reasons for low identity is the absence of the actual donor in the NR database; accordingly, progress in genome sequencing efforts is likely to enable the discovery and confirmation of many new HGTs.

Acknowledgements
This work was supported by National Science Centre grants #2017/25/B/NZ2/01880 to A.M., #2019/33/B/ST6/00737 to P.G., #2017/25/B/NZ8/00473 to J.P.

Author Contributions

M.C, P.G., J.P. and A.M. designed the study, M.C. and P.G. developed the algorithms, M.C. implemented the software, M.C. and A.M. prepared the dataset and performed genome

analyses, M.C. performed the statistical analyses. All authors have interpreted the data and wrote the manuscript.

Data Availability

Phylogenetic trees of supported and unsupported HGT are deposited under 10.5281/zenodo.7461853, all accessions and assemblies are listed in Supplementary Table S1.

References

Abe, Kimihiro, Nobuhiko Nomura, and Satoru Suzuki. 2020. "Biofilms: Hot Spots of Horizontal Gene Transfer (HGT) in Aquatic Environments, with a Focus on a New HGT Mechanism." *FEMS Microbiology Ecology* 96 (5). https://doi.org/10.1093/femsec/fiaa031.

Alexander, William G., Jennifer H. Wisecaver, Antonis Rokas, and Chris Todd Hittinger. 2016. "Horizontally Acquired Genes in Early-Diverging Pathogenic Fungi Enable the Use of Host Nucleosides and Nucleotides." *Proceedings of the National Academy of Sciences of the United States of America* 113 (15): 4116–21.

Alexandre, Flávia Sampaio, Larissa Venturini Della Flora, Ivanildo Guilherme Henrique, Daniely Camila da Silva, Andreza Pereira Mercedes, Aline Cardoso Silva, Afonso Silva de Oliveira, et al. 2021. "Arbuscular Mycorrhizal Fungi (Rhizophagus Clarus) and Rhizobacteria (Bacillus Subtilis) Can Improve the Clonal Propagation and Development of Teak for Commercial Plantings." *Frontiers in Plant Science*. https://doi.org/10.3389/fpls.2021.628769.

Alqarihi, Abdullah, Teclegiorgis Gebremariam, Yiyou Gu, Marc Swidergall, Sondus Alkhazraji, Sameh S. M. Soliman, Vincent M. Bruno, et al. 2020. "GRP78 and Integrins Play Different Roles in Host Cell Invasion during Mucormycosis." *mBio* 11 (3). https://doi.org/10.1128/mBio.01087-20.

Altenhoff, Adrian M., Jeremy Levy, Magdalena Zarowiecki, Bartłomiej Tomiczek, Alex Warwick Vesztrocy, Daniel A. Dalquen, Steven Müller, et al. 2019. "OMA Standalone: Orthology Inference among Public and Custom Genomes and Transcriptomes." *Genome Research* 29 (7): 1152–63.

Ambrose, Karen V., Albrecht M. Koppenhöfer, and Faith C. Belanger. 2014. "Horizontal Gene Transfer of a Bacterial Insect Toxin Gene into the Epichloë Fungal Symbionts of Grasses." *Scientific Reports* 4 (July): 5562.

Anselmetti, Yoann, Nadia El-Mabrouk, Manuel Lafond, and Aïda Ouangraoua. 2021. "Gene Tree and Species Tree Reconciliation with Endosymbiotic Gene Transfer." *Bioinformatics*. https://doi.org/10.1093/bioinformatics/btab328.

Aramaki, Takuya, Romain Blanc-Mathieu, Hisashi Endo, Koichi Ohkubo, Minoru Kanehisa, Susumu Goto, and Hiroyuki Ogata. 2020. "KofamKOALA: KEGG Ortholog Assignment Based on Profile HMM and Adaptive Score Threshold." *Bioinformatics* 36 (7): 2251–52.

Armenteros, José Juan Almagro, Konstantinos D. Tsirigos, Casper Kaae Sønderby, Thomas Nordahl Petersen, Ole Winther, Søren Brunak, Gunnar von Heijne, and Henrik Nielsen. 2019. "SignalP 5.0 Improves Signal Peptide Predictions Using Deep Neural Networks." *Nature Biotechnology*. https://doi.org/10.1038/s41587-019-0036-z.

Beg, Marijan, Juliette Taka, Thomas Kluyver, Alexander Konovalov, Min Ragan-Kelley, Nicolas M. Thiery, and Hans Fangohr. 2021. "Using Jupyter for Reproducible Scientific

Workflows." *Computing in Science & Engineering*. https://doi.org/10.1109/mcse.2021.3052101.

Bisong, Ekaba. 2019. "Matplotlib and Seaborn." *Building Machine Learning and Deep Learning Models on Google Cloud Platform*. https://doi.org/10.1007/978-1-4842-4470-8_12.

Chang, Lulu, Haiqin Chen, Xin Tang, Jianxin Zhao, Hao Zhang, Yong Q. Chen, and Wei Chen. 2021. "Advances in Improving the Biotechnological Application of Oleaginous Fungus Mortierella Alpina." *Applied Microbiology and Biotechnology* 105 (16-17): 6275–89.

Chang, Ying, Sishuo Wang, Satoshi Sekimoto, Andrea L. Aerts, Cindy Choi, Alicia Clum, Kurt M. LaButti, et al. 2015. "Phylogenomic Analyses Indicate That Early Fungi Evolved Digesting Cell Walls of Algal Ancestors of Land Plants." *Genome Biology and Evolution* 7 (6): 1590–1601.

Clutterbuck, A. John, and A. John Clutterbuck. 2011. "Genomic Evidence of Repeat-Induced Point Mutation (RIP) in Filamentous Ascomycetes." *Fungal Genetics and Biology*. https://doi.org/10.1016/j.fgb.2010.09.002.

Coletta, Alain, John W. Pinney, David Y. Weiss Solís, James Marsh, Steve R. Pettifer, and Teresa K. Attwood. 2010. "Low-Complexity Regions within Protein Sequences Have Position-Dependent Roles." *BMC Systems Biology*. https://doi.org/10.1186/1752-0509-4-43.

Coyte, Katharine Z., Cagla Stevenson, Christopher G. Knight, Ellie Harrison, James P. J. Hall, and Michael A. Brockhurst. 2022. "Horizontal Gene Transfer and Ecological Interactions Jointly Control Microbiome Stability." *PLoS Biology* 20 (11): e3001847.

Dyrka, Witold, Marina Lamacchia, Pascal Durrens, Bostjan Kobe, Asen Daskalov, Matthieu Paoletti, David J. Sherman, and Sven J. Saupe. 2014. "Diversity and Variability of NOD-like Receptors in Fungi." *Genome Biology and Evolution* 6 (12): 3137–58.

E., F. Y., Y. E. F., and G. Udny Yule. 1925. "A Mathematical Theory of Evolution Based on the Conclusions of Dr. J. C. Willis, F.R.S." *Journal of the Royal Statistical Society*. https://doi.org/10.2307/2341419.

El-Gebali, Sara, Jaina Mistry, Alex Bateman, Sean R. Eddy, Aurélien Luciani, Simon C. Potter, Matloob Qureshi, et al. 2019. "The Pfam Protein Families Database in 2019." *Nucleic Acids Research*. https://doi.org/10.1093/nar/gky995.

Erdős, Gábor, Mátyás Pajkos, and Zsuzsanna Dosztányi. 2021. "IUPred3: Prediction of Protein Disorder Enhanced with Unambiguous Experimental Annotation and Visualization of Evolutionary Conservation." *Nucleic Acids Research* 49 (W1): W297–303.

Feurtey, Alice, and Eva H. Stukenbrock. 2018. "Interspecific Gene Exchange as a Driver of Adaptive Evolution in Fungi." *Annual Review of Microbiology* 72 (September): 377–98.

Forslund, Kristoffer, Cecile Pereira, Salvador Capella-Gutierrez, Alan Sousa da Silva, Adrian Altenhoff, Jaime Huerta-Cepas, Matthieu Muffato, et al. 2018. "Gearing up to Handle the Mosaic Nature of Life in the Quest for Orthologs." *Bioinformatics* 34 (2): 323–29.

Gabaldón, Toni. 2020. "Patterns and Impacts of Nonvertical Evolution in Eukaryotes: A Paradigm Shift." *Annals of the New York Academy of Sciences* 1476 (1): 78–92.

Gao, Shan, Scott E. Gold, Jennifer H. Wisecaver, Yong Zhang, Li Guo, Li-Jun Ma, Antonis Rokas, and Anthony E. Glenn. 2019. "Genome-Wide Analysis of Fusarium Verticillioides Reveals Inter-Kingdom Contribution of Horizontal Gene Transfer to the Expansion of Metabolism." *Fungal Genetics and Biology: FG & B* 128 (July): 60–73.

Grishin, N. V. 1995. "Estimation of the Number of Amino Acid Substitutions per Site When the Substitution Rate Varies among Sites." *Journal of Molecular Evolution* 41 (5): 675–79.

Grüll, M. P., M. E. Mulligan, and A. S. Lang. 2018. "Small Extracellular Particles with Big

Potential for Horizontal Gene Transfer: Membrane Vesicles and Gene Transfer Agents." *FEMS Microbiology Letters*. https://doi.org/10.1093/femsle/fny192.

Hansberg, Wilhelm, Rodolfo Salas-Lizana, and Laura Domínguez. 2012. "Fungal Catalases: Function, Phylogenetic Origin and Structure." *Archives of Biochemistry and Biophysics* 525 (2): 170–80.

Harris, Charles R., K. Jarrod Millman, Stéfan J. van der Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, et al. 2020. "Array Programming with NumPy." *Nature* 585 (7825): 357–62.

Horton, Paul, Keun-Joon Park, Takeshi Obayashi, Naoya Fujita, Hajime Harada, C. J. Adams-Collier, and Kenta Nakai. 2007. "WoLF PSORT: Protein Localization Predictor." *Nucleic Acids Research* 35 (Web Server issue): W585–87.

Huerta-Cepas, Jaime, François Serra, and Peer Bork. 2016. "ETE 3: Reconstruction, Analysis, and Visualization of Phylogenomic Data." *Molecular Biology and Evolution* 33 (6): 1635–38.

Ianiri, Giuseppe, Marco A. Coelho, Fiorella Ruchti, Florian Sparber, Timothy J. McMahon, Ci Fu, Madison Bolejack, et al. 2020. "HGT in the Human and Skin commensalMalassezia: A Bacterially Derived Flavohemoglobin Is Required for NO Resistance and Host Interaction." *Proceedings of the National Academy of Sciences*. https://doi.org/10.1073/pnas.2003473117.

Irwin, Nicholas A. T., Alexandros A. Pittis, Thomas A. Richards, and Patrick J. Keeling. 2022. "Systematic Evaluation of Horizontal Gene Transfer between Eukaryotes and Viruses." *Nature Microbiology* 7 (2): 327–36.

Jaramillo, Vinicio D. Armijos, Vinicio D. Armijos Jaramillo, Serenella A. Sukno, and Michael R. Thon. 2015. "Identification of Horizontally Transferred Genes in the Genus Colletotrichum Reveals a Steady Tempo of Bacterial to Fungal Gene Transfer." *BMC Genomics*. https://doi.org/10.1186/1471-2164-16-2.

Jarnot, Patryk, Joanna Ziemska-Legiecka, Marcin Grynberg, and Aleksandra Gruca. 2022. "Insights from Analyses of Low Complexity Regions with Canonical Methods for Protein Sequence Comparison." *Briefings in Bioinformatics* 23 (5). https://doi.org/10.1093/bib/bbac299.

Keeling, Patrick J., and Jeffrey D. Palmer. 2008. "Horizontal Gene Transfer in Eukaryotic Evolution." *Nature Reviews. Genetics* 9 (8): 605–18.

Kellner, M., A. Burmester, A. Wöstemeyer, and J. Wöstemeyer. 1993. "Transfer of Genetic Information from the Mycoparasite Parasitella Parasitica to Its Host Absidia Glauca." *Current Genetics* 23 (4): 334–37.

Koski, Liisa B., and G. Brian Golding. 2001. "The Closest BLAST Hit Is Often Not the Nearest Neighbor." *Journal of Molecular Evolution*. https://doi.org/10.1007/s002390010184.

Krassowski, Tadeusz, Aisling Y. Coughlan, Xing-Xing Shen, Xiaofan Zhou, Jacek Kominek, Dana A. Opulente, Robert Riley, et al. 2018. "Evolutionary Instability of CUG-Leu in the Genetic Code of Budding Yeasts." *Nature Communications* 9 (1): 1887.

Lage, Jean-Luc Da, Jean-Luc Da Lage, Manfred Binder, Aurélie Hua-Van, Štefan Janeček, and Didier Casane. 2013. "Gene Make-up: Rapid and Massive Intron Gains after Horizontal Transfer of a Bacterial α-Amylase Gene to Basidiomycetes." *BMC Evolutionary Biology*. https://doi.org/10.1186/1471-2148-13-40.

Lamarca, Alessandra P., Beatriz Mello, and Carlos G. Schrago. 2022. "The Performance of Outgroup-Free Rooting under Evolutionary Radiations." *Molecular Phylogenetics and Evolution* 169 (April): 107434.

Libeskind-Hadas, Ran, Yi-Chieh Wu, Mukul S. Bansal, and Manolis Kellis. 2014. "Pareto-Optimal Phylogenetic Tree Reconciliation." *Bioinformatics* 30 (12): i87–95.

Li, Meng, Jinjie Zhao, Nianwu Tang, Hang Sun, and Jinling Huang. 2018. "Horizontal Gene

Transfer From Bacteria and Plants to the Arbuscular Mycorrhizal Fungus Rhizophagus Irregularis." *Frontiers in Plant Science*. https://doi.org/10.3389/fpls.2018.00701.

Linard, Benjamin, Ingo Ebersberger, Shawn E. McGlynn, Natasha Glover, Tomohiro Mochizuki, Mateus Patricio, Odile Lecompte, et al. 2021. "Ten Years of Collaborative Progress in the Quest for Orthologs." *Molecular Biology and Evolution* 38 (8): 3033–45.

Ma, Li-Jun, H. Charlotte van der Does, Katherine A. Borkovich, Jeffrey J. Coleman, Marie-Josée Daboussi, Antonio Di Pietro, Marie Dufresne, et al. 2010. "Comparative Genomics Reveals Mobile Pathogenicity Chromosomes in Fusarium." *Nature*. https://doi.org/10.1038/nature08850.

Marcet-Houben, Marina, and Toni Gabaldón. 2010. "Acquisition of Prokaryotic Genes by Fungal Genomes." *Trends in Genetics: TIG* 26 (1): 5–8.

McDaniel, Lauren D., Elizabeth Young, Jennifer Delaney, Fabian Ruhnau, Kim B. Ritchie, and John H. Paul. 2010. "High Frequency of Horizontal Gene Transfer in the Oceans." *Science* 330 (6000): 50.

Mehrabi, Rahim, Ali H. Bahkali, Kamel A. Abd-Elsalam, Mohamed Moslem, Sarrah Ben M'Barek, Amir Mirzadi Gohari, Mansoor Karimi Jashni, Ioannis Stergiopoulos, Gert H. J. Kema, and Pierre J. G. M. de Wit. 2011. "Horizontal Gene and Chromosome Transfer in Plant Pathogenic Fungi Affecting Host Range." *FEMS Microbiology Reviews*. https://doi.org/10.1111/j.1574-6976.2010.00263.x.

Mier, Pablo, Lisanna Paladin, Stella Tamana, Sophia Petrosian, Borbála Hajdu-Soltész, Annika Urbanek, Aleksandra Gruca, et al. 2020. "Disentangling the Complexity of Low Complexity Proteins." *Briefings in Bioinformatics* 21 (2): 458–72.

Minh, Bui Quang, Heiko A. Schmidt, Olga Chernomor, Dominik Schrempf, Michael D. Woodhams, Arndt von Haeseler, and Robert Lanfear. 2020. "IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era." *Molecular Biology and Evolution* 37 (5): 1530–34.

Mondo, Stephen J., Richard O. Dannebaum, Rita C. Kuo, Katherine B. Louie, Adam J. Bewick, Kurt LaButti, Sajeet Haridas, et al. 2017. "Widespread Adenine N6-Methylation of Active Genes in Fungi." *Nature Genetics* 49 (6): 964–68.

Murphy, Chelsea L., Noha H. Youssef, Radwa A. Hanafy, M. B. Couger, Jason E. Stajich, Yan Wang, Kristina Baker, et al. 2019. "Horizontal Gene Transfer as an Indispensable Driver for Evolution of Neocallimastigomycota into a Distinct Gut-Dwelling Fungal Lineage." *Applied and Environmental Microbiology* 85 (15). https://doi.org/10.1128/AEM.00988-19.

Naito, Mizue, Joseph B. Morton, and Teresa E. Pawlowska. 2015. "Minimal Genomes of Mycoplasma-Related Endobacteria Are Plastic and Contain Host-Derived Genes for Sustained Life within Glomeromycota." *Proceedings of the National Academy of Sciences of the United States of America* 112 (25): 7791–96.

Nakamura, Tsukasa, Kazunori D. Yamada, Kentaro Tomii, and Kazutaka Katoh. 2018. "Parallelization of MAFFT for Large-Scale Multiple Sequence Alignments." *Bioinformatics*. https://doi.org/10.1093/bioinformatics/bty121.

Naranjo-Ortiz, Miguel A., and Toni Gabaldón. 2020. "Fungal Evolution: Cellular, Genomic and Metabolic Complexity." *Biological Reviews of the Cambridge Philosophical Society* 95 (5): 1198–1232.

NCBI Resource Coordinators. 2017. "Database Resources of the National Center for Biotechnology Information." *Nucleic Acids Research* 45 (D1): D12–17.

Nee, S., R. M. May, and P. H. Harvey. 1994. "The Reconstructed Evolutionary Process." *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 344 (1309): 305–11.

Nguyen, Kim B., Anju Sreelatha, Eric S. Durrant, Javier Lopez-Garrido, Anna Muszewska, Małgorzata Dudkiewicz, Marcin Grynberg, et al. 2016. "Phosphorylation of Spore Coat

Proteins by a Family of Atypical Protein Kinases." *Proceedings of the National Academy of Sciences of the United States of America*.

Page, R. D., and M. A. Charleston. 1997. "From Gene to Organismal Phylogeny: Reconciled Trees and the Gene Tree/species Tree Problem." *Molecular Phylogenetics and Evolution* 7 (2): 231–40.

Pawlowska, Teresa E., Maria L. Gaspar, Olga A. Lastovetsky, Stephen J. Mondo, Imperio Real-Ramirez, Evaniya Shakya, and Paola Bonfante. 2018. "Biology of Fungi and Their Bacterial Endosymbionts." *Annual Review of Phytopathology* 56 (August): 289–309.

Persson, Emma, Mateusz Kaduk, Sofia K. Forslund, and Erik L. L. Sonnhammer. 2019. "Domainoid: Domain-Oriented Orthology Inference." *BMC Bioinformatics* 20 (1): 523.

Põlme, Sergei, Kessy Abarenkov, R. Henrik Nilsson, Björn D. Lindahl, Karina Engelbrecht Clemmensen, Havard Kauserud, Nhu Nguyen, et al. 2020. "FungalTraits: A User-Friendly Traits Database of Fungi and Fungus-like Stramenopiles." *Fungal Diversity*. https://doi.org/10.1007/s13225-020-00466-2.

Poptsova, Maria S., and J. Peter Gogarten. 2010. "Using Comparative Genome Analysis to Identify Problems in Annotated Microbial Genomes." *Microbiology* 156 (Pt 7): 1909–17.

Priest, Shelby J., Vikas Yadav, and Joseph Heitman. 2020. "Advances in Understanding the Evolution of Fungal Genome Architecture." *F1000Research* 9 (July). https://doi.org/10.12688/f1000research.25424.1.

Richards, Thomas A., Guy Leonard, Darren M. Soanes, and Nicholas J. Talbot. 2011. "Gene Transfer into the Fungi." *Fungal Biology Reviews*. https://doi.org/10.1016/j.fbr.2011.04.003.

Richards, Thomas A., Darren M. Soanes, Peter G. Foster, Guy Leonard, Christopher R. Thornton, and Nicholas J. Talbot. 2009. "Phylogenomic Analysis Demonstrates a Pattern of Rare and Ancient Horizontal Gene Transfer between Plants and Fungi." *The Plant Cell* 21 (7): 1897–1911.

Richards, Thomas A., and Nicholas J. Talbot. 2013. "Horizontal Gene Transfer in Osmotrophs: Playing with Public Goods." *Nature Reviews Microbiology*. https://doi.org/10.1038/nrmicro3108.

Robinson, Aaron J., Geoffrey L. House, Demosthenes P. Morales, Julia M. Kelliher, La Verne Gallegos-Graves, Erick S. LeBrun, Karen W. Davenport, et al. 2021. "Widespread Bacterial Diversity within the Bacteriome of Fungi." *Communications Biology* 4 (1): 1168.

Rodríguez-Ezpeleta, Naiara, Henner Brinkmann, Béatrice Roure, Nicolas Lartillot, B. Franz Lang, and Hervé Philippe. 2007. "Detecting and Overcoming Systematic Errors in Genome-Scale Phylogenies." *Systematic Biology* 56 (3): 389–99.

Rokas, Antonis, and Sean B. Carroll. 2008. "Frequent and Widespread Parallel Evolution of Protein Sequences." *Molecular Biology and Evolution* 25 (9): 1943–53.

Schönknecht, Gerald, Andreas P. M. Weber, and Martin J. Lercher. 2014. "Horizontal Gene Acquisitions by Eukaryotes as Drivers of Adaptive Evolution." *BioEssays: News and Reviews in Molecular, Cellular and Developmental Biology* 36 (1): 9–20.

Shi-Kunne, Xiaoqian, Mathijs van Kooten, Jasper R. L. Depotter, Bart P. H. J. Thomma, and Michael F. Seidl. 2019. "The Genome of the Fungal Pathogen Verticillium Dahliae Reveals Extensive Bacterial to Fungal Gene Transfer." *Genome Biology and Evolution* 11 (3): 855–68.

Sjölander, Kimmen, Ruchira S. Datta, Yaoqing Shen, and Grant M. Shoffner. 2011. "Ortholog Identification in the Presence of Domain Architecture Rearrangement." *Briefings in Bioinformatics* 12 (5): 413–22.

Soliman, Sameh S. M., Clara Baldin, Yiyou Gu, Shakti Singh, Teclegiorgis Gebremariam, Marc Swidergall, Abdullah Alqarihi, et al. 2021. "Mucoricin Is a Ricin-like Toxin That Is Critical for the Pathogenesis of Mucormycosis." *Nature Microbiology*. https://doi.org/10.1038/s41564-020-00837-0.

Steel, M., and A. McKenzie. 2001. "Properties of Phylogenetic Trees Generated by Yule-Type Speciation Models." *Mathematical Biosciences* 170 (1): 91–112.

Tabima, Javier F., Ian A. Trautman, Ying Chang, Yan Wang, Stephen Mondo, Alan Kuo, Asaf Salamov, Igor V. Grigoriev, Jason E. Stajich, and Joseph W. Spatafora. 2020. "Phylogenomic Analyses of Non-Dikarya Fungi Supports Horizontal Gene Transfer Driving Diversification of Secondary Metabolism in the Amphibian Gastrointestinal Symbiont,." *G3* 10 (9): 3417–33.

Tan, Ge, Matthieu Muffato, Christian Ledergerber, Javier Herrero, Nick Goldman, Manuel Gil, and Christophe Dessimoz. 2015. "Current Methods for Automated Filtering of Multiple Sequence Alignments Frequently Worsen Single-Gene Phylogenetic Inference." *Systematic Biology* 64 (5): 778–91.

Torres-Martínez, Santiago, and Rosa M. Ruiz-Vázquez. 2017. "The RNAi Universe in Fungi: A Varied Landscape of Small RNAs and Biological Functions." *Annual Review of Microbiology* 71 (September): 371–91.

Uehling, Jessie, Aurélie Deveau, and Mathieu Paoletti. 2017. "Do Fungi Have an Innate Immune Response? An NLR-Based Comparison to Plant and Animal Immune Systems." *PLoS Pathogens* 13 (10): e1006578.

Uehling, J., A. Gryganskyi, K. Hameed, T. Tschaplinski, P. K. Misztal, S. Wu, A. Desirò, et al. 2017. "Comparative Genomics of Mortierella Elongata and Its Bacterial Endosymbiont Mycoavidus Cysteinexigens." *Environmental Microbiology* 19 (8): 2964–83.

Van Etten, Julia, and Debashish Bhattacharya. 2020. "Horizontal Gene Transfer in Eukaryotes: Not If, but How Much?" *Trends in Genetics*. https://doi.org/10.1016/j.tig.2020.08.006.

Venice, Francesco, Alessandro Desirò, Gladstone Silva, Alessandra Salvioli, and Paola Bonfante. 2020. "The Mosaic Architecture of NRPS-PKS in the Arbuscular Mycorrhizal Fungus Gigaspora Margarita Shows a Domain With Bacterial Signature." *Frontiers in Microbiology*. https://doi.org/10.3389/fmicb.2020.581313.

Virtanen, Pauli, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, et al. 2020. "SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python." *Nature Methods*. https://doi.org/10.1038/s41592-019-0686-2.

Voigt, Kerstin, Timothy Y. James, Paul M. Kirk, André L. C. M. de A. Santiago, Bruce Waldman, Gareth W. Griffith, Minjie Fu, et al. 2021. "Early-Diverging Fungal Phyla: Taxonomy, Species Concept, Ecology, Distribution, Anthropogenic Impact, and Novel Phylogenetic Proposals." *Fungal Diversity*, September, 1–40.

Vossenberg, Bart T. L. H. van de, Sven Warris, Hai D. T. Nguyen, Marga P. E. van Gent-Pelzer, David L. Joly, Henri C. van de Geest, Peter J. M. Bonants, Donna S. Smith, C. André Lévesque, and Theo A. J. van der Lee. 2019. "Comparative Genomics of Chytrid Fungi Reveal Insights into the Obligate Biotrophic and Pathogenic Lifestyle of Synchytrium Endobioticum." *Scientific Reports* 9 (1): 8672.

Wade, Taylor, L. Thiberio Rangel, Soumya Kundu, Gregory P. Fournier, and Mukul S. Bansal. 2020. "Assessing the Accuracy of Phylogenetic Rooting Methods on Prokaryotic Gene Families." *PLoS One* 15 (5): e0232950.

Wisecaver, Jennifer H., and Antonis Rokas. 2015. "Fungal Metabolic Gene Clustersâ€"caravans Traveling across Genomes and Environments." *Frontiers in Microbiology*. https://doi.org/10.3389/fmicb.2015.00161.

Wisecaver, Jennifer H., Jason C. Slot, and Antonis Rokas. 2014. "The Evolution of Fungal Metabolic Pathways." *PLoS Genetics* 10 (12): e1004816.

Wootton, J. C., and S. Federhen. 1996. "Analysis of Compositionally Biased Regions in Sequence Databases." *Methods in Enzymology* 266: 554–71.

Wootton, John C., and Scott Federhen. 1993. "Statistics of Local Complexity in Amino Acid

Sequences and Sequence Databases." *Computers & Chemistry*. https://doi.org/10.1016/0097-8485(93)85006-x.

Wright, Peter E., and H. Jane Dyson. 2015. "Intrinsically Disordered Proteins in Cellular Signalling and Regulation." *Nature Reviews. Molecular Cell Biology* 16 (1): 18–29.

Wurlitzer, Jacob M., Aleksa Stanišić, Ina Wasmuth, Sandra Jungmann, Dagmar Fischer, Hajo Kries, and Markus Gressler. 2021. "Bacterial-Like Nonribosomal Peptide Synthetases Produce Cyclopeptides in the Zygomycetous Fungus Mortierella Alpina." *Applied and Environmental Microbiology*. https://doi.org/10.1128/aem.02051-20.

Yanai, Itai, Yuri I. Wolf, and Eugene V. Koonin. 2002. "Evolution of Gene Fusions: Horizontal Transfer versus Independent Events." *Genome Biology* 3 (5): research0024.

Zhang, Qiangqiang, Xiaoxuan Chen, Chuan Xu, Hong Zhao, Xing Zhang, Guohong Zeng, Ying Qian, et al. 2019. "Horizontal Gene Transfer Allowed the Emergence of Broad Host Range Entomopathogens." *Proceedings of the National Academy of Sciences of the United States of America* 116 (16): 7982–89.

Zhao, Hong, Chuan Xu, Hsiao-Ling Lu, Xiaoxuan Chen, Raymond J. St. Leger, and Weiguo Fang. 2014. "Host-to-Pathogen Gene Transfer Facilitated Infection of Insects by a Pathogenic Fungus." *PLoS Pathogens*. https://doi.org/10.1371/journal.ppat.1004009.