Trial-by-trial fluctuations in amygdala activity track motivational enhancement of

desirable sensory evidence during perceptual decision-making

Abbreviated Title: Amygdala contributions to motivated seeing

Ren Paterson, Yizhou Lyu, Yuan Chang Leong*

Department of Psychology, University of Chicago, Chicago, IL 60637

* Correspondence should be addressed to Yuan Chang Leong at ycleong@uchicago.edu

Number of pages: 34

Number of figures: 4

Number of words for Abstract: 228; Introduction: 587; Discussion: 1421

The authors declare no competing financial interests.

Acknowledgments: We would like to thank members of the Motivation and Cognition

Neuroscience Lab for their feedback on this work, and Howard Nusbaum for helpful comments

on a previous version of this manuscript.

Abstract

People are biased towards seeing outcomes that they are motivated to see. For example, sports fans

of opposing teams often perceive the same ambiguous foul in favor of the team they support. Here,

we test the hypothesis that amygdala-dependent allocation of visual attention facilitates

motivational biases in perceptual decision-making. Human participants were rewarded for

correctly categorizing an ambiguous image into one of two categories while undergoing fMRI. On

each trial, we used a financial bonus to motivate participants to see one category over another. The

reward maximizing strategy was to perform the categorization task accurately, but participants

were biased towards categorizing the images as the category we motivated them to see. Heightened

amygdala activity preceded motivation consistent categorizations, and participants with higher

amygdala activation exhibited stronger motivational biases in their perceptual reports. Trial-by-

trial amygdala activity was associated with stronger enhancement of neural activity encoding

desirable percepts in sensory cortices, suggesting that amygdala-dependent effects on perceptual

decisions arose from biased sensory processing. Analyses using a drift diffusion model provide

converging evidence that trial-by-trial amygdala activity was associated with stronger motivational

biases in the accumulation of sensory evidence. Prior work examining biases in perceptual

decision-making have focused on the role of frontoparietal regions. Our work highlights an

important contribution of the amygdala. When people are motivated to see one outcome over

another, the amygdala biases perceptual decisions towards those outcomes.

Significance Statement

Forming accurate perceptions of the environment is essential for adaptive behavior. People

however are biased towards seeing what they want to see, giving rise to inaccurate perceptions and

erroneous decisions. Here, we combined behavior, modeling, and fMRI to show that the bias

towards seeing desirable percepts is related to trial-by-trial fluctuations in amygdala activity. In

particular, during moments with higher amygdala activity, sensory processing is biased in favor of

desirable percepts, such that participants are more likely to see what they want to see. These

findings highlight the role of the amygdala in biasing visual perception, and shed light on the

neural mechanisms underlying the influence of motivation and reward on how people decide what

they see.

Introduction

People are biased towards seeing percepts that they are motivated to see, a phenomenon known as 'wishful seeing' (Dunning and Balcetis, 2013). For example, sports fans of opposing teams watching the same game often perceive an ambiguous foul differently, with each group of fans judging the foul in favor of the team they support. When motivated to see one interpretation of reality over another, people become less objective and are more prone to perceptual errors (Voss et al., 2008; Bromberg-Martin and Sharot, 2020; Leong et al., 2021). Individuals who score higher on trait paranoia are also more likely to exhibit wishful seeing (Rossi-Goldthorpe et al., 2021), suggesting that extreme cases of wishful seeing might contribute to biased beliefs about the world that are the hallmark of many psychiatric disorders (Kube and Rozenkrantz, 2021). Why are people more likely to see what they are motivated to see?

To formulate a mechanistic understanding of wishful seeing, we can draw on the extensive neurophysiological and behavioral modeling work characterizing how perceptual decisions are determined from noisy sensory information (e.g., Heekeren et al., 2008; Shadlen and Kiani, 2013). In particular, perceptual decisions are thought to involve the temporal accumulation of sensory evidence from the environment. When the level of sensory evidence reaches a threshold, the individual commits to a decision. Recent work suggests that wishful seeing is driven by the selective enhancement of neural activity encoding desirable percepts (Leong et al., 2019). This biases sensory evidence accumulation in favor of percepts one is motivated to see, giving rise to motivationally biased perceptual decisions. While prior work has shown that motivation enhances the neural representation of desirable percepts in sensory cortices, how this enhancement occurs is not known. Consequently, the neural mechanisms that give rise to wishful seeing remain underspecified.

The amygdala has extensive connections to the visual cortex (Freese and Amaral, 2005),

and is well-positioned to mediate wishful seeing. Human neuroimaging and non-human primate

electrophysiology studies have found that the amygdala plays a central role in biasing attention

towards affectively salient aspects of the environment (Pessoa and Adolphs, 2010; Pourtois et al.,

2013; Peck and Salzman, 2014a; Mather et al., 2016). For example, fluctuations in amygdala

activity correlate with the enhanced neural representation of images that had acquired affective

salience due to having been previously paired with an electrical shock (Lim et al., 2009). We

hypothesize that wanting to see a percept ascribes affective significance to the associated sensory

features, and the amygdala is similarly involved in enhancing the neural representation of these

features. Past studies, however, have not examined the role of the amygdala in contexts where

participants are motivationally biased to see one percept over another. It is thus unclear if and how

the amygdala biases perception towards desirable percepts.

In the current work, we conducted a series of pre-registered analyses on a previously

published dataset (Leong et al., 2019) to test the hypothesis that the amygdala facilitates wishful

seeing by enhancing sensory evidence accumulation in favor of desirable percepts. We first

assessed if amygdala activity was associated with motivationally biased perceptual decisions.

Next, we tested if amygdala activity was associated with stronger neural representations of

percepts participants were motivated to see. Finally, we incorporated participants' trial-by-trial

amygdala activity, perceptual decisions and response times into a computational model to test the

hypothesis that amygdala activity was specifically associated with motivational biases in evidence

accumulation. Together, our work takes a convergent approach that combines behavioral, neural

and modeling measures to characterize the neural mechanisms underlying motivational biases in

perceptual decision-making.

Materials and Methods

Participants

Thirty-three participants provided informed consent and were compensated between $30-50

depending on their performance in the task. Data were discarded from 3 participants due to

excessive head motion (i.e., >3 mm in any direction) during >1 scanning sessions, yielding a final

sample of 30 participants (17 males, 13 females; age range, 18-43 years; mean age, 22.3 years).

These data have been previously reported in Leong et al. (2019).

Stimuli

Stimuli consisted of grayscale images created by combining a scene image and face image in

varying proportions. Face images were frontal photographs taken from the Chicago Face Database

(Ma et al., 2015), and consisted of half female and half male faces with neutral expressions. Scene

images consisted of half outdoor and half indoor scenes. For each participant, six stimulus sets

were created. Each set contained 40 images: $1 \times 100\%$ scene, $3 \times 65\%$ scene, $5 \times 60\%$ scene,

$7 \times 55\%$ scene, $8 \times 50\%$ scene, $7 \times 45\%$ scene, $5 \times 40\%$ scene, $3 \times 35\%$ scene and $1 \times 0\%$ scene.

Images with more than 50% scene were considered "more scene" while images with less than 50%

scene were considered "more face". Stimuli were presented to participants using MATLAB

(Mathworks) and the Psychophysics Toolbox (Brainard, 1997).

Experimental Task

Participants were told that they would be performing a visual categorization task with a teammate

("Cooperation" condition) and an opponent ("Competition" condition) (Fig. 1). At the beginning

of each trial, either the teammate or the opponent would make a bet on whether the upcoming

image would consist of more face or more scene (i.e. whether the face image or scene image was

of higher intensity; "Motivation Cue", presented for 4 seconds). Participants were then presented

with a face-scene composite image (see *Stimuli*). Participants would earn 40 cents if the

teammate's bet was correct (e.g., if the teammate bet "more face" and the image actually consists

of more face) and lose 40 cents if the teammate's bet was wrong. In contrast, participants would

lose 40 cents if the opponent's bet was correct and earn 40 cents if the opponent's bet was wrong.

As such, participants would be motivated to see the image as the category that the teammate had

bet on and the category that the opponent had bet against.

Participants were then asked to indicate whether the image consists of more face or more

scene and received a 10-cent reward if they correctly categorized the image. If participants did not

respond in 4 seconds, the trial would time out and participants would lose the opportunity to earn

the reward. The image remained on screen for the entire 4 seconds regardless of when participants

made their response. Participants then indicated how confident they were in their categorization

on a 1-5 rating scale. After a variable inter-trial-interval (ITI) of 2-4 seconds, participants

proceeded to the next trial. Participants performed 2 fMRI runs of the Cooperation condition and

2 fMRI runs of the Competition condition in an interleaved order that was counterbalanced across

participants. Each run was approximately 8 minutes long and consisted of 40 trials.

Importantly, the 40-cent bonus associated with the teammate or opponent's bet depended

solely on the objective category of the image (i.e. whether the image actually contained more face

or more scene) and was not affected by participants' responses. For example, if the teammate bet

that the upcoming image contains more face but the image objectively contained more scene,

participants would lose the 40 cents regardless of how they categorized the image. As such,

participants would earn the most amount of money from the experiment if they ignored the bets

and categorized the images accurately based on what they saw to earn the maximum reward from

correctly categorizing each image. Participants were told that both the teammate and the opponent

did not have *a priori* information about the images and were making blind bets. Participants were

also told that neither the teammate or opponent would be informed of their responses, and were

thus not pressured to conform to the bets. Unbeknownst to participants, the teammate's and

opponent's bets were pseudo-randomized to be correct on half of the trials.

The instructions were delivered verbally by the experimenter and also in written form

on screen at the beginning of the experiment. It was intuitive to participants that the bet

depended on the objective category of the images rather than their categorizations, and that

they should categorize each image accurately to earn maximum reward. In the post-experiment

survey, all participants responded that the task instructions were clear and that they had

understood the task correctly.

Figure 1. Experimental design. Each trial begins with a teammate (Cooperation condition) or an
opponent (Competition condition) making a bet on whether the upcoming image would consist of
more face or more scene. Participants earn 40 cents if the teammate's bet is correct, and lose 40
cents otherwise. Conversely, they lose 40 cents if the opponent's bet is correct and earn 40 cents
otherwise. Participants are then presented with a face-scene composite image, and have to
categorize whether the image has more face or more scene. Finally, participants rated how
confident they were in their categorization. Images have been replaced with cartoon images to
comply with *bioRxiv* policy.

Motivational Bias

In earlier work, we had found that motivational bias and response times did not differ between the

Competition and Cooperation conditions - participants were just as likely and quickly to categorize

the image as the category that the teammate had bet on as they were to categorize it as the category

the opponent had bet against (Leong et al., 2019). Therefore, in the current study, we collapsed

across the two conditions based on whether participants were motivated to see more face (i.e. when

the teammate bet on more face or when the opponent bet on more scene) or more scene (i.e. when

the teammate bet on more scene or when the opponent bet on more face).

We modeled participants' data using a generalized linear mixed effects model (GLMM)

using the glmer function from the lme4 package in R:

168   response ~ % scene + motivation + (motivation | subj)

where *response* denotes the perceptual report (face = 0, scene = 1), *motivation* is the category that

participants were motivated to see (face = 0, scene = 1), and *% scene* is the proportion of scene in

an image. The model included *motivation* as a random effect to account for random variation

between subjects. Importantly, the estimate of the random effect of a given participant provides a

measure of the effect of *motivation* on that participants' perceptual categorizations. For each

participant, we added the random effect to the corresponding fixed effect to obtain an individual

difference measure of motivational bias.

We used a robust regression model to examine the relationship between motivational bias

and amygdala activation (see fMRI analyses) using the rlm function from the MASS package in R

with default settings (M-estimation and Huber weighting). Robust regression has been shown to

be less sensitive to the influence of outliers than Pearson correlation and ordinary least squares

regression (Venables and Ripley, 2013). Statistical significance of regression coefficients were

assessed using a robust F test as implemented in the sfsmisc package in R.

MRI Data Acquisition and Preprocessing

A 3T General Electric scanner was used to collect the MRI data. Functional images were obtained

with T2*-weighted echo planar imaging (EPI) pulse sequences. Each volume comprised 46

transverse slices. Volumes were acquired in interleaved order, with the following imaging

parameters: repetition time, 2 s; echo time, 25 ms; flip angle, 77º; voxel size, 2.9 mm^3. A T1-

weighted pulse sequence was used to acquire anatomical images at the beginning of the session

with the following parameters: repetition time, 7.2 ms; echo time, 2.8 ms; flip angle, 12º; voxel

size, 1mm^3.

Preprocessing and analysis of EPI images were performed using FSL/FEAT v6.05 (FMRIB

software library, FMRIB, Oxford, UK) and included motion correction (rigid-body realignment of

functional volumes to the first volume), slice-timing correction, high-pass filtering of the data with

a 100 ms cut-off, and spatial smoothing using a Gaussian kernel with a full-width at half-maximum

of 4 mm. For multivoxel classification analyses, our classifier was trained and tested in each

participant's native space (see fMRI analyses). For all other analyses, we first registered the

functional volumes to participants' anatomical images (6 d.f.) and then registered the volumes to

a template brain in the Montreal Neurological Institute (MNI) space (affine transformation with

12 d.f.).

Regions of interest (ROI) definition

We defined a bilateral amygdala ROI as the voxels that are estimated to have greater than 0.5

probability as being part of the amygdala according to the Harvard-Oxford Subcortical Structural

Atlas. To measure sensory representations, we defined an occipito-temporal ROI by combining

the bilateral occipital and temporal lobe masks from the Harvard-Oxford Structural Atlas. This

mask includes the visual cortex as well as the ventral temporal areas important for object

recognition (Grill-Spector, 2003). We defined additional ROIs from an atlas derived from applying

independent component analysis to resting-state data (Shirer et al., 2012). Specifically, the anterior

insula and dorsal anterior cingulate cortex (dACC) ROIs were obtained from the set of regions

labeled the "anterior salience network", while the dorsolateral prefrontal cortex (DLPFC) and

inferior parietal lobule (IPL) ROIs were obtained from the set of regions labeled the "executive

control network".

fMRI Analyses

We ran a general linear model (GLM) analysis contrasting neural activity on motivation consistent

trials (i.e. trials where participants categorized the image as the desirable category) against that on

motivation inconsistent trials (i.e. trials where participants categorized the image as the less

desirable category). The two trial types were modeled as separate regressors, each created by

convolving a stimulus function, consisting of an impulse response at the time of stimulus onsets,

with a Gamma hemodynamic response function (HRF). For each participant, a separate model was

estimated for each run. Response times (convolved with HRF) and extended motion parameters

(i.e. standard motion parameters, their temporal derivatives, and the squares of both the standard

motion parameters and the temporal derivatives; not convolved with HRF) were included as

covariates of no interest. The results were then combined across the four runs in a fixed effects

analysis to obtain a statistical parametric map for each participant. We then used FSL's *featquery*

function to extract the average *z*-statistic of the contrast in the amygdala ROI for each participant,

and ran a one-sample t-test against zero to assess if amygdala activity was higher on motivation

consistent trials than on motivation inconsistent trials.

To examine if individual differences in amygdala activation tracked motivational biases in

categorizations, we ran a second set of GLM analyses to estimate the average magnitude of

amygdala activation for each participant. The modeling procedure was the same as before, except

that motivation consistent and motivation inconsistent trials were not modeled separately. Instead,

a single regressor was created by convolving a stimulus function, consisting of an impulse response

at the time of stimulus onset of all trials, with a HRF. For each participant, we used FSL's *featquery*

function to extract the average beta weights in the amygdala ROI as an estimate of the extent to

which that participant's amygdala was activated by the task. We then used a robust regression

analysis to assess if individual differences in amygdala activation were associated with the

magnitude of motivational bias in categorizations (see *Motivational Bias*).

To examine the time course of amygdala activation and its relationship to motivation

consistent responses, we extracted the time course in the amygdala ROI from the preprocessed

BOLD data of each participant. The time courses were z-scored across each run, shifted by 4

seconds to correct for haemodynamic lag, then time locked to the stimulus onset of each trial. For

each time point from -8 seconds prior to stimulus onset to 8 seconds post stimulus onset, we ran a

generalized linear mixed effects model examining the effect of amygdala activity on whether

participants made a motivation consistent categorization:

243   MotCon ~ amygdala + (amygdala | subj)

where *MotCon* denotes whether participants made a motivation consistent or inconsistent response

(motivation consistent = 1, motivation inconsistent = 0) and amygdala denotes amygdala activity

at a given time point.

At the end of the experimental task, participants performed two runs of a localizer task

where they were presented with unambiguous face and scene images one at a time (5 blocks of 15

images of each category; each image presented for 2 seconds with a 2-second interval). Participants

had to indicate whether each face was male or female, and whether each scene was indoors or

outdoors. We trained an L1-regularized logistic regression classifier (C = 1) to classify whether

the participant was seeing a face or scene based on the patterns of activity in the occipito-temporal

ROI. We then applied the trained classifier to the single-trial activation patterns in the experimental

task. Specifically, all trials in a run were simultaneously estimated in a single GLM, with each trial

modeled using a separate regressor consisting of an impulse response at the time of stimulus onset

convolved with a HRF (i.e. a "Least-Squares All" model; Mumford et al., 2014). The beta weights

of each trial were then used as inputs for the trained classifier. For each trial, the classifier outputs

the probability that the participant was viewing a scene vs. a face based on the beta weights in the

occipito-temporal ROI, thus providing us with a measure of the relative strength of scene vs. face

selective neural activity for that trial.

We estimated anticipatory amygdala activity on each trial using a Least-Squares All GLM

that modeled each trial as a separate regressor created by convolving a stimulus function,

consisting of an impulse response at the time of cue onset, with a HRF. The resulting beta weights

obtained for each regressor were then averaged within the amygdala ROI as the estimate of

anticipatory amygdala activity for that trial. A linear mixed effects model was used to test the

relationship between trial-by-trial anticipatory amygdala activity and sensory representations:

scene vs. face activity ~ % scene + motivation + amygdala + motivation*amygdala +

268   (motivation + amygdala + motivation*amygdala | subj)

where *scene vs. face activity* is the relative strength of scene vs. face-selective neural activity as

indexed by the classifier output, *% scene* is the amount of scene in an image, *motivation* is the

category participants were motivated to see (face = 0, scene = 1), and *amygdala* is amygdala

activation at the time of the cue. The coefficient for the interaction term, *motivation*amygdala*,

would thus capture the extent to which motivational effects on sensory-related activity depended

on anticipatory amygdala activation. To unpack the interaction effect, we divided each

participant's data into trials with high and low amygdala activity based on a median split, and

examined the effect of motivation on *scene vs. face activity* for the two trial types separately:

scene vs. face activity ~ % scene + motivation + (motivation | subj)

In a series of exploratory analyses, we also examined if motivational effects on scene and face-

selective activity was moderated by activity in the anterior insula, dorsal anterior cingulate cortex

(dACC), dorsolateral prefrontal cortex (DLPFC), and inferior parietal lobule (IPL). All models

were estimated using the lmer function in the lme4 package in R, with P values computed from t-

tests with Satterthwaite approximation for the degrees of freedom as implemented in the lmerTest

package.

Drift Diffusion Model

We fit a drift diffusion model to participants' data using the HDDM toolbox with default priors

(Wiecki et al., 2013). The model assumes that participants' decisions are determined by the

stochastic accumulation of sensory evidence towards one of two decision thresholds. We estimated

model parameters that determined the starting point of the decision process (parameter *z*), the rate

of evidence accumulation (i.e. *drift rate*, parameter *v*), the distance between the two decision

thresholds (parameter *a*) and time not related to the decision process *(e.g.,* time for stimulus

encoding or motor response; *non-decision time*, parameter *t*). We modeled the "scene" decision

threshold as the top boundary (i.e. scene threshold = *a*) and the "face" threshold as the bottom

boundary (i.e. face threshold = *0*). HDDM implements hierarchical Bayesian estimation, which

assumes that parameters for individual participants were randomly drawn from a group-level

distribution. Individual participant parameters and group-level parameters were jointly estimated

using Markov Chain Monte Carlo sampling (20,000 samples; burn-in=2,000 samples; thinning=2).

To account for outliers generated by a process other than that assumed by the model (e.g.,

accidental button press, lapses in attention), we estimated a mixture model where 5% of trials were

assumed to be distributed according to a uniform distribution. We chose not to include inter-trial

variability parameters as simulation studies have shown that they can compromise the reliability

and accuracy of the estimates of individual parameters (Boehm et al., 2018). A model that included

parameters for inter-trial variability in the starting point, drift rate and non-decision time yielded

qualitatively similar results (see *Results*).

The drift rate was allowed to vary trial-by-trial as a function of the stimulus, the category

participants were motivated to see, and amygdala activity:

306
307    $$v = \beta_{v1}\, \%scene \; + \; \beta_{v2}\, motivation + \beta_{v3}\, amygdala \; + \; \beta_{v4}\, amygdala * motivation + \beta_{v0}$$

308
where *% scene* refers to the percentage scene in an image, *motivation* refers to the category

participants were motivated to see (face = -1; scene = 1), and *amygdala* refers to amygdala

activation at cue onset estimated from a GLM (see fMRI analysis). The coefficient for the

amygdala*motivation interaction reflects the extent to which the effect of motivation on the drift

rate depended on amygdala activation. Specifically, a positive value of β_{v4} would indicate that the

drift bias towards the motivation consistent category was stronger when amygdala activation was

higher. To assess if β_{v4} was positive, we extracted the posterior distribution of β_{v4} estimated from

participants' data and computed the proportion of the distribution that was greater than 0. This

proportion denotes the probability the amygdala activation moderated motivational effects on the

drift rate.

Within the same model, the starting point was also allowed to vary trial-by-trial as a

function of the motivation consistent category and the amygdala activation at cue onset:

$$z = \frac{1}{1 + exp - (\beta_{z1} motivation\ +\ \beta_{z2} amygdala\ +\ \beta_{z3} amygdala * motivation\ +\ \beta_{z4})}$$

Here, the coefficient for the *amygdala*motivation* interaction reflects the extent to which the effect

of motivation on the starting point depended on amygdala activation. Specifically, a positive value

of β_{z3} would indicate that the starting point shifts more towards the motivation consistent category

when amygdala activation was higher. To assess if β_{z3} was positive, we extracted the posterior

distribution of β_{z3} estimated from participants' data and computed the proportion of the

distribution that was greater than 0. This proportion denotes the probability the amygdala

activation moderated motivational effects on the starting point.

Model convergence was assessed using the Gelman-Rubin $\hat{R}$ statistic. Model convergence

metrics, posterior means and 95% credible intervals of model parameters are reported in Extended

Table 4.1.

Preregistration

We pre-registered how we defined the amygdala and occipitotemporal ROIs. We had also pre-

registered the following hypotheses: (1) participants with stronger amygdala responses will exhibit

greater motivational biases in their perceptual categorizations, (2) participants would be more

biased in their categorizations on trials with high amygdala activity, (3) there would be stronger

motivational enhancement of desirable sensory representations on trials with high amygdala

activity, and (4) there was be stronger motivational biases in evidence accumulation, as estimated

by the DDM, on trials with high amygdala activity. The pre-registration can be viewed here:

https://osf.io/2vbcn/?view_only=030ad17d15724af2bb5f6f02ca5b34e7

Results

Thirty participants were scanned using fMRI while they were presented with visually ambiguous

images consisting of a face image and a scene image superimposed with one another. Participants

had to categorize whether they saw more face or more scene in each image, and received a

monetary reward for each correct categorization. At the beginning of each trial, we motivated

participants to see one category over the other with a cue indicating that they would win extra

money if the image was of one category (the "desirable" percept) and lose some of their earnings

if the image was of the other category (the "less desirable" percept) (see *Materials and Methods*).

The bonus or loss depended on the objective category of the image (i.e. whether the image actually

contained more face or more scene) and not participants' categorizations. As such, participants

would earn the most money from the task if they categorized the images accurately and were not

influenced by what they were motivated to see. In earlier work, we had shown that participants

were nevertheless more likely to categorize the images as the desirable percept, and that this bias

was associated with enhanced neural activity selective to the desirable percept in the ventral visual

stream (Leong et al., 2019).

Amygdala activity was associated with motivational bias in perceptual decisions

We divided trials into trials where participants categorized the image as the desirable percept

(motivation consistent trials) and trials where they categorized the image as the less desirable

percept (motivation inconsistent trials). We then ran a GLM that contrasted neural activity on

motivation consistent trials against that on motivation inconsistent trials. In a bilateral amygdala

ROI, the average z-statistic of the contrast was significantly above zero ($t(29) = 2.96, P = 0.006;$

Fig. 2A, 2B), indicating that amygdala activity was higher when participants made motivation

consistent categorizations than when they made motivation inconsistent categorizations.

To examine the time course of amygdala activation, we extracted the amygdala activity

time course of each trial and used generalized linear mixed effects models to assess the relationship

between each time point and participants' categorizations. The time courses were first shifted by

4 seconds to account for hemodynamic lag, and then time-locked to image onset. Amygdala

activity was marginally associated with making a motivation consistent response at the time of the

motivational cue (i.e. when the teammate or opponent made their bet; $b = 0.059$, $SE = 0.035$, $z =$

1.71, $p = 0.087$). This relationship was significant 2 seconds later ($b = 0.10$, $SE = 0.037$, $z = 2.80$,

$p = 0.0051$), and remained significant until the image appeared on screen ($b = 0.112$, $SE = 0.036$,

$z = 3.13$, $p = 0.0018$; Fig. 2C). As the relationship between amygdala activity and motivational

consistent responses was present prior to image onset, the results suggest that the heightened

amygdala activity did not reflect activation in response to participants having seen the desirable

category. Instead, they suggest that the amygdala activation predisposes participants towards

categorizing the image as the desirable category.

Next, we assessed if individual differences in motivational bias were associated with

amygdala activation. We first fit a generalized linear mixed effects model to participants' data to

jointly estimate the effect of motivation and % scene in the image on perceptual categorizations.

For a given participant, the random effect of motivation reflects the extent to which that participant

was more likely to make motivation consistent categorizations. We added each participant's

random effect to the fixed effect of motivation on perceptual categorizations to obtain an individual

difference measure of motivational bias. For each participant, we also ran a GLM to estimate the

amygdala response to the task (see Methods). Using a robust regression analysis, we found that

participants who exhibited stronger motivational biases also had stronger amygdala activation

(Fig. 2C, $b = 4.3$, $SE = 0.86$, $F(1, 28) = 5.6$, $P = 0.025$; Fig. 2D).

In summary, motivation consistent categorizations were associated with higher amygdala

activity, this amygdala activation preceded image onset, and participants who exhibited stronger

motivational biases in their categorizations had a stronger amygdala response to the task. Together,

these results suggest that heightened amygdala activity biased participants towards making

motivationally consistent perceptual decisions.

Figure 2. Amygdala activity was associated with motivational bias in perceptual decisions.
A. Amygdala ROI defined using the Harvard-Oxford Subcortical Structural Atlas. **B.** Amygdala
activity was higher when participants made motivation consistent categorizations (MotCon) than
when they made motivation inconsistent categorizations (MotIncon). Data points denote average
z-statistic of MotCon > MotIncon contrast in the amygdala ROI for individual participants. Error
bars indicate s.e.m. **C.** Amygdala activation preceded image onset. For each timepoint, we ran a
generalized linear mixed effects model estimating the effect of amygdala activity on whether
participants would make a motivation consistent response. Plot shows estimates of regression
coefficients with standard errors. Time points have been corrected for haemodynamic lag by
shifting the BOLD data by 4s. †$P < 0.1$, **$P < 0.01$. **D.** Participants with stronger motivational
biases in their categorizations had on average higher amygdala responses.

Anticipatory amygdala activity was associated with stronger motivational enhancement of category-selective neural activity

The preceding analyses indicated amygdala activity was associated with motivational biases in perceptual decisions. Did amygdala activity also moderate motivational enhancement of sensory representations? To answer this question, we used multivariate pattern analysis to quantify face- and scene-selective activity in the occipitotemporal cortex. We first trained a logistic regression classifier on BOLD data from a localizer task to categorize whether participants were seeing a face or a scene based on the pattern of activity in a occipitotemporal cortex ROI (see Methods). We then applied the trained classifier to data from the experimental task to measure the level of face and scene-selective activity on each trial.

In previous work, we had found that motivation enhances sensory representations in the occipitotemporal cortex — for a given image, face-selective activity was higher when participants were motivated to see more face, while scene-selective activity was higher when participants were motivated to see more scene (Leong et al., 2019). Here, we test if this motivational enhancement was dependent on amygdala activity. Given our earlier analyses indicating that amygdala activation preceded image onset, we focused on amygdala activity at the time of the cue (i.e. when participants were informed which category would win them more money). As the amygdala is known to respond to faces (Rutishauser et al., 2015; Visser et al., 2021), extracting the amygdala response to the cue also allowed us to isolate amygdala activation that is not contaminated by the response to the face-scene composite images.

We modeled each trial as a separate regressor in a GLM. Each regressor was created by convolving an impulse response at the time of cue onset with the HRF. The resulting beta weights were then averaged within the amygdala ROI to obtain amygdala activation at the time of the cue

for each trial. We then used a linear mixed effects model to examine the relationship between

category-selective activity, motivation and amygdala activity. There was a significant interaction

between amygdala activity and motivation on category-selective activity in the occipitotemporal

cortex ($t(37) = 2.77$, $P < 0.009$, $b = 0.039$, $SE = 0.014$). To unpack this interaction, we divided

each participant's data into trials with high and low amygdala activity. Motivation enhanced

category-selective activity on trials with high amygdala activity ($t(30) = 2.22$, $P = 0.034$, $b = 0.034$,

$SE = 0.015$), but not on trials with with low amygdala activity ($t(33) = 0.49$, $P = 0.63$, $b = 0.0077$,

$SE = 0.016$; Fig. 3A), suggesting that motivational enhancement of sensory representations was

dependent on amygdala activity.

Prior work implicates a network of frontoparietal regions in biasing perceptual decisions,

including the dorsolateral prefrontal cortex (DLPFC), the dorsal anterior cingulate cortex (dACC),

and the intraparietal lobule (IPL) (Domenech and Dreher, 2010; Summerfield and Koechlin, 2010;

Mulder et al., 2012, 2014; White et al., 2012). In a series of exploratory analyses, we tested if these

regions were associated with the motivational enhancement of sensory representations. We also

considered the role of the anterior insula, which previous work had found to be involved in the

accumulation of sensory evidence during perceptual decision-making (Ho et al., 2009). Activity

in these regions did not moderate motivational effects on face and scene-selectively regions,

suggesting that these regions were not involved in the motivational enhancement of desirable

perceptual representations (DLPFC: $t(20) = -0.19$, $P = 0.852$, $b = -0.001$, $SE = 0.008$; dACC: $t(24)$

$= 0.89$, $P = 0.385$, $b = 0.011$, $SE = 0.012$; IPL: $t(39) = 0.62$, $P = 0.551$, $b = 0.005$, $SE = 0.009$;

anterior insula: $t(33) = 1.25$, $P = 0.221$, $b = 0.017$, $SE = 0.013$; Fig. 3B).

Figure 3. Amygdala activity was associated with stronger motivational enhancement of category-selective neural activity in occipitotemporal cortex. A. Classifier probability that the participant was seeing a scene rather than a face based on the pattern of BOLD activity in the occipito-temporal ROI, which we used as a proxy for category-selective activity. On trials with high amygdala activity, scene-selective activity was higher when participants were motivated to see more scene than when they were motivated to see more face. On trials with low amygdala activity, motivation had no effect on category-selective activity. Error bars indicate s.e.m. **B.** Regression coefficients of the motivation x activity interaction on classifier probability for the amygdala, DLPFC, dACC, IPL and anterior insula. The interaction was significant only in the amygdala. Error bars indicate 95% confidence intervals.

Amygdala activity was specifically associated with motivational biases in evidence

accumulation

The drift diffusion model (DDM) provides a formal framework to examine the decision processes

underlying perceptual decision-making (Forstmann et al., 2016). The model assumes that

perceptual decisions arise from the noisy accumulation of sensory evidence from the external

environment. When the level of evidence reaches a decision threshold, the corresponding decision

is made. The starting point and rate of evidence accumulation, as well as the decision thresholds

and time not related to decision processes (i.e. *non-decision time*, e.g., time needed to translate the

decision into a motor response), are determined by model parameters that can be estimated by

fitting the model to participants' data (Fig. 4A). In previous work, we had found that motivation

biases perceptual decisions by (1) shifting the starting point towards the decision threshold of the

desirable category, thereby reducing the amount of sensory evidence needed to make a motivation

consistent response, and (2) by biasing sensory evidence in favor of the desirable category (Leong

et al., 2019). To understand the amygdala's contribution to motivation biases, we assessed if either

or both of these biasing mechanisms were dependent on trial-by-trial fluctuations in amygdala

activity.

We took a linear regression approach to examine the effects of the amygdala on

motivational biases in the starting point and rate of evidence accumulation. Specifically, the

starting point (parameter z), was allowed to vary as a function of the motivation consistent

category, amygdala activation at the time of the cue, and the interaction between the two

(amygdala*motivation interaction; see Methods). The regression coefficient on the interaction

term reflects the extent to which motivational effects on the starting point depended on amygdala

activity. A positive coefficient would indicate that the starting point is more strongly biased

towards the decision threshold of the motivation consistent category on trials with higher amygdala

activity. When we fit the model to participants' behavioral data, the posterior distribution of the

regression coefficient of the interaction term was centered around zero ($P(z_{amygdala*motivation}>0)$ =

0.384, mean = -0.002, 95% credible interval = -0.020 to 0.018; Fig. 4B), indicating that

motivational effects on the starting point did not depend on amygdala activity.

In the same model, we allowed the rate of evidence accumulation (i.e. *"drift"* rate,

parameter *v*) to vary as a function of the percentage scene in the image, the motivation consistent

category, amygdala activation at the time of the cue, and the interaction between the motivation

consistent category and amygdala activation. Here, the regression coefficient on the interaction

term reflects the extent to which motivational effects on the drift rate depended on amygdala

activity. A positive coefficient would indicate a stronger bias in evidence accumulation in favor of

the motivation consistent category on trials with higher amygdala activity. When we fit the model

to participants' behavioral data, the posterior distribution of the regression coefficient of the

interaction term indicated a high probability that the coefficient was positive

($P(v_{amygdala*motivation}>0)$ = 0.978, mean = 0.079, 95% credible interval = 0.002 to 0.160; Fig. 4B).

These results suggest on trials with high amygdala activity, evidence accumulation is biased

towards the decision threshold of the motivation consistent category, resulting in an increased bias

towards making motivation consistent responses.

In exploratory analyses, we fit a model that included inter-trial variability parameters for

the drift rate, starting point and non-decision-time, which allows for random trial-by-trial

variability in model parameters. Consistent with our earlier results, we found that trial-by-trial

amygdala activity moderated motivational biases in the drift rate ($P(v_{amygdala*motivation}> 0)$ = 0.997,

mean = 0.138, 95% credible interval = 0.040 to 0.243) but not the starting point

($P(z_{amygdala*motivation} > 0) = 0.124$, mean = -0.015, 95% credible interval = -0.039 to 0.011),

suggesting that our results were not due to random trial-by-trial fluctuations in model parameters.

Figure 4. Modeling Results. A. Schematic diagram of the drift diffusion model (DDM). The
DDM assumes that perceptual decisions arise from the stochastic accumulation of sensory
evidence over time towards one of two decision thresholds. z: starting point, v: drift rate, a:
decision threshold, t_0: sensory delay, t_1: time needed for response execution, the sum of t_0 and t_1
constitutes non-decision time. Motivation can bias perceptual decisions by biasing both the starting
point and drift rate (red and blue lines). **B.** Posterior distribution of the regression coefficient of
the amygdala*motivation interaction on the starting point. The distribution is centered around 0,
indicating that motivational effects on the starting point did not depend on amygdala activity. **C.**
Posterior distribution of the regression coefficient of the amygdala*motivation interaction on the
drift rate. The distribution skews positive, indicating the motivational effects on the drift rate were
dependent on amygdala activity.

Parameter	Prior	Coefficient	Estimate	Gelman-Rubin $\hat{R}$
a	$a \sim Gamma(2, 0.3)$	a	2.264[2.166, 2.366]	1.000
t	$t \sim N(0.5, 0.15)$	t	0.650[0.609, 0.694]	1.000
z	$z = invlogit(N(0.03, 0.04))$	$z_{intercept}$	0.486[0.468, 0.505]	1.000
		$z_{amygdala}$	0.007[-0.011, 0.026]	1.001
		$z_{motivation}$	0.022[0.007, 0.037]	1.000
		$z_{amygdala*motivation}$	-0.002[-0.020, 0.018]	1.002
v	$v = N(0.06, 0.2)$	$v_{intercept}$	0.191[0.065, 0.314]	1.000
		$v_{stimulus}$	0.920[0.833, 1.010]	1.000
		$v_{amygdala}$	0.087[-0.003, 0.179]	1.000
		$v_{motivation}$	0.085[0.002, 0.167]	1.000
		$v_{amygdala*motivation}$	0.081[0.002, 0.160]	1.003

523
Extended Table 4.1. Priors, posterior estimates, and model convergence metrics of model
parameters. Estimates indicate the posterior means, with square brackets denoting 95% credible
intervals. Model convergence was assessed using the Gelman-Rubin $\hat{R}$ statistic. $\hat{R}$ of all parameters
were less than 1.1, suggesting that there were no issues with model convergence. a: decision
threshold, t: non-decision time, z: starting point; v: drift rate, Gamma: a gamma distribution
parameterized by mean and standard deviation, N: denotes a normal distribution parameterized by
mean and standard deviation, invlogit: inverse logit transform.

Discussion

Psychophysical studies have previously shown that stimulus features that have been repeatedly

paired with reward tend to "capture" attention and are preferentially processed by the visual system

(Anderson et al., 2011; Le Pelley et al., 2015). Relatedly, fMRI studies have found that reward

value modulates sensory-related neural activity, including activity in both early visual areas

(Serences, 2008; Anderson, 2017) and activity in the object-selective ventral temporal cortex

(Hickey and Peelen, 2015; Barbaro et al., 2017; Leong, Radulescu et al., 2017). These findings

highlight the pervasive influence of reward on sensory representations in the brain. Building on

this body of work, we recently demonstrated that using financial incentives to motivate participants

to see a particular interpretation of an ambiguous image enhances neural activity encoding stimulus

features associated with said interpretation (Leong et al., 2019). This selective enhancement biases

participants towards seeing perceptual outcomes that they are motivated to see. The neural

mechanisms driving the selective enhancement of desirable representations, however, remain

unclear.

In the current work, we show that the motivational enhancement of desirable percepts is

dependent on amygdala activity. In particular, motivation enhanced neural activity selective to

desirable perceptual features, but only on trials with high amygdala activity. Heightened amygdala

activity preceded participants categorizing an image as the category they were motivated to see,

and participants with stronger amygdala activation also exhibited strong motivational biases in

their perceptual categorizations. To understand the decision processes affected by the amygdala,

we fit a DDM to participants' data and trial-by-trial amygdala activity. The model fits suggest that

increased amygdala activity biases sensory evidence accumulation in favor of desirable percepts.

Together, the current findings provide converging behavioral, neural and modeling evidence of

the amygdala's role in biasing perceptual decisions towards desirable perceptual outcomes.

Historically, the amygdala has been associated with the processing of fear and threat-

related stimuli (LeDoux, 2000, c.f. Visser et al., 2021). More recent perspectives have emphasized

a broader role of the amygdala in processing motivationally salient information, including

information relevant for predicting and obtaining rewards (Pessoa and Adolphs, 2010; Pourtois et

al., 2013). For example, activity in the amygdala has been shown to encode the appetitive value of

stimuli (Gottfried et al., 2003; Jenison et al., 2011; Lichtenberg et al., 2017; Shanahan et al., 2021).

In a series of studies with non-human primates, Peck and colleagues have also found that amygdala

activity tracks spatial attention towards stimuli associated with reward (Peck et al., 2013; Peck and

Salzman, 2014a, 2014b). Our findings are consistent with this broader view of amygdala function

that considers its role in processing reward-relevant information. Notably, our results suggest that

the amygdala is not only sensitive to the value of reward-relevant stimuli, but also flexibly

enhances the neural representation of what is valuable at a given moment, which in turn gives rise

to a perceptual bias towards percepts that are desirable.

The amygdala can enhance sensory representations both via direct connections to sensory

areas (Freese and Amaral, 2005; Gschwind et al., 2012), as well as via indirect pathways involving

frontoparietal regions and brainstem neuromodulatory systems (Lim et al., 2009; Pourtois et al.,

2013). Recent theories have proposed that the amygdala interacts with the locus-coeruleus

norepinephrine (LC-NE) system to selectively amplify the representation of physically and

affectively salient stimuli (Markovic et al., 2014; Mather et al., 2016). According to these theories,

the amygdala enhances cortical processing of salient stimuli through its direct projections to

sensory cortices. At the same time, the amygdala recruits the LC which then releases NE into the

cortex. NE released in sensory areas increase the gain of sensory neurons selective to salient

stimuli, such that they are more likely to fire to target inputs (Berridge and Waterhouse, 2003). In

addition, NE released in the amygdala further enhances the saliency signal, resulting in even

greater amplification of the neural response to salient stimuli (Gallagher and Holland, 1994).

Amygdala activity has indeed been found to be associated with reward-driven changes in

pupil dilation (Watanabe et al., 2019), a known correlate of LC-NE activity (Murphy et al., 2014;

Joshi et al., 2016). Using a variant of the current task, we recently demonstrated that motivational

biases in sensory evidence accumulation were associated with increased pupil dilation (Leong et

al., 2021). Thus, there is reason to hypothesize that the amygdala-dependent motivational

enhancement of sensory representations observed in the current study were mediated by the LC-

NE system. In particular, wanting to see a percept could have imbued affective salience to the

associated perceptual features. The amygdala then amplifies the neural response to these features

through coordinated action with the LC-NE system. The small size and location of the LC present

challenges to imaging the LC using fMRI (Liu et al., 2017). Recent studies have been able to take

advantage of novel approaches to functionally and anatomically localize the LC (de Gee et al.,

2017; Grueschow et al., 2021). Future work can use these approaches to simultaneously measure

activity in the amygdala, LC and sensory regions to further characterize the neural circuitry

underlying motivational biases in perceptual decisions.

Prior work examining reward-related biases in perceptual decision-making have focused

on the role of frontoparietal regions (Mulder et al., 2014). Here, we did not find evidence of the

involvement of frontoparietal regions in enhancing desirable sensory evidence, suggesting that our

results arose from distinct biasing mechanisms. We note that in earlier studies that assign

asymmetric rewards to perceptual options, participants would earn the larger reward only if they

correctly categorized a stimulus as the category associated with larger reward (Summerfield and

Koechlin, 2010; Mulder et al., 2012). In such settings, participants would earn more reward over

the course of the experiment if they biased their responses towards the category associated with

larger reward (Bogacz et al., 2006; Feng et al., 2009; Fan et al., 2018). Thus, reward-related biases

in these earlier experiments have largely been interpreted as a shift in response strategy to

maximize reward on the task and are not necessarily related to biases in sensory evidence

accumulation. Consistent with this idea, these studies have found that reward biases perceptual

responses without a corresponding effect on sensory representations.

In our task, however, the additional reward associated with the motivation consistent

category was not contingent on participants' responses - participants won or lost the bonus based

on the objective category of the image, regardless of their subjective categorizations. Hence, the

bias observed in our task was unlikely to have resulted from a shift in response strategy to

maximize reward. Instead, it was likely driven by participants wanting to see one percept over

another. Using a drift diffusion model, we had previously found that this motivational bias reflects

both an *a priori* response bias towards outcomes participants want to see (i.e. a bias in the starting

point of evidence accumulation), and a perceptual bias in how sensory evidence accumulates over

time (i.e. a bias in the drift rate) (Leong et al., 2019). The current results indicate that amygdala

activity was specifically associated with the perceptual bias and not the response bias. Thus, the

amygdala-dependent biases in perceptual decisions are potentially distinct from the response-

related biases associated with frontoparietal regions documented in earlier studies.

In everyday life, people are rarely neutral observers indifferent to different perceptual

outcomes. They are motivated to see some outcomes over others, and exhibit a bias to report seeing

those outcomes over possible alternatives. The current work suggests that this bias is associated

with the amygdala-dependent enhancement of desirable neural representations in sensory cortices.

Together, these results expand our understanding of the neural mechanisms underlying

motivational influences on perceptual decision-making. Notably, similar motivational biases have

been documented across a broad array of decision-making contexts (Hughes and Zaki, 2015). In

one recent study, participants were asked to judge whether they were in a desirable state (i.e. an

environment with greater rewards than loses) or an undesirable state (i.e. an environment with low

probability of reward) by sampling evidence that was probabilistically associated with each state

(Gesiarz et al., 2019). The authors found that participants accumulated evidence in a manner that

was biased towards the belief they were in a desirable state, much like the bias towards desirable

visual percepts observed in our study. Could the amygdala be involved in biases in evidence

accumulation beyond sensory perception? Given that the amygdala has also been implicated in

value-based learning and decision-making (Jenison et al., 2011; Lichtenberg et al., 2017; Prévost

et al., 2013), this is certainly possible. We believe that exploring the role of the amygdala in

mediating motivational biases in other reasoning and evaluative processes would be a fruitful

direction for future research.

References

Anderson BA (2017) Reward processing in the value-driven attention network: reward signals tracking cue identity and location. Social Cognitive and Affective Neuroscience 12:461–467.

Anderson BA, Laurent PA, Yantis S (2011) Value-driven attentional capture. PNAS 108:10367–10371.

Barbaro L, Peelen MV, Hickey C (2017) Valence, Not Utility, Underlies Reward-Driven Prioritization in Human Vision. J Neurosci 37:10438–10450.

Berridge CW, Waterhouse BD (2003) The locus coeruleus–noradrenergic system: modulation of behavioral state and state-dependent cognitive processes. Brain Research Reviews 42:33–84.

Boehm U, Annis J, Frank MJ, Hawkins GE, Heathcote A, Kellen D, Krypotos A-M, Lerche V, Logan GD, Palmeri TJ, van Ravenzwaaij D, Servant M, Singmann H, Starns JJ, Voss A, Wiecki TV, Matzke D, Wagenmakers E-J (2018) Estimating across-trial variability parameters of the Diffusion Decision Model: Expert advice and recommendations. Journal of Mathematical Psychology 87:46–75.

Bogacz R, Brown E, Moehlis J, Holmes P, Cohen JD (2006) The Physics of Optimal Decision Making: A Formal Analysis of Models of Performance in Two-Alternative Forced-Choice Tasks. Psychological Review 113:700–765.

Brainard DH (1997) The Psychophysics Toolbox. Spatial Vis 10:433–436.

Bromberg-Martin ES, Sharot T (2020) The Value of Beliefs. Neuron 106:561–565.

de Gee JW, Colizoli O, Kloosterman NA, Knapen T, Nieuwenhuis S, Donner TH (2017) Dynamic modulation of decision biases by brainstem arousal systems. eLife 6:e23232.

Domenech P, Dreher J-C (2010) Decision Threshold Modulation in the Human Brain. J Neurosci 30:14305–14317.

Dunning D, Balcetis E (2013) Wishful Seeing: How Preferences Shape Visual Perception. Curr Dir Psychol Sci 22:33–37.

Fan Y, Gold JI, Ding L (2018) Ongoing, rational calibration of reward-driven perceptual biases. eLife 7:e36018.

Feng S, Holmes P, Rorie A, Newsome WT (2009) Can Monkeys Choose Optimally When Faced with Noisy Stimuli and Unequal Rewards? PLOS Computational Biology 5:e1000284.

Forstmann BU, Ratcliff R, Wagenmakers E-J (2016) Sequential Sampling Models in Cognitive Neuroscience: Advantages, Applications, and Extensions. Annual Review of Psychology 67:641–666.

Freese JL, Amaral DG (2005) The organization of projections from the amygdala to visual cortical areas TE and V1 in the macaque monkey. Journal of Comparative Neurology 486:295–317.

Gallagher M, Holland PC (1994) The amygdala complex: multiple roles in associative learning and attention. PNAS 91:11771–11776.

Gesiarz F, Cahill D, Sharot T (2019) Evidence accumulation is biased by motivation: A computational account. PLOS Computational Biology 15:e1007089.

Gottfried JA, O'Doherty J, Dolan RJ (2003) Encoding Predictive Reward Value in Human Amygdala and Orbitofrontal Cortex. Science 301:1104–1107.

Grill-Spector K (2003) The neural basis of object perception. Current Opinion in Neurobiology
13:159–166.
Grueschow M, Stenz N, Thörn H, Ehlert U, Breckwoldt J, Brodmann Maeder M, Exadaktylos
AK, Bingisser R, Ruff CC, Kleim B (2021) Real-world stress resilience is associated with the
responsivity of the locus coeruleus. Nat Commun 12:2275.
Gschwind M, Pourtois G, Schwartz S, Van De Ville D, Vuilleumier P (2012) White-Matter
Connectivity between Face-Responsive Regions in the Human Brain. Cerebral Cortex
22:1564–1576.
Heekeren HR, Marrett S, Ungerleider LG (2008) The neural systems that mediate human
perceptual decision making. Nature Reviews Neuroscience 9:467–479.
Hickey C, Peelen MV (2015) Neural Mechanisms of Incentive Salience in Naturalistic Human
Vision. Neuron 85:512–518.
Ho TC, Brown S, Serences JT (2009) Domain General Mechanisms of Perceptual Decision
Making in Human Cortex. J Neurosci 29:8675–8687.
Hughes BL, Zaki J (2015) The neuroscience of motivated cognition. Trends in Cognitive
Sciences 19:62–64.
Jenison RL, Rangel A, Oya H, Kawasaki H, Howard MA (2011) Value Encoding in Single
Neurons in the Human Amygdala during Decision Making. J Neurosci 31:331–338.
Joshi S, Li Y, Kalwani RM, Gold JI (2016) Relationships between Pupil Diameter and Neuronal
Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. Neuron 89:221–234.
Kube T, Rozenkrantz L (2021) When Beliefs Face Reality: An Integrative Review of Belief
Updating in Mental Health and Illness. Perspect Psychol Sci 16:247–274.
Le Pelley ME, Pearson D, Griffiths O, Beesley T (2015) When goals conflict with values:
Counterproductive attentional and oculomotor capture by reward-related stimuli. Journal of
Experimental Psychology: General 144:158–171.
LeDoux JE (2000) Emotion Circuits in the Brain. Annu Rev Neurosci 23:155–184.
Leong YC, Dziembaj R, D'Esposito M (2021) Pupil-Linked Arousal Biases Evidence
Accumulation Toward Desirable Percepts During Perceptual Decision-Making. Psychol Sci
32:1494-1509
Leong YC, Hughes BL, Wang Y, Zaki J (2019) Neurocomputational mechanisms underlying
motivated seeing. Nature Human Behaviour 3:962–973.
Leong YC, Radulescu A, Daniel R, DeWoskin V, Niv Y (2017) Dynamic Interaction between
Reinforcement Learning and Attention in Multidimensional Environments. Neuron 93:451–
463.
Lichtenberg NT, Pennington ZT, Holley SM, Greenfield VY, Cepeda C, Levine MS & Wassum,
KM (2017). Basolateral amygdala to orbitofrontal cortex projections enable cue-triggered
reward expectations. J Neurosci, 37:8374-8384.
Lim S-L, Padmala S, Pessoa L (2009) Segregating the significant from the mundane on a
moment-to-moment basis via direct and indirect amygdala contributions. PNAS 106:16841–
16846.

Liu KY, Marijatta F, Hämmerer D, Acosta-Cabronero J, Düzel E, Howard RJ (2017) Magnetic
resonance imaging of the human locus coeruleus: A systematic review. Neuroscience &
Biobehavioral Reviews 83:325–355.
Ma DS, Correll J, Wittenbrink B (2015) The Chicago face database: A free stimulus set of faces
and norming data. Behav Res 47:1122–1135.
Markovic J, Anderson AK, Todd RM (2014) Tuning to the significant: Neural and genetic
processes underlying affective enhancement of visual perception and memory. Behavioural
Brain Research 259:229–241.
Mather M, Clewett D, Sakaki M, Harley CW (2016) Norepinephrine ignites local hotspots of
neuronal excitation: How arousal amplifies selectivity in perception and memory. Behavioral
and Brain Sciences 39
Mulder MJ, van Maanen L, Forstmann BU (2014) Perceptual decision neurosciences – A model-
based review. Neuroscience 277:872–884.
Mulder MJ, Wagenmakers E-J, Ratcliff R, Boekel W, Forstmann BU (2012) Bias in the Brain: A
Diffusion Model Analysis of Prior Probability and Potential Payoff. J Neurosci 32:2335–
2343.
Mumford JA, Davis T, Poldrack RA (2014) The impact of study design on pattern estimation for
single-trial multivariate pattern analysis. NeuroImage 103:130–138.
Murphy PR, O'Connell RG, O'Sullivan M, Robertson IH, Balsters JH (2014) Pupil diameter
covaries with BOLD activity in human locus coeruleus. Human Brain Mapping 35:4140–
4154.
Peck CJ, Lau B, Salzman CD (2013) The primate amygdala combines information about space
and value. Nat Neurosci 16:340–348.
Peck CJ, Salzman CD (2014a) The Amygdala and Basal Forebrain as a Pathway for
Motivationally Guided Attention. J Neurosci 34:13757–13767.
Peck CJ, Salzman CD (2014b) Amygdala neural activity reflects spatial attention towards stimuli
promising reward or threatening punishment. eLife 3:e04478.
Pessoa L, Adolphs R (2010) Emotion processing and the amygdala: from a "low road" to "many
roads" of evaluating biological significance. Nature Reviews Neuroscience 11:773–782.
Pourtois G, Schettino A, Vuilleumier P (2013) Brain mechanisms for emotional influences on
perception and attention: What is magic and what is not. Biological Psychology 92:492–512.
Prévost C, McNamee D, Jessup RK, Bossaerts P, O'Doherty JP (2013) Evidence for Model-
based Computations in the Human Amygdala during Pavlovian Conditioning. PLOS
Computational Biology 9:e1002918.
Rossi-Goldthorpe RA, Leong YC, Leptourgos P, Corlett PR (2021) Paranoia, self-deception and
overconfidence. PLOS Computational Biology 17:e1009453.
Rutishauser U, Mamelak AN, Adolphs R (2015) The primate amygdala in social perception –
insights from electrophysiological recordings and stimulation. Trends in Neurosciences
38:295–306.
Serences JT (2008) Value-Based Modulations in Human Visual Cortex. Neuron 60:1169–1181.

Shadlen MN, Kiani R (2013) Decision Making as a Window on Cognition. Neuron 80:791–806.

Shanahan LK, Bhutani S, Kahnt T (2021) Olfactory perceptual decision-making is biased by
motivational state. PLOS Biology 19:e3001374.

Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD (2012) Decoding subject-driven
cognitive states with whole-brain connectivity patterns. Cereb Cortex 22:158–165.

Summerfield C, Koechlin E (2010) Economic Value Biases Uncertain Perceptual Choices in the
Parietal and Prefrontal Cortices. Frontiers in Human Neuroscience 4:208.

Venables WN, Ripley BD (2013) Modern Applied Statistics with S-PLUS. Springer Science &
Business Media.

Visser RM, Bathelt J, Scholte HS, Kindt M (2021) Robust BOLD responses to faces but not to
conditioned threat: challenging the amygdala's reputation in human fear and extinction
learning. J Neurosci.

Voss A, Rothermund K, Brandtstädter J (2008) Interpreting ambiguous stimuli: Separating
perceptual and judgmental biases. Journal of Experimental Social Psychology 44:1048–1056.

Watanabe N, Bhanji JP, Ohira H, Delgado MR (2019) Reward-Driven Arousal Impacts
Preparation to Perform a Task via Amygdala–Caudate Mechanisms. Cerebral Cortex
29:3010–3022.

White CN, Mumford JA, Poldrack RA (2012) Perceptual Criteria in the Human Brain. J
Neurosci 32:16716–16724.

Wiecki TV, Sofer I, Frank MJ (2013) HDDM: Hierarchical Bayesian estimation of the Drift-
Diffusion Model in Python. Front Neuroinform 7:14