Methylation pattern of *nc886* in non-human mammals

Daria Kostiniuk[1¥], Hely Tamminen[1¥], Pashupati P. Mishra[2], Saara Marttila[1,3#] Emma Raitoharju[1#]*

[1]Molecular Epidemiology, Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

[2]Department of Clinical Chemistry, Pirkanmaa Hospital District, Fimlab Laboratories, and Finnish Cardiovascular Research Center, Tampere, Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

[3]Gerontology Research Center, Tampere University, Finland

[¥] and [#] Equal contribution

***Corresponding Author**: Emma Raitoharju, PhD, Molecular Epidemiology, Tampere University, Tampere, Finland. Tel: +358 50 318 7676.

Email: emma.raitoharju@tuni.fi

Abstract

Background: In humans, the *nc886* locus is a polymorphically imprinted metastable

epiallele. Periconceptional conditions have an effect on the methylation status of *nc886*, and

further, this methylation status is associated with health outcomes in later life, in line with the

Developmental Origins of Health and Disease (DOHaD) hypothesis. Animal models would

offer opportunities to study the associations between periconceptional conditions, *nc886*

methylation status and metabolic phenotypes further. Thus, we set out to investigate the

methylation pattern of the *nc886* locus in non-human mammals.

Data: We obtained DNA methylation data from the data repository GEO for mammals,

whose *nc886* gene included all three major parts of *nc886* and had sequence similarity of

over 80% with the human *nc886*. Our final sample set consisted of DNA methylation data

from humans, chimpanzees, bonobos, gorillas, orangutangs, baboons, macaques, vervets,

marmosets and guinea pigs.

Results: In human data sets the methylation pattern of *nc886* locus followed the expected

bimodal distribution, indicative of polymorphic imprinting. In great apes, we identified a

unimodal DNA methylation pattern with 50% methylation level in all individuals and in all

subspecies. In Old World monkeys, the between individual variation was greater and

methylation on average was close to 60%. In guinea pigs the region around the *nc886*

homologue was non-methylated. Results obtained from the sequence comparison of the

CTCF binding sites flanking the *nc886* gene support the results on the DNA methylation

data.

Conclusions: Our results indicate that unlike in humans, *nc886* is not a polymorphically

imprinted metastable epiallele in non-human primates or in guinea pigs, thus implying that

animal models are not applicable for *nc886* research. The obtained data suggests that the

nc886 region may be classically imprinted in great apes, and potentially also in Old World

monkeys, but not in guinea pigs.

Key words: nc886, vtRNA2-1, primates, imprinting, DNA methylation

Background

In mammalian genomic imprinting, only one parental allele is expressed, while gene expression from the other allele is suppressed in a parent-of-origin-dependent manner. The expression of imprinted genes in general has been associated with fetal and placental growth and suggested to a have a role in the development of cardiometabolic diseases in adulthood [1–3]. Imprinting arose relatively recently at most loci— while only a few imprinted genes in Eutherians are also imprinted in marsupials, to date no imprinting has been reported in the egg-laying monotreme mammals [4]. Genetic imprinting is best described in mice, and while the mouse is an informative proxy for human imprinted gene regulation, less than half of the 100 human imprinted genes have been shown to be similarly imprinted in mice (https://www.geneimprint.com/site/home). Distinct differences in placental evolution, physiology, and reproductive biology of the primate and murine groups may be responsible for these differences.

During gametogenesis and fertilization, the original DNA methylation pattern of imprinted genes is erased, and parent of origin-based methylation pattern is established. While most imprinted genes are located in clusters that are regulated by insulators or long noncoding RNAs [5], some unclustered imprinted genes can be regulated by differential promoter methylation [3]. Parental imprints are maintained after fertilization through these mechanisms despite extensive reprogramming of the mammalian genome [3]. A common feature of imprinted genes is insulators, such as CCCTC binding factor (CTCF) binding sites, which block the enhancers from interacting with gene promoters and/or act as barrier to the spread of transcriptionally repressive condensed chromatin [6].

Due to the important developmental roles of the imprinted genes, imprinting is tightly fixed across human populations, with all individuals displaying monoallelic expression of

imprinted genes. An exception to this is the *non-coding 886* (*nc886*, also known as *VTRNA2-*

1), which is the only polymorphically imprinted metastable epiallele that has been repeatedly

described in humans in the literature. Non-coding RNA 886 is encoded in chromosome

5q31.1, from a 1.9-kb long, differentially methylated region (DMR), the boundaries of which

are marked by two CTCF binding sites [7,8]. This DMR has been shown to present maternal

imprinting in ~75% of individuals in several populations [7,9,10]. This means that while in

all individuals the paternal allele is unmethylated, in approximately 75% of individuals the

maternal allele is methylated (individuals present a 50% methylation level at *nc886* locus and

are hemi-methylated) and in the remaining 25% of individuals the maternal allele is

unmethylated (individuals present a 0% methylation level at the *nc886* locus and are non-

methylated).

The *nc886* gene codes for a 102nt long, non-coding RNA, which is then cleaved into two

short RNAs (hsa-miR-886-3p/nc886-3p [23 nt] and hsa-miR-886-5p/nc886-5p [24–25 nt])

[11–13]. There is no consensus on whether the effects of *nc886* expression is mediated by the

102 nt long hairpin structure or the nc886-3p and -5p molecules, as the short molecules have

been indicated to function as miRNAs, while the hairpin loop has been shown to inhibit

protein kinase R (PKR) [11,14].

The periconceptional environment has been suggested to affect DNA methylation patterns in

maternal alleles[15], including the *nc886* epiallele [7,8,10]. Season of conception, maternal

age and socioeconomic status have been linked to changes in the proportion of non- and

hemi-methylated offspring [7,8,10]. On the other hand, lower levels of *nc886* methylation

have been linked to cleft palate [16], and a non-methylated *nc886* epiallele has been

associated with an elevated childhood BMI [17]. The methylation status of this epiallele has

also been associated with allergies [18], asthma [19], infections [20], and inflammation [21].

We and others have also shown that both the *nc886* methylation status and RNA expression

are associated with indicators of glucose metabolism [10,22].

These results indicate that *nc886* could mediate the association between periconceptional

conditions and later metabolic health, in line with the Developmental Origins of Health and

Disease (DOHaD) hypothesis (aka the Barker hypothesis) [23]. More detailed analysis on

periconceptional conditions and *nc886* methylation status and investigations between *nc886*

and metabolic phenotypes, with less cofounding factors, would be needed to confirm this

hypothesis. As carcinogenesis [24] and pluripotency induction [10] affect the DNA

methylation pattern in *nc886* locus, *in vitro* work has its limitations. Animal models could be

a feasible option for this research. Unfortunately, rodents do not harbor the *nc886* gene,

limiting the use of traditional model organisms [7]. Thus, this study was set up to investigate

1) which animals have *nc886* gene, 2) whether this gene is surrounded by similar CTFC

elements as the human homolog and 3) whether the methylation status of the *nc886* region

suggest polymorphic imprinting in non-human mammals.

Materials and methods

The presence of *nc886* gene was investigated in ensemble, in 65 amniota vertebrates Mercator-Pecan collection and 24 primates EPO-extended collection [25]. To select species for further investigation, we required the *nc886* gene have 80% sequence similarity with the human homolog and to present the sequences for nc886-3p and nc886-5p RNAs, as well as the loop structure, previously shown to mediate the binding of PKR [11,26] (Supplementary figures S1 and S2). The existence and sequence similarity of the centromeric (chr5:135415115-135415544) and telomeric CTCF (chr5:135418124-135418523) binding site flanking *nc886* gene were also investigated in species shown to harbor intact *nc886* gene. If homologous CTCF-binding sites were not discovered, CTCFBSDB 2.0 [27] was utilized to predict possible non-homologous sites. Interactions of the *nc886* flanking CTCF-sites were also investigated using K562 CTCF CiA-PET Interactions data in genome browser [28].

For species harboring the *nc886* gene, we investigated the Gene Expression Omnibus (GEO) repository [29] for available DNA methylation data, with both general and binomial name of the species. DNA methylation data was available in apes from chimpanzees (*Pan troglodytes*, n=83; GSE136296 [30] and n=5; GSE41782 [31]), bonobos (*Pan paniscus*, n=6; GSE41782 [31]), gorillas (*Troglodytes gorilla*, n=6; GSE41782 [31]) and orangutangs (*Pongo spp.*, n=6; GSE41782[31]). In Old World monkeys, data was obtained from baboons (*Papio spp.* n=28; GSE103287 [32]), rhesus macaques (*Macaca mulatta,* n=10; GSE103287 [32]), vervets (*Chlorocebus aethiops*, n=10; GSE103287 [32]) and in New World monkeys, from marmosets (*Callithrix jacchus*, n=6; GSE103287 [32]). From primates, only data from blood or femur was utilized. In addition to primates, we obtained DNA methylation data from guinea pig hippocampi (*Cavia porcellus,* n=36; GSE98549 [33]). As reference, we utilized data from human (*Homo sapiens*) blood (n=1658; GSE105018 [34]), femur (n=48; GSE64490 [35]) and hippocampus (n=33; GSE72778 [36]).

Methylation profiling data obtained by high throughput sequencing (guinea pigs, GSE98549) was processed as follows. Quality of the paired-end reads in all the samples was assessed using FastQC [37] and MultiQC [38]. Paired-end fastq files were trimmed using Trimmomatic-0.39 with a sliding window of size 4 set to remove bases with phred score lower than 20 [39]. The trimmed samples were analyzed using Bismark-0.23.0 tools [40]. The reads were aligned to the guinea pig genome (cavPor3). Duplicate alignments, which can arise for example by PCR amplification, were removed. Methylation information was extracted from the alignment result files using Bismark's methylation extractor. DNA methylation values for CpGs inside the gene were first inspected and then a wider region (±2000nt) around the gene was investigated.

Primate DNA methylation data from GSE41782, GSE105018, GSE64490, GSE72778 (profiled with Illumina 450K) and GSE136296 (profiled with Illumina EPIC) were available as processed data and was used as such. Primate DNA methylation data from GSE103271, GSE103280, GSE103286, which are subseries of GSE103332 (profiled with Illumina EPIC), were available as raw data, and were normalized by using minfi quantile normalization for each species separately.

From primate data, the 14 CpGs in the *nc886* DMR previously reported to show bimodal methylation pattern in humans were retrieved [7,10]. In all the primate species, for which methylation data was available, the sequence on the binding site of the Illumina probes was investigated. Only data from sites with the CG-sequence intact in the species in question were further utilized. Also probes, which had mutations in the probe binding site and methylation pattern clearly distinct from neighboring sites, were discarded. Similar process was repeated for 50 probes in paternally expressed 10 (*PEG10*) previously shown to be imprinted [41]. *PEG10* was used as evolutionally conserved reference for a classically imprinted gene [4].

Results and Discussion

nc886 **gene in non-human mammals**

Human *nc886* has been suggested to be an evolutionarily young gene, producing a 102 nt long

RNA, which is then ineffectively cleaved to two miRNA-like RNAs [11]. In line with

previous reports [7]*, nc886* gene, with intact short RNA coding sequences and the hairpin

loop, can be found in primates, in guinea pig (*Cavia porcellus*), Eurasian red squirrel (*Sciurus*

vulgaris), and Alpine marmot (*Marmota marmota*), with two of the latter having insertions in

the centromeric end of the gene (Figure 1, Supplementary figure S1).

Upon further inspection of primate *n886* gene, almost 100% sequence similarity was

identified in apes (*hominoidea*). The sequence similarity was high (over 98%) between

humans and the investigated Old World monkeys (*Cercopithecidae*), while less (91-93%)

similarity can be seen between humans and New World monkeys (*Ceboidea*) and even less

(84-89%) between humans and tarsiers (*Tarsiidae*) or lemurs (*Lemuroidea*). Sequences

coding for the short nc886 RNAs are identical within Old World anthropoids (*Catarrhini).*

Differences between human and New World monkey and prosimian *nc886* sequences can be

found both in the short nc886 RNA and the hairpin coding regions, but the most significant

differences can be found in the centromeric end of the gene (Figure 1 and Supplementary

figure S2).

CTCF binding site sequence in mammals with *nc886* gene

In humans, the *nc886* locus is flanked by two CTCF binding sites [7]. The telomeric CTCF

binding site can be found in most vertebrates, with the binding sequence being identical in all

species showed to have the *nc886* gene. The telomeric CTCF binding site was shown to

interact with a CTCF binding site (chr5:135222814-135223707) locating near the *IL9* gene in

the CTCF CiA-PET data. This prediction is in line with our previous finding indicating that

genetic polymorphisms near *IL9* gene are associated with the expression of nc886 RNAs

[10]. Together these results suggest that the evolutionally conserved telomeric CTCF binding

site of *nc886* interacts with CTCF binding sites near *IL9* gene, possibly forming a

topologically associating domain (TAD), or an interaction within one (sub-TAD), and

bringing the suggested enhancer area near the *nc886* gene [42]. On the contrary, the

centromeric CTCF binding site is present only in primates and even in primates, the binding

sequence cannot be identified in marmosets. For the centromeric CTCF binding site no

interactions were detected according to CTCF CiA-PET data. There are also changes in the

CTFC binding sequence in gorillas (position 9), in all Old World monkeys (position 4) and

also in New World monkeys (position 14). According to the CTCFBSDB 2.0, the guinea pig

genome does not harbor any non-homologous predicted CTCF binding sites in the

centromeric side of the *nc886* gene.

Guinea pigs present a non-methylated *nc886* locus

One of the aims of this study was to investigate whether the information gathered from the

nc886 gene and DMR from cell culture and population studies could be supplemented with

research on animal models. In line with a previous report [7] we identified this gene only in

primates, guinea pigs and few members of the squirrel family, of which guinea pig was the

most promising candidate as a model organism. In data from Boureau et al. [33], in guinea

pig hippocampi the whole *nc886/vtRNA2-1* gene was non-methylated (Supplementary figure

S3A). The surrounding *nc886* region (Scaffold DS562872.1: 24,622,179-24,622,280) +/-

2000 nt was mostly unmethylated, with only 2% of the reads in the region being methylated.

It should be noted that the number of reads in the region in the data utilized was low (max

number of reads=17, average number of reads=7). *nc886* methylation pattern in human

hippocampi presented the expected bimodal distribution, and thus the discovered methylation

pattern in the guinea pig hippocampi was most likely not due to the selection of tissue

(Supplementary figure S3B). This identified lack of methylation in the *nc886* locus is

compatible with the absence of the telomeric CTCF binding site, as CTCF binding sites can

delineate the boundaries of an imprinted region [43]. These results thus suggest that guinea

pigs are not suitable model organisms for the investigation of establishment of *nc886*

methylation status.

Imprinted *nc886* region in great apes

The blood of chimpanzees, gorillas, bonobos and orangutans presented beta-values close to

0.5 with a unimodal distribution in the *nc886* region (Figure 2). This methylation pattern

closely resembles the methylation pattern in the known maternally imprinted gene *PEG10*

(Supplementary figure S4). The methylation levels are also very similar to those presented in

hemi-methylated humans (Figure 2). In these data sets, that include more than 110 apes, we

did not identify any individuals with methylation level close to 0 in the *nc886* locus, whereas

in humans 25% of the population present a methylation level close to 0 at this locus [7,10] . If

the prevalence on non- and hemi-methylated individuals in apes was similar to humans,

already 11 individuals would present at least one non-methylated individual with 95%

probability. Of individual species, we had the largest dataset for chimpanzees (n=83 in

GSE13629631 and n=5 in GSE41782). Again, assuming the same proportion of non-

methylated individuals as in humans (25%), probability of not identifying any non-

methylated chimpanzees in a population of 88 individuals is extremely low, $1.01*10^{-11}$. These

results imply that the *nc886* locus is not polymorphically imprinted in apes.

Within species standard deviation of the methylation in a probe is on average around 0.04,

which is comparable to that seen in *PEG10* in apes (SD on average 0.04) and in human blood

(0.02), indicating good data quality. In all apes the methylation levels of *nc886* region

resembled those seen in *PEG10,* which is an evolutionally conserved maternally imprinted

gene [4]. It is thus reasonable to suggest that *nc886* could be classically imprinted in other

great apes, excluding humans.

***nc886* region methylation patterns in Old World monkeys**

The patterns of *nc886* region methylation are very similar in all of the Old World monkeys.

The median methylation level is close to 0.60, higher as compared to apes. The

interindividual variation is larger, especially in baboons, than in apes, with the average SD

within a probe being 0.11. The SD within a probe is also higher in probes locating in *PEG10*

in baboons, where the within probe SD is on average 0.07. As the between individual

variation in methylation levels of a known evolutionally conserved imprinted gene is also

higher, this suggests a technical bias in the data, potentially due to the use of Illumina

Infinium 450K and EPIC methylation assays, that are designed for humans. All methylation

data available for Old World monkeys was from femur, but as in the human reference data set

form femur samples both *nc886* and *PEG10* present similar methylation patterns as in blood

(Figure 3 and Supplementary figure S5), this phenomenon most likely is not caused by the

tissue of origin. Regardless of the precise methylation levels of the Old World monkeys, in

the 48 individuals we did not identify any presenting a non-methylated methylation pattern in

nc886 region, probability of which is $1.0*10^{-6}$, when assuming similar distribution as in

humans. This implies that similar to non-human great apes, the *nc886* locus is not

polymorphically imprinted in Old World monkeys.

nc886 region methylation patterns in New World monkeys

Data from the six marmosets is not conclusive, as we found that only 3 of the probes in *nc886*

region bound to areas with no great sequence differences. The methylation levels of two of

these probes were around 0.8 and one 0.32, showing no indications of imprinting, while the

methylation beta values in the probes locating in the *PEG10* are on average 0.48 in

marmosets (Figure 3 and Supplementary figure S5). The lack of imprinting of any kind in

marmosets is further supported by the finding that they lack the centromeric CTCF binding

sequence, which is thought to have an important role in insulating the DMR [7].

Limitations of the study

Our study is purely descriptive in nature. In guinea pigs, the shallowness of the sequencing

data limits the ability to make conclusions of the methylation pattern. In non-human primates,

utilizing methylation arrays that have been designed for humans raises questions on data

quality, especially in marmosets. Concerns over data quality are however mitigated by the

observed methylation pattern in well-established imprinted gene, *PEG10*, as well as the

consistency of observed results across different species and data sets.

Conclusions

We describe here an analysis on the methylation status of *nc886* region in non-human

mammals. A genetic locus, with more than 80% similarity to human *nc886* gene can be found

in primates, guinea pigs and some members of the squirrel family. We obtained DNA

methylation data for 8 different non-human primate species and for guinea pigs, and in none

of these species could we observe methylation pattern indicative of similar polymorphic

imprinting, as could be observed, and has been reported [7,9,10], in humans. The observed

methylation pattern in apes and in Old World monkeys implies that the *nc886* region might

be classically imprinted, although these findings have to be interpreted with caution, as apart

from chimpanzees the sample number were low, and in Old World monkeys the variation

between individuals was notable. In guinea pigs, the most feasible potential model organism

of those harboring the *nc886* locus, the data indicated that the locus is completely

unmethylated. It is noteworthy that only primates, whose genome also contained the

centromeric CTCF binding sequence flanking the *nc886* gene, had methylation levels

indicative of genetic imprinting.

As such, it appears there are no animal models suited to study the establishment of the

methylation pattern of the polymorphically imprinted metastable epiallele *nc886*. Further

studies on how this kind of unusual metastable developed, and how it links to the

periconceptional conditions and later life health traits, are thus restricted to *in vitro* and

population studies.

Declarations

Ethics approval:

Datasets used in this study were retrieved from Gene Expression Omnibus (GEO,

www.ncbi.nlm.nih.gov/geo/) repository. For human data sets, informed consent or "Consent

for Autopsy" were given by all participants. For animal studies all protocols were approved

by the local ethical committees and/or the samples were collected during standard veterinary

checks or routine necropsies. Details can be found from the original publications[30–36].

Consent for publication: Not applicable

Availability of data and materials: All methylation data utilized are available in GEO,

under accession numbers GSE136296, GSE41782, GSE103287, GSE98549, GSE105018,

GSE64490 and GSE72778.

Competing interests: The authors declare that they have no competing interests

Funding: This study has been supported by the Academy of Finland (https://www.aka.fi/en/)

(grants 330809 and 338395 (E.R.)); the Tampere University Hospital Medical Funds

(shorturl.at/gsyV0) (grants 9X047, 9S054, and 9AB059 for E.R.); the Yrjö Jahnsson

Foundation (https://www.yjs.fi/en/) (S.M. and E.R); Orion Research Foundation

(https://www.orion.fi/en/rd/orion-research-foundation/) (E.R.); the Foundation of Clinical

Chemistry (https://www.juhaniahonlaaketieteensaatio.fi/) (E.R), Laboratoriolääketieteen

edistämissäätiö sr. (https://www.labes.fi/) (E.R.) and the Paulo Foundation

(https://www.paulo.fi/in-english) (S.M). The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions: S.M. and E.R. designed the research; D.K., H.T., P.P.M., S.M. and

E.R. performed research; S.M. and E.R. provided resources; D.K., H.T., P.P.M., S.M. and

E.R. curated data; S.M. and E.R wrote the original draft manuscript; D.K., H.T., P.P.M., S.M.

and E.R. reviewed and edited the manuscript. Funding was acquired by S.M and E.R. All

authors have read and agreed upon the published version of the manuscript.

Acknowledgements: The authors wish to thank Jooseppi Järvinen (MSc) for his contribution

to the probability calculations, Nina Mononen (PhD) and Binisha Mishra (MSc) for their

valuable input on the manuscript and to acknowledge all the researchers who contributed

towards the collection of datasets utilized in this study.

References:

1. Smith FM, Garfield AS, Ward A. Regulation of growth and metabolism by imprinted

genes. Cytogenetic and Genome Research. 2006. doi:10.1159/000090843

2. Millership SJ, Van de Pette M, Withers DJ. Genomic imprinting and its effects on

postnatal growth and adult metabolism. Cellular and Molecular Life Sciences. 2019.

doi:10.1007/s00018-019-03197-z

3. Barlow DP, Bartolomei MS. Genomic imprinting in mammals. Cold Spring Harbor

Perspectives in Biology. 2014;6. doi:10.1101/cshperspect.a018382

4. Renfree MB, Suzuki S, Kaneko-Ishino T. The origin and evolution of genomic

imprinting and viviparity in mammals. Philosophical Transactions of the Royal

Society B: Biological Sciences. 2013. doi:10.1098/rstb.2012.0151

5. Varmuza S, Mann M. Genomic imprinting - defusing the ovarian time bomb. Trends in

Genetics. 1994;10. doi:10.1016/0168-9525(94)90212-7

6. Lewis A, Reik W. How imprinting centres work. Cytogenetic and Genome Research.

2006. doi:10.1159/000090818

7. Carpenter BL, Zhou W, Madaj Z, DeWitt AK, Ross JP, Grønbæk K, et al. Mother–

child transmission of epigenetic information by tunable polymorphic imprinting.

Proceedings of the National Academy of Sciences of the United States of America.

2018. doi:10.1073/pnas.1815005115

8. Silver MJ, Kessler NJ, Hennig BJ, Dominguez-Salas P, Laritsky E, Baker MS, et al.

Independent genomewide screens identify the tumor suppressor VTRNA2-1 as a

human epiallele responsive to periconceptional environment. Genome Biology. 2015.

doi:10.1186/s13059-015-0660-y

9.   Carpenter BL, Remba TK, Thomas SL, Madaj Z, Brink L, Tiedemann RL, et al.

Oocyte age and preconceptual alcohol use are highly correlated with epigenetic

imprinting of a noncoding RNA (nc886). Proceedings of the National Academy of

Sciences of the United States of America. 2021;118. doi:10.1073/pnas.2026580118

10.   Marttila S, Viiri LE, Mishra PP, Kühnel B, Matias-Garcia PR, Lyytikäinen L-P, et al.

Methylation status of nc886 epiallele reflects periconceptional conditions and is

associated with glucose metabolism through nc886 RNAs. Clinical Epigenetics.

2021;13. doi:10.1186/s13148-021-01132-3

11.   Fort RS, Garat B, Sotelo-Silveira JR, Duhagon MA. vtRNA2-1/nc886 produces a

small RNA that contributes to its tumor suppression action through the microRNA

pathway in prostate cancer. Non-coding RNA. 2020. doi:10.3390/ncrna6010007

12.   Kong L, Hao Q, Wang Y, Zhou P, Zou B, Zhang Y xiang. Regulation of p53

expression and apoptosis by vault RNA2-1-5p in cervical cancer cells. Oncotarget.

2015. doi:10.18632/oncotarget.4948

13.   Lee K, Kunkeaw N, Jeon SH, Lee I, Johnson BH, Kang GY, et al. Precursor miR-886,

a novel noncoding RNA repressed in cancer, associates with PKR and modulates its

activity. RNA. 2011. doi:10.1261/rna.2701111

14.   Lee YS. A Novel Type of Non-coding RNA, nc886, Implicated in Tumor Sensing and

Suppression. Genomics & Informatics. 2015. doi:10.5808/gi.2015.13.2.26

15.   Steegers-Theunissen RPM, Twigt J, Pestinger V, Sinclair KD. The periconceptional

period, reproduction and long-term health of offspring: The importance of one-carbon

metabolism. Human Reproduction Update. 2013. doi:10.1093/humupd/dmt041

16.   Gonseth S, Shaw GM, Roy R, Segal MR, Asrani K, Rine J, et al. Epigenomic profiling

of newborns with isolated orofacial clefts reveals widespread DNA methylation

changes and implicates metastable epiallele regions in disease risk. Epigenetics. 2019.

doi:10.1080/15592294.2019.1581591

17.   van Dijk SJ, Peters TJ, Buckley M, Zhou J, Jones PA, Gibson RA, et al. DNA

methylation in blood from neonatal screening cards and the association with BMI and

insulin sensitivity in early childhood. International Journal of Obesity. 2018.

doi:10.1038/ijo.2017.228

18.   Yu S, Zhang R, Liu G, Yan Z, Hu H, Yu S, et al. Microarray analysis of differentially

expressed microRNAs in allergic rhinitis. American Journal of Rhinology and Allergy.

2011. doi:10.2500/ajra.2011.25.3682

19.   Suojalehto H, Lindström I, Majuri ML, Mitts C, Karjalainen J, Wolff H, et al. Altered

microRNA expression of nasal mucosa in long-term asthma and allergic rhinitis.

International Archives of Allergy and Immunology. 2014. doi:10.1159/000358486

20.   Sharbati J, Lewin A, Kutz-Lohroff B, Kamal E, Einspanier R, Sharbati S. Integrated

microrna-mrna-analysis of human monocyte derived macrophages upon

mycobacterium avium subsp. hominissuis infection. PLoS ONE. 2011.

doi:10.1371/journal.pone.0020258

21.   Asaoka T, Sotolongo B, Island ER, Tryphonopoulos P, Selvaggi G, Moon J, et al.

MicroRNA signature of intestinal acute cellular rejection in formalin-fixed paraffin-

embedded mucosal biopsies. American Journal of Transplantation. 2012.

doi:10.1111/j.1600-6143.2011.03807.x

22. Lin CH, Lee YS, Huang YY, Tsai CN. Methylation status of vault rna 2-1 promoter is

a predictor of glycemic response to glucagon-like peptide-1 analog therapy in type 2

diabetes mellitus. BMJ Open Diabetes Research and Care. 2021;9.

doi:10.1136/bmjdrc-2020-001416

23. Barker DJP, Osmond C. INFANT MORTALITY, CHILDHOOD NUTRITION, AND

ISCHAEMIC HEART DISEASE IN ENGLAND AND WALES. The Lancet. 1986.

doi:10.1016/S0140-6736(86)91340-1

24. Romanelli V, Nakabayashi K, Vizoso M, Moran S, Iglesias-Platas I, Sugahara N, et al.

Variable maternal methylation overlapping the nc886/vtRNA2-1 locus is locked

between hypermethylated repeats and is frequently altered in cancer. Epigenetics.

2014. doi:10.4161/epi.28323

25. Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, et al. The

Human Genome Browser at UCSC. Genome Research. 2002;12.

doi:10.1101/gr.229102

26. Bou-Nader C, Gordon JM, Henderson FE, Zhang J. The search for a PKR code—

differential regulation of protein kinase R activity by diverse RNA and protein

regulators. RNA. 2019. doi:10.1261/rna.070169.118

27. Ziebarth JD, Bhattacharya A, Cui Y. CTCFBSDB 2.0: A database for CTCF-binding

sites and genome organization. Nucleic Acids Research. 2013;41.

doi:10.1093/nar/gks1165

28.   Rosenbloom KR, Sloan CA, Malladi VS, Dreszer TR, Learned K, Kirkup VM, et al.

409   ENCODE Data in the UCSC Genome Browser: Year 5 update. Nucleic Acids

410   Research. 2013;41. doi:10.1093/nar/gks1172

411   29.   Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, et al. NCBI

412   GEO: Archive for functional genomics data sets - Update. Nucleic Acids Research.

413   2013;41. doi:10.1093/nar/gks1193

414   30.   Guevara EE, Lawler RR, Staes N, White CM, Sherwood CC, Ely JJ, et al. Age-

415   associated epigenetic change in chimpanzees and humans. Philosophical Transactions

416   of the Royal Society B: Biological Sciences. 2020;375. doi:10.1098/rstb.2019.0616

417   31.   Hernando-Herraez I, Prado-Martinez J, Garg P, Fernandez-Callejo M, Heyn H,

418   Hvilsom C, et al. Dynamics of DNA Methylation in Recent Human and Great Ape

419   Evolution. PLoS Genetics. 2013;9. doi:10.1371/journal.pgen.1003763

420   32.   Housman G, Quillen EE, Stone AC. Intraspecific and interspecific investigations of

421   skeletal DNA methylation and femur morphology in primates. American Journal of

422   Physical Anthropology. 2020;173. doi:10.1002/ajpa.24041

423   33.   Boureau L, Constantinof A, Moisiadis VG, Matthews SG, Szyf M. The DNA

424   methylation landscape of enhancers in the Guinea pig hippocampus. Epigenomics.

425   2018;10. doi:10.2217/epi-2017-0064

426   34.   Hannon E, Knox O, Sugden K, Burrage J, Wong CCY, Belsky DW, et al.

427   Characterizing genetic and environmental influences on variable DNA methylation

428   using monozygotic and dizygotic twins. PLoS Genetics. 2018.

429   doi:10.1371/journal.pgen.1007544

35. Horvath S, Mah V, Lu AT, Woo JS, Choi OW, Jasinska AJ, et al. The cerebellum ages

slowly according to the epigenetic clock. Aging. 2015;7. doi:10.18632/aging.100742

36. Horvath S, Langfelder P, Kwak S, Aaronson J, Rosinski J, Vogt TF, et al.

Huntington's disease accelerates epigenetic aging of human brain and disrupts DNA

methylation levels. Aging. 2016;8. doi:10.18632/aging.101005

37. Andrews S. FastQC - A quality control tool for high throughput sequence data.

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/. Babraham Bioinformatics.

2010.

38. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: Summarize analysis results for

multiple tools and samples in a single report. Bioinformatics. 2016;32.

doi:10.1093/bioinformatics/btw354

39. Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina

sequence data. Bioinformatics. 2014;30. doi:10.1093/bioinformatics/btu170

40. Krueger F, Andrews SR. Bismark: A flexible aligner and methylation caller for

Bisulfite-Seq applications. Bioinformatics. 2011;27. doi:10.1093/bioinformatics/btr167

41. Hernandez Mora JR, Tayama C, Sánchez-Delgado M, Monteagudo-Sánchez A, Hata

K, Ogata T, et al. Characterization of parent-of-origin methylation using the Illumina

Infinium MethylationEPIC array platform. Epigenomics. 2018;10. doi:10.2217/epi-

2017-0172

42. Kentepozidou E, Aitken SJ, Feig C, Stefflova K, Ibarra-Soria X, Odom DT, et al.

Clustered CTCF binding is an evolutionary mechanism to maintain topologically

associating domains. Genome Biology. 2020;21. doi:10.1186/s13059-019-1894-x

43.   Dowen JM, Fan ZP, Hnisz D, Ren G, Abraham BJ, Zhang LN, et al. Control of cell

453   identity genes occurs in insulated neighborhoods in mammalian chromosomes. Cell.

454   2014;159. doi:10.1016/j.cell.2014.09.030

Figure legends

Figure 1. Schematic presentation of the similarity of the *nc886* gene and flanking CTCF binding sites. Sequence similarity decreases with the diluting color intensity. Guinea pigs lack the whole centromeric flanking CTCF binding site, whilst the marmosets have a region with sequence similarity, but lack the binding sequence of the CTCF. For detailed sequence comparisons, see Supplementary figures S1 and S2.

Figure 2. Blood DNA methylation beta values in *nc886* region in great apes (*Hominidae*). In each graph, one dot represents one individual. In humans a bimodal methylation pattern can be detected, in line with the expected population distribution of hemi-methylated (75% of the population, methylation level ~0.5) and non-methylated individuals (25% of the population, methylation level close to 0 [10]). All of the other species present a unimodal methylation pattern, with all individual having methylation beta-values near 0.5 across the *nc886* locus.

Figure 3. Femur DNA methylation beta values in the *nc886* region in humans and in monkeys. In humans a bimodal methylation pattern can be detected, similar to blood and in line with expected population distribution of non- and hemi-methylated individuals [10]. In Old World monkeys the between individual variation is bigger than in apes. No individuals presenting a non-methylated *nc886* region are detected. In marmosets only 3 probes were considered to provide reliable methylation values and none of them present methylation levels near 0.5.

Supplementary files

480   Supplementary file 1 (.docx) containing Supplementary Figures S1-S5 with captions.