Long-term labeling and imaging of synaptically-connected neuronal networks *in vivo* using double-deletion-mutant rabies viruses

Lei Jin[1,2], Heather A. Sullivan[1], Mulangma Zhu[1], Thomas K. Lavin[1], Makoto Matsuyama[1], Xin Fu[1,3], Nicholas E. Lea[1], Ran Xu[1], YuanYuan Hou[1], Luca Rutigliani[1], Maxwell Pruner[1], Kelsey R. Babcock[1], Jacque Pak Kan Ip[3,4,5], Ming Hu[3,4,6], Tanya L. Daigle[7], Hongkui Zeng[7], Mriganka Sur[3,4], Guoping Feng[1,3,8] & Ian R. Wickersham[1]*

1. McGovern Institute for Brain Research, Massachusetts Institute of Technology, Cambridge, MA, USA
2. Present address: Lingang Laboratory, Shanghai, China
3. Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, MA, USA
4. Picower Institute for Learning and Memory, Massachusetts Institute of Technology, Cambridge, MA, USA
5. Present address: School of Biomedical Sciences, The Chinese University of Hong Kong, Hong Kong, China
6. Present address: Department of Neuroscience, Baylor College of Medicine, Houston, TX, USA
7. Allen Institute for Brain Science, Seattle, WA, USA
8. Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA.
*Corresponding author (wickersham@mit.edu).

Abstract

Monosynaptic tracing is a widely-used technique for mapping neural circuitry, but its cytotoxicity has confined it primarily to anatomical applications. Here we present a second-generation system for labeling direct inputs to targeted neuronal populations with minimal toxicity, using double-deletion-mutant rabies viruses. Spread of the viruses requires expression of both deleted viral genes *in trans* in postsynaptic source cells; suppressing this expression with doxycycline following an initial period of viral replication reduces toxicity to postsynaptic cells. Longitudinal two-photon imaging *in vivo* indicated that over 90% of both presynaptic and source cells survived for the full twelve-week course of imaging. *Ex vivo* whole-cell recordings at 5 weeks postinfection showed that the second-generation system perturbs input and source cells much less than does the first-generation system. Finally, two-photon calcium imaging of labeled networks of visual cortex neurons showed that their visual response properties appeared normal for 10 weeks, the longest we followed them.

Main

Monosynaptic tracing, or the labeling of neurons in direct synaptic contact with a targeted neuronal population using a deletion-mutant neurotropic virus, has become an important technique in neuroscience since its introduction in 2007[1], because it is the primary available means of brain-wide identification of directly connected neurons. Its core principles are (i) selective infection of a targeted neuronal group of interest with a deletion-mutant neurotropic virus (which in almost all implementations to date is a rabies virus with the envelope glycoprotein gene "G" deleted from its genome) and (ii) complementation of the deletion *in trans* in the targeted starting cells, so that the virus can fully replicate within them and spread, as wild-type rabies virus does, retrogradely (in the central nervous system[2]) across those neurons' input synapses to neurons directly presynaptic to them. This system has

contributed to many significant discoveries about the synaptic organization of many systems in the
mammalian nervous system[3-16].

However, with a few notable exceptions[14-21], monosynaptic tracing has primarily served simply as an
anatomical tool for static identification of connected neurons, because the first-generation, "ΔG" rabies
viral vectors that it is based on are swiftly cytotoxic[22-24]. This toxicity stems from the fact that, while
deletion of the glycoprotein gene, as intended, prevents the virus from spreading between neurons in the
absence of transcomplementation, it leaves the viral transcription and replication machinery intact, so
that the virus still rapidly expresses viral genes at high levels and replicates the viral core to high copy
numbers[25], perturbing endogenous gene expression[26,27], inhibiting host cell protein synthesis[28], and
killing most infected neurons within approximately two weeks[22,23].

Several efforts have been made to engineer less-toxic or nontoxic monosynaptic tracing systems. A first-
generation system based on the CVS-N2c strain of rabies virus appears to have lower toxicity than does
the widely-used SAD B19 strain[24]. More recently, a paper in *Cell* reported that adding a destabilization
domain to the C terminus of the viral nucleoprotein rendered the virus nontoxic, allowing monosynaptic
tracing "with no adverse effects on neural physiology, circuit function, and circuit-dependent
computations"[29]. We have since shown that those results were probably obtained using reversion
mutants that had lost the intended C-terminal addition[30], although we also showed that the technique
may be salvageable[30], and the authors of the original study have persevered with their approach[31]. From
our own laboratory, we have introduced second-generation, "ΔGL" rabies viral vectors, which have the
viral polymerase gene "L" (for "large" protein) deleted along with G, and showed that they do not appear
to perturb the structure or function of labeled neurons[23]. However, no one has previously shown that
these vectors can spread between neurons. Although a baculovirus-based complementation system has
recently been reported, it was not shown to work either *in vivo* or otherwise[32].

Here we show that second-generation (ΔGL) rabies viral vectors can be used for monosynaptic tracing
of inputs to genetically-defined populations of neurons, with the double deletion complemented by
expression of both deleted viral genes *in trans* in the postsynaptic cells.
76

Results

Construction of a monosynaptic tracing system based on ΔGL rabies viruses

We planned for the second-generation monosynaptic tracing system to use the same pseudotyping
strategy[1] used in the first-generation one for targeting the initial RV infection to the postsynaptic starting
cells (or "source cells", if defined as cells expressing the complementing gene(s) and therefore able to
support viral replication). We therefore began by making versions of our previously-introduced second-
generation (ΔGL) rabies viral vectors[23] packaged with the avian retroviral envelope protein EnvA, for
selective infection of cells engineered to express EnvA's receptor, TVA[33,34].

Whereas the first-generation system typically relies on an AAV "helper virus" for expression of the deleted
glycoprotein gene *in trans* in the source cells[35-38], the polymerase gene that is also deleted in the second-
generation vectors is ~6.4 kb, too large[39-42] for a straightforward extension of this approach. We also
wanted expression of both the polymerase and the glycoprotein to be regulatable using the tetracycline
system[43], because we anticipated that sustained expression and resulting viral replication could be toxic
to source cells.

We therefore generated a knock-in mouse line, "TRE-CB" (short for "TRE-tight-mCardinal-P2A-B19L"),
in which the genes for the RV polymerase (SAD B19 strain) and the red fluorophore mCardinal[44] are
under the control of the tetracycline response element in the TIGRE locus[45-47]. To interface with this TRE-
driven polymerase allele, we developed a helper virus combination (now published for use with first-
generation RV[38,48,49]) based on the tet system. Specifically, one AAV, "AAV1-syn-FLEX-splitTVA-EGFP-
tTA" is Cre-dependent[50] (FLEX/DIO design[51]) and expresses TVA, EGFP[52,53], and the tetracycline
transactivator (tTA), while a second AAV, "AAV1-TREtight-mTagBFP2-B19G", expresses the RV
glycoprotein and the blue fluorophore mTagBFP2[54] under the control of the tetracycline response element.
Our intent was that, when the helper viruses were used in the TRE-CB mouse with Cre expression in
starting cells, the tTA expressed by the first AAV would drive expression both of G from the second AAV
and of L from the knock-in allele. Note that the use of tTA instead of rtTA makes this a Tet-Off system,
so that the genes are expressed by default, with optional administration of a tetracycline analog (e.g.,
doxycycline) acting to suppress their expression.

ΔGL viruses, by design, express genes at very low levels and are therefore intended to express
recombinases, which allow many downstream applications even when expressed at low levels[23]. The
viruses in this study therefore encoded either Cre (codon-optimized for mouse[55]) or Flpo (a codon-
optimized version of the yeast recombinase Flp[56,57]). Another necessary component of the system was
therefore a means of reporting Cre or Flpo activity, for which we used the Ai14[58] and Ai65F[47] mouse lines,
in which recombination causes tdTomato expression in cells expressing Cre or Flp, respectively.

Based on our extensive prior experiments titrating these helper viruses for use with first-generation rabies
virus to achieve high transsynaptic spread efficiency in Cre mice but low background without Cre[48,49], for
this second-generation system we used the same helper virus concentrations and seven-day interval
between AAV and RV injections as we have previously described for the first-generation one, only varying
the survival time after RV injection.

Labeling inputs to corticostriatal projection neurons: Cre vs Flpo

We targeted corticostriatal cells in primary somatosensory cortex (S1) using retrograde infection by
rAAV2-retro[59] injected into dorsolateral striatum (Fig. 1), with helper AAVs and subsequently RVΔGL
injected into S1. In one design (Fig. 1a), the rAAV2-retro expressed Cre, the helper AAVs were Cre-
dependent as described above, and the RVΔGL expressed Flpo; these injections were done in the Ai65F
reporter line (Flp-dependent tdTomato) crossed to TRE-CB. In the other design (Fig. 1b), the rAAV2-retro
expressed Flpo, the helper AAVs were Flpo-dependent (i.e., the first helper virus contained orthogonal
FRT sites instead of lox sites), and the RVΔGL expressed Cre instead of Flpo; these injections were done
in the Ai14 reporter line (Cre-dependent tdTomato) crossed to TRE-CB. Titers of AAV and RV vectors
were matched across the two designs (see Methods). In order to test the effect of using the Tet-Off
mechanism to suppress G and L expression after an initial period of unrestricted viral replication, in some
conditions we switched the mice to food containing doxycycline two weeks after rabies virus injection
("dox (food)" conditions), while in others (the "dox (inj+food)" conditions) mice also received doxycycline
by intraperitoneal injection for three days beginning at the same time as dox food was begun. Two to five
weeks after rabies virus injection, mice were perfused and the results examined by confocal microscopy,
with transsynaptic spread efficiency quantified by counting labeled neurons in contralateral cortex and
ipsilateral thalamus.

With the Flpo-expressing RVΔGL, when all components were included, there were labeled neurons in
the input regions at two different survival times (2 and 5 weeks after RV injection), and doxycycline
administration resulted in slightly more spread, though not significantly (Fig. 1c, 1e, and 1g. see
Supplementary File S1 for all cell counts and statistical comparisons). This spread of the ΔGL virus to
input regions, increasing with time, was despite the fact that we were unable to see clear mCardinal
signal, suggesting very low expression of L under the conditions implemented (Extended Data Fig. S1).
First-generation control experiments using RVΔG-Flpo with the 7-day survival time that is typical for the
first-generation system[48,60] labeled significantly more cells in both input regions than any condition using
the second-generation RVΔGL-Flpo (Fig. 1e, and see Extended Data Fig. S2 for images from control
conditions).

The second-generation system using the RVΔGL expressing Cre instead of Flpo resulted in more labeled
neurons in input regions than the one using RVΔGL expressing Flpo (Fig. 1d, 1f, and 1h), although the
comparisons between the Cre and Flpo conditions were underpowered due to high variance and modest
n (=4 mice per condition throughout our study) and did not achieve statistical significance (see
Supplementary File S1 for all cell counts and statistical comparisons).

Control experiments in which either Cre or G was omitted resulted in almost no labeled cells in input
regions in either contralateral cortex or thalamus (Fig. 1e-h & Extended Data Fig. S2. See Supplementary
File S1 for all cell counts and statistical comparisons). This indicates that the apparent transsynaptic
spread was not due to trivial confounding effects, such as direct retrograde infection by the TVA-
expressing helper AAV followed by infection of the resulting TVA-expressing axons by the rabies virus,
or simply direct retrograde infection by residual RV coated with its native glycoprotein. For more
discussion of such possible confounds, see Lavin et al.[48].

Crucially, control experiments in which L was omitted (by using double transgenics without the TRE-CB
allele) also resulted in no label in input regions (Fig. 1e-h & Extended Data Fig. S2). This indicates that
complementation with both G and L is required for the ΔGL virus to replicate within the source cells and
spread to presynaptic ones. See Supplementary File S1 for all counts and statistical comparisons.

Note that although the first-generation RV∆G-Cre at two weeks postinjection had labeled more input
neurons on average than any other condition (**f**), almost no source cells were found at this timepoint with
either of the first-generation combinations (none in six out of eight mice, and two cells each in two mice;
see Extended Data Fig. S3, panels (**a**) and (**c**), and Supplementary File S1), suggesting that the first-
generation system had killed nearly all source cells by that point.

Representative whole-brain image series of labeled inputs to corticostriatal neurons using the Cre-
expressing first- and second-generation rabies viral vectors are shown in Extended Data Fig. S4.
166
167 Mapping inputs to dopaminergic and parvalbuminergic neurons in Cre mouse lines

We tested the system in two widely-used Cre mouse lines: DAT-IRES-Cre[61] and PV-Cre[62], crossed to
TRE-CB and the Flp reporter line Ai65F (Fig. 2 and Extended Data Figs. S5 & S6). In DAT-IRES-Cre, in
which Cre is expressed in dopaminergic cells in the midbrain, we injected the viruses in substantia nigra,
pars compacta (SNc); in PV-Cre, in which Cre is expressed in parvalbumin-expressing cells in cortex and
elsewhere, we injected primary somatosensory cortex (S1).

In both mouse lines, with or without doxycycline, we found spread of rabies virus to regions known to
provide input to the respective populations: in DAT-IRES-Cre, we found labeled neurons in striatum (Fig.
2b) and cortex; in PV-Cre, we found labeled neurons in thalamus (Fig. 2e) and other cortical areas,
among other locations. In both mouse lines, just as with the corticostriatal experiments described above,
there were more neurons at a later time point (five weeks following rabies virus injection) than at an earlier
one (three weeks) (see Supplementary Files S2 and S3 for all cell counts and statistical comparisons).
Because pilot studies had suggested that survival times of one or two weeks did not result in substantial
spread, and that survival times of seven weeks and higher did not result in much more spread than at
five weeks, we only compared three- and five-week survival times for this set of quantitative experiments.

A somewhat surprising result from these experiments was that label in input regions in both mouse lines
and at both survival times was somewhat higher when doxycycline was administered beginning two
weeks after rabies virus injection, although these comparisons were again underpowered and not
statistically significant. This was contrary to our expectation that this intervention, which was designed to
reduce toxicity to source cells by shutting off viral replication after an initial period, could also reduce the
efficiency of transsynaptic spread.

Comparison to the first-generation system in the same Cre lines, however, showed that this
implementation of the second-generation one appeared to be far less efficient: matched experiments
using a first-generation virus, RV∆G-Flpo (and in mice without the TRE-CB allele), with a seven-day
survival time, labeled 1.3-1.5 orders of magnitude (~21x in DAT-IRES-Cre, ~33x in PV-Cre) more
neurons in input regions (Fig. 2c & 2f).

The results of the control experiments with both mouse lines were as expected: omission of any of G, L,
or Cre resulted in almost no labeled cells in input regions (Fig. 2 & Extended Data Figs. S5 & S6; see
Supplementary Files S2 and S3 for all counts and statistical comparisons).
196
197 Mapping inputs to dopaminergic neurons in a Flpo mouse line

Because the Flp-dependent version of the system using RV∆GL-Cre resulted in more labeled input
neurons than the Cre-dependent one using RV∆GL-Flpo in the corticostriatal experiments (Fig. 1), we
tested the Flp-dependent system in DAT-P2A-Flpo mice (crossed to TRE-CB and Ai14), as shown in Fig.
3. We again found the expected patterns of label at the injection site and in the striatum and other input
regions, and controls in which either G or Flpo were omitted resulted in very few (0-5) labeled cells in
striatum (see Supplementary File S4 for counts and statistical comparisons). Without doxycycline, the
numbers of labeled striatal cells were higher with the Flp-dependent system in this mouse line than with
the corresponding Cre-dependent condition in DAT-IRES-Cre mice, although this difference was not
statistically significant (p=0.134); with doxycycline, however, the numbers were lower than in the
corresponding Cre-dependent condition, a difference which was significant. The first-generation system
again labeled more striatal cells than did the second-generation system, although in this case the
difference was much smaller (mostly because the efficiency of this Flp-dependent implementation of the
first generation system was significantly less than that of the Cre-dependent one) and the comparison
was not quite statistically significant (p=0.051). See also Extended Data Fig. S7 for whole-brain image
series for RV∆G-Cre and RV∆GL-Cre, and see Extended Data Fig. S8 for higher-resolution images of

label in several input regions (amygdala, external globus pallidus, and nucleus accumbens) for both ∆G
and ∆GL viruses expressing Flpo and Cre.
215
Longitudinal two-photon imaging of labeled networks

We conducted several experiments to assay the survival and physiological status of neurons labeled with
the second-generation monosynaptic tracing system. In previous work, we used longitudinal two-photon
microscopy to show that neurons labeled with a second-generation (∆GL) RV stayed alive for as long as
we imaged them (in stark contrast to neurons labeled with a first-generation (∆G) RV) and also that it did
not appear to perturb their visual response properties[23]. Because we had only used the Cre-expressing
versions of ∆G and ∆GL viruses for the longitudinal two-photon imaging work from our previous
publications[23,30], for the two-photon work in the present paper we began by conducting a similar set of
experiments using the Flpo-expressing RV∆GL to confirm that it was similarly nontoxic (Extended Data
Fig. S11). We found that it was: in contrast to the first-generation version, RV∆G-Flpo, which killed over
96% of labeled neurons by 4 weeks postinjection, the second-generation RV∆GL-Flpo left nearly all
labeled neurons alive for as long as we imaged them (16 weeks) (specifically, 85 of the 86 neurons seen
at 4 weeks postinjection, when the numbers of visibly ∆GL-Flpo-labeled neurons had stabilized – cf. our
similar findings with ∆GL-Cre in Chatterjee et al.[23] – were still found at 16 weeks in the two mice in which
we were able to image that long. See Supplementary File S5 for counts). This confirmed that the Flpo-
expressing and Cre-expressing ∆GL viruses are similarly nontoxic to labeled cells, consistent with our
expectation that, regardless of the transgene, the large reduction in gene expression level caused by
deletion of L would eliminate the cytotoxicity caused by the high gene expression and intracellular
replication of wild-type and ∆G rabies viruses. These findings suggested that second-generation
monosynaptic tracing should be quite nonperturbative to labeled presynaptic neurons (which are only
infected with the ∆GL viruses and not the helper viruses), except perhaps to the extent that toxicity to the
postsynaptic source cells might perturb cells presynaptic to them by trophic or other network effects.

We then conducted an imaging experiment to assay how toxic the full second-generation monosynaptic
tracing system was to neurons, including the source cells, in which G and L were provided (Fig. 4). Similar
to the experiments presented in Fig. 2d-f, we injected the Cre-dependent helper virus combination into
visual cortex of PV-Cre x TRE-CB x Ai65F mice, followed by RV∆GL-Flpo(EnvA) seven days later. For
this set of experiments, we additionally implanted a headmount and optical window (see Methods), and
we imaged fields of view at or near the injection sites on a two-photon microscope repeatedly over 14
consecutive sessions beginning at two days postinjection and ending at 12 weeks (Fig 4a). Just as for
the previous experiments, half of the mice were given doxycycline beginning at two weeks postinjection.

We found that, with or without doxycycline, the majority of tdTomato-labeled cells at the injection sites
survived for the full 12 weeks of imaging in most mice. For the mice that did not receive doxycycline, and
that therefore had G and L expressing throughout the 12-week experiment, 63.4% of labeled cells
survived through the final imaging session. For the mice that did receive doxycycline, however, 94.4% of
labeled neurons survived (Fig. 4b-h; see also Extended Data Fig. S12; see also Supplementary File S6
for counts of total and surviving cells). Again, despite doxycycline administration, transsynaptic spread to
other cortical areas was still widespread in these mice (Fig. 4k).

Although the overall cell survival rate was high, we anticipated that it would be lower for source neurons,
in which G and L are provided and viral replication is therefore restored. In order to determine the survival
rate of source cells, the ideal approach would have been to image both tdTomato (reporting activity of
the Flpo expressed by the rabies virus) and mTagBFP2 (coexpressed with G by the second helper virus):
neurons that expressed both fluorophores would be both infected by the RV∆GL and expressing G (and
presumably L, because both depend on tTA expression) and therefore competent hosts for the RV to
replicate within and spread to cells presynaptic to them. However, because the two-photon microscope
to which we had access was unfortunately incapable of producing short-wavelength light at sufficient
power to allow imaging of mTagBFP2 *in vivo*, we had to rely on indirect measures to estimate the source
cells' survival rates. Note that EGFP, which was coexpressed with TVA, is not a good indication of
whether a cell is a source cell, because, although some of the cells expressing TVA would have been
directly infected by the EnvA-enveloped rabies virus, only a subset of such cells also express G (as
reported by mTagBFP2: see Supplementary File S3) and L and are therefore able to support replication
and spread of the rabies virus; in any case, at the helper virus concentrations used[48,49], the intrinsic EGFP
signal is quite weak and needs to be amplified with immunostaining in order to be clearly discernible.

Following longitudinal two-photon imaging of tdTomato-labeled cells over the full 12 weeks, therefore, we
perfused the mice, sectioned the brains in a plane approximately parallel to the imaging plane, then
imaged the same regions using confocal microscopy. This allowed us to image tdTomato as well as
mTagBFP2 (for animals that had not received doxycycline), then align the confocal images with the two-
photon ones from the 12-week timepoint to identify the surviving double-labeled source cells (Fig. 4i-j).
For those animals that did receive doxycycline, the blue channel contained little signal, as expected (Fig.
4j); our second goal for the confocal imaging, however, was to confirm that transsynaptic spread in these
mice used for two-photon imaging had occurred, which it had (Fig. 4k). For both dox and no-dox
conditions, we made the assumption that all cells that had died were source cells, based on our finding
that cells infected solely with RVΔGL-Flpo without complementation almost always survive (Extended
Data Fig. S11; see also Chatterjee et al.[23] for extensive characterization of the nontoxicity of ΔGL RV).

In the mice that did not receive doxycycline, in which the surviving source cells (i.e., those expressing
red and blue fluorophores) were visible by confocal microscopy (Fig. 4i), we estimated the source cell
survival rate in each mouse as the ratio of the number of cells in the field of view expressing both red and
blue fluorophores divided by the sum of that number plus the number of cells that died in that field of view
(Fig. 4l & 4n; see Supplementary File S6 for cell counts and calculations). For these no-doxycycline mice,
we obtained a survival rate of 53.2% of source cells.

In the mice that did receive doxycycline, in which surviving source cells were not identifiable as such by
postmortem confocal microscopy because they no longer expressed mTagBFP2 (Fig. 4j), we made the
assumption that source cells made up the same percentage of total tdTomato-labeled neurons in the
fields of view in the dox mice as in the no-dox ones, because the experimental conditions were identical
apart from the switch to doxycycline two weeks after RV injection. This gave us an estimated number of
total source cells for each mouse; the difference between that and the (small) number of neurons that
had disappeared over the course of the two-photon imaging gave us an estimated number of surviving
source cells. The ratio of the estimated number of surviving source cells to the estimated total number of
source cells was our estimate of the source cell survival rate (Fig. 4m & 4n; see Supplementary File S6
for cell counts and calculations). For these doxycycline mice, we obtained an estimated survival rate of
92.3% of source cells.
296
Ex vivo whole-cell recordings from labeled input and source neurons

In order to further assay the degree to which the second-generation monosynaptic tracing system
perturbs input and source neurons, we performed whole-cell recordings on neurons in acute brain slices
made from mice in which inputs to parvalbumin-expressing cortical neurons had been labeled (Fig. 5).
As with the anatomical experiments described previously (Fig. 2), we injected Cre-dependent helper
viruses in primary sensory cortex of PV-Cre x TRE-CB x Ai65F mice, followed by either RVΔG-Cre(EnvA)
or RVΔGL-Cre(EnvA) (or no rabies virus, for AAV-only controls) seven days later. Beginning 35 days
after rabies virus injection, we made slices from the injected brains and conducted whole-cell recordings
from labeled input cells and source cells to measure their membrane properties.

In brains labeled by the second-generation system, we found that the membrane properties of input cells
did not differ from those of unlabeled neurons in the AAV-only control mice, and that source cells differed
from "source cells" in the AAV-only control mice (i.e., cells labeled by the helper AAVs) in two of the five
measured membrane properties.

By contrast, in brains labeled by the first-generation system, those input cells that remained at the time
of recording (5 weeks after infection) differed from unlabeled control neurons in three out of five measured
membrane properties (cf. Chatterjee et al.[23]), and no remaining source cells could be found at all. This
marked cytotoxicity of the first-generation system to source cells after 5 weeks is unsurprising but
illustrates a major difference between the first- and second-generation systems.
315
Longitudinal two-photon calcium imaging of cortical neurons' responses to visual stimuli

Finally, we conducted experiments designed to provide proof of concept for long-term functional imaging
of synaptically-connected networks of neurons (Fig. 6). We began by making a Flp-dependent jGCaMP7s
reporter AAV and conducted pilot studies in PV-Cre mice to determine a dilution that would not itself
cause mass cytotoxicity due to overexpression of jGCaMP7s[47,63] (Extended Data Fig. S13). We then
conducted an experiment similar in design to the structural imaging study, except that the rabies virus
was mixed with the jGCaMP7s AAV before injection and headmount/coverslip implantation (Fig. 6a). We

then imaged labeled neurons at or near the injection sites over a series of sessions spanning ten weeks.
We imaged the jGCaMP7s signal while the awake mice viewed a series of drifting grating stimuli (Fig.
6b-c); we also imaged tdTomato fluorescence in the same fields of view to confirm that jGCaMP7s
expression was restricted to RV-labeled neurons (Fig. 6c). As shown in Fig. 6d-h, as well as Extended
Data Figs. S13 and S15, we observed clear orientation tuning curves in many of the labeled neurons,
with visual responses appearing to remain stable across repeated imaging sessions over several months
(Fig 6d-h, and Extended Data Fig. S15). Following the full course of two-photon imaging, we conducted
postmortem sectioning and confocal imaging to confirm that viral spread had occurred in the imaged mice
(Extended Data Fig. S16).
332

333   Discussion

We have shown here that it is possible to use nontoxic double-deletion-mutant rabies viral vectors to
label direct inputs to neurons targeted based on either their extrinsic projections or their gene expression
(using Cre or Flp lines). We have shown that use of the Tet-Off system results in relatively little toxicity
even to the source cells. Using whole-cell recordings from labeled neurons, we found that input cells'
membrane properties were not significantly different from those of unlabeled control neurons, and source
cells were perturbed in two out of four measured membrane properties; this represents a considerable
improvement over the first-generation system, which, at the five-week survival time we examined, left
input cells perturbed and source cells dead. We have further used two-photon functional imaging of live
second-generation monosynaptic tracing *in vivo* to show that apparently normal visual response
properties of labeled neurons can be recorded for at least ten weeks.

It is perhaps surprising that so many source cells (over 90%) appeared to survive when doxycycline was
administered beginning at two weeks postinjection (Fig. 4). This was not entirely accidental: we designed
the system so that the level and time course of L expression could be titrated as necessary to reduce
toxicity to source cells. However, it was undoubtedly fortunate that administering doxycycline beginning
two weeks after ΔGL rabies virus injection turned out to be sufficient to preserve the majority of source
cells (at least for cortical cells in the PV-Cre mice that we tested this in). It was even more fortunate that
this intervention did not appear to reduce the efficiency of transsynaptic spread in either of the Cre lines
or in the corticostriatal experiments, although these comparisons were underpowered and not statistically
significant.

As we have mentioned throughout, many of the comparisons of the efficiency of the various *in vivo*
conditions in our study were underpowered, given to the high variance we found in most conditions and
the modest number of mice we used per condition (n=4) because of the large numbers of conditions that
we tested. For example, for the corticostriatal experiments shown in Fig. 1, comparing the ΔGL-Cre and
ΔGL-Flpo at 5 weeks without dox, the mean numbers of labeled cells in contralateral cortex were 36.5
and 20.5, respectively, with a p-value of 0.20 (Supplementary File S1; numbers are totals across every
sixth section; see Methods). Given the pooled standard deviation of 22.38, the number of mice needed
to adequately power this comparison would have been 31 in each group. Similarly, for the labeled
thalamic neurons in the same groups, we found means of 45 with ΔGL-Cre and 16 with ΔGL-Flpo, with a
p-value of 0.14. The pooled standard deviation for these counts was 33.86, indicating that 22 mice per
group would have been required for adequate power. We speculate that the high variability seen in many
of the conditions may be in part due to our injecting the helper viruses and rabies virus in two successive
surgeries, as is typical (and in the case of the corticostriatal experiments, a third injection – AAV2-retro
in striatum – was made). It may be that simply mixing the helper AAVs and rabies virus and injecting
them together (as some have successfully done[64]) would decrease variability, but we have not tested this
approach so far.

A drawback of the second-generation system introduced here is its slow time course, necessitating multi-
week incubation times (Figs. 1-3). We presume that this slow time course is due to low L expression from
the TRE-CB allele given the low level of tTA expressed by the intentionally-dilute Cre- or Flp-dependent
helper virus[48,49]; this hypothesis is consistent with our inability to detect clear mCardinal signal by confocal
microscopy (Extended Data Fig. S1) and perhaps also explains the high percentage of surviving source
cells that we found. While a slow time course is inconvenient, it may be the necessary price of preserving
the source cells.

An interesting finding from the current study was that using an RVΔGL expressing Cre instead of one
expressing Flpo resulted in increased numbers of labeled cells under some conditions (Figs. 1-3),

although again these comparisons were not statistically significant. Our working hypothesis is that such a difference, if real, could be due to a difference in inefficiency of the two recombinases[57,65,66], with Cre more faithfully reporting the actual spread of the virus than Flpo when the recombinases are expressed at low levels (e.g., by ΔGL rabies viruses). A relative inefficiency of Flpo could also, however, cause lower efficiency of the Flp-dependent helper virus combination itself relative to the Cre-dependent one; this is consistent with the first-generation system's lower efficiency in the Flp-dependent corticostriatal experiments than in the Cre-dependent ones (Fig. 1), and similarly in DAT-P2A-Flpo mice (Fig. 3) than in DAT-IRES-Cre mice (Fig. 2). In this study we simply matched the titers of the Flp-dependent combination to those used for the Cre-dependent one, but titration and further optimization of the Flp-dependent helper virus might improve performance.

We note that, even apart from the rabies virus and its complementing gene products, several of the components of the system have the potential to cause cytotoxicity, including tTA[47,67,68], GCaMPs[47,63], and even AAV itself[69]. These can be expected to contribute to the toxicity to source cells, perhaps making it more remarkable that we found that so many of them survived, albeit with some perturbation to their membrane properties (Fig. 5). The toxicity to source cells may also depend on the cell type, both because neuron types can be differentially vulnerable to adverse conditions[70,71] and because differential tropism for AAV1[49] (or whatever other serotype might be used for the helper viruses) will change the dosage of tTA, G, L, TVA (and therefore the initial multiplicity of infection by the injected RV), and the fluorophores.

The presynaptic cells, on the other hand, are less burdened than the postsynaptic ones: because we have found that ΔGL RV by itself (in combination with tdTomato expression in Ai14 or Ai65F mice) does not kill neurons, the vast majority of labeled presynaptic cells can be expected to survive. It seems possible that residual toxicity to the source cells postsynaptic to them could cause some remodeling or toxicity via retrograde transneuronal effects[72]; however, our electrophysiological recordings did not detect any perturbation of the input cells' membrane properties.

While the second-generation system is more complicated than the first-generation one in that it requires two additional transgenic alleles (for expressing L and for reporting activity of the recombinase expressed by the RV), in practice this is not overly burdensome: we have found that the TRE-CB mice can be readily maintained as double homozygotes with the Ai14 or Ai65F alleles, so that production of triply (Figs. 2-6) or doubly (Fig. 1) heterozygous mice is achieved by a single cross with either a recombinase-expressing (e.g., PV-Cre) or wild-type mouse.

Perhaps importantly, our finding that ΔGL rabies viruses do not spread when not complemented by L expression *in trans*, even when G is provided *in trans*, indicates that deletion of either L or G alone is sufficient to prevent transsynaptic spread. This suggests the possibility of a third-generation monosynaptic tracing system, based on single-deletion-mutant ΔL viruses[73] complemented by expression *in trans* of just L. Because of the reduced complexity of complementing with (once again) only a single gene, such a system may have advantages over the second-generation one described here.

References

1. Wickersham, I.R. *et al.* Monosynaptic restriction of transsynaptic tracing from single, genetically targeted neurons. *Neuron* **53**, 639-47 (2007).
2. Zampieri, N., Jessell, T.M. & Murray, A.J. Mapping sensory circuits by anterograde transsynaptic transfer of recombinant rabies virus. *Neuron* **81**, 766-78 (2014).
3. Miyamichi, K. *et al.* Cortical representations of olfactory input by trans-synaptic tracing. *Nature* **472**, 191-6 (2011).
4. Schwarz, L.A. *et al.* Viral-genetic tracing of the input-output organization of a central noradrenaline circuit. *Nature* (2015).
5. Stephenson-Jones, M. *et al.* A basal ganglia circuit for evaluating action outcomes. *Nature* **539**, 289-293 (2016).
6. Beier, K.T. *et al.* Rabies screen reveals GPe control of cocaine-triggered plasticity. *Nature* **549**, 345-350 (2017).
7. Szonyi, A. *et al.* Median raphe controls acquisition of negative experience in the mouse. *Science* **366** (2019).
8. Takahashi, T.M. *et al.* A discrete neuronal circuit induces a hibernation-like state in rodents. *Nature* **583**, 109-114 (2020).
9. Rossi, L.F., Harris, K.D. & Carandini, M. Spatial connectivity matches direction selectivity in visual cortex. *Nature* **588**, 648-652 (2020).

10. Jin, H., Fishman, Z.H., Ye, M., Wang, L. & Zuker, C.S. Top-Down Control of Sweet and Bitter Taste in the
Mammalian Brain. *Cell* **184**, 257-271 e16 (2021).
11. Yao, S. *et al.* A whole-brain monosynaptic input connectome to neuron classes in mouse visual cortex.
bioRxiv, 2021.09.29.459010 (2021).
12. Schmidt, E.R.E. *et al.* A human-specific modifier of cortical connectivity and circuit function. *Nature*
(2021).
13. Wu, X., Morishita, W., Beier, K.T., Heifets, B.D. & Malenka, R.C. 5-HT modulation of a medial septal
circuit tunes social memory stability. *Nature* **599**, 96-101 (2021).
14. Yonehara, K. *et al.* Spatially asymmetric reorganization of inhibition establishes a motion-sensitive circuit.
Nature **469**, 407-10 (2011).
15. Wertz, A. *et al.* PRESYNAPTIC NETWORKS. Single-cell-initiated monosynaptic tracing reveals layer-
specific cortical network modules. *Science* **349**, 70-4 (2015).
16. Geiller, T. *et al.* Local circuit amplification of spatial selectivity in the hippocampus. *Nature* (2021).
17. Rancz, E.A. *et al.* Transfection via whole-cell recording in vivo: bridging single-cell physiology, genetics
and connectomics. *Nature neuroscience* (2011).
18. Velez-Fort, M. *et al.* A Circuit for Integration of Head- and Visual-Motion Signals in Layer 6 of Mouse
Primary Visual Cortex. *Neuron* **98**, 179-191 e6 (2018).
19. Yonehara, K. *et al.* The first stage of cardinal direction selectivity is localized to the dendrites of retinal
ganglion cells. *Neuron* **79**, 1078-85 (2013).
20. Rompani, S.B. *et al.* Different Modes of Visual Integration in the Lateral Geniculate Nucleus Revealed by
Single-Cell-Initiated Transsynaptic Tracing. *Neuron* **93**, 1519 (2017).
21. Schwarz, M.K. & Remy, S. Rabies virus-mediated connectivity tracing from single neurons. *J Neurosci*
Methods **325**, 108365 (2019).
22. Wickersham, I.R., Finke, S., Conzelmann, K.K. & Callaway, E.M. Retrograde neuronal tracing with a
deletion-mutant rabies virus. *Nature Methods* **4**, 47-9 (2007).
23. Chatterjee, S. *et al.* Nontoxic, double-deletion-mutant rabies viral vectors for retrograde targeting of
projection neurons. *Nat Neurosci* **21**, 638-646 (2018).
24. Reardon, T.R. *et al.* Rabies Virus CVS-N2c(DeltaG) Strain Enhances Retrograde Synaptic Transfer and
Neuronal Viability. *Neuron* **89**, 711-24 (2016).
25. Albertini, A.A., Ruigrok, R.W. & Blondel, D. Rabies virus transcription and replication. *Advances in virus*
research **79**, 1-22 (2011).
26. Prosniak, M., Hooper, D.C., Dietzschold, B. & Koprowski, H. Effect of rabies virus infection on gene
expression in mouse brain. *Proc Natl Acad Sci U S A* **98**, 2758-63 (2001).
27. Huang, K.W. & Sabatini, B.L. Single-Cell Analysis of Neuroinflammatory Responses Following Intracranial
Injection of G-Deleted Rabies Viruses. *Front Cell Neurosci* **14**, 65 (2020).
28. Komarova, A.V. *et al.* Rabies virus matrix protein interplay with eIF3, new insights into rabies virus
pathogenesis. *Nucleic Acids Res* **35**, 1522-32 (2007).
29. Ciabatti, E., Gonzalez-Rueda, A., Mariotti, L., Morgese, F. & Tripodi, M. Life-Long Genetic and Functional
Access to Neural Circuits Using Self-Inactivating Rabies Virus. *Cell* **170**, 382-392 e14 (2017).
30. Jin, L. *et al.* "Self-inactivating" rabies viruses are susceptible to loss of their intended attenuating
modification. *Proceedings of the National Academy of Sciences* **120**, e2023481120 (2023).
31. Lee, H. *et al.* Combining long-term circuit mapping and network transcriptomics with SiR-N2c. *Nat*
Methods (2023).
32. Zhang, S. *et al.* Light-Guided Rabies Virus Tracing for Neural Circuit Analysis. *bioRxiv*,
2023.03.04.531104 (2023).
33. Young, J.A., Bates, P. & Varmus, H.E. Isolation of a chicken gene that confers susceptibility to infection
by subgroup A avian leukosis and sarcoma viruses. *J Virol* **67**, 1811-6 (1993).
34. Bates, P., Young, J.A. & Varmus, H.E. A receptor for subgroup A Rous sarcoma virus is related to the low
density lipoprotein receptor. *Cell* **74**, 1043-51 (1993).
35. Wall, N.R., Wickersham, I.R., Cetin, A., De La Parra, M. & Callaway, E.M. Monosynaptic circuit tracing in
vivo through Cre-dependent targeting and complementation of modified rabies virus. *Proceedings of the*
National Academy of Sciences of the United States of America (2010).
36. Watabe-Uchida, M., Zhu, L., Ogawa, S.K., Vamanrao, A. & Uchida, N. Whole-brain mapping of direct
inputs to midbrain dopamine neurons. *Neuron* **74**, 858-73 (2012).
37. Kohara, K. *et al.* Cell type-specific genetic and optogenetic tools reveal hippocampal CA2 circuits. *Nat*
Neurosci **17**, 269-79 (2014).
38. Liu, K. *et al.* Lhx6-positive GABA-releasing neurons of the zona incerta promote sleep. *Nature* **548**, 582-
587 (2017).
39. Lai, Y., Yue, Y. & Duan, D. Evidence for the failure of adeno-associated virus serotype 5 to package a
viral genome > or = 8.2 kb. *Mol Ther* **18**, 75-9 (2010).
40. Wu, Z., Yang, H. & Colosi, P. Effect of genome size on AAV vector packaging. *Molecular therapy : the*
journal of the American Society of Gene Therapy **18**, 80-6 (2010).

41.   Dong, B., Nakai, H. & Xiao, W. Characterization of genome integrity for oversized recombinant AAV
497   vector. *Molecular therapy : the journal of the American Society of Gene Therapy* **18**, 87-92 (2010).

498   42.   Hirsch, M.L., Wolf, S.J. & Samulski, R.J. Delivering Transgenic DNA Exceeding the Carrying Capacity of
499   AAV Vectors. *Methods Mol Biol* **1382**, 21-39 (2016).

500   43.   Gossen, M. & Bujard, H. Tight control of gene expression in mammalian cells by tetracycline-responsive
501   promoters. *Proc Natl Acad Sci U S A* **89**, 5547-51 (1992).

502   44.   Chu, J. *et al.* Non-invasive intravital imaging of cellular differentiation with a bright red-excitable
503   fluorescent protein. *Nat Methods* **11**, 572-8 (2014).

504   45.   Zeng, H. *et al.* An inducible and reversible mouse genetic rescue system. *PLoS genetics* **4**, e1000069
505   (2008).

506   46.   Madisen, L. *et al.* Transgenic mice for intersectional targeting of neural sensors and effectors with high
507   specificity and performance. *Neuron* **85**, 942-58 (2015).

508   47.   Daigle, T.L. *et al.* A Suite of Transgenic Driver and Reporter Mouse Lines with Enhanced Brain-Cell-Type
509   Targeting and Functionality. *Cell* **174**, 465-480 e22 (2018).

510   48.   Lavin, T.K., Jin, L. & Wickersham, I.R. Monosynaptic tracing: a step-by-step protocol. *J Chem Neuroanat*,
511   101661 (2019).

512   49.   Lavin, T.K., Jin, L., Lea, N.E. & Wickersham, I.R. Monosynaptic Tracing Success Depends Critically on
513   Helper Virus Concentrations. *Front Synaptic Neurosci* **12**, 6 (2020).

514   50.   Sternberg, N. & Hamilton, D. Bacteriophage P1 site-specific recombination. I. Recombination between
515   loxP sites. *J Mol Biol* **150**, 467-86 (1981).

516   51.   Atasoy, D., Aponte, Y., Su, H.H. & Sternson, S.M. A FLEX switch targets Channelrhodopsin-2 to multiple
517   cell types for imaging and long-range circuit mapping. *The Journal of neuroscience : the official journal of*
518   *the Society for Neuroscience* **28**, 7025-30 (2008).

519   52.   Cormack, B.P., Valdivia, R.H. & Falkow, S. FACS-optimized mutants of the green fluorescent protein
520   (GFP). *Gene* **173**, 33-8 (1996).

521   53.   Thastrup, O., Tullin, S., Poulsen, L.K. & Bjorn, S.P. Fluorescent proteins. (2001).

522   54.   Subach, O.M., Cranfill, P.J., Davidson, M.W. & Verkhusha, V.V. An enhanced monomeric blue
523   fluorescent protein with the high chemical stability of the chromophore. *PloS one* **6**, e28674 (2011).

524   55.   Koresawa, Y. *et al.* Synthesis of a new Cre recombinase gene based on optimal codon usage for
525   mammalian systems. *Journal of biochemistry* **127**, 367-72 (2000).

526   56.   Sadowski, P.D. The Flp recombinase of the 2-microns plasmid of Saccharomyces cerevisiae. *Prog*
527   *Nucleic Acid Res Mol Biol* **51**, 53-91 (1995).

528   57.   Raymond, C.S. & Soriano, P. High-efficiency FLP and PhiC31 site-specific recombination in mammalian
529   cells. *PloS one* **2**, e162 (2007).

530   58.   Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system for the whole
531   mouse brain. *Nature neuroscience* **13**, 133-40 (2010).

532   59.   Tervo, D.G. *et al.* A Designer AAV Variant Permits Efficient Retrograde Access to Projection Neurons.
533   *Neuron* **92**, 372-382 (2016).

534   60.   Zhang, Y. *et al.* Targeting thalamic circuits rescues motor and mood deficits in PD mice. *Nature* **607**, 321-
535   329 (2022).

536   61.   Backman, C.M. *et al.* Characterization of a mouse strain expressing Cre recombinase from the 3'
537   untranslated region of the dopamine transporter locus. *Genesis* **44**, 383-90 (2006).

538   62.   Hippenmeyer, S. *et al.* A developmental switch in the response of DRG neurons to ETS transcription
539   factor signaling. *PLoS Biol* **3**, e159 (2005).

540   63.   Steinmetz, N.A. *et al.* Aberrant Cortical Activity in Multiple GCaMP6-Expressing Transgenic Mouse Lines.
541   *eNeuro* **4**(2017).

542   64.   Pouchelon, G. *et al.* The organization and development of cortical interneuron presynaptic circuits are
543   area specific. *Cell Rep* **37**, 109993 (2021).

544   65.   Kranz, A. *et al.* An improved Flp deleter mouse in C57Bl/6 based on Flpo recombinase. *Genesis* **48**, 512-
545   20 (2010).

546   66.   Ringrose, L. *et al.* Comparative kinetic analysis of FLP and cre recombinases: mathematical models for
547   DNA binding and recombination. *J Mol Biol* **284**, 363-84 (1998).

548   67.   Han, H.J. *et al.* Strain background influences neurotoxicity and behavioral abnormalities in mice
549   expressing the tetracycline transactivator. *J Neurosci* **32**, 10574-86 (2012).

550   68.   Hoyng, S.A. *et al.* Developing a potentially immunologically inert tetracycline-regulatable viral vector for
551   gene therapy in the peripheral nerve. *Gene Ther* **21**, 549-57 (2014).

552   69.   Xiong, W. *et al.* AAV cis-regulatory sequences are correlated with ocular toxicity. *Proc Natl Acad Sci U S*
553   *A* **116**, 5785-5794 (2019).

554   70.   Hedreen, J.C. & Folstein, S.E. Early loss of neostriatal striosome neurons in Huntington's disease. *J*
555   *Neuropathol Exp Neurol* **54**, 105-20 (1995).

556   71.   Burke, R.E. & Baimbridge, K.G. Relative loss of the striatal striosome compartment, defined by calbindin-
557   D28k immunostaining, following developmental hypoxic-ischemic injury. *Neuroscience* **56**, 305-15 (1993).

558   72.   Fricker, M., Tolkovsky, A.M., Borutaite, V., Coleman, M. & Brown, G.C. Neuronal Cell Death. *Physiol Rev*
559   **98**, 813-880 (2018).
560   73.   Jin, L. *et al.* Single-deletion-mutant, third-generation rabies viral vectors allow nontoxic retrograde
561   targeting of projection neurons with greatly increased efficiency. *bioRxiv*, 2022.02.23.481706 (2022).
562

Fig. 1: Second-generation monosynaptic tracing of inputs to corticostriatal neurons. a-b, Corticostriatal neurons were retrogradely targeted by an AAV2-retro expressing either Cre or Flpo injected in dorsolateral striatum, with a Cre- or Flp-dependent helper virus combination injected in S1. EnvA-enveloped ΔGL rabies virus (RV) expressing either Flpo or Cre was injected in S1 seven days later. The mice were crosses of the "TRE-CB" line described here (providing RV polymerase *in trans*) with tdTomato reporter lines to report activity of the rabies viruses (either Ai65F for Flp or Ai14 for Cre). For the "dox" conditions, mice were switched to doxycycline-containing food 14 days after rabies virus injection, to suppress expression of the viral polymerase and glycoprotein after viral spread; the "dox (inj+food)" mice also received intraperitoneal doxycycline injections for three days beginning 14 days after rabies virus injection. Brain images adapted from the Allen Mouse Brain Reference Atlas (atlas.brain-map.org). **c,** Images of cells labeled with the Flpo-expressing RV. Whereas at 2 weeks few cells were found in input regions, by 5 weeks substantial numbers of labeled cells were found in ipsilateral secondary motor and somatosensory cortices, in contralateral S1, and ipsilateral thalamus, with or without doxycycline (note that doxycycline suppresses expression of mTagBFP2 as well as the viral genes G and L). Considerably fewer input cells were labeled with this version of the second-generation system than with the corresponding first-generation one (panel **e** and Extended Data Fig. S2). Scale bars: 200 μm, apply to all images. **d,** With the Cre-expressing RV and Flp-dependent helper viruses, labeled cells were found in the same regions but in greater numbers. **e-f,** Counts of labeled cells in contralateral S1 and ipsilateral thalamus for RVΔGL-Flpo (**e**) and RVΔGL-Cre (**f**). Counts are the total number of cells found in a series of every sixth 50-μm section spanning each brain, so that the true number of labeled neurons in the entirety of each brain would be approximately six times the number shown here. Note that, although the first-generation Flp-dependent system at two weeks had on average labeled more neurons in input regions than any other condition (**f**), almost no source cells were found in the first-generation conditions at two weeks (see Extended Data Fig. S3), suggesting that almost all source cells had died by that point, consistent with our previous findings regarding the toxicity of first-generation rabies viral vectors. **g-h,** Same data as in **e-f** but with the first-generation counts omitted for clearer display of the ΔGL data and with the ranges of the y-axes of the Flpo and Cre versions set to the same value for each input region for easier comparison of the Flpo vs Cre results. In all experiments with the second-generation system, spread of RVΔGL was greater at five weeks than three weeks, and omission of L, G, or Cre/Flpo resulted in trivial numbers of labeled neurons in input regions (see Extended Data Fig. S2 for counts and representative images for these control conditions). The Flpo-expressing ΔGL RV labeled far fewer cells than did the first-generation version; however, the Cre-expressing ΔGL RV labeled more cells than the Flpo-expressing version in all experimental conditions, although these differences were not statistically significant and were underpowered due to the high variance and low n (=4 mice) for each condition (see Supplementary File S1 for counts and statistical comparisons; see also Extended Data Fig. S4 for example whole-brain image series of mice labeled with each virus combination).

602
603
Fig. 2: Second-generation monosynaptic tracing in Cre mice. a, Experimental design for labeling
inputs to dopaminergic cells in the substantia nigra, pars compacta (SNc). The Cre-dependent helper
virus combination was injected into the SNc of triple transgenic mice (DAT-IRES-Cre x TRE-CB x Ai65F);
EnvA-enveloped ΔGL rabies virus expressing Flpo was injected seven days later. Brain images adapted
from the Allen Mouse Brain Reference Atlas (atlas.brain-map.org). **b,** Example of labeled cells five weeks
after rabies virus injection, without doxycycline. **b.i-v,** Injection site in SNc, with many cells co-expressing
tyrosine hydroxylase (indicating the dopaminergic cells), EGFP (expressed by the first helper virus),
mTagBFP2 (expressed by the second helper virus), and tdTomato (reporting activity of the Flpo-
expressing rabies virus). Scale bar in **iii**: 100 μm, applies to all images. **b.vi,** Medium spiny neurons in
striatum labeled by the ΔGL rabies virus. **c,** Counts of labeled striatal cells (total found in a series of every
sixth 50-μm section spanning each brain) in all conditions tested. Omitting any of L, G, or Cre resulted in
almost no labeled cells in striatum. Administration of doxycycline to suppress G and L expression after
the first two weeks, as well as longer survival time (5 weeks vs 3 weeks), both appeared to result in
increased transsynaptic spread, although most comparisons were not statistically significant (see
Supplementary File S2 for all counts and statistical comparisons). The first-generation system (with a
much shorter survival time) labeled many more cells than all conditions tested for this implementation of
the second-generation one. **d,** Experimental design for labeling inputs to parvalbumin-expressing cells in
primary somatosensory cortex (S1) in PV-Cre x TRE-CB x Ai65F mice. **e,** Example of labeled cells five
weeks after rabies virus injection, without doxycycline. **e.i-v,** Injection site in S1, with many cells
coexpressing parvalbumin, reporters of both helper viruses, and tdTomato reporting activity of the rabies

virus. In addition to these source cells, many putatively presynaptic cells expressing tdTomato are
present. Scale bar in **iii**: 100 µm, applies to all images. **e.vi,** Relay neurons in ipsilateral thalamus labeled
by the ΔGL rabies virus. **f,** Counts of labeled thalamic cells (total found in a series of every sixth 50-µm
section spanning each brain) in all conditions tested. Again, omitting any of L, G, or Cre resulted in little
or no transsynaptic spread, provision of doxycyline appeared to increase transsynaptic spread somewhat,
and longer survival time resulted in more spread, although comparisons were underpowered and not
statistically significant (see Supplementary File S3 for all statistical comparisons and p-values). The first-
generation system again labeled many more cells than did this implementation of the second-generation
system, potentially due to low efficiency of recombination by Flpo in these reporter mice.
633

634
635
Fig. 3: Second-generation monosynaptic tracing in DAT-P2A-Flpo mice. a, The design of the
experiment was similar to that shown in Fig. **2a** but using the Flp-dependent helper virus combination,
and Cre-expressing rabies virus, in DAT-P2A-Flpo x TRE-CB x Ai14 mice. Brain image adapted from the
Allen Mouse Brain Reference Atlas (atlas.brain-map.org). **b,** Counts of labeled striatal neurons (total
found in a series of every sixth 50-μm section spanning each brain) in all conditions tested. **c,** Example
images of labeled cells five weeks after rabies virus injection, without doxycycline. Many cells at the
injection site in SNc were found to co-express tyrosine hydroxylase and the fluorophores expressed by
the helper viruses, as well as tdTomato, reporting activity of the Cre-expressing rabies virus. Medium
spiny neurons in striatum were labeled by the ΔGL rabies virus. **d,** Example of a doxycycline (injection +
food) mouse; numbers of striatal input neurons were lower in this condition than without doxycycline,
although comparisons were again underpowered and not statistically significant. **e,** Example label from
the first-generation system. More cells were labeled on average than in the second-generation conditions,
although again the differences were not statistically significant. **f-g,** Example of control conditions in which
either G (**f**) or Flpo (**g**) was omitted. Scale bars: 100 μm, apply to all images.

650
651
Fig. 4: Longitudinal structural two-photon imaging *in vivo*: second-generation monosynaptic
tracing preserves >90% of postsynaptic source cells. a, Schematic of virus injection and longitudinal
two-photon imaging. The helper AAVs, followed by RVΔGL-Flpo(EnvA), were injected in primary visual
cortex of PV-Cre x TRE-CB x Ai65F mice, then the injection site was imaged on a two-photon microscope
repeatedly for 12 weeks after RV injection. Brain image adapted from the Allen Mouse Brain Reference
Atlas (atlas.brain-map.org). **b-c,** Representative two-photon images at different timepoints of tdTomato-
labelled neurons in the injection site in V1. The 'No dox' group (b) was fed with regular food throughout,
whereas the 'Dox' group (c) was fed with doxycycline food (200 mg/kg) starting at two weeks after RV
injection until perfusion at week 12, and also received intraperitoneal injections of 100 mg/kg doxycycline
every 12 hours for three days, beginning two weeks after RV injection, in order to suppress rabies viral
polymerase and glycoprotein expression. Arrows indicate example cells that are alive up to one timepoint
(white arrows) but that are missing subsequently (green arrows). In this example, a total of 46 out of 215
cells were lost in the 'No dox' group, whereas 12 out of 305 cells were lost in the 'Dox' group. Scale bar
= 100 μm. **d,** Cell counts from all 14 timepoints in one of the 'No dox' mice. This is the same mouse as
shown in **b**. Bar height for each color indicates the number of cells that were first visible at the
corresponding timepoint that are still visible at the timepoint shown on the X axis; for example, the dark
green bars indicate the number of cells that were first visible on day 10 that are identifiable at each
timepoint from 10 days onward. **e,** Cell counts from all 14 timepoints in one of the 'Dox' mice. This is the
same mouse as shown in **c**. **f,** Total number of imaged cells and number of cells present at the last
imaging timepoint for each of the 'No dox' mice. **g,** Total number of imaged cells and number of cells
present at the last imaging timepoint for each of the 'Dox' mice. Administration of doxycycline greatly
reduces cell attrition. **h,** Survival rates of cells that appeared at each time point for each mouse in both
'Dox' and 'No dox' groups. Each data point represents the fraction of cells that first became visible at that
timepoint that were still present at the last (12-week) timepoint. **i,** Aligned 12-week two-photon image (top
left) and postmortem confocal images (tdTomato, mTagBFP2, and merged) of an FOV from a 'No dox'

mouse (the same mouse as used for panel **b**). Cells found by confocal imaging to have expressed both
tdTomato (reporting rabies virus activity) and mTagBFP2 (reporting G expression) were considered to be
source cells, along with the cells that died during the 12 weeks of two-photon imaging. Scale bar: 20 µm.
j, Aligned 12-week two-photon image (left) and postmortem confocal image of an FOV from a 'Dox'
mouse (the same mouse as used for panel **c**). Because doxycycline shut off mTagBFP2 expression along
with expression of G and L, it was not possible for us to directly determine the surviving number of source
cells in 'Dox' mice; for this group, we used an indirect estimate, as shown in panel **m** below. Scale bar:
20 µm. **k,** Large-scale confocal image of labeled neurons in multiple visual cortical areas in the same
'Dox' mouse. The green rectangle indicates the FOV shown in **j**. This demonstrates that the transsynaptic
spread of ΔGL virus was intact in the mice used for these imaging experiments, just as in the mice used
for the similar experiments shown in previous figures. **l,** Counts for each of the 'No dox' mice of the total
number of imaged tdTomato+ cells (red), the number of cells that disappeared (black), and the number
of cells found by postmortem confocal imaging to have expressed both tdTomato and mTagBFP2 and
which therefore were considered surviving source cells (magenta). The fraction of surviving source cells
for each of these 'No dox' mice was calculated as the ratio (magenta)/(black + magenta). **m,** Counts for
each of the 'Dox' mice of the total number of imaged tdTomato+ cells (red), the number of cells that
disappeared (black), and the *assumed number of total source cells* (magenta) based on the assumption
that the same fraction of total tdTomato+ cells were source cells in the 'Dox' mice as in the otherwise-
identical 'No dox' mice. The fraction of surviving source cells for each of these 'Dox' mice was estimated
as the ratio (magenta - black)/(magenta). All counts and calculations used for estimating source cell
survival rates are given in Supplementary File S6. **n,** Estimated rates of source cell survival for 'Dox' and
'No dox' groups. Administration of doxycycline beginning at two weeks postinjection to suppress
expression of the two deleted viral genes markedly increases cell survival rate from 53.2% ('No dox') to
92.3% ('Dox').

701
702
Fig. 5: Membrane properties of source cells and input cells labeled by first- and second-
generation monosynaptic tracing. a-b, Experimental design. Cre-dependent helper viruses were
injected in S1 of either PV-Cre x Ai65F x TRE-CB het/het/het (ΔGL group) or PV-Cre x Ai65F het/het (ΔG
and AAV-only groups); 7 days later, either RVΔGL-Flpo(EnvA) or RVΔG-Flpo(EnvA) was injected, with
an AAV-only control group receiving no rabies virus injection. Beginning 35 days after RV injection and
continuing for up to 10 days later, brain slices were prepared, and labeled neurons ("input cells" defined
as pyramidal neurons expressing tdTomato alone, or "source cells" defined by neurons expressing both
tdTomato and mTagBFP2) were targeted for whole-cell recordings. For AAV-only mice, "source cells"
denotes neurons expressing mTagBFP2, and "input cells" denotes randomly-selected unlabeled
pyramidal neurons nearby. Brain image adapted from the Allen Mouse Brain Reference Atlas (atlas.brain-
map.org). **c-h,** Membrane properties of input cells (as defined above) in the three experimental groups.
c-f, Resting membrane potential (RMP), membrane capacitance (Cm), rheobase, and input resistance
(Rm) of input neurons. With the second-generation system, all four of the measured membrane properties
remain normal (not statistically different from the corresponding AAV-only control measurements) 5
weeks after injection of ΔGL rabies virus. With the first-generation system, the few labeled neurons found
at this long timepoint had capacitance and input resistance significantly different from controls. ΔGL, 26
cells from 4 mice; AAV control, 13 cells from 3 mice; ΔG, 22 cells from 3 mice; RMP, one-way ANOVA,
F(2, 58) = 0.35, p = 0.71. Capacitance, one-way ANOVA, F(2, 58) = 4.924, p = 0.011, Dunnett's multiple
comparisons test, ΔGL vs. AAV control, p = 0.93, ΔG vs. AAV control, p = 0.023, *p < 0.05. Rheobase,
one-way ANOVA, F(2, 58) = 1.81, p = 0.17. Input resistance, one-way ANOVA, F(2, 58) = 4.13, p = 0.021,
Dunnett's multiple comparisons test, ΔGL vs. AAV control, p > 0.99, ΔG vs. AAV control, #p = 0.060. **g,**
Representative traces of the action potential firing of input cells in response to 250 pA current injection.
h, Plot of action potential firing frequency in response to sequential current step injection showed
increased and normal excitability in the principal neurons labeled by ΔG and ΔGL rabies virus,
respectively. Two-Way ANOVA, F(2, 52) = 5.61, p = 0.0062. Dunnett's multiple comparisons test, ΔGL
vs. AAV control, p = 0.94, ΔG vs. AAV control, p = 0.031, *p < 0.05. **i-n,** Membrane properties of source
cells (as defined above) in the ΔGL and AAV-only groups. Note that no surviving source cells could be
found in the ΔG group at this long (5-week) survival time. **i-l,** Resting membrane potential (RMP),
membrane capacitance (Cm), rheobase, and input resistance (Rm) of source cells (as defined above) in
the two groups (ΔGL and AAV-only) in which extant source cells could be found. Source cells in the ΔGL
group showed more negative resting membrane potential and higher input resistance. AAV control, 19
cells from 3 mice, ΔGL, 10 cells from 4 mice). RMP, unpaired t-tests (two-tailed), p = 0.016, *p < 0.05.
Capacitance, unpaired t-tests (two-tailed), p = 0.95. Rheobase, unpaired t-tests (two-tailed), p = 0.47.
Input resistance, unpaired t-tests (two-tailed), p = 0.040, *p < 0.05. **m,** Representative traces of the action
potential firing of source cells in response to 450 pA current injection. **n,** Action potential firing frequency
in response to sequential current step injection showed similar excitability in the source cells of the ΔGL
and AAV-only groups. Two-way ANOVA, F(1, 27) = 1.42, p = 0.24.

740
741
Fig. 6: Functional two-photon imaging of transsynaptically-labeled neurons' visual response
properties over ten weeks. a, The experimental design was similar to that shown in Fig. 4 but with a
Flp-dependent jGCaMP7s AAV injected along with the rabies virus. Brain image adapted from the Allen
Mouse Brain Reference Atlas (atlas.brain-map.org). **b,** Following the virus injections, the injection site
was imaged on a two-photon microscope while the awake mice were presented with drifting grating
stimuli of different orientations and temporal frequencies, repeatedly for 10 weeks after RV injection. **c,**
Representative two-photon field of view of neurons expressing jGCaMP7s (green channel) and tdTomato
(red channel). Scale bar: 50 μm. **d,** Tuning curves of a jGCaMP7s-expressing neuron obtained with
drifting gratings presented at 12 directions of motion and 5 temporal frequencies (TF), repeated 10 times
(mean Δ F/F ± s.e.m.) at three different time points (left: week 3; middle: week 6; right: week 10). **e,**
Fraction of cells tuned at six different timepoints in two mice. The number of tuned neurons does not
decline with time, suggesting intact response properties despite labeling by RVΔG. **f,** Tuning patterns at
week 10 of 11 jGCaMP7s-expressing neurons showing clear preferred direction tuning, as wells as the
tuning pattern of an untuned neuron, representative of roughly half of imaged cells. **g,** Direction tuning
patterns of six individual cells recorded at multiple time points (from week 1 to week 10). The top two
cells became visible at week one; the middle two appeared at week two, and the bottom two cells
appeared at week three. Tuning patterns remain stable over the entire imaging period. **h,** Single-cell
fluorescence time courses for 25 cells, showing activity over the first 120 s of visual stimulation. Cells are
ranked in descending order of total activity. Scale bar: 10 s.
761

Methods

All experiments involving animals were conducted according to NIH guidelines and approved by the MIT Committee for Animal Care. Mice were housed 1-5 per cage under a normal light/dark cycle for all experiments.

Cloning

pAAV-syn-F14F15S-jGCaMP7s (Addgene 178514) was made by cloning a Flp-dependent FLEX arrangement of mutually-incompatible FRT sites[74] into pAAV-synP-FLEX-EGFP-B19G (Addgene 59333) followed by the jGCaMP7s[75] gene from pGP-CMV-jGCaMP7s (Addgene 104463).

pAAV-syn-F14F15S-splitTVA-EGFP-tTA_v2 (Addgene 183352) was made similarly but with the tricistronic open reading frame from pAAV-syn-FLEX-splitTVA-EGFP-tTA (Addgene 100798) except for the use of FRT sites instead of lox ones (and with a 2-bp frameshift of the in-frame FRT site to prevent creation of a premature stop codon). Note that this plasmid is a replacement for pAAV-synP-F14F15S-splitTVA-EGFP-tTA, which was described in an earlier preprint of this manuscript.

pAAV-syn-FLEX-tTA (Addgene 178516) was made by cloning a codon-optimized tet transactivator gene[76] into pAAV-synP-FLEX-EGFP-B19G (Addgene 59333).

pAAV-syn-Flpo (Addgene 174378) and pAAV-syn-mCre (Addgene 178515) were made by replacing the FLEX cassette from pAAV-synP-FLEX-EGFP-B19G (Addgene 59333) with mouse-codon-optimized genes for Flp[57] and Cre[55].

pCAG-hypBase was made by synthesizing a fragment encoding the "hyperactive" piggyBac transposase iPB7[77] and cloning into the EcoRI & NotI sites of pCAG-GFP[78] (Addgene 11150).

The piggyBac plasmid pB-CAG-B19L-IRES-mCherry-WPRE-BGHpA (Addgene 178518) was made by cloning the CAG promoter from pCAG-B19G (Addgene 59921), the SAD B19 L gene, the EMCV IRES[79], the mCherry[80] gene, and the woodchuck post-transcriptional regulatory element and bovine growth hormone polyadenylation signal from pCSC-SP-PW-GFP (Addgene 12337), into PB-CMV-MCS-EF1-Puro (System Biosciences #PB510B-1).

The "TLoop"-style[81] lentiviral transfer vectors pLV-TTBG (Addgene 115232) and pLV-TTBL (Addgene 115233) were made by replacing the CMV promoter and EGFP gene in pCSC-SP-PW-GFP (Addgene 12337) with the leaky tetracycline response element from pAAV-FLEX-hGTB (Addgene 26196) followed by a tricistronic open reading frame consisting of the genes for the "improved tetracycline transactivator" itTA[82], mTagBFP2[54], and either the glycoprotein or polymerase gene, respectively, of the rabies virus SAD B19 strain, separated by P2A elements.

Mouse strains

Mice used in this study were crosses (triple transgenic for DAT-IRES-Cre and PV-Cre experiments, double transgenic for corticostriatal experiments, double or single transgenic (either Cre-negative or L-negative) for control experiments) of the following strains, all in a C57BL/6J (Jackson Laboratory 000664) background:

PV-Cre[62], DAT-IRES-Cre[61], Ai14[58], and DAT-P2A-Flpo[83] were purchased from Jackson Laboratory (catalog #s 017320, 006660, 007914, and 035436).

The Flp-dependent tdTomato reporter line Ai65F was obtained in our case by crossing the Cre- and Flp-dependent tdTomato double-reporter line Ai65D[46] (Jackson Laboratory 021875) to the Cre deleter line Meox2-Cre[84] (Jackson Laboratory 003755), then breeding out the Meox2-Cre allele. An equivalent Ai65F line, made using a different Cre deleter line, was described in Daigle et al. '18[47] and is now available from Jackson Laboratory with catalog # 032864).

The L-expressing mouse line TRE-CB (TRE-tight-mCardinal-P2A-B19L, being distributed by Jackson Laboratory with catalog # 036974) was generated by cloning the genes for mCardinal[44] and the rabies virus (SAD B19) polymerase, separated by a picornavirus 2A element[85], into "Ai62(TITL-tdT) Flp-in replacement vector" (Addgene 61576), to make the Flp-in construct "TRE-mCardinal-P2A-B19L Flp-in vector" (Addgene # 178519). The lox-stop-lox sequence present in the original Flp-in vector was removed to make the new construct, so that the tetracycline response element TRE-tight[86] drives mCardinal-P2A-

813 B19L directly with no dependence on Cre recombination. This plasmid was then used for FLP-mediated
814 targeted insertion into the TIGRE locus[45] in Ai99 embryonic stem cells as described[47]. Clones verified as
815 containing the correct insert were used for production of knock-in mice by the Mouse ES Cell &
816 Transgenic Facility at the Koch Institute for Integrative Cancer Research at MIT. All subsequent breeding
817 steps were with mice of a C57BL/6J background.

Production of lentiviral vectors for making cell lines

820 Lentiviral vectors were made as described[87] but with the VSVG expression vector pMD2.G (Addgene
821 12259) as the envelope plasmid and with the following transfer vectors:

822 pLV-TTBL (described above), to make the L-expressing lentiviral vector LV-TTBL(VSVG),

823 pLV-U-TVA950 (Addgene 115236), to make the TVA-expressing lentiviral vector LV-U-TVA950(VSVG).

Production of cell lines

To make the L-expressing cell line BHK-B19L, BHK-21 cells (ATCC# CCL-10) were transfected with
pCAG-hypBase and pB-CAG-B19L-IRES-mCherry-WPRE-BGHpA; the cells were expanded into two 15c
plates, then resuspended and sorted on a BD FACS Aria to collect the brightest 5% of mCherry-
expressing cells.

BHK-B19L-TVA950 cells were made by infecting BHK-B19L cells (described above) with LV-U-
TVA950(VSVG) at a multiplicity of infection of approximately 100. Cells were then expanded and frozen
in medium containing 10% DMSO for subsequent use.

The BHK-EnvA2-TTBL2 cell line, expressing both the "EnvARGCD" fusion protein[1] and the RV
polymerase, was made by infecting BHK-EnvA2 cells[88] with the L-expressing lentiviral vector LV-
TTBL(VSVG) (described above) at a multiplicity of infection of approximately 100. The cells were
expanded into three 15 cm plates, then sorted on a BD FACS Aria to collect those cells with the highest
2% of both green (EnvA) and blue (L) fluorescence. After expanding the collected cells again, we
observed that blue fluorescence was so dim as to be difficult to detect by eye; we therefore sorted the
cells a second time, again retaining cells with the brightest 2% fluorescence in both channels. The twice-
sorted cells, now called BHK-EnvA2-TTBL2, were expanded again and frozen in medium containing 10%
DMSO for subsequent use.

Production and titering of adeno-associated viruses

AAV genome plasmids pAAV-syn-FLEX-splitTVA-EGFP-tTA (Addgene 100798) and pAAV-TREtight-
mTagBFP2-B19G (Addgene 100799) have been described previously[38]. These genomes were packaged
in serotype 1 AAV capsids by Addgene (catalog numbers 52473-AAV1, 100798-AAV1, and 100799-
AAV1). The two helper AAVs from Addgene were diluted in Dulbecco's phosphate-buffered saline (DPBS)
(Thermo, 14190-250) to final titers of 7.22E+10 genome counts (gc)/mL for AAV1-syn-FLEX-splitTVA-
EGFP-tTA and 6.50E+11 gc/mL for AAV-TREtight-mTagBFP2-B19G, then combined in a 50/50 ratio by
volume before injection. AAVs from Addgene were initially thawed and aliquoted in 20x 5ul aliquots; to
make working dilutions, a 5ul aliquot of undiluted virus was thawed, diluted in DPBS (Thermo Fisher
14190-250) to the desired working dilutions (1:20 or 1:200)[48,49]. pAAV-syn-FLEX-tTA (described above)
was packaged as serotype 1 by the UNC Vector Core with titer of 2.51E+12 gc/mL.

rAAV2-retro-syn-Cre, rAAV2-retro-syn-Flpo, AAV1-syn-F14F15S-jGCaMP7s, and AAV1-syn-F14F15S-
sTpEptTA_v2 were made by transfecting 8x15c plates (per virus) of HEK 293T cells using Xfect
transfection reagent (Takara 631318). Briefly, plates were treated with poly-L-lysine solution were
transfected with pAAV vector, pHelper (Cellbiolabs VPK-421), and the appropriate rep/cap plasmid
("rAAV2-retro helper" (Addgene 81070) or pAAV-RC1 (Cellbiolabs VPK-421)). 4 hours after transfection,
the medium from plates was aspirated and replaced with 15mL DMEM (Thermo #11995-081) containing
2% fetal bovine serum (FBS) (HyClone SH30071.02) and antibiotic/antimycotic (Thermo Fisher 15240-
062). 72 hours after transfection, the supernatants were collected from each plate and replaced again
with 15mL DMEM with 2% FBS and antibiotic/antimycotic. Supernatants were stored in 50mL tubes at
4°C. 120 hours after transfection, cells and supernatants were collected and transferred into sterile 50mL
tubes. Cells were pelleted by centrifugation and washed with DPBS, and supernatants were transferred
into a sterile bottle for PEG precipitation. Cells were lysed by four freeze/thaw cycles and stored at -80°C

until final thaw and purification steps. 50mL of cold 40% polyethylene glycol 8000 (PEG) in 2.5M NaCl
was added to 250mL clarified supernatants and mixed thoroughly. This solution was incubated at 4°C
overnight, then transferred into sterile 50mL tubes and centrifugated at 4000 RCF for 30 minutes @ 4°C.
Supernatants were discarded and pellets resuspended in a total of 1mL 150mM NaCl, 50mM Tris buffer.
For final purification steps, cell pellets were combined with PEG-precipitated virus, treated with
Benzonase nuclease (250U/uL) for 30 minutes at 37°C, then centrifugated at 7188 RCF for 30 minutes
at 4°C. The clarified lysate was transferred to Optiseal tubes (Beckman #361625) containing layered
gradients of iodixanol (15%, 25%, 40%, and 54% iodixanol solutions). Tubes were ultracentrifugated in
a Beckman 70 Ti fixed-angle rotor at 70,000 rpm for 1hr 40min at 15°C on a WX 80+ centrifuge (Thermo
Scientific). The layers containing virus were carefully removed from between the 40% and 54% iodixanol
layers, filtered through 0.22um polyethersulfone (PES) membrane syringe filters, and concentrated in
Amicon Ultra-15 Centrifuge Units (Millipore #UFC910008), with three full buffer exchanges with DPBS to
remove all residual iodixanol. Each batch of virus was concentrated to a final volume of 250-300 µL total,
aliquoted in 5 µl aliquots and stored at -80°C, and titered by quantitative PCR.
880
Production of rabies viruses

EnvA-enveloped ΔG rabies viruses were made as described previously[30,88,89], using genome plasmids
pRVΔG-4Flpo (Addgene 122050) and pRVΔG-4Cre (Addgene 98034).

EnvA-enveloped ΔGL rabies viruses were rescued in 15 cm plates as described[23] using genome plasmids
pRVΔGL-4Flpo (Addgene 98040) and pRVΔGL-4Cre (Addgene 98039). Supernatants from rescue plates
were passaged on 15 cm plates of HEK 293T/17 cells (ATCC) transfected with pLV-TTBG and pLV-TTBL
(described above) using Xfect transfection reagent (Takara 631318) according to the manufacturer's
protocol. Supernatants were titered on reporter cells[23] and used to infect BHK-EnvA2-TTBL2 cells
(described above) at a multiplicity of infection of approximately two. Supernatants from these producer
cells were collected and purified as described[88,89].
891
Titering of rabies viruses

Recombinase-expressing rabies viruses were titered primarily on the reporter cell lines 293T-FLEX-BC,
293T-F14F15S-BC, 293T-FLEX-BC-TVA and 293T-F14F15S-BC-TVA in infectious units per ml (iu/ml)
as described[30] but using three-fold serial dilutions, instead of ten-fold dilutions, for higher precision. For
the corticostriatal experiment, in order to directly compare viral titers in a way that did not depend on the
efficacy of the encoded recombinases, we made the BHK-B19L-TVA950 cell line (described above),
which expresses both TVA, to allow infection by EnvA-enveloped viruses, and L, in order to allow
intracellular replication of ΔGL viruses so that infected cells could be clearly labeled using immunostaining
against the rabies virus nucleoprotein. BHK-B19L-TVA950 cells, as well as BHK-B19L cells, were
infected with three-fold serial dilutions of RVΔGL-Cre(EnvA) and RVΔGL-Flpo(EnvA. Three days after
infection, cells were fixed with 4% paraformaldehyde in PBS and immunostained with a FITC-conjugated
anti-nucleocapsid monoclonal antibody blend (Light Diagnostics Rabies DFA, EMD Millipore, catalog #
5100) diluted 1:100 in 1% BSA, 0.1% Triton in PBS. Immunolabeled cells were analyzed by flow
cytometry to determine titers as described[88].

The titers of the rabies viruses were determined to be as follows:
RVΔGL-Cre(EnvA):
on 293T-FLEX-BC-TVA950:   1.28E+09 iu/ml
on 293T-FLEX-BC (no TVA):   1.48E+05 iu/ml
on BHK-B19L-TVA950 with anti-N staining: 6.13E+08 iu/ml
RVΔGL-Flpo(EnvA):
on 293T-F14F15S-BC-TVA950:   2.94E+08 iu/ml
on 293T-F14F15S-BC (no TVA):   3.26E+05 iu/ml
on BHK-B19L-TVA950 with anti-N staining: 7.67E+08 iu/ml
RVΔG-Cre(EnvA):
on 293T-FLEX-BC-TVA950:   1.24E+10 iu/ml
on 293T-FLEX-BC (no TVA):   4.02E+05 iu/ml
RVΔG-Flpo(EnvA):
on 293T-F14F15S-BC-TVA950:   4.49E+09 iu/ml
on 293T-F14F15S-BC (no TVA):   4.81E+05 iu/ml

RVΔG-4Flpo(B19G):
922   on 293T-F14F15S-BC: 1.86E+10 iu/ml
923   RVΔGL-Flpo(B19G):
924   on 293T-F14F15S-BC: 1.86E+09 iu/ml
925

Stereotaxic injections

Stereotaxic injections were made in anesthetized adult mice of both sexes using a stereotaxic instrument (Stoelting Co., 51925) and a custom injection apparatus consisting of a hydraulic manipulator (Narishige, MO-10) with headstage coupled via custom adaptors to a wire plunger advanced through pulled glass capillaries (Drummond, Wiretrol II) back-filled with mineral oil and front-filled with viral vector solution. We have described this injection system in detail previously[48].

For the corticostriatal experiments (Fig. 1), 200 nl of AAV2-retro-syn-Flpo (1.16E+13 GC/mL; Ai14 or Ai14 x TRE-CB mice) or AAV2-retro-synP-mCre (1.48E+13 GC/mL, diluted to 1.16E+13 GC/mL for matching to the Flpo version; Ai65F or Ai65F x TRE-CB mice) was injected into dorsolateral striatum (AP = +0.74 mm w.r.t. bregma, LM = 2.25 mm w.r.t. bregma, DV = -2.30 mm w.r.t the brain surface); in the same surgery, 250nl of helper virus mixture (AAV1-syn-F14F15S-sTpEptTA_v2 (diluted to 7.19E+10 GC/mL; Ai14 or Ai14 x TRE-CB mice) or AAV1-syn-FLEX-splitTVA-EGFP-tTA (diluted to 7.22E+10 GC/mL; Ai65F or Ai65F x TRE-CB mice) mixed with AAV1-TREtight-mTagBFP2-B19G (diluted to 6.50E+11) in a 50/50 ratio by volume) was injected into S1BF, layer 5 DLS projection region (AP = -1.55 mm w.r.t. bregma, LM = 3.00 mm w.r.t. bregma, DV = -0.75 mm w.r.t the brain surface). 7 days after AAV injection, 250nl of RVΔGL-Cre(EnvA) (6.13E+08 iu/ml as titered by nucleoprotein staining on L-expressing cells; see above) or RVΔGL-Flpo(EnvA) (7.67E+08 iu/ml, diluted to 6.13E+08 iu/ml) or RVΔG-Flpo(EnvA) (4.49E+09 iu/ml) or RVΔG-Cre(EnvA) (1.24E+10 iu/ml, diluted to 4.49E+09 iu/ml) was injected at the same cortical injection site as the helper viruses. For no-G controls, DPBS was included instead of AAV1-TREtight-mTagBFP2-B19G.

For the anatomical experiments in Cre lines (Fig. 2), 300 nl of helper AAV mixture (AAV1-syn-FLEX-splitTVA-EGFP-tTA (diluted to 7.22E+10 GC/mL) mixed with AAV1-TREtight-mTagBFP2-B19G (diluted to 6.50E+11 GC/mL) in a 50/50 ratio by volume; note that these are the same dilutions as we have recommended previously[48,49] but retitered by qPCR in our own lab) was injected into either primary somatosensory cortex or substantia nigra pars compacta (see below for stereotaxic coordinates). Injection coordinates for substantia nigra pars compacta were: anteroposterior (AP) = -3.00 mm with respect to (w.r.t.) bregma, lateromedial (LM) = +1.50 mm w.r.t bregma, dorsoventral (DV) = -4.20 mm w.r.t the brain surface); injection coordinates for somatosensory cortex were: AP = -0.58 mm w.r.t. bregma, LM = 3.00 mm w.r.t. bregma, DV = -1.00 mm w.r.t the brain surface. Seven days after AAV injection, 500 nl (PV-Cre and DAT-IRES-Cre) of RVΔGL-Flpo(EnvA) (7.67E+08 iu/mL) was injected at the same injection site as the AAVs. The experiments in DAT-P2A-Flpo mice (Fig. 3) were done similarly but using the Flp-dependent AAV-syn-F14F15S-EGFP-tTA_v2 (diluted to 7.19E+10 GC/mL) and RVΔG-Cre(EnvA) (1.24E+10 iu/ml, diluted to 4.49E+09 iu/ml).

For longitudinal two-photon imaging for comparison of RVΔG-Flpo(B19G) and RVΔGL-Flpo(B19G), 200 nL RVΔG-Flpo(B19G) (1.86E+10 iu/ml) or RVΔGL-Flpo(B19G) (1.86E+09 iu/ml) was injected into V1 (AP = -2.70 mm w.r.t. bregma, LM = 2.50 mm w.r.t. bregma, DV = -0.26 mm w.r.t. brain surface) into Ai65F mice. Glass windows composed of a 3mm-diameter glass coverslip (Warner Instruments CS-3R) glued (Optical Adhesive 61, Norland Products) to a 5mm-diameter glass coverslip (Warner Instruments CS-5R) were then affixed over the craniotomy with Metabond (Parkell), and custom stainless steel headplates (eMachineShop) were affixed to the skulls around the windows.

For longitudinal two-photon structural imaging of live monosynaptic tracing at the injection site, a 3 mm craniotomy was opened over primary visual cortex (V1). 300 nl of AAV1-syn-FLEX-splitTVA-EGFP-tTA (diluted to 7.22E+10 GC/mL; Ai65F or Ai65F x TRE-CB mice) mixed with AAV1-TREtight-mTagBFP2-B19G (diluted to 6.50E+11) in a 50/50 ratio by volume was injected into V1 (AP = -2.70 mm w.r.t bregma, LM = 2.50 mm w.r.t. bregma, DV = -0.26 mm w.r.t. brain surface), followed by implantation of windows as described above. Seven days after injection of helper AAVs, the coverslips were removed and 100 nl of RVΔGL-Flpo (EnvA) (7.67E+08 iu/mL) was injected at the same site. Coverslips were reapplied and custom stainless steel headplates (eMachineShop) were affixed to the skulls around the windows.

For functional imaging experiments, the V1 injection coordinates were AP = -2.45 mm w.r.t bregma, LM = 2.00 mm w.r.t bregma, DV = -0.26 mm w.r.t. brain surface, and the RVΔGL-Flpo(EnvA) (7.67E+08

976   iu/mL) was mixed in a 50/50 ratio by volume with AAV1-syn-F14F15S-jGCaMP7s (5.44E+12 GC/ml,
977   diluted 1:10 in DPBS) before injection of 200 nl of the mixture 7d following the helper AAV injection.
978

Doxycycline administration

'No-dox' mice were fed with regular rodent chow throughout, while 'dox (food)' mice were switched to
chow containing doxycycline 200 mg/kg (Fisher Scientific, 14-727-450) beginning two weeks after RV
injection and maintained on doxycycline chow until perfusion, in order to suppress rabies viral polymerase
and glycoprotein expression. 'Dox (inj+food)' mice, including those used for the structural two-photon
imaging experiment, also received intraperitoneal injections of 100 mg/kg doxycycline every 12 hours for
three days, beginning two weeks after RV injection.
986

In vivo two-photon imaging and image analysis

For longitudinal two-photon imaging of live monosynaptic tracing, injection sites were imaged on a
Prairie/Bruker Ultima IV In Vivo two-photon microscope driven by a Spectra Physics Mai-Tai Deep See
laser with a mode locked Ti:sapphire laser emitting at a wavelength of 1020 nm for tdTomato. Mice were
reanesthetized and mounted via their headplates to a custom frame, with ointment applied to protect their
eyes and with a handwarmer maintaining body temperature. One field of view was chosen in each mouse
in the area of maximal fluorescent labelling. The imaging parameters were as follows: image size 1024
X 1024 pixels (565.1 µm x 565.1 µm), 0.360 Hz frame rate, dwell time 2.0 µs, 1x optical zoom, Z-stack
step size 1 µm. Image acquisition was controlled with Prairie View 5.4 software. Laser power exiting the
20x water-immersion objective (Zeiss, W plan-apochromat, NA 1.0) varied between 20 and 65 mW
depending on focal plane depth (Pockels cell value was automatically increased from 450 at the top
section of each stack to 750 at the bottom section). For the example images of labeled cells, maximum
intensity projections (stacks of 150-400 µm) were made with Fiji software. Cell counting was performed
with the ImageJ Cell Counter plugin. When doing cell counting, week 1 tdTomato labelled cells were
defined as a reference; remaining week 1 cells were the same cells at later time point that align with week
1 reference cells but the not-visible cells at week 1 (the dead cells). Plots of cell counts were made with
Origin 7.0 software (OriginLab, Northampton, MA) or Prism 9 (GraphPad Software, San Diego, California).

For functional two-photon imaging of source cells, injection sites in left-hemisphere V1 was chosen as
the imaging area. This imaging was performed using the same microscope (5.356-Hz frame rate, 1024
X 128 pixels, 565.1 µm x 565.1 µm, dwell time 0.8 µs, 1x optical zoom, scan angle 45 degree) with the
same objective and laser (at 920 nm) as in the structural imaging experiments. Laser power at the
objective ranged from 10 to 65 mW. Calcium imaging data were acquired in supragranular layers (100 to
200 µm deep). The animal was awake, head-fixed and the animal body is free to lay into a customized
tube with some holes for air circulation. No behavioral training or reward was given. In order to find the
same FOVs in subsequent weeks, surface vasculature map was draw on a notebook for each mouse
under microscope with light source (Leica M50) and put '*' marker near the location of FOVs in this map
for reference when doing the first imaging session. The cell body images from averaged data taken in
the first session provided a template for fine alignment. Visual stimuli were generated in Matlab (R2015R
version) with custom software based on Psychtoolbox (http://psychtoolbox.org) and shown on the same
LCD screen as in the widefield mapping experiments. Each condition consisted of 2 s of a full-field sine-
wave grating drifting in one direction, presented at 80% contrast with spatial frequency of 0.04 cycles/
degree, followed by 2 s of uniform mean luminance (gray). All permutations of 12 directions (30° steps)
and 5 temporal frequencies (1, 2, 4, 8 and 15 Hz) were shown, in randomized order. The complete set
was repeated 10 times, for a total stimulation period of 40 min per FOV per session. Cells were then
manually segmented, and single-cell fluorescence traces were extracted by averaging the fluorescence
of all pixels masking the soma by ImageJ (versions: 2-0-0-rc-69) software. The mean ΔF/F over the full
2 s of each stimulus condition was used to calculate orientation tuning curves, with background
fluorescence (F) in Δ F/F taken as the value of the trace immediately preceding a condition, averaged
over all conditions. The raw calcium traces from cells within individual FOVs (not across FOVs, given
different imaging conditions across animals and time points) were sorted by mean fluorescence. For
'tuned' cells in Figure 4 panel e, the counts are based on all imaged neurons' individual tuning curves,
plotted in MATLAB; any cell showing response to a preferred orientation (including narrowly tuned
neurons and broadly tuned neurons) at any temporal frequency (1Hz, 2Hz, 4Hz, 8Hz, or 15Hz) was
counted manually as a tuned cell (see Supplementary File S7 for counts of tuned and untuned cells).
031

Brain slice electrophysiology

Coronal brain slices containing the S1 were collected from 12 to 32 week old PV-Cre x Ai65F x TRE-CB het/het/het (ΔGL group) or PV-Cre x Ai65F het/het (ΔG and AAV-only groups) for ex vivo patch clamp recordings. Mice were anesthetized with isoflurane, perfused with ice cold cutting solution, and decapitated using scissors. Brains were extracted and immersed in ice-cold (0–4 °C) sucrose cutting solution containing (in mM) 252 sucrose, 26 NaHCO3, 2.5 KCl, 1.25 NaH2PO4, 1 CaCl2, 5 MgCl2 and 10 glucose, which was oxygenated with 95% O2 and 5% CO2. The brains were trimmed and coronal brain slices (300 μm) were sectioned using a vibratome (VT1200, Leica). After sectioning, slices for patch clamp recordings were transferred to a holding chamber containing oxygenated patch clamp recording medium (aCSF) containing (in mM): 126 NaCl, 2.5 KCl, 1.25 NaH2PO4, 1.3 MgCl2, 2.5 CaCl2, 26 NaHCO3, and 10 glucose, where they were maintained at 32 °C for 30 min before decreasing the chamber temperature to ~20 °C. Slices were transferred one at a time from the holding chamber to a submerged recording chamber mounted on the fixed stage of an Olympus BX51WI fluorescence microscope equipped with differential interference contrast (DIC) illumination. The slices in the recording chamber were continuously perfused at a rate of 2 ml/min with recording aCSF at room temperature and continuously aerated with 95% O2/5% CO2. Whole-cell patch clamp recordings were performed in the fluorescently-labeled PV+ neurons and principal neurons in the S1 and neighboring S2 region. The presynaptic principal neurons to PV cells in the AAV control group were randomly selected based on the size, morphology, and accommodating firing pattern. Glass pipettes with a resistance of 4-8 MΩ were pulled from borosilicate glass (ID 1.2 mm, OD 1.65 mm) on a horizontal puller (Sutter P-97) and filled with an intracellular patch solution containing (in mM): 130 potassium gluconate, 10 HEPES, 10 phosphocreatine Na2, 4 Mg-ATP, 0.4 Na-GTP, 5 KCl, 0.6 EGTA; pH was adjusted to 7.25 with KOH and the solution had a final osmolarity of 293 mOsm. Series resistance was continuously monitored. Cells were discarded when the series resistance changed more than 20%. Data were acquired using a Multiclamp 700B amplifier, a Digidata 1440 A analog/digital interface, and pClamp 10 software (Molecular Devices). Data were analyzed with Clampfit 10 (Molecular Devices). The membrane capacitance (Cm) and Input resistance (Rm) were calculated from a membrane seal test conducted in voltage-clamp mode, in which 100-ms, 5-mV voltage steps were delivered at a frequency of 5 Hz. In order to evaluate the action potential rheobase, a 1-s, positive current was delivered in current-clamp mode at 10pA steps. Rheobase was defined as the minimum current required to depolarize the membrane potential for action potential firing. Finally, to evaluate the relationship between injected current and firing frequency, we delivered a series of 0.5s current pulses in current-clamp at 50pA steps (50-250pA for principal cells, 50-450pA for PV cells). Statistical comparisons were conducted with unpaired Student's t-test or with a one- or two-way ANOVA followed by a post hoc Dunnett's test as appropriate ($p < 0.05$ with a two-tailed analysis was considered significant).

Perfusions and histology

1 to 12 weeks (depending on experiment; see main text) after injection of rabies virus, mice were transcardially perfused with 4% paraformaldehyde in phosphate-buffered saline. Brains not used for two-photon imaging were postfixed overnight in 4% paraformaldehyde in PBS on a shaker at 4°C and cut into 50 μm coronal (S1 injections) or parasagittal (SNc injections) sections on a vibrating microtome (Leica, VT-1000S). Sections were collected sequentially into 6 tubes containing cryoprotectant, so that each tube contained a sixth of the collected tissue. For coronal sections, 15 rounds of section were collected. For sagittal sections, 12 rounds were collected. Brains from mice used for two-photon imaging were postfixed in 4% paraformaldehyde/30% sucrose in PBS for two days on a shaker at 4°C, then cut into 50 μm sections on a freezing microtome in a plane approximately tangential to the surface of the brain at the imaged location. Sections were immunostained as described[48] with the following primary antibodies (as applicable) at the following respective dilutions: chicken anti-GFP (Aves Labs GFP-1020) 1:500, guinea pig anti-parvalbumin (Synaptic Systems 195004) 1:1000, sheep anti-tyrosine hydroxylase (Millipore AB1542)) 1:1000, with secondary antibodies donkey anti-chicken Alexa Fluor 488 (Jackson Immuno 703-545-155) 1:200, donkey anti-guinea pig, AlexaFluor 647 conjugated (Jackson Immuno 706-605-148) 1:200, and donkey anti-sheep, AlexaFluor 647 conjugated (Jackson Immuno 713-605-147)) 1:200. Sections were mounted with Prolong Diamond Antifade mounting medium (Thermo Fisher P36970) and imaged on a confocal microscope (Zeiss, LSM 900).

Cell counts and statistical analysis

Labeled neurons in striatum (in DAT-IRES-Cre mice), contralateral cortex and thalamus (in PV-Cre mice
and for the corticostriatal experiments), and at the injection sites were counted either manually or with
the automated Cell Counter function in ImageJ, in every sixth 50-μm section on an epifluorescence
microscope (Zeiss Imager.Z2). Coronal sections (corticostriatal experiments and PV-Cre) included
sections between 1.2mm and -3.3mm relative to Bregma. Sagittal sections (DAT-IRES-Cre) covered
sections 3.6mm to 0.0mm relative to Bregma. Cells expressing EGFP, mTagBFP2, or both, along with
tdTomato, were counted by adding separate labels to each and then looking for overlapping cells. Groups
were compared using single factor ANOVAs. Symbols denoting significance levels on graphs are as
follows: *** = $p<0.001$; ** = $0.001 \leq p < 0.01$; * = $0.01 \leq p < 0.05$; n.s. = $p \geq 0.05$.
097

Reporting Summary

099 Further information on research design is available in the Nature Research Reporting Summary linked to
100 this paper.
101

Data Availability

103 All cell counts and statistical analyses are provided in Supplementary Information. The novel plasmids
104 described in this paper have been deposited with Addgene with the accession numbers given in Methods.
105 The TRE-CB mouse line is available from the Jackson Laboratory (accession number 036974).
106

References for Methods

74.   Turan, S., Kuehle, J., Schambach, A., Baum, C. & Bode, J. Multiplexing RMCE: versatile extensions of
the Flp-recombinase-mediated cassette-exchange technology. *J Mol Biol* **402**, 52-69 (2010).
75.   Dana, H. *et al.* High-performance calcium sensors for imaging activity in neuronal populations and
microcompartments. *Nat Methods* **16**, 649-657 (2019).
76.   Shimshek, D.R. *et al.* Enhanced odor discrimination and impaired olfactory memory by spatially controlled
switch of AMPA receptors. *PLoS biology* **3**, e354 (2005).
77.   Yusa, K., Zhou, L., Li, M.A., Bradley, A. & Craig, N.L. A hyperactive piggyBac transposase for mammalian
applications. *Proc Natl Acad Sci U S A* **108**, 1531-6 (2011).
78.   Matsuda, T. & Cepko, C.L. Electroporation and RNA interference in the rodent retina in vivo and in vitro.
Proceedings of the National Academy of Sciences of the United States of America **101**, 16-22 (2004).
79.   Gallardo, H.F., Tan, C. & Sadelain, M. The internal ribosomal entry site of the encephalomyocarditis virus
enables reliable coexpression of two transgenes in human primary T lymphocytes. *Gene Ther* **4**, 1115-9
(1997).
80.   Shaner, N.C. *et al.* Improved monomeric red, orange and yellow fluorescent proteins derived from
Discosoma sp. red fluorescent protein. *Nature biotechnology* **22**, 1567-72 (2004).
81.   Cetin, A.H. & Callaway, E.M. Optical Control of Retrogradely Infected Neurons Using Drug Regulated
"TLoop" Lentiviral Vectors. *J Neurophysiol* (2014).
82.   Krestel, H.E. *et al.* A genetic switch for epilepsy in adult mice. *The Journal of neuroscience : the official
journal of the Society for Neuroscience* **24**, 10568-78 (2004).
83.   Kramer, D.J. *et al.* Generation of a DAT-P2A-Flpo mouse line for intersectional genetic targeting of
dopamine neuron subpopulations. *Cell Rep* **35**, 109123 (2021).
84.   Tallquist, M.D. & Soriano, P. Epiblast-restricted Cre expression in MORE mice: a tool to distinguish
embryonic vs. extra-embryonic gene function. *Genesis* **26**, 113-5 (2000).
85.   Kim, J.H. *et al.* High cleavage efficiency of a 2A peptide derived from porcine teschovirus-1 in human cell
lines, zebrafish and mice. *PLoS one* **6**, e18556 (2011).
86.   Backman, C.M., Zhang, Y., Hoffer, B.J. & Tomac, A.C. Tetracycline-inducible expression systems for the
generation of transgenic animals: a comparison of various inducible systems carried in a single vector. *J
Neurosci Methods* **139**, 257-62 (2004).
87.   Wickersham, I.R. *et al.* Lentiviral vectors for retrograde delivery of recombinases and transactivators.
Cold Spring Harb Protoc **2015**, 368-74 (2015).
88.   Wickersham, I.R., Sullivan, H.A. & Seung, H.S. Production of glycoprotein-deleted rabies viruses for
monosynaptic tracing and high-level gene expression in neurons. *Nature protocols* **5**, 595-606 (2010).
89.   Wickersham, I.R. & Sullivan, H.A. Rabies viral vectors for monosynaptic tracing and targeted transgene
expression in neurons. *Cold Spring Harb Protoc* **2015**, 375-85 (2015).
142

Acknowledgements

We thank Aurora Burds-Connor and Noranne Enzer of the Preclinical Modeling Facility at the Koch
Institute for Integrative Cancer Research at MIT for generation of the TRE-CB line from ES cells. We
thank Karel Svoboda for advice on titrating jGCaMP6s expression for reduced toxicity. We thank Ayano
Matsushima for examination of labeled regions in the DAT-P2A-Flpo mice. Research reported in this
publication was supported by BRAIN Initiative awards U01MH106018, U01MH114829, U19MH114830,
and RF1MH120017 from the National Institute of Mental Health.
150

151 Author contributions

L.J. contributed molecular biology, cell culture, mouse colony management, virus injections, two-photon
imaging, histology, microscopy, data analysis, and preparation of figures. H.A.S. produced viruses and
conducted cell culture assays. M.Z. contributed perfusions, histology, microscopy, cell counts, data
analysis, and mouse colony management. T.K.L. contributed virus injections, perfusions, histology,
microscopy, and mouse colony management. M.M., C.X. and Y.H contributed cloning and molecular
biology. N.E.L., M.P., & K.R.B. contributed mouse colony management, assisted with molecular
biology. L.R. contributed cell counts. T.L.D. and H.Z. contributed embryonic stem cell targeting for
making the TRE-CB mouse line. J.I., M.H., and M.S. assisted with establishing the structural and
functional two-photon imaging. X.F and G.F. planned slice electrophysiology experiments; X.F.
conducted slice electrophysiology experiments and identified the problem with the original Flp-
dependent helper AAV that was used for the original version of the manuscript. I.R.W. designed
experiments and wrote the paper with input from L.J., H.A.S., M.Z., and other authors.
164

165 Competing interests

166 I.R.W. is a consultant for Monosynaptix, LLC, advising on design of neuroscientific experiments.
167

168 Additional information

169 Supplementary information is available for this paper.
170 A preprint version of this paper is available on bioRxiv.
171 Correspondence and requests for materials should be addressed to I.R.W. (wickersham@mit.edu).
172

Extended Data Fig. S1: Intrinsic mCardinal fluorescence was not detected by confocal imaging, suggesting very low L expression in the conditions implemented. Representative confocal images of injection sites in PV-Cre x Ai65F x TRE-CB mice, showing lack of clear mCardinal fluorescence induced by either of the laser lines (561 nm or 640 nm) flanking the excitation maximum on the Zeiss LSM 900 confocal microscope. Because of the lack of fluorescence in the channel otherwise used for AlexaFluor 647, specifically, we were able to use that channel for immunostaining for PV and TH (Fig. 1 and Extended Data Figs. wasFigS2 & wasFigS3). **a,** Injection sites of a FLEX AAV expressing tTA without a fluorophore, and **b,** injection sites of the FLEX AAV expressing TVA, EGFP, and tTA, with survival times of 3 and 4 weeks in each case. Left: EGFP channel; center-left: tdTomato channel; center-right: AlexaFluor 647 channel; right: "best case" for imaging mCardinal: excitation with 561 nm and collecting all emitted light between 565 nm and 700 nm. Scale bars: 200 µm, apply to all panels.

186
187
188 **Extended Data Fig. S2: Representative images of control experiments for corticostriatal**
189 **experiments.** See Fig. 2 for explanation of experimental design and quantification. **a-b,** First-
190 generation controls (7d and 14d survival times). **c-d,** Controls using second-generation viruses: no L,
191 no G, and no Cre/Flpo controls (all without doxycycline), as well as the 'dox (inj+food)' condition for
192 comparison. Scale bars: 200 μm, apply to all images.
193

194
195
Extended Data Fig. S3: Counts of corticothalamic "source" and "starter" cells and ratios of
contralateral and thalamic cells to source and starter cells. Counts are totals found in every sixth
50-μm section spanning each brain. **a-b,** Results using RVΔGL-Flpo(EnvA). **a,** Counts of source cells

(left), defined as cells coexpressing tdTomato (marking activity of Flpo) and mTagBFP2 (marking
expression of G), and ratios of these numbers of source cells to numbers of tdTomato+ cells in
contralateral cortex (middle) and thalamus (right). Note the near-absence of source cells in the first-
generation conditions at two weeks, suggesting that most had died by this time point. Note also that the
low numbers of source cells in the second-generation conditions with doxycycline could reflect either
cell death or simply the intended suppression of mTagBFP2 expression in these mice. For mice in
which no source cells were found (either because they had died or because mTagBFP2 was
suppressed by doxycycline), no ratio is included on the graphs. **b,** Counts and ratios of starting cells,
defined here as cells coexpressing tdTomato (marking activity of Flpo) and EGFP (marking expression
of TVA and tTA). **c-d,** Results using RVΔGL-Cre(EnvA). **c,** Counts and ratios of source cells, defined
here as cells coexpressing tdTomato (marking activity of Cre) and mTagBFP2 (marking expression of
G). **d,** Counts and ratios of starting cells, defined here as cells coexpressing tdTomato (marking activity
of Cre) and EGFP (marking expression of TVA and tTA).
212

213   *(images in external file)*
214
215   **Extended Data Fig. S4: Series of whole-brain images of labeled inputs to corticostriatal neurons**
216   **using first- and second-generation rabies viral vectors expressing Cre.** Images are tiled confocal
217   images of every sixth 50-µm coronal section spanning most of the rostrocaudal extents of the brains.

219
Extended Data Fig. S5: Representative images of second-generation monosynaptic tracing of
inputs to dopaminergic midbrain neurons using RV∆GL-Flpo. See Fig. 1 for explanation of
experimental design and for quantification. **a,** 3 weeks, no doxycycline, **b,** 3 weeks, with doxycycline, **c,**
5 weeks, with doxycycline. Control conditions shown here: **d,** first-generation (∆G) system, **e,** no L, **f,** no
G, **g,** no Cre, no doxycycline, **h,** no Cre, with doxycycline. Note that omitting L gave very similar results
to omitting G.
226

228
Extended Data Fig. S6: Representative images of second-generation monosynaptic tracing of
inputs to parvalbumin-expressing cortical interneurons. See Fig. 1 for explanation of experimental
design and for quantification. **a,** 3 weeks, no doxycycline, **b,** 3 weeks, with doxycycline (note that there
is still blue fluorescence in this image, suggesting that, at one week after the mice were switched to dox
food, the doxycycline has not completely suppressed G and presumably L expression), **c,** 5 weeks, with
doxycycline. Control conditions shown here: d) first-generation (ΔG) system, e) no L, f) no G, g) no Cre,
no doxycycline, h) no Cre, with doxycycline. Again, omitting L and omitting G gave very similar results.
236

237   *(images in external file)*
238
239   **Extended Data Fig. S7: Whole-hemisphere image series of labeled inputs to dopaminergic**
240   **neurons using first- and second-generation rabies viral vectors expressing Cre.** Images are tiled
241   confocal images of every sixth 50-µm parasagittal section spanning the injected hemispheres.

242
243
244 **Extended Data Fig. S8: Examples of extrinsic inputs to dopaminergic cells labeled with first- and**
245 **second-generation RV vectors expressing Flpo and Cre.**
246

247
248
Extended Data Fig. S9: Counts of dopaminergic and parvabuminergic source cells and ratios of
striatal and thalamic cells to starting cells in Cre mice. Counts are totals found in every sixth 50-μm
section spanning each brain. **a-b,** Results using RVΔGL-Flpo(EnvA) in DAT-IRES-Cre. **a,** Counts of
source cells (left), defined as cells coexpressing tdTomato (marking activity of Flpo) and mTagBFP2
(marking expression of G), and ratios of these numbers of source cells to numbers of tdTomato+ cells
in striatum (right). Note that the low numbers of source cells in the second-generation conditions with
doxycycline could reflect either cell death or simply the intended suppression of mTagBFP2 expression
in these mice. Note also that there is no graph for the ratio of labeled striatal cells to source cells for the
"5 weeks dox (inj + food)" because no source cells were found for that condition (i.e., doxycycline
appears to have fully suppressed mTagBFP2 expression). **b,** Counts and ratios of starter cells, defined
here as cells coexpressing tdTomato (marking activity of Flpo) and EGFP (marking expression of TVA
and tTA). **c-d,** Results using RVΔGL-Flpo(EnvA) in PV-Cre. **c,** Counts and ratios of starting cells,
defined here as cells coexpressing tdTomato (marking activity of Flpo) and mTagBFP2 (marking
expression of G). Note that again there is no graph for the ratio of labeled contralateral cells to source
cells for the "5 weeks dox (inj + food)" because no source cells were found for that condition. **d,** Counts
and ratios of starting cells, defined here as cells coexpressing tdTomato (marking activity of Flpo) and
EGFP (marking expression of TVA and tTA).
266

267
268
Extended Data Fig. S10: Counts of dopaminergic and parvabuminergic source cells and ratios
of striatal and thalamic cells to starting cells using RVΔGL-Cre(EnvA) in DAT-P2A-Flpo mice.
Counts are totals found in every sixth 50-µm section spanning each brain. **a,** Counts of source cells
(left), defined as cells coexpressing tdTomato (marking activity of Cre) and mTagBFP2 (marking
expression of G), and ratios of these numbers of source cells to numbers of tdTomato+ cells in striatum
(right). Note that the low numbers of source cells in the second-generation condition with doxycycline
could reflect either cell death or simply the intended suppression of mTagBFP2 expression in these
mice. **b,** Counts and ratios of starter cells, defined here as cells coexpressing tdTomato (marking
activity of Cre) and EGFP (marking expression of TVA and tTA).
278

279
280
Extended Data Fig. S11: Second-generation rabies virus encoding Flpo does not kill most
labeled neurons for at least 16 weeks. a, Representative images of longitudinal two-photon structural
imaging fields of view (FOVs) from one week (left) to 4 weeks (right) after injection of a first-generation
rabies viral vector encoding Flpo (RVΔG-Flpo). The viruses in this experiment were coated with the
native RV glycoprotein for direct (TVA-independent) infection of neurons. Images are of the same FOV
at different time points. This first-generation virus killed almost all infected cells in this FOV within 4
weeks: only two labeled cells (circled) survived until the 4-week timepoint. Scale bar: 100 μm. **b,**
Representative images of longitudinal two-photon structural imaging FOVs from one week (left) to 12
weeks (right) after injection of a second-generation rabies viral vector encoding Flpo (RVΔGL-Flpo).
Images are of the same FOV at different time points. All labeled cells are still present at 16 weeks, the
last timepoint of imaging. Two example labeled cells are circled as fiducial markers. Scale bar: 100 μm.

c, Fraction of visible labeled cells over time, relative to the number visible at one week after RVΔG-Flpo
or RVΔGL-Flpo injection; connected sets of dots represent counts obtained from the same FOV within
the same mouse at the different time points (for RVΔG-Flpo: 1 week to 4 weeks; for RVΔGL-Flpo: 1
week to 16 weeks). Cells infected by RVΔG-Flpo (red) have almost entirely disappeared by 4 weeks
postinjection. The number of cells labeled by RVΔGL-Flpo (cyan) increases up to 4 weeks postinfection
and then remains constant for as long as the brains were imaged.

Extended Data Fig. S12: Additional examples of longitudinal two-photon imaging of second-generation monosynaptic tracing *in vivo*. a-b, Example images of tdTomato-labelled neurons at the injection site in V1 over all imaging sessions, from 2 days to 12 weeks, in two mice. The 'No dox' mice (**a**) were fed with regular food throughout, while the 'Dox' mice (**b**) were fed with food containing doxycycline (200 mg/kg) starting at two weeks after RV injection until perfusion at week 12, in order to suppress expression of the rabies virus polymerase and glycoprotein genes. White arrows show example cells that are last seen at that time point, with green arrows indicating the former positions of those now-missing cells at the next time point. In these examples, a total of 20 out of 141 cells were lost in the 'No dox' mouse, whereas 5 out of 98 cells were lost in the 'Dox' mouse. Scale bar: 100 µm. **c-d,** Counts of tdTomato-labeled cells appearing at each timepoint in individual mice. Each connected set of dots represents the numbers of the cells that appeared at one timepoint that are still present at the subsequent timepoints.

Extended Data Fig. S13: Titration of jGCaMP7s AAV concentration to minimize toxicity from
jGCaMP7s overexpression. a, Representative two-photon images of neurons expressing tdTomato
and jGCaMP7s, with the Flp-dependent AAV1 expressing jGCaMP7s injected at a higher concentration
(2.15x diluted from maximum), with doxycycline administered. Top row: 2 weeks after injection. Bottom
row: the same FOV at 8 weeks after injection. Numerous cells visible at 2 weeks are no longer seen at
8 weeks. Scale bar: 100 um. **b,** Representative two-photon images from another mouse, in which the
Flp-dependent AAV1 expressing jGCaMP7s had been injected at a lower concentration (4.64x diluted
from maximum), with doxycycline administered. Top row: 2 weeks after injection. Bottom row: the same
FOV at 8 weeks after injection. Almost all cells seen at 2 weeks are still present at 8 weeks. Scale bar:
100 μm. **c,** Counts of jGCaMP7s- and tdTomato-expressing neurons infected with high-concentration
(left graph) and low-concentration (middle graph) jGCaMP7s AAV. Each data point was obtained from
one FOV from each animal. Virus dilution factor and doxycycline condition are marked on the top of
each line; for example, '10x, Dox' means the mouse received the 10-fold dilution of jGCaMP7s AAV
and doxycycline food two weeks after the second injection. The surviving cell percentage is shown in
the right graph; there is a significant decrease in the high concentration of jGCaMP7s group
(***$p=0.00036<0.001$, one-way ANOVA test).

328
329
Extended Data Fig. S14: Additional examples of two-photon functional imaging of labeled cells
in mouse visual cortex. a, Representative two-photon images of neurons expressing jGCaMP7s and
tdTomato at 3 weeks. Scale bar: 100 µm. **b,** Direction tuning curves of three different jGCaMP7s-
expressing neurons obtained with drifting gratings presented at 12 directions of motion and 5 temporal
frequencies (TF), repeated 10 times (mean ΔF/F ± s.e.m.) at 8 weeks; color coding and axis labels are
as in Fig. 4. **c,** Single-cell fluorescence time courses for 85 cells over the first 480 s of visual
stimulation, showing robust spontaneous and evoked activity. Scale bar: 50 s.

337
338
Extended Data Fig. S15: jGCaMP7s signals and tuning curves of two example V1 neurons over
multiple imaging sessions. a, jGCaMP7s signals and tuning curves of a "week 1" cell (i.e., in which
jGCaMP7s and tdTomato fluorescence could be detected seven days after RVΔGL-Flpo(EnvA) and
AAV1-syn-F14F15S-jGCaMP7s injection). These data were obtained with drifting gratings presented at
12 directions of motion and 5 temporal frequencies (TF), repeated 10 times (tuning curve: mean ΔF/F ±
s.e.m; jGCaMP7s signals: mean ΔF/F) in three different imaging sessions (left: week 1; middle: week 5;
right: week 10). **b,** jGCaMP7s signals and tuning curves of a "week 2" cell (the same cell as used for
Fig. 4d).

347
348
Extended Data Fig. S16: Example confocal images of injection site and input regions in mouse
used for longitudinal functional calcium imaging. See Fig. 4a for diagram of experimental design.
These images show that inclusion of the jGCaMP7s AAV did not prevent successful monosynaptic
tracing. **a-d,** Injection site in V1. **a,** tdTomato, **b),** jCaMP7s, **c,** mTagBFP2, **d,** merge. Scale bar: 100
µm, applies to all panels. **e-p,** Inputs to parvalbumin-expressing V1 neurons are found in many different
brain regions: other visual areas (**e-g**, **i**, **o-p**), contralateral cortex (**h**), somatosensory areas (**j-k**),
thalamus (**l**), and retrosplenial areas (**m-n**). VISa, anterior visual area; VISam, anteromedial visual
area; VISp: Primary visual area; VISpm: Posteromedial visual area; CC-VISam: Contralateral cortex -
Anteromedial visual area; SSs: Supplemental somatosensory area; SSp: Primary somatosensory area;
LP: Lateral posterior nucleus of the thalamus; RSPd: Retrosplenial area, dorsal part; RSPv:
Retrosplenial area, ventral part; VISal: Anterolateral visual area; VISrl: Rostrolateral visual area; VISli:
Laterointermediate visual area; VISl: Lateral visual area.

BD FACSDiva 8.0

361
362 **Extended Data Fig. S17: Representative example of FACS data for titering viruses.** This example,
363 included as stipulated by the Nature Portfolio Reporting Summary, shows FACS analysis data for one

dilution of RV∆GL-Flpo(EnvA), used to infect HEK 293T cells which were immunostained for the rabies virus nucleoprotein using a blend of FITC-conjugated monoclonal antibodies (see Methods). The middle histogram in the second row shows the characteristic bimodal distribution (less distinct with second-generation RV vectors than with first-generation ones) with the uninfected cells in the mode on the left and infected ones in the one on the right. Gates are set by comparison with negative control wells of uninfected cells.

Supplementary File S1: Counts and statistics for corticostriatal experiments.

Supplementary File S2: Counts and statistics for DAT-IRES-Cre experiments.

Supplementary File S3: Counts and statistics for PV-Cre experiments.

Supplementary File S4: Counts and statistics for DAT-P2A-Flpo experiments.

Supplementary File S5: Counts for ∆G-Flpo and ∆GL-Flpo comparison.

Supplementary File S6: Counts and calculations for estimation of starting cell survival rates.

Supplementary File S7: Counts of tuned and untuned cells in functional imaging experiments.