Large genetic divergence underpins cryptic local adaptation across ecological and evolutionary gradients

Morgan M. Sparks[1,*], Joshua C. Kraft[2], Kliffi M. Subida Blackstone[1], Gordon G. McNickle[2,3], Mark R. Christie[1,4]

[1]Department of Biological Sciences, Purdue University; 915 W. State St., West Lafayette, Indiana 47907-2054 USA

[2]Department of Botany and Plant Pathology, Purdue University; 915 W. State St., West Lafayette, Indiana 47907-2054 USA

[3]Purdue Center for Plant Biology, Purdue University; 915 W. State St., West Lafayette, Indiana 47907-2054 USA

[4]Department of Forestry and Natural Resources, Purdue University; 715 W. State St., West Lafayette, Indiana 47907-2054 USA

*corresponding author: Department of Biological Sciences, Purdue University; 915 W. State St., West Lafayette, Indiana 47907-2054 USA; msparks1309@gmail.com

Keywords: countergradient variation, cogradient variation, genetic assimilation, maladaptive plasticity, perfect compensation, environmentally covarying local adaptation

Abstract

Environmentally covarying local adaptation is a form of cryptic local adaptation in which the

covariance of the genetic and environmental effects on a phenotype obscures the divergence

between locally adapted genotypes. Here, we systematically document the magnitude and drivers

of the genetic effect (V_G) for two forms of environmentally covarying local adaptation: counter-

and cogradient variation. Using a hierarchical Bayesian meta-analysis, we calculated the overall

effect size of V_G as 1.05 and 2.13 for populations exhibiting countergradient or cogradient

variation, respectively. These results indicate that the genetic contribution to phenotypic

variation represents a 1.05 to 2.13 standard deviation change in trait value between the most

disparate populations depending on if populations are expressing counter- or cogradient

variation. We also found that while there was substantial variance among abiotic and biotic

covariates, the covariates with the largest mean effects were temperature (2.41) and gamete size

(2.81). Our results demonstrate the pervasiveness and large genetic effects underlying

environmentally covarying local adaptation in wild populations and highlights the importance of

accounting for these effects in future studies.

Introduction

How and when populations can persist via adaptation to local environmental conditions remains a central question in ecology and evolutionary biology [1]. Environmental gradients, where abiotic and biotic factors such as temperature, salinity, or predator type vary in a predictable manner, provide a framework to formulate and test questions about the pattern and magnitude of local adaptation. Pairing environmental gradients with common garden or reciprocal transplant studies, especially when there are many populations distributed along these gradients, can lead to insights into how local adaptation covaries with specific environmental factors [2,3].

Shared patterns of phenotypic variation along ecological gradients spurred notable research on the environmental and genetic determinants of body size and morphology and led to the establishment of eco-geographical "rules" such as Bergmann's, Allen's, and Hesse's rules and subsequent derivations therein [4–9]. Foundational work by Levins [2,10] revealed a pattern whereby fruit flies distributed along an elevation gradient expressed a similar body size in their wild home environments, leading to the conclusion that the trait was canalized. However, a subsequent study revealed the higher elevation populations were much smaller than lowland populations in common environments—expressing disparate phenotypes across all common environments, a response believed to be related to desiccation—suggesting these populations were locally adapted to different elevations. In concert, these findings provided the first formal evidence of environmentally covarying local adaptation.

The variance components of phenotypic plasticity are commonly expressed with reaction norm plots, where the trait value of a genotype is plotted as a function of multiple environmental values [11]. In a reaction norm, the slope of a genotype's response is representative of phenotypic plasticity (the effect of the environment) and the difference between genotype-

specific reaction norms at each environmental variable is representative of the genotypic

difference (the genetic effect). When different genotypes express the same reaction norm (*i.e.*,

overlapping lines, Fig. 1a, plasticity alone) this is representative of a purely environmentally

driven response where distinct genotypes would express the same trait values (or phenotype)

along an environmental gradient. Alternatively, populations may be locally adapted, such that the

reaction norms—namely, fitness or fitness related traits—cross when comparing populations in

their respective home and away environments, which is indicative of a genotype by environment

interaction (Fig. 1b, Genetic x Environment). Finally, populations may exhibit a form of cryptic

local adaptation, where the genetic and environmental effects covary with one another, either

negatively or positively [12]. In this scenario, the reaction norms for populations may have the

same or similar slopes thus maintaining consistent rank order in their trait values (Fig.1, co- and

countergradient variation). For example, a high latitude population may have consistently larger

body sizes across environments than a low latitude population, despite both expressing plasticity

in body size. When the genetic portion of phenotypic variance opposes the environmental

portion, it is called countergradient variation (negative covariance; Fig. 1c), and when those two

factors act in the same direction it is termed cogradient variation (positive covariance; Fig. 1d)

[2,13]. Thus, when comparing among wild populations, phenotypic traits appear more similar

than they would if populations were compared in a common environment for countergradient

variation, and vice versa for cogradient variation.

Counter- and cogradient variation constitute forms of cryptic local adaptation because

their true genetic effect on phenotypic variation is masked by the environment [12–15] (Fig.1).

Countergradient variation can be particularly difficult to identify in wild populations because the

environmental influence on a phenotype can conceal the genetic influence, such that the signal of

local adaptation only appears when individuals are reared in common environments. This

response has been termed perfect compensation (or exact compensation), when wild-type

phenotypes are equal across a gradient in their home environments (Fig. 1c) (*sensu* Conover *et*

al. [12]). Compensation can be thought of as a measure of how similar phenotypic values are

across populations in their home environments—and as an indicator of whether selection may

favor similar or disparate phenotypes along an environmental gradient.

While Levins [10] was the first to formally describe countergradient variation—which

was followed by early research by Berven and others [16,17] in *Ranid* frogs—it wasn't until two-

and-a-half decades later that interest in environmentally covarying local adaptation grew

substantially. Research with an eastern North American fish, Atlantic silverside (*Menidia*

menidia), revealed that populations had similar body sizes along a latitudinal gradient that

spanned from the Gulf of Maine to the Floridian Atlantic Coast. Using a common garden

approach, the researchers found that the northern populations were consistently larger than

southern populations, regardless of the temperature in which they were raised (*e.g.*, Fig. 1c). This

finding led to substantial follow up research investigating the proximate and ultimate factors

driving the trend [18,19], namely faster growth and more efficient metabolism in response to

size-dependent winter mortality. A theoretical review by the group outlined the evolutionary

significance of countergradient variation and described 20 prior studies with results consistent

with countergradient variation, though generally without the original authors formally describing

their observations as such [13]. The early work highlighted in that review and subsequent

research substantiate a large body of evidence for co- and countergradient variation across

diverse taxa including plants [20], bivalves [21], fishes [22–25], and insects [26–28], and across

ecological gradients such as temperature [29,30], salinity [31], carotenoid availability [32,33],

and urbanization [34].

Despite the attention paid to counter- and cogradient variation, there have been few

studies that describe the magnitude and extent of these types of local adaptation, nor has there

been much investigation into the broader abiotic and biotic factors that might contribute to their

occurrence [12,35,36]. Without understanding the extent of environmentally covarying local

adaptation, it is difficult to make predictions about how locally adapted populations may interact

with a changing environment or how cryptic local adaptation may aid or hinder conservation

goals. Here we use a Bayesian meta-analytical framework with hierarchical models to describe

the general effect size of local adaptation resulting from counter- and cogradient variation as they

appear in the literature and to investigate what biotic and abiotic factors are associated with the

magnitude of adaptation. The effect sizes we calculate represent a direct estimate of the standard

deviation of the mean difference in trait values for the local populations being measured–or more

generally, they are the genetic effect of environmentally covarying local adaptation. With respect

to the classic representation of phenotypic variance [37]

$V_P = V_E + V_G + V_{(GxE)} + 2(cov_{GE})$ (1)

this study estimates V_G or the total phenotypic variance driven by genetic differences in

populations that exhibit environmentally covarying local adaptation (*e.g.*, $cov_{GE} < 0$;

countergradient variation, or $cov_{GE} > 0$;cogradient variation, Fig. 1c,d; Fig. S1).

Counter- and cogradient variation as phenomena, represent the negatively or positively

covarying contributions of V_E and V_G in equation 1 (cov_{GE}). To directly measure this covariation

studies are limited to reciprocal transplants or common gardens that recreate home environments

106 [36,38]. Additionally, while cov_{GE} may be large, little research has explored how the different

107 phenotypic variance components in equation 1 are structured along gradients in wild populations

108 [38]. While V_G represents both the full effects of genetic change on phenotypic variance

109 (including drift and gene flow), clinally covarying V_G of fitness related traits covary specifically

110 because of selection and the measure of V_G in this specific scenario is a suitable representation of

111 the magnitude of local adaptation in these populations. As such, the goals of our analyses were to

112 *i)* quantify the genetic variance component of countergradient and cogradient variation as an

113 effect size, *ii)* describe these effect sizes within and across taxa, phenotypic traits, and different

114 gradient types, and *iii)* quantify the effect size of compensation—or how similar countergradient

115 populations are in their home environments—to assess if there are clear trends for phenotypic

116 similarity or dissimilarity along gradients.

117 **Methods**

118 *Literature Review and Selection*

119 Overall, we collected 858 studies that investigated counter- and/or cogradient variation. To

120 generate a full list of studies, we first analyzed studies indicated as showing counter- or

121 cogradient variation in the qualitative analysis from Conover *et al.* [12]. We then searched the

122 Web of Science topic field using the terms "counter*gradient variation" in May of 2018 and

123 "co*gradient variation" (while excluding the previous search, as the wildcard renders them

124 redundant) in June of 2019, resulting in 384 and 34 results, respectively. In August 2019, we also

125 included studies citing the Conover *et al.* [12] review, as well as the earlier review article by

126 Conover & Schultz [13]—682 studies total (some of which were redundant with studies in the

127 prior search resulting in a total of 858 studies). These methods are discussed in more depth in the

128 Supplementary Methods.

To be further included in our analysis, a study had to meet a set of qualifying criteria.

Specifically, a study had to be designed to include some form of a common environment—this

may have been in a common garden study or a reciprocal transplant using wild populations or

populations recently brought into lab conditions. We did not include examples from domestic

populations or lab induced selection. Additionally, studies must have included two or more

common environments to determine the signal of local adaptation (countergradient, cogradient,

or others such as GxE that were ultimately not included in this study), as well as two or more

populations to compare the genotypic difference of the phenotypic response between

populations. Finally, studies were determined to show counter- or cogradient based on plotted

reaction norms to verify environmental and genetic effects covaried—*i.e.*, populations in away

environments minimized (countergradient) or maximized genetic variance (cogradient) (positive

or negative covariance, Fig. 1c,d)—and genotypes maintained their rank order across

environments (Fig. S1). Occasionally, studies investigated these distinctions across multiple

species in the place of populations and such studies were discarded.

Statistical Methods

To calculate effect sizes, we took data from both tables and figures in manuscripts. For data in

the form of figures, we used the software WebPlotDigitizer [39] to extract the relevant data.

Briefly, the program works by importing a photo of the respective figure and the user manually

sets the scales for the x- and y-axes based on the figure values and then adds digital points with

their cursor to the data of interest. The digital points are then translated to numeric data via the

imaging algorithm.

The statistic used to calculate effect size was Hedge's *d*, which, for this study, was a

standardized mean difference with a correction for small sample sizes[40,41] to avoid bias in the

test statistic [42,43]. We calculated standardized mean difference (synonymous with effect size

in this paper) by comparing the most disparate populations at each treatment level. For example,

if we had three populations collected from an elevational gradient at 500m, 1000m, and 1500m

in common gardens with three temperature treatments (*e.g.*, 10, 15, and 20 °C, as was often the

case where latitude or elevation were proxies for temperature clines), then we would collect the

mean and variance for every population in every treatment, but calculate the effect size

(standardized mean difference) by using the values from the 500m and 1500m populations at

each treatment. Occasionally, not all populations were used in all treatments, in these scenarios

we used only the treatments for the most extremely distributed (with respect to the gradient—

e.g., very highest and lowest elevations) populations. Some studies conducted multiple

experiments measuring the same trait. If multiple experiments measuring the same trait were

recorded, we calculated the effect sizes for each.

In all, we calculated 422 total effect sizes from 1204 individual data points extracted

from 83 studies. Many studies showed counter- or cogradient patterns in multiple traits and/or

trials within a given study. We collected all these data and controlled for their non-independence

using a robust, Bayesian hierarchical modeling approach (discussed below). Hedge's *d* was

calculated using the *metafor* package in R (version 3.6.3) [44] with the *escalcl()* function [45].

Additional metadata was collected from each study and included features such as year the study

was published, the species investigated, and the distance in meters (for elevational gradients) and

kilometers (for latitudinal gradients). The full list of these covariates is available in our

supplemental metadata (Supplementary Data) and were collected to be used as covariates in the

mixed-effects model (Table 1).

To estimate overall effect size and the effect of covariates, Bayesian linear mixed models

fit with random intercepts were used in the R package *brms* [46,47], which is a high-level

interface in R for the Stan modeling language [48]. Linear mixed-effects models were chosen

due to the hierarchical nature of this data set, where a study may have effect sizes estimated for

multiple traits in multiple experiments. For this approach, we fit the multiple estimates of effect

size for a trait in an experiment by using an estimate from each experimental level (Fig. S1c)—

for instance, using the prior example, an estimate for each of 10, 15, and 20 °C. This approach

also allows each treatment to have different effect sizes and accounts for variability driven by

Genotype x Environment interactions ($V_{(GxE)}$ eq.1). As such, nested random intercepts effect was

used to control for this non-independence. We originally set out to nest traits in experiments in

studies in our random effects but model parameterization necessary to overcome divergent

transitions (specifically, max tree depth in Stan) extended model run time to be computationally

intractable. Random effects were instead reduced to traits nested in studies.

Similarly, phylogenetic relationships may provide another source of non-independence,

and was controlled for by using a phylogenetic random intercepts effect for which we created a

variance-covariance matrix [49]. To do so, we generated a pseudo-phylogeny using the *taxize*

package in R [50]. Using the *classification()* function, we extracted the full taxonomy of a given

organism in our dataset from the NCBI database. We then used the *class2tree()* function to

generate a phylogenetically accurate tree in terms of taxonomic rank (*e.g.*, all species within a

genus would have the same distance from one another, all genera within a family would have the

same distance—a full explanation of the method is available in the Supplementary Methods).

The function not only clusters taxa on their specific taxonomic rank but also within the

taxonomic clade, allowing for more accurate distance measures. The tree created using this

method was transformed into a matrix and then used as a custom variance-covariance component

in our hierarchical model. The mathematical representation of our random effects in all models is

shown in our Supplementary Materials (Supplementary Methods). We used moderately

informative priors for our hierarchical models [51]—except for the metaregression, where a

stronger prior was used to rein in variance—lists of priors and discussion of their role in

controlling for potential bias are available in Supplementary Methods (Tables S1-S4, diagnostic

plots Figures S2-5). Because we used a Bayesian framework, the following effects are presented

with the means and 95 percent credible intervals (indicated as 95% CI) of the posterior

distributions, unless otherwise noted.

The overall effect size for both co- and countergradient variation was estimated using

random-effects only models, which is the convention when assuming study effects may vary in

meta-analytical approaches [52]. We included the covariates of trait, gradient, and taxonomic

class in our metaregression countergradient model to assess how different biotic and abiotic

factors may contribute to the estimate of countergradient variation genetic variance effect size.

Compensation Analysis

Finally, we analyzed whether populations appeared to be under-, over-, or perfectly

compensating (*sensu* Conover *et al.* [12]) when comparing their home environment phenotypes.

To do so, we used the most extreme treatments and paired them with the polarly (as in opposites,

not necessarily geographically) distributed populations used in the study. For example, if we

consider three populations of fish from a range of saline conditions—freshwater, brackish, and

marine—then, the most saline treatment in the experiment would be considered the home

environment of the marine population and the least saline the home environment of the

freshwater fish. We then compared the trait values of the two polarly distributed populations and,

using the previous example, if the trait value was larger for the most saline population than that

of the least saline population the relationship was classified as overcompensating, close to even

was classified as perfectly compensating, and negative was classified as undercompensating. To

standardize these data, we again used Hedge's d, because the values of different traits are

generally measured on different scales [53]. Because there are no quantitative definitions of

compensation, our results were conservatively classified as overcompensating if values were >

0.5, perfectly compensating if values were between -0.5 and 0.5, and undercompensating if

values were < -0.5. Because compensation measurements were computed as Hedge's $d,$ these

were meant to reflect a broad distribution around 0 for perfect compensation (what amounts to a

medium effect size in both positive and negative directions, [54]) that encapsulates the inherent

measurement error in the value.

Results

Random Effects Models

The result of our random effects models that estimated overall effect size for countergradient

variation was 1.05 (95% CI = 0.30-2.49) and for cogradient variation 2.13 (95% CI = 0.35-7.00)

(Fig. 2, non-truncated in Fig. S6). The heterogeneity parameter τ—the full random effect

standard deviation—was modeled with its own prior distribution and resulted in 2.19 (95% CI =

1.79-2.82) for countergradient variation and 9.33 (95% CI = 4.00-22.45) for cogradient variation.

We similarly included a random effect for phylogenetic relatedness which resulted in an estimate

of 2.54 SD (95% CI = 1.04-5.28) for countergradient variation and 3.71 SD (95% CI = 1.03-

14.82) for cogradient variation (Tables S5, S6).

Metaregression

We also implemented a mixed effects metaregression with our countergradient dataset to estimate the effect size for trait classification, gradient classification, and taxonomic class effect. The mean effect, posterior distribution and 95% credible intervals for each modeled covariate are presented in Table 1 and those same values are represented as the main effects estimated marginal means with the posterior distribution and credible intervals in Figure 3 (non-truncated in Fig. S7). For each covariate, these results illustrate similar mean values with large and overlapping credible intervals. While variability was large across all covariates, the mean coefficient estimates of the posterior distributions for temperature (mean = 2.41, 95% CI = 0.70-6.15) and gamete size (mean = 2.81, 95% CI = 0.52-8.68) were much larger than the global estimate of 1.19. By contrast, the estimate for urbanization (mean = 0.74, 95% CI = 0.13-2.46)—had lower credible intervals near zero and was much lower than other estimates. These values are represented as main effect marginal means in Figure 3. The among study standard deviation was 1.97 (95% CI = 1.22-2.79), while the among trait (nested in study) standard deviation was 2.26 (95% CI = 1.82-2.91), and the standard deviation for phylogenetic random effect was 3.41 (95% CI = 1.13-6.46), indicating more variance was accounted for among traits in studies and by phylogeny, than among studies.

Compensation Analysis

The results of our compensation analysis were plotted to represent the conceptual relationship between home phenotypes of populations exhibiting countergradient variation (*sensu* Conover *et al.* [12]) (Fig. 4). Our data provide support for far more frequent over- and under compensation than perfect compensation, meaning that populations in our study generally have divergent home phenotypes as opposed to identical phenotypes along a gradient. We considered any value

between -0.5 and 0.5 as perfectly compensating (or a range of 1 effect size around 0 for a

conservative estimate), values less than -0.5 undercompensating, and values larger than 0.5

overcompensating. We consider this classification conservative because it allows for a medium

effect size in the positive or negative direction from zero [52], despite the theoretical description

of perfect compensation being a value equal to zero [12]. The distribution of these values in our

data set were overcompensating = 38.6%, undercompensating = 36.0%, and perfectly

compensating = 25.4%. Effect sizes of compensation ranged from -70.38 to 36.45 but 5% and

95% quantiles were -27.74 and 6.49, respectively.

Discussion

The ecological and evolutionary importance of co- and countergradient variation has been

recognized for three decades [13,18]. Here, we find that the genetic effect of populations

expressing counter- and cogradient variation are 1.05 and 2.13, respectively. Our measure of

standardized mean difference, Hedges' *d,* translates the difference between populations into units

of standard deviation. Thus, our results indicate that the standardized mean difference between

the most disparate populations in our study—*i.e.*, the genetic effect of environmentally covarying

local adaptation—is between one and two standard deviations change in trait value. In other

words, populations that express the same trait value in their home environments would diverge

by one or two standard deviations in a common environment due to their genetic differences

depending on whether they were expressing counter- or cogradient variation, respectively.

Additionally, our results suggest populations expressing cogradient variation may express twice

the genetic divergence as those expressing countergradient variation, but the results for

cogradient variation are based on small sample set with significant variance. Our findings

demonstrate that the genetic divergence among locally adapted, clinally distributed populations

is large but also obscured as a function of the covariance between environmental and genetic

effects and their interaction with phenotype.

We used a metaregression to investigate the effect sizes of abiotic and biotic covariates of

countergradient variation. Covariates were statistically indistinguishable from one another due to

high variance in their estimates, though many showed mean values that were large with

distributions away from 0 (Effect sizes estimates of 1-3, Table 1, Fig. 3). Particularly, we

estimated large effect sizes for gamete size (2.81), and temperature (2.41). For gamete size,

multiple studies in fishes showed effects greater than 2.5 and as high as 13.41 [55]. The tradeoff

between size and number of eggs produced by female fishes is a well-established trend in

ecology [56,57], and particularly in marine fishes where there are strong links between egg size

and latitude [58]. The large effect size estimate for gamete size from our metaregression then is

less surprising given the strong selection on gamete size across species and environments, but

also suggests that despite the tradeoff between size and number of eggs there is benefit to larger

eggs moving poleward. While there were additional large study-specific effects for temperature

(4 instances > 4.5), we separated temperature and temperature related gradients based on the

explicit distinction made by authors. Temperature is a fundamental selective agent for wild

organisms, while latitudinal, elevational, and shade cover gradients serve as proxies for

temperature—all three had estimates at or near one standardized mean difference. The proxy

measures then likely fail to capture the full evolutionary dynamics driving environmentally

covarying local adaptation at the same resolution as that caused by temperature.

As far as the variability in our metaregression estimates, we find that the inter-effect

variability (τ^2, or the variability of our random effects) generally swamped the covariate-specific

effects estimates (Table 1). The variability in this random effect is also a measure of GxE (the

variability of multiple effect sizes for a trait in a study, Fig. Sc) and indicates that when

environmentally covarying local adaptation is present, this adaptation corresponds with a large

population-specific (or genotype-specific) GxE influence. While the lack of predictive results in

our metaregression may be indicative that our data is not robust enough for differences among

covariates to emerge, we think it is more likely that these results suggest that countergradient

variation, when it does occur, is an ecologically ubiquitous phenomenon for species distributed

along environmental gradients. Furthermore, we suspect that species or populations experiencing

similar agents of selection—for example, those along latitudinal, elevational, or temperature

gradients—may achieve adaptation through different mechanisms (as indicated by the large GxE

influence) and therefore contributing to the variability observed in our models. In other words,

while the phenotypic solution may appear similar across species and gradients, the processes

driving the genetic divergence of the phenotypes of different populations is idiosyncratic. The

conclusion that covariate specific effects are idiosyncratic is further supported by the lack of

pattern with respect to latitudinal distance and temperature difference regressions analyses,

neither of which were predictive (Fig. S8, Table S7). Thus, while populations almost always

showed a large effect of environmentally covarying local adaptation—metaregression estimates

were 2.41 and 0.99 for temperature and latitudinal gradients, respectively—the magnitude did

not scale with temperature or physical distance based on separate analyses (Fig. S8, Table S7).

This may be due in part to microgeographic variation—in which large adaptive phenotypic

divergence occurs over small spatial scales—as multiple studies reviewed as part of this work

indicated microgeographic variation as driver of phenotypic patterns across the landscape

[59,60].

Even though the overall effect size of cogradient variation was large, only 15 out of 858

studies were consistent with cogradient variation and included enough information to calculate

effect sizes. This paucity may reflect an overall lack of published research investigating and

demonstrating cogradient variation (see also, [12]). However, in a meta-analysis of reciprocal

transplant studies, Stamp & Hadfield [36] found 60% of traits exhibited a cogradient pattern. Our

results (*i.e*., the magnitude of effect size), in conjunction with the work of Stamp & Hadfield [36]

suggest that, while rarely reported, cogradient variation constitutes an ecologically meaningful

phenomenon. We further hypothesize cogradient variation may suffer from underreporting

because it's natural sign mirrors environmentally induced plasticity that may not drive

researchers to investigate populations using common gardens. While cogradient variation may be

underreported, standalone cases such as that by Trussell & Etter [61] can show both the large

effect of cogradient variation and how home phenotypic differences can disguise that effect. In a

reciprocal transplant study, the researchers measured shell thickness in marine snail populations,

finding that, when in their home environments, populations expressed a 3.18 standardized mean

difference in shell thickness. But, when considered in shared environments, that difference

reduced to 2.07 units. Notably, this result amounts to a substantial effect size difference due to

genetic variation alone, but is ~2/3 value if the populations had only been observed solely in

their home environments. More importantly, if the researchers had only investigated these

populations in their home environments without common gardens, a likely inference for the

phenotypes would have been a strong plastic response, which, while true, did not account for

even half of phenotypic variation in home environments.

Our compensation analysis for countergradient variation revealed that populations in their

home environments were more commonly over- or undercompensasting, as opposed to perfectly

compensating (Fig. 4). In other words, the genetic effect of countergradient variation was

generally not equal to that of the environmental effect. This result was clear with conservative

bounds for perfect compensation (-0.5 to 0.5 standardized mean difference). Reducing those

bounds to the less conservative values of -0.2 to 0.2 (considered to be a small effect size [54]) for

perfect compensation resulted in only 11.7 percent of studies demonstrating perfect

compensation. While we caution that these results may not necessarily point to clear biological

trends about how common and in what scenarios over- and under-compensation may occur, it

does suggest that either: 1. selection for maintaining optimal phenotypes across broad

environmental gradients may not be the norm (*sensu* [12,13]), or 2. the response to selection may

not be strong enough to effectively minimize phenotypic variation across populations. If the

latter, the response to selection may be minimized by insufficient adaptive genetic variation or

countervailing selection on correlated traits specific to each local population [62].

Our results are indicative of a response to selection contributing to strong genetic

divergence in trait values distributed along environmental clines and reinforce the evolutionary

importance of counter and co-gradient variation. As such, negative or positively environmentally

covarying selection is a strong candidate as a mechanism for driving parapatric speciation—if

the selection pressure is consistent across the gradient—or peripatric speciation—if selection is

strongest at the poles. Genetic differentiation may also be expected by further anthropogenic

change, where habitat fragmentation breaks up corridors of gene flow within clines [63], or

climate change disrupts habitat envelopes at the polar ends. Both scenarios have significant

implications for how populations exhibiting environmentally covarying local adaptation are

managed. This clinal breakdown may also lead to underestimation of the importance of co- and

countergradient variation, as range-wide patterns may have already been disrupted causing

researchers to undercount their occurrence or underestimate their historical magnitude if the

edges of clines are lost. Additionally, environmentally covarying adaptation should be

considered in conservation and management efforts, as cryptic local adaptation may lead to

undesirable management outcomes if ignored, or good candidates for translocation (*e.g.*, genetic

rescue [64]) if well understood.

Conclusion

Our findings illustrate that environmentally covarying local adaptation is pervasive across

multiple gradients in a diverse set of taxa. With this meta-analysis, we quantitatively

characterized the genetic effect on phenotypic divergence for environmentally covarying local

adaptation and find that the result is one to two full standard deviations from the mean trait value

of a population. While the genetic effect of co- and countergradient variation may be large, the

variability of response observed across studies is also quite high, as indicated in the values of our

random effects, indicating strong GxE influences. Moreover, when analyzing the mean trait

difference between home phenotypes of populations exhibiting countergradient variation, we

find no support for a strong signal of perfect compensation, indicating that selection for optimal

phenotypes across a gradient can or does not perfectly minimize variation in mean phenotypic

trait values. Much like local adaptation more generally, environmentally covarying local

adaptation necessitates close consideration when working with populations distributed across

ecological gradients. We conclude that environmentally covarying local adaptation is ubiquitous

with a large effect size and argue that this form of cryptic local adaptation deserves continued

attention with theoretical, experimental, and genomic approaches.

Acknowledgements

We thank K. Lotterhos, N. Emery, and C. Oakley for providing helpful discussion regarding environmentally covarying local adaptation. K. Horvath offered useful suggestions about instituting a phylogenetic random effect in our models, D. Pascall provided valuable insights on implementing the pseudo-phylogeny, and R. Swihart gave useful direction on Bayesian model setup. We also thank A. Harder, A. Lee, and A. Nalesnik for their helpful comments on a past version of this manuscript. This work was supported, in part, by the Purdue Ecology and Evolutionary Biology Waser Fellowship to MMS and by funding from the National Science Foundation (OCE-1924505 ; DEB-1856710) to MRC.

References

1. Blanquart F, Kaltz O, Nuismer SL, Gandon S. 2013 A practical guide to measuring local adaptation. *Ecol. Lett.* **16**, 1195–1205. (doi:https://doi.org/10.1111/ele.12150)

2. Levins R. 1968 *Evolution in Changing Environments: Some Theoretical Explorations*. Princeton University Press.

3. Hargreaves AL, Germain RM, Bontrager M, Persi J, Angert AL. 2020 Local adaptation to biotic interactions: A meta-analysis across latitudes. *Am. Nat.* **195**, 395–411. (doi:10.1086/707323)

4. Bergmann C. 1848 *Über die Verhältnisse der Wärmeökonomie der Thiere zu ihrer Grösse*.

5. Allen JA. 1877 The influence of physical conditions in the genesis of species. *Radical review* **1**, 108–140.

6. Atkinson D, Sibly RM. 1997 Why are organisms usually bigger in colder environments?

Making sense of a life history puzzle. *Trends Ecol. Evol.* **12**, 235–239. (doi:10.1016/S0169-

5347(97)01058-6)

7. Blanckenhorn WU, Demont M. 2004 Bergmann and Converse Bergmann Latitudinal Clines

in Arthropods: Two Ends of a Continuum? *Integr. Comp. Biol.* **44**, 413–424.

(doi:10.1093/icb/44.6.413)

8. Gaston KJ, Chown SL, Evans KL. 2008 Ecogeographical rules: elements of a synthesis. *J.*

Biogeogr. **35**, 483–500. (doi:10.1111/j.1365-2699.2007.01772.x)

9. Tian L, Benton MJ. 2020 Predicting biotic responses to future climate warming with classic

ecogeographic rules. *Curr. Biol.* **30**, R744–R749. (doi:10.1016/j.cub.2020.06.003)

10. Levins R. 1969 Thermal Acclimation and Heat Resistance in Drosophila Species. *Am. Nat.*

103, 483–499. (doi:10.1086/282616)

11. West-Eberhard MJ. 2003 *Developmental Plasticity and Evolution*. OUP USA.

12. Conover DO, Duffy TA, Hice LA. 2009 The covariance between genetic and environmental

influences across ecological gradients: reassessing the evolutionary significance of

countergradient and cogradient variation. *Ann. N. Y. Acad. Sci.* **1168**, 100–129.

(doi:10.1111/j.1749-6632.2009.04575.x)

13. Conover DO, Schultz ET. 1995 Phenotypic similarity and the evolutionary significance of

countergradient variation. *Trends Ecol. Evol.* **10**, 248–252. (doi:10.1016/S0169-

5347(00)89081-3)

14. Schlichting CD. 2008 Hidden Reaction Norms, Cryptic Genetic Variation, and Evolvability.

Ann. N. Y. Acad. Sci. **1133**, 187–203. (doi:10.1196/annals.1438.010)

15. Kinnison MT, Hairston Jr. NG, Hendry AP. 2015 Cryptic eco-evolutionary dynamics. *Ann.*

N. Y. Acad. Sci. **1360**, 120–144. (doi:10.1111/nyas.12974)

16. Berven KA, Gill DE, Smith-Gill SJ. 1979 Countergradient selection in the green frog, *Rana*

clamitans. *Evolution* **33**, 609–623. (doi:10.1111/j.1558-5646.1979.tb04714.x)

17. Berven KA. 1982 The genetic basis of altitudinal variation in the wood frog *Rana sylvatica*

II. An experimental analysis of larval development. *Oecologia* **52**, 360–369.

(doi:10.1007/BF00367960)

18. Conover DO, Present TMC. 1990 Countergradient variation in growth rate: compensation for

length of the growing season among Atlantic silversides from different latitudes. *Oecologia*

83, 316–324. (doi:10.1007/BF00317554)

19. Conover DO, Arnott SA, Walsh MR, Munch SB. 2005 Darwinian fishery science: lessons

from the Atlantic silverside (*Menidia menidia*). *Can. J. Fish. Aquat. Sci.* **62**, 730–737.

(doi:10.1139/f05-069)

20. Chapin FS, Oechel WC. 1983 Photosynthesis, respiration, and phosphate absorption by

Carex aquatilis ecotypes along latitudinal and local environmental gradients. *Ecology* **64**,

743–751. (doi:10.2307/1937197)

21. Dittman DE. 1997 Latitudinal compensation in oyster ciliary activity. *Funct. Ecol.* **11**, 573–

578. (doi:10.1046/j.1365-2435.1997.00127.x)

22. DiMichele L, Westerman ME. 1997 Geographic variation in development rate between

populations of the teleost *Fundulus heteroclitus*. *Mar. Biol.* **128**, 1–7.

(doi:10.1007/s002270050062)

23. Imsland AK, Foss A, Nevdal G, Cross T, Bonga SW, Ham EA, Stefansson SO. 2000
Countergradient variation in growth and food conversion efficiency of juvenile turbot. *J.*
Fish Biol. **57**, 1213–1226. (doi:10.1111/j.1095-8649.2000.tb00482.x)

24. Billerbeck JM, Schultz ET, Conover DO. 2000 Adaptive variation in energy acquisition and
allocation among latitudinal populations of the Atlantic silverside. *Oecologia* **122**, 210–219.
(doi:10.1007/PL00008848)

25. Jonassen TM, Imsland AK, Fitzgerald R, Bonga SW, Ham EV, Naevdal G, Stefansson MO,
Stefansson SO. 2000 Geographic variation in growth and food conversion efficiency of
juvenile Atlantic halibut related to latitude. *J. Fish Biol.* **56**, 279–294. (doi:10.1111/j.1095-
8649.2000.tb02106.x)

26. Blanckenhorn WU. 1991 Life-history difference in adjacent water strider populations:
phenotypic plasticity or heritable responses to stream temperature? *Evolution* **45**, 1520.
(doi:10.2307/2409899)

27. Armbruster P, Conn JE. 2006 Geographic variation of larval growth in North American
Aedes albopictus (Diptera: Culicidae). *Ann. Entomol. Soc. Am.* **99**, 1234–1243.
(doi:10.1603/0013-8746(2006)99[1234:GVOLGI]2.0.CO;2)

28. Pöykkö H, Tammaru T. 2010 Countergradient vs. cogradient variation in growth and
diapause in a lichen-feeding moth, *Eilema depressum* (Lepidoptera: Arctiidae). *J. Evol. Biol.*
23, 1278–1285. (doi:10.1111/j.1420-9101.2010.01990.x)

29. Yamahira K, Conover DO. 2002 Intra- vs. interspecific latitudinal variation in growth:
Adaptation to temperature or seasonality? *Ecology* **83**, 1252. (doi:10.2307/3071940)

30. Kokita T. 2003 Potential latitudinal variation in egg size and number of a geographically

widespread reef fish, revealed by common-environment experiments. *Mar. Biol.* **143**, 593–

601. (doi:10.1007/s00227-003-1104-x)

31. Secor DH, Gunderson TE, Karlsson K. 2000 Effect of temperature and salinity on growth

performance in anadromous (Chesapeake Bay) and nonanadromous (Santee-Cooper) strains

of striped bass *Morone saxatilis*. *Copeia* **2000**, 291–296.

32. Craig JK, Foote CJ, Wood CC. 2005 Countergradient variation in carotenoid use between

sympatric morphs of sockeye salmon (*Oncorhynchus nerka*) exposes nonanadromous

hybrids in the wild by their mismatched spawning colour: spawning colour identifies hybrids

between sympatric morphs. *Biol. J. Linn. Soc. Lond.* **84**, 287–305. (doi:10.1111/j.1095-

8312.2005.00430.x)

33. Grether GF. 2005 Environmental change, phenotypic plasticity, and genetic compensation.

Am. Nat. **166**, 115–123. (doi:10.1086/432023)

34. Tüzün N, Beeck LO de, Brans KI, Janssens L, Stoks R. 2017 Microgeographic

differentiation in thermal performance curves between rural and urban populations of an

aquatic insect. *Evol. Appl.* **10**, 1067–1075. (doi:10.1111/eva.12512)

35. Hendry AP. 2017 *Eco-evolutionary Dynamics*. Princeton University Press.

36. Stamp MA, Hadfield JD. 2020 The relative importance of plasticity versus genetic

differentiation in explaining between population differences; a meta-analysis. *Ecol. Lett.* **23**,

1432–1441. (doi:https://doi.org/10.1111/ele.13565)

37. Falconer DS, Mackay TFC. 1996 Introduction to quantitative genetics. Longmans Green,

Harlow, Essex, UK.

38. Albecker MA, Trussell GC, Lotterhos KE. 2022 A novel analytical framework to quantify

co-gradient and countergradient variation. *Ecol. Lett.* **25**, 1521–1533.

(doi:10.1111/ele.14020)

39. Rohatgi A. 2020 *WebPlotDigitizer*. See https://automeris.io/WebPlotDigitizer.

40. Hedges LV. 1981 Distribution theory for Glass's estimator of effect size and related

estimators. *J. Educ. Behav. Stat.* **6**, 107–128. (doi:10.2307/1164588)

41. Rosenberg MS. 2013 Moment and least-squares based approaches to meta-analytic

inference. In *Handbook of Meta-analysis in Ecology and Evolution* (eds J Koricheva, J

Gurevitch, K Mengersen), pp. 108–124. Princeton University Press.

42. Hamman EA, Pappalardo P, Bence JR, Peacor SD, Osenberg CW. 2018 Bias in meta-

analyses using Hedges' d. *Ecosphere* **9**, e02419. (doi:10.1002/ecs2.2419)

43. Borenstein M, editor. 2009 *Introduction to meta-analysis*. Chichester, U.K: John Wiley &

Sons.

44. R Core Team. 2020 *R: A language and environment for statistical computing*. R Foundation

for Statistical Computing, Vienna, Austria. See https://www.R-project.org/.

45. Viechtbauer W. 2010 Conducting meta-Analyses in R with the metafor package. *J. Stat.*

Softw. **36**. (doi:10.18637/jss.v036.i03)

46. Bürkner P-C. 2017 brms: An R package for Bayesian multilevel models using Stan. *J. Stat.*

Softw. **80**, 1–28. (doi:10.18637/jss.v080.i01)

47. Bürkner P-C. 2018 Advanced Bayesian multilevel modeling with the R package brms. *R J.*

10, 395–411.

48. Stan Development Team. 2021 *Stan Modeling Language*. See https://mc-

stan.org/users/documentation/.

49. Lajeunesse MJ, Rosenberg MS, Jennions MD. 2013 Phylogenetic nonindependence and

meta-analysis. In *Handbook of Meta-analysis in Ecology and Evolution* (eds J Koricheva, J

Gurevitch, K Mengersen), pp. 284–299. Princeton University Press.

50. Chamberlain SA, Szöcs E. 2013 taxize: taxonomic search and retrieval in R. *F1000Res* **2**,

191. (doi:10.12688/f1000research.2-191.v2)

51. Williams DR, Rast P, Bürkner P-C. 2018 Bayesian meta-analysis with weakly informative

prior distributions. *PsyArXiv* (doi:10.31234/osf.io/7tbrm)

52. Mengersen K, Schmid CH, Jennions MD, Gurevitch J. 2013 Statistical models and

approaches to inference. In *Handbook of Meta-analysis in Ecology and Evolution* (eds K

Mengersen, J Gurevitch, J Koricheva), pp. 89–107. Princeton University Press.

53. Safran R *et al.* 2012 A robust new metric of phenotypic distance to estimate and compare

multiple trait differences among populations. *Curr. Zool.* **58**, 426–439.

(doi:10.1093/czoolo/58.3.426)

54. Sawilowsky S. 2009 New effect size rules of thumb. *J. Mod. Appl. Stat. Methods* **8**.

(doi:10.22237/jmasm/1257035100)

55. Kinnison MT, Unwin MJ, Hendry AP, Quinn TP. 2001 Migratory costs and the evolution of

egg size and number in introduced and indigenous salmon populations. *Evolution* **55**, 1656–

1667. (doi:10.1111/j.0014-3820.2001.tb00685.x)

56. Smith CC, Fretwell SD. 1974 The optimal balance between size and number of offspring.

Am. Nat. **108**, 499–506. (doi:10.1086/282929)

57. Parker GA, Begon M. 1986 Optimal egg size and clutch size: effects of environment and

maternal phenotype. *Am. Nat.* **128**, 573–592. (doi:10.1086/284589)

58. Kasimatis K, Riginos C. 2016 A phylogenetic analysis of egg size, clutch size, spawning

mode, adult body size, and latitude in reef fishes. *Coral Reefs* **35**, 387–397.

(doi:10.1007/s00338-015-1380-1)

59. Freidenburg LK, Skelly DK. 2004 Microgeographical variation in thermal preference by an

amphibian: microgeographical thermal preference. *Ecol. Lett.* **7**, 369–373.

(doi:10.1111/j.1461-0248.2004.00587.x)

60. Skelly DK. 2004 Microgeographic countergradient variation in the wood frog, *Rana*

Sylvatica. *Evolution* **58**, 160–165. (doi:https://doi.org/10.1111/j.0014-3820.2004.tb01582.x)

61. Trussell GC, Etter RJ. 2001 Integrating genetic and environmental forces that shape the

evolution of geographic variation in a marine snail. In *Microevolution Rate, Pattern, Process*

(eds AP Hendry, MT Kinnison), pp. 321–337. Dordrecht: Springer Netherlands.

(doi:10.1007/978-94-010-0585-2_20)

62. Lande R, Arnold SJ. 1983 The measurement of selection on correlated characters. *Evolution*

37, 1210–1226. (doi:10.2307/2408842)

63. Christie MR, Knowles LL. 2015 Habitat corridors facilitate genetic resilience irrespective of

species dispersal abilities or population sizes. *Evol. Appl.* **8**, 454–463.

(doi:10.1111/eva.12255)

64. Whiteley AR, Fitzpatrick SW, Funk WC, Tallmon DA. 2015 Genetic rescue to the rescue.

Trends Ecol. Evol. **30**, 42–49. (doi:10.1016/j.tree.2014.10.009)

Figure Captions

Figure 1: Conceptual illustration of four types of phenotypic responses for two populations of a marine fish distributed along the Atlantic coast of the United States. Each row represents four related, but different responses: a) environment alone (phenotypic plasticity without local adaptation), b) a genetic by environment response, c) countergradient variation, d) and cogradient variation. The first two columns illustrate how each of the four responses manifest in different environments—shown is the expected phenotypic responses of both populations in their home environments ("wild") and if they were exposed to a reciprocal transplant experiment in their respective home habitats ("reciprocal transplant"). We also include the reaction norm for each response, where the base of the arrow indicates a population in its home environment and the arrowhead its away environment. Notice that for the counter-gradient variation example, both populations are the same size in the wild (*i.e.*, perfect compensation), illustrating one way that local adaptation can be cryptic.

Figure 2: Posterior distributions of effect size estimates from random-effects meta-analysis for counter- and cogradient variation. The circle indicates the mean and the diamond the median of the posterior distribution (overlapping in countergradient) and the solid, black line, the 95% credible intervals. The mean and credible interval values are also indicated with text on the right. A total of 557 of the largest values (of 50,000 in each distribution) were removed because they extended beyond the x-axis margin (see Fig. S6 for all data points). Purple represents the posterior distribution for the effect size of countergradient variation and green, cogradient variation. The dashed line illustrates the mean value of the posterior distribution for countergradient (purple) and cogradient (green) variation.

Figure 3: Posterior distributions of estimated marginal means for the covariates in a

metaregression analysis for countergradient variation. Yellow distributions depict taxonomic

class, green distributions depict environmental gradients, and purple distributions depict trait

covariates. The black circle indicates the mean and grey diamond the median of the posterior

distribution. The thick solid, horizontal black line depicts the 50% credible intervals for each

covariate and the grey line the 95% credible intervals, some of which extend beyond the x-axis

limit (the full 95% CI is indicated with text to the right). The x-axis limit was set to 5 to better

visualize the behavior around the bulk of the posterior, as such 8,949 of the largest values (of

60,000 thinned to 6,000 in each distribution—198,000 total) were removed because they

extended beyond the x-axis margin (all values can be viewed in Figure S7). The dashed line

illustrates each posterior distribution's relationship to the study-wide effect size of 1.05

estimated from the random effects model and shown in Fig. 2.

Figure 4: Conceptual representation of two countergradient populations in their home

environments and the reaction norms of those populations in the home and away environments.

Also shown are the results of our compensation analysis. a) Represents an overcompensating

relationship where the northern population is larger than the southern population in all

environments and the slope of the dashed line connecting their home phenotype (arrow bases)

has a negative sign as it decreases from the larger northern population to the smaller southern

population. b) Represents a perfectly compensating relationship, where phenotypes are equal in

their respective home environments and thus the slope is flat. c.) Represents an

undercompensating relationship, where the southern population is larger than the northern

population in their respective home environments and the slope of the dashed line connecting

their home phenotype (arrow bases) has a positive sign as it increases from the smaller northern

population to the larger southern population. d.) represents the results of our compensation

analysis where overcompensating was defined as an effect size > 0.5, perfect as $0.5 \leq$ and ≥ -0.5,

and undercompensating < -0.5. Of these three responses, our data was distributed as 38.6%

overcompensating, 25.4% perfectly compensating, and 36.0% undercompensating. Points on the

left are the calculated effect sizes and are connected to a point at 0 to show a slope to match the

conceptual panels. The larger figure is bounded at -15 and 7 (removing 7 values) to better show

the results near 0 and perfect compensation, whereas the inset figure includes all data.

 Table 1

	Estimate	Est.Error	Q2.5	Q97.5
Intercept	1.19	1.23	0.16	4.34
Body size	1.93	1.12	0.6	4.83
Carotenoid concentration	0.87	0.84	0.13	3.08
Ciliary activity	1.25	1.39	0.16	4.71
Developmental rate	1.59	0.76	0.58	3.46
Gamete size	2.81	2.18	0.52	8.68
Growth rate	1.19	0.53	0.47	2.5
Metabolic rate	1.14	0.57	0.4	2.54
Phenology	1.29	1.42	0.15	4.89
Reproductive rate	1.56	1.4	0.25	5.26
Thermal response	1.39	1.39	0.18	5.02
Elevation	1.51	1.43	0.22	5.34
Latitude	0.99	0.45	0.38	2.08
Migration distance	1.85	1.87	0.25	6.89
Photoperiod	1.66	1.63	0.25	5.88
Predator presence	1.34	1.22	0.21	4.66
Salinity	1.02	1.04	0.14	3.73
Shade cover	1.08	0.78	0.24	3.03
Soil phosphate	1.82	1.97	0.22	7.12
Temperature	2.41	1.44	0.7	6.15
Urbanisation	0.74	0.65	0.13	2.46
Wave action	1.57	1.32	0.3	5.05
Amphibia	1.41	1.13	0.27	4.42
Anthozoa	1.35	1.41	0.17	4.94
Asteraceae	1.31	1.46	0.15	5.12
Bivalvia	1.24	1.31	0.14	4.65
Gastropoda	1.15	1.09	0.18	4.02
Insecta	1.22	0.98	0.25	3.92
Liliopsida	1.83	2.02	0.21	7.21
Malacostraca	1.35	1.21	0.24	4.75
Reptilia	1.31	1.13	0.25	4.12

SD Paper Number	1.97	0.39	1.22	2.79
SD Paper Number:Trait	2.26	0.28	1.82	2.91
SD Species	3.41	1.35	1.13	6.46

629 **Table 1:** Results of Bayesian metaregression mixed model with effect size fit as a function of

630 trait, gradient, and organism class. Random effects included trait nested in paper and paper alone,

631 as well as a phylogenetic species effects. Shown is the intercept and the effect of each level of a

632 covariate along with their estimated error and 95% credible intervals.

Figure 1

Figure 2

Figure 3

	Estimated Marginal Mean
Actinopterygii	1.27 [0.21, 3.95]
Amphibia	1.72 [0.15, 7.63]
Anthozoa	1.61 [0.11, 6.95]
Asteraceae	1.5 [0.11, 6.92]
Bivalvia	1.51 [0.09, 6.69]
Gastropoda	1.35 [0.11, 6.18]
Insecta	1.42 [0.15, 5.64]
Liliopsida	2.06 [0.15, 9.43]
Malacostraca	1.62 [0.14, 7.13]
Reptilia	1.56 [0.14, 6.61]
Carotenoid availability	1.2 [0.2, 3.89]
Elevation	1.76 [0.12, 8.02]
Latitude	1.04 [0.19, 3.11]
Migration distance	2.13 [0.15, 9.29]
Photoperiod	1.89 [0.13, 8.55]
Predator presence	1.56 [0.11, 7.22]
Salinity	1.16 [0.08, 5.29]
Shade cover	1.19 [0.13, 4.56]
Soil phosphate level	1.95 [0.14, 8.52]
Temperature	2.58 [0.38, 8.96]
Urbanisation	0.79 [0.08, 3.2]
Wave action	1.72 [0.17, 7.24]
Body shape	1.13 [0.19, 3.75]
Body size	1.95 [0.28, 6.69]
Carotenoid concentration	0.93 [0.07, 4.2]
Ciliary activity	1.33 [0.09, 5.9]
Developmental rate	1.59 [0.28, 5.05]
Gamete size	2.94 [0.26, 11.11]
Growth rate	1.19 [0.22, 3.59]
Metabolic rate	1.14 [0.2, 3.68]
Phenology	1.33 [0.09, 6.03]
Reproductive rate	1.7 [0.12, 7.58]
Thermal response	1.5 [0.1, 7.13]

Figure 4

a) **Overcompensating**

b) **Perfectly compensating**

c) **Undercompensating**

d)