**Cryptic local adaptation is pervasive across environmental and evolutionary gradients**

Morgan M. Sparks[1,*], Joshua C. Kraft[2], Kliffi M. Subida Blackstone[1], Gordon G. McNickle[2,3],

Mark R. Christie[1,4]

1. Department of Biological Sciences, Purdue University; 915 W. State St., West Lafayette,

Indiana 47907-2054 USA

2. Department of Botany and Plant Pathology, Purdue University; 915 W. State St., West

Lafayette, Indiana 47907-2054 USA

3. Purdue Center for Plant Biology, Purdue University; 915 W. State St., West Lafayette, Indiana

47907-2054 USA

4. Department of Forestry and Natural Resources, Purdue University; 715 W. State St., West

Lafayette, Indiana 47907-2054 USA

*corresponding author: Morgan M. Sparks; Department of Biological Sciences, Purdue
University; 915 W. State St., West Lafayette, Indiana 47907-2054 USA;
msparks1309@gmail.com; 360-791-1369

**Keywords:** countergradient variation, cogradient variation, genetic assimilation, maladaptive
plasticity, perfect compensation, phenotypic plasticity

**Abstract**

Cryptic local adaptation—where an environmental effect masks the magnitude of the genetic

contribution to a phenotype—has long been a topic of interest in ecology and evolution. Here,

we systematically document the magnitude and drivers of two common forms of cryptic local

adaptation—counter- and cogradient variation. Using a hierarchical Bayesian meta-analysis, we

calculated the overall effect size as 1.03 for countergradient variation and 0.90 for cogradient

variation. This result indicates that the genetic effect of cryptic local adaptation is approximately

equal to a one standard deviation change in trait value between the most disparate populations.

We also found that the abiotic and biotic covariates with the largest mean effects were

temperature (2.50) and gamete size (2.78), although there was substantial variance. Our results

demonstrate the pervasiveness and large effect of cryptic local adaptation in wild populations

and underscores the importance of accounting for these effects in future studies.

45
46

## Introduction

The persistence of populations via adaptation to local environmental conditions remains a central issue in ecology and evolutionary biology (Blanquart *et al.* 2013). Environmental gradients, where abiotic factors such as temperature, salinity, or nutrient availability vary in a predictable manner, provide a framework to formulate and test questions about the pattern and magnitude of local adaptation. Pairing environmental gradients with common garden or reciprocal transplant studies, especially when there are many populations distributed along these gradients, can lead to insights into how local adaptation covaries with specific environmental factors (Levins 1968; Hargreaves *et al.* 2020).

Shared patterns of phenotypic variation along ecological gradients spurred notable research into the environmental and genetic determinants of body size and morphology and led to the establishment of eco-geographical "rules" such as Bergmann's, Allen's, and Hesse's rules and the subsequent deviations therein (Bergmann 1848; Allen 1877; Atkinson & Sibly 1997; Blanckenhorn 2004; Gaston *et al.* 2008; Tian & Benton 2020). Foundational work by Levins (1968, 1969) revealed a pattern whereby fruit flies along an elevation gradient expressed a similar body size in their wild home environments, leading to the conclusion that the trait was canalized. However, a subsequent study revealed the higher elevation populations were much smaller than lowland populations in common environments—a response believed to be related to desiccation—revealing these populations were locally adapted to different elevations and their environmental conditions and expressed disparate phenotypes across all common environments. In concert, these findings provided the first evidence of cryptic local adaptation.

Local adaptation is commonly expressed with reaction norm plots, where the trait value of a genotype is plotted as a function of multiple environmental values (West-Eberhard 2003). In

a reaction norm, the slope of a genotype's response is representative of phenotypic plasticity (the

effect of the environment) and the difference between genotype-specific reaction norms at each

environmental variable is representative of the genotypic difference (the genetic effect). When

different genotypes express the same reaction norm (i.e., overlapping lines, Fig. 1a, plasticity

alone) this is representative of a purely environmentally driven response where distinct

genotypes would express the same trait values (or phenotype) along an environmental gradient.

Alternatively, populations may be locally adapted, such that the reaction norms cross when

comparing different common environments, which is indicative of a genotype by environment

interaction (Fig. 1b, Genetic x Environment). Finally, populations may exhibit cryptic local

adaptation, where the genetic and environmental effects covary with one another, either

negatively or positively (Conover *et al.* 2009). In this scenario, the reaction norms for

populations may have the same or similar slopes thus maintaining consistent rank order in their

trait values (Fig.1, co- and countergradient variation). For example, a high latitude population

may have consistently larger body sizes across environments than a low latitude population,

despite both expressing plasticity in body size. When the genetic portion of phenotypic variance

opposes with the environmental portion, it is called countergradient variation (Fig. 1c), and when

those two factors act in the same direction it is termed cogradient variation (Fig. 1d) (Levins

1968; Conover & Schultz 1995). That is, when comparing among wild populations, phenotypic

traits appear more similar than they would if populations were compared in a common

environment for countergradient variation, and vice versa for cogradient variation.

Co- and countergradient variation constitute forms of cryptic local adaptation because

their true genetic effect on phenotypic variation is masked by the environment (Conover &

Schultz 1995; Schlichting 2008; Conover *et al.* 2009; Kinnison *et al.* 2015) (Fig.1).

Countergradient variation can be particularly difficult to identify in wild populations because the

environmental influence on a phenotype can conceal the genetic influence, such that the signal of

local adaptation only appears when individuals are reared in common environments. This

response has been termed perfect compensation (or exact compensation), when wild-type

phenotypes are roughly equal across a gradient in their home environments (Fig. 1c) (sensu

Conover *et al.* 2009 ). Compensation can be thought of as a measure of how similar phenotypic

values are across populations in their home environments—and as an indicator of whether

selection may favor similar or disparate phenotypes along an environmental gradient.

While Levins (1969) was the first to formally describe countergradient variation—which

was followed by early research by Berven and others (Berven *et al.* 1979; Berven 1982) in *Ranid*

frogs—it wasn't until a two-and-a-half decades later than interest in cryptic local adaptation

grew substantially. This research was led by a group investigating a coastal, eastern North

American fish, Atlantic silverside (*Menidia menidia*), after they discovered that populations had

similar body sizes along a latitudinal gradient that spanned from the Gulf of Maine to the

Floridian Atlantic Coast. Using a common garden approach, they found that the northern

populations were consistently larger than southern populations, regardless of the temperature in

which they were raised (*e.g.*, Fig. 1c). This finding led to substantial follow up research

investigating the proximate and ultimate factors driving the trend (Conover & Present 1990;

Conover *et al.* 2005), namely faster growth and more efficient metabolism in response to size-

dependent winter mortality. After a theoretical review outlining the evolutionary significance of

countergradient variation (Conover & Schultz 1995), subsequent researchers added to the

evidence for co- and countergradient variation across diverse taxa including plants, bivalves,

fishes, and insects (*e.g.*, *plants*, Chapin & Oechel 1983; *bivalves*, Dittman 1997; *fishes*,

DiMichele & Westerman 1997; Imsland *et al.* 2000; Jean M. Billerbeck *et al.* 2000; Jonassen *et*

*al.* 2000; *insects*, Blanckenhorn 1991; Armbruster & Conn 2006; Pöykkö & Tammaru 2010), and

across ecological gradients such as temperature, salinity, carotenoid availability, and

urbanization (*e.g., temperature*, Yamahira & Conover 2002; Kokita 2003; *salinity*, Secor *et al.*

2000; *carotenoid availability*, Craig *et al.* 2005; Grether 2005; and *urbanisation*, Tüzün *et al.*

2017).

Despite the attention paid to counter- and cogradient variation, there have been few

studies that describe the magnitude and extent of these types of local adaptation, nor has there

been much investigation into the broad abiotic and biotic factors that might contribute to their

occurrence (Conover *et al.* 2009; Hendry 2017; Stamp & Hadfield 2020). Without understanding

the extent of cryptic local adaptation, it is difficult to make predictions about how populations

expressing cryptic local adaptation may interact with a changing environment or how local

adaptation may aid or hinder conservation goals. Here we use a Bayesian meta-analytical

framework with hierarchical models to describe the general effect size of both counter- and

cogradient variation as they appear in the literature and to investigate what biotic and abiotic

factors are associated with their occurrence. The goals of these analyses are to *i)* quantify the

effect size of countergradient and cogradient variation, *ii)* describe these effect sizes within and

across taxa, phenotypic traits, and different environmental gradients, and *iii)* quantify the effect

size of compensation for the home environment phenotypes to assess if there are clear trends for

phenotypic similarity or dissimilarity along gradients.

**Methods**

*Literature Review and Selection*

Overall, we collected 858 studies that likely investigated counter- and/or cogradient variation. To

generate a full list of studies, we first analyzed studies indicated as showing counter- or

cogradient variation in the qualitative analysis from Conover *et al.* (2009). We then searched

Web of Science topic field using the terms "counter*gradient variation" in May of 2018 and

"co*gradient variation" (while excluding the previous search, as the wildcard renders them

redundant) in June of 2019, resulting in 384 and 34 results, respectively. In August 2019, we also

included studies citing the Conover *et al.* (2009) review, as well as the earlier review article by

Conover & Schultz (1995)—682 studies total (some of which were redundant with studies in the

prior search). These methods are discussed in more depth in the Supplementary Methods.

To be included in our analysis, a study had to meet a set of qualifying criteria.

Specifically, a study had to be designed to include some form of a common environment—this

may have been in a common garden study or a reciprocal transplant. Additionally, studies must

have included two or more common environments to determine the signal of local adaptation

(countergradient, cogradient, or others such as GxE that were ultimately not included in this

study), as well as two or more populations to compare the genotypic difference of the phenotypic

response between populations. Occasionally, studies investigated these distinctions across

multiple species in the place of populations and such studies were discarded. Furthermore, each

study must have included some mean estimate for a population phenotypic value (or data that

could be used to calculate a mean), a measure of variation that could be transformed into units of

standard deviation, and sample sizes for these statistics. When studies reported ranges an

estimate for SD was made by subtracting the maximum value from the minimum value and then

dividing by four as in Davidson *et al.* (2011).

*Statistical Methods*

To calculate effect sizes, we took data from both tables and figures in manuscripts. For data in

the form of figures, we used the software WebPlotDigitizer (Rohatgi 2020) to extract the

relevant data. Briefly, the program works by importing a photo of the respective figure and the

user manually sets the scales for the x and y-axes based on the figure values and then adds digital

points with their cursor to the data of interest. The digital points are then translated to numeric

data via the imaging algorithm.

The statistic used to calculate effect size was Hedge's $d$, which is a standardized mean

difference with a correction for small sample sizes (Hedges & Olkin 1985; Rosenberg *et al.*

2013). We calculated standardized mean difference (synonymous with effect size in this paper)

by comparing the most extreme populations at each treatment level. For example, if we had three

populations collected from an elevational gradient at 500m, 1000m, and 1500m in common

gardens with three temperature treatments (e.g., 10, 15, and 20 °C) (as was often the case where

temperature was a proxy for latitude or elevation), then we would calculate the mean and

variance for every population in every treatment, but calculate the effect size (standardized

mean difference) by using the values from the 500m and 1500m populations at each treatment.

Occasionally, not all populations were used in all treatments, in these scenarios we used only the

treatments for the most extremely distributed (with respect to the gradient—*e.g.*, very highest

and lowest elevations) populations. The multiple values for a given trait in a given study (for

example, the mean effect size of those that were calculated at 10, 15, and 20 °C in our prior

example) were averaged as a single value (Borenstein 2009). Some studies conducted multiple

experiments measuring the same trait. If multiple experiments measuring the same trait were

recorded, we calculated the mean effect sizes for each experiment independently.

In all, we calculated 422 total effect sizes and 119 averaged effect sizes from 1204

individual data points extracted from 83 studies. Many studies showed counter- or cogradient

patterns in multiple traits and/or trials within a given study. We collected all these data and

controlled for their non-independence using a robust, Bayesian hierarchical modeling approach

(discussed below). Hedge's *d* was calculated using the *metafor* package in program R (version

3.6.3) (R Core Team 2020) with the *escalcl()* function (Viechtbauer 2010). Additional metadata

was collected from each study and included features such as year the study was published, the

species investigated, and the distance in meters (for elevational gradients) and kilometers (for

latitudinal gradients), the full list of which are available in our data file metadata tab

(Supplementary Data). These data were collected to be used as covariates in the mixed-effects

model (Table 1).

To estimate overall effect size and the effect of covariates, Bayesian linear mixed models

fit with random intercepts were used in the R package *brms* (Bürkner 2017, 2018), which is a

high-level interface in R for the Stan modeling language (Stan Development Team 2021). Linear

mixed-effects models were chosen due to the hierarchical nature of this data set, where a study

may have effect sizes estimated for multiple traits in multiple experiments. As such, a nested

random intercepts effect was used to control for this non-independence. We originally set out to

nest experiment within trait within study as the most appropriate reflection of our data's non-

independence, but that structure proved to be too computationally intensive because model

parameterization necessary to overcome divergent transitions (specifically, max tree depth in

Stan) extended model run time beyond available compute power on our computational cluster.

As such, random effects were reduced to traits nested in studies.

Similarly, phylogenetic relationships may provide another source of non-independence,

and was controlled for by using a phylogenetic random intercepts effect for which we created the

variance-covariance matrix (Lajeunesse *et al.* 2013). To do so, we generated a pseudo-phylogeny

using the *taxize* package in R (Chamberlain & Szöcs 2013). Using the *classification()* function,

we extracted the full taxonomy of a given organism in our dataset from the NCBI database. We

then used the *class2tree()* function to generate a phylogenetically accurate tree in terms of

taxonomic rank (*e.g.*, all species within a genus would have the same distance from one another,

all genera within a family would have the same distance—a full explanation of the method is

available in the Supplementary Methods). The function not only clusters taxa on their specific

taxonomic rank but also within the taxonomic clade, allowing for more accurate distance

measures. The tree created using this method was made into a matrix and then used as a custom

variance-covariance component in our hierarchical model. The mathematical representation of

our random effects in all models is shown in our Supplementary Materials (Supplementary

Methods).

We used weakly to moderately informative priors for our hierarchical models (Williams

*et al.* 2018)—except for the metaregression, where a stronger prior was used to reign in

variance—lists of priors are available in Supplementary Materials (Tables S1-S3). For our

models, we used four chains, with thousands of burn-ins, tens-of-thousands of iterations for each

chain, with thinning set to 1 to 10 depending on the number of iterations in the model, resulting

in >5,000 estimates for our posterior distributions (exact values are available in Table S4). Trace

plots and posterior predictive checks are included in Figures S1-4.

Because we were interested in the magnitude of our effect and not the sign (*e.g.*, when

calculating an effect size for developmental rate a countergradient response would be quicker

time to hatch relative to other populations and thus a negative number), we used log transformed

absolute values of effect sizes as our response variable in order to use a Student's t-distribution

and normally distributed priors (other candidate distribution types available in *brms,* such as

lognormal, disallow weighting on variance, which is needed for a meta-analysis). As such, back-

transformed values tended to stack near zero, but could not drop below zero, and distributions

tended to have a long tail as numbers increased. Because we used a Bayesian framework, the

following effects are presented with the means and 95 percent credible intervals (indicated as

95% CI) of the posterior distributions, unless otherwise noted.

The overall effect size for both co- and countergradient variation was estimated using

random-effects only models, which is the convention when assuming study effects may vary in

meta-analytical approaches (Mengersen *et al.* 2013). We included the covariates of trait,

gradient, publication year, and taxonomic class in our metaregression countergradient model to

assess how different biotic and abiotic factors may contribute to the estimate of countergradient

variation effect size. This analysis was not performed with cogradient variation because of the

lack of diversity of data to include many covariates in our modeling approach.

*Compensation Analysis*

Finally, we analyzed whether populations appeared to be under-, over-, or perfectly

compensating (*sensu* Conover *et al.* 2009) when comparing their home environment phenotypes.

To do so, we used the most extreme treatments and paired them with the polarly (as in opposites,

not necessarily geographically) distributed populations used in the study. For example, if we

consider three populations of fish from a range of saline conditions—freshwater, intermediary,

and marine—then, the most saline treatment in the experiment would be considered the home

environment of the marine population and the least saline the home environment of the

freshwater fish (unless other treatments were noted to be reflective of home environments or

chosen to be specifically extreme). We then compared the trait values of the two polarly

distributed populations and using the previous example, if the trait value was larger for the most

saline population than that of the least saline population the relationship was classified as

overcompensating, close to even was classified as perfectly compensating, and negative was

classified as undercompensating. To standardize these data, we again used Hedge's $d$, because

the values of different traits are generally measured on different scales (Safran $et$ $al.$ 2012).

Importantly, compensation was not assessed with a model, as our analysis was meant to be

exploratory and because there are no quantitative definitions of compensation. The results were

classified as overcompensating if values were >0.5, perfectly compensating if values were

between -0.5 and 0.5, and undercompensating if values were < -0.5. Because compensation

measurements were computed as Hedge's $d,$ these were meant to reflect a broad distribution

around 0 for perfect compensation (what amounts to a medium effect size in both positive and

negative directions, Sawilowsky 2009) that encapsulates the inherent measurement error in the

value.

**Results**

In all, our systematic review considered 858 studies. We included 83 studies that either

demonstrated co-, countergradient variation, and met the requirements for inclusion in our

analysis. From those, we calculated 1,204 individual trait values, and 1,024 were for traits that

were exhibiting countergradient variation and 180 for cogradient variation.

*Random Effects Models*

The result of our random effects models that estimated overall effect size for countergradient

variation was 1.03 (95% CI = 0.56-1.59) and for cogradient variation 0.90 (95% CI = 0.22-2.50)

(Fig. 2, non-truncated in Fig. S5). The heterogeneity parameter τ—the full random effect

standard deviation—was modeled with its own prior distribution and resulted in 2.32 (95% CI =

1.73-3.02) for countergradient variation and 2.96 (95% CI = 1.13-6.69) for cogradient variation.

We similarly included a random effect for phylogenetic relatedness which resulted in an estimate

of 1.37 SD (95% CI = 1.01-2.26) for countergradient variation and 2.73 SD (95% CI = 1.03-

8.36) for cogradient variation (Tables S5, S6).

*Metaregression*

We also implemented a mixed effects metaregression with our countergradient dataset to

estimate the effect size for trait classification, gradient classification, and taxonomic class effect.

The mean effect, posterior distribution and 95% credible intervals for each modeled covariate are

presented in Table 1 and those same values are represented as the main effects estimated

marginal means with the posterior distribution and credible intervals in Figure 3 (non-truncated

in Fig. S6). For each covariate, these results illustrate similar mean values with large and

overlapping credible intervals. While variability was large across all covariates, the mean

coefficient estimates of the posterior distributions for temperature (mean = 2.50, 95% CI = 0.74-

6.19) and gamete size (mean = 2.78, 95% CI = 0.54-8.74) were much larger than the global

estimate of 1.01. By contrast, the smallest value for urbanization (mean = 0.75, 95% CI = 0.14-

2.46)—which had lower credible intervals near zero and was much lower than other estimates.

The between trait (nested in study) standard deviation was 2.28 (95% CI = 1.90-2.86) and the

standard deviation for phylogenetic random effect was 1.51 (95% CI = 1.01-2.77). Note, these

estimates are slightly different than those presented in Figure 3, which are the main effect

estimated marginal means derived from the model, rather than model coefficients.

*Compensation Analysis*

The results of our compensation analysis were plotted to represent the conceptual relationship

between home phenotypes of populations exhibiting countergradient variation (*sensu* Conover et

al. 2009) (Fig. 4). Our data provide support for far more frequent over- and under compensation

than perfect compensation, meaning that populations in our study generally have disparate home

phenotypes as opposed to equal ones.  We considered any value between -0.5 and 0.5 as

perfectly compensating (or a range of 1 effect size around 0 for a conservative estimate), values

less than -0.5 undercompensating, and values larger than 0.5 overcompensating. We consider this

classification conservative because it allows for a medium effect size in the positive or negative

direction from zero (Sawilowsky 2009), despite the theoretical description of perfect

compensation being a value equal to zero (Conover et al. 2009). The distribution of these values

in our data set were overcompensating = 38.6%, undercompensating = 36.0%, and perfectly

compensating = 25.4%. Effect sizes ranged from -70.38 to 36.45 but 5% and 95% quantiles were

-27.74 and 6.49, respectively.

**Discussion**

The ecological and evolutionary importance of co- and countergradient variation has been

recognized for three decades (Conover & Present 1990; Conover & Schultz 1995). Here, we find

that the overall effect size of co- and countergradient variation are 1.03 and 0.90, respectively.

Our measure of standardized mean difference, Hedges' *d,* essentially translates the difference

between populations into units of standard deviation. Thus, our results indicate that the

standardized mean difference between the most disparate populations in our study—*i.e.*, the

genetic effect of cryptic local adaptation—is approximately equal to a one standard deviation

change in trait value. This result demonstrates that genetic differentiation among cryptic, locally

adapted populations is large but also obscured as a function of the covariance between

environmental and genetic effects and their interaction with phenotype.

We used a metaregression to investigate the effect sizes of abiotic and biotic covariates of

countergradient variation; covariates were statistically indistinguishable from one another due to

high variance in their estimate, though many showed values that were large with distributions

away from 0 (Effect sizes estimates of 1-3, Table 1, Fig. 3). Instead, we find that the inter-effect

variability ($\tau^2$, or the variability of our random effects) generally swamped the covariate-specific

effects estimates in our metaregression (Table 1). While the lack of predictive results in our

metaregression may be indicative that our data is not robust enough for differences among

covariates to emerge, we think it is more likely that these results suggest that countergradient

variation, when it does occur, is an ecologically ubiquitous phenomenon for species distributed

along environmental gradients. Furthermore, we suspect the mechanism driving such adaptation

is different among populations and gradients and therefore differs in its effect contributing to the

variability observed in our models. In other words, while the phenotypic solution may appear

similar across species and gradients, the processes driving the genetic divergence of the

phenotypes of different populations is idiosyncratic. This conclusion is further supported by the

lack of pattern with respect to latitudinal distance and temperature difference regressions

analyses, neither of which were predictive (Fig. S7, Table S7). Thus, while populations almost

always showed a large effect of cryptic local adaptation—model covariates were 2.50 and 0.99,

respectively—the magnitude did not scale with temperature or physical distance.

Despite the fact that the overall effect size of cogradient variation was large, our results

for cogradient variation were only based on 15 studies. This paucity reflects an overall lack of

published research investigating and demonstrating cogradient variation (see also, Conover *et al.*

2009). However, in a meta-analysis of reciprocal transplant studies, Stamp & Hadfield (2020)

found 60% of traits exhibited a cogradient pattern. Our results (*i.e.*, the magnitude of effect size),

in conjunction with the work of Stamp & Hadfield (2020) suggest that, while rarely reported,

cogradient variation constitutes an ecologically meaningful phenomenon. We further hypothesize

cogradient variation may suffer from underreporting because it's natural sign mirrors

environmentally induced plasticity that may not drive researchers to investigate populations

using common gardens. While cogradient variation may be underreported, work such as that by

Trussell & Etter (2001) show both the large effect of cogradient variation and how home

phenotypic differences can disguise that effect. In a reciprocal transplant study, the researchers

measured shell thickness in marine snail populations, finding that, when in their home

environments, populations expressed a 3.18 standardized mean difference in shell thickness. But,

when considered in shared environments—or the genetic difference due to cogradient

variation—that difference reduced to 2.07. Notably, this amounts to a substantial effect size

difference due to genetic variation alone, but is ~2/3 value if we observed the populations only in

their home environments. More importantly, if the researchers had only investigated these

populations in their home environments without common gardens, the obvious interpretation for

the phenotypes would have been a strong plastic response, which, while true, did not account for

even half of phenotypic variation in home environments.

Our compensation analysis for countergradient variation revealed that populations in their

home environments were more commonly over- or undercompensasting, as opposed to perfectly

compensating (Fig. 4). In other words, the genetic effect of countergradient variation was

generally not equal to that of the environmental effect. This result was clear with conservative

bounds for perfect compensation (-0.5 to 0.5 standardized mean difference). Reducing those

bounds to the less conservative values of -0.2 to 0.2 (considered to be a small effect size,

Sawilowsky 2009) for perfect compensation resulted in only 11.7 percent of studies

demonstrating perfect compensation. While we caution that these results may not necessarily

point to clear biological trends about how common and in what scenarios over- and under-

compensation may occur, it does suggest that either: 1. selection for maintaining optimal

phenotypes across broad environmental gradients may not be the norm (*sensu* Conover &

Schultz 1995; Conover *et al.* 2009), or 2. the response to selection may not be strong enough to

effectively minimize phenotypic variation across populations. If the latter, the response to

selection may be minimized by insufficient adaptive genetic variation or countervailing selection

on correlated traits specific to each local population (Lande & Arnold 1983).

One of the confounding aspects of investigating clinal trends in locally adapted

populations is the often non-continuous nature of their phenotypic values. For example, the non-

continuous nature of population responses is particularly clear in Hice *et al.* (2012), in which

they investigated 37 sites across the entire range of Atlantic silversides and found an overall

trend of countergradient variation in growth rate, but multiple breakpoints along the cline. The

results presented in Hice *et al.* (2012), constitutes one of the few attempts to characterize cryptic

local adaptation across the full environmental gradient for a species or set of populations.

However, if the full distribution of populations is ignored or nor correctly accounted for, such

sample designs may present a false signal of cryptic local adaptation. Few studies in this analysis

included measures of genetic differentiation among the populations they were measuring. For

example, hypothetical meta-populations distributed along a gradient such as latitude with little or

no gene flow between them may experience divergent selection on the same trait but from

different modes of selection and could appear as cryptic local adaptation.

The observation that change in treatment temperature or latitude was not predictive of

effect size may be explained by the heterogeneity of environments (Fig. S7). A useful example

of why this result may occur is microgeographic variation, in which large adaptive phenotypic

divergence occurs over small spatial scales where environmental factors may be more disparate

than initially predicted. Indeed, microgeographic variation was observed in some of the studies

that were reviewed as part of this study. Researchers found that in temperature choice trials,

wood frogs (*Rana sylvatica*) captured from ponds with more forest canopy, preferred higher

temperatures (Freidenburg & Skelly 2004) across a single 3800-hectare forest. Companion work

from a common garden with two temperature treatments, found that frogs from more shaded

ponds developed faster than those from more open ponds (Skelly 2004). While broad patterns in

temperature may make for useful categorizations of clines, the habitat level specifics with respect

to local adaptation may be more complicated.

Clinal variation has been the subject of numerous neutral and quantitative genetic studies

(Richter-Boix *et al.* 2010, 2013; Hangartner *et al.* 2012; Frei *et al.* 2014), however, the pace of

genomic investigation into cryptic local adaptation lags behind other forms of local adaptation.

In fact, fundamental theory about countergradient variation often assumes directional selection

drives the maladapted population/s opposite of the environmental effect to achieve the negative

covariance that characterizes countergradient variation (Grether 2005; Crispo 2007). But, work

by Wilder *et al.* (2020), found substantial genomic divergence at specific chromosomal sites

among northern and southern populations of Atlantic silverside, despite these being continuously

distributed populations with substantial gene flow and low overall differentiation. These most

differentiated genomic sites were largely driven by a single population, with the warmest

temperatures, distinguishing it from all the other more northern populations. These results

suggest that strong selection maintains these regions, despite genomic homogenization from gene

flow. This research is only beginning to scratch the surface of understanding the functional

processes at play at the genomic level in counter- and cogradient variation, and what tradeoffs

may exist in shaping the development of local adaptation.

**Conclusion**

Our findings illustrate that cryptic local adaptation is pervasive across multiple environmental

gradients in a diverse set of taxa. With this meta-analysis, we quantitatively characterized the

genetic effect on phenotypic divergence in cryptic local adaptation and find that the result is at or

close to one full standard deviation from the mean trait value of a population. While the effect of

co- and countergradient variation may be large, the variability of response observed across

studies is also quite high, as indicated in the values of our random effects. Moreover, when

analyzing the mean trait difference between home phenotypes of populations exhibiting

countergradient variation, we find no support for a strong signal of perfect compensation,

indicating that selection for optimal phenotypes across a gradient does not or cannot minimize

variation in mean phenotypic trait values. We conclude that the cryptical local adaptation is both

ubiquitous and has a large effect size and argue that cryptic local adaptation deserves continued

attention in theoretical, experimental, and applied approaches. Undesired consequences can

occur when cryptic local adaptation is ignored, (*e.g.,* Philipp & Whitt 1991) and could lead to

especially problematic outcomes in conservation and/or translocation-oriented applications.

Much like local adaptation more generally, cryptic local adaptation necessitates close

consideration when working with populations distributed across ecological gradients.

**Acknowledgements**

We thank N. Emery and C. Oakley for their helpful discussion regarding cryptic local adaptation.

K. Horvath offered useful suggestions about instituting a phylogenetic random effect in our

models, D. Pascall provided valuable insights on implementing the pseudo-phylogeny, and R.

Swihart gave useful direction on Bayesian model setup. We also thank A. Harder, A. Lee, and A.

Nalesnik for their helpful comments on a past version of this manuscript. This work was

supported, in part, by funding from the National Science Foundation (OCE-1924505 ; DEB-

1856710) to MRC.

**Data Archiving statement**

Code for these analyses is posted on the GitHub account of MMS (https://github.com/morgan-

sparks/CnGV-CoGV-Meta-Analysis) and cloned at the GitHub account of MRC. Data and model

files will be stored permanently at DataDryad.

**Author Contributions**

MMS and MRC designed the research, MMS, JRK, and KMSB collected data, MMS performed

the data analysis, and MMS, GGM, and MRC wrote the manuscript with input from all authors.

**References**

Allen, J.A. (1877). The influence of physical conditions in the genesis of species. *Radical*

*review*, 1, 108–140.

Armbruster, P. & Conn, J.E. (2006). Geographic Variation of Larval Growth in North American

*Aedes albopictus* (Diptera: Culicidae). *Annals of the Entomological Society of America*,

99, 1234–1243.

Atkinson, D. & Sibly, R.M. (1997). Why are organisms usually bigger in colder environments?

Making sense of a life history puzzle. *Trends in Ecology & Evolution*, 12, 235–239.

Bergmann, C. (1848). *Über die Verhältnisse der Wärmeökonomie der Thiere zu ihrer Grösse*.

Berven, K.A. (1982). The genetic basis of altitudinal variation in the wood frog Rana sylvatica

II. An experimental analysis of larval development. *Oecologia*, 52, 360–369.

Berven, K.A., Gill, D.E. & Smith-Gill, S.J. (1979). Countergradient selection in the green frog,

*Rana clamitans*. *Evolution*, 33, 609–623.

Blanckenhorn, W.U. (1991). Life-History Difference in Adjacent Water Strider Populations:

Phenotypic Plasticity or Heritable Responses to Stream Temperature? *Evolution*, 45,

1520.

Blanckenhorn, W.U. (2004). Bergmann and Converse Bergmann Latitudinal Clines in

Arthropods: Two Ends of a Continuum? *Integrative and Comparative Biology*, 44, 413–

424.

Blanquart, F., Kaltz, O., Nuismer, S.L. & Gandon, S. (2013). A practical guide to measuring

local adaptation. *Ecology Letters*, 16, 1195–1205.

Borenstein, M. (Ed.). (2009). *Introduction to meta-analysis*. John Wiley & Sons, Chichester,

U.K.

Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal

of Statistical Software*, 80, 1–28.

Bürkner, P.-C. (2018). Advanced Bayesian Multilevel Modeling with the R Package brms. *The R

Journal*, 10, 395–411.

Chamberlain, S.A. & Szöcs, E. (2013). taxize: taxonomic search and retrieval in R. *F1000Res*, 2,

191.

Chapin, F.S. & Oechel, W.C. (1983). Photosynthesis, respiration, and phosphate absorption by

*Carex aquatilis* ecotypes along latitudinal and local environmental gradients. *Ecology*,

64, 743–751.

Conover, D. & Present, T. (1990). Countergradient variation in growth rate: Compensation for

length of the growing season among Atlantic silversides from different latitudes.

*Oecologia*, 83, 316–324.

Conover, D.O., Arnott, S.A., Walsh, M.R. & Munch, S.B. (2005). Darwinian fishery science:

lessons from the Atlantic silverside (*Menidia menidia*). *Canadian Journal of Fisheries*

*and Aquatic Sciences*, 62, 730–737.

Conover, D.O., Duffy, T.A. & Hice, L.A. (2009). The covariance between genetic and

environmental influences across ecological gradients: reassessing the evolutionary

significance of countergradient and cogradient variation. *Ann. N. Y. Acad. Sci.*, 1168,

100–129.

Conover, D.O. & Schultz, E.T. (1995). Phenotypic similarity and the evolutionary significance

of countergradient variation. *Trends Ecol. Evol.*, 10, 248–252.

Craig, J.K., Foote, C.J. & Wood, C.C. (2005). Countergradient variation in carotenoid use

between sympatric morphs of sockeye salmon (*Oncorhynchus nerka*) exposes

nonanadromous hybrids in the wild by their mismatched spawning colour. *Biological*

*Journal of the Linnean Society*, 84, 287–305.

Crispo, E. (2007). The Baldwin effect and genetic assimilation: Revisiting two mechanisms of

evolutionary change mediated by phenotypic plasticity. *Evolution*, 61, 2469–2479.

Davidson, A.M., Jennions, M. & Nicotra, A.B. (2011). Do invasive species show higher

phenotypic plasticity than native species and, if so, is it adaptive? A meta-analysis.

*Ecology Letters*, 14, 419–431.

DiMichele, L. & Westerman, M.E. (1997). Geographic variation in development rate between

populations of the teleost *Fundulus heteroclitus*. *Marine Biology*, 128, 1–7.

Dittman, D.E. (1997). Latitudinal compensation in oyster ciliary activity. *Functional Ecology*,

11, 573–578.

Frei, E.R., Hahn, T., Ghazoul, J. & Pluess, A.R. (2014). Divergent selection in low and high

elevation populations of a perennial herb in the Swiss Alps. *Alp Botany*, 124, 131–142.

Freidenburg, L.K. & Skelly, D.K. (2004). Microgeographical variation in thermal preference by

an amphibian: Microgeographical thermal preference. *Ecology Letters*, 7, 369–373.

Gaston, K.J., Chown, S.L. & Evans, K.L. (2008). Ecogeographical rules: elements of a synthesis.

*Journal of Biogeography*, 35, 483–500.

Grether, G.F. (2005). Environmental change, phenotypic plasticity, and genetic compensation.

*The American Naturalist*, 166, 115–123.

Hangartner, S., Laurila, A. & Räsänen, K. (2012). Adaptive divergence in Moor frog (*Rana*

*arvalis*) populations along an acidification gradient: Inferences from Qst–Fst correlations.

*Evolution*, 66, 867–881.

Hargreaves, A.L., Germain, R.M., Bontrager, M., Persi, J. & Angert, A.L. (2020). Local

adaptation to biotic interactions: A meta-analysis across latitudes. *The American*

*Naturalist*, 195, 395–411.

Hedges, L.V. & Olkin, I. (1985). Statistical methods for meta-analysis. Academic Press.

Hendry, A.P. (2017). *Eco-evolutionary Dynamics*. Princeton University Press.

Hice, L., Duffy, T., Munch, S., & David Conover. (2012). Spatial scale and divergent patterns of

variation in adapted traits in the ocean. *Ecology Letters*, 15, 568–575.

Imsland, A.K., Foss, A., Nevdal, G., Cross, T., Bonga, S.W., Ham, E.A., *et al.* (2000).

Countergradient variation in growth and food conversion efficiency of juvenile turbot.

*Journal of Fish Biology*, 57, 1213–1226.

Lande, R., & Arnold, S. J. (1983). The measurement of selection on correlated characters.

*Evolution*, 1, 1210-1226.

Jean M. Billerbeck, Eric T. Schultz & David O. Conover. (2000). Adaptive variation in energy

acquisition and allocation among latitudinal populations of the Atlantic silverside.

*Oecologia*, 122, 210–219.

Jonassen, T.M., Imsland, A.K., Fitzgerald, R., Bonga, S.W., Ham, E.V., Naevdal, G., *et al.*

(2000). Geographic variation in growth and food conversion efficiency of juvenile

Atlantic halibut related to latitude. *Journal of Fish Biology*, 56, 279–294.

Kinnison, M.T., Hairston Jr., N.G. & Hendry, A.P. (2015). Cryptic eco-evolutionary dynamics.

*Annals of the New York Academy of Sciences*, 1360, 120–144.

Kokita, T. (2003). Potential latitudinal variation in egg size and number of a geographically

widespread reef fish, revealed by common-environment experiments. *Marine Biology*,

143, 593–601.

Lajeunesse, M.J., Rosenberg, M.S. & Jennions, M.D. (2013). Phylogenetic nonindependence and

meta-analysis. In: *Handbook of Meta-analysis in Ecology and Evolution* (eds. Koricheva,

J., Gurevitch, J. & Mengersen, K.). Princeton University Press, pp. 284–299.

Levins, R. (1968). *Evolution in Changing Environments: Some Theoretical Explorations*.

Princeton University Press.

Levins, R. (1969). Thermal acclimation and heat Resistance in drosophila species. *The American*

*Naturalist*, 103, 483–499.

Mengersen, K., Jennions, M.D. & Schmid, C.H. (2013). Statistical models for the meta-analysis

of nonindependent Data. In: *Handbook of Meta-analysis in Ecology and Evolution* (eds.

Mengersen, K., Koricheva, J. & Gurevitch, J.). Princeton University Press, pp. 255–283.

Philipp, D.P. & Whitt, G.S. (1991). Survival and growth of northern, Florida, and reciprocal Fl

hybrid largemouth bass in central Illinois. *Transactions of the American Fisheries*

*Society*, 120, 58–64.

Pöykkö, H. & Tammaru, T. (2010). Countergradient vs. cogradient variation in growth and

diapause in a lichen-feeding moth, *Eilema depressum* (Lepidoptera: Arctiidae). *Journal*

*of Evolutionary Biology*, 23, 1278–1285.

R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation

for  Statistical Computing, Vienna, Austria.

Richter-Boix, A., Quintela, M., Kierczak, M., Franch, M. & Laurila, A. (2013). Fine-grained

adaptive divergence in an amphibian: genetic basis of phenotypic divergence and the role

of nonrandom gene flow in restricting effective migration among wetlands. *Molecular*

*Ecology*, 22, 1322–1340.

Richter-Boix, A., Teplitsky, C., Rogell, B. & Laurila, A. (2010). Local selection modifies

phenotypic divergence among *Rana temporaria* populations in the presence of gene flow.

*Molecular Ecology*, 19, 716–731.

Rohatgi, A. (2020). *WebPlotDigitizer*. Version 4.1. https://automeris.io/WebPlotDigitizer/

Rosenberg, M.S., Rothstein, H.R. & Gurevitch, J. (2013). Effect Sizes: Conventional choices and

calculations. In: *Handbook of Meta-analysis in Ecology and Evolution* (eds. Gurevitch, J.,

Koricheva, J. & Mengersen, K.). Princeton University Press, pp. 61–71.

Safran, R., Flaxman, S., Kopp, M., Irwin, D.E., Briggs, D., Evans, M.R., *et al.* (2012). A robust

new metric of phenotypic distance to estimate and compare multiple trait differences

among populations. *Current Zoology*, 58, 426–439.

Sawilowsky, S. (2009). New effect size rules of thumb. *Journal of Modern Applied Statistical*

*Methods*, 8, 597-599.

Schlichting, C.D. (2008). Hidden reaction norms, cryptic genetic variation, and evolvability.

*Annals of the New York Academy of Sciences*, 1133, 187–203.

Secor, D.H., Gunderson, T.E. & Karlsson, K. (2000). Effect of temperature and salinity on

growth performance in anadromous (Chesapeake Bay) and nonanadromous (Santee-

Cooper) strains of striped bass *Morone saxatilis*. *Copeia*, 2000, 291–296.

Skelly, D.K. (2004). Microgeographic countergradient variation in the wood grog, *Rana*

*sylvatica*. *Evolution*, 58, 160–165.

Stamp, M.A. & Hadfield, J.D. (2020). The relative importance of plasticity versus genetic

differentiation in explaining between population differences; a meta-analysis. *Ecology*

*Letters*, 23, 1432–1441.

Stan Development Team. (2021). *Stan Modeling Language*.

Tian, L. & Benton, M.J. (2020). Predicting biotic responses to future climate warming with

classic ecogeographic rules. *Current Biology*, 30, R744–R749.

Trussell, G.C. & Etter, R.J. (2001). Integrating genetic and environmental forces that shape the

evolution of geographic variation in a marine snail. In: *Microevolution Rate, Pattern,*

*Process*, Contemporary Issues in Genetics and Evolution (eds. Hendry, A.P. & Kinnison,

M.T.). Springer Netherlands, Dordrecht, pp. 321–337.

Tüzün, N., Beeck, L.O. de, Brans, K.I., Janssens, L. & Stoks, R. (2017). Microgeographic

differentiation in thermal performance curves between rural and urban populations of an

aquatic insect. *Evolutionary Applications*, 10, 1067–1075.

Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of*

*Statistical Software*, 36.

West-Eberhard, M.J. (2003). *Developmental Plasticity and Evolution*. OUP USA.

Wilder, A.P., Palumbi, S.R., Conover, D.O. & Therkildsen, N.O. (2020). Footprints of local

adaptation span hundreds of linked genes in the Atlantic silverside genome. *Evolution*

*Letters*, 4, 430–443.

Williams, D.R., Rast, P. & Bürkner, P.-C. (2018). Bayesian meta-analysis with weakly

informative prior distributions. *PsyArXiv*. doi:10.31234/osf.io/7tbrm

Yamahira, K. & Conover, D.O. (2002). Intra- vs. interspecific latitudinal variation in growth:

adaptation to temperature or seasonality? *Ecology*, 83, 1252.

**Table Captions**

**Table 1:** Results of Bayesian metaregression mixed model with effect size fit as a function of

trait, gradient, and organism class. Random effects included trait nested in paper and paper alone,

as well as a phylogenetic species effects. Shown is the intercept and the effect of each level of a

covariate along with their estimated error and 95% credible intervals.

**Figure Captions**

**Figure 1:** Conceptual illustration of four types of phenotypic responses for two populations of a marine fish distributed along the Atlantic coast of the United States. Each row represents four related, but different responses: a) environment alone (phenotypic plasticity without local adaptation), b) a genetic by environment response, c) countergradient variation, d) and cogradient variation. The first two columns illustrate how each of the four responses manifest in different environments—shown is the expected phenotypic responses of both populations in their home environments ("wild") and if they were exposed to a reciprocal transplant experiment in their respective home habitats ("reciprocal transplant"). We also include the reaction norm for each response, where the base of the arrow indicates a population in its home environment and the arrowhead its away environment. Notice that for the counter-gradient variation example, both populations are the same size in the wild (i.e., perfect compensation), illustrating one way that local adaptation can be cryptic.

**Figure 2**: Posterior distributions of effect size estimates from random-effects meta-analysis for counter- and cogradient variation. The circle indicates the mean and the diamond the median of the posterior distribution (overlapping in countergradient) and the solid, black line, the 95% credible intervals. The mean and credible interval values are also indicated with text on the right. A total of 140 of the largest values (of 50,000 in each distribution) were removed because they extended beyond the x-axis margin (see Fig. S5 for all data points). Purple represents the posterior distribution for the effect size of countergradient variation and green, cogradient variation. The dashed line illustrates an effect size of 1.

**Figure 3:** Posterior distributions of estimated marginal means for the covariates in a

metaregression analysis for countergradient variation. Yellow distributions depict taxonomic

class, green distributions depict environmental gradients, and purple distributions depict trait

covariates. The black circle indicates the mean of the posterior distribution (medians omitted

because they overlapped with means). The thick solid, horizontal black line depicts the 50%

credible intervals for each covariate and the grey, thinner line the 95% credible intervals (also

indicated with text to the right). A total of the 314 largest values (of 60,000 thinned to 6,000 in

each distribution) were removed because they extended beyond the x-axis margin (all points can

be viewed in Figure S6). The dashed line illustrates each posterior distribution's relationship to

the study-wide effect size of 1.03 estimated from the random effects model and shown in Fig. 2.

**Figure 4:** Conceptual representation of two countergradient populations in their home

environments and the reaction norms of those populations in the home and away environments.

Also shown are the results of our compensation analysis. a) Represents an overcompensating

relationship where the northern population is larger than the southern population in all

environments and the slope of the dashed line connecting their home phenotype (arrow bases)

has a negative sign as it decreases from the larger northern population to the smaller southern

population. b) Represents a perfectly compensating relationship, where phenotypes are equal in

their respective home environments and thus the slope is flat. c.) Represents an

undercompensating relationship, where the southern population is larger than the northern

population in their respective home environments and the slope of the dashed line connecting

their home phenotype (arrow bases) has a positive sign as it increases from the smaller northern

population to the larger southern population. d.) represents the results of our compensation

analysis where overcompensating was defined as an effect size > 0.5, perfect as $0.5 \leq$ and $\geq$ -0.5,

and undercompensating < -0.5. Of these three responses, our data was distributed as 38.6%

overcompensating, 25.4% perfectly compensating, and 36.0% undercompensating. Points on the

left are the calculated effect sizes and are connected to a point at 0 to show a slope to match the

conceptual panels. The larger figure is bounded at -15 and 7 (removing 7 values) to better show

the results near 0 and perfect compensation, whereas the inset figure includes all data.

681 **Table 1**

| | Estimate | Est.Error | Q2.5 | Q97.5 |
|---|---|---|---|---|
| Intercept | 1.09 | 0.74 | 0.27 | 2.98 |
| Body size | 1.72 | 0.91 | 0.56 | 4.03 |
| Carotenoid concentration | 0.98 | 0.94 | 0.15 | 3.37 |
| Ciliary activity | 1.03 | 1.05 | 0.13 | 3.79 |
| Developmental rate | 1.48 | 0.68 | 0.55 | 3.17 |
| Gamete size | 2.78 | 2.27 | 0.54 | 8.74 |
| Growth rate | 1.16 | 0.51 | 0.47 | 2.41 |
| Metabolic rate | 1.43 | 0.69 | 0.52 | 3.12 |
| Phenology | 1.27 | 1.38 | 0.15 | 5.16 |
| Reproductive rate | 1.82 | 1.73 | 0.28 | 5.97 |
| Thermal response | 1.25 | 1.22 | 0.18 | 4.52 |
| Elevation | 1.08 | 0.83 | 0.24 | 3.2 |
| Latitude | 0.99 | 0.44 | 0.38 | 2.1 |
| Migration distance | 1.65 | 1.72 | 0.23 | 6.11 |
| Photoperiod | 1.6 | 1.5 | 0.24 | 5.43 |
| Predator presence | 1.49 | 1.37 | 0.23 | 5.24 |
| Salinity | 1.01 | 0.91 | 0.15 | 3.47 |
| Shade cover | 1.19 | 0.89 | 0.26 | 3.59 |
| Soil phosphate | 1.86 | 1.92 | 0.23 | 7.09 |
| Temperature | 2.5 | 1.43 | 0.74 | 6.19 |
| Urbanisation | 0.75 | 0.67 | 0.14 | 2.46 |
| Wave action | 1.38 | 1.14 | 0.26 | 4.29 |
| Amphibia | 1.23 | 0.66 | 0.4 | 2.87 |
| Anthozoa | 1.02 | 0.97 | 0.15 | 3.55 |
| Asteraceae | 1.24 | 1.41 | 0.14 | 4.76 |
| Bivalvia | 1.04 | 1.12 | 0.14 | 3.8 |
| Gastropoda | 1.07 | 0.78 | 0.26 | 2.98 |
| Insecta | 0.97 | 0.53 | 0.36 | 2.35 |
| Liliopsida | 1.85 | 1.89 | 0.23 | 7.08 |
| Malacostraca | 1.1 | 0.74 | 0.28 | 3.07 |
| Reptilia | 1.21 | 0.78 | 0.32 | 3.21 |

| | | | | |
|---|---|---|---|---|
| SD Paper Number | 2.31 | 0.37 | 1.65 | 3.11 |
| SD Paper Number:Trait | 2.28 | 0.25 | 1.9 | 2.86 |
| SD Species | 1.51 | 0.49 | 1.01 | 2.77 |

|  | **Wild** | **Reciprocal Transplant** | **Reaction Norm** |
|---|---|---|---|
| a) Environment alone (plasticity) | | North / South | Body size — North / South |
| b) Genetic x Environment | | North / South | Body size — North / South |
| c) Countergradient Variation | | North / South | Body size — North / South |
| d) Cogradient Variation | | North / South | Body size — North / South |

| | |
|---|---|
| Actinopterygii | 1.14 [0.43, 2.46] |
| Amphibia | 1.34 [0.36, 3.54] |
| Anthozoa | 1.13 [0.14, 4.35] |
| Asteraceae | 1.32 [0.13, 5.13] |
| Bivalvia | 1.14 [0.13, 4.51] |
| Gastropoda | 1.15 [0.23, 3.48] |
| Insecta | 1.05 [0.31, 2.81] |
| Liliopsida | 1.99 [0.22, 7.84] |
| Malacostraca | 1.21 [0.25, 3.74] |
| Reptilia | 1.32 [0.27, 3.86] |
| Carotenoid availability | 1.05 [0.35, 2.52] |
| Elevation | 1.05 [0.2, 3.33] |
| Latitude | 0.91 [0.38, 1.8] |
| Migration distance | 1.63 [0.19, 6.41] |
| Photoperiod | 1.57 [0.2, 5.88] |
| Predator | 1.49 [0.18, 5.5] |
| Salinity | 1.01 [0.12, 3.81] |
| Shade cover | 1.15 [0.21, 3.66] |
| Soil phosphate level | 1.82 [0.19, 7.48] |
| Temperature | 2.35 [0.67, 5.81] |
| Urbanisation | 0.7 [0.12, 2.33] |
| Wave action | 1.33 [0.23, 4.41] |
| Body shape | 0.98 [0.32, 2.35] |
| Body size | 1.5 [0.45, 3.75] |
| Carotenoid concentration | 0.9 [0.11, 3.49] |
| Ciliary activity | 0.95 [0.1, 3.77] |
| Developmental rate | 1.28 [0.47, 2.76] |
| Gamete size | 2.48 [0.43, 8.43] |
| Growth rate | 1.01 [0.39, 2.08] |
| Metabolic rate | 1.24 [0.43, 2.8] |
| Phenology | 1.14 [0.11, 4.88] |
| Reproductive rate | 1.69 [0.19, 6.36] |
| Thermal response | 1.15 [0.14, 4.42] |

Effect size

a) **Overcompensating**

Body size

North  South

b) **Perfectly compensating**

Body size

North  South

c) **Undercompensating**

Body size

North  South

d)