Title: Mathematical expressions describing enzyme velocity and inhibition at high enzyme

concentration

Author: Agustin Hernandez

Affiliation:

Unit for Integrated Research on Tropical Biodiversity – BIOTROP

Department of Genetics and Evolution

Centre for Biological and Health Sciences

Federal University of São Carlos

São Carlos (SP) Brazil

e-mail: agustin.hernandez@ufscar.br

Running title: Rate and inhibition at high enzyme concentration

Keywords: Continuous Fraction, Fast Equilibrium, Quasi-Steady-State, Dixon plots,

Cornish-Bowden plots, Catalyst Saturation

ABSTRACT

Enzyme behaviour is typically characterised in the laboratory using very diluted solutions of enzyme. However, *in vivo* processes usually occur at $[S_T] \approx [E_T] \approx K_m$. Furthermore, the study of enzyme action usually involves analysis and characterisation of inhibitors and their mechanisms. However, to date, there have been no reports proposing mathematical expressions that can be used to describe enzyme activity at high enzyme concentration apart from the simplest single substrate, irreversible case. Using a continued fraction approach, equations can be easily derived to apply to the most common cases in monosubstrate reactions, such as irreversible or reversible reactions and small molecule (inhibitor or activator) kinetic interactions. These expressions are simple and can be understood as an extension of the classical Michaelis-Menten equations. A first analysis of these expressions permits to deduce some differences at high *vs* low enzyme concentration, such as the greater effectiveness of allosteric inhibitors compared to catalytic ones. Also, they can be used to understand catalyst saturation in a reaction. Although they can be linearised following classical approaches, these equations also show some differences that need to be taken into account. The most important one may be the different meaning of line intersection points in Dixon plots. All in all, these expressions may be useful tools for the translation *in vivo* of *in vitro* experimental data or for modelling *in vivo* and biotechnological processes.

INTRODUCTION

Enzymes in the laboratory are usually assayed in conditions that are considered close to the *in vivo* conditions but, actually, they may differ significantly in some aspects from those governing their action in the cell. This includes variations in pH, crowding and ionic strength, among others. Particularly, it is very frequent to observe that, *in vivo*, enzymes are catalysing reactions in conditions where the concentration of substrate is close to that of the enzyme. On the contrary, in the laboratory, a typical enzyme assay is done in conditions where the concentration of the enzyme is as low as possible. This difference is a major one to understand the real activity of enzymes *in vivo*, because although it is usually assumed there exists a linear relationship between enzyme concentration and activity [1] that is not necessarily the case. The usual approaches for prediction of the initial velocity of an enzyme reaction assume either a fast substrate-enzyme binding equilibrium (FE), as originally proposed by Michaelis and Menten [2] or, more often, a quasi-steady state for the variation in the concentration of the enzyme-substrate complex (QSS), as the modification introduced by Briggs and Haldane [3]. In addition, both approaches assume that the concentration of the enzyme is negligible compared to that of the substrate. This is an important assumption when deriving the rate expressions and the one that dictates how enzyme assays are done in the laboratory. Several mathematical approaches have been proposed to derive expressions that could be valid in conditions where the concentration of the enzyme and substrate are close [4–7]. Moreover, a so-called inverse Michaelis-Menten expression has been proposed for interfacial enzyme kinetics [8]. However, to the best of this researcher`s knowledge, none yet have dealt with situations different from the simplest irreversible single substrate reaction.

In addition to an interest in describing the biochemistry of living organisms unperturbed, in pharmacology and biochemistry it is especially important the capacity to describe the effect of small molecules on enzyme activity, such as inhibitors and activators. Again, small molecules are assayed and characterised using low enzyme concentrations, but since target enzymes in the cell are in much closer ratios with their substrates, the effect of those small molecules *in vivo* may depart from the behaviour found *in vitro*.

From the point of view of the experimental biochemist, the mathematical expressions describing enzyme action *in vivo*, or in a biotechnological setting, should ideally use parameters that can be easily estimated in the laboratory. In addition, the possibility of

creating plots that can show differences between assay or living conditions can help in the

interpretation of results. From the point of view of teaching biochemistry, the use of

uncomplicated algebra and calculus is a bonus when explaining the derivation of those

expressions.

In this study, I present a series of mathematical expressions useful to describe the velocity

of monosubstrate enzyme reactions in different conditions when the concentration of the

enzyme is near that of the substrate. Those can be used with estimates of parameters

obtained in the laboratory at low enzyme-to-substrate concentration ratios to obtain

biologically relevant conclusions.

METHODS

Let be [S], [P] and [E] the respective free concentrations of substrate, product and enzyme

in the reaction depicted in Scheme 1. The microscopic kinetic constants are numbered

using odd figures for those representing forward reactions with respect to product

formation, while reverse reactions receive corresponding even numbering. Initial

conditions for the reaction (t=0) are assumed and, therefore, [P] ≈ 0 is also assumed,

leading to irreversibility of the reaction. Although incorrect, for simplicity`s sake only, that

last assumption is depicted by the corresponding microscopic kinetic constant as showing

zero value in the scheme. The presence of high concentrations of enzyme has been

approached by considering that the following mass conservation expression needs to be

taken into account: $[S_T] = [S] + [ES]$, where $[S_T]$ stands for total amount of substrate in the

reaction. In the case of a reversible reaction, $[P_T] = [P] + [EP]$ was also contemplated.

These conservation laws are considered in addition to the usual expression for the

conservation of enzyme, $[E_T] = [E] + [ES]$, where $[E_T]$ stands for the total amount of

enzyme in the system. Assuming QSS conditions for the concentration of the enzyme-

substrate complex ([ES]), the following expression can be derived (a detailed process of

derivation is shown in Supplemental Material 1A):

(1) $\dfrac{d[P]}{dt} = 0 = [ES]^2 - [ES] \cdot \left(K_m + [E_T] + [S_T] \right) + [S_T] \cdot [E_T]$

Solving this quadratic expression by regular means leads to

(2) $[ES] = \frac{1}{2}\left(\left(K_m + [S_T] + [E_T] \right) \pm \sqrt{\left(K_m + [S_T] + [E_T] \right)^2 - 4 \cdot [S_T] \cdot [E_T]} \right)$

However, a simpler approach to approximate (1) can be taken by using continued fractions (CF). This method uses simple algebra and has been used extensively in other areas [9, 10]. In general, a quadratic expression where the coefficient for the quadratic term is one, i.e. an expression of the type:

(3) $0 = x^2 + bx + c$

can be approximated for both negative and positive sign solutions as:

Positive sign solution (4) $x = b - \dfrac{c}{x}$

Negative sign solution (5) $x = \dfrac{c}{b - x}$

Recursive substitution of the variable x in the left side of the equality with the whole expression on that same side permits increasing degrees of approximation (for an example, see Supplemental Material 1B and Figure 1A). A first degree approximant can be obtained by neglecting the variable x in the left hand side of the equality. In this study, first degree approximants were used only. On the other hand, the negative sign solution of (2) is the only one valid in chemistry. This can be exemplified by the fact that, in the case of the positive sign, when $[S_T] = 0$, the value of $[ES]$ is non-zero:

(6) $[S_T] = 0, [ES] = K_m + [E_T]$

The CF approach for the negative sign solution has been used with (1) and all other reaction schemes treated in this study. Detailed derivation of the expressions can be found in Supplemental Material 1 and 2.

RESULTS

Derivation of expressions for monosubstrate irreversible reactions

The CF approximation to solving $[ES]$ in (1) under QSS assumption leads to the expression (Supplemental Material 1, B):

(7) $[ES] \simeq \dfrac{[S_T] \cdot [E_T]}{K_m + [E_T] + [S_T]}$

Expression (7) can be reduced to either that obtained using standard Michaelis-Menten

assumptions (sMM) [3]

(7b) $[ES] = \dfrac{[E_T][S]}{[S] + K_m}$

or the inverse Michaelis-Menten (iMM) [8]

(7c) $[ES] = \dfrac{[E][S_T]}{[E] + K_m}$

in cases where $[E_T] << [S_T]$ or $[E_T] >> [S_T]$, respectively. Therefore, (7) can be understood

as an extended Michaelis-Menten (eMM) bridging two extremes. Indeed, a comparison

between the values estimated for [ES] using one as a token value for K_m show that

expression (7) produces approximate values of [ES] ($[ES]_a$) that show a similar behaviour

to the real value throughout the range of $[E_T]$ and $[S_T]$ values (Figure 1A), although they

underestimate the true value of [ES]. Greater accuracy can be obtained by higher degree

approximants or a wise use of the sMM and iMM (Figure 1A and 1B). Nevertheless, a first

degree approximant already provides a much closer approximation to the value of [ES]

than sMM or iMM approaches when the whole range of $[S_T]$ and $[E_T]$ is taken into account

(Figure 1A and 1B). The relative value of the error for (7) was found never to exceed 0.4

(Figure 1B), while that for sMM and iMM increased exponentially as conditions departed

from the assumed ones for their derivation.

When $[E_T]$ was assumed constant at a very low concentration (e.g. $[E_T] = 0.001 \cdot K_m$), it

could be observed that $[ES]_a$ from both sMM and eMM produced curves that overlapped

the values obtained using the exact solution (expression (2)) over the range $0 \geq [S_T] \leq$

$10 \cdot K_m$, while iMM generated gross overestimations (Figure 1C). As $[E_T]$ is increased, the

curves generated by sMM and eMM separated from the true value of [ES]. For example, if

$1 = [E_T] = K_m$, sMM produced moderate overestimations while eMM underestimated the

value of [ES] (Figure 1D). These deviations tended to asymptotically converge with the

true value of [ES] at high $[S_T]$. This behaviour was exacerbated if greater $[E_T]$ were

contemplated (Figure 1E). A similar phenomenon can be observed if $[S_T]$ is kept constant

and curves are created over ranges of $[E_T]$ similar to those used in Figures 1C to 1E for

$[S_T]$ (data not shown). However, in that case, sMM is the expression providing gross

overestimations, while iMM produces moderate overestimations, and eMM keeps on

producing moderate underestimations of the true value of [ES].

QSS assumption may not be appropriate in all cases. Nevertheless, the CF approach can

also be used using FE assumptions and, in the present case, leads to the similar

expression (supplemental material 1C):

(8) $[ES] \simeq \dfrac{[S_T] \cdot [E_T]}{K_s + [E_T] + [S_T]}$

Expressions (7) and (8) can be used to provide rate laws for a monosubstrate irreversible

reaction under the conditions stated (Table 1). In particular, under QSS assumption:

(9) $\dfrac{d[P]}{dt} = \dfrac{V_{max} \cdot [S_T]}{K_m + [E_T] + [S_T]}$

As it is easily observed in (9), the catalytic efficiency, i.e. the value of the kinetic constant

for the reaction when it shows first order kinetics, is not V_{max}/K_m but $V_{max}/(K_m+[E_T])$. This

also leads to the observation that K_m does not coincide with either the concentration of

substrate providing $1/2V_{max}$ or the abscissa value of the intersection point between the

lines describing the velocity of the reaction when first order and when zero order. In both

cases, it is easily demonstrated that the value for both is $K_m+[E_T]$ (Figure 2A).

Reversible monosubstrate reactions

Most chemical reactions in a cell are reversible. The CF approach, assuming FE

conditions and two intermediate enzyme complexes ($[ES]$ and $[EP]$), can be used to

calculate the velocity of a reversible enzyme-catalysed reaction in the presence of both

substrate and product (Table 1, Supplemental Material 1D). The expression derived is

similar to the one derived assuming $[E_T] \ll [S_T]$. Further, if $[E_T] \approx 0$, the eMM expression

can be reduced to the sMM one. However, if expression in Table 1 is rearranged to obtain

a single denominator, the additional term in $[E_T]$ becomes very complex:

(10)

202
$$v = \frac{(V_1[S_T]([E_T]+K_P)-V_2[P_T]([E_T]+K_S))}{[E_T]([E_T]+[S_T](1+\frac{V_1}{V_2})+[P_T](1+\frac{V_2}{V_1})+K_S+K_P)+[S_T]K_P(1+\frac{V_1}{V_2})+[P_T]K_S(1+\frac{V_2}{V_1})+[S_T][P_T]\frac{(V_1+V_2)^2}{(V_1V_2)}+K_S K_P}$$

This tendency to complexity is shared with other approaches. Thus, in the hands of this

researcher, assuming other conditions, such as QSS, leads to expressions that depart

from the simplicity scope of this work (not shown). Nevertheless, those may still describe

the process accurately.

Integrated form of the QSS, eMM expression for irreversible monosubstrate

reactions.

Expression (9) is amenable to integration and, as expected, it also provides an expression

similar to that obtained using standard assumptions (Table 1 and Supplementary Material

1E):

(11) $V_{max} \cdot t = \left[S_T^0 \right] - \left[S_T^t \right] + \left(K_m + [E_T] \right) ln\left(\frac{\left[S_T^0 \right]}{\left[S_T^t \right]} \right)$

Needless to say that, since there is necessarily no changes in the total amount of enzyme

over time, no expression for the integrated form of (9) using $[E_T]$ as a variable is

meaningful.

Reactions at constant $[S_T]$

Similar to the iMM, velocity estimations by eMM at constant $[S_T]$ shows a hyperbolic

behaviour with respect to the amount of $[E_T]$ in the system (Fig. 2A). Therefore, under

those conditions, there exists a maximum asymptotic catalytic velocity attainable that can

be defined as

(12) $C_{max} = \left[S_T \right] k_3$

Similarly, we could define as Relative Catalytic Velocity (RCV) the *ad-hoc* parameter

(13) $RCV = \frac{v_e - v_u}{C_{max}}$

Where v_e is the experimentally observed enzyme-catalysed velocity, v_u is the velocity

observed in the absence of enzyme under the same conditions, and C_{max} the maximum

catalysed velocity attainable at the set [S_T]. Under most circumstances, v_u is several

orders of magnitude smaller than both v_e and C_{max}; therefore, RCV may be safely

approximated by the ratio v_e/C_{max}. Hence, RCV should obey:

(14) $RCV = \dfrac{[E_T]}{[E_T] + [S_T] + K_m}$

From this, it is possible to demonstrate that, at constant concentration of [S_T], the

concentration of enzyme needed to attain a certain proportion of C_{max} (RCV) would be:

(15) $[E_T] = \dfrac{RCV}{1 - RCV}\left(K_m + [S_T]\right)$

This expression shows that, while RCV is small, the amount of [E_T] necessary to attain it

increases in a near-linear fashion $([E_T] \approx RCV(K_m + [S_T]))$, but as the values of RCV

progress, the values for [E_T] increase exponentially (Figure 2B).

An example for the use of these expressions could be yeast soluble pyrophosphatase

isoform 1 (Ipp1p). This is a well characterised enzyme that hydrolyses Mg•pyrophosphate

into two orthophosphate molecules in a *bona fide* irreversible reaction. The K_m of that

enzyme is typically estimated in the micromolar range, 8 µM being reported earlier [11].

The concentration of the enzyme can be estimated as ~2 µM from the values reported for

its median abundance (42812 molecules/cell) and average cell volume (42 µm^3) [12]. The

pyrophosphate concentration in the cell is not precisely known, but probably ranges

between 1 and 100 µM [13], with 10 µM being a safe option. Assuming these conditions to

be representative of the cell ones, Ipp1p would be working at RCV ≈ 0.1. Similarly, figures

ranging from 0.001 (Pyk2p) to 0.3 (Fba1p) can easily be obtained for many of the enzymes

involved in yeast glycolysis [14].

Linearisations of eMM expressions

Although largely superseded by non-linear regression for parameter estimation,

linearisations of the rate equations are still useful to show graphically differences between

cases or reaction conditions. The CF and QSS-derived rate law (9), and its analogous

obtained under FE assumption (Table 1), can also be linearised provided either [E_T] or [S_T]

are kept constant (Table 1 and Figure 2C and 2D). Among these linearisations are the

most commonly ones, namely Lineweaver-Burk and Hanes linearisations. Also, similar to

those linearisations, the slopes and intersection points are symmetrical, i.e., the
intersection point from a Lineweaver-Burk plot is equivalent to the slope from a Hanes plot
and vice-versa.

Inhibition and activation expressions.
CFs were also used to derive expressions for the most common linear inhibition
mechanisms under QSS assumptions (Supplementary Material 2). However, in this case, it
was also assumed that $[I] >> [E_T] \approx [S_T]$, which means that $[I_T] \approx [I]$. This assumption
simplifies the expressions and may well be a real situation in the cell.
In all cases, the expressions derived were similar to those obtained under the sMM
assumption of negligible concentration of enzyme (Table 2). Noticeably, the enzyme
concentration in the denominator also formed a term with the concentration of inhibitor and
its equilibrium constant in the case of allosteric inhibition mechanisms (uncompetitive and
mixed uncompetitive-competitive). When $[ES]_a$ was plotted versus the whole range of $[S_T]$
and $[E_T]$, it was observed that a competitive inhibitor at the same conditions of
concentration and binding constant values, was much less effective than inhibitors with
mechanisms that implied allosteric interactions (Figure 3A). This was also observed in
plots with sMM expressions but it was less dramatic (data not shown). Further, using eMM
expressions, a comparison between the expected behaviour of those types of inhibitors at
$[E_T] >> K_m$ and $[E_T] \approx K_m$ revealed that, in the case of $[E_T] >> K_m$, at $[S_T]$ up to *ca 2x*K_m,
competitive inhibitors are more effective than uncompetitive inhibitors. This was not
surprising since sMM expressions already revealed that behaviour (data not shown).
However, at $[E_T] \approx K_m$, uncompetitive inhibitors were more effective than competitive
inhibitors in the whole $[S_T]$ range (Figure 2B and 2C). On the other hand, mixed
competitive-uncompetitive inhibitors showed an intermediate situation and were
consistently predicted more effective than competitive inhibitors at any concentrations of
$[E_T]$ and $[S_T]$, in agreement with known behaviour revealed by sMM expressions (data not
shown).

Non-linear inhibition, activation and linearisation of eMM inhibition expressions
Similar to the previous eMM equations and to those obtained under sMM assumptions,
eMM equations describing inhibition can be linearised to provide Lineweaver-Burk or
Hanes plots (Figure 4). Further, similar to the non-inhibition situation, the values for the
slopes and intercepts look transposed when comparing those two types of linearisation.

Furthermore, the CF approach can be extended to the nonlinear types of inhibition or
activation (Table 2). In the case of nonlinear small molecule-enzyme interactions,
Lineweaver-Burk and Hanes linearisations provide straight lines similar to those obtained
with sMM expressions. Moreover, when Dixon (DX) or Cornish-Bowden (CB) plots are
used, bent lines are produced, in agreement with what is observed using sMM
expressions.

Being probably the most informative, these types of plots were also checked for
consistency with classical expressions in the case of linear inhibition. CB plots from eMM
expressions produced straight lines that, in the case of allosteric inhibitors, intersected in
the second or third quadrant or on the negative side of the abscissa (depending on the
value of α); on the contrary, in the case of competitive inhibitors, parallel lines were
observed (Figure 5). This is similar to what is described for sMM expressions [15]. The
absolute value of the abscissa at the intersection point of the lines was estimated to be
equal to the allosteric equilibrium constant for the binding of the inhibitor to the enzyme
(Figure 5 and Table 3). All this is identical to what is found using sMM expressions.
However, important differences were observed in Dixon plots. Intersection of the lines did
not follow the pattern expected from sMM expressions. Thus, uncompetitive inhibition
equations showed convergent lines on DX plots that crossed in the third quadrant (Figure
5B) while parallel lines are typical from sMM equations. In appearance, competitive
inhibition, mixed uncompetitive-competitive inhibition and non-competitive inhibition
expressions produce DX plots that agree with their sMM counterparts. However, the
abscissa absolute values for the line intersection points are all dependent on the values of
K_m and $[E_T]$ in the system (Table 3). Only in the case of non-competitive inhibition, the
abscissa absolute value corresponds to K_i, as it is the case of sMM expressions.

DISCUSSION

Derivation of expressions predicting the velocity of monosubstrate enzyme-catalysed
reactions was found possible using a simple algebraic approach. Further, the expressions
were similar to those already in use (sMM) and could be reduced to those under the
assumption of negligible concentration of $[E_T]$. This helps them to be understood as easy
to use extensions of the classical expressions. Nevertheless, these equations only provide
approximations to the real value, and, therefore, care should be taken that assumptions
are valid when using them. In any case, the expression proposed to estimate the

concentration of the ES complex in a monosubstrate irreversible reaction (7) is identical to

those proposed earlier [4,5,16,17] and was found to approximate accurately the true value

of [ES] from (2) provided $\left(K_m + [E_T] + [S_T] \right) \gg [ES]$. It is true that expressions (7) and (9) are

not new but, to the best of this researcher's knowledge, this is the first report were the

continued fraction approach was used to obtain it. This is a simple approach that is

capable of different degrees of accuracy, if needed. Also, its simplicity may help students

to understand how expressions are derived and experimental researchers to get familiar

with them. Furthermore, a wide range of expressions are made available and not only an

equation describing the simplest monosubstrate, irreversible case. That should open the

possibility for their use in a wide variety of conditions and studies.

Some previous proposals were based on macroscopic parameters different from those

experimental biochemists are used to obtain [18]. While the estimation of different

parameters may just be a question of use and learning, undoubtedly the possibility of

using the same approaches and parameters to extrapolate enzyme behaviour in

conditions difficult to mimic in the test tube may be considered an advantage.

The introduction of [E_T] as a reactant in the denominator of the equations in this study

leads to the concept of enzyme saturation in a reaction. This means that increasing the

concentration of enzyme in a system above a certain amount may not result in meaningful

increases in velocity. Therefore, equation (14) can be a useful tool to understand the

cellular effort needed to modulate cell concentrations of metabolites, the feasibility in each

context of gene regulation of protein expression or to evaluate the investment needed to

attain an acceptable velocity in a biotechnological setting.

Linearisations are used extensively by biochemists to show results and point to differences

in enzyme action. The expressions presented in this work can also be linearised using, at

least, some of the common approaches in use for sMM. They provide lines that, in most

cases, behave just like the sMM ones. However, as other linearisations are tested,

exceptions, like those observed in Dixon plots, may appear.

One particularity of these linearisations for a simple irreversible reaction is that, at least

theoretically, it could be possible to estimate the value of k_3 (or k_{cat}) in non-pure samples,

such as extracts. That determination would need sufficient different amounts of extract

assayed for activity maintaining a constant [S_T]. The velocities obtained could be used in a

plot of [E_T]/v *vs* [E_T], where [E_T] can be expressed as fold over the sample having the

smallest amount of extract. The reciprocal of the slope of that straight line would equal

$[S_T]k_3$. Since $[S_T]$ is a known parameter in the assay, k_3 could be estimated. Nevertheless,

maintaining the QSS assumption valid in those experiments may well be the major hurdle.

Optimally, to ensure accuracy of the estimation, several of the determinations should need

to fall in the non-linear part of the curve. In other words, those determinations should need

to be made in conditions where $[S_T]$ is close to that of $[E_T]$ and, thus, it can be very difficult

to assume $[P] \approx 0$ or that variations in $[ES]$ are negligible during the assay. In any case, in

special cases, it might be a possibility.

Although similar to the classical expressions, the equations presented here already

revealed some differences that may be important to understand enzyme and inhibitor

action *in vivo*. The value of K_m in sMM it is often described as (and sometimes used as a

definition) the concentration of substrate providing $1/2V_{max}$. On the other hand, the

catalytic efficiency (the slope of the line describing first order kinetics), is a parameter often

used to compare catalysis on different substrates and even isoenzyme performance.

However, eMM equations showed that, in a general setting, $[E_T]$ needs to be taken into

account in those cases. While the issue of the K_m may not have important consequences

in experimental biochemistry and the values of the catalytic efficiencies at $[E_T] = 0$ could

still be used to compare different catalytic situations, other differences may be more

relevant. For example, the increase in inhibitory power of uncompetitive inhibitors that is

only observed at $[E_T] \approx [S_T] \approx K_m$. That difference, together with the general differences in

effect between allosteric and catalytic inhibitors predicted by these expressions, may be

useful to understand inhibitor action *in* vivo or to direct the search for new drugs in a more

selective way.

Finally, these expressions cover only a small part of the situations that can be faced in

relation to enzyme kinetics. Among the most important ones not covered by this work are

multisubstrate reactions and allosteric or cooperative enzymes. Efforts are in place now to

deduce equations able to attend to those cases. On the whole, the expressions proposed

here may be useful tools for the translation *in vivo* of *in vitro* experimental data or for

modelling *in vivo* and biotechnological processes.

Acknowledgements

The author thanks Ms E. A. de Magalhães for her help in this study. This work is dedicated

to late Drs D. T. Cooke and D. T. Clarkson for lifelong teaching through example.

The author declares not to have received funds for the development of this study and no

conflicts of interest.

REFERENCES

[1] Acerenza L & Kacser H (1990) Enzyme kinetics and metabolic control. A method to test and quantify the effect of enzymic properties on metabolic variables. *Biochem J* **269**, 697–707.

[2] Michaelis L & Menten MML (2013) The kinetics of invertin action. 1913. *FEBS Lett* **587**, 2712–2720.

[3] Briggs GE & Haldane JB (1925) A Note on the Kinetics of Enzyme Action. *Biochem J* **19**, 338–339.

[4] Cha S & Cha C-JM (1965) Kinetics of Cyclic Enzyme Systems. *Mol Pharmacol* **1**, 178–189.

[5] Borghans JA, de Boer RJ & Segel LA (1996) Extending the quasi-steady state approximation by changing variables. *Bull Math Biol* **58**, 43–63.

[6] Bajzer Z & Strehler EE (2012) About and beyond the Henri-Michaelis-Menten rate equation for single-substrate enzyme kinetics. *Biochem Biophys Res Commun* **417**, 982–985.

[7] Kargi, F. (2009) Generalized rate equation for single-substrate enzyme catalyzed reactions. *Biochemical and biophysical research communications* **382**.

[8] Kari J, Andersen M, Borch K & Westh P (2017) An Inverse Michaelis–Menten Approach for Interfacial Enzyme Kinetics. *ACS Catal* **7**, 4904–4914.

[9] Hayden TL (1965) Continued fraction approximation to functions. *Numer Math* **7**, 292–309.

[10] Rosen D & Shallit J (1978) A Continued Fraction Algorithm for Approximating All Real Polynomial Roots. *Mathematics Magazine* **51**, 112–116.

[11] Baykov AA & Avaeva SM (1974) Yeast inorganic pyrophosphatase: studies on metal binding. *Eur J Biochem* **47**, 57–66.

[12] Ho B, Baryshnikova A & Brown GW (2018) Unification of Protein Abundance Datasets Yields a Quantitative Saccharomyces cerevisiae Proteome. *Cell Syst* **6**, 192-205.e3.

[13] Heinonen JK (2001) *Biological Role of Inorganic Pyrophosphate* Kluwer Academic Publishers, Dordrecht.

[14] Smallbone K, Messiha HL, Carroll KM, Winder CL, Malys N, Dunn WB, Murabito E, Swainston N, Dada JO, Khan F, Pir P, Simeonidis E, Spasić I, Wishart J, Weichart D, Hayes NW, Jameson D, Broomhead DS, Oliver SG, Gaskell SJ, McCarthy JEG, Paton

NW, Westerhoff HV, Kell DB & Mendes P (2013) A model of yeast glycolysis based on

a consistent kinetic characterisation of all its enzymes. *FEBS Lett* **587**, 2832–2841.

[15] Cornish-Bowden A (1974) A simple graphical method for determining the inhibition

constants of mixed, uncompetitive and non-competitive inhibitors. *Biochem J* **137**,

143–144.

[16] Schnell S & Maini PK (2000) Enzyme kinetics at high enzyme concentration. *Bull Math

Biol* **62**, 483–499.

[17] Pedersen MG & Bersani AM (2010) Introducing total substrates simplifies theoretical

analysis at non-negligible enzyme concentrations: pseudo first-order kinetics and the

loss of zero-order ultrasensitivity. *J Math Biol* **60**, 267–283.

[18] Patsatzis DG & Goussis DA (2019) A new Michaelis-Menten equation valid

everywhere multi-scale dynamics prevails. *Math Biosci* **315**, 108220.

Supporting Information

Supplementary Materials 1.

Derivation of the equations for the simple cases: The exact solution for irreversible QSS

monosubstrate reaction, single monosubstrate irreversible FE eMM equation, single

monosubstrate irreversible QSS eMM equation, reversible single monosubstrate FE eMM

equation, and integrated QSS eMM equation.

Supplementary Materials 2

Derivation of the eMM equations for inhibition and activation under QSS. Competitive

linear inhibition, uncompetitive linear inhibition, mixed competitive-uncompetitive linear

inhibition, non-competitive linear inhibition, and general inhibition/activation model.

Table 1. Comparison of velocity expressions for monosubstrate enzyme-catalysed reactions.

	sMM	eMM	Linearisations (eMM)
FE (Irr.)	$v = \dfrac{V_{max} \cdot [S_T]}{[S_T] + K_S}$ (16)	$v = \dfrac{V_{max} \cdot [S_T]}{[E_T] + [S_T] + K_S}$ (9)	$\dfrac{1}{v} = \dfrac{1}{[S_T]} \cdot \dfrac{[E_T] + K_S}{V_{max}} + \dfrac{1}{V_{max}}$ (17) $\dfrac{1}{v} = \dfrac{1}{[E_T]} \cdot \dfrac{[S_T] + K_S}{C_{max}} + \dfrac{1}{C_{max}}$ (18) $\dfrac{[S_T]}{v} = [S_T]\dfrac{1}{V_{max}} + \dfrac{[E_T] + K_S}{V_{max}}$ (19) $\dfrac{[E_T]}{v} = [E_T]\dfrac{1}{C_{max}} + \dfrac{[S_T] + K_S}{C_{max}}$ (20)
FE (Rev.)	$v = \dfrac{V_1[S]}{[S]\left(1 + \dfrac{V_1}{V_2}\right) + K_S} - \dfrac{V_2[P]}{[P]\left(1 + \dfrac{V_2}{V_1}\right) + K_P}$ (21)	$v = \dfrac{V_1[S_T]}{[E_T] + [S_T]\left(1 + \dfrac{V_1}{V_2}\right) + K_S} - \dfrac{V_2[P_T]}{[E_T] + [P_T]\left(1 + \dfrac{V_2}{V_1}\right) + K_P}$ (22)	
QSS (Irr.)	$v = \dfrac{V_{max} \cdot [S_T]}{[S_T] + K_m}$ (23)	$v = \dfrac{V_{max} \cdot [S_T]}{[E_T] + [S_T] + K_m}$ (24)	$\dfrac{1}{v} = \dfrac{1}{[S_T]} \cdot \dfrac{[E_T] + K_m}{V_{max}} + \dfrac{1}{V_{max}}$ (25) $\dfrac{1}{v} = \dfrac{1}{[E_T]} \cdot \dfrac{[S_T] + K_m}{C_{max}} + \dfrac{1}{C_{max}}$ (26) $\dfrac{[S_T]}{v} = [S_T]\dfrac{1}{V_{max}} + \dfrac{[E_T] + K_m}{V_{max}}$ (27) $\dfrac{[E_T]}{v} = [E_T]\dfrac{1}{C_{max}} + \dfrac{[S_T] + K_m}{C_{max}}$ (28)
QSS (Irr.)	$V_{max} \cdot t = [S_T^0] - [S_T^t] + K_m ln\left(\dfrac{[S_T^0]}{[S_T^t]}\right)$ (29)	$V_{max} \cdot t = [S_T^0] - [S_T^t] + (K_m + [E_T]) ln\left(\dfrac{[S_T^0]}{[S_T^t]}\right)$ (11)	$\dfrac{([S_T^0] - [S_T^t])}{t} = V_{max} - (K_m + [E_T]) ln\left(\dfrac{[S_T^0]}{[S_T^t]}\right)\left(\dfrac{1}{t}\right)$ (30)

455 FE: Fast Equilibrium assumption, QSS: Quasi-steady state assumption, Irr.: Irreversible reaction, Rev.: Reversible reaction, sMM:
456 standard Michaelis-Menten, eMM: extended Michaelis-Menten
457

Table 2. Comparison of velocity expressions for monosubstrate, irreversible, enzyme-catalysed reactions affected by a non-covalent modifier (inhibitor or activator). All expressions derived assuming QSS.

Mechanism		sMM	eMM
Linear inhibition	Competitive	$$v = \dfrac{V_{max}[S_T]}{[S_T] + K_m\left(1 + \dfrac{[I]}{K_i}\right)} \quad (31)$$	$$v = \dfrac{V_{max}[S_T]}{[E_T] + [S_T] + K_m\left(1 + \dfrac{[I]}{K_i}\right)} \quad (32)$$
	Uncompetitive	$$v = \dfrac{V_{max}[S_T]}{[S_T]\left(1 + \dfrac{[I]}{\alpha K_i}\right) + K_m} \quad (33)$$	$$v = \dfrac{V_{max}[S_T]}{([E_T] + [S_T])\left(1 + \dfrac{[I]}{\alpha K_i}\right) + K_m} \quad (34)$$
	Mixed Competitive-Uncompetitive	$$v = \dfrac{V_{max}[S_T]}{[S_T]\left(1 + \dfrac{[I]}{\alpha K_i}\right) + K_m\left(1 + \dfrac{[I]}{K_i}\right)} \quad (35)$$	$$v = \dfrac{V_{max}[S_T]}{([E_T] + [S_T])\left(1 + \dfrac{[I]}{\alpha K_i}\right) + K_m\left(1 + \dfrac{[I]}{K_i}\right)} \quad (36)$$
	Non-Competitive	$$v = \dfrac{V_{max}[S_T]}{([S_T] + K_m)\left(1 + \dfrac{[I]}{K_i}\right)} \quad (37)$$	$$v = \dfrac{V_{max}[S_T]}{([E_T] + [S_T] + K_m)\left(1 + \dfrac{[I]}{K_i}\right)} \quad (38)$$
General Activation or Inhibition	All	$$v = \dfrac{V_{max}[S_T]\left(1 + \dfrac{\beta[A]}{\alpha K_a}\right)}{[S_T]\left(1 + \dfrac{[A]}{\alpha K_a}\right) + K_m\left(1 + \dfrac{[A]}{K_a}\right)} \quad (39)$$	$$v = \dfrac{V_{max}[S_T]\left(1 + \dfrac{\beta[A]}{\alpha K_a}\right)}{([E_T] + [S_T])\left(1 + \dfrac{[A]}{\alpha K_a}\right) + K_m\left(1 + \dfrac{[A]}{K_a}\right)} \quad (40)$$

sMM: standard Michaelis-Menten, eMM: extended Michaelis-Menten

464 Table 3. Abscissa values for the intersection points of lines created using eMM linear
465 inhibition expressions in Dixon and Cornish-Bowden plots.

	Dixon (1/v *vs* [I])	Cornish-Bowden ([S$_T$]/v *vs* [I])
Competitive	$-K_i \dfrac{[E_T]+K_m}{K_m}$	No intersection
Uncompetitive	$-\alpha K_i \dfrac{[E_T]+K_m}{[E_T]}$	$-\alpha K_i$
Mixed Competitive-Uncompetitive	$-\alpha K_i \dfrac{[E_T]+K_m}{[E_T]+\alpha K_m}$	$-\alpha K_i$
Non-Competitive	$-K_i$	$-K_i$

466
467

 Figure Legends

 Figure 1. Estimation of enzyme-substrate complex concentration using different

 mathematical expressions. **A** Approximated [ES] ([ES]$_a$) using the exact solution

 (expression (2) in main text; blue surface), extended Michaelis-Menten (eMM; expression

 (7) in main text; green surface), a second degree approximant (eMM, expression (19) from

 Supplemental Material 1; magenta surface), standard Michaelis-Menten (expression (7b)

 in main text, cyan surface) and inversed Michaelis-Menten (expression (7c) in main text;

 red surface). All surfaces calculated using K_m = 1. **B** Estimated error relative to the real

 value estimated with the exact solution and the extended Michaelis-Menten (eMM;

 expression (7); green surface), standard Michaelis-Menten (expression (7b); cyan surface)

 and inversed Michaelis-Menten (expression (7c); red surface). All surfaces calculated

 using K_m = 1. **C** Departure of [ES]$_a$ from the exact solution at a constant low enzyme

 concentration ([E$_T$] =0.001 x K_m). Solid black line: exact solution (expression (2) in main

 text); Dot-dashed black line: extended Michaelis-Menten (eMM; expression (7)); Dotted

 red line: standard Michaelis-Menten (expression (7b)); Dashed red line: inversed

 Michaelis-Menten (expression (7c)). **D** Departure of [ES]$_a$ from the exact solution at a

 constant enzyme concentration similar to K_m ([E$_T$] = K_m). Line identities as in C. **E**

 Departure of [ES]$_a$ from the exact solution at a constant high enzyme concentration ([E$_T$] =

 100 x K_m). Line identities as in C. All lines in C, D, and E calculated using K_m =1. Please

 note that eMM and sMM lines in panel C are obscured by the overlapping line

 corresponding to the exact solution.

 Figure 2. Hyperbolic and linear plots of eMM expressions for monosubstrate irreversible

 reactions. **A**. Direct plots of the estimated velocities (equation (9)) at constant [E$_T$] (Dot-

dashed black line) or constant $[S_T]$ (Dashed red line). Both lines calculated using $K_m = 1$.

Value for the concentration of the constant reactant was set to one (1). To help

visualisation, $k_3 = 1$ was used for constant $[E_T]$ while $k_3 = 0.5$ was used for constant $[S_T]$.

Other parameter values: $[E_T] = [S_T] = K_m = 1$. Cat. eff.: catalytic efficiency. **B** Concentration of

$[E_T]$ as a function of Relative Catalytic Velocity (RCV). Line calculated using expression

(14) from main text and $K_m = [S_T] = 10$. **C** Hanes plots of eMM rate expressions. Lines

drawn using $k_3 = K_m = [E_T] = 1$ (dot-dashed black line, expression (27)) or $k_3 = K_m = [S_T] = 1$

(dashed red line, expression (28)). **D** Lineweaver-Burk plots of eMM rate expressions.

Lines drawn using $k_3 = K_m = [E_T] = 1$ (dot-dashed black line, expression (25)) or $k_3 = K_m =$

$[S_T] = 1$ (dashed red line, expression (26)).

Figure 3. Behaviour of enzyme inhibition expressions obtained using CF approach. **A**

Approximated [ES] ($[ES]_a$) over $[S_T]$ and $[E_T]$. $[ES]_a$ obtained using extended Michaelis-

Menten (eMM; expression (7); green surface), competitive inhibition (expression (32);

magenta surface), uncompetitive inhibition (expression (34); blue surface) and mixed

competitive-uncompetitive/non-competitive inhibition (expressions (36) and (38); cyan

surface). **B** Velocity estimated from the different types of inhibition at a constant low

concentration of $[E_T] = 0.001$ x K_m. Solid black line, no inhibition (expression (9)); dashed

red line, competitive inhibition (expression (32)); dot-dashed black line, uncompetitive

inhibition (expression (34)); solid red line, mixed/non-competitive inhibition (expressions

(36) and (38)). **C** Velocity estimated from the different types of inhibition at $[E_T] = K_m$. Solid

black line, no inhibition (expression (9)); dashed red line, competitive inhibition (expression

(32)); dot-dashed black line, uncompetitive inhibition (expression (34)); solid red line,

mixed/non-competitive inhibition (expressions (36) and (38)). All surfaces and lines

calculated using $k_3 = K_m = 1$, $[I] = 10$ and $K_i = \alpha K_i = 9$.

Figure 4. Plots of eMM and sMM linearised equations for total (linear) and partial (non-linear) inhibition mechanisms. All plots correspond to mixed uncompetitive-competitive inhibition as a general representative of all mechanisms. Linear inhibition corresponds to expressions (35) (sMM) and (36) (eMM); non-linear inhibition corresponds to expressions (39) (sMM) and (40) (eMM); values of parameters for all lines as follows: $K_m = [E_T] = 1$, $k_3 = 0.75$, $K_i = 10$, $\alpha K_i = 5$ and $\beta = 0$ (linear) or $\beta = 0.25$ (non-linear). **A** Lineweaver-Burk plots. Solid black line, eMM non-linear inhibition; dotted black line, eMM linear inhibition; dash-dotted red line, sMM non-linear inhibition; dotted red line, sMM linear inhibition. **B** Hanes plots. Lines as in A. **C** Dixon plots, Lines as in A. **D** Cornish-Bowden plots. Lines as in A.

Figure 5. Dixon and Cornish-Bowden plots for eMM linear inhibition mechanisms. All lines drawn using equations (32) (competitive), (34) (uncompetitive), (36) (mixed competitive-uncompetitive) and (38) (non-competitive) with the following parameters: $K_m = [E_T] = 1$, $k_3 = 2$, $K_i = 10$, $\alpha K_i = 5$. Solid red lines, $[S_T] = 0.5$; dash-dotted black lines $[S_T] = 10$. Panels **A** and **E**, competitive inhibition; panels **B** and **F**, uncompetitive inhibition, panels **C** and **G**, mixed competitive-uncompetitive inhibition; panels **D** and **H**, non-competitive inhibition. Panels **A** to **D**, Dixon plots; panels **E** to **H**, Cornish-Bowden plots.

Scheme 1

$$E+S \underset{k_2}{\overset{k_1}{\rightleftharpoons}} ES \overset{k_3}{\underset{k_4=0}{\longrightarrow}} E+P$$

Figure 1

Figure 2

Figure 3

A

B

K_m = 1
k_3 = 1
[E_T] = 0.001
K_i = αK_i = 9
[I] = 10

C

K_m = 1
k_3 = 1
[E_T] = 1
K_i = αK_i = 9
[I] = 10

Figure 4

Figure 5