Title:

I-M-150847, a novel GLP-1 and GIP receptor dual agonist, improves glycemic control and reduces obesity in the rodent model of type 2 diabetes and obesity.

Authors:

Rathin Bauri [1, 2], Shilpak Bele, [1, 2], Jhansi Edelli[1], Neelesh C. Reddy[3], Sreenivasulu Kurukuti [4], Tom Devasia[2], Ahamed Ibrahim[5], Vishal Rai[3], Prasenjit Mitra[1,6†]

[1.] Dr. Reddy's Institute of Life Sciences, University of Hyderabad Campus, Hyderabad, Telangana, India, 500046

[2.] Manipal Academy of Higher Education, Manipal, Karnataka, India -576104.

[3] Indian Institute of Science Education and Research, Bhopal, 462066

[4] Department of Animal Biology, University of Hyderabad, India 500046

[5] Division of Lipid Chemistry, National Institute of Nutrition Hyderabad, India 500007.

[6†] corresponding author

Correspondence:
prasenjit.mitra01604@gmail.com

Running Title: A novel incretin receptor dual agonist ameliorates diabesity.

Abstract:

We report the discovery of a novel unimolecular glucagon-like peptide-1 (GLP-1) and glucose-dependent insulinotropic polypeptide (GIP) receptor dual agonist that exhibits potent glycemic control and weight loss in diet-induced obese mice. The dual agonism of GLP-1 and GIP receptor was achieved by replacing the tryptophan cage of exendin-4 with the C- terminal undecapeptide sequence of oxyntomodulin along with a single amino acid substitution from histidine to tyrosine at the amino terminus of the peptide. The structural modification places lysine 30 of the novel incretin agonist in frame with the corresponding lysine residue in the native GIP sequence. The novel incretin receptor dual agonist, named I-M-150847, induces rapid redistribution of GLP-1R at the plasma membrane following activation ensuring the maintenance of the receptor in a sensitized state. I-M-150847 promotes glucose-stimulated insulin exocytosis in cultured pancreatic beta cells and augments insulin-stimulated glucose uptake in mouse adipocytes. Chronic administration of I-M-150847 enhances insulin sensitivity, improves glycemic control, and achieves significant weight loss relative to the control or exendin-4-treated DIO-mice demonstrating the therapeutic efficacy of dual agonist in ameliorating type 2 Diabetes and Obesity.

Significance statement.

• Replacement of the Trp-cage with the C- terminal oxyntomodulin undecapeptide

along with the tyrosine substitution at the amino terminus converts the selective

GLP-1R agonist exendin-4 to a novel GLP-1R and GIPR dual agonist I-M-

150847.

• The GLP-1R and GIPR dual agonist I-M-150847 induces the expeditious

redistribution of GLP-1R at the plasma membrane following initial activation

thereby maintaining the receptor in a sensitized state.

• The incretin receptor dual agonist I-M-150847 enhances insulin sensitivity and

delivers superior glycemic control and weight loss compared to exendin-4 in the

rodent model of diabetes and obesity.

1   **Graphical Abstract:**

Rapid Resensitization of GLP-1R

Retention of GIPR function upon chronic exposure

Increased Insulinotropic Response

Enhanced Insulin Sensitivity

Reduction of Obesity

Introduction:

Incretin hormones, glucagon-like peptide-1 (GLP-1), and glucose-dependent insulinotropic polypeptide (GIP) lower the postprandial glucose as they potentiate glucose-stimulated insulin release from the pancreatic beta cells (Drucker, 2006),(Ranganath, 2008). The therapeutic efficacy of GLP-1 is presently well established and GLP-1R agonists (GLP-1RAs) have emerged as the medicine of choice for the treatment of type 2 Diabetes (T2DM) and obesity (Amori et al., 2007; Andersen et al., 2018; Astrup et al., 2009). However, not all patients receiving GLP-1RA therapy can attain long-term glycemic control and desired reduction of obesity necessitating a complementary therapeutic approach to attain proper metabolic benefits (Samms et al., 2020). The reduction of insulin resistance in the case of GLP-1RA therapy is primarily due to weight loss (Fonseca et al., 2019). In contrast, long-acting GIPR agonists (LAGIPRAs) contribute to enhanced insulin sensitivity through their action on adipose tissues (Samms et al., 2021; Varol et al., 2014). Consequently, to avail the benefits of both the incretin receptor agonists, hybrid peptides were designed incorporating residues from GLP-1 and GIP to confer dual GLP-1R and GIPR agonist activity in a single molecule (Bastin and Andreelli, 2019; DiMarchi, 2018; Finan et al., 2013). The dual agonists exhibit greater efficacy as compared to GLP-1R monoagonists in preclinical settings (Coskun et al., 2018; Finan et al., 2013) as well as in clinical trials (Frias et al., 2018),(Rizvi and Rizzo, 2022),(Jastreboff et al., 2022),(Rosenstock et al., 2021)

The approach to regulating energy balance through dual agonism exists in nature's repository (Pocai, 2012; Pocai et al., 2009). Oxyntomodulin, an endogenous dual

agonist of GLP-1R and Glucagon receptor (GCGR) displays superior bodyweight and glucose lowering effects than GLP-1R monoagonists upon enhancement of its plasma stability through chemical modifications (Kerr et al., 2010),(Muppidi et al., 2016). In a parallel approach, GIP residues were incorporated into the oxyntomodulin sequence to achieve a GIPR, GLP-1R, and GCGR triagonist response in a single hybrid molecule for the development of a potent multi-receptor agonist (Bhat et al., 2013).

In our present report, we employed a structural approach for the rational design of GLP-1R and GIPR dual agonists. The existing crystal structure reveals that the peptide ligands form alpha-helix as they bind to the extracellular domain of the incretin receptors. The affinity and specificity of the respective ligand and the receptor are determined by the interactions between the amino acid residues of the ligand and the binding site of the receptor (Parthier et al., 2007; Runge et al., 2008; Underwood et al., 2010). Our data reveal that the replacement of the tryptophan cage (trp-cage) of Exendin-4 tyrosine substituted at the amino terminus (ex-4Y1) with the C terminal undecapeptide sequence of Oxyntomodulin places the lysine 30 of the novel hybrid agonist in frame with the corresponding lysine residue in the native GIP sequence thereby converting the peptide to a GLP-1R-GIPR dual agonist. The novel incretin peptide exhibits potent GLP-1R and GIPR dual agonism with significantly diminished activity towards glucagon receptors. Sustained GLP-1R agonism from endosomes was earlier reported from our laboratory (Kuna et al., 2013), (Girada et al., 2017), In the present study, we demonstrate that I-M-150847- mediated GLP-1R signaling, contrary to exendin-4, is short-lived and facilitates rapid redistribution of GLP-1R at the plasma membrane following activation thereby maintaining the receptor in a sensitized state.

Our data shows that I-M-150847 enhanced insulin sensitivity upon chronic exposure in

cultured mouse adipocytes. In vivo, I-M-150847 displays superior glycemic control,

enhanced insulin sensitivity, and greater weight loss compared to exendin-4 highlighting

its promising efficacy for the treatment of T2DM and obesity

Results:

Generation of potent GLP-1R and GIPR dual agonists:

Our primary objective was to design a GLP-1R and GIPR dual agonist while minimizing

its effect on GCGR. The crystal structure of ligand coupled to GLP-1R and GIPR

extracellular domain (Parthier et al., 2007; Runge et al., 2008; Underwood et al., 2010),

structure-activity relationships of incretins (Finan et al., 2013; Patterson et al., 2013;

Yang et al., 2020), the similarities in their primary sequences as well as the sequence

differences in the middle and the C terminal region of the respective agonists guided us

to rationally design the hybrid peptide with potent GLP-1R and GIPR dual activity in a

single molecule. The unimolecular GLP-1R and GIPR dual agonism was achieved in

three steps. In the first step, at the C-terminus of the 9^{th} amino acid residue of I-M-

150844 that corresponds to the N terminal sequence of exendin-4, we added GIP-

specific amino acids Tyr^{10}, Ser^{11}, Ile^{12}, Ala^{13}, Met^{14}, Asp^{15}, Lys^{16}, Ile^{17}, His^{18}, Gln^{19},

Gln^{20}, Asp^{21}, Phe^{22}, Val^{23}, Asn^{24}, Trp^{25}, and Leu^{26} followed by GLP-1 specific residues

Val^{27}, lys^{28}, Gly^{29}, Arg^{30} and Gly^{31} at the C terminal end **(Table 1, Figure 1-Figure**

supplement 1). The resultant peptide I-M-150845 exhibits 76.78± 6.32% GLP-1R

activity, 86.86% ± 13.36 GIPR activity, and 11.5 ± 0.99 % GCGR activity measured

relative to liraglutide, GIP, and oxyntomodulin in the corresponding GLP-1R. GIPR and

GCGR luciferase reporter assays. In the next step, we replaced the GLP-1 specific

residues Val[27], lys[28], Gly[29], Arg[30,] and Gly[31] of I-M-150845 with the C- terminal

oxyntomodulin residues Met[27], Asn[28], Thr[29], Lys[30], Arg[31], Asn[32], Arg[33], Asn[34], Asn[35],

Ile[36], and Ala[37] to generate a new incretin agonist I-M-150846. The peptide showed

65.65 ± 6.7 % GIPR activity but a drastically attenuated GLP-1R activity **(Table 1,**

Figure 1- Figure supplement 2). To retain the GLP-1R activity, we removed the GIP-

specific residues Tyr[10], Ser[11], Ile[12], Ala[13], Met[14], Asp[15], Lys[16], Ile[17], His[18], Gln[19], Gln[20],

Asp[21], Phe[22], Val [23], Asn [24] from I-M-150846 and replaced them with exendin-4 specific

residues Leu[10], Ser [11], Lys[12], Gln[13], Met[14], Glu[15], Glu[16], Glu[17], Ala[18], Val[19], Arg[20], Leu[21],

Phe[22], Ile[23,] and Glu[24]. The novel incretin peptide I-M-150847 derived by this

replacement exhibited 122.95±2.04% GLP-1R activity, 108.09 ± 5.17 % GIPR activity,

and 14.93 ±1.68 % GCGR activity as described in human GLP-1R, GIPR, and GCGR

cell-based cAMP reporter assay using liraglutide, GIP and Oxyntomodulin as a positive

control in respective assays (**Table1, Figure 1-Figure supplement 3, Fig 1A, B, C**). To

ascertain the functional characteristics of I-M-150847, we compared its potency with

exendin F1 (Jones et.al 2018), and Tirzepatide, (Willard et.al 2020) in HEK 293T cell

lines expressing GLP-1R. As our data shows, I-M-150847, like exendin F1 and

Tirzepatide, is a full agonist of GLP-1R in cAMP accumulation assay (EC_{50} 16.35 nM)

(**Fig 1D).** In HEK 293T cells expressing GIPR, I-M-150847 also displays a full agonist

activity though its potency of cAMP accumulation (EC_{50} 28.90 nM) is right-shifted

relative to Tirzepatide (EC_{50} 0.609 nM) **(Fig 1E).** In contrast, we observed a drastically

reduced activity of I-M150847 towards GCGR in the HEK GCGR cAMP accumulation

assay **(Figure1- Figure supplement 4)**. The data thus demonstrates the potent GLP-

1R and GIPR dual agonism of the hybrid peptide I-M-150847 while exhibiting minimal

activity towards GCGR. In cultured pancreatic beta cells, I-M-150847 shows full agonist

activity and comparable potency to exendin-4 which underlines its efficacy in the

physiological milieu **(Fig 1F)**. LC/MS analysis, and purity, of the C terminal amidated

version of I-M-150847, are provided in **Figure1-Figure supplement 5,** and **Figure 1-**

Figure supplement 6.

I-M-150847 elicits short-term GLP-1R-mediated cAMP generation and facilitates

expeditious redistribution of the functional receptor at the plasma membrane.

GLP-1RAs exhibit variable efficiencies of internalization and trafficking (Jones et al.,

2018). We first reported the prolonged internalization of GLP-1R upon activation that is

associated with sustained GLP-1R mediated cAMP signaling upon treatment with GLP-

1- conjugated to tetramethyl rhodamine (GLP-1Tmr), liraglutide, and exendin-4 (Kuna et

al., 2013), (Girada et al., 2017). In our present study, we compared the trafficking and

signaling efficacy of I-M-150847 relative to GLP-1R monoagonist exendin-4 in cultured

pancreatic beta cells. We observed that, unlike exendin-4, cAMP generation upon

treatment with I-M-150847 is short-lived, as it maximizes at 10 minutes of activation and

recedes abruptly at 30 min after the agonist treatment **(Fig 2A).** Along with the short-

term cAMP generation, our data reveals contrasting post-endocytic sorting of GLP-1R

upon I-M-150847 treatment relative to exendin-4 **(Fig 2B).** At 0 min, we observed an

annular ring of GLP-1R GFP at the plasma membrane and a diffused distribution at the

cytoplasm **(Fig 2B i).** At 10 minutes following the treatment with exendin-4, there is a

complete loss of the annular ring as the GLP-1R GFP signal appears as punctate dots

in the cytoplasm demonstrating complete internalization of the receptor after activation

with the agonist. In contrast, when treated with I-M-150847 for 10 minutes, we observed

the presence of GLP-1R -GFP both as an annular ring at the plasma membrane and as

punctate dots in the cytoplasm (**Fig 2B ii)**. Following 30-minute treatment with exendin-

4, the punctate dots representing GLP-1R GFP is prevalent in the cytoplasm whereas in

I-M150847 treated cells, we observe the prominent annular ring at the plasma

membrane as well as a few GLP-1R GFP punctate dots at the cytoplasm mostly

distributed at the periphery near the plasma membrane **(Fig 2B iii).** The punctate GLP-

1R GFP dots are retained in the cytoplasm after 60 minutes of exendin-4 treatment;

however, in I-M-150847 treated cells, we observe the majority of GLP-1R -GFP

fluorescence at the plasma membrane as annular ring; the punctate GLP-1R GFP dots

are progressively much less in number with time and have a limited distribution in the

proximity of the plasma membrane **(Fig 2B iv).** The data thus demonstrates that I-M-

150847 exhibits reduced internalization and /or faster recycling of GLP-1R compared to

exendin-4 as the maximum GLP-1R- GFP signal is observed back at the plasma

membrane on a much shorter time course following activation with I-M-150847.

We first reported lysosome as one of the destinations of GLP-1R following

internalization of the receptor (Kuna et al., 2013) which was supported by subsequent

studies using immunoelectron microscopy (Jones et al., 2018). However endocytic

destination of GLP-1R was later found to be ligand specific as the receptor has been

shown to recycle back to the plasma membrane upon treatment with Exendin F1 (Jones

et al., 2018). In our present study, we compared the association of GLP-1R with

lysosomes at 45 minutes post-treatment with exendin-4 or I-M-150847. We observed

partial co-localization of GLP-1R -GFP with lysotracker red (colocalization index,

Pearson r=0.46 ±0.03) upon treatment with exendin-4 at 45 minutes post internalization

of the receptor **(Fig 2C).** In contrast, the localization of GLP-1R to lysosome was

significantly reduced upon treatment with I-M-150847 (colocalization index, Pearson

r=0.32 ±0.23; **p<0.01, n=5) **(Fig 2C).** To functionally assess the GLP-1R trafficking we

next carried out the receptor resensitization experiment **(Fig 2D).** The pancreatic beta

cells were treated with exendin-4 or I-M-150847 for 3 minutes followed by a wash with

KRB buffer. After 45 minutes of treatment, the cells were rechallenged with respective

agonists for 10 minutes and cAMP accumulation was evaluated. As **Fig 2D** shows,

cultured pancreatic beta cells with prior exposure to I-M-150847 respond more

efficiently to GLP-1R mediated cAMP generation relative to the cells that are pre-

exposed to exendin-4 during agonist rechallenge. We reported that I-M-150847-

mediated cAMP generation subsided 30 minutes post-treatment **(Fig 2A).** Aligned to

these findings, cAMP generation is attenuated to the basal level 45 minutes after

treatment with I-M-150847. Thereafter, upon the I-M-150847 rechallenge, we observed

an increase in cAMP generation (from 61.30 ± 11.96 to 342.3 ±61.68 pmol/mg protein,

5.58-fold increase, **p<0.01, one-way ANOVA, Tukey's multiple comparison test, n=3).

The increase of cAMP generation upon rechallenge signifies prompt redistribution of the

functional GLP-1R at the plasma membrane after initial activation. In contrast, similar

exendin-4 treatment resulted in a sustained cAMP generation that decreases from

255.3± 54.91 to 143.8 ± 33.39 pmol/mg protein, (n=3) after agonist rechallenge. The

data taken together demonstrate that the I-M-150847 treatment ensures efficient

receptor recycling that preserves a higher level of functional GLP-1R at the plasma

membrane relative to exendin-4 sufficient to initiate de novo second messenger

signaling upon I-M-150847 re-exposure. **Fig 2E** explains the mechanism of GLP-1R

trafficking upon activation by I-M-150847 vis-à-vis exendin-4.

Prolonged I-M-150847 exposure has a contrasting functional response in cultured

pancreatic beta cells and mouse adipocytes

We next evaluated whether the rapid redistribution at the plasma membrane has any

functional advantage over prolonged internalization of GLP-1R following activation by

agonists upon continuous exposure. As **Fig 3A** reveals, I-M-150847 significantly

enhances GSIS relative to exendin-4 after 30 minutes of incubation (33.01±1.94 fold

over basal upon I-M-150847 compared to 24.04 ±0.71 fold over basal upon exendin-4

treatment, **p<0.01, n=3) demonstrating its superior insulinotropic property. However,

upon prolonged exposure, we observed blunting of GSIS both upon exendin-4 and I-M-

150847 treatment **(Fig 3A).** The data thus underlines that GLP-1R induced GSIS in

BRIN-BD11 pancreatic beta cells is independent of the GLP-1R sorting dynamics as

both prolonged internalizations upon exendin-4-mediated activation or rapid

redistribution at the plasma membrane following I-M-150847 treatment has a similar

impact on GLP-1R desensitization upon chronic agonist exposure. The GSIS response

in BRIN-BD11 pancreatic beta cells is mostly GLP-1R specific; as we observed a 6.95-

fold enrichment of GLP-1R expression over GIPR in BRIN-BD11 pancreatic beta cells

(Figure 3- Figure supplement 1) In contrast, in cultured mouse adipocytes where

GIPR is predominant (McIntosh et al., 2012), the I-M-150847 treatment increases the

cAMP generation from 31.6±3.7 pmol to 72.44±2.535 pmol per mg protein, (***p< 0.001,

one-way ANOVA Tukey's multiple comparison test n=3) upon chronic exposure **(Fig**

3B). Since enhanced cAMP generation is indicative of increased lipolysis and

subsequent blunting of insulin sensitivity, we assessed insulin-stimulated glucose

uptake in 3T3L1 adipocytes following 16 h treatment with exendin-4 or I-M-150847.

Contrary to our postulation, we observed an increase in insulin-stimulated glucose

uptake upon I-M-150847 treatment (from 1.385±0.009 fold to 1.91±0.05 fold,

***$p<0.001$, one-way ANOVA, Tukey's multi comparison test n=3) **(Fig 3C)** thereby

demonstrating the efficacy of the incretin receptor dual agonist I-M-150847 in enhancing

insulin sensitivity in cultured adipocytes. The data thus signifies that the functional

outcome of chronic exposure to I-M-150847 is different in mouse adipocytes and

cultured pancreatic beta cells. The data further demonstrates that the enhancement of

insulin sensitivity upon I-M-150847 treatment is independent of the agonist-induced

cAMP response in cultured mouse adipocytes.

I-M-150847 outperforms Exendin-4 in imparting glycemic benefits and reducing

obesity.

To explore whether these in vitro observations translate to better regulation of glucose

homeostasis and weight loss in the rodent model of diabesity, we compared the effect

of I-M-150847 and exendin-4 in regulating hyperglycemia and obesity in C57BL/6 mice

fed on the high-fat diet. We carried out twice-daily dosing of 100nmol/ kg body weight

of exendin-4 or I-M-150847 for 28 days and assessed the metabolic parameters as

described in the flow diagram **(Fig 4A).** As our data reveals, 10 days of exendin-4

treatment reduces fasting blood glucose from 176.5 ±11.31 mg/dL glucose to 130 ±3.02

mg/dL compared to vehicle control (***$p<0.001$, one-way ANOVA, Tukey's multi

comparison test n=6) **(Fig 4B (ii)).** In comparison, after 10 days of I-M-150847

treatment, fasting blood glucose was reduced from 183 ±14.02 mg/dL glucose to 101.7

±2.17 mg/dL glucose (***$p<0.001$, one-way ANOVA, Tukey's multi comparison test,

n=6) **(Fig 4B (iii)**)). The results thus reveal that after 10 days of treatment, I-M-150847

outperforms exendin-4 in reducing fasting blood sugar in DIO mice (mean blood glucose

101.7±2.17 mg/dL in I-M-150847 treated group versus 130 ±3.02 mg/dL exendin-4

treated group, ***$p<0.001$, one-way ANOVA, Tukey's multi comparison test, n=6 **(Fig**

4B (iv)). A comparative glucose tolerance test on the 20th day of agonist treatment also

demonstrates the superiority of I-M-150847 over exendin-4 in executing glycemic

control. As **Fig 4C** reveals, glucose excursion is superior upon I-M-150847 treatment

relative to exendin-4 mediated therapy (mean AUC 20809 ± 312.1, in exendin-4 treated

group, versus mean AUC 13218 ± 164.5, in I-M-150847 treated group; $p<0.001$, one-

way ANOVA, Tukey's multi comparison test, n=6). To explore whether enhanced

glucose disposal is associated with increased insulin sensitivity, we carried out the

insulin tolerance test. I-M-150847 treatment, in our study, displayed better insulin

sensitivity compared to exendin-4 treatment (AUC 6149 ± 125.7 upon exendin-4

treatment, versus AUC 5134 ± 39.54 upon I-M-150847 treatment; ***$p<0.001$, one-way

ANOVA, Tukey's multi comparison test, n=6) as demonstrated in **Fig 4D**. The data,

taken together, underlines the superiority of the dual agonist I-M-150847 over exendin-4

mediated therapy in regulating glucose homeostasis.

Based on the evidence of enhanced glycemic benefits, we explored whether I-M-

150847 provides appreciable weight loss in DIO mice. As **Fig 4E** shows, I-M-150847

causes a significant decrease in cumulative food intake compared to vehicle control

(*$p<0.05$, one-way ANOVA, Tukey's multiple comparison test, n=6 for each group) after

16 days of treatment. The decrease in cumulative food intake upon I-M-150847

treatment is significant compared to exendin-4 from the 7th day of treatment (*p<0.05,

two-way ANOVA, Bonferroni post-tests) that maximizes on the 16th day of the study

paradigm **(Fig 4E, ***p<0.001, two-way ANOVA, Bonferroni post-tests)**. Associated with

the decreased food intake, we observed appreciable weight loss in I-M-150847 treated

group that is significantly superior compared to the group treated with exendin-4 or

vehicle **(Fig 4F,** ***p<0.001, one-way ANOVA, Tukey's multiple comparison test, n=6 for

each group**)**. Further, I-M-150847 treatment causes a significant reduction in liver

weight compared to exendin-4 mediated therapy and vehicle control (1.712 ± 0.05

grams in I-M-150847 treated group versus 2.47± 0.08 grams in the control group

receiving normal saline as the vehicle, *** p<0.001, one-way ANOVA, Tukey's multiple

comparison test, n=6 for each group and 2.027± 62.37 grams in exendin-4 treated

group p<0.05, one-way ANOVA Tukey's multiple comparison test, n=6 for each group.

Fig 4G(i)). We observed a significant reduction of inguinal WAT both in I-M-150847 and

exendin-4 treated mice (152.7±27.94 grams and 140 ±14.96 grams in respective I-M-

150847 treated and exendin-4 treated groups compared to 337 ±0.45 gram in DIO mice

receiving normal saline as vehicle control, ***p<0.001, one-way ANOVA, Tukey's

multiple comparison test, n=6 for each group). Similarly, there is a significant reduction

in epididymal WAT (341.8±100.5 grams and 431.8 ±42.80 grams in I-M-150847 treated

and exendin-4 treated group compared to 930.3 ±44.40 grams in the control DIO mice

receiving normal saline as the vehicle,***p<0.001, one-way ANOVA, Tukey's multiple

comparison test, n=6 for each group), retroperitoneal WAT(68.0± 16.71 grams and

127.8 ± 15.72 grams upon respective I-M-150847 and exendin-4 treatment compared to

259.2±27.93 grams in control DIO mice receiving normal saline as vehicle ***p<0.001

and **p<0.01, one-way ANOVA, Tukey's multiple comparison test, n=6 for each group)

and mesenteric WAT (40.5± 3.905 grams and 54.67± 6.307 grams upon respective I-M-

150847 and exendin-4 treatment compared to 98.33± 9.2 grams ***p<0.001 and

**p<0.01, one-way ANOVA, Tukey's multiple comparison test, n=6 for each group),

thereby demonstrating prominent reduction of adiposity both in I-M-150847- treated and

exendin-4 treated DIO mice. The data, taken together, highlight the efficacy of GLP-1R

GIPR dual agonist I-M-150847 in achieving improved glucose homeostasis and

enhanced insulin sensitivity associated with the significant reduction of body weight,

food intake, and adiposity in the DIO rodent model.

Discussion:

Multi-receptor agonism of incretins has recently emerged as a preferred therapy to

achieve superior glycemic control and reduction of obesity (Gimeno et al., 2020; Samms

et al., 2021). The concept of unimolecular GLP-1R and GIPR dual agonism first

described by Finan et.al demonstrated the efficacy of hybrid peptides in reducing

diabesity more effectively than the conventional GLP-1R agonists (Finan et al., 2013).

Consistent with these findings, the dual GLP-1R, and GIPR agonist Tirzepatide

(LY3298176) developed by Eli Lilly revealed better therapeutic ability than the most

potent GLP-1R agonist semaglutide in reducing HbA1c, body weight, and serum

triglycerides in SURPASS phase 3 clinical trials (Frias et al., 2021; Khoo and Tan, 2021;

Min and Bain, 2021; Rosenstock et al., 2021; Slomski, 2021) thereby promising superior

long-term glycemic control and significant reduction of obesity (Coskun et al., 2018),

(Baggio and Drucker, 2021; Bailey, 2021).

The majority of the unimolecular incretin dual agonists reported to date were designed

by intermixing GLP-1 and GIP amino acids in a single peptide using either glucagon

(Finan et al., 2013), oxyntomodulin ((Bhat et al., 2013; Muppidi et al., 2016),(Yang et al.,

2020), (Zhao et al., 2019), or GIP as the base peptide (Willard et al., 2020). The incretin

receptor dual agonist I-M-150847 was designed based on the structure-based activation

model of the Class B GPCRs (Hoare, 2005) taking clues from the conformational

rearrangement upon ligand binding that initiates the second messenger signaling

(Hoare, 2005; Parthier et al., 2007; Runge et al., 2008). The data on the crystal

structure of GLP-1, exendin-4, and GIP bound to respective GLP-1R and GIPR

extracellular domain (ECD) (Parthier et al., 2007; Runge et al., 2008; Underwood et al.,

2010) provided us the critical insights for the designing of the middle and the C terminal

region of the hybrid agonist. We focused on the Lys^{30} of GIP that aligns with the

corresponding amino acid residue of oxyntomodulin and experimentally determined the

activity by cAMP reporter assays. We replaced the tryptophan cage (Trp-cage) at the C

terminus of exendin-4 (Lee et al., 2018; Neidigh et al., 2001; Runge et al., 2008; Wolff et

al., 2021) despite its presence in Tirzepatide (Kegg Drug D11360) and stapled GLP-1R

dual agonist O14(Yang et al., 2020) as previous studies reported the redundancy of the

motif for the preservation of high binding affinity for GLP-1R (Lee et al., 2018).

We initiated structure-activity relationship (SAR) analysis with the synthesis of I-M-

150845 (**Table 1, Figure 1 -Figure supplement 1**) upon the addition of GLP-1 and

GIP-specific amino acids to the first 9 amino acid residues of the peptide I-M-150844

that corresponds to the N-terminal sequence of exendin-4. I-M-150845 showed dual

GLP-1R and GIPR activity in cAMP reporter assays (**Table 1**). From the structural

perspective, the amino acid residues Phe^{22}, Val^{23}, Leu^{26}, and Leu^{27} of the amphipathic

alpha-helix of GIP that forms-minute hydrophobic interactions with GIPR ECD are

present in the hybrid peptide I-M-150845. In addition, we retained Asp^{15}, Gln^{19}, Gln^{20},

Leu^{26}, and leu^{27}, the amino acids that form an extensive hydrogen-bonding network with

several residues of GIPR ECD in I-M-150845 to provide GIPR-specific activity. In

contrast, the amino acids of the exendin-4/GLP-1 amphipathic alpha-helix that support

the hydrophilic interactions with GLP-1R ECD are absent in I-M-150845. Of the 7 amino

acids of exendin-4, Ala^{18}, Val^{19}, Phe ^{22}, Ile ^{23}, Trp^{25}, Leu^{26}, and Pro^{31} that form

hydrophobic interaction with GLP-1R ECD, only Phe^{22} and Leu^{26} are retained in hybrid

dual agonist I-M-150845. We hypothesized that Val^{27} at the I-M-150845 has a

compensatory influence as it establishes hydrophobic contact as well as hydrogen

bonding with GLP-1R ECD in the receptor-ligand crystal complex (Underwood et al.,

2010). To experimentally ascertain the importance of Val^{27} for GLP-1R specific activity,

we synthesized a new incretin agonist I-M-150846 where the C- terminal amino acids

Val^{27}-Gly^{31} of I-M-150845 are substituted with 11- amino acid C terminus (Met^{27}-Ala^{37})

of oxyntomodulin (**Figure 1 Figure supplement 2**). The substitution removes Val^{27} but

introduces Lys^{30} which in GIP establishes an intricate hydrogen-bonding network with

several residues in the C terminal part of GIPR ECD (Parthier et al., 2007). The incretin

agonist I-M-150846 revealed a complete loss of function at GLP-1R while retaining

GIPR activity **(Table 1).** We then replaced GIP-specific Tyr^{10}-Asn^{24} of I-M-150846 with

amino acid residues Leu^{10}- Glu^{24} of exendin-4 (Figure **1- Figure supplement 3**) that

forms superior helical architecture with GLP-1R ECD (Runge et al., 2008). The resultant

incretin agonist I-M-150847 showed robust GLP-1R and GIPR dual agonism signifying

that in the absence of Trp cage, the presence of the amino acid residues Leu[10]- Glu[24] of

exendin-4 in the hybrid agonist peptide is critical to establish a stable interaction with

the GLP-1R ECD thereby restoring the functional activity towards GLP-1R as measured

by cAMP reporter assay. The data also implies that Phe[22], Leu[26], and Lys[30] are the

crucial amino acids that confer binding to the GIPR ECD required for the GIPR-specific

activity of I-M-150847.

Variable propensities of internalization and recycling of GLP-1R determine the potential

of GLP1-RAs in delivering insulinotropic response (Jones et al., 2018). We first reported

sustained cAMP signaling following prolonged internalization of GLP-1R upon treatment

of BRIN-BD11 pancreatic beta cells with GLP-1Tmr, exendin-4, and liraglutide (Kuna et

al., 2013),(Girada et al., 2017),(Bele et al., 2020). Endocytic sorting of GLP-1R to

lysosome upon treatment with GLP-1Tmr and exendin-4 was reported in cultured

pancreatic beta cells(Kuna et al., 2013),(Jones et al., 2018). In contrast, exendin F1, an

exendin -4 analog, elicits a fast-recycling response of GLP-1R and provides better

glycemic benefits compared to other GLP-1R monoagonists (Jones et al., 2018). In our

present study, we observed prompt redistribution of functional GLP-1R at the plasma

membrane following I-M-150847 treatment in contrast to exendin-4 -mediated prolonged

receptor internalization. Our data shows that the I-M-150847- mediated cAMP

generation is discrete as we observe an abrupt attenuation of receptor-mediated cAMP

accumulation 30 minutes after the treatment with I-M-150847 which is in contrast to the

exendin-4 mediated activation where continuous cAMP accumulation is observed till an

hour after the agonist exposure. In this context, the gain of cAMP generating response following agonist rechallenge that is specific for I-M-150847- mediated activation is indicative of reduced internalization and or faster GLP-1R recycling that may account for the higher distribution of the sensitized receptor at the plasma membrane following I-M-150847 treatment.

Functional consequences of the preservation of incretin receptors in a resensitized state are presently unclear. In our study, we observe superior insulinotropic activity upon acute treatment with I-M150847 compared to exendin-4. However, GSIS is blunted in the case of chronic exposure to exendin-4 or I-M-150847 signifying homologous desensitization of the incretin receptor both in the case of prolonged internalization in exendin-4 - treated cells as well as upon faster recycling or reduced internalization that we observe upon I-M-150847 treatment. In contrast, in the case of mouse adipocytes that express GIPR (McIntosh et al., 2012), we observe retention of insulin sensitivity as well as cAMP accumulation response after chronic exposure to I-M-150847.GIPR internalization and desensitization upon treatment with the cognate agonist have been earlier documented (Ismail et al., 2015). In this context, the functional response of I-M-150847 in mouse adipocytes reveals retention of GIPR in a sensitized state upon chronic exposure to the agonist supporting the downstream signaling cascade that regulates the cellular metabolism. The data is also indicative of the cell type variability of incretin receptor desensitization as has been reported in the case of the M3 muscarinic receptor(Torrecilla et al., 2007), (Kelly et al., 2008).

The efficiency of I-M-150847 in eliciting insulin sensitivity in adipocytes translates into better in vivo glycemic benefits as we observe significantly improved reduction of fasting

blood glucose and better glucose tolerance in DIO mice upon I-M-150847 treatment.

We also observed a significant reduction of food intake and weight loss in I-M-150847

treated mice relative to the group administered with exendin-4 or vehicle. However,

whether this greater anorexic response upon I-M-150847 treatment is a consequence of

the preservation of GLP-1R and or GIPR at a sensitized state at the feeding centers in

the hypothalamus deserves attention in future research.

A key question regarding incretin receptor dual agonism is the contribution of GIPR in

the attainment of superior glycemic control. Several reports highlight the benefits of

LAGIPRA in enhancing insulin sensitivity through activation of the Akt signaling pathway

that is associated with the storage and oxidation of glucose in adipose tissues, (Samms

et al., 2021; Samms et al., 2020). However, similar attributes of improved glucose

tolerance and insulin sensitivity have been reported in contrasting mouse models

representing GIPR overexpression (Kim et al., 2012) and adipose tissue-specific GIPR

knockout (Joo et al., 2017). Our data with 3T3-L1 adipocytes demonstrate a significant

increase in insulin-stimulated glucose uptake upon chronic I-M-150847 treatment

despite the increase in the cAMP generation which is indicative of increased lipolysis.

The data is in agreement with the previous report of increased lipoprotein lipase activity

upon GIP treatment in cultured preadipocytes (Eckel et al., 1979) as well as the

enhancement of insulin sensitivity in epididymal WAT upon LAGIPRA treatment in DIO

mice due to increased Akt phosphorylation (Varol et al., 2014). Thus, while it is likely

that GIPR knock-out mice showed enhanced insulin sensitivity due to reduced GIPR -

mediated lipolysis and consequent less free fatty acid (FFA) generation, GIPR

overexpression or treatment with LAGIPRA or GLP-1R GIPR dual agonists may

augment insulin sensitivity through activation of metabolic pathways that are associated

with glucose and FFA metabolism in the adipose tissue. We note that a detailed

mechanistic study is required to fully test the hypothesis.

Collectively, the preclinical data presented in this study demonstrate I-M-150847 as a

novel GLP-1R and GIPR dual agonist that promotes expeditious resensitization of GLP-

1R and preserves the functional attributes of GIPR upon chronic exposure in adipocytes

thereby augmenting better insulinotropic response and enhancing insulin sensitivity

consequentially improving the systemic glycemic control. Further, I-M-150847 reduces

food intake, ensures significant body weight loss, and decreases adiposity highlighting

the therapeutic potential of the novel hybrid agonist in ameliorating T2DM and obesity.

1   **Materials and Methods:**

2   **Key resources table:**

3   Following reagents are used in this study.

REAGENT	SOURCE	IDENTIFIER
CHEMICALS, PEPTIDES, AND RECOMBINANT PROTEINS		
RPMI 1640 HiGlutaXL	Himedia	AL-060G
DMEM HIGH GLUCOSE	Himedia	AL-219A
FBS Heat Inactivated (US ORIGIN)	Himedia	RM-10409
TRYPSIN EDTA (10X)	Himedia	TCL-034
Penicillin Streptomycin solution (100X)	Himedia	A001
Gentamycin	Himedia	A005
DMSO	SIGMA	M81802
LIPOFECTAMINE 3000	THERMO	L3000001
IBMX	SANTA CRUZ	SC-201188
GLUCOSE	SIGMA	G-8270
PHOSPHATE BUFFER SALINE	INVITROGEN	70011044
BETA MERCAPTOETHANOL	SIGMA	M6250
RNAiso Plus	Takara	9108
BOVINE SERUM ALBUMIN	SIGMA	A2153
FAT-FREE BSA (FRACTION V)	SIGMA	5479
STEADYLITE Plus	PERKIN ELMER	6066751
ATP	SIGMA	A7699
PROTEASE INHIBITOR COCKTAIL	CALBIOCHEM	539131
LIRAGLUTIDE (VICTOZA)	NOVO NORDISK	Acylated GLP-1R agonist
PHOSPHATASE INHIBITOR	CALBIOCHEM	524625
RNA Later solution	AMBION	AM7020
Triton-X 100	SIGMA	T8787
Dr. Trust Glucometer and strips	Dr. Trust USA	9001
IN-M-150845	Designed in-house, synthesized, and, purified by GL BIOCHEM, CHINA	Procured through Biotech Desk India
IN-M-150846		
IN-M-150847		
Exendin F1	Dr. Ben Jones, Imperial College, London	
Oxyntomodulin	Tocris	2094
GIP	Genscript	I
2-nitro-phenyl beta galactopyranoside	SIGMA	N1127
Paraformaldehyde	SIGMA	158127

Vectashield	Vector Labs	H-1200-10
Plasmid GLP-1R GFP	Mitra Lab	Kuna et.al
Plasmid CRE Luciferase	Richard Day, Indiana University	Kuna et.al 2013(Kuna et al., 2013)
Plasmid GIPR	Bernard Thorens	ADDGENE#14942(Gremlich et al., 1995)
Plasmid GCGR	Brian Finan	Novo Nordisk

CRITICAL COMMERCIAL ASSAYS		
BCA PROTEIN ASSAY KIT	SANTA CRUZ	sc-200629
INSULIN ELISA KIT (RAT/MOUSE)	MERCK	EZRMI-13K
cAMP ELISA kit	ENZO	ADI-900-067
TB Green® Premix Ex Taq™ II	Takara	RR820B
PRIMESCRIPT cDNA synthesis kit	Takara	RR014B
PLASMID ISOLATION KIT	INVITROGEN	K210005

OLIGONUCLEOTIDES	
RT-qPCR Primers	IDT

EXPERIMENTAL MODELS/ CELL LINES		
BRIN-BD11 CELL LINE	ECACC	10033003
HEK-293T CELL LINE	ATCC	CRL-3216
3T3L1 preadipocytes	ATCC	CL-173
C57BL/6 MICE	Jeeva Life Sciences, Hyderabad	JLS-000664
Animal Feed (HIGH FAT DIET/CHOW DIET)	NIN, Hyderabad	

Methods:

Peptide Synthesis:

Synthesis of the 11- amino acid residue peptide I-M-150844 was carried out in Biotage

Syro I automated peptide synthesizer using Wang resin and following Fmoc chemistry.

The amino acid that was first loaded to the resin is Phenylalanine and following

installation, the capping of residual functional groups on the resin was carried out by

treatment with 33% acetic anhydride and pyridine. Before loading of next amino acid,

the Fmoc group of the amino acid was deprotected through treatment by 40% piperidine

/ DMF and the incorporation of the next amino acid was carried out using coupling

reagent HBTU (0.5 mmol) and DIPEA/NMP (2 mmol). The peptide was cleaved from the

resin by mixing it with TFA/H_2O/TIPS (95:2.5:2.5, 4 ml) and TFA was removed by

evaporation on a rotatory evaporator. The crude mixture was triturated (diethyl ether)

and purified by preparative HPLC using Agilent column ZORBAX 300SB-C18 (5 µm,

9.4*250 mm).

The other peptides I-M-150845, I-M-150846, and I-M-150847 were designed in-house

and outsourced for chemical synthesis and purification to GL Biochem China. The C

terminal amidated version of I-M-150847 was synthesized by Aurigene Pharmaceutical

Services, India.

Circular dichroism spectroscopy:

Determination of the secondary structure of I-M-150847 was carried out by using

Circular Dichroism Instrument (JASCO, J1500, Easton MD) (Kelly et al., 2005;

Waterhous and Johnson, 1994). The spectra from 200 to 260 nm were taken with a

data interval of 0.5nm. A bandwidth of 1.00nm with a 100 nm/min scanning rate was

used to determine the property.10μM of I-M-150847 was used for the study and 50mM

Phosphate buffer serves as the blank background. The data were represented as

Ellipticity (mdeg) vs wavelength (nm) at the Y and X-axis respectively.

Cell Culture:

Pancreatic beta cells, BRIN-BD11 (cat no.10033003) were procured from the European

Collection of Authenticated Cell Cultures. The cells were cultured in RPMI media 1640

HiGlutaXL with 10% heat-inactivated fetal bovine serum (US origin) supplemented with

1mM sodium pyruvate, 50μM β-mercaptoethanol,10mg/mL gentamycin, 100U/mL

penicillin, and 100mg/mL streptomycin at an incubation temperature of 37º C with 5%

CO_2 following the method of Kuna et. al (Kuna et al., 2013). Human Embryonic Kidney

cell lineage (HEK 293T) cells were procured from ATCC (CRL3216). The cells were

cultured in DMEM-Hi Glucose media supplemented with 10% heat-inactivated FBS,

10mg/mL gentamycin, 100U/mL penicillin, and 100mg/mL streptomycin at an incubation

temperature of 37º C with 5% CO_2, cultured following the protocol recommended by

ATCC. 3T3 L1 mouse preadipocytes (ATCC (CL-173)) were sub-cultured using DMEM

supplemented with 10% newborn calf serum and 100U/mL penicillin, and 100mg/mL

streptomycin at an incubation temperature of 37º C with 5% CO_2 following ATCC

protocol. The preadipocytes were differentiated into adipocytes following the protocol of

Mitra et.al (Mitra et al., 2004)

GLP-1R, GIPR, GCGR activation:

Each peptide was tested for its ability to activate GLP-1R, GIPR, or GCGR in a cell-based cAMP-responsive element luciferase reporter assay following the method of Kuna et.al (Kuna et al., 2013). Briefly, HEK 293T cells were cultured in a 60mm tissue culture treated dish. Upon 70% confluency, the cells were transiently transfected with cAMP-responsive element luciferase reporter plasmid, beta-galactosidase plasmid, and GIP-R or GLP-1 R, or GCGR plasmid in a 3:1:3 ratio. The transient transfection was carried out using lipofectamine 3000 following the manufacturer's protocol. After 4 hours of transfection, the cells were seeded in a tissue culture grade 96 well plate at a density of 50000 cells / well. After overnight incubation, the cells were treated with different concentrations of agonists for the different cell-based assays. After 4 hours of treatment, the media was aspirated and cells were lysed and the luminescence was measured using Steady lite Plus reagent (Perkin Elmer Life and Analytical Science, Waltham, MA) after 15 minutes of shaking at 300 rpm at room temperature in dark. 2-nitro-phenyl beta galactopyranoside (Sigma) was used to carry out a beta-galactosidase assay for correcting the inter-well transfection variability.

cAMP accumulation assay:

The accumulation of cAMP was measured in cultured BRIN-BD11 at different time points upon agonist treatment. 1μM I-M-150847 and exendin-4 were tested for the time-course of cAMP generation using a direct cAMP ELISA kit (Enzo Life Sciences, Cat. No. ADI-900-163). The assay was carried out following the protocol of Girada et. al (Girada

et al., 2017) with modifications. Briefly, BRIN-BD11 cells were seeded in 24 well-treated

plates at a density of 0.1 million cells/well. After overnight incubation, the cells were

incubated again for 60 minutes in the KRB-BSA medium (with 1.1mM glucose) after

which the cells were washed and treated with 1µM I-M-150847 or Exendin-4 for 3

minutes, and then the treatment medium was washed off and replaced with KRB-BSA

medium with 200µM IBMX and 5.5mM glucose and incubated for 1, 10,30, and 60

minutes after which the cells were lysed and processed for the cAMP ELISA assay.

Protein quantification was carried out using BCA following kit protocol.

To assess chronic cAMP generation in 3T3-L1 adipocytes, the cells were treated with

1µM I-M-150847 or exendin-4 overnight in serum-free media following which they are

washed and incubated with KRB buffer with 0.1%BSA containing 1.1mM glucose. The

cells were then exposed to a treatment medium comprising fresh KRB-BSA buffer with

5.5mM glucose, 200µM IBMX, and 1µM I-M-150847 or exendin-4 for 3 minutes, and

following wash incubated in KRB buffer with 5.5mM glucose and 200µM IBMX for 30

minutes. The cells were then lysed and processed for cAMP ELISA assay. Protein

quantification was carried out using BCA following the kit protocol.

Agonist rechallenge assay:

Agonist rechallenge was carried out following the protocol by Jones et.al (Jones et al.,

2018) with modifications. Briefly, BRIN-BD11 cells were seeded in 24 well tissue culture

treated plates at a density of 0.1 million cells/ well. After overnight incubation in the

complete medium, the cells were incubated for 60 minutes in the KRB-BSA medium

with 1.1mM glucose after which the cells were washed and treated with 1µM I-M-

150847 or Exendin-4 in a KRB-BSA medium containing 200µM IBMX and 5.5mM

glucose for 3 minutes. The treatment medium was then washed off and replaced with a

fresh KRB-BSA medium containing 5.5mM glucose and 200µM IBMX and incubated for

45 minutes. The cells were re-challenged with agonists for 10-minutes followed by lysis

and subsequently processed for the cAMP ELISA assay. Protein quantification was

carried out using BCA following kit protocol.

GLP-1R internalization and post-endocytic sorting:

Assessment of GLP-1R internalization was carried out in BRIN-BD11 pancreatic beta

cells and monitored using confocal microscopy. The cells were transfected with GLP-

1R-GFP using lipofectamine 3000 and seeded in six-well plates on 22*22mm sterile

cover glass coated with poly-L-lysine. 48 hours after transfection, the cells were washed

with sterile PBS and incubated with I-M-150847 (1µM) or Exendin-4 (1µM) in 200µL of

Krebs-HEPES buffer for 60 min at 4°C in a humidified chamber in the dark. Cells were

then washed in PBS and incubated at 37°C for 1 min, 10 minutes, 30 minutes, and 60

minutes after which they are fixed in 4% paraformaldehyde and imaged using a Zeiss

LSM 880 confocal laser scanning microscope (Carl Zeiss, Oberkochen, Germany)

equipped with HXP 120V light sources. Pinhole diameter was maintained at 1 airy unit.

Fields were selected using a 20X objective and image acquisition was carried out using

a 63 X oil immersion objective lens.

The post-endocytic sorting of GLP-1R after activation was carried out following the

method of Kuna et.al with modifications (Kuna et al., 2013). Briefly, BRIN-BD11 cells

were transfected with GLP-1R-GFP, and 60 hours after transfection were treated with

1µM I-M-150847 or Exendin-4 and incubated at 4°C, in a dark humidified chamber for

one hour. The cells were then washed with ice-cold PBS and treated with Lysotracker

red (100nM) for 45 minutes in RPMI-1640 medium, washed with PBS, and fixed in 4%

paraformaldehyde. The coverslips were mounted on glass slides by using Vectashield

mounting media and the imaging was carried out in Zeiss LSM 880 confocal laser

scanning microscope (Carl Zeiss, Oberkochen, Germany) equipped with HXP 120V

light sources with a pinhole diameter of 1 airy unit. The emission was measured at 500-

550nm bandpass for GFP and 575-615 nm for lysotracker red. The images were

acquired by using a 63X oil immersion objective lens with 2X optical zoom and analyzed

using Zenlite 2011 program.

Glucose- stimulated Insulin secretion:

Insulin secretion was assessed using a Millipore rat/ mouse insulin ELISA kit (Merck,

EZRMI-13K) following the method of Asalla et.al.(Asalla et al., 2016) Briefly, BRIN-

BD11 cells were seeded in 24 well tissue culture plates at a density of 0.1 million cells /

well. In the next day, the cells were treated with Kreb's Ringer Bicarbonate buffer (KRB)

[115 mM NaCl, 4.7 mM KCl, 1.28 mM $CaCl_2$, 1.2 mM KH_2PO_4, 1.2 mM $MgSO_4$, 10 mM

$NaHCO_3$, 0.1% (wt./vol) BSA, pH 7.4] for 1h without glucose. Glucose-stimulated insulin

secretion (GSIS) was assessed in the presence of indicated concentrations of glucose

and incretin receptor agonists for 30 min and the supernatant was collected,

centrifuged, and processed for the ELISA as per the manufacturer's protocol. The cells

were lysed with 0.1% Triton-X 100 and processed for protein quantification to normalize

the insulin secretion. The data were expressed as fold-over basal insulin secretion (pg

insulin/µg protein). The insulin secretion at 0.1mM glucose is considered basal

secretion.

For assessment of GSIS during prolonged agonist treatment, BRIN-BD 11 cells were

treated with 1µM agonists in a 2% FBS-RPMI medium for 6 hours. The cells were then

washed with KRB buffer and incubated in KRB-0.1%BSA for 1 hour. The incubation

medium was replaced with a fresh KRB-BSA medium in the presence of 16.7mM

glucose and 1µM agonists for 30 minutes following which the supernatant was collected

for insulin ELISA and the cells were lysed to determine the protein to normalize the

insulin secretion. 10 µL of the sample supernatant was used for the insulin ELISA

assay. The quantity of insulin was measured as the picogram (pg) of insulin per

milligram (mg) of protein and expressed as a fold increase over the basal insulin

secretion. The insulin secretion at 0.1mM glucose is considered basal secretion.

Glucose uptake assay:

Glucose uptake was carried out on the 9[th] day differentiated 3T3-L1 adipocytes with a

fluorescent D- glucose analog 2-[N-(7-nitrobenz-2-oxa-1,3diazol-4-yl) amino]-2-deoxy-D

glucose (NBDG) following the method of Zou et.al (Zou et al., 2005). Briefly, 3T3 L1

adipocytes were treated with 1µM of I-M-150847 or Exendin-4 overnight in serum-free

DMEM. The cells were then washed and incubated with KRB buffer containing 0.1% BSA

for 60 minutes. Post incubation, the cells were treated with 100nM of insulin for 30 minutes after which 150 μM of 2-NBDG was added for 20 minutes. The reaction was stopped using ice-cold PBS, and the cells were lysed with the lysis buffer containing 0.1% Triton X 100 in dark on an orbital shaker. The lysates were transferred to the Corning 96 well polystyrene Black microplate and read in triplicate at 485/535 nm excitation/emission in a multi-mode plate reader.

Total RNA isolation, cDNA preparation, and qRT PCR

The RNA extraction was carried out by phenol-chloroform extraction protocol from cultured BRIN-BD11 pancreatic beta cells.1 μg of total RNA was used for first-strand cDNA synthesis. Takara Prime script cDNA synthesis kit was used to carry out the two-step cDNA synthesis following the manufacturer's protocol. The mRNA for GLP-1R and GIPR was amplified and quantified by quantitative real-time PCR using Quant Studio 5 (A&B Biosystems). 2XSYBR master mix (Takara Life Sciences) reagent was used for the assessment of the relative abundance of mRNAs measured using the $2^{-\Delta\Delta CT}$ method (Livak and Schmittgen, 2001). Normalization was accomplished using GAPDH as an invariant reference. Primers used for amplification are as follows: GLP1R *f*: TATTGGCTCATCATACGCTTG; GLP1R*r:* GTCTGCATTTGATGTCGGTCT; GIPR *f*: CGAAGTCAAAGCCATTTGGT; GIPR *r*: CCAGCCTTAGTCGGTAGTCG; Beta actin *f*: CCCATACCCACCATCACACC; Beta actin *r*: GAGAGGGAAATCGTGCGTGAC.

Animal Experiments:

Animal experiments were performed following the guidelines of Control and Supervision

of Experiments on Animals (CPCSEA) upon permission from the Institutional Ethical

Committee Hyderabad, India (UH/IAEC/KS/2021-1/36).C57BL/6 male mice of 5 weeks

of age were procured from Jeeva Life Sciences, Hyderabad (Cat No: JLS-000664). Only

male mice were used for the study since female mice are mostly protected from diet-

induced obesity (Pettersson et al., 2012).

Pharmacological and Metabolism studies:

The mice were group-housed or single-housed on a 12:12 h light-dark cycle at 22˚-24˚C

and were fed ad-libitum on a diabetogenic diet (Bele et al., 2020) which is a high-

sucrose diet with 58% kcal from fat for 75 days. The mice were then randomized based

on body weight, food intake, and blood glucose before the initiation of the study and

were divided into 3 groups (n=6 / group): control receiving normal saline as a vehicle,

comparator group receiving 100 nmol/kg body weight exendin-4 twice daily and test

group receiving 100nmol/kg body weight I-M-150847 twice daily through subcutaneous

administration for 4 weeks. Food intake and body weight were measured every day after

the first injection of agonists. Fasted blood glucose was measured at the study initiation,

on the 10th day of treatment, and at the termination of the experiment.

Glucose tolerance test:

On the 20th day of treatment, DIO mice were subjected to fasting for 5 hours and

following which the intraperitoneal injection of glucose (2g/kg body weight) (D-glucose

[Sigma] 40% w/v in 0.9% normal saline) was administered. Glucose level was

monitored by a glucometer upon collecting blood from the tail vein before (0) and after

(15, 30, 60, 90, and 120 minutes) after glucose administration.

Insulin tolerance test:

On the 24th day of treatment, the animals fasted for 4 hours after which the animals

received an intraperitoneal injection of insulin (0.75U/kg body weight, Humulin R, Eli

Lilly, and Company). The blood from the tail vein was collected before (0) and after (15,

30, 60, and 90 minutes) insulin administration, and blood glucose was monitored using

a glucometer.

Tissue weight measurement:

For the measurement of tissue weight, animals were sacrificed on the 28th day of

treatment and mesenteric, retroperitoneal, inguinal, and epididymal WAT were collected.

Blood and Liver were also collected during the sacrifice and kept at -80° C for further

processing.

Statistical analysis:

Statistical analyses were performed with GraphPad Prism 6.0 and presented as means

± SEM. The analysis of the experiments was performed with one-way ANOVA (Tukey's

multiple comparison test) and Student's t-test or two-way ANOVA Bonferroni posttests.

P values lower than 0.05 were considered significant.

Ethics:

The animal studies were carried out following the guidelines of Control and Supervision

of Experiments on Animals (CPCSEA) upon permission from the Institutional Ethical

Committee Hyderabad, India (UH/IAEC/KS/2021-1/36).

Author Contributions:

RB and SB jointly carried out the experiments with the help of JE and equally

contributed to the study. SK, TD, and AI provided the necessary support for conducting

experiments. NC and VR synthesized some of the peptides and carried out physical

characterizations. RB, SB, and PM carried out the in-silico analysis of the structure of

novel incretin peptides. PM conceptualized the project and supervised the experiments,

designed the peptides, analyzed and interpreted the data, wrote the manuscript upon

discussion with RB and SB, and take the responsibility for the integrity and accuracy of

the data. All authors have full access to the data related to the manuscript and

approved the final version of the manuscript for publication.

Acknowledgments:

The work has been supported by Department of Science and Technology Grants;

EMR/2016/007057 provided to PM, VR, and AI. Fellowship of RB and SB is supported

by the Council of Scientific and Industrial Research (CSIR No: 09/0985 (0032)/2019-

EMRI) and ICMR senior research fellowship (No 2017-3891-CMB/BMS). SB and RB are

registered at the Manipal Academy of Higher Education (MAHE) for their doctoral

degree. Exendin F1 is a generous gift from Dr. Ben Jones, Imperial College London.

Declaration of interest:

The authors declare no conflicts of interest.

References:

Amori, R.E., Lau, J., and Pittas, A.G. (2007). Efficacy and safety of incretin therapy in type 2 diabetes:
systematic review and meta-analysis. JAMA *298*, 194-206.
Andersen, A., Lund, A., Knop, F.K., and Vilsboll, T. (2018). Glucagon-like peptide 1 in health and disease.
Nat Rev Endocrinol *14*, 390-403.
Asalla, S., Girada, S.B., Kuna, R.S., Chowdhury, D., Kandagatla, B., Oruganti, S., Bhadra, U., Bhadra, M.P.,
Kalivendi, S.V., Rao, S.P., et al. (2016). Restoring Mitochondrial Function: A Small Molecule-mediated
Approach to Enhance Glucose Stimulated Insulin Secretion in Cholesterol Accumulated Pancreatic beta
cells. Sci Rep *6*, 27513.
Astrup, A., Rossner, S., Van Gaal, L., Rissanen, A., Niskanen, L., Al Hakim, M., Madsen, J., Rasmussen,
M.F., Lean, M.E., and Group, N.N.S. (2009). Effects of liraglutide in the treatment of obesity: a
randomised, double-blind, placebo-controlled study. Lancet *374*, 1606-1616.
Baggio, L.L., and Drucker, D.J. (2021). Glucagon-like peptide-1 receptor co-agonists for treating
metabolic disease. Mol Metab *46*, 101090.
Bailey, C.J. (2021). Tirzepatide: a new low for bodyweight and blood glucose. Lancet Diabetes Endocrinol
9, 646-648.
Bastin, M., and Andreelli, F. (2019). Dual GIP-GLP1-Receptor Agonists In The Treatment Of Type 2
Diabetes: A Short Review On Emerging Data And Therapeutic Potential. Diabetes Metab Syndr Obes *12*,
1973-1985.
Bele, S., Girada, S.B., Ray, A., Gupta, A., Oruganti, S., Prakash Babu, P., Rayalla, R.S., Kalivendi, S.V.,
Ibrahim, A., Puri, V., et al. (2020). MS-275, a class 1 histone deacetylase inhibitor augments glucagon-like
peptide-1 receptor agonism to improve glycemic control and reduce obesity in diet-induced obese mice.
Elife *9*.
Bhat, V.K., Kerr, B.D., Flatt, P.R., and Gault, V.A. (2013). A novel GIP-oxyntomodulin hybrid peptide acting
through GIP, glucagon and GLP-1 receptors exhibits weight reducing and anti-diabetic properties.
Biochem Pharmacol *85*, 1655-1662.
Coskun, T., Sloop, K.W., Loghin, C., Alsina-Fernandez, J., Urva, S., Bokvist, K.B., Cui, X., Briere, D.A.,
Cabrera, O., Roell, W.C., et al. (2018). LY3298176, a novel dual GIP and GLP-1 receptor agonist for the
treatment of type 2 diabetes mellitus: From discovery to clinical proof of concept. Mol Metab *18*, 3-14.
DiMarchi, R.D. (2018). "Let's Stay Together"; GIP and GLP-1 dual agonism in the treatment of metabolic
disease. Mol Metab *18*, 1-2.
Drucker, D.J. (2006). The biology of incretin hormones. Cell Metab *3*, 153-165.
Eckel, R.H., Fujimoto, W.Y., and Brunzell, J.D. (1979). Gastric inhibitory polypeptide enhanced lipoprotein
lipase activity in cultured preadipocytes. Diabetes *28*, 1141-1142.
Finan, B., Ma, T., Ottaway, N., Muller, T.D., Habegger, K.M., Heppner, K.M., Kirchner, H., Holland, J.,
Hembree, J., Raver, C., et al. (2013). Unimolecular dual incretins maximize metabolic benefits in rodents,
monkeys, and humans. Sci Transl Med *5*, 209ra151.
Fonseca, V.A., Capehorn, M.S., Garg, S.K., Jodar Gimeno, E., Hansen, O.H., Holst, A.G., Nayak, G., and
Seufert, J. (2019). Reductions in insulin resistance are mediated primarily via weight loss in subjects with
type 2 diabetes on semaglutide. J Clin Endocrinol Metab.

Frias, J.P., Davies, M.J., Rosenstock, J., Perez Manghi, F.C., Fernandez Lando, L., Bergman, B.K., Liu, B.,
Cui, X., Brown, K., and Investigators, S.-. (2021). Tirzepatide versus Semaglutide Once Weekly in Patients
with Type 2 Diabetes. N Engl J Med *385*, 503-515.
Frias, J.P., Nauck, M.A., Van, J., Kutner, M.E., Cui, X., Benson, C., Urva, S., Gimeno, R.E., Milicevic, Z.,
Robins, D., et al. (2018). Efficacy and safety of LY3298176, a novel dual GIP and GLP-1 receptor agonist,
in patients with type 2 diabetes: a randomised, placebo-controlled and active comparator-controlled
phase 2 trial. Lancet *392*, 2180-2193.
Gimeno, R.E., Briere, D.A., and Seeley, R.J. (2020). Leveraging the Gut to Treat Metabolic Disease. Cell
Metab *31*, 679-698.
Girada, S.B., Kuna, R.S., Bele, S., Zhu, Z., Chakravarthi, N.R., DiMarchi, R.D., and Mitra, P. (2017). Galphas
regulates Glucagon-Like Peptide 1 Receptor-mediated cyclic AMP generation at Rab5 endosomal
compartment. Mol Metab *6*, 1173-1185.
Gremlich, S., Porret, A., Hani, E.H., Cherif, D., Vionnet, N., Froguel, P., and Thorens, B. (1995). Cloning,
functional expression, and chromosomal localization of the human pancreatic islet glucose-dependent
insulinotropic polypeptide receptor. Diabetes *44*, 1202-1208.
Hoare, S.R. (2005). Mechanisms of peptide and nonpeptide ligand binding to Class B G-protein-coupled
receptors. Drug Discov Today *10*, 417-427.
Ismail, S., Dubois-Vedrenne, I., Laval, M., Tikhonova, I.G., D'Angelo, R., Sanchez, C., Clerc, P., Gherardi,
M.J., Gigoux, V., Magnan, R., et al. (2015). Internalization and desensitization of the human glucose-
dependent-insulinotropic receptor is affected by N-terminal acetylation of the agonist. Mol Cell
Endocrinol *414*, 202-215.
Jastreboff, A.M., Aronne, L.J., Ahmad, N.N., Wharton, S., Connery, L., Alves, B., Kiyosue, A., Zhang, S., Liu,
B., Bunck, M.C., et al. (2022). Tirzepatide Once Weekly for the Treatment of Obesity. N Engl J Med *387*,
205-216.
Jones, B., Buenaventura, T., Kanda, N., Chabosseau, P., Owen, B.M., Scott, R., Goldin, R.,
Angkathunyakul, N., Correa, I.R., Jr., Bosco, D., et al. (2018). Targeting GLP-1 receptor trafficking to
improve agonist efficacy. Nat Commun *9*, 1602.
Joo, E., Harada, N., Yamane, S., Fukushima, T., Taura, D., Iwasaki, K., Sankoda, A., Shibue, K., Harada, T.,
Suzuki, K., et al. (2017). Inhibition of Gastric Inhibitory Polypeptide Receptor Signaling in Adipose Tissue
Reduces Insulin Resistance and Hepatic Steatosis in High-Fat Diet-Fed Mice. Diabetes *66*, 868-879.
Kelly, E., Bailey, C.P., and Henderson, G. (2008). Agonist-selective mechanisms of GPCR desensitization.
Br J Pharmacol *153 Suppl 1*, S379-388.
Kelly, S.M., Jess, T.J., and Price, N.C. (2005). How to study proteins by circular dichroism. Biochim
Biophys Acta *1751*, 119-139.
Kerr, B.D., Flatt, P.R., and Gault, V.A. (2010). Effects of gamma-glutamyl linker on DPP-IV resistance,
duration of action and biological efficacy of acylated glucagon-like peptide-1. Biochem Pharmacol *80*,
396-401.
Khoo, B., and Tan, T.M. (2021). Surpassing insulin glargine in type 2 diabetes with tirzepatide. Lancet.
Kim, S.J., Nian, C., Karunakaran, S., Clee, S.M., Isales, C.M., and McIntosh, C.H. (2012). GIP-
overexpressing mice demonstrate reduced diet-induced obesity and steatosis, and improved glucose
homeostasis. PLoS One *7*, e40156.
Kuna, R.S., Girada, S.B., Asalla, S., Vallentyne, J., Maddika, S., Patterson, J.T., Smiley, D.L., DiMarchi, R.D.,
and Mitra, P. (2013). Glucagon-like peptide-1 receptor-mediated endosomal cAMP generation promotes
glucose-stimulated insulin secretion in pancreatic beta-cells. Am J Physiol Endocrinol Metab *305*, E161-
170.

Lee, J.G., Ryu, J.H., Kim, S.M., Park, M.Y., Kim, S.H., Shin, Y.G., Sohn, J.W., Kim, H.H., Park, Z.Y., Seong, J.Y., et al. (2018). Replacement of the C-terminal Trp-cage of exendin-4 with a fatty acid improves therapeutic utility. Biochem Pharmacol *151*, 59-68.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402-408.

McIntosh, C.H., Widenmaier, S., and Kim, S.J. (2012). Glucose-dependent insulinotropic polypeptide signaling in pancreatic beta-cells and adipocytes. J Diabetes Investig *3*, 96-106.

Min, T., and Bain, S.C. (2021). The Role of Tirzepatide, Dual GIP and GLP-1 Receptor Agonist, in the Management of Type 2 Diabetes: The SURPASS Clinical Trials. Diabetes Ther *12*, 143-157.

Mitra, P., Zheng, X., and Czech, M.P. (2004). RNAi-based analysis of CAP, Cbl, and CrkII function in the regulation of GLUT4 by insulin. J Biol Chem *279*, 37431-37435.

Muppidi, A., Zou, H., Yang, P.Y., Chao, E., Sherwood, L., Nunez, V., Woods, A.K., Schultz, P.G., Lin, Q., and Shen, W. (2016). Design of Potent and Proteolytically Stable Oxyntomodulin Analogs. ACS Chem Biol *11*, 324-328.

Neidigh, J.W., Fesinmeyer, R.M., Prickett, K.S., and Andersen, N.H. (2001). Exendin-4 and glucagon-like-peptide-1: NMR structural comparisons in the solution and micelle-associated states. Biochemistry *40*, 13188-13200.

Parthier, C., Kleinschmidt, M., Neumann, P., Rudolph, R., Manhart, S., Schlenzig, D., Fanghanel, J., Rahfeld, J.U., Demuth, H.U., and Stubbs, M.T. (2007). Crystal structure of the incretin-bound extracellular domain of a G protein-coupled receptor. Proc Natl Acad Sci U S A *104*, 13942-13947.

Patterson, J.T., Li, P., Day, J.W., Gelfanov, V.M., and Dimarchi, R.D. (2013). A hydrophobic site on the GLP-1 receptor extracellular domain orients the peptide ligand for signal transduction. Mol Metab *2*, 86-91.

Pettersson, U.S., Walden, T.B., Carlsson, P.O., Jansson, L., and Phillipson, M. (2012). Female mice are protected against high-fat diet induced metabolic syndrome and increase the regulatory T cell population in adipose tissue. PLoS One *7*, e46057.

Pocai, A. (2012). Unraveling oxyntomodulin, GLP1's enigmatic brother. J Endocrinol *215*, 335-346.

Pocai, A., Carrington, P.E., Adams, J.R., Wright, M., Eiermann, G., Zhu, L., Du, X., Petrov, A., Lassman, M.E., Jiang, G., et al. (2009). Glucagon-like peptide 1/glucagon receptor dual agonism reverses obesity in mice. Diabetes *58*, 2258-2266.

Ranganath, L.R. (2008). The entero-insular axis: implications for human metabolism. Clin Chem Lab Med *46*, 43-56.

Rizvi, A.A., and Rizzo, M. (2022). The Emerging Role of Dual GLP-1 and GIP Receptor Agonists in Glycemic Management and Cardiovascular Risk Reduction. Diabetes Metab Syndr Obes *15*, 1023-1030.

Rosenstock, J., Wysham, C., Frias, J.P., Kaneko, S., Lee, C.J., Fernandez Lando, L., Mao, H., Cui, X., Karanikas, C.A., and Thieu, V.T. (2021). Efficacy and safety of a novel dual GIP and GLP-1 receptor agonist tirzepatide in patients with type 2 diabetes (SURPASS-1): a double-blind, randomised, phase 3 trial. Lancet *398*, 143-155.

Runge, S., Thogersen, H., Madsen, K., Lau, J., and Rudolph, R. (2008). Crystal structure of the ligand-bound glucagon-like peptide-1 receptor extracellular domain. J Biol Chem *283*, 11340-11347.

Samms, R.J., Christe, M.E., Collins, K.A., Pirro, V., Droz, B.A., Holland, A.K., Friedrich, J.L., Wojnicki, S., Konkol, D.L., Cosgrove, R., et al. (2021). GIPR agonism mediates weight-independent insulin sensitization by tirzepatide in obese mice. J Clin Invest *131*.

Samms, R.J., Coghlan, M.P., and Sloop, K.W. (2020). How May GIP Enhance the Therapeutic Efficacy of GLP-1? Trends Endocrinol Metab *31*, 410-421.

Slomski, A. (2021). Tirzepatide Safely Reduces Blood Glucose, Weight in Type 2 Diabetes. JAMA *326*, 691.

Torrecilla, I., Spragg, E.J., Poulin, B., McWilliams, P.J., Mistry, S.C., Blaukat, A., and Tobin, A.B. (2007). Phosphorylation and regulation of a G protein-coupled receptor by protein kinase CK2. J Cell Biol *177*, 127-137.

Underwood, C.R., Garibay, P., Knudsen, L.B., Hastrup, S., Peters, G.H., Rudolph, R., and Reedtz-Runge, S. (2010). Crystal structure of glucagon-like peptide-1 in complex with the extracellular domain of the glucagon-like peptide-1 receptor. J Biol Chem *285*, 723-730.

Varol, C., Zvibel, I., Spektor, L., Mantelmacher, F.D., Vugman, M., Thurm, T., Khatib, M., Elmaliah, E., Halpern, Z., and Fishman, S. (2014). Long-acting glucose-dependent insulinotropic polypeptide ameliorates obesity-induced adipose tissue inflammation. J Immunol *193*, 4002-4009.

Waterhous, D.V., and Johnson, W.C., Jr. (1994). Importance of environment in determining secondary structure in proteins. Biochemistry *33*, 2121-2128.

Willard, F.S., Douros, J.D., Gabe, M.B., Showalter, A.D., Wainscott, D.B., Suter, T.M., Capozzi, M.E., van der Velden, W.J., Stutsman, C., Cardona, G.R., et al. (2020). Tirzepatide is an imbalanced and biased dual GIP and GLP-1 receptor agonist. JCI Insight *5*.

Wolff, M., Gast, K., Evers, A., Kurz, M., Pfeiffer-Marek, S., Schuler, A., Seckler, R., and Thalhammer, A. (2021). A Conserved Hydrophobic Moiety and Helix-Helix Interactions Drive the Self-Assembly of the Incretin Analog Exendin-4. Biomolecules *11*.

Yang, P.Y., Zou, H., Amso, Z., Lee, C., Huang, D., Woods, A.K., Nguyen-Tran, V.T.B., Schultz, P.G., and Shen, W. (2020). New Generation Oxyntomodulin Peptides with Improved Pharmacokinetic Profiles Exhibit Weight Reducing and Anti-Steatotic Properties in Mice. Bioconjug Chem *31*, 1167-1176.

Zhao, L., Wang, B., Wang, L., Zhao, X., Chen, Z., and Sun, L. (2019). Design, screening and biological evaluation of novel fatty acid chain-modified oxyntomodulin-based derivatives with prolonged glucose-lowering ability and potent anti-obesity effects. Org Biomol Chem *17*, 7760-7771.

Zou, C., Wang, Y., and Shen, Z. (2005). 2-NBDG as a fluorescent indicator for direct glucose uptake measurement. J Biochem Biophys Methods *64*, 207-215.

Figure legends:

Figure 1: Discovery of I-M-150847 a GLP-1R GIPR dual agonist **A**. Screening of peptide analogs in HEK 293 cells expressing GLP-1R for the cAMP generation as measured by luciferase reporter assay. Y-axis represents relative Luciferase units normalized by β-galactosidase expression. The data expressed as fold over the basal is presented as means ± SD. Lira: Liraglutide; Oxm: Oxyntomodulin, EX-4: Exendin-4. **B.** Screening of peptide analogs in HEK 293 cells expressing GIPR for the cAMP generation as measured by luciferase reporter assay.**C.** Screening of peptide analogs in HEK 293 cells expressing GCGR for the cAMP generation as measured by luciferase

reporter assay. **D.** Dose-response curve of stimulation of cAMP generation of I-M-

150847 vis-à-vis Exendin-F1 and Tirzepatide in HEK 293 cells expressing GLP-1R as

measured by cAMP responsive element luciferase reporter assay. EC_{50} represented as

mean ±SE of three independent experiments.**E.** Dose-response curve of stimulation of

cAMP generation of I-M-150847 vis-à-vis Tirzepatide in HEK 293 cells expressing

GIPRR as measured by cAMP responsive element luciferase reporter assay. EC_{50} is

represented as the mean ±SE of two independent experiments carried out in duplicate.

F. Dose-response curve of stimulation of cAMP generation of I-M-150847 relative to

exendin-4 in BRIN-BD11 pancreatic beta cells as measured by cAMP responsive

element luciferase reporter assay. EC_{50} is represented as the mean ±SEM of three

independent experiments carried out in duplicate.

Figure 2: I-M-150847 facilitates prompt redistribution of GLP-1R at the plasma

membrane following activation. **A.** The time-course of cAMP accumulation in BRIN-

BD11 pancreatic beta cells following activation by I-M-150847 or exendin-4. The cells

were incubated with 1μM I-M-150847 or exendin-4 and 3 minutes after the incubation

the excess ligand was washed with KRB buffer. The cAMP was measured at 1 min, 10,

30, and 60 min after KRB wash using Direct cAMP Enzyme Immunoassay. The data are

expressed as mean ±SE of three independent experiments *$p<0.05$; ***$p<0.001$, one

way-ANOVA Tukey's multiple comparison test, comparing cAMP generation upon

agonist treatment relative to basal cAMP accumulation. **B.** Confocal images studying

GLP-1R internalization at different time points upon exendin-4 or I-M-150847 treatment,

representative image from 4 independent experiments, n=15 cells each time point, scale

bars 10μM. **C.** Colocalization of GLP-1R -GFP with lysotracker red 45 min after

activation with exendin-4 or I-M-150847 as evidenced in confocal microscopy,

representative image from 2 independent experiments, n=5 for each experiment. Image

quantification was carried out in Image J (JACoP plugin) and co-localization was

measured by Pearson correlation coefficient (r) represented as mean ± SE; **$p < 0.01$,

n=5, student's t-test (unpaired) comparing correlation coefficient (Pearson r) between

exendin-4 and I-M-150847 treated cells. **D;(i)** Schematic diagram of the agonist

rechallenge in BRIN-BD11 pancreatic beta cells. The cells were subjected to treatment

with exendin-4 or I-M-150847 for 3 minutes following which the cells were washed and

incubated for another 45-minute following which they are rechallenged with exendin-4 or

I-M-150847 and cAMP generation was measured; **(ii)** Agonist rechallenge experiment;

the y-axis represents cAMP accumulation (pmol/mg protein), the data is expressed as

mean ± sem; *$p < 0.05$, ***$p < 0.001$, one-way ANOVA, n=3 independent experiments,

Tukey's multiple comparison test comparing cAMP accumulation by agonists pre and

post rechallenge. **E:** Schematic illustration of GLP-1R trafficking upon activation by

exendin-4 or I-M-150847.

Figure 3: Functional response of I-M-150847 in cultured pancreatic beta cells and

mouse adipocytes. **A. (i)** Schematic of GSIS in BRIN-BD11 pancreatic beta cells upon

acute and chronic treatment with exendin-4 or I-M-150847. **(ii)** Acute and prolonged

GSIS upon exendin-4 or I-M-150847 treatment. Y axis represents insulin secretion

(pg./mg protein) expressed as the fold increase over basal insulin secretion at 0.1mM

glucose. The data are expressed as mean ± SE **$p < 0.01$, one-way ANOVA, n=3

Tukey's multiple comparison test comparing GSIS upon 30 min treatment (acute mode)

with exendin-4 and I-M-150847; ns=non-significant comparing GSIS upon 6h

(prolonged) exendin-4 or I-M-150847 treatment with insulin secretion at 16.7mM

glucose. **B (i)** Schematic of I-M-150847 or exendin-4 mediated cAMP generation upon

overnight pre-exposure with agonists. **(ii)** The cAMP generation upon exendin-4 or I-M-

150847 treatment following overnight pre-exposure with agonists. The y axis represents

cAMP accumulation (pmol/mg protein), the data are expressed as mean ± SE;

***p<0.001, one-way ANOVA, n=3, Tukey's multiple comparison test comparing cAMP

generation between exendin-4 and I-M-150847 treatment, ns=non-significant comparing

prolonged cAMP generation upon agonist treatment with untreated control cells. **C(i)**

Schematic of insulin-stimulated glucose uptake upon overnight pre-exposure with

exendin-4 or I-M-150847 in 3T3-L1 adipocytes; **(ii)** Insulin-stimulated glucose uptake

upon overnight pre-exposure with exendin-4 or I-M-150847 in 3T3-L1 adipocytes; the

data are expressed as mean ± SE; ***p<0.001, n=3, one-way ANOVA, Tukey's multiple

comparison test.

Figure 4: Comparison of I-M-150847 and exendin-4 in providing glycemic benefits and

reducing obesity in DIO mice. **A.** Schematic diagram of the diet and the treatment

paradigm as well as the pharmacological parameters measured upon the treatment of I-

M-150847 and exendin-4. **B.** Measurement of the fasting blood glucose at 0 day versus

the 10th day of treatment; **(i)** Fasting blood glucose in control mice on high fat diet at 0

day and 10th day of treatment relative to the group fed on the chow diet, results are

mean ± SE, *** p<0.001 one-way ANOVA, Tukey's multiple comparison test, n=6 each

group comparing blood glucose between chow diet and high fat diet -fed group; **(ii)**

Comparative fasting blood glucose in DIO mice upon exendin-4 administration at 0 day

and 10th day of treatment; , *** p<0.001, one-way ANOVA, Tukey's multiple comparison

test n=6, comparing fasting blood glucose (mean ± SE)   at 0 day versus the 10th

day**;(iii)** Comparative fasting blood glucose in DIO mice upon I-M-150847 administration

at 0 day and 10th day of treatment; *** p<0.001, one-way ANOVA, Tukey's multiple

comparison test n=6, comparing fasting blood glucose (mean ± SE) at 0 day versus the

10th day; **(iv)** Fasting blood sugar in I-M-150847 treated mice as compared to exendin-4

treated group and the group receiving normal saline as vehicle, results are expressed

as mean ± SE; *** p<0.001, one-way ANOVA, Tukey's multiple comparison test, n=6

for each group, comparing fasting blood glucose on the 10th day between control mice

receiving normal saline versus exendin-4 treated and I-M-150847 treated group. **C.**

Intraperitoneal glucose tolerance (IPGTT) in DIO mice receiving normal saline as the

vehicle (control), exendin-4 or I-M-150847 100nmol/kg bodyweight twice daily through

subcutaneous administration, the data are expressed as mean ± SE ***p<0.001, *p<

0.05,(exendin-4 versus I-M-150847 treated group) as assessed by two-way ANOVA,

Bonferroni (n=6, for each group) at different time points; glucose lowering during IPGTT

as assessed by the area under the curve (AUC, mean ± SE, arbitrary units) in the

vehicle, exendin-4 and I-M-150847 treated group, ***p<0.001, one-way ANOVA,

Tukey's multiple comparison test, n=6 each group comparing the vehicle, exendin-4,

and I-M-150847 treated mice. **D**. Insulin tolerance test (ITT) in DIO mice receiving

normal saline as the vehicle (control), exendin-4 or I-M-150847 100 nmol/kg body

weight twice daily through subcutaneous administration, data are expressed as mean ±

SE,   **p<0.01, (exendin-4 and I-M-150847 treated group) assessed by two-way

ANOVA, Bonferroni posttests (n=6, for each group) at different time points; ns=

nonsignificant; glucose lowering upon insulin administration assessed by the area under

the curve (AUC, mean ± SE, arbitrary units) in the vehicle, exendin-4 and I-M-150847

treated group, ***$p<0.001$, one-way ANOVA, Tukey's multiple comparison test, n=6

each group comparing AUC of ITT in the vehicle, exendin-4 and I-M-150847 treated

mice. **E.** Cumulative food intake in mice treated with vehicle, exendin-4, and I-M-150847

measured every day; the results are expressed as mean ± SE, *$p<0.05$ one-way

ANOVA, Tukey's multiple comparison test, n=6 for each group; *$p<0.05$, ***$p<0.001$, the

statistical significance of the difference in cumulative food intake (I-M-150847 versus

exendin-4 treated group, mean ± SE, n=6 for each group) at specific time points as

assessed by two-way ANOVA, Bonferroni posttests. **F.** Body weight gain in mice

receiving subcutaneous administration of normal saline (vehicle), exendin-4 or I-M-

150847 (n=6 each group). Results are expressed as mean ±SE, ***$p<0.001$ one-way

ANOVA, Tukey's multiple comparison test comparing % body weight gain among

different groups. **$p<0.01$, ***$p<0.001$, the % bodyweight gain at specific time points (I-

M-150847 versus exendin-4) was analyzed by two-way ANOVA, Bonferroni posttests

(n=6 per group). **G.** Tissue weight of mice treated with normal saline (vehicle), exendin-

4, and I-M-150847 (n=6 per group), (i) liver, (ii) inguinal WAT, (iii) epididymal WAT, (iv)

retroperitoneal WAT and (v) mesenteric WAT. Results are expressed as mean ±SE;

**$p<0.01$, ***$p<0.001$, one-way ANOVA, Tukey's multi comparison test (n=6 for each

group).

Table 1: In vitro GLP-1R, GIPR, and GCGR activity of peptide analogs

No	Peptide	Constituent amino acids	GLP-1R		GIPR		GCGR	
			% Liraglutide activity	SD	% GIP activity	SD	% Oxm activity	**SD**
1	**Liraglutide**	HAEGTFTSDVSSYLEGQAAKcEFIAWLVKGRG	**100**	**6.7**	24.08	2.13	10.90	0.13
2	**GIP**	YAESTFISDYSIAMDKIHQQDFVNWLLAQKGKK NDWKHNITQ	31.86	3.8	**100**	**6.6**	23.39	1.74
3.	**Exendin 4**	HGEGTFTSDLSKQMEEEAVRLFIEWLKNGGPSS GAPPPS-NH2	**140.24**	**3.15**	**28.20**	**1.17**	21.45	1.49
4	**Oxyntomo dulin**	HSQGTFTSDYSKYLDSRRAQDFVQWLMNTKRN RNNIA	**96.26**	**0.96**	30.76	1.68	100	1.4
5	**I-M-150844**	HGEGTFTSDVF	21.66	2.69	15.55	2.37	6.49	0.45
6.	**I-M-150845**	YGEGTFTSDYSIAMDKIHQQDFVNWLVKGRG (3563.9)	**76.78**	**6.32**	**86.9**	**13.4**	11.5	0.9
7	**I-M-150846**	YGEGTFTSDYSIAMDKIHQQDFVNWLMNTKRNR NNIA (4379.8)	38.21	0.6	**65.65**	**6.7**	7.14	0.4
8	**I-M-150847**	YGEGTFTSDLSKQMEEEAVRLFIEWLMNTKRNR NNIA (4392.88)	**122.95**	**2.0**	**108.0**	**5.1**	14.93	1.68
9	**Basal**	**Phosphate Basal Saline**	29.96	1.5	16.43	**5.0**	8.12	0.2

Table legend: The activity of the peptide analogs in vitro in GLP-1R, GIPR, and GCGR

assay system as measured by cyclic AMP responsive element luciferase reporter

assay. % Liraglutide activity is defined as the normalized relative luciferase units

generated upon treatment of HEK –GLP-1R expressing cells with 1µM liraglutide. %

GIP activity is defined as the normalized relative luciferase units generated upon

treatment of HEK –GIPR expressing cells with 10µM GIP. % GCGR activity is defined

as the normalized relative luciferase units generated upon treatment of HEK –GCGR

expressing cells with 10µM Oxyntomodulin. Oxm =Oxyntomodulin. The numbers in

brackets represent the molecular mass of the peptide. The data represent the means ±

SD of the experiment carried out in duplicate in each heterologous assay system. The

amino acids that correspond to Liraglutide, exendin-4, oxyntomodulin, and GIP are

colored green, black, blue, and red respectively.

Figure 1.

	I-M-150847	Exendin-F1	Tirzepatide
log(agonist) vs. response -- Variable slope			
Best-fit values			
BOTTOM	1.109	1.199	1.374
TOP	4.142	4.157	4.660
LOGEC50	1.214	1.487	-0.01463
HILLSLOPE	0.4697	0.7744	1.563
EC50	16.35	30.71	0.9669
Span	3.033	2.958	3.286

	I-M-150847	Tirzepatide
log(agonist) vs. response -- Variable slope		
Best-fit values		
BOTTOM	1.327	1.416
TOP	3.888	5.242
LOGEC50	1.461	-0.2148
HILLSLOPE	1.021	0.3552
EC50	28.90	0.6098

	Exendin-4	IM150847
log(agonist) vs. response -- Variable slope		
Best-fit values		
Bottom	1.478	1.332
Top	5.626	5.776
LogEC50	0.09472	1.157
HillSlope	0.3943	0.4830
EC50	1.244	14.35
Span	4.148	4.444

Figure 2

Figure 3.

Figure 4.

Figure 1- Figure supplement 1 : I-M-150845

A.

B.

C.

Mol probity score 0.5
Ramachandran Favored: 100%
Ramachandran outliers : 0%
C-Beta Deviations : 0

Figure1-Figuresupplement 1: Design of the peptide I-M-150845

A. Amino acid constituents of peptide I-M-150845. The amino acids common to

exendin-4, GLP-1, and GIP are colored black, green, and red respectively.

B. Cartoon of the predicted secondary structure of I-M-150845 as modeled by

Swiss- model protein structure homology modeling server. C=loop or irregular,

H=α helix, T= Hydrogen bonded turns.

C. Ramachandran plot for I-M-150845 using Mol Probity version 4.4 in Swiss-model

protein structure homology modeling server.

Fig 1-Figure supplement 2 : I-M-150846

A.

```
 1  2  3  4  5  6  7  8  9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37
 Y  G  E  G  T  F  T  S  D  Y  S  I  A  M  D  K  I  H  Q  Q  D  F  V  N  W  L  M  N  T  K  R  N  R  N  N  I  A
```

B.

```
C C T T H H H H H H H H H H H H H H H H H H H H H H H H H H H H H T T T T C -
Y G E G T F T S D Y S I A M D K I H Q Q D F V N W L M N T K R N R N N I A
```

C.

Ramachandran plot

Molprobity score : 1.38
Ramachandran favored 91.18%
Ramachandran outliers 2.94 A4Gly
C Beta Deviations: 1, A36 ILE

Figure1- Figure supplement 2: Design of the peptide I-M-150846

A. Amino acid constituents of peptide I-M-150846. The amino acids common to

exendin-4, GIP, and oxyntomodulin are colored black, red, and blue respectively.

B. Cartoon of the predicted secondary structure of I-M-150846 as modeled by

Swiss- model protein structure homology modeling server. C=loop or irregular,

H=α helix, T= Hydrogen bonded turns.

C. Ramachandran plot for I-M-150846 using Mol Probity version 4.4

Figure 1-Figure supplement 3 : I-M-150847

A.

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37

YGEGTFTSD L S K Q M E E E A V R LF I E W LMN T KRNR NN I A

B.

D. Ramachandran plot

Molprobity score: 2.00
Ramachandran favored: 84.85%
Ramachandran outliers: 9.09% A4GLY,
A34ASN,A29THR
C beta Deviations : 2, A29 THR, A30LYS

C.

Figure 1-Figure supplement 3: Design and physical characterization of peptide I-

M-150847.

A.　Amino acid constituents of I-M-150847. The amino acids common to exendin-4,

GIP, and oxyntomodulin are colored black, red, and blue respectively

B.　Cartoon of the predicted secondary structure of I-M-150847 as modeled by

Swiss- model protein structure homology modeling server. C=loop or irregular,

H=α helix, T= Hydrogen bonded turns. G= 3-helix (3_{10}- helix), S= bend.

C.　CD analysis of I-M-150847.

D.　Ramachandran plot for I-M-150847 using Mol Probity version 4.4

Figure 1-Figure supplement 4

HEK GCGR assay

log(agonist) vs. response	
Best-fit values	
BOTTOM	0.9120
TOP	3.919
LOGEC50	1.730
EC50	53.75
Span	3.007

16 **Figure-1 Figure supplement 4:**

17 Dose response curve of I-M-150847 mediated cAMP response in HEK GCGR cell line
18 measured by cAMP-responsive element luciferase reporter assay. Data are represented
19 as mean ± SE of 3 independent experiments

Figure 1-figure supplement 5

Figure 1 Figure supplement 5:

LC-MS analysis of peptide I-M-150847 amide.

Figure 1-Figure supplement 6

Column	: Luna 5µm C18(2) 100A(250 x 4.6mm)
Mobile Phase	: A= 0.1%TFA (Aq) ,B=ACN
Gradient	:0/2,2/2,15/70,20/95,25/100,30/100,32/2,42/2
Flow	:1.0ml/min
Temperature	:25.0 C

1 PDA Multi 1/220nm 4nm

PeakTable

Ch1 220nm 4nm

Peak#	Ret. Time	Area	Area %
1	15.189	24311	0.806
2	15.328	58590	1.943
3	15.487	2870294	95.201
4	15.648	26346	0.874
5	15.680	35435	1.175
Total		3014977	100.000

2 **Figure 1-Figure supplement 6:**

3 RP-HPLC of peptide I-M-150847 amide

Figure 3 –Figure supplement 1

3 **Figure -3 Figure supplement 1:**

4 Relative mRNA expression of GLP-1R GIPR measured by the $2^{-\Delta\Delta CT}$ method (Livak and
5 Schmittgen, 2001). Data are expressed as mean ± SE; n=3; ***p<0.001 comparing
6 GLP-1R and GIPR expression in BRIN-BD11 pancreatic beta cells.

2