IL-21 signaling promotes early dissemination of KSHV infection in B lymphocytes

Nedaa Alomari[1], Farizeh Aalam[1], Romina Nabiee[1], Jesus Ramirez Castano[1] and Jennifer Totonchy‡

[1]Biomedical and Pharmaceutical Sciences, Chapman University, Irvine, CA USA

‡Corresponding Author

Jennifer Totonchy

Chapman University School of Pharmacy

9410 Jeronimo Road

Irvine, CA

Phone: 714-516-5438

Email: totonchy@chapman.edu

Short Title: IL-21 and KSHV infection of B cells

1 Abstract

Kaposi's sarcoma-associated herpesvirus (KSHV) extensively manipulates the host immune

system and the cytokine milieu, and cytokines are known to influence the progression of KSHV-

associated diseases. However, the precise role of cytokines in the early stages of KSHV infection

remains undefined. Here, using our unique model of KSHV infection in tonsil lymphocytes, we

investigate the influence of host cytokines on the establishment of KSHV infection in B cells.

Our data demonstrate that KSHV manipulates the host cytokine microenvironment during early

infection and susceptibility generally associated with downregulation of multiple cytokines.

However, we show that IL-21 signaling promotes KSHV infection by promoting both plasma

cell numbers and increasing KSHV infection in plasma cells. Our data reveal that IL-21

producing T cells, particularly Th17/Tc17 and central memory CD8+ T cells may represent

immunological factors that modulate host-level susceptibility to KSHV infection. These results

suggest that IL-21 plays a significant role in the early stages of KSHV infection in the human

immune system and may represent a novel mechanism to be further explored in the context of

preventing KSHV transmission.

16 Author Summary

Very little is known about how KSHV is transmitted and how it initially establishes infection in a

new human host and this lack of information limits our ability to prevent KSHV-associated cancers

by limiting its person-to-person transmission. Saliva is thought to be the primary route of person-

to-person transmission for KSHV, making the tonsil a likely first site for KSHV replication in a

new human host. In particular, the tonsil is likely to be the first place KSHV is able to enter B

cells, which are thought to be a major site of persistent infection. Our previous work identified

plasma cells as a highly targeted cell type in early KSHV infection in cultured cells from human

tonsil. In this study, we show that the human cytokine IL-21 promotes both overall KSHV infection

and the establishment of infection in plasma cells. We also investigate the immunological

mechanisms underlying this effect. Our results demonstrate that IL-21 and IL-21-producing cells

are a novel factor that influences the initial establishment of KSHV infection in humans.

Introduction

Kaposi's Sarcoma Herpesvirus (KSHV) is a lymphotropic gamma-herpesvirus, originally

discovered as the causative agent of Kaposi Sarcoma (KS) [1]. KS is a highly proliferative tumor

derived from lymphatic endothelial cells [2]. KSHV is also associated with the B cell

lymphoproliferative diseases, Primary Effusion Lymphoma (PEL) and Multicentric Castleman's

Disease (MCD) [3, 4], as well as the inflammatory disorder KSHV inflammatory cytokine

syndrome (KICS) [5]. KSHV is linked to 1% of all human tumors, and the World Health

Organization (WHO) has classified it as class I carcinogen [6, 7]. KSHV infection is asymptomatic

in most healthy individuals, and KSHV-associated malignancies arise primarily in

immunocompromised patients. Indeed, KS remains one of the most common cancers in people

living with HIV/AIDS [8].

The geographical distribution of KSHV is not ubiquitous. KSHV infection is endemic in sub-

Saharan Africa and in the Mediterranean basin. KSHV prevalence is also high in subpopulations

in other parts of the world such as men who have sex with men (MSM). Saliva is the only secretion

where KSHV DNA is commonly detected [9], and, based on this, person-to-person transmission

of KSHV is thought to occur via saliva. The oral lymphoid tissues are rich in KSHV target cell

types including lymphatic endothelial cells and B cells, and are therefore a likely site for the initial

establishment of KSHV infection in a new human host. However, the exact mechanisms for KSHV

transmission and how environmental, behavioral and host factors influence transmission and early

infection events remain to be established. This gap in our understanding dramatically affects our

ability to find efficient strategies to decrease the transmission or influence host-level susceptibility

to KSHV infection. One of the main factors that has been linked to KSHV-associated

lymphoproliferations is cytokine dysregulation [10]. However, their contribution to the early

stages of KSHV infection and whether the cytokine milieu in the oral cavity contributes to host-

level susceptibility to KSHV infection is unclear. KSHV-infected cells in KSHV-associated

lymphoproliferative diseases generally display plasma cell or plasmablast features [11], and our

recent work showed that KSHV targets plasma cells in early infection in tonsil B lymphocytes

[12]. Whether the KSHV-infected plasma cells we observe arise from direct infection or

differentiation from a precursor B cell subset remains to be established. In either case, it is likely

that cytokines play a role in supporting the survival and/or differentiation of KSHV-infected

plasma cells.

IL-21 is a pleiotropic cytokine that has diverse effects on B cell, T cell, macrophage, monocyte,

and dendritic cell biology. It is produced mainly by natural killer T (NKT) cells and CD4+ T cells,

including follicular helper (TFH) cells [13]. The IL-21 receptor is expressed by several immune

cells, including B and T cells and is comprised of a unique IL-21R subunit and the common

cytokine receptor γ chain (CD132), which is also part of the receptor for IL-2, IL-4, IL-7, IL-9,

and IL-15 [14]. IL-21 plays a critical role in B cell activation and expansion [15], and plays a

critical role in B cell differentiation to immunoglobulin (Ig)-secreting plasma cells. The regulation

of maturation of B cells into plasma cell is driven by the several transcription factors including

Blimp1 and Bcl6 [16], which can both be induced by IL-21 signaling, indicating that IL-21 is an

important regulator of plasma cell differentiation [17, 18].

There are few studies to date examining the contribution of IL-21 to KSHV infection and KSHV-

associated disease. However, IL-21 is detected in interfollicular areas in MCD patients [19].

Moreover, IL-21 induces differentiation of B-lymphoblastoid cell lines (BCLs) into late

plasmablasts/early plasma cells and regulates the expression of many latent proteins in

EBV[+] Burkitt lymphoma cell lines [20, 21]. Importantly, IL-21 signaling is required for the

establishment of MHV-68 infection and the generation for MHV-68 infected long-lived plasma

cells in mice [22].

In this study, we use our well-established tonsil lymphocyte infection model to explore whether

KSHV alters cytokine secretion early in infection and whether cytokine levels have an effect on

the establishment of KSHV infection. We identify IL-21 as a factor that specifically influences

KSHV infection in plasma cells, which we previously characterized as a highly targeted cell type

in early infection [12]. We demonstrate that IL-21 signaling increases plasma cell levels in our *ex*

vivo model system and that this effect promotes overall KSHV infection. We explore the

mechanisms of IL-21 signaling by establishing which B cell types are responding IL-21 and what

T cell subsets are producing IL-21 in our model.

We determine that expression of IL-21 receptor (IL-21R) on naïve and classical memory T cells

is correlated with increased KSHV infection in IL-21 treated cultures, and we demonstrate that

both KSHV infection and IL-21 treatment modulates IL-21R expression on B cells. We examine

IL-21 secretion by T cell subsets in our tonsil lymphocyte cultures and demonstrate that the

magnitude and distribution of IL-21 secretion varies considerably based on tonsil donor, but is not

significantly modulated by KSHV infection. Interestingly, RoRγT+ T cells (both CD4+ and CD8+)

contributed substantially to IL-21 secretion and were positively correlated with KSHV infection

of plasma cells at 3 dpi. Moreover, IL-21 secretion by CD8+ central memory T cells was positively

correlated with both overall KSHV infection and plasma cell targeting.

These results identify IL-21 signaling as a factor that influences the establishment of KSHV

infection in B lymphocytes and, together with our previous work, underscores the importance of

plasma cell biology in the initial establishment of KSHV infection in the oral lymphoid tissues.

Based upon this work, we conclude that specific IL-21 secreting T cell subsets represent an

important susceptibility factor for KSHV transmission.

Results

Host cytokines influence the establishment of KSHV infection B lymphocytes

Despite the critical interplay between KSHV and host cytokine signaling, little is known about

whether cytokines influence host susceptibility to KSHV infection. In fact, the roles of

proinflammatory cytokines during KSHV infection have been studied mostly in naturally-infected

human B cell lines derived from PEL [23, 24]. In order to examine whether cytokines alter the

early stages of KSHV infection in the tonsil, we quantitated the levels of 13 cytokines in the

supernatants of Mock and KSHV-infected tonsil lymphocyte cultures at 3 days post-infection

using a bead-based multiplex immunoassay. This dataset includes 33 independent infections using

24 unique tonsil specimens. IL-6, IFNg, TNFa and IL-22 were the most prevalent cytokines in our

cultures based on the median values overall (Fig 1A, panel order). IL-6 was the only cytokine

significantly induced in KSHV-infected cultures compared to Mock cultures (Fig 1A) and the

magnitude of IL-6 induction by KSHV infection is far greater than the effect of infection on any

other cytokine (Fig 1B). This result is consistent with our previous results in cultures containing

only naïve B lymphocytes [25]. Our data also reveals statistically significant reductions in IL-5

and IL-4 concentrations in KSHV-infected cultures compared to Mock cultures (Fig 1A), but these

changes are very small compared to those seen with IL-6 (Fig 1B). Interestingly, IFNg

concentrations were highly affected by KSHV infection, but there were sample-specific

differences in whether this effect was positive or negative (Fig 1B).

In order to determine whether cytokines affect the establishment of KSHV infection, we examined

whether the concentration of cytokines in the supernatants of KSHV-infected cultures is correlated

with the level of infection in B lymphocytes (based on GFP reporter expression) in the same culture

by flow cytometry analysis (Fig 1C & D). On a per-sample level, many individual cytokines

(notably IL-6 and IFNg) were induced or repressed independent of susceptibility. However, there

is a cluster of highly susceptible samples (ND19, ND40, ND41, ND32, ND4) in which multiple

cytokines are repressed in the KSHV-infected cultures (Fig 1C). Pairwise comparisons between

overall GFP level in B lymphocytes and the level of each cytokine in the KSHV-infected cultures

revealed universally negative correlations between overall KSHV infection and cytokine levels

with lower cytokine levels observed in more susceptible samples. These negative correlations were

statistically significant for IL-2, IL-9, IL-10, TNFa, IL-4 and IL-22 (Fig 1D). Since plasma cells

were identified as a highly targeted cell type in our previous study [12], we examined the

correlations between cytokine levels in KSHV-infected cultures and infection in the CD138+

plasma cell subset. This analysis revealed negative correlations similar to those seen with overall

infection with IL-13, IL-9, IFNg, TNFa and IL-22 levels showing statistically significant negative

correlations with plasma cell infection. However, in this analysis IL-21 levels showed a significant

positive   correlation   with   plasma   cell   infection   (Fig   1E).   Taken   together,   these   data

demonstrate that (1) KSHV infection influences the production of multiple cytokines in our *ex vivo*

infection model, (2) lower cytokine levels and/or repression of cytokines during infection are

generally associated with higher susceptibility to KSHV infection, (3) several individual cytokines

show significant negative associations with susceptibility to KSHV infection and (4) IL-21 is

positively correlated with KSHV infection of plasma cells. Overall, these data suggest that distinct

inflammatory responses in each tonsil specimen contribute to variable susceptibility to KSHV

infection.

IL-21 supplementation increases KSHV infection in tonsil B lymphocytes

Because IL-21 production was positively correlated with plasma cell infection in our initial dataset

(Fig 1E), we wanted to examine the impact of manipulating IL-21 levels on the establishment of

KSHV infection. To do this, we performed Mock infection or KSHV infection in 12 unique tonsil

samples and supplemented the resulting cultures with varying concentrations of recombinant IL-

21. At 3 dpi, we analyzed these cultures for GFP+ B lymphocytes by flow cytometry to assess the

magnitude of KSHV infection (Fig 2A & B). Although the specimens included in this data set had

high variability in their baseline susceptibility, we can see increased infection in response to IL-

21 treatment, and the effect seems to be particularly strong in the more susceptible samples (Fig

2A). Normalization of the data to each specimen's untreated control reveals that at 10/12 samples

show increased infection upon treatment with 100pg/ml of IL-21. Importantly, most of these

concentrations were higher than what was observed in our initial dataset quantitating native

cytokine secretion in our culture system (Fig 1A), which may explain why we didn't observe an

association of IL-21 secretion with overall infection in that data (Fig 1D). We then repeated these

supplementation experiments with only the 100pg/ml dose of recombinant IL-21 in an additional

12 tonsil specimens and examined both overall infection and subset-specific responses in these

cultures at 3 dpi using our B cell immunophenotyping panel. Similar to the initial dataset, this

analysis shows increased infection in response to recombinant IL-21 in the majority of tonsils, and

the difference in GFP+ B lymphocytes was statistically significant in IL-21 treatment compared to

control (p=0.02, F=6.4) (Fig 2C).

IL-21 increases plasma cell frequency and susceptibility in primary human tonsil B

lymphocytes

In order to determine whether IL-21 signaling influences the overall distribution of B cell subsets

in both Mock and KSHV infected cultures, we performed B cell immunophenotyping. Analysis of

overall subset frequencies, with all subsets represented as a fraction of viable CD19+ B cells, at 3

dpi revealed that most subsets did not change with either KSHV infection or IL-21 treatment. We

did observe a statistically significant increase in CD20+ B cells and plasmablasts in the KSHV-

infected conditions independent of IL-21 treatment. Naïve B cells showed significant main effects

for both infection and treatment, with both KSHV infection and IL-21 treatment reducing naïve B

cells in the culture, but the interaction was not significant. Importantly, this analysis revealed a

highly significant increase in total plasma cell frequency associated with both IL-21 treatment and

KSHV infection with a significant interaction of the two variables. This effect seems to be

restricted to CD20+ plasma cells in the mock conditions and distributed between both CD20+ and

CD20- plasma cells in the KSHV-infected cultures. However, both CD20+ and CD20- plasma

cells showed statistically significant interactions between infection and treatment (Fig 2D, Table

1). Post hoc paired T tests revealed significant differences with IL-21 treatment on total plasma

cells (p=0.0002) and CD20+ plasma cells (p=0.002) for the KSHV-infected conditions only. The

IL-21 effect on naïve B cell frequencies was significant for both Mock (p=0.01) and KSHV

infected conditions (p=0.003). Moreover, in IL-21 treated conditions there was a significant

difference between Mock and KSHV cultures for total plasma cells (p=0.0003), CD20+ plasma

cells (p=0.02) and naïve B cells (p=0.006). One interpretation of this data is that IL-21 can

independently drive plasma cell differentiation of B cells in our culture system resulting in

decreased naïve cells and increased plasmablast and plasma cells, and KSHV infection acts

synergistically with IL-21 to potentiate the same effect. We next wanted to determine whether IL-

21 treatment altered the B cell subset-specific distribution of KSHV infection. For this analysis,

we quantitated the percent of each B cell subset that was GFP+ to determine the within-subsets

distribution of KSHV infection in control or IL-21 treated cultures (Fig 2E). In this analysis, we

observed a significant increase in plasma cell targeting with IL-21 treatment (p=0.02, F=6.6). We

next examined whether increased frequency of plasma cells or increased plasma cell targeting was

directly correlated with the observed increase in overall KSHV infection in the IL-21 treated

conditions (Fig 2F). These results reveal a significant linear correlation between total GFP and

plasma cell frequency (r=0.7, p=0.007) and a weaker correlation between total GFP and the

frequency of GFP+ cells within the plasma cell subset (r=0.56, p=0.04).

Taken together this data shows that IL-21 treatment promotes the establishment of KSHV infection

in human tonsil lymphocytes and that this increased infection is correlated with both increased

plasma cell frequencies and increased plasma cell infection. Thus, our results suggest that IL-21

signaling facilitates the establishment of KSHV infection by driving the proliferation, survival or

differentiation of plasma cells in our tonsil lymphocyte cultures. The observation that naïve B cell

frequencies are significantly decreased in KSHV-infected, IL-21-treated cultures (Fig 2D)

supports the latter conclusion that increased plasma cell frequencies are a result of B cell

differentiation in response to IL-21 signaling.

Table 1: Statistically significant effects from two-way repeated measures ANOVA analysis

in Fig 2D

Variable	Effect	DFn	DFd	F	p	ges
GFP	Treatment	1	13	6.425	0.025	0.081
GFP	Infection	1	13	102.896	1.52e-07	0.708
GFP	Tx:Inf	1	13	7.635	0.016	0.084
Plasma cells	Treatment	1	13	18.596	0.000844	0.163
Plasma cells	Infection	1	13	18.575	0.000848	0.166
Plasma cells (PC)	Tx:Inf	1	13	22.779	0.000364	0.09
Naive	Treatment	1	13	13.643	0.003	0.099
Naive	Infection	1	13	8.462	0.012	0.069
Plasmablast	Infection	1	13	6.023	0.029	0.02
CD20- PC	Tx:Inf	1	13	12.645	0.004	0.043
All CD20+CD19+	Infection	1	13	6.234	0.027	0.016
CD20+ PC	Treatment	1	13	4.858	0.046	0.028
CD20+ PC	Tx:Inf	1	13	7.798	0.015	0.012

<u>**Neutralization of IL-21 inhibits KSHV infection in primary tonsil B lymphocytes**</u>

We next wanted to determine whether neutralization of the natively-secreted IL-21 in our tonsil

lymphocyte cultures would affect the establishment of KSHV infection. To do this we performed

infections with Mock or KSHV-infection in 11 unique tonsil specimens, included varying

concentrations of an IL-21 neutralizing antibody in the resulting cultures, and assessed the

magnitude and distribution of KSHV infection at 3 dpi by flow cytometry. These results revealed

decreased KSHV infection in the presence of IL-21 neutralizing antibodies (Fig 3A). One-way

repeated measures ANOVA revealed a significant effect of IL-21 neutralization on GFP in KSHV

infected cultures (p=0.00001, F=9.4) and post-hoc Dunnett test revealed significance at the

100µg/ml dose (p=0.03). When each sample was normalized to its untreated control, we observed

that 9/11 samples had decreased infection in the presence of 100µg/ml IL-21 neutralizing antibody

and this increased to 10/11 samples at higher antibody doses (Fig 3B).

When we examined the overall frequency of B cell subsets at 3dpi in these experiments, we

observed statistically significant effects of IL-21 neutralization on a number of B cell subsets, but

there were no significant interaction effects with IL-21 neutralization and KSHV infection using

two-way repeated measures ANOVA (Fig 3C). Specifically, IL-21 neutralization increased IgG+

(p=0.04, F=5.0), memory (p=0.03, F=3.4), and double negative (p=0.02, F=3.8) B cells and

decreased IgM+ (p=0.0003, F=8.6) and transitional (p=0.01, F=4.1) subsets. Post-hoc paired T-

tests showed a statistically significant decrease of on both overall (p=0.02) and CD20+ (p=0.005)

plasma cell frequencies at the 200µg/ml dose in the mock infected cultures only. The observation

that the effect of IL-21 neutralization on plasma cell frequencies is restricted to mock infected

cultures is interesting in the context of our IL-21 supplementation data where we observed

significant main effects of both IL-21 and KSHV infection on plasma cell frequencies as well as a

significant interaction between the two factors (Fig 2D & Table 1). The two data sets taken together

support several interesting conclusions: (1) IL-21 affects plasma cell frequencies independent of

KSHV infection; evidenced by opposite significant effects of IL-21 supplementation and

neutralization in mock cultures, (2) KSHV infection affects plasma cell frequencies independent

of IL-21 signaling; evidenced by significant main effect of infection in supplemented cultures and

a lack of inhibition in KSHV-infected neutralized cultures, and (3) IL-21 and KSHV can

synergistically affect plasma cell frequencies; evidenced by the significant interaction effect and

the significant increase in plasma cell frequencies in KSHV-infected cultures that are

supplemented with IL-21.

We next wanted to determine whether neutralization of IL-21 influences the frequency of KSHV

infection within B cell subsets. One-way repeated measures ANOVA revealed a significant effect

of neutralization on infection of transitional B cells ($p=0.01$, $F=4.0$) but paired T tests were not

significant for any individual dose. When we plotted this data normalized to the per-sample, per-

subset infection rate of the untreated control cultures, we observed that neutralization was

associated with lower levels of infection in transitional, plasmablast, and CD20+ PC populations

(Fig 3D). We hypothesized that if IL-21 signaling is affecting overall infection by contributing

substantially to differentiation of KSHV-infected cells, we might observe a correlation between

overall infection and the contribution of specific subsets to infection within neutralized cultures

compared to control cultures. If this hypothesis is correct, subsets whose differentiation is

important to the establishment of infection would accumulate with IL-21 neutralization and this

accumulation would correlate with decreased overall levels of infection in the same cultures. To

examine this, we calculated the change in IL-21 neutralized cultures compared to control cultures

(neutralized-control) for both overall infection and the contribution of each subset to infection

(between subsets frequency of GFP), and performed correlation analysis using Pearson's method.

This analysis reveals that decreased overall infection with IL-21 neutralization was significantly

correlated with a decreased contribution of plasma cells (r=0.6, p=0.0002), CD20+ plasma cells

(r=0.6, p=0.0002), and transitional B cells (r=0.5, p=0.003) and an increased proportion of infected

germinal center cells (r=-0.4, p=0.02) (Fig 3E). Interestingly, these correlations were driven more

by sample-specific differences (indicated by point color) compared to dose response of

neutralization (indicated by point shape). This data could indicate that IL-21 signaling increases

the overall establishment of KSHV infection in tonsil lymphocytes by driving differentiation of

KSHV-infected germinal center cells into transitional and CD20+ plasma cells. Our previous

studies demonstrated that plasma cells display a mixture of lytic and latent KSHV infection [12].

Therefore, we wanted to determine whether the increase in overall KSHV infection with IL-21

treatment and decrease in infection with IL-21 neutralization is due to IL-21-mediated alterations

in KSHV lytic reactivation. To examine this, we performed RT-PCR for LANA (latent) and K8.1

(lytic) on total RNA from untreated, IL-21 supplemented or IL-21 neutralizing antibody treated,

KSHV-infected cultures from 8 unique tonsil specimens. GAPDH was used as a housekeeping

gene and normalizing factor for the viral gene expression data. This data is consistent with our

previous data showing a mix of lytic and latent transcripts in infected lymphocytes [12]. These

data reveal no significant influence of either supplementation or neutralization on lytic gene

expression (Fig 3F). In the majority of samples K8.1 expression remained unchanged or changes

were also reflected in LANA transcripts (Fig 3G).

IL21 receptor distribution in primary human tonsil B lymphocytes affects the magnitude

and distribution of KSHV infection

Our data presented thus far demonstrates that IL-21 signaling has a positive effect on the overall

establishment of KSHV infection (Fig 2C and 3B) and this increase in overall infection is related

to both the absolute number of plasma cells (Fig 2D&F and 3C), and the establishment of infection

in plasma cells (Fig 2E&F and 3E). Moreover, our data suggests that the increase in plasma cell

numbers and targeting may be due to differentiation of new plasma cells via a process that requires

IL-21 signaling to germinal center B cells (Fig 3E). To further address the early stages of the IL-

21 response during infection, we examined expression of the IL21 receptor in primary human

tonsil B lymphocytes at baseline (day 0) in each tonsil specimen. We observed that IL-21 receptor

expression is rare on B cells in tonsil at less than 3% of total viable B cells in most samples (Fig

4A). The distribution of IL-21 receptor positive cells among B cell sub-populations is broad, but

IL-21 receptor-expressing B cells are most likely to have an MZ-like or plasmablast

immunophenotype (Fig 4B) and on a per-sample basis, either plasmablast or MZ-like subsets

dominated the IL-21R positive cells in most tonsil samples (Fig 4C)

In order to determine whether IL-21 receptor expression at baseline influenced the establishment

of KSHV infection, we aggregated the untreated conditions from both the supplementation and the

neutralization experiments and examined correlations between baseline IL21R distribution and

KSHV infection based on overall GFP. This analysis revealed that the proportion of plasmablasts

within IL21R+ B cells is significantly correlated with overall susceptibility to KSHV infection

(r=0.81, p=0.0007) (Fig 4D). In experiments where we supplemented cultures with IL-21, the

positive effect of IL-21 treatment on overall KSHV infection at 3 dpi (Fig 2C) is correlated with

the baseline frequency of IL-21 receptor expression on naïve B cells (r=0.82, p=0.01) and classical

memory B cells (r=0.7, p=0.05) (Fig 4E). The significant correlation between the IL-21 response

and baseline IL-21 receptor expression is particularly interesting and consistent with our

hypothesis that differentiation of naïve B cells into plasma cells contributes to the increase in

overall infection we see with IL-21 treatment (Fig 2D).

When we correlated baseline IL21R expression with the positive effect of IL-21 treatment on total

plasma cell numbers in KSHV-infected cultures (Fig 2D), no significant positive correlations were

found (Fig 4F). This lack of correlation suggests that the plasma cell response to IL-21 at 3 dpi in

KSHV infected cultures may be a product of IL-21 receptor up-regulation in response to infection

instead of intrinsic baseline levels of IL-21 on plasma cells or plasma cell precursors in our tonsil

lymphocyte cultures. Indeed, modulation of IL-21 receptor expression by KSHV infection could

be one mechanism for the synergistic promotion of plasma cell numbers we observe with both IL-

21 treatment and infection (Fig 2D).

In order to examine this hypothesis, we analyzed IL-21R expression on B cell subsets at 3 dpi in

our culture system with or without IL-21 supplementation to determine whether KSHV and/or IL-

21 can modulate the response to IL-21 during infection. These results reveal that the majority of

effects on IL-21R at 3 dpi are present in both Mock and KSHV-infected cultures, indicating they

are a product of the culture system and not driven by KSHV. However, the proportion of

plasmablasts within IL21R+ cells was significantly different comparing Mock to KSHV-infected

cultures without IL-21 treatment (p=0.03) and this difference was increased with the combination

of KSHV infection and IL-21 treatment (Fig 4G). This result may indicate that infection is

affecting IL21R expression on plasmablasts or that KSHV is infecting a precursor B cell and

driving differentiation of these cells into IL21R+ plasmablasts. Taken together with the fact that

IL-21R expression on plasmablasts at day 0 is highly correlated with overall infection (Fig 4D),

these data strongly suggest that plasmablasts play a role in the response of KSHV infection to IL-

21 treatment. Although not statistically significant, a similar trend can be observed for IL21R+

plasma cells while the converse trend is seen for IL21R positive germinal center cells (Fig 4G),

these observations are consistent with the data from our IL21 neutralization studies that suggests

germinal center cells may be differentiating into plasma cells in response to IL-21 signaling (Fig

3E), potentially via a plasmablast intermediate phenotype.

<u>Characterization of T cell subsets producing IL-21 in primary human tonsil B lymphocytes</u>

We next wanted to determine the source of native IL-21 secretion in our culture system, and

determine whether the production of IL-21 is affected by KSHV infection. To accomplish this, we

utilized an additional immunophenotyping panel for T cell subsets (Table 2 and Supplemental Fig

1) and performed intracellular cytokine staining (ICCS) to identify IL-21 producing T cells at 3

dpi in Mock and KHSV-infected cultures from 14 unique tonsil samples. This data reveals that IL-

21 secretion in T cells is highly variable between tonsil lymphocyte cultures, but is not significantly

affected by KSHV infection (Fig 5A). More of the IL-21+ cells were CD4+ T cells vs. CD8+ T

cells and this distribution was also not affected by KSHV infection (Fig 5B). When we examined

the subset-level distribution of IL-21 secretion within T cells, we observed that within CD4+ cells

CD45RO+, central memory, TFH and RoRγT+ cells were contributing most to IL-21 secretion.

Among CD8+ T cells, CD45RA+, stem cell memory, central memory and RoRγT+ cells were the

highest contributors to IL-21 secretion (Fig 5C). When we examined whether KSHV infection

altered the contribution of T cell subsets to IL-21 secretion in these cultures, we found that the

contribution of CD4+ CD45RA+, CD4+ RoRγT+ and CD8+ central memory subsets was

significantly decreased in KSHV-infected cultures vs. Mock cultures (Fig 5D). However, KSHV

infection did not significantly change the overall levels and subset distribution of T cells within

these cultures, indicating these are biological changes within T cell subsets and not due to changes

in the T cell population during infection (Supplemental Fig 2).

Table 2: Lineage definitions for lymphocyte subsets used in the study	
B Lymphocytes	
Subset	**Molecular Markers**
Plasma	CD19$^+$**,** CD20$^{+/-}$, CD138$^{+(Mid\ to\ High)}$, CD38$^-$
Transitional	CD19$^+$**,** CD138$^-$, CD38^{Mid}, IgD$^{+\ (Mid\ to\ High)}$
Plasmablast	CD19$^+$**,** CD138$^-$, CD38^{High}, IgD$^{+\ /-\ (mostly\ -)}$
Germinal Center	CD19$^+$**,** CD138$^-$, CD38^{Mid}, IgD$^-$
Naïve	CD19$^+$**,** CD138$^-$, CD38^{Low}, CD27$^-$, IgD$^{+\ (Mid\ to\ High)}$
Marginal Zone Like (MZ-Like)	CD19$^+$**,** CD138$^-$, CD38^{Low}, CD27$^{+\ (Mid\ to\ High)}$, IgD$^{+\ (Mid\ to\ High)}$
Memory	CD19$^+$**,** CD138$^-$, CD38^{Low}, CD27$^{+\ (Mid\ to\ High)}$, IgD$^-$
Double Negative	CD19$^+$**,** CD138$^-$, CD38^{Low}, CD27$^-$, IgD$^-$
T lymphocytes	
Subset	**Molecular Markers**
CD4+	CD19$^-$, CD4$^{+(Mid\ to\ High)}$**,** CD8$^-$
CD8+	CD19$^-$, CD4$^-$, CD8$^{+\ (Mid\ to\ High)}$
Naïve	CD19$^-$, CD4+ or CD8+, CCR7$^{+(High)}$, CD45RA$^{+(Mid\ to\ High)}$, CD45RO$^-$, CD28$^+$, CD95$^-$
Stem Cell Memory	CD19$^-$, CD4+ or CD8+, CCR7$^{+(High)}$, CD45RA$^{+(Mid\ to\ High)}$, CD45RO$^-$, CD28$^+$, CD95$^{+\ (Low\ to\ Mid)}$
Central Memory	CD19$^-$, CD4+ or CD8+, CCR7$^+$, CD45RA$^-$ CD45RO$^{+(Mid\ to\ High)}$, CD28$^{+\ (Mid\ to\ High)}$
Transitional Memory	CD19$^-$, CD4+ or CD8+, CCR7$^-$, CD45RA$^-$ CD45RO$^{+(Mid)}$, CD28$^{+\ (Mid\ to\ High)}$
Effector Memory	CD19$^-$, CD4+ or CD8+, CCR7$^-$, CD45RA$^-$ CD45RO$^{+(Mid)}$, CD28$^-$
Terminal Effector Memory	CD19$^-$, CD4+ or CD8+, CCR7$^-$, CD45RA$^-$ CD45RO$^-$, CD28$^-$
TEMRA CD4+ Cells	CD19$^-$, CD4+ or CD8+, CCR7$^-$, CD45RA$^{+(High)}$**,** CD45RO$^-$, CD28$^-$
Tfh	CD19-, CD4+, CD8- , PD-1+, CXCR5+, CD127+, Intracellular BCL-6+
Treg	CD19-, CD4+, CD8- , CD25+, CD127+, Intracellular FoxP3+
Th17	CD19-, CD4+, CD8- , Intracellular ROR gamma T+

IL-21 secretion by CD8+ central memory T cells influences both overall KSHV infection and plasma cell targeting

We wanted to determine whether IL-21 secretion by any particular T cell subset was correlated with susceptibility to KSHV infection in our experiments. To do this, we performed B cell immunophenotyping analysis to determine the extent and distribution of KSHV infection in the same cultures where ICCS was performed on T cell subsets. When we examined correlations between IL-21 secretion by T cell subsets and overall KSHV infection in B cells we found that only IL-21+ CD8+ central memory cells were significantly correlated (r=0.57, p=0.03) (Fig 6A). Interestingly, when we examined correlations between IL-21+ T cell subsets and the B cell subset-specific distribution of KSHV infection at 3dpi, we found that the positive correlation between IL-21+ CD8+ central memory T cells is coupled with a highly significant positive correlation to the targeting of plasma cells (r=0.82, p=0.0003) (Fig 6B & 6C). In addition, both CD4+ and CD8+ IL-21+ T cells that express RoRγT+ (the Th17/Tc17-defining transcription factor) were significantly correlated with plasma cell targeting by KSHV (Fig 6B & 6D). These correlations were stronger for CD4+ cells and were coupled with a negative correlation with KSHV-infection of naïve T cells (Fig 6B). When we examined whether baseline (day 0) frequencies of T cell subsets influenced the magnitude and distribution of KSHV infection at 3dpi we found several interesting correlations (1) frequencies of CD8+ central memory cells significantly correlated with KSHV targeting of plasma cell subsets (Fig 6E and 6F), (2) baseline frequencies of CD4+ RoRγT+ cells significantly correlated with infection of CD20+ plasma cells (Fig 6D) and (3) both CD4+ and CD8+ RoRγT+ were negatively correlated with KSHV infection of naïve B cells at 3dpi (Fig 6D). Taken together, these results support a hypothetical model in which CD8+ central memory T cells secrete IL-21, which supports the differentiation, survival or targeting of plasma cells, resulting in

higher plasma cell infection frequencies and also an overall increase in dissemination of KSHV

within B lymphocytes. Our data also support the conclusion that RoRγT+ T cells are participating

in KSHV targeting of plasma cells via IL-21 signaling, possibly by mediating differentiation of

naïve B cells, but this effect does not have a significant influence on overall infection in this data

set.

Discussion

Our results presented in this study indicate that KSHV can influence cytokine production in tonsil-

derived lymphocytes and that the host inflammatory state contributes to the dramatic variation in

susceptibility we observe among our tonsil lymphocyte specimens [12]. This result is not

surprising considering dysregulation of the inflammatory environment is a hallmark of all KSHV-

associated malignancies [24, 26, 27]. However, the role of the baseline inflammatory environment

in the oral cavity as a potentially modifiable susceptibility factor for the acquisition of KSHV

infection in humans is an interesting consideration stemming from these results that deserves

further study.

Our previous work demonstrated that plasma cells are highly targeted during early KSHV

infection, and subsequent results from our group have shown that manipulations which increase

plasma cell numbers also increase overall KSHV infection in B cells [28]. This latter observation

suggested that, in addition to being a highly targeted cell type, plasma cells are playing an active

role in early infection events that ultimately influences the success of initial dissemination for

KSHV within the B cell compartment. The results presented herein are consistent with that

conclusion, and mechanistically extend our understanding of how the immune microenvironment

influences the establishment of infection in plasma cells. Specifically, we uncovered a critical role

for IL-21 signaling in this process.

While the precise role of IL-21 and IL-21R in human KSHV disease is poorly characterized, IL-

21 has been studied in the pathogenesis of chronic lymphocytic choriomeningitis virus (LCMV)

infection, influenza virus, and, perhaps most relevant to this study, murine gammaherpesvirus 68

(MHV68) [22, 29, 30]. Collins and Speck recently used IL-21R knockout mice to demonstrate that

IL-21 signaling is critical for the establishment of MHV68 latency specifically in B cells.

Interestingly, this study showed that the mechanisms of decreased infection were related to

decreases in both germinal center and plasma cell frequencies as well as decreased infection in

both the germinal center and plasma cell compartment at later timepoints post-infection [22],

suggesting a critical mechanism for IL-21 in MHV68 transit of the germinal center and

differentiation of follicular-derived plasma cells. These results are generally consistent with our

current findings showing that IL-21 signaling promotes KSHV infection and dissemination in

tonsil via promotion of both plasma cell frequencies and plasma cell targeting (Fig 2&3),

potentially via differentiation through a germinal center intermediate (Fig 3E). Despite this, our

data presented herein do not adequately address the question of whether plasma cells are

differentiating in response to KSHV infection and IL-21 signaling, or whether the effect we see is

increased survival or expansion of existing plasma cells. Studies are currently ongoing to address

this question more directly, and to examine the influence of viral and cellular factors on B cell

differentiation in our model system.

Moreover, both IL-21 and IL-6, which is highly induced in our KSHV-infected cultures (Fig 1),

are involved in the generation of RoRγT+ T cells via STAT3 signaling. The resulting Th17 cells

produce IL-17A which is another cytokine that promotes the establishment of chronic MHV68

infection via promotion of the MHV68-mediated germinal center response [31] and is

mechanistically linked to suppression of T cell-intrinsic IRF-1 [32]. These results are particularly

interesting in light of our current findings showing that IL-21 secretion from, and baseline levels

of, RoRγT+ T cells correlate with the early targeting of plasma cells during KSHV infection (Fig

6B and 6D), suggesting that the Th17/Tc17 environment in the tonsil may be a critical factor

influencing donor-specific susceptibility to KSHV infection. Indeed, as an important site for

mucosal immunity in the oral cavity, the Th17/Tc17 environment in tonsil is highly dynamic and

physiologically important. In fact, Th17 cells play a major role in host defenses against several

pathogens and immunopathogenesis [33, 34]. Many studies have shown that certain parasites

modulate the immune response by inducing Th17 [35, 36]. Previous finding suggest that parasite

infection is linked with KSHV infection in Uganda [37]. Thus, the parasite burden in sub-Saharan

Africa may modulate susceptibility to KSHV infection via manipulating Th17/Tc17 frequencies.

Consistent with the MHV68 literature, our current results mechanistically implicate germinal

center cells in these observations. However, although our *ex vivo* model of KSHV infection in

primary lymphocytes is a powerful tool, it certainly does not recapitulate the complex interactions

that are needed for a functional germinal center reaction, so further examination of these particular

mechanisms will require the utilization of an alternative model system, such as a humanized

mouse. Interestingly, neither IL-21 secretion from, nor baseline levels of, CXCR5+, PD1+ TFH

cells correlated with increased KSHV infection or increased plasma cell targeting in our study.

This was surprising given that TFH are generally considered the canonical IL-21 producing cells

in secondary lymphoid organs [38]. This observation may indicate that IL-21 acts support plasma

cell numbers during infection via an extrafollicular pathway, which is consistent with literature

implicating extrafollicular maturation of KSHV-infected B cells in the pathogenesis of MCD [39].

We observed that IL-21 secretion by CD8+ T cells with a central memory immunophenotype

(CCR7+CD45RO+CD28+) was significantly correlated with both overall KSHV infection and

plasma cell targeting, and baseline levels of CD8+ central memory T cells were also correlated

with plasma cell targeting at 3 dpi (Fig 6). Interestingly, recent studies have shown that IL-6

regulates IL-21 production in CD8+ T cells in a STAT3-dependent manner, and that CD8+ T cells

induced in this way can effectively provide help to B cells [40]. Thus, the induction of human IL-

6 during KSHV infection may modulate the function of CD8+ T cells in a way that favors the

establishment and dissemination of KSHV infection within the lymphocyte compartment

independent of traditional CD4+ helper T cells, which would be an interesting dynamic in the

context of CD4+ T cell immunosuppression associated with HIV infection where KSHV-mediated

malignancies are common.

Material and Methods

Ethics Statement. Human specimens used in this research were de-identified prior to receipt, and

thus were not subject to IRB review as human subjects research.

Reagents and Cell Lines. CDw32 L cells (CRL-10680) were obtained from ATCC and were

cultured in DMEM supplemented with 20% FBS (Sigma Aldrich) and Penicililin/Streptomycin/L-

glutamine (PSG/Corning). For preparation of feeder cells CDw32 L cells were trypsinized and

resuspended in 15 ml of media in a petri dish and irradiated with 45 Gy of X-ray radiation using a

Rad-Source (RS200) irradiator. Irradiated cells were then counted and cyropreserved until needed

for experiments. Cell-free KSHV.219 virus derived from iSLK cells [39] was a gift from Javier G.

Ogembo (City of Hope). Human tonsil specimens were obtained from the National Disease

Research Interchange (NDRI; ndriresource.org). Human fibroblasts for viral titering were derived

from primary human tonsil tissue and immortalized using HPV E6/E7 lentivirus derived from

PA317 LXSN 16E6E7 cells (ATCC CRL-2203). Antibodies for flow cytometry were from BD

Biosciences and Biolegend and are detailed below.   Recombinant human IL-21 was from

Preprotech (200-21) and IL-21 neutralizing antibody was from R&D Systems (991-R2).

***Isolation of primary lymphocytes from human tonsils*.** De-identified human tonsil specimens

were obtained after routine tonsillectomy by NDRI and shipped overnight on wet ice in

DMEM+PSG. All specimens were received in the laboratory less than 24 hours post-surgery and

were kept at 4˚C throughout the collection and transportation process.   Lymphocytes were

extracted by dissection and maceration of the tissue in RPMI media. Lymphocyte-containing

media was passed through a 40µm filter and pelleted at 1500rpm for 5 minutes. RBC were lysed

for 5 minutes in sterile RBC lysing solution (0.15M ammonium chloride, 10mM potassium

bicarbonate, 0.1M EDTA). After dilution to 50ml with PBS, lymphocytes were counted, and

pelleted. Aliquots of 5(10)7 to 1(10)8 cells were resuspended in 1ml of freezing media containing

90% FBS and 10% DMSO and cryopreserved until needed for experiments.

***Infection of primary lymphocytes with KSHV*.** Lymphocytes were thawed rapidly at 37˚C, diluted

dropwise   to   5ml   with   RPMI   and   pelleted.   Pellets   were   resuspended   in   1ml

RPMI+20%FBS+100µg/ml DNaseI+ Primocin 100µg/ml and allowed to recover in a low-binding

24 well plate for 2 hours at 37˚C, 5% CO2. After recovery, total lymphocytes were counted and

naïve B cells were isolated using Mojosort Naïve B cell isolation beads (Biolegend 480068) or

Naïve B cell Isolation Kit II (Miltenyi 130-091-150) according to manufacturer instructions.

Bound cells (non-naïve B and other lymphocytes) were retained and kept at 37°C in RPMI+20%

FBS+ Primocin 100μg/ml during the initial infection process. $1(10)^6$ Isolated naïve B cells were

infected with iSLK-derived KSHV.219 (dose equivalent to the ID20 at 3dpi on human fibroblasts)

or Mock infected in 400ul of total of virus + serum free RPMI in 12x75mm round bottom tubes

via spinoculation at 1000rpm for 30 minutes at 4°C followed by incubation at 37°C for an

additional 30 minutes. Following infection, cells were plated on irradiated CDW32 feeder cells in

a 48 well plate, reserved bound cell fractions were added back to the infected cell cultures, and

FBS and Primocin (Invivogen) were added to final concentrations of 20% and 100μg/ml,

respectively and recombinant cytokines or neutralizing antibodies were also added at this stage,

depending upon the specific experiment. Cultures were incubated at 37°C, 5% CO2 for the

duration of the experiment. At 3 days post-infection, cells were harvested for analysis by flow

cytometry and supernatants were harvested, clarified by centrifugation for 15 minutes at 15,000

rpm to remove cellular debris, and stored at -80°C for analysis.

Bead-based immunoassay for supernatant cytokines. Clarified supernatants were thawed on ice

and 25μl of each was added to a 13-plex LEGENDplex (Biolegend) bead-based immunoassay

containing capture beads for the following analytes: IL-5, IL-13, IL-2, IL-9, IL-10, IL17A, IL-

17F, IL-6, IL-21, IL-22, IL-4, TNF-α, and IFN-γ. These assays were performed according to the

manufacturer's instructions, data was acquired for 5000 beads per sample (based on approximately

300 beads per analyte recommended by the manufacturer) using a BD FACS VERSE flow

cytometry analyzer and cytokine concentrations in the experimental supernatants was calculated

from standard curves using the LEGENDPlex software.

Flow cytometry analysis of baseline lymphocyte subsets and KSHV infection. Approximately

$5(10)^6$ lymphocytes per condition were harvested into a 96- well round bottom plate at day 0

(baseline) or at 3 days post-infection at 1500 rpm for 5 minutes. Cells were resuspended in 100µl

PBS containing zombie violet fixable viability stain (BL Cat# 423113) and incubated on ice for

15 minutes. After incubation, cells were pelleted and resuspended in 100ul PBS, containing the

following: 2% FBS and 0.5% BSA (FACS Block) was added to the wells. Cells were pelleted at

1500rpm 5 minutes and resuspended in 200ul FACS Block for 10 minutes on ice. Cells were

pelleted at 1500rpm for 5 minutes and resuspended in 50µl of PBS with 0.5% BSA and 0.1%

Sodium Azide (FACS Wash), **For B cell frequencies** 10µl BD Brilliant Stain Buffer Plus and

antibodies as follows: IgD-BUV395 (2.5µl/test BD 563823), CD77-BV510 (2.0 µl/ test BD

563630), CD138- BV650 (2µl/test BD 555462), CD27-BV750 (2µ/test BD 563328), CD19-

PerCPCy5.5 (2.0µl/test BD 561295), CD38-APC (10µl/test BD 560158), CD20-APCH7 (2ul/test

BL 302313), IgM (2µl/test BL 314524), IgG (2µl/test BD 561298), IgE (2µl/test BD 744319) and

IL-21 receptor (2µl/test BD 330114). **For baseline T cell frequencies**. For baseline T cell

frequencies $0.5(10)^6$ cells from baseline uninfected total lymphocyte samples were stained and

analyzed as above with phenotype antibody panel as follows: CD95-APC (2µl, Biolegend

305611), CCR7-PE (2µl, BD 566742), CD28-PE Cy7 (2µl, Biolegend 302925), CD45RO-FITC

(3µl, Biolegend 304204), CD45RA-PerCP Cy5.5 (2µl, 304121), CD4-APC H7 (2µl, BD 560158),

CD19-V510 (3µl, BD 562953), CD8-V450 (2.5µl, BD 561426). and incubated on ice for 15

minutes. After incubation, 150µl FACS Wash was added. Cells were pelleted at 1500rpm for 5

minutes followed by two washes with FACS Wash. Cells were collected in 200µl FACS Wash for

flow cytometry analysis. Cells were analyzed using an LSR Fortessa X-20 cell analyzer (BD

Biosciences). BD CompBeads (51-90-9001229) were used to calculate compensation for all

antibody stains and methanol-fixed Namalwa cells (ATCC CRL1432) +/- KSHV were used to

calculate compensation for GFP and the fixable viability stain. Flow cytometry data was analyzed

using FlowJo software and exported for quantitative analysis in R as described below.

ICCS for IL-21 secretion. At 3dpi, cultures were treated for 6 hours with [4 ul for every 6ml of

cel culture] monensin to block cytokine secretion. Following incubation, approximately 1 million

cells were harvested and viability and surface staining for T cell lineage markers was performed

as described above. After the final wash, cells were fixed for 10 minutes in BD cytofix/cytoperm

(51-2090KZ), pelleted and further treated for 10 minutes with cytofix/cytoperm+10% DMSO

(superperm) to more effectively get intracellular antibodies into the nucleus. Intracellular

antibodies, as follows, were diluted in 1x BD Permwash (51-2091KZ) and left on fixed cells

overnight at 4˚C. RoR-γT-BV421 (563282, 5µl/test), FoxP3-BB700 (566527, 5 µl/test), IL-21-

APC (513007, 5 µl/test), BLC6-BV711 (561080, 5µl/test). Cells were then washed twice with 1x

permwash and analyzed as described above.

RT-PCR. At 3 days post infection, $1(10)^6$ lymphocytes were harvested into an equal volume of

Trizol and DNA/RNA shield (Zymo Research R110-250). Total RNA was extracted using using

Zymo Directzol Microprep (Zymo Research R2060) according to manufacturer instructions. RNA

was eluted in 10µl H2O containing 2U RNase inhibitors and a second DNase step was performed

for 30 minutes using the Turbo DNA-Free kit (Invitrogen AM1907M) according to manufacturer

instructions. One-step RT-PCR cDNA synthesis and preamplification of GAPDH, LANA and

K8.1 transcripts was performed on 15ng of total RNA using the Superscript III One-step RT-PCR

kit (ThermoFisher 12574026).

Duplicate no RT (NRT) control reactions were assembled for each sample containing only

Platinum Taq DNA polymerase (Thermofisher 15966005) instead of the Superscript III RT/Taq

DNA polymerase mix. After cDNA synthesis and 20 cycles of target pre-amplification, 2μl of pre-

amplified cDNA or NRT control reaction was used as template for multiplexed real-time PCR

reactions using TaqProbe 5x qPCR MasterMix -Multiplex (ABM MasterMix-5PM), 5% DMSO,

primers at 900nM and probes at 250nM against target genes. All primer and probe sequences used

in these assays have been previously published [12]. Real time PCR was performed using a 40-

cycle program on a Biorad real time thermocycler. Data is represented as quantitation cycle (Cq)

and assays in which there was no detectable Cq value were set numerically as Cq = 41 for analysis

and data visualization. The expression of each gene was normalized to that of a housekeeping

gene *GAPDH*.

Statistical Analysis. The indicated data sets and statistical analysis were performed in Rstudio

software using ggplot2 [41], ggcorrplo [42] , and tidyverse [43] packages. Statistical analysis was

performed using rstatix [44] package. Specific methods of statistical analysis including Anova,

independent t-test and Pearson correlations and resulting values for significance and correlation

are detailed in the corresponding figure legends.

Figure Legends

Figure 1: KSHV alters cytokine secretion and cytokines affect the establishment of infection.

33 replicate infections using 24 unique tonsil specimens were performed using KSHV.219

infection of naïve B lymphocytes followed by reconstitution of the total lymphocyte environment

and culture on CDW32 feeder cells. At 3 dpi, cells were collected for flow cytometry analysis for

infection (GFP) and B cell subsets using our previously-characterized immunophenotyping panel

and supernatants were collected for analysis of cytokines by multiplex immunoassay (Biolegend

Legendplex) **(A)** cytokine production in Mock and KSHV-infected cultures showing individual

sample quantities and means (red diamonds, top panels) and matched Mock and KSHV samples

to show trends of induction/repression (bottom panels) Statistical analysis was performed by one-

way repeated measures ANOVA. p=0.01 F=7.06 for IL-5, p=0.0001 F=14 for IL-6, p=0.5 F=4.3

for IL-4   **(B)** Data as in (A) showing the level of induction or repression of each cytokine

comparing KSHV to matched Mock cultures **(C)** Induction or repression of all cytokines (left y-

axis) on a per-sample basis ordered based on overall susceptibility based on percentage of GFP+

B lymphocytes in the same culture (right y-axis, red diamonds) **(D)** Pairwise correlations using

Pearson method between cytokine concentration (y-axis) and overall infection (x-axis) in KSHV-

infected lymphocyte cultures **(E)** Pairwise correlations using Pearson method between cytokine

concentration (y-axis) and Percent GFP+ within CD138+ (x-axis) in KSHV-infected lymphocyte

cultures. For panels A, B, D and E colors indicate individual tonsil specimens and can be compared

between panels.

Figure 2: IL-21 supplementation increases overall KSHV infection and plasma cell

frequencies. Lymphocytes from 12 tonsil donors were infected with KSHV.219 and cultured with

indicated doses of recombinant human IL-21 and analyzed at 3dpi by flow cytometry (A) the dose

effect of IL-21 supplementation on GFP+ viable B lymphocytes. (B) data as in (A) normalized to

the untreated control for each specimen. (C) 14 additional tonsil donors analyzed as in (A) with

only 100pg/ml IL-21 treatment. Red diamonds indicate group means. p=0.02, F=6.4 via one-way

repeated measures ANOVA. (D) tonsil lymphocyte specimens from (C) were stained for B cell

immunophenotypes and analyzed by flow cytometry. Red boxes indicate subsets with significant

main effects of, and/or interactions between, KSHV infection and IL-21 treatment by two-way

repeated measures ANOVA analysis. See Table 1 for p-values and F statistics from this analysis.

(E) immunophenotyping data was used to determine the within-subset frequency of GFP+ cells in

the KSHV-infected conditions with or without IL-21 treatment. Top panels show individual

sample quantities and means (red diamonds) and bottom panels show trends of

increased/decreased subset targeting on a per-sample basis. Colored points denote unique tonsil

specimens and can be compared between panels D and E. (F) Pearson correlation between overall

GFP+ B cells in KSHV-infected, IL-21 treated cultures as in (C) and the level of plasma cells

(right) and infection of plasma cells (left). Blue line is linear model regression and grey shading

indicates 95% confidence interval.

Figure 3: IL-21 neutralization inhibits the establishment of KSHV infection. Naïve B cells

from 11 unique tonsil specimens were Mock or KSHV-infected and indicated concentrations of

IL-21 neutralizing antibody was added to the resulting total lymphocyte cultures. Cultures were

analyzed at 3 dpi for B lymphocyte immunophenotypes and the distribution of KSHV infection

via GFP expression. Total GFP+ viable B lymphocytes represented as (A) raw percentages or (B)

normalized to the untreated control for each tonsil sample. For (A) one-way repeated measures

ANOVA shows p=0.00001, F=9.4 for the main effect of IL-21 neutralization and Dunnett's test

reveals p=0.03 at the 100µg/ml dose. (C) frequencies of B cell subsets in the cultures. Red boxes

indicate significant results as follows: Two-way repeated measures ANOVA on the raw data

showed significant effects of IL-21 treatment on B subset frequencies; IgG+ (p=0.04, F=5.0),

memory (p=0.03, F=3.4), and double negative (p=0.02, F=3.8), IgM+ (p=0.0003, F=8.6),

transitional (p=0.01, F=4.1). Post-hoc paired T-tests showed significant effect of 200µg/ml

neutralizing antibody on total plasma cells (p=0.02) and CD20+ plasma cells (p=0.005) in Mock

cultures only. (D) Effect of indicated doses of IL-21 neutralizing antibody on KSHV infection of

B cell subsets. Data is normalized to each sample's untreated control level for each B cell subset.

One-way repeated measures ANOVA on the raw data reveals a significant effect on infection of

transitional B cells (p=0.01, F=4.0). (E) Correlation between the effect of IL-21 neutralization on

overall infection (x-axis) and the contribution of each subset to KSHV infection in the same

condition (y-axis). Shapes indicate doses in this panel (circle=100µg/ml, triangle=200µg/ml,

square=400µg/ml). Subsets with statistically significant correlations are indicated by red boxes

with statistics from Pearson's linear correlation as follows: CD20+ plasma cells (r=0.6, p=0.0002),

plasma cells (r=0.6, p=0.0003), transitional (r=0.5, p=0.004), germinal center (r=-0.4, p=0.02).

For panels C-E colors indicate unique tonsil specimens and can be compared between these panels

and red diamonds indicate the mean value for all tonsil specimens. RT-PCR analysis of KSHV

transcripts at 3 dpi in 8 unique tonsil specimens with either IL-21 supplementation at 100 pg/ml

(Fig 2) or IL-21 neutralizing antibody at 100µg/ml with LANA (latent) and K8.1 (lytic) transcript

targets. No RT controls were used to determine that RT-PCR signal is not due to DNA

contamination (F) Cq values for viral targets normalized to the within-sample Cq for GAPDH (G)

GAPDH-normalized values further normalized to the within-sample value for the untreated

control.

Figure 4: IL21 receptor distribution in primary human tonsil B lymphocytes affects the

magnitude and distribution of KSHV infection. B cell immunophenotyping analysis including

IL-21R was performed at baseline (Day 0) for 10 unique tonsil specimens. (A) total percentage of

IL-21R+ within viable CD19+ B cells. (B) Percent of individual B cell subsets within IL-21+ B

cells. Red diamonds indicate the mean value for all tonsil specimens and (C) distribution of B cell

subsets within IL-21+ on a per-tonsil basis. (D) correlation analysis of baseline IL-21 receptor

distribution within subsets and the overall susceptibility of the same tonsil specimen to KSHV

infection at 3 dpi. Pearson correlation coefficients and p-values are denoted on panels where the

result is statistically significant. (E) correlation analysis of baseline IL-21 receptor distribution

with the effect of IL-21 supplementation on overall KSHV infection in the same tonsil specimens.

Pearson correlation coefficients and p-values are denoted on panels where the result is statistically

significant. (F) correlation analysis of baseline IL-21 receptor distribution with the IL-21 mediated

increase in total plasma cell numbers at 3dpi in KSHV-infected cultures in the same tonsil

specimens. (G) Distribution of IL-21 receptor at day 0 (baseline) or 3dpi within Mock,

Mock+100ng/ml IL-21, KSHV or KSHV+ 100ng/ml IL-21 conditions. Red diamonds indicate the

mean values for each condition and significant differences were assessed via paired T-test.

Figure 5: Characterization of T cell subsets producing IL-21 in primary human tonsil B

lymphocytes. T cells were analyzed by surface immunophenotyping, intracellular transcription

factor staining and ICCS for IL-21 secretion in Mock and KSHV-infected total lymphocyte

cultures at 3 dpi in 14 unique tonsil specimens. (A) Total IL-21+ viable non-B cells (B) percent of

CD4+ or CD8+ T cells within IL-21+. For (A) and (B) red diamonds indicate the mean value for

the condition and colors indicate specific tonsil specimens and can be compared between the

panels. (C) Immunophenotypic analysis of T cell subsets (Table 2) within IL-21+ in Mock (red)

and KSHV (blue) cultures. (D) Data as in (C) but normalized to show the induction or repression

of the T cell subset within IL-21+ cells (KSHV-Mock). *p=0.05; **p=0.04; ***p=0.003

comparing Mock and KSHV-infected via paired T test pre-normalization.

Figure 6: IL-21 secretion by CD8+ central memory T cells influences both overall KSHV

infection and plasma cell targeting. Lymphocyte cultures from the experiments shown in Fig 5

were further analyzed for B cell subsets and the magnitude and distribution of KSHV infection.

(A) correlation analysis of total GFP+ within viable, CD19+ B lymphocytes at 3 dpi with total IL-

21 secretion in non-B cells or the contribution of individual T cell subsets to IL-21 secretion.

Pearson's correlation coefficient and p-value is shown on panels that have statistically significant

correlations. (B) correlogram of Pearson correlations between the distribution of KSHV infection

within B cell subsets (y-axis) and total IL-21 or the contribution of individual T cell subsets to IL-

21 secretion (x-axis). Pearson's r values with an absolute value greater than or equal to 0.53 are

statistically significant for this dataset. (C) scatterplot and linear model fit for the correlation

between GFP+ plasma cells and IL-21 secretion by CD8+ central memory T cells (D) Scatterplots

and linear model fits for the correlations between GFP+ plasma cells and secretion of IL-21 by

RoRγT+ CD4+ (left) or CD8+ (right) T cells. (E) correlogram of Pearson correlations between the

distribution of KSHV infection within B cell subsets (y-axis) and baseline (Day 0) levels of T cell

subsets. Pearson's r values with an absolute value greater than or equal to 0.53 are statistically

significant for this dataset. (F) scatterplot and linear model fit for the correlation between GFP+

plasma cells and baseline frequency of CD8+ central memory T cells. For A, C, D and F grey

shading indicates 95% confidence intervals.

Supplemental Figure 1: Representative gating scheme for T cell immunophenotyping using

lineage definitions as detailed in Table 2.

Supplemental Figure 2: Total T cell subset frequencies in Mock and KSHV-infected cultures as

in Figure 5.

References

1. Chang, Y., et al., *Identification of herpesvirus-like DNA sequences in AIDS-associated*
Kaposi's sarcoma. Science, 1994. **266**(5192): p. 1865-1869.
2. Ablashi, D.V., et al., *Spectrum of Kaposi's sarcoma-associated herpesvirus, or human*
herpesvirus 8, diseases. Clinical microbiology reviews, 2002. **15**(3): p. 439-464.
3. Cesarman, E., et al., *Kaposi's sarcoma–associated herpesvirus-like DNA sequences in*
AIDS-related body-cavity–based lymphomas. New England Journal of Medicine, 1995.
332(18): p. 1186-1191.
4. Soulier, J., et al., *Kaposi's sarcoma-associated herpesvirus-like DNA sequences in*
multicentric Castleman's disease [see comments]. Blood, 1995. **86**(4): p. 1276-1280.
5. Uldrick, T.S., et al., *An interleukin-6-related systemic inflammatory syndrome in patients*
co-infected with Kaposi sarcoma-associated herpesvirus and HIV but without
Multicentric Castleman disease. Clinical Infectious Diseases, 2010. **51**(3): p. 350-358.
6. Bouvard, V., et al., *A review of human carcinogens--Part B: biological agents.* The
Lancet. Oncology, 2009. **10**(4): p. 321-322.
7. Parkin, D.M., *The global health burden of infection-associated cancers in the year 2002.*
International journal of cancer, 2006. **118**(12): p. 3030-3044.
8. Ganem, D., *KSHV and the pathogenesis of Kaposi sarcoma: listening to human biology*
and medicine. The Journal of clinical investigation, 2010. **120**(4): p. 939-949.
9. Casper, C., et al., *Frequent and asymptomatic oropharyngeal shedding of human*
herpesvirus 8 among immunocompetent men. The Journal of infectious diseases, 2007.
195(1): p. 30-36.
10. Alomari, N. and J. Totonchy, *Cytokine-Targeted Therapeutics for KSHV-Associated*
Disease. Viruses, 2020. **12**(10): p. 1097.
11. Carbone, A., et al., *Understanding pathogenetic aspects and clinical presentation of*
primary effusion lymphoma (PEL) through its derived cell lines. AIDS (London,
England), 2010. **24**(4): p. 479.
12. Aalam, F., et al., *Analysis of KSHV B lymphocyte lineage tropism in human tonsil reveals*
efficient infection of CD138+ plasma cells. PLoS pathogens, 2020. **16**(10): p. e1008968.
13. Parrish-Novak, J., et al., *Interleukin-21 and the IL-21 receptor: novel effectors of NK and*
T cell responses. Journal of leukocyte biology, 2002. **72**(5): p. 856-863.
14. Parrish-Novak, J., et al., *Interleukin 21 and its receptor are involved in NK cell expansion*
and regulation of lymphocyte function. Nature, 2000. **408**(6808): p. 57-63.
15. Kuchen, S., et al., *Essential role of IL-21 in B cell activation, expansion, and plasma cell*
generation during CD4+ T cell-B cell collaboration. The journal of immunology, 2007.
179(9): p. 5886-5896.
16. Konforte, D., N. Simard, and C.J. Paige, *IL-21: an executor of B cell fate.* The Journal of
Immunology, 2009. **182**(4): p. 1781-1787.
17. Ozaki, K., et al., *Regulation of B cell differentiation and plasma cell generation by IL-21,*
a novel inducer of Blimp-1 and Bcl-6. The Journal of Immunology, 2004. **173**(9): p.
5361-5371.

18.   Calame, K.L., K.-I. Lin, and C. Tunyaplin, *Regulatory mechanisms that determine the*
development and function of plasma cells. Annual review of immunology, 2003. **21**(1): p.
205-230.
19.   Yajima, H., et al., *Loss of interleukin-21 leads to atrophic germinal centers in*
multicentric Castleman's disease. Annals of hematology, 2016. **95**(1): p. 35-40.
20.   Konforte, D. and C.J. Paige, *Identification of cellular intermediates and molecular*
pathways induced by IL-21 in human B cells. The Journal of Immunology, 2006. **177**(12):
p. 8381-8392.
21.   Konforte, D., N. Simard, and C.J. Paige, *Interleukin-21 regulates expression of key*
Epstein–Barr virus oncoproteins, EBNA2 and LMP1, in infected human B cells.
Virology, 2008. **374**(1): p. 100-113.
22.   Collins, C.M. and S.H. Speck, *Interleukin 21 signaling in B cells is required for efficient*
establishment of murine gammaherpesvirus latency. PLoS pathogens, 2015. **11**(4): p.
e1004831.
23.   Gasperini, P. and G. Tosato, *Targeting the mammalian target of Rapamycin to inhibit*
VEGF and cytokines for the treatment of primary effusion lymphoma. Leukemia, 2009.
23(10): p. 1867-1874.
24.   Gasperini, P., S. Sakakibara, and G. Tosato, *Contribution of viral and cellular cytokines*
to Kaposi's sarcoma-associated herpesvirus pathogenesis. Journal of leukocyte biology,
2008. **84**(4): p. 994-1000.
25.   Totonchy, J., et al., *KSHV induces immunoglobulin rearrangements in mature B*
lymphocytes. PLoS pathogens, 2018. **14**(4): p. e1006967.
26.   Chang, J., et al., *Induction of Kaposi's sarcoma-associated herpesvirus from latency by*
inflammatory cytokines. Virology, 2000. **266**: p. 17-25.
27.   Chang, J., et al., *Inflammatory cytokines and the reactivation of Kaposi's sarcoma-*
associated herpesvirus lytic replication. Virology, 2000. **266**(1): p. 17-25.
28.   Palmerin, N., et al., *Suppression of DC-SIGN and gH Reveals Complex, Subset-Specific*
Mechanisms for KSHV Entry in Primary B Lymphocytes. Viruses 2021, 13, 1512. 2021, s
Note: MDPI stays neutral with regard to jurisdictional claims in published ….
29.   Schmitz, I., et al., *IL-21 restricts virus-driven Treg cell expansion in chronic LCMV*
infection. PLoS pathogens, 2013. **9**(5): p. e1003362.
30.   Pallikkuth, S., et al., *Upregulation of IL-21 receptor on B cells and IL-21 secretion*
distinguishes novel 2009 H1N1 vaccine responders from nonresponders among HIV-
infected persons on combination antiretroviral therapy. The Journal of Immunology,
2011. **186**(11): p. 6173-6181.
31.   Jondle, C., et al., *Gammaherpesvirus Usurps Host IL-17 Signaling To Support the*
Establishment of Chronic Infection. Mbio, 2021. **12**(2): p. e00566-21.
32.   Jondle, C., et al., *T Cell-Intrinsic Interferon Regulatory Factor 1 Expression Suppresses*
Differentiation of CD4+ T Cell Populations That Support Chronic Gammaherpesvirus
Infection. Journal of Virology, 2021. **95**(20): p. e00726-21.
33.   Gaddi, P.J. and G.S. Yap, *Cytokine regulation of immunopathology in toxoplasmosis.*
Immunology and cell biology, 2007. **85**(2): p. 155-159.
34.   Kelly, M.N., et al., *Interleukin-17/interleukin-17 receptor-mediated signaling is*
important for generation of an optimal polymorphonuclear response against Toxoplasma
gondii infection. Infection and immunity, 2005. **73**(1): p. 617-621.

762 35. Shainheit, M.G., et al., *The pathogenic Th17 cell response to major schistosome egg*
763   *antigen is sequentially dependent on IL-23 and IL-1β.* The Journal of Immunology, 2011.
764   **187**(10): p. 5328-5335.
765 36. Wen, X., et al., *Dynamics of Th17 cells and their role in Schistosoma japonicum infection*
766   *in C57BL/6 mice.* PLoS neglected tropical diseases, 2011. **5**(11): p. e1399.
767 37. Wakeham, K., et al., *Parasite infection is associated with Kaposi's sarcoma associated*
768   *herpesvirus (KSHV) in Ugandan women.* Infectious agents and cancer, 2011. **6**(1): p. 1-7.
769 38. Song, W. and J. Craft, *T follicular helper cell heterogeneity: time, space, and function.*
770   Immunological reviews, 2019. **288**(1): p. 85-96.
771 39. Totonchy, J., *Extrafollicular activities: perspectives on HIV infection, germinal center-*
772   *independent maturation pathways, and KSHV-mediated lymphoproliferation.* Current
773   opinion in virology, 2017. **26**: p. 69-73.
774 40. Yang, R., et al., *IL-6 promotes the differentiation of a subset of naive CD8+ T cells into*
775   *IL-21–producing B helper CD8+ T cells.* Journal of Experimental Medicine, 2016.
776   **213**(11): p. 2281-2291.
777 41. Wickham, H., *Ggplot2 : elegant graphics for data analysis*. Use R! 2009, New York:
778   Springer. viii, 212 p.
779 42. Kassambara, A., *ggcorrplot: Visualization of a Correlation Matrix using 'ggplot2'.* 2019.
780 43. Lander, J.P., *R for everyone : advanced analytics and graphics*. Second edition ed. The
781   Addison Wesley data and anlytics series. 2017, Boston: Addison-Wesley. xxiv, 528
782   pages.
783 44. Kassambara, A., *rstatix: Pipe-Friendly Framework for Basic Statistical Tests.* 2020.
784

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6